Peroxiredoxin 2 Regulates DAF-16/FOXO Mediated Mitochondrial Remodelling in Response to Exercise that is Disrupted in Ageing

Qin Xia[1,2], Penglin Li[1,2], José C. Casas-Martinez[1,2], Antonio Miranda-Vizuete[,3], Emma McDermott[4], Peter Dockery[4], Katarzyna Goljanek-Whysall[1,2,5] and Brian McDonagh[1,2*]

[1]Discipline of Physiology, School of Medicine, [2]Apoptosis Research Centre, University of Galway, Ireland
[3]Instituto de Biomedicina de Sevilla, Hospital Universitario Virgen del Rocío/CSIC/Universidad de Sevilla, Sevilla, Spain
[4]Centre for Microscopy and Imaging, Discipline of Anatomy, School of Medicine, University of Galway, Ireland
[5]Institute of Lifecourse and Medical Sciences, University of Liverpool, UK

[*]Corresponding author: brian.mcdonagh@universityofgalway.ie

Abstract:
Ageing is associated with mitochondrial dysfunction and increased oxidative stress. Exercise generates endogenous reactive oxygen species (ROS) and promotes rapid mitochondrial remodelling. We investigated the role of Peroxiredoxin 2 (PRDX-2) in mitochondrial adaptations to exercise and ageing using *Caenorhabditis elegans* as a model system. PRDX-2 was required for the mitochondrial remodelling in response to exercise mediated by DAF-16 nuclear localisation. Employing an acute exercise and recovery cycle, we demonstrated exercise-induced mitochondrial ER contact sites (MERCS) assembly and mitochondrial remodelling dependent on PRDX-2 and DAF-16 signalling. There was increased mitochondrial fragmentation, elevated ROS and an altered redox state of PRDX-2 concomitant with impaired DAF-16 nuclear localisation during ageing. Similarly, the *prdx-2* mutant strain exhibited increased mitochondrial fragmentation and a failure to activate DAF-16 required for mitochondrial fusion. Collectively, our data highlight the critical role of PRDX-2 in orchestrating mitochondrial remodelling in response to a physiological stress by regulating DAF-16 nuclear localisation.

Keywords:

Ageing, Peroxiredoxin 2, Exercise, *C. elegans*, Mitochondrial ER Contact Sites, DAF-16

Graphical abstract:

Highlights:

- Exercise generates ROS and promotes mitochondrial remodelling dependent on DAF-16.
- Exercise induces mitochondrial ER contact site assembly and mitochondrial dynamics.
- Ageing and loss of PRDX-2 results in disrupted mitochondrial fusion.
- The redox state of PRDX-2 determines appropriate DAF-16 nuclear localisation.

1. Introduction

Ageing results in a gradual decline in physiological function, leading to compromised overall health and increased susceptibility to diseases and mortality [1]. Age-related disorders, such as diabetes, sarcopenia and neurodegenerative conditions, pose a significant and growing public health challenge [2, 3]. Exercise is recognised as a nonpharmacological intervention to mitigate the adverse effects of ageing. Physical exercise is essential for preserving and enhancing skeletal muscle function, ameliorating age-related muscle atrophy and providing systemic benefits against various age-related diseases [4, 5]. During exercise, skeletal muscle mitochondria undergo significant structural remodelling to meet energy and nutritional requirements, impacting cellular metabolism and signalling [6, 7]. The decline in mitochondrial function is a hallmark of ageing [8] and the accumulation of dysfunctional mitochondria with age has been observed in *C. elegans* [9]. Damaged or dysfunctional components of mitochondria can be selectively removed via fission by the transfer to the lysosome for degradation or mitophagy [10]. Another critical mechanism for mitochondrial health is fusion, where individual mitochondria merge to form a singular, enlarged structure [11]. The maintenance of mitochondrial quality, encompassing processes such as mitochondrial biogenesis, fission and fusion, is critically important in governing both the structure and functionality of mitochondria. In *C. elegans*, mitochondrial fission involves DRP-1 (DRP1 in mammals), while fusion requires FZO-1 (MFN1/2 in mammals) for outer membrane fusion and EAT-3 (OPA1 in mammals) for inner membrane fusion [12]. Overexpression of DRP-1 leads to mitochondrial hyper-fragmentation, whereas suppression of DRP-1 results in fragmented mitochondrial matrices but fused outer mitochondrial membranes [13]. Mitochondrial fusion mutant strains (*eat-3* and *fzo-1*) also display increased mitochondrial fragmentation [14].

The Forkhead Box O (FOXO) family of transcription factors play crucial roles in regulating diverse cellular processes such as oxidative stress response, mitochondrial function and cellular metabolism [15]. In *C. elegans*, the FOXO ortholog, DAF-16, exhibits heightened nuclear localisation in response to elevated reactive oxygen species (ROS) and following a swimming exercise [16, 17]. Moreover, DAF-16 has been reported to regulate mitochondrial morphology [18]. The mitochondrial proteases SPG-7 and PPGN-1 target the mitochondrial fusion protein EAT-3, DAF-16 nuclear localisation results in transcriptional repression of *spg-7* and *ppgn-1*, allowing the negative regulation of EAT-3 to be alleviated and thereby promoting mitochondrial fusion [18].

Mitochondrial morphology is also thought to determine fuel substrate use, whereby an increase in mitochondrial fragmentation increases fatty acid oxidation [19]. It was demonstrated that fragmented mitochondria promote Carnitine O-palmitoyltransferase 1 (CPT1) regulated long chain fatty acid oxidation [19]. Our previous proteomic analysis following a swim exercise in wild type (N2) *C. elegans* demonstrated an increased abundance of proteins involved in fatty acids oxidation and decreased abundance of fatty acid anabolism and the most upregulated protein was CPT-1 [20]. These results indicated that under conditions of bioenergetic stress such as during exercise, there was a promotion

of mitochondrial fatty acid oxidation and increased mitochondrial fragmentation.

Mitochondria and endoplasmic reticulum (ER) are key regulatory hubs in maintaining cellular homeostasis and have a synergistic relationship that determines their function and adaptability to the cellular environment. Communication between these organelles is mediated by Mitochondrial ER contact sites (MERCS), allowing the exchange of metabolites, lipids and calcium [21, 22]. MERCS are dynamic and relatively stable structures between mitochondria and ER (< 50 nm) that remodel in response to cellular signalling events, that can affect the function of both organelles. Recent evidence would suggest that during acute ER stress, there is activation of the adaptive Unfolded Protein Response (UPR), resulting in ER and MERCS remodelling [23]. MERCS have been identified as regulating the sites of mitochondrial fission and there is some evidence to suggest they also play a role in mitochondrial fusion [22, 24]. During fission, the ER constricts mitochondria at the site of fission using Drp1 [25]. Furthermore, the sites of mitochondrial fission have distinct ROS signatures that may regulate formation of MERCS, fission at the periphery or tip results in mitochondrial fragments destined for degradation while midzone fission is preferential for mitochondrial dynamics [26]. Disruption of mitochondrial dynamics has been reported in a wide variety of age-related diseases including neurodegeneration and sarcopenia, associated with an accumulation of dysfunctional mitochondria [27]. An increase in MERCS formation has been proposed to be involved in cell senescence and in models of neurodegenerative disease [28, 29]. In skeletal muscle, sarcomeres are surrounded by mitochondria and sarco/endoplasmic reticulum, essential for Ca^{2+} regulation. However, decreased MERCS formation has been reported with age [30]. Energy stress and subsequent AMPK activation has been demonstrated in cell models to promote autophagy and MERCS formation [31]. Introducing an exercise protocol that promotes a mild ER stress response and induces mitochondrial remodelling, will ultimately result in an improved bioenergetic profile. During exercise, there is a site-specific elevation of endogenous ROS in skeletal muscle [32, 33]. However, it has been reported in a number of different studies that there is a chronic basal elevation of ROS in muscle with age [34, 35]. The acute elevation in ROS following a physiological stress such as exercise activates specific signalling pathways including NRF2 and NF-κB [36, 37], promoting a beneficial adaptive response. However, the majority of prior investigations mainly focused on static redox states, with limited exploration of dynamic responses in both young and elderly individuals to redox stress. The Peroxiredoxins (PRDXs) family, constitute up to 1% of cellular protein content and are generally considered as a group of antioxidant enzymes with peroxidase activity [38]. Peroxiredoxin 2 (PRDX-2) functions as a peroxidase to mitigate ROS during stress and has been demonstrated to be required for the beneficial adaptation to exercise in *C. elegans* [20]. Understanding the specific role of PRDX-2 in regulating mitochondrial dynamics during exercise and ageing is essential.

In this study, the nematode *C. elegans* was utilised as a model system to investigate the intricate interplay between ageing and exercise, with a specific focus on elucidating the role of PRDX-2 in mitochondrial dynamics. We demonstrated that exercise promoted mitochondrial dynamics, MERCS formation and fatigue in *C. elegans*. A 24-hour recovery period was adequate to increase mitochondrial fusion in young worms. Old worms exhibited

fragmented mitochondria, no or delayed recovery response, an accumulation of basal ROS, an increase in hyperoxidised Peroxiredoxins and decreased fitness. The *prdx-2* mutant strain had increased mitochondrial fragmentation but did not activate mitophagy and displayed reduced physical fitness. Additionally, the acute swim exercise induced nuclear localisation of DAF-16 in the adult N2 strain promoting mitochondrial fusion. However, this response was not observed in the ageing N2 strain or the *prdx-2* mutant strain, which did not stimulate mitochondrial fusion as a result of a failure to stimulate DAF-16 dependent signalling. Our data results demonstrate blunted mitochondrial remodelling and an accumulation of ROS with age associated with disruption in the redox state of PRDX-2, highlighting the crucial role of PRDX-2 in coordinating mitochondrial adaptation in response to exercise and ageing.

2 Results

2.1 Redox mediated adaptation to exercise diminishes with age through a PRDX-2 dependent pathway

To explore the adaptive response to ageing and exercise, we employed *C. elegans*, a model organism that has been validated as an effective platform for studying physiological responses to exercise [20, 39, 40]. A 90-minute swimming exercise has been demonstrated to elicit muscle fat consumption, increase locomotory fatigue and elevate mitochondrial ROS levels [41]. Therefore, we conducted a single 90-minute swim exercise session at distinct developmental stages, adult (D4), late middle age (D8) and old (D12) worms (Fig. 1a) in N2 wild type strain. Initially, worms were subjected to diverse forms of oxidative stress, including exposure to paraquat (PQ) and arsenite (AS) following an exercise regimen and subsequent recovery. A discernible reduction in the adaptive response capacity was observed with progressive ageing. The resistance to both PQ and AS decreased immediately after acute exercise (Fig. 1b-c). Subsequently, a recovery was observed 24 hours post-exercise in the younger D4 and D8 worms, contrasting with the absence of a recovery in D12 worms (Suppl. Fig. 1a-h). The mean lifespan of each condition is detailed in Suppl. Table 1. These results demonstrate a decreased capacity to withstand stress with age and older worms are not able to recover following a physiological exercise stress.

Given the previously demonstrated involvement of PRDX-2 in redox-mediated adaptation to chronic exercise in our prior investigations [20], we conducted qPCR and Western blot analyses following a regimen of acute exercise and subsequent recovery. Initially, we examined the mRNA expression levels of the Peroxiredoxin genes in *C. elegans*, encompassing *prdx-2*, *prdx-3* and *prdx-6*, in response to the exercise and recovery protocol. Intriguingly, our observations unveiled a distinctive sensitivity pattern, with only *prdx-2* exhibiting an adaptive response to both acute exercise and the ageing processes. Specifically, following acute exercise and post-exercise in D4 worms, *prdx-2* but not *prdx-3* or *prdx-6* demonstrated an elevated expression level (Fig. 1d-f). There was no significant change of *prdx-2* expression in D8 and D12 worms in response to exercise. Furthermore, a gradual increase in the expression of *prdx-2* was observed in control conditions at D8 and D12 during the ageing process (Fig. 1d). The ratio of PRDX-2 Dimer:Monomer was analysed using non-reducing gel electrophoresis. Exercise resulted in a pronounced shift in the dimerisation of PRDX-2 in D4 worms but returned to control levels following a recovery period (Fig. 1g). D8 and D12 worms had increased expression of PRDX-2 but decreased % PRDX-2 dimer formation compared to D4 worms (Fig. 1g,h). There was also an increase in hyperoxidised Peroxiredoxins in D12 worms compared to D4 and D8 worms, which increased further following the recovery period (Fig. 1i). These findings demonstrate that resistance to oxidative stress declines with ageing and exercise was associated with distinct changes in the sensitivity of the redox state of PRDX-2 to exercise and age.

Figure 1 Decreased survival with exposure to oxidative stress, alterations in Peroxiredoxin mRNA and PRDX-2 in response to ageing and exercise. Schematic of acute swimming protocol for *C. elegans* at different ages (a). Decreased survival to paraquat and arsenite following acute exercise as well as ageing (b-c). Effects of ageing and exercise on mRNA levels of *prdx-2* (d), *prdx-3* (e), *prdx-6* (f), percentage dimer (D)/total (D+M) ratio of PRDX-2 (g), protein levels of PRDX-2 (h) and PRDX-SO$_2$/SO$_3$ (i). Graphs are the normalised relative means ± SEM and all experiments were performed with n = 60 for oxidative stress assays, n = 4 for expression of mRNA/protein and *p*-value of < 0.05 was considered as statistically significant *(*p* < 0.05). *p* values (b: D4: C vs E = 0.0034, E vs R = 0.001; D8: C vs E = 0.0435; D12: C vs E = 0.022, C vs R = 0.0076; D4 vs D8: = 0.0266; D4 vs D12 < 0.0001; D8 vs D12 = 0.0181. c: D4: C vs E = 0.0114, E vs R = 0.0009; D8: C vs E = 0.0143, E vs R = 0.0079; D12: C vs E = 0.031; C vs R = 0.0048; D4 vs D8: = 0.0136; D4 vs D12 < 0.0001; D8 vs D12 = 0.0023. d: D4: C vs E = 0.017, C vs R = 0.0485; D4 vs D8 = 0.0445; D4 vs D12 = 0.0006. g: D4: C vs E = 0.0325; E vs R = 0.0432; D12: C vs R = 0.0173, E vs R = 0.0402; D4 vs D8 = 0.0237; D4 vs D12 = 0.0009. h: D4 vs D8 = 0.0063; D4 vs D12 = 0.0271. i: D12: E vs R = 0.0475; D4 vs D12 = 0.002;

D8 vs D12 = 0.0044)

2.2 PRDX-2 is required for mitochondrial adaptation in response to exercise

In order to determine the role of PRDX-2 in mitochondrial adaptation in response to exercise, we performed MitoTracker, MitoSOX and DCFDA staining using N2 and *prdx-2* (gk169) mutant strains throughout a structured exercise and recovery regimen at distinct developmental stages. A progressive reduction in MitoTracker Red staining intensity, indicative of mitochondrial membrane potential, was noted at D8 and D12 worms relative to D4 worms control groups in both N2 and *prdx-2* mutant strains (Fig. 2a). Subsequent analysis revealed an immediate increase in staining intensity and mitochondrial membrane potential post-acute swim, with recovery to control levels observed 24 hours after exercise in D4 and D8 worms in the N2 strain. This trend was absent in old D12 wild type N2 strain. MitoTracker red staining decreased in the *prdx-2* mutant strain and unlike the N2 strain, there was no increase in staining at any age following the exercise protocol. Notably, a gradual decline in the recovery rate (Δ24h-0h) was observed with age in both N2 and *prdx-2* mutant strains, with N2 consistently exhibiting a superior recovery rate compared to the *prdx-2* mutant strain across all developmental stages (Fig. 2a). To assess the adaptive response to exercise-induced ROS, we utilised DCFDA staining as a proxy for overall cellular ROS and MitoSOX staining as an indicator for mitochondrial ROS. We observed an elevation in DCFDA and MitoSOX intensity at D4 and D8 post-exercise in both N2 and *prdx-2* mutant strains (Fig. 2b, Suppl. Fig. 2a). However, this increase was transient in the N2 strain and returned to basal levels following the recovery period but remained elevated in the *prdx-2* mutant strain at D4 and D8. There was a progressive escalation of DCFDA and MitoSOX intensity with advancing age compared to D4 worms in both strains. Remarkably, in old worms at D12, an immediate increase of DCFDA and MitoSOX intensity was not observed directly after exercise, instead a significant increase was noted 24 hours later (Fig. 2b, Suppl. Fig. 2a). The *prdx-2* mutant strain was crossed with a reporter for SKN-1 activation, CL2166 *dvIs19[(pAF15)gst-4p::GFP::NLS] III* to assess SKN-1 (ortholog of NRF2) activation during both exercise and ageing process. The staining analysis revealed a gradual elevation of SKN-1 activity at D8 and D12 in both N2 and *prdx-2* mutant strains (Suppl. Fig. 2b). Furthermore, there was an increase in SKN-1 activity 24 hours post-acute exercise at D4, although this effect was not observed at D8 or D12 in the N2 strain. Intriguingly, a decrease in SKN-1 activity was noted during the recovery phase in *prdx-2* mutants at D8 and D12. Consistent with MitoTracker staining results, the SKN-1 recovery rate exhibited a decline with ageing in both N2 and *prdx-2* mutant strains, the N2 strain demonstrated a superior recovery rate compared to the *prdx-2* mutant throughout the lifespan (Suppl. Fig. 2b). Collectively, the data indicate an acute increase in ROS and mitochondrial membrane potential following exercise in the N2 strain that returns to baseline levels following a recovery in D4 and D8 worms. There was a gradual increase in overall ROS with age and a decline in mitochondrial membrane potential. The *prdx-2* mutant strain has reduced mitochondrial membrane potential and exercise induced an increase in ROS at D4 and D8, which did not return to baseline following a recovery period. The results highlight the essential role of PRDX-2 in mediating the adaptive response to exercise.

Figure 2 Ageing and loss of PRDX-2 result in decreased membrane potential and increased ROS resulting in disrupted adaptation to exercise. MitoTracker red staining as an indicator of mitochondrial membrane potential (a) and DCFDA staining for intracellular ROS (b) following acute exercise at different stages, scale bar = 275 μm. Graphs are the normalised relative means ± SEM and all experiments were performed with n = 30-40 and *p*-value of < 0.05 was considered as statistically significant *(*p* < 0.05). *p* values (a: in N2 worms: D4: C vs E < 0.0001, C vs R < 0.0001, E vs R = 0.0002; D8: C vs E < 0.0001, E vs R < 0.0001; D12: C vs R < 0.0001, E vs R < 0.0001; D4 vs D12 < 0.0001; D8 vs D12 < 0.0001; in *prdx-2* worms: D4: C vs R = 0.0183; D8: C vs R = 0.0079; D12: C vs E = 0.0361, C vs R < 0.0001, T vs R = 0.0086; D4 vs D12 = 0.0001; D8 vs D12 = 0.005; recovery rate: N2: D4 vs D8 = 0.0035; D4 vs D12 = 0.0007; *prdx-2*: D4 vs D12 = 0.002; D8 vs D12 = 0.0022; N2 vs *prdx-2*: D4 < 0.0001; D8 = 0.0067; D12 = 0.0295. b: in N2 worms: D4: C vs R < 0.0001, E vs R < 0.0001; D8: C vs E < 0.0001, C vs R < 0.0001; D12: C vs R < 0.0001; E vs R < 0.0001; D4 vs D8 < 0.0001; D4 vs D12 < 0.0001; D8 vs D12 < 0.0001. in *prdx-2* worms: D4: C vs E = 0.005; D8: C vs E < 0.0001, C vs R < 0.0001; D12: C vs E = 0.001, C vs R = 0.0035; D4 vs D8 < 0.0001; D4 vs D12 < 0.0001; D8 vs D12 < 0.0001. recovery rate: N2: D4 vs D12 < 0.0001; D8 vs D12 < 0.0001; *prdx-2*: D4 vs D12 = 0.0297; N2 vs *prdx-2*: D12 = 0.0002)

2.3 Loss of PRDX-2 results in fragmented mitochondria and disrupted mitochondrial dynamics

To elucidate the impact of PRDX-2 on mitochondria in response to exercise, the *prdx-2* mutant strain was crossed with the muscle mitochondrial reporter strain SJ4103 *zcIs14 [myo-3::GFP(mit)]* for monitoring mitochondrial morphology and IR2539 *unc-119(ed3); Ex[pmyo-3TOMM-20::Rosella;unc-119(+)]* to determine mitophagy. Mitochondrial morphology was quantified within the body-wall muscle of *C. elegans* (Fig.3a), a tissue displaying numerous parallels to mammalian skeletal muscle [42]. This analysis entailed the classification of mitochondrial morphology into five distinct categories, each indicative of a stepwise escalation in the levels of mitochondrial fragmentation and disorganisation (Fig. 3b). The majority of body wall muscles exhibited plentiful and interconnected mitochondria in D4 N2 worms. At D8 and D12, there was an increase in mitochondrial fragmentation and progressive impairment in mitochondrial connectivity (Fig. 3a). Furthermore, acute exercise induced mitochondrial fragmentation in body wall muscles, with subsequent restoration of filamentous mitochondria observed 24 hours post-exercise in D4 and D8 worms, but not in D12 worms (Fig. 3b). The *prdx-2* mutant strain exhibits an accelerated ageing phenotype [20, 43] and a highly fragmented mitochondrial network was apparent at D4 that further deteriorates with age, suggesting a critical role for PRDX-2 in preserving mitochondrial integrity during both acute exercise and ageing process. The evaluation of mitochondrial turnover involved the use of a mitophagy reporter strain *[myo-3p tomm-20::Rosella]*. This Rosella biosensor consists of a pH-stable RFP fused with a pH-sensitive GFP. Under basal conditions, the mitochondrial network displays red and green fluorescence. In the context of mitophagy, as mitochondria undergo transportation to the acidic lysosomal environment, the GFP fluorescence undergoes quenching, while the RFP fluorescence remains stable [44]. The GFP/RFP ratio was quantified to assess mitophagy [20]. A heightened ratio, indicative of a decreased mitophagy level, was discerned in D8 and D12 worms compared to D4 worms. Interestingly, the 90-minute swim regimen induced mitophagy and recovery was evident 24 hours later in D4 and D8 worms, whereas D12 worms did not exhibit a change in mitophagy (Fig. 3c). The obvious alterations of GFP/RFP during ageing and exercise were not observed in *prdx-2* mutant strain at any age (Fig. 3c). These findings demonstrate that ageing results in an increase in mitochondrial fragmentation and decreased mitophagy. Exercise promotes mitochondrial fragmentation and mitophagy but the recovery was inhibited with age. The results highlight that loss of PRDX-2 results in increased mitochondrial fragmentation and potentially disrupted mitochondrial fusion along with a failure to activate mitophagy with exercise.

Figure 3 Ageing and loss of PRDX-2 results in mitochondrial fragmentation and disrupted mitophagy. Representative images of *myo-3::gfp* reporter strain for mitochondrial morphology (a,b) and *myo3-p::tomm20::Rosella* reporter strain for mitophagy (c) following acute exercise at different ages, scale bar = 50 µm. Graphs are the normalised relative means ± SEM and all experiments were performed with n = 130-150 for mitochondria reporter, n = 30-45 for mitophagy reporter and *p*-value of < 0.05 was considered as statistically significant *(p < 0.05). p values (b: in N2 worms: D4: C vs E = 0.03, E vs R = 0.0456; D8: C vs E = 0.041, E vs R = 0.0478; D4 vs D8 = 0.0004; D8 vs D12 = 0.0007; in *prdx-2* worms: D4 vs D12 = 0.0135. c: in N2 worms: C vs E < 0.0001, E vs R < 0.0001; D8: C vs E = 0.0063; E vs R < 0.0001; D4 vs D12 = 0.0194; D8 vs D12 = 0.0004)

2.4 Exercise induces DAF-16 nuclear localisation and promotes an acute increase in genes regulating mitochondrial morphology that is absent in the *prdx-2* mutant strain

To elucidate the molecular consequences of alterations in mitochondrial adaptation during

exercise and ageing process, the *prdx-2* mutant strain was crossed with DAF-16 reporter strains OH16024 *daf-16(ot971[daf-16::GFP]) I* and TJ356 *zIs356 [daf-16p::daf-16a/b::GFP + rol-6(su1006)]*. Moreover, mRNA levels of genes associated with mitochondrial morphology were quantified in both the N2 and *prdx-2* mutant strains. The distribution of DAF-16::GFP was quantified within the body-wall muscle of *C. elegans*, specified into three categories: cytosolic, intermediate and nuclear (Fig. 4a). The acute exercise induced DAF-16 nuclear localisation in D4 and D8 worms, but not in D12 worms (Fig. 4a, Suppl. Fig. 3a). The *prdx-2* mutant did not induce DAF-16 nuclear localisation following exercise at any stage. Previous studies have demonstrated that *prdx-2* mutants at L4 stage have elevated resistance to sodium arsenite, as a result of increased DAF-16 nuclear localisation [45]. We confirmed these results at the L4 stage but at D4 there was decreased resistance to sodium arsenite and DAF-16 nuclear localisation (Suppl. Fig, 3b,c). Inhibition of *spg-7* and *ppgn-1* expression by DAF-16 nuclear localisation, alleviates the negative regulation of mitochondrial fusion protein EAT-3 by mitochondrial proteases SPG-7 and PPGN-1 [18]. There was decreased expression of *ppgn-1* and increased expression of *eat-3* in N2 at adult D4 and adult D8 following exercise, but not in the *prdx-2* mutant (Fig. 4b-d). Furthermore, the expression levels of *dct-1* (ortholog of BNIP3 mitophagy receptor), *cpt-1* (ortholog of CPT1, a marker of long chain fatty acid oxidation and regulator of mitochondrial morphology) and *skn-1* exhibited a notable increase either immediately or 24 hours post-exercise, particularly in D4 worms (Fig. 4e-g). In contrast, D12 worms did not display a similar elevation in the expression of any of these genes following exercise. Notably, the *prdx-2* mutant strain did not demonstrate an observable upregulation of these genes (*dct-1*, *cpt-1* and *skn-1*) following exercise at any age (Fig. 4e-g). Moreover, it was observed that the expression levels of *eat-3* and *skn-1* exhibit a gradual increase with advancing age (Suppl. Fig. 3d,i). The results demonstrate that exercise induces DAF-16 nuclear localisation and expression of genes promoting mitochondrial remodelling, these changes are absent in old worms and the *prdx-2* mutant strain.

Figure 4 Ageing and loss of PRDX-2 adversely impacts the DAF-16 nuclear localisation and the expression of genes regulating mitochondrial dynamics and UPR in response to exercise. Representative images of the OH16024 strain DAF-16::GFP distribution (a) and mRNA level of *eat-3* (b), *ppgn-1* (c), *spg-1* (d), *dct-1* (e), *cpt-1* (f), *skn-1* (g), *tom-20* (h), *hsp-4* (i) and *hsp-6* (j) following acute exercise at different stages. Graphs are the normalised relative means ± SEM and all experiments were performed with n = 130-150 for DAF-16 reporter, n = 4 for mRNA expression and p-value of < 0.05 was considered as statistically significant *($p < 0.05$). p values (a: in N2 worms: D4: C vs E $<$ 0.0001, E vs R = 0.046, C vs R = 0.0182; D8: C vs E < 0.0001, E vs R = 0.0238. b: in N2 worms: D4: C vs R = 0.0157, D8: C vs E = 0.0059, C vs R = 0.0066; in *prdx-2* worms: D8: C vs R = 0.0409; D12: C vs E = 0.0019, C vs R = 0.0029. c: in N2 worms: D4: C vs E = 0.0004, C vs R < 0.0001; D8: C vs E = 0.0034, C vs R = 0.005. e: in N2 worms: D4: C vs E = 0.0073, C vs R = 0.0305. f: in N2 worms: D4: C vs E = 0.0063. g: in N2 worms: D4: C vs E = 0.0062, E vs R = 0.0055. h: in N2 worms: D4: E vs R = 0.0242. i: in *prdx-2* worms: D4: C vs R = 0.0467; D8: E vs R =0.0258. j: in N2 worms: D4: C vs E = 0.0449, E vs R = 0.0337; in *prdx-2* worms: D8: E vs R = 0.0108)

2.5 PRDX-2 is required for exercise induced UPR activation and mitochondrial ER remodelling

Mitochondria and the ER are essential regulatory centres in cellular homeostasis, engaging in a synergistic relationship. Communication between these organelles is facilitated by

MERCS, which have been identified as key regulators of both mitochondrial fission and fusion [21, 22, 24]. The activation of the adaptive Unfolded Protein Response (UPR) is linked with remodelling of ER and MERCS [23]. Expression of *hsp-4* (reflection of ER-associated UPR) and *hsp-6* (indication of mitochondrial UPR) was quantified following exercise and during the ageing process. Results indicated a heightened expression of *hsp-6* following exercise at D4 in the N2 strain (Fig. 4j). Furthermore, the expression levels of *hsp-4* increased at D8 and D12 compared to D4 in the N2 strain (Suppl. Fig. 3k). Notably, the *prdx-2* mutant strain did not demonstrate an upregulation of *tom-20* and *hsp-6* following exercise at any age (Fig. 4h, j). The link between activation of the UPR following ER stress has been suggested to regulate MERCS formation [46]. Transmission electron microscopy (TEM) imaging of mitochondria and ER was employed to quantify mitochondrial morphology and MERCS formation in response to the cellular adaptations to exercise [47]. In N2 worms, the acute exercise induces a reduction in mitochondrial length and aspect ratio (length/width), implying a transition in muscle mitochondrial morphology from rod-shaped to oval-shaped, but this alteration was not observed in *prdx-2* mutant strain (Fig. 5a). Following 5 days of exercise, there was a notable increase observed in mitochondrial length, area and aspect ratio in the N2 strain (Fig.5b). These findings suggest an augmentation in mitochondrial mass as a consequence of the chronic exercise regimen. In the comparison between adult D4 and D8 worms, it was observed that *prdx-2* mutants exhibited a more fragmented mitochondria relative to N2 worms, consistent with the highly fragmented mitochondrial network in the *prdx-2* mutant strain (Fig. 3a). Notably, both acute and chronic exercise interventions resulted in a significant reduction in the MERC distance, representing a closer coupling between mitochondria and ER in the N2 strain. Furthermore, a notable increase in the frequency of Endoplasmic Reticulum-Mitochondria Contact Compartments (ERMICCs: calculated as MERC length divided by the product of mitochondrial perimeter and MERC distance) [48] was observed following both acute and chronic exercise interventions in the N2 strain but not in the *prdx-2* mutant strain (Fig. 5a-b). These findings substantiate the presence of a substantial and dynamic interplay between mitochondria and the ER in response to exercise. Together, these findings suggest that PRDX-2 plays a key role in regulating UPR activation and MERCS formation following exercise.

Figure 5 Exercise influences mitochondrial ER contact sites in N2 but not in the *prdx-2* mutant. Mitochondria ER contact sites change following acute exercise (a) and chronic exercise (b). Graphs are the normalised relative means ± SEM and all experiments were performed with n = 30-45 and p-value of < 0.05 was considered as statistically significant *(p < 0.05). *p* values (a: mitochondrial length: N2: C vs E = 0.0231, N2 vs *prdx-2* = 0.0184. aspect ratio: N2: C vs E = 0.0317. mitochondrial area: N2 vs *prdx-2* = 0.0126. MERC distance: N2: C vs E = 0.0046. ERMICC: N2: C vs E < 0.0001. b: mitochondrial length: N2: C vs E = 0.0142; N2 vs *prdx-2* = 0.0162. aspect ratio: N2: C vs E < 0.0001. mitochondrial area: N2:C vs E = 0.0203; N2 vs *prdx-2* = 0.0225. MERC length: N2: C vs E < 0.0001; N2 vs *prdx-2* < 0.0001. MERC distance: N2:C vs E = 0.001. ERMICC: N2:C vs E < 0.0001)

2.6 PRDX-2 is required for increased fitness in response to exercise

To investigate the physiological functional implications of exercise and ageing in both N2 wild-type strain and *prdx-2* mutant strain that has an altered mitochondrial response capacity, CeleST [49] was implemented to quantify the activity patterns in these strains throughout the cycles of exercise and recovery at various stages of ageing. Parameters quantified included activity index, wave initiation rate, travel speed and brush stroke that are indicative of beneficial physical fitness, whereas body wave number, asymmetry,

stretch and curling represent frailty [40]. In both strains, a consistent decline was observed in the activity index, travel speed, wave initiation rate and brush stroke with advancing age (Fig. 6a,b, Suppl. Fig. 4a,b). Simultaneously, there was an increase in body wave number, stretch, asymmetry and curling, exhibiting more pronounced effects in older worms, particularly at the D12 stage (Fig. 6c,d, Suppl. Fig. 4c,d). Moreover, acute exercise induced alterations in these parameters, reverting to baseline in D4 and D8 worms following exercise, but not in old D12 N2 worms. Importantly the *prdx-2* mutant strain did not recover following the exercise intervention at any age. In the comparative analysis of recovery rates in N2 at different ages, D4 worms exhibited superior recovery rates in travel speed, body wave number, stretch and wave initiation rate compared to D12 worms (Fig. 6f-h, Suppl. Fig. 4e). Additionally, N2 demonstrated a more robust recovery rate in activity index, body wave number, stretch and wave initiation rate than *prdx-2* mutant, particularly notable at the D4 stage (Fig. 6e,g,h, Suppl. Fig. 4e). Furthermore, comprehensive quantification of these parameters over all stages revealed that the *prdx-2* mutant strain displayed impaired locomotory activity (Suppl. Fig. 4i). The data showed that physical fitness decreased with ageing and PRDX-2 is required for the enhanced locomotory activity or fitness in response to exercise.

Figure 6 Ageing results in a decline in physical fitness and *prdx-2* mutants cannot recover following an exercise intervention. Activity index (a), travel speed (b), body wave number (c) and stretch (d) following acute exercise at different stages; the recovery

rate of these parameters compared to normal condition (e-h). Graphs are the normalised relative means ± SEM and all experiments were performed with n = 30-40 and p-value of < 0.05 was considered as statistically significant *($p < 0.05$). p values (a: in N2 worms: D4: C vs E = 0.0032, E vs R = 0.0017; D8: C vs E = 0.0027, E vs R = 0.0088; D12: C vs E = 0.0405; D4 vs D12 = 0.0073; in *prdx-2* worms: D4: C vs E = 0.0262; D8: C vs E = 0.0231; D12: C vs E = 0.0291, C vs R = 0.0362; D4 vs D12 = 0.0289. b: in N2 worms: D4: C vs E = 0.006, E vs R = 0.001; D8: C vs E = 0.0098, E vs R = 0.0148; D12: C vs E = 0.008, C vs R = 0.0346. in *prdx-2* worms: D8: C vs E = 0.0441; D12: C vs E = 0.0263; D4 vs D12 = 0.0008. c: in N2 worms: D4: E vs R = 0.0365; D12: C vs E = 0.0084; C vs R = 0.0097; D4 vs D12 < 0.0001; D8 vs D12 = 0.0058; in *prdx-2* worms: D4: C vs E = 0.0394; D8: C vs E = 0.0002, C vs R = 0.0022; D12: C vs E = 0.0091, C vs R = 0.01; D4 vs D12 = 0.0003. d: in N2 worms: D4: C vs E = 0.0067, E vs R = 0.0177; D8: C vs E = 0.0064, E vs R = 0.0069; D12: C vs E = 0.0016, C vs R = 0.0267; D4 vs D8 = 0.036; D4 vs D12 < 0.0001; in *prdx-2* worms: D4: C vs E = 0.0098; D8: C vs E = 0.0467, C vs R = 0.0479; D12: C vs E = 0.0279; D4 vs D12 < 0.0001; D8 vs D12 = 0.0153; e: N2 vs *prdx-2*: D4 = 0.0165. f: in N2 worms: D4 vs D12 = 0.0217. g: in N2 worms: D4 vs D12 = 0.0003; D8 vs D12 = 0.0018; N2 vs *prdx-2*: D4 = 0.009; D8 = 0.0001. h: in N2 worms: D4 vs D12 = 0.0151; D8 vs D12 = 0.0095; N2 vs *prdx-2*: D4 = 0.0268; D8 = 0.00221)

In summary, the data presented demonstrate the adaptive rapid mitochondrial remodelling observed following acute exercise and ageing. PRDX-2 is sensitive to changes in the redox environment following exercise and required for exercise induced mitochondrial adaptations. Employing a cycle of acute exercise and recovery period at different stages in *C. elegans*, we observed that acute exercise initially induced mitochondrial fragmentation and mitophagy followed by mitochondrial fusion in young animals, but these responses decreased with age ultimately influencing physical fitness. Notably, the cycle of exercise and recovery failed to elicit these changes in the *prdx-2* mutant strain, evidenced by the failure to restore the redox environment, induce DAF-16 nuclear localisation and mitochondrial fusion, resulting in decreased physiological activity following exercise. Mechanistically, our results demonstrate that PRDX-2 is required for the nuclear localisation of DAF-16 following exercise and subsequent changes in mitochondrial morphology. Collectively, our data identify the indispensable role of PRDX-2 in orchestrating mitochondrial dynamics in response to a physiological stress by regulating DAF-16.

3 Discussion

In this study, we employed the nematode *C. elegans* as a model to explore the mechanistic role of PRDX-2 in regulating mitochondrial dynamics during exercise and ageing. Our data demonstrate that exercise induced alterations in the redox environment, resulting in increased DAF-16 nuclear localisation and MERCS formation dependent on PRDX-2, that regulates mitochondrial remodelling and physiological activity. These signalling mechanisms are disrupted in ageing and with the loss of PRDX-2, where there is a failure to restore the redox environment and results in reduced mitochondrial fusion and increased mitochondrial fragmentation due to an inability to stimulate DAF-16 nuclear localisation.

Mitochondrial quality is intricately regulated by mitochondrial biogenesis and selective degradation, which collectively contribute to the maintenance of mitochondrial mass, morphology and size [7, 50]. The dysregulation of mitochondrial biogenesis and impaired mitophagy are both recognised as contributing factors to compromised mitochondrial function in skeletal muscle during the ageing process [50-52]. Our previous studies have explored the effects of chronic swimming exercise on enhancing mitochondrial respiration, promoting mitochondrial biogenesis and regulating mitochondrial dynamics [20]. Recent research indicates that a cycle of mitochondrial fragmentation is induced during a single exercise session, followed by fusion after the recovery period [42]. This observation is corroborated by our current findings, which demonstrate that acute swimming triggers mitophagy, increases mitochondrial membrane potential and rapid changes in mitochondrial morphology. Furthermore, mitochondrial adaptations in response to exercise return to normal levels following a one-day recovery period. However, this cycle of mitochondrial remodelling was not observed in ageing worms as a result of an altered redox state of PRDX-2 and chronically elevated ROS stress resulting in disrupted DAF-16 nuclear localisation and downstream signalling.

A single swim session induces locomotory fatigue, elevated mitochondrial ROS and increased mitochondrial metabolic rate [41]. Our data demonstrates that exercise-induced ROS are associated with specific changes in the redox state of PRDX-2, increased DAF-16 nuclear localisation and mitochondrial remodelling that increases physical fitness and sensitivity to oxidative stress in young worms. However, these alterations were delayed or not observed in old worms. Recent studies have proposed a theory termed "redox-stress response capacity (RRC)" to elucidate this phenomenon, suggesting that cellular signalling and homeostasis maintenance are regulated through ROS-mediated adaptive responses or a hormesis effect [53]. Furthermore, the decline in RRC over time was termed as redox-stress response resistance (RRR), which offers an explanation for the reduction in RRC observed during ageing [54]. Interestingly, the mechanisms of RRC and RRR are mediated by PRDX-2 and hyperoxidation of Peroxiredoxins [54]. Decreased H_2O_2 sensitivity of PRDX-2 and elevated PRDX-2 hyperoxidation have been observed in ageing human fibroblasts and *C. elegans* [54]. These findings align with our own observations, as evidenced by reversible changes in the redox state of PRDX-2 following exercise only in young worms and a corresponding increase in hyperoxidised Peroxiredoxins in old worms. Furthermore,

our previous redox proteomic analysis indicated that PRDX-2 regulates the induction of a redox relay for cellular adaptive response to exercise [20].

Peroxiredoxin 2, outcompetes other proteins for activation with physiological concentrations of H_2O_2 due to its abundance and kinetic reactivity [55]. However, Peroxiredoxin 2 has also been explored as involved in a redox relay for the transfer of oxidative equivalents to target proteins via reversible modification of Cysteine residues on redox-sensitive proteins [20, 54, 56]. STAT3, NRF2, and FOXO (DAF-16) are key transcription factors involved in the regulation of mitochondrial function and muscle myogenesis [17, 20, 57]. Peroxiredoxin 2 has been identified to interact with STAT3, forming a redox relay in response to a short bolus of H_2O_2 treatment [58]. Peroxiredoxin 1 has also been implicated in binding and regulating the activity of FOXO3 [59]. Likewise, the nuclear localisation of DAF-16 can be induced by ROS, with evidence suggesting that this process is regulated by its redox-sensitive cysteine residues forming a mixed disulphide with a redox signalling cascade [16]. However, the precise relationship between PRDX-2 and DAF-16 remains ambiguous in the current understanding of cellular signalling pathways.

DAF-16 is the only ortholog of the FOXO transcription factors in *C. elegans* and vital for preserving cellular homeostasis, it is required for the lifespan extension reported in *daf-2* mutant strains [15]. FOXO1 and FOXO3 govern glucose metabolism, lipid homeostasis and mitochondrial function primarily through their nuclear accumulation [60]. Elevated levels of ROS, originating from three mitochondrial mutants (*clk-1*, *isp-1* and *nuo-6*), prompt the nuclear translocation of DAF-16, thereby enhancing longevity [17]. The oxidation of Cysteine residues within FOXO/DAF-16 is necessary for its nuclear localisation and subsequent activation in response to ROS [16]. A previous study demonstrated that DAF-16 exhibits increased nuclear localisation in response to elevated ROS levels as well as following a swimming protocol [17]. Moreover, it is noteworthy that the heightened nuclear localisation of DAF-16 promotes mitochondrial fusion by promoting EAT-3 expression through the inhibition of mitochondrial proteases *spg-7* and *ppgn-1* [18]. Our findings corroborate that acute exercise promotes DAF-16 nuclear accumulation, leading to an increase in mitochondrial dynamics. Furthermore, we observed increased levels of the mitophagy receptor *dct-1*, mitochondrial fusion regulator *eat-3* and decreased *ppgn-1* following acute exercise. Surprisingly, these changes were not observed in the *prdx-2* mutant strain, suggesting that PRDX-2 is required for maintaining mitochondrial quality via regulating DAF-16 in response to exercise.

The altered redox state of PRDX-2 following exercise can potentially trigger the downstream activation of SKN-1, which has been reported via the p38 MAPK cascade [61]. It is widely recognised that changes occur in the transcriptome, proteome and metabolome during the ageing process [9]. RNA sequencing data from isolated tissues of ageing *C. elegans* demonstrated elevated expression levels of *skn-1*, *daf-16* and *prdx-2* in body wall muscle [62]. Similarly, it was demonstrated elevated levels of *daf-16* and *skn-1* in adult D7 worms compared to adult D1 worms [63]. These findings are consistent with the results

presented here, which demonstrate heightened SKN-1 reporter activity and increased PRDX-2 (including hyperoxidised) protein levels at D12 compared to D4 worms. The *prdx-2* mutant strain displays an accelerated ageing phenotype [20, 43], characterised by a highly fragmented mitochondrial network evident at D4, which is aggravated with age. These findings suggest a potential role for PRDX-2 in maintaining mitochondrial integrity during both acute exercise and the ageing process. The depletion of PRDX-2 has previously been reported to enhance the activation of DAF-16 in response to low concentrations of arsenite [45]. These experiments were performed at the L4 stage in the *prdx-2* mutant strain. The accelerated ageing phenotype characteristic of the *prdx-2* mutant strain and altered redox environment could have resulted in the increased nuclear localisation of DAF-16 observed at this early developmental age.

In summary, our results demonstrate that there is blunted mitochondrial remodelling with age which is associated with an altered redox environment. PRDX-2 is required for mitochondrial adaptations in response to exercise through the regulation of the intracellular redox environment and appropriate DAF-16 nuclear localisation. Our data demonstrate that during ageing, there are elevated levels of ROS, increased mitochondrial fragmentation, reduced survival capabilities and decreased locomotory activity. Employing a cycle of acute exercise and recovery period model at various stages in *C. elegans*, acute exercise induces mitochondrial fragmentation and subsequent mitochondrial fusion through the activation of DAF-16, ultimately influencing physical fitness. Notably in the *prdx-2* mutant strain, the cycle of exercise and recovery failed to induce the observed changes compared to the N2 strain. The *prdx-2* mutant strain had an absence of alterations in mitochondrial membrane potential, mitochondrial dynamics as well as an inability to resolve the altered redox environment and reestablish mitochondrial adaptation and physiological activity following exercise. The redox state of PRDX-2 is sensitive to the redox environment and decreased reversible modification of the dimer ratio of PRDX-2 during ageing correlated with decreased survival and physical fitness in *C. elegans*. There are several limitations to the work presented in this study. We acknowledge that mitochondrial mass and morphology could affect the uptake of MitoTracker Red and MitoSOX [64] used to estimate mitochondrial membrane potential and mitochondrial ROS generation both of which are altered as a result of ageing and exercise. Furthermore, a physical interaction between PRDX-2 and DAF-16 was not demonstrated and future research should encompass additional investigations for a comprehensive understanding of this relationship. Despite these limitations, this study demonstrates the pivotal role of PRDX-2 in regulating DAF-16 nuclear localisation and mitochondrial remodelling during exercise and ageing.

4. Reagents and Resources:

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
rabbit anti-Peroxiredoxin 2 (*C. elegans*)	Elizabeth Veal lab [65]	N/A
rabbit anti-Peroxiredoxin-SO$_3$	Abcam	Cat# ab16830, RRID:AB_443491
IRDye 800CW Goat anti-Rabbit IgG	LI-COR Biosciences	Cat# 926-32211, RRID:AB_621843
Bacterial and Virus Strains		
E. coli: Strain OP50	Caenorhabditis Genetics Center	N/A
Chemicals, Peptides and Recombinant Proteins		
Acetic acid	Sigma	Cat# A6283
Acrylamide	Sigma	Cat# A3699
Agar	Sigma	Cat# A1296
APS	Sigma	Cat# A3678
Arsenite	Sigma	Cat# S7400
Beta-mercaptoethanol	Sigma	Cat# M7522
Boric acid	Sigma	Cat# B0252
Bradford Reagent	Bio-Rad	Cat# 5000006
Bromophenol blue	Sigma	Cat# 114391
BSA	Sigma	Cat# A3059
CaCl$_2$	Sigma	Cat# C1016
Chloroform	Sigma	Cat# C0549
Cholesterol	Sigma	Cat# C8667
dATP	Thermo Fisher Scientific	Cat# R0141
dCTP	Thermo Fisher Scientific	Cat# R0151
dGTP	Thermo Fisher Scientific	Cat# R0161
DTT	Sigma	Cat# Y00147
dTTP	Thermo Fisher Scientific	Cat# R0171
EDTA	Sigma	Cat# ED2SS
Ethanol	Sigma	Cat# E7023
Glycerol	Sigma	Cat# G6279
Glycine	Sigma	Cat# G8898
Isopropanol	Sigma	Cat# I9516
K$_2$HPO$_4$	Sigma	Cat# 3786
KH$_2$PO$_4$	Sigma	Cat# P9791
Levamisole	Sigma	Cat# L9756
Methanol	Sigma	Cat# 34860
MgSO$_4$	Sigma	Cat# M7506

MyTaq ™ Reaction Buffer Red	Sigma	Cat# R2523
Na_2HPO_4	Sigma	Cat# S0876
NaCl	Sigma	Cat# S9888
NaClO	Sigma	Cat# 1056142500
NaOH	Sigma	Cat# S5881
NEM	Sigma	Cat# E3876
Nitrocellulose Blotting membrane	GE Healthcare Life Sciences	Cat# 1060003
Nystatin	Sigma	Cat# N3503
Paraquat	Sigma	Cat# 856177
Peptone	Sigma	Cat# 91249
Ponceau S Staining Solution	Thermo Fisher Scientific	Cat# A40000279
Protease Inhibitor Cocktail	Sigma	Cat# P8340
Proteinase K	Sigma	Cat# 03115828001
Random hexamer	Sigma	Cat# 100026484
Ribolock	Thermo Fisher Scientific	Cat# EO0381
RNase free water	Sigma	Cat# W4502
RT buffer	Sigma	Cat# Y02321
SDS	Sigma	Cat# L3771
Superscript II	Sigma	Cat# 100004925
SYBR Green	Qiagen	Cat# 339347
TEMED	Sigma	Cat# T9281
Triton^{TM} X-100	Sigma	Cat# T8787
Trizma® base	Sigma	Cat# T1503
TRIzol Reagent	Life Technologies	Cat# 15596018
UltraPure ^{TM} Agarose	Sigma	Cat# 16500
Critical Commercial Assays		
MitoTracker^{TM} Red CMXRos	Thermo Fisher Scientific	Cat# M7512
MitoSOX^{TM} Red mitochondrial superoxide indicator	Thermo Fisher Scientific	Cat# M36008
Carboxy-H_2DCFDA	Thermo Fisher Scientific	Cat# C2938
Experimental Models: Organisms/Strains		
C. elegans: wild type	Caenorhabditis Genetics Center	N2
C. elegans: prdx-2(gk169) II	Elizabeth A Veal and Caenorhabditis Genetics Center	VC289
C. elegans: dvIs19[(pAF15)gst-4p::gfp] III	Caenorhabditis Genetics Center	CL2166
C. elegans: zcIs14 [myo-3::gfp(mi t)]	Caenorhabditis Genetics Center	SJ4103
C. elegans: *I (unc-119(ed3);*	Tavernarakis Lab	IR2539

Ex[myo-3p tomm20::Rosella;unc-119(+)]		
C. elegans: *zIs356 [daf-16p::daf-16a/b::GFP + rol-6(su1006)]*	Antonio Lab	TJ356
C. elegans: *daf-16(ot971[daf-16::GFP]) I.*	Antonio Lab	OH16024
C. elegans: *prdx-2 (gk169) II; dvIs19[(pAF15)gst-4p::GFP::NLS] III*	This paper	*prdx-2* x CL2166
C. elegans: *prdx-2 (gk169) II; zcIs14 [myo-3::GFP(mit)]*	This paper	*prdx-2* x SJ4103
C. elegans: *prdx-2(gk169) II; unc-119(ed3); Ex[pmyo-3TOMM-20::Rosella;unc-119(+)]*	This paper	*prdx-2* x IR2539
C. elegans: *prdx-2(gk169) II; zIs356 [daf-16p::daf-16a/b::GFP + rol-6 (su1006)]*	This paper	*prdx-2* x TJ356
C. elegans: *prdx-2(gk169) II; daf-16(ot971[daf-16::GFP]) I.*	This paper	*prdx-2* x OH16024
Oligonucleotides		
Primer: prdx-2 OF: CGCTCCTCCTAAACGTTGTAGTC OR: CTAGGGATACGGGGGAAATTAG IR: GGATATTGAAACCCAACGGCAACG	Sigma	genotyping
Primer: *prdx-2* Forward: CATTCCAGTTCTCGCTGAC Reverse: ATGATGAAGAGTCCACGGA	Sigma	qPCR
Primer: *prdx-3* Forward: GTTCCGTTCTCTTGGAGCTG Reverse: CTTGTTGAAATCAGCGAGCA	Sigma	qPCR
Primer: *prdx-6* Forward: GGAGAACAATGCTGATGC Reverse: ATCTGAACATGGCGTTTGC	Sigma	qPCR
Primer: *skn-1* Forward: GGACAACAGAATCCCAAAGG Reverse: TCAGGACGTCAACAGCAGAC	Sigma	qPCR
Primer: CDC-42 Forward: AGCCATTCTGGCCGCTC	Sigma	qPCR

TCG Reverse: GCAACCGCTTCTCGTTT GGC		
Primer: *eat-3* Forward: AGAGCATCGAAACCGG ATGG Reverse: GCGTCAGCATAGCTTCT TCG	Sigma	qPCR
Primer: *ppgn-1* Forward: ATGCTTCTACACCGCTC CAC Reverse: GTGGAAATCTGCGAGC ACT	Sigma	qPCR
Primer: *spg-7* Forward: CCGTTGTCGTTTGAGA CACC Reverse: CGGCGAAGTGCGTTCA TTAC	Sigma	qPCR
Primer: *cpt-1* Forward: AACGACGTACCGGACG ATTC Reverse: CCGAGGTAGCCATCACC AAA	Sigma	qPCR
Primer: *dct-1* Forward: GAAAATACACCTCCAA AGACTGTCCG Reverse: GATGTTTGATTAGTTTC CGGCAAACAG	Sigma	qPCR
Primer: *hsp-4* Forward: ACGACCACAATCGTCT CAGTCC Reverse: CTTCGTCAGTGAGCTTT CCTCC	Sigma	qPCR
Primer: *hsp-6* Forward: GAACCGGAAAGGAAC AACAGATCG Reverse: GCAATCTTGGTTCGGAG AGCCTC	Sigma	qPCR
Primer: *tomm-20* Forward: CGGCTACTGCATTTACT TCGA Reverse: TCATTGCCTGCTGCAGC TGGA	Sigma	qPCR
Software and Algorithms		
Prism 7	GraphPad Software	RRID:SCR_002798
ImageJ	NIH	RRID:SCR_003070
Image Studio Lite	Image Studio Lite	RRID:SCR_013715

CeLeSt	Christophe Restif et al., 2014 [49]	N/A
OASIS 2	Online Application for Survival Analysis 2	N/A

5. Materials and methods

5.1 *C. elegans* strains

C. elegans were cultured on NGM plates seeded with E. Coli (OP50) at 20 °C. *C. elegans* stains N2 wild type, VC289 (*prdx-2(gk169) II*) strain, CL2166 (*dvIs19[(pAF15)gst-4p::gfp] III*) and SJ4103 (*zcIs14 [myo-3::gfp(mit)]*) strains were obtained from the Caenorhabditis Genetics Center (CGC); IRE2539 (*Ex[pmyo-3 tomm-20::Rosella;unc-119(+)]*) was a gift from the Tavernarakis lab University of Crete, Greece; TJ356 (*zIs356 [daf-16p::daf-16a/b::GFP + rol-6(su1006)*) and OH16024 (*daf-16(ot971[daf-16::GFP]) I*) were obtained from Instituto de Biomedicina de Sevilla. In this paper, the *prdx-2* mutant strain was crossed with the reporter strains: SKN-1 reporter CL2166, DAF-16 reporter OH16024/TJ356, mitochondrial GFP reporter SJ4103 and mitophagy reporter IR2539.

5.2 *C. elegans* swimming exercise

For the different stages of worms, adult D1 worms were washed off plates and the larvae were removed daily to obtain worms at different ages (adult D4, D8, D12). For the acute exercise and control groups, 30 worms at least were transferred to unseeded NGM plates with or without M9 buffer, then worms were allowed to swim or crawl for a duration of 90 minutes at D4, D8 and D12 separately [41]. The recovery group underwent a 24-hour recovery period following the exercise regimen. For the chronic swimming exercise, adult D1 worms underwent 90-minute swimming sessions twice daily over a period of 5 days.

5.3 Oxidative stress survival assays

For the paraquat and sodium arsenite stress assays, 10 worms were selected for each group with a total of 60 worms per condition. These worms were carefully placed in individual wells of a 96-well plate containing either 100mM Paraquat or 2.5mM Sodium Arsenite, both diluted in M9 buffer. Worm death was determined by the absence of response to a gentle tap with a picker and survival was evaluated at intervals of 2 hours [66].

5.4 Western blotting

Protein samples from *C. elegans* were quantified using the Bradford reagent subsequent to homogenisation in an alkylating lysis buffer. 20 µg of protein per each condition was applied to 12% reducing or non-reducing SDS PAGE gels. The protein transfer was accomplished via a semi-dry blotter and the membrane underwent Ponceau S staining for normalisation. Following washing with TBS-T, the membranes were subjected to blocking in 5% milk within TBS-T for 1 hour at room temperature. Following the blocking step, the membranes were exposed to primary antibodies (rabbit anti-Peroxiredoxin 2 and rabbit anti-Peroxiredoxin-SO$_3$) at a dilution of 1:1000 in 5% milk and this incubation persisted overnight. Subsequently, membranes were washed three times with TBS-T, each lasting 10 minutes. Membranes underwent incubation with the secondary antibody at a dilution of 1:10,000 in TBS-T, conducted in the absence of light, for a duration of 1 hour. Image acquisition was performed using the Odyssey Fc imaging system (Li-Cor). The subsequent

analysis of blot quantification and normalisation was performed employing Image Studio Lite.

5.5 qPCR

RNA isolation from *C. elegans* was conducted utilizing TRIzol, following the methodology described in [67]. Subsequent to RNA isolation, cDNA synthesis and real-time qPCR were executed employing established protocols. For mRNA synthesis, 500 ng of RNA was mixed with 1 µl of random hexamers and incubated at 65 °C for 10 minutes. Then, the mixture was combined with a master mix including 4 µl of RT buffer, 2 µl of DTT, 1 µl of dNTP, 1 µl of Superscript II and 1 µl of Ribolock, followed by incubation at 42 °C for 60 minutes. Subsequent qPCR analysis was conducted using the SYBR Green Master Mix in a 10 µl reaction volume. The quantification of gene expression, relative to the housekeeping gene CDC42, was determined utilising the delta Ct method.

5.6 Imaging of *C. elegans*

For the staining of *C. elegans*, a total of 30-45 worms per condition were subjected to the procedure immediately after the acute exercise or following a 24-hour recovery period. The worms were incubated in 2.5 µM MitoTracker Red CMXRos for 10 minutes, 10 µM MitoSOX Red for 1 hour, or 25 µM H2DCFDA for 1 hour. After the probe incubation, the worms were transferred to NGM plates for 2-3 hours in the dark room to prevent accumulation of the stain in the guts. Subsequently, the worms were immobilised by 20 mM levamisole and imaged using EVOS at 10× magnification [20].

For the SKN-1 reporter, utilising the CL2166 (*dvIs19 [(pAF15)gst-4p::gfp III]*) strain, a cohort of 30-45 worms at different ages underwent acute swimming. Post-exercise, the worms were immobilised on unseeded NGM plates for imaging using EVOS microscopy at a ×10 magnification. The assessment of green fluorescence intensity throughout the entire body of each worm was conducted using ImageJ [20].

For the assessment of mitochondrial dynamics in the body wall muscle, the SJ4103 strain (*zcIs14 [myo-3::gfp(mit)]*) was employed, using 30-45 worms subjected or not to acute swimming. Post-exercise, immobilisation of the worms was achieved using an agar slide covered with a coverslip. Subsequent imaging of mitochondria within the body wall muscles, specifically the region between the pharynx and vulva, was conducted under EVOS M7000 microscopy at a magnification of ×60. A total of 130-150 images were acquired and subsequently categorised into five distinct classes: Class 1, signifying highly abundant and well-networked mitochondria with minimal or no blebbing; Class 2, characterized by highly abundant mitochondria with network gaps and some blebbing; Class 3, featuring less abundant mitochondria with network gaps and increased blebbing; Class 4, indicating sparse mitochondria with some blebbing; and Class 5, denoting very sparse mitochondria [39].

To evaluate mitophagy, the IR2539 strain (*unc-119(ed3); Ex[pmyo-3 tomm-20::Rosella;unc-119(+)]*) was used. A group consisting of 30-45 worms underwent

imaging at 60× magnification using the EVOS M7000 microscope. The ratio of green to red fluorescence intensity within a representative head region of each individual worm was subsequently determined using ImageJ [20].

To determine the nuclear localisation of DAF-16, the OH16024 strain (*daf-16(ot971[daf-16::GFP]) I.*) and TJ356 strain (*zIs356 [daf-16p::daf-16a/b::GFP + rol-6(su1006)]*) were used. 30-45 worms were immobilised using an agar slide covered with a coverslip. Subsequent imaging of DAF-16 nuclear localisation within the body wall muscles, specifically the region between the pharynx and vulva, was conducted under EVOS M7000 microscopy at a magnification of ×60. A total of 130-150 images were acquired and subsequently categorized into three distinct classes: nuclear, where the DAF-16::GFP distribution was visible in the nuclear of body wall muscle; intermediate, where the DAF-16::GFP distribution was not completely visible, showing punctate fluorescence in the cytosolic; and cytosolic, where the DAF-16::GFP distribution was in the cytosolic [18].

5.7 Transmission Electron Microscopy

C. elegans strains were immobilised with osmium tetroxide. Subsequently, the specimens underwent a dehydration process through a graded series of ethanol concentrations (30%, 50%, 70%, 90% and finally 100%), with each step lasting 2×15 minutes. Following the last 100% ethanol dehydration step, acetone was used to replace ethanol in a 2×20-minute procedure. Subsequently, the samples were immersed in a 50:50 mixture of resin and acetone for 4 hours, followed by placement in a 75:25 mixture on a rotator overnight. The following day, samples were transferred into 100% resin and rotated for 5-6 hours. Upon completion of this infiltration step, the specimens were relocated into the appropriate embedding mould, filled with fresh 100% resin and then subjected to an oven at 65 °C for 48 hours for polymerisation. Finally, the moulded samples were sectioned to prepare them for imaging. 30-50 captured images were acquired at a magnification of 25,000 X. Using Image J according to [47], mitochondrial length (mitochondrial longitudinal distance), mitochondrial width (mitochondrial lateral distance), aspect ratio (ratio of mitochondrial length/ width), mitochondrial area (mitochondrial area size), MERC length (contacted length between mitochondria and ER), MERC distance (contacted distance between mitochondria and ER) and ERMICC (ratio of MERC length to product of mitochondrial perimeter and MERC distance),.

5.8 CeleST

To evaluate the swimming proficiency of *C. elegans*, CeleST analysis was performed following the acute exercise and the revery period. A minimum of 30 worms, arranged in 4-5 individuals per trial, were positioned on a glass substrate within a 10 mm ring. Subsequently, 60-second video recordings were captured at a rate of 16 frames per second, utilizing a Nikon LV-TV microscope set at 1× magnification and equipped with an OPTIKA C-P20CM camera [20, 40].

5.9 Statistical analysis

Images obtained from Western blot and microscopy of *C. elegans* were subjected to semi-automated quantification using ImageJ and Image Studio Lite and manual corrections were applied as needed to ensure accuracy in the quantification process. For the MitoTracker, MitoSOX, DCFDA and CL2166 strain staining, the quantification involved a total of 30-45 worms, with the fluorescence intensity measured per worm for each specific experimental condition. For SJ4103 strain staining, an evaluation was conducted on 130-150 images captured in the region between the pharynx and vulva and assessment was carried out based on the five predefined categories described earlier. For TJ356 and OH16024 strain, 130-150 images of body wall muscle were captured and quantified based on the DAF-16::GFP distribution categorized as nuclear, intermediate and cytosolic. For the IR2539 strain staining, the assessment involved determining the ratio of green fluorescence to red fluorescence for each individual worm. The specific details regarding the statistical analyses conducted are provided in the respective figure legend. To assess differences between two groups, a student t-test was employed. For comparisons involving more than two groups, either one-way or two-way ANOVA was utilised. The log-rank test was employed for comparing tolerance in the oxidative stress assay. The distribution into multiple classes was analysed using the chi-square test. A significance threshold of p-value < 0.05 was considered statistically significant. All graphical representations and statistical analyses were performed using GraphPad Prism 7. The source data used for figure generation in this manuscript are available in the source data file.

Acknowledgements:

We would like to sincerely thank Elizabeth Veal lab for providing the anti-PRDX-2 antibody and VC289 *prdx-2* mutant strain, the Tavernarakis lab for providing the IR2539 mitophagy reporter strain and the facilities and scientific and technical assistance of the Anatomy Imaging and Microscopy Facility at the University of Galway (https://imaging.universityofgalway.ie/imaging/) for electron microscopy work. QX (202006370047) and PL (202206370063) studentships are funded by the Chinese Scholarship Council (CSC), JCCM studentship is funded by the College of Nursing Medicine and Health Sciences, University of Galway.

Competing Interests:

The authors declare no competing interests.

Author Contributions:

Conceptualisation, BMcD and QX; methodology QX, PL, JCCM, KW, AMV, EM, PD and BMcD; investigation, QX, PLL, JCCM, EM and BMcD; resources, KW, AVM, PD and BMcD; writing-original draft QX and BMcD; writing-review and editing, QX, PL, JCCM, AMV, EM, PD, KW and BMcD; supervision, KW and BMcD.

References

Lopez-Otin, C., *et al.* (2013) The hallmarks of aging. *Cell* 153, 1194-1217

Longo, M., *et al.* (2019) Diabetes and aging: from treatment goals to pharmacologic therapy. *Frontiers in Endocrinology* 10, 45

Hou, Y., *et al.* (2019) Ageing as a risk factor for neurodegenerative disease. *Nat Rev Neurol* 15, 565-581

Hood, D.A., *et al.* (2019) Maintenance of skeletal muscle mitochondria in health, exercise, and aging. *Annual Review of Physiology* 81, 19-41

Pedersen, B.K. and Saltin, B. (2015) Exercise as medicine - evidence for prescribing exercise as therapy in 26 different chronic diseases. *Scand J Med Sci Sports* 25 Suppl 3, 1-72

Yan, Z., *et al.* (2012) Exercise training-induced regulation of mitochondrial quality. *Exercise and sport sciences reviews* 40, 159

Hood, D.A., *et al.* (2019) Maintenance of Skeletal Muscle Mitochondria in Health, Exercise, and Aging. *Annu Rev Physiol* 81, 19-41

López-Otín, C., *et al.* (2023) Hallmarks of aging: An expanding universe. *Cell*

Spanoudakis, E. and Tavernarakis, N. (2023) Age-associated anatomical and physiological alterations in Caenorhabditis elegans. *Mechanisms of Ageing and Development*, 111827

Youle, R.J. and Van Der Bliek, A.M. (2012) Mitochondrial fission, fusion, and stress. *Science* 337, 1062-1065

Adebayo, M., *et al.* (2021) Mitochondrial fusion and fission: The fine-tune balance for cellular homeostasis. *FASEB journal: official publication of the Federation of American Societies for Experimental Biology* 35, e21620

Campbell, D. and Zuryn, S. (2023) The mechanisms and roles of mitochondrial dynamics in C. elegans. In *Seminars in Cell & Developmental Biology*, Elsevier

Labrousse, A.M., *et al.* (1999) C. elegans dynamin-related protein DRP-1 controls severing of the mitochondrial outer membrane. *Molecular cell* 4, 815-826

Machiela, E., *et al.* (2020) Disruption of mitochondrial dynamics increases stress resistance through activation of multiple stress response pathways. *The FASEB Journal* 34, 8475

Rodriguez-Colman, M.J., *et al.* (2024) FOXO transcription factors as mediators of stress adaptation. *Nat Rev Mol Cell Biol* 25, 46-64

Putker, M., *et al.* (2013) Redox-dependent control of FOXO/DAF-16 by transportin-1. *Molecular cell* 49, 730-742

Senchuk, M.M., *et al.* (2018) Activation of DAF-16/FOXO by reactive oxygen species contributes to longevity in long-lived mitochondrial mutants in Caenorhabditis elegans. *PLoS Genetics* 14, e1007268

Chaudhari, S.N. and Kipreos, E.T. (2017) Increased mitochondrial fusion allows the survival of older animals in diverse C. elegans longevity pathways. *Nature communications* 8, 182

Ngo, J., *et al.* (2023) Mitochondrial morphology controls fatty acid utilization by changing CPT1 sensitivity to malonyl-CoA. *The EMBO Journal*, e111901

Xia, Q., *et al.* (2023) Peroxiredoxin 2 is required for the redox mediated adaptation to exercise. *Redox biology* 60, 102631

Giacomello, M. and Pellegrini, L. (2016) The coming of age of the mitochondria–ER contact: a matter of thickness. *Cell Death & Differentiation* 23, 1417-1427

Murley, A. and Nunnari, J. (2016) The emerging network of mitochondria-organelle contacts.

Molecular cell 61, 648-653

van Vliet, A.R., *et al.* (2017) The ER stress sensor PERK coordinates ER-plasma membrane contact site formation through interaction with filamin-A and F-actin remodeling. *Molecular cell* 65, 885-899. e886

Cieri, D., *et al.* (2018) SPLICS: a split green fluorescent protein-based contact site sensor for narrow and wide heterotypic organelle juxtaposition. *Cell Death & Differentiation* 25, 1131-1145

Quintana-Cabrera, R. and Scorrano, L. (2023) Determinants and outcomes of mitochondrial dynamics. *Molecular Cell*

Kleele, T., *et al.* (2021) Distinct fission signatures predict mitochondrial degradation or biogenesis. *Nature* 593, 435-439

Sharma, A., *et al.* (2019) Causal roles of mitochondrial dynamics in longevity and healthy aging. *EMBO reports* 20, e48395

Ziegler, D.V., *et al.* (2021) Calcium channel ITPR2 and mitochondria–ER contacts promote cellular senescence and aging. *Nature Communications* 12, 720

Morgado-Cáceres, P., *et al.* (2022) The aging of ER-mitochondria communication: A journey from undifferentiated to aged cells. *Frontiers in Cell and Developmental Biology* 10, 946678

Tubbs, E., *et al.* (2018) Disruption of mitochondria-associated endoplasmic reticulum membrane (MAM) integrity contributes to muscle insulin resistance in mice and humans. *Diabetes* 67, 636-650

Hu, Y., *et al.* (2021) The AMPK-MFN2 axis regulates MAM dynamics and autophagy induced by energy stresses. *Autophagy* 17, 1142-1156

Sakellariou, G.K., *et al.* (2013) Studies of mitochondrial and nonmitochondrial sources implicate nicotinamide adenine dinucleotide phosphate oxidase (s) in the increased skeletal muscle

superoxide generation that occurs during contractile activity. *Antioxidants & redox signaling* 18, 603-621

Murphy, M.P., *et al.* (2011) Unraveling the biological roles of reactive oxygen species. *Cell metabolism* 13, 361-366

Sakellariou, G.K., *et al.* (2017) Redox homeostasis and age-related deficits in neuromuscular integrity and function. *J Cachexia Sarcopenia Muscle*

Palomero, J., *et al.* (2013) Aging increases the oxidation of dichlorohydrofluorescein in single isolated skeletal muscle fibers at rest, but not during contractions. *Am J Physiol Regul Integr Comp Physiol* 305, R351-358

Vasilaki, A., *et al.* (2006) Adaptive responses of mouse skeletal muscle to contractile activity: the effect of age. *Mechanisms of ageing and development* 127, 830-839

Done, A.J. and Traustadóttir, T. (2016) Nrf2 mediates redox adaptations to exercise. *Redox biology* 10, 191-199

Rhee, S.G. and Kil, I.S. (2017) Multiple functions and regulation of mammalian peroxiredoxins. *Annual review of biochemistry* 86, 749-775

Laranjeiro, R., *et al.* (2019) Swim exercise in Caenorhabditis elegans extends neuromuscular and gut healthspan, enhances learning ability, and protects against neurodegeneration. *Proceedings of the National Academy of Sciences* 116, 23829-23839

Restif, C., *et al.* (2014) CeleST: computer vision software for quantitative analysis of C. elegans swim behavior reveals novel features of locomotion. *PLoS computational biology* 10, e1003702

Laranjeiro, R., *et al.* (2017) Single swim sessions in C. elegans induce key features of mammalian exercise. *BMC biology* 15, 1-17

Campos, J.C., *et al.* (2023) Exercise preserves physical fitness during aging through AMPK and mitochondrial dynamics. *Proceedings of the National Academy of Sciences* 120, e2204750120

Oláhová, M., *et al.* (2008) A redox-sensitive peroxiredoxin that is important for longevity has tissue-and stress-specific roles in stress resistance. *Proceedings of the national academy of sciences* 105, 19839-19844

Palikaras, K., *et al.* (2015) Coordination of mitophagy and mitochondrial biogenesis during ageing in C. elegans. *Nature* 521, 525-528

Olahova, M. and Veal, E.A. (2015) A peroxiredoxin, PRDX-2, is required for insulin secretion and insulin/IIS-dependent regulation of stress resistance and longevity. *Aging Cell* 14, 558-568

Fan, Y. and Simmen, T. (2019) Mechanistic connections between endoplasmic reticulum (ER) redox control and mitochondrial metabolism. *Cells* 8, 1071

Lam, J., *et al.* (2021) A universal approach to analyzing transmission electron microscopy with ImageJ. *Cells* 10, 2177

Naon, D., *et al.* (2016) Critical reappraisal confirms that Mitofusin 2 is an endoplasmic reticulum-mitochondria tether. *Proc Natl Acad Sci U S A* 113, 11249-11254

Restif, C., *et al.* (2014) CeleST: computer vision software for quantitative analysis of C. elegans swim behavior reveals novel features of locomotion. *PLoS Comput Biol* 10, e1003702

Sharma, A., *et al.* (2019) Causal roles of mitochondrial dynamics in longevity and healthy aging. *EMBO Rep* 20, e48395

Goljanek-Whysall, K., *et al.* (2020) miR-181a regulates p62/SQSTM1, parkin, and protein DJ-1 promoting mitochondrial dynamics in skeletal muscle aging. *Aging Cell* 19, e13140

Shally, A. and McDonagh, B. (2020) The redox environment and mitochondrial dysfunction in

age-related skeletal muscle atrophy. *Biogerontology* 21, 461-473

Meng, J., *et al.* (2017) The decay of Redox-stress Response Capacity is a substantive characteristic of aging: Revising the redox theory of aging. *Redox Biology* 11, 365-374

Meng, J., *et al.* (2023) Redox-stress response resistance (RRR) mediated by hyperoxidation of peroxiredoxin 2 in senescent cells. *Science China Life Sciences*, 1-15

Lim, J.B., *et al.* (2015) Analysis of the lifetime and spatial localization of hydrogen peroxide generated in the cytosol using a reduced kinetic model. *Free Radical Biology and Medicine* 89, 47-53

Amponsah, P.S., *et al.* (2021) Peroxiredoxins couple metabolism and cell division in an ultradian cycle. *Nature Chemical Biology* 17, 477-484

Sala, D., *et al.* (2019) The Stat3-Fam3a axis promotes muscle stem cell myogenic lineage progression by inducing mitochondrial respiration. *Nature communications* 10, 1796

Sobotta, M.C., *et al.* (2015) Peroxiredoxin-2 and STAT3 form a redox relay for H2O2 signaling. *Nature chemical biology* 11, 64-70

Hopkins, B.L., *et al.* (2018) A peroxidase peroxiredoxin 1-specific redox regulation of the novel FOXO3 microRNA target let-7. *Antioxidants & redox signaling* 28, 62-77

Lundell, L.S., *et al.* (2019) Regulation of glucose uptake and inflammation markers by FOXO1 and FOXO3 in skeletal muscle. *Molecular metabolism* 20, 79-88

De Haes, W., *et al.* (2014) Metformin promotes lifespan through mitohormesis via the peroxiredoxin PRDX-2. *Proceedings of the National Academy of Sciences* 111, E2501-E2509

Wang, X., *et al.* (2022) Ageing induces tissue-specific transcriptomic changes in Caenorhabditis elegans. *The EMBO journal* 41, e109633

Li, S.T., *et al.* (2019) DAF-16 stabilizes the aging transcriptome and is activated in mid-aged Caenorhabditis elegans to cope with internal stress. *Aging Cell* 18, e12896

Murphy, M.P., *et al.* (2022) Guidelines for measuring reactive oxygen species and oxidative damage in cells and in vivo. *Nature metabolism* 4, 651-662

Olahova, M., *et al.* (2008) A redox-sensitive peroxiredoxin that is important for longevity has tissue- and stress-specific roles in stress resistance. *Proc Natl Acad Sci U S A* 105, 19839-19844

Bar-Ziv, R., *et al.* (2020) Measurements of physiological stress responses in C. elegans. *JoVE (Journal of Visualized Experiments)*, e61001

Green, M.R. and Sambrook, J. (2020) Total RNA extraction from Caenorhabditis elegans. *Cold Spring Harbor Protocols* 2020, pdb. prot101683

Supplementary Figure 1 Decay of resistance to paraquat and arsenite with ageing and exercise. Decreased survival to paraquat following acute exercise and ageing (a-d). Decreased survival to arsenite following acute exercise and ageing (e-h). The Log-rank (Mantel-Cox) test was employed to compare survival between distinct groups. *p* values (a: C vs E *p*= 0.0034, E vs R *p*= 0.001; b: C vs E *p*= 0.0435; c: C vs E *p*= 0.022, C vs R *p*= 0.0076; d: D4 vs D8: *p*= 0.0266; D4 vs D12: *p* < 0.0001; D8 vs D12: *p*= 0.0181. e: C vs E *p*= 0.0114, E vs R *p*= 0.0009; f: C vs E *p*= 0.0143, E vs R *p*= 0.0079; g: C vs E *p*= 0.031; C vs R *p*= 0.0048; h: D4 vs D8: *p*= 0.0136; D4 vs D12: *p* < 0.0001; D8 vs D12: *p*= 0.0023)

Supplementary Figure 2 Ageing and the loss of PRDX-2 directly affect MitoSOX staining and SKN-1 activation. MitoSOX red staining of worms for mitochondrial ROS (a) and *gst-4p::gfp* SKN-1 transcriptional reporter (b) following acute exercise at different stages, scale bar = 275 μm. Graphs are the normalised relative means ± SEM and all experiments were performed with n = 30-40 and *p*-value of < 0.05 was considered as statistically significant *(*p* < 0.05), one-way or two-way ANOVA was used for significance between groups (a–b). *p* values (a: in N2 worms: D4: C vs E < 0.0001, E vs R = 0.0004; D8: C vs E = 0.0094; D12: C vs R < 0.0001, T vs R < 0.0001; D4 vs D8 < 0.0001; D4 vs D12 < 0.0001; D8 vs D12 = 0.0004; in *prdx-2* worms: D4: C vs E < 0.0001, C vs R = 0.0004; D8: C vs E = 0.0163, C vs R = 0.0315; D12: C vs T < 0.0001, E vs R = 0.0001; D4 vs D8 < 0.0001; D4 vs D12 < 0.0001; D8 vs D12 = 0.0332; recovery rate: N2: D4 vs D12 < 0.0001; D8 vs D12 < 0.0001; *prdx-2*: D4 vs D12 < 0.0001; D8 vs D12 = 0.0013; N2 vs *prdx-2*: D4 = 0.0453; D8 = 0.0486; D12 = 0.0211. b: in N2 worms: D4: C vs R < 0.0001, E vs R = 0.0217; D4 vs D8 < 0.0001; D4 vs D12 < 0.0001; D8 vs D12 < 0.0001. in *prdx-2* worms: D8: C vs R = 0.0024; D12: C vs E = 0.0014, C vs R = 0.045; D4 vs D8 = 0.0137; D4 vs D12 < 0.0001; D8 vs D12 = 0.0197. recovery rate: N2: D4 vs D8 = 0.0035; D4 vs D12 = 0.0007; *prdx-2*: D4 vs D8 = 0.0448; N2 VS *prdx-2*: D4 = 0.0018; D8 = 0.0001)

Supplementary Figure 3 Ageing and loss of PRDX-2 in adult *C. elegans* adversely impacts DAF-16 nuclear localisation and mitochondrial dynamics in response to exercise. Representative images of the TJ356 strain DAF-16::GFP distribution at adult (a) and L4 (b) stage in *C. elegans*. Increased survival to arsenite in *prdx-2* mutant compared to N2 at L4 stage (c). mRNA level of *eat-3* (d), *ppgn-1* (e), *spg-7* (f), *dct-1* (g), *cpt-1* (h), *skn-1* (i), *tom-20* (j), *hsp-4* (k) and *hsp-6* (l) at different stages. Graphs are the normalised relative means ± SEM and all experiments were performed with n = 130-150 for DAF-16 reporter, n=60 for arsenite survival assays, n = 4 for mRNA expression and *p*-value of < 0.05 was considered as statistically significant *(*p* < 0.05). *p* values (a: in N2 worms: D4: C vs E < 0.0001, E vs R < 0.0001; D8: C vs E < 0.0001, E vs R < 0.0001. b: N2 vs *prdx-2* =

0.0172. c: N2 L4 vs *prdx-2* L4 = 0.0094; N2 D4 vs *prdx-2* D4 = 0.0186. d: in N2 worms: D4 vs D8 = 0.0274; D4 vs D12 = 0.0007. N2 vs *prdx-2*: D8 = 0.0188, D12 = 0.0135. g: N2 vs prdx-2: D12 = 0.0052. i: in N2 worms: D4 vs D8 < 0.0001, D4 vs D12 < 0.0001, D8 vs D12 = 0.0355; in *prdx-2*: D4 vs D8 < 0.0001, D4 vs D12 < 0.0001. N2 vs *prdx-2*: D12 = 0.0011. j: in *prdx-2* worms: D4 vs D12 = 0.0126; N2 vs *prdx-2*: D12 = 0.0013. k: in N2 worms: D4 vs D8 = 0.002, D4 vs D12 < 0.0001; in *prdx-2* worms: D4 vs D8 = 0.0002, D8 vs D12 = 0.0154; N2 vs *prdx-2*: D12: < 0.0001)

Supplementary Figure 4 Ageing and loss of PRDX-2 undermines physical fitness following exercise. Wave initiation rate (a), brush stroke (b), asymmetry (c) and curling (d) following acute exercise at different stages; the recovery rate of these parameters compared to normal condition (e-h). Physical fitness overall 12 days of N2 and *prdx-2* worms (i). Graphs are the normalised relative means ± SEM and all experiments were performed with n = 30-40 and *p*-value of < 0.05 was considered as statistically significant *(*p* < 0.05). *p* values (a: in N2 worms: D4: C vs E = 0.027, E vs R = 0.0037; D8: C vs E = 0.0019, E vs R = 0.0278; D12: C vs E = 0.0063, C vs R = 0.0056; D4 vs D12 < 0.0001; D8 vs D12 = 0.014; in *prdx-2* worms: D4: C vs E = 0.0141; D8: C vs E = 0.022; D12: C vs R = 0.0179; D4 vs D8 = 0.0196; D4 vs D12 = 0.0001. b: in N2 worms: D4 vs D12 = 0.0433. in *prdx-2* worms: D8: C vs E = 0.0432; D4 vs D8 = 0.0318; D4 vs D12 < 0.0001; D8 vs D12 = 0.0133. c: in N2 worms: D4 vs D8 = 0.0164; D4 vs D12 < 0.0001; D8 vs D12 < 0.0001; in *prdx-2* worms: D8: C vs E = 0.016, C vs R = 0.0036; D4 vs D8 = 0.017; D4 vs D12 < 0.0001; D8 vs D12 < 0.0001. d: in N2 worms: D8: C vs E = 0.0161, E vs R = 0.0303; D12: C vs E = 0.0277; D4 vs D12 < 0.0001; D8 vs D12 = 0.013; in *prdx-2* worms: D4: C vs E = 0.0369; D12: C vs E = 0.014; D4 vs D8 = 0.0327; D4 vs D12 < 0.0001; D8 vs D12 = 0.0087. e: in N2 worms: D4 vs D12 = 0.0066; N2 vs *prdx-2*: D4 = 0.0126. g: N2 vs *prdx-2*: D8 = 0.001. i: wave initiation rate: N2 vs *prdx-2* = 0.0007; travel speed: N2 vs *prdx-2* < 0.0001; brush stroke: N2 vs *prdx-2* = 0.015; body wave number: N2 vs *prdx-2* < 0.0001; stretch: N2 vs *prdx-2* < 0.0001; curling: N2 vs *prdx-2* = 0.0098)

Figure	age	Condition	n	Mean lifespan	Log-rank test, *p*-value
Fig 1b	D4	Control	60	21.1 ± 1.14	
		Exercise	60	17.33 ± 1.05	compared to D4 C, p= 0.0034
		Recovery	60	21.83 ± 1.19	compared to D4 E, p= 0.0011
	D8	Control	60	17.57 ± 1.19	compared to D4 C, p= 0.0266
		Exercise	60	15.8 ± 1.08	compared to D8 C, p= 0.0435
		Recovery	60	16.5 ± 1.09	compared to D8 E, p= 0.2379
	D12	Control	60	14.7 ± 1.11	compared to D8 C, p= 0.0181
		Exercise	60	11.83 ± 0.93	compared to D12 C, p= 0.022
		Recovery	60	11.27 ± 0.89	compared to D12 C, p= 0.0076
Fig 1c	D4	Control	60	28.2 ± 1.97	
		Exercise	60	23.37 ± 1.34	compared to D4 C, p= 0.0114
		Recovery	60	30.83 ± 2	compared to D4 E, p= 0.0009
	D8	Control	60	23.77 ± 1.34	compared to D4 C, p= 0.0136
		Exercise	60	19.67 ± 1.19	compared to D8 C, p= 0.0143
		Recovery	60	24.03 ± 1.12	compared to D8 E, p= 0.0079
	D12	Control	60	17.77 ± 1.21	compared to D8 C, p= 0.0023
		Exercise	60	14.6 ± 1.12	compared to D12 C, p= 0.0231
		Recovery	60	13.93 ± 1.08	compared to D12 C, p= 0.0048
Suppl Fig 3c	L4	N2	60	19.3 ± 1.23	
		prdx-2	60	22.9 ± 1.43	compared to N2 L4, p= 0.0094
	D4	N2	60	26.47 ± 1.57	
		prdx-2	60	21.77 ± 1.43	compared to N2 D4, p= 0.0186

Mean lifespan and statistical analysis were determined by OASIS2 platform, Kaplan-Meier curve was performed using GraphPad Prism 7.

Supplementary Table 1 Data from oxidative stress assays related to Figure 1b,c and Suppl Figure 3c.