Sex dependent effects of amyloidosis on functional network 'hub' topology is associated with downregulated neuronal gene signatures in the APP*swe*/PSEN1dE9 double transgenic mouse

Zachary D. Simon[1,2], Karen N. McFarland[3,5], Todd E. Golde[5], Paramita Chakrabarty[2,4], Marcelo Febo[1,2]

1) Department of Psychiatry, University of Florida, Gainesville, FL, USA

2) Department of Neuroscience, University of Florida, Gainesville, FL, USA

3) Department of Neurology, University of Florida, Gainesville, FL, USA

4) Center for Translational Research in Neurodegenerative Disease, University of Florida, Gainesville, FL, USA

5) Department of Pharmacology and Chemical Biology; Center for Neurodegenerative Disease, Emory University, Atlanta, GA, USA

Short title: Sex dependent effects of amyloidosis

Address correspondence to:

Marcelo Febo, PhD,

Email: febo@ufl.edu,

Phone: 352-294-4911

Address: Department of Psychiatry, University of Florida, 1149 S. Newell Dr., Bldg. 059, Room L4-100F, Gainesville, Florida 32611, USA

Abstract

Background: Extracellular amyloid-β (Aβ) impairs brain-wide functional connectivity, although mechanisms linking Aβ to broader functional network processing remain elusive.

Objective: Here, we evaluated the effects of Aβ on fear memory and functional connectome measures in male and female mice.

Methods: Middle-aged (9-11mo) double transgenic APP-PS1 mice and age and sex-matched controls were tested on a fear conditioning protocol and then imaged at 11.1 Tesla. Brains were harvested and processed for analysis of Aβ plaques and Iba1 immunolabeling in cortex, hippocampus, and basolateral amygdala. Additional RNA sequencing data from separate age, strain, and sex matched mice were analyzed for differentially expressed genes (DEGs) and weighted gene co-expression networks.

Results: In both male and female mice, we observed increased functional connectivity in a dorsal striatal/amygdala network due to Aβ. Increased functional connectivity within this network was matched by increases in AβPP gene expression, Aβ and Iba1 immunolabeling, and an upregulated cluster of DEGs involved in the immune response. Conversely, the network measure representing node 'hubness', eigenvector centrality, was increased in prefrontal cortical brain regions, but only in female APP-PS1 mice. This female specific-effect of amyloid was associated with downregulation of a cluster of DEGs involved in cortical and striatal GABA transmission, anxiogenic responses, and motor activity, in female APP-PS1 mice, but not males.

Conclusions: Our results contribute to a growing literature linking between Aβ, immune activation and functional network connectivity. Furthermore, they reveal effects of Aβ on gene expression patterns in female mice that may contribute to amyloidosis-induced dysregulation of non-cognitive circuitry.

Keywords: Amyloid, sex, gene networks, functional connectivity networks, fear memory

Introduction

Early onset Alzheimer's disease (AD) has an estimated prevalence of 1-5%, with the
remaining 95% of all cases representing late onset sporadic forms [1]. Approximately 200,000
individuals in the US suffer symptoms as early as 30-40 years of age, compared to non-familial
AD which emerges at or after 65 [2]. Studies of isolated familial cases have led to the discovery of
several heritable amyloid-β protein precursor (AβPP) mutations and over 200 highly penetrant
mutations in presenilin1 (PSEN1) that are causative for a percentage of early onset cases [3, 4].
Familial AD involves mutation-specified amino acid substitutions in N- or C-terminal domain
residues or Aβ regions of AβPP that cause aberrant AβPP processing, increase amyloid-β (Aβ)
levels and aggregation and extracellular Aβ plaque formation [3]. The Swedish AβPP (AβPP*swe*)
mutations, for example, produces amino acid substitutions (K670N, M671L) near the beta-
secretase cleavage site of AβPP, leading to increased production of a 42 amino acid long Aβ
peptide (Aβ42) [4]. Deletion of exon-9 in the PSEN1 gene sequence impairs gamma-secretase
activity, which results in formation of aggregation prone Aβ42 [4]. Aβ is thought to mediate
functional deficits in neurotransmission, including changes adversely impacting the activity of
both excitatory and inhibitory neurons [5, 6]. Understanding these and other Aβ-linked functional
changes, at the genetic, neuronal and neural circuit scales, is fundamental to uncovering
targetable early pathology and neural signatures driving behavioral and cognitive changes in
early onset and sporadic forms of AD.

Many reports on changes in excitatory and inhibitory neurons and synapses in AD brains
are extrapolated from postmortem assessments [6-9]. These neuronal changes, and their broader
effects on network activity in AD, has been further explored in mice harboring human familial
AβPP mutations [10, 11] and resting state functional magnetic resonance imaging studies (fMRI) [12].
Functional neuroimaging studies of early onset AD have reported widespread functional
connectivity declines across multiple well establish networks [13], particularly in default mode
network (DMN) regions [14]. This includes lower connectome measures of network strength,
efficiency and clustering in frontoinsular, parietal, temporal and occipital cortices [15], and reduced
eigenvector centrality in bilateral medial temporal gyrus, right posterior hippocampal gyrus, and
right postcentral gyrus [16]. Interestingly, there is some disagreement between studies in terms of
the specific regions affected and extent of reductions in functional connectivity and connectome
measures. Pini *et al* reported a greater reduction in hippocampal connectivity to a group of
cortical memory related regions in late onset versus early onset AD [17]. Adriaanse *et al* observed

broad functional connectivity declines in early onset AD compared to late onset, the latter mostly

had reductions in the DMN compared to controls [13]. Yet, others have reported only DMN

reductions relative to controls in early onset AD [14]. All of the above studies use similarly aged

subjects, around the age of 60 and matched control groups. Although pathology of AD is highly

characterized by amyloidogenic mechanisms related to AβPP and PSEN mutations, there are

other disease contributing genes differentially expressed in this population [18, 19]. Moreover, there

are significant sex differences in the genetic architecture conferring resilience to AD, particularly

cognitive resilience [20]. How this translates to functional brain changes in males versus females

in response to AD pathologies remains an area of active research. Thus, the oligogenic

contributions to AD, and important biological factors such as sex, warrant consideration in

experiments to examine mechanistic links between Aβ and brain functional connectivity.

Functional neuroimaging studies in transgenic rodents harboring AβPP, PSEN1 and other

AD causative or moderate-to-high risk mutations, selectively or in combination, can help clarify

specific gene-pathology-brain functional connectivity interactions. A number of animal fMRI

studies have begun to address Aβ-functional connectivity interactions, and these shown results

consistent with some human studies of early onset AD [21-24]. Mouse fMRI studies have revealed

nuanced interactions between Aβ aggregation and functional connectivity outcomes [23], which

calls for attention to how AβPP is processed and how it interacts with other AD genes. Here we

report new findings in the APP-PS1 mouse [25] to expand on the existing literature on the

relationship between Aβ and functional connectome changes. Further, we report a potentially

important link between female-specific effects of Aβ on functional connectome node eigenvector

centrality and brain gene expression. The differentially expressed genes specifically affected by

Aβ in females include several genes in involved in striatal inhibitory gamma amino butyric acid

(GABA) transmission.

117   **Methods**

Subjects and data

Mice were housed in age- and sex-matched groups of 3-4 in a temperature- and humidity-

controlled room, inside conventional air filtered cages (dimensions: 29 x 18 x 13 cm) with food

and water available *ad libitum* (vivarium lights on from 07:00-19:00 hours). Transgenic (TG)

cohorts were generated in house by breeding heterozygous male mice bearing amyloid-β

protein precursor (AβPP) with the Swedish familial mutation and exon 9 deficient presenilin-1

(PSEN1) and hybrid C57BL6/J/6xC3H F1 females (Envigo)[25]. APP-PS1 mice develop gliosis

and plaque accumulation starting at 6mo and cognitive and behavioral impairments as early as

8 mo [26]. Control mice (NTG) were littermate non-transgenic mice and all groups contained male

and female mice. Experiments started when mice were 10-11 months of age(mo), which

corresponds to high amyloidosis, gliosis, with previously reported impairments in cognitive

function and other non-cognitive behaviors [26]. The behavior and imaging experiment group

sizes were 13 NTG controls (including 4 female NTG) and 15 TG mice (including 7 female TG).

The tissue histology group sizes differed between regions based on slice quality. Groups were

8-9 NTG controls (including 3-4 female NTG) and 11-13 TG mice (including female 5-7 TG).

Forebrain RNA sequence data of the APP/PS1 mice were obtained from the AD Knowledge

Portal synapse.org repository (Project SynID: syn2580853, title: "The Tau and AβPP Mouse

Model (TAUAPPms) Study"). The dataset included 9-12 mo NTG (n=15, including 7 NTG

females) and 12 mo TG mice (n=11, including 5 TG females). All procedures received prior

approval from the Institutional Animal Care and Use Committee of the University of Florida and

follow all applicable NIH guidelines.

Fear conditioning

Fear conditioning was performed two weeks before fMRI studies began. The fear

conditioning system and methods used were previously described [27]. The fear conditioning

cages were housed in sound attenuation chambers. The chambers had computer-controlled

lighting, white noise generator, a fan, and control units. These included audio speaker controls

for tone generation, shock controllers for the internal cage floors, and infrared and white lighting

controllers. The internal operant cage was made of translucent Plexiglas with a steel frame and

steel shock grid floor connected to a computer controlled current stimulator (Med-Associates,

Inc. St. Albans, VT). Activity within the fear conditioning cages was captured by a monochrome

video camera. All cage accessories were controlled by PC running Video Freeze Software™.

Camera lens brightness, gain, and shutter settings were verified and adjusted before collecting

data and were kept constant across all mice. For each session, the detected motion-sensitive

signals were calibrated with the mouse inside the cage and NIR lights on. The estimated motion

index was set to the same threshold level (10 a.u.) across all mice prior to data exporting or

analyses. Videos were acquired at 30 frames per second (fps) with a minimum freeze duration

of 30 frames. For training, on the first day a 180 second baseline was followed by four

presentations of a 20 second tone (90 dB, 5kHz, 50ms risetime) with the presentation of

moderate level current to the entire grid floor for 1 second (0.9mA) (inescapable shock) on the final second before the tone ended. Each of the four tone-shock presentations were separated by 190 seconds and the entire session lasted 17 minutes. On days 2 and 3 (recall tests at 24 and 48 h of training day 1, respectively), the same protocol was carried out in the absence of the electric shock. On day 3 (modified context test), the cage environment was modified by placing a plastic smooth white floor and diagonal walls, which covered the grid floor, light, and speaker. For each of the time epochs corresponding to the 4 tone-shock pairings, a component comprising a 20 second pre-shock tone interval and a 40 second post-shock interval were used to quantify percent component freezing (per day there were four 60 second components summed to provide total percent freezing per day). Motion-index was used as a measure of locomotor activity.

Magnetic resonance imaging instrumentation

Images were collected on a magnetic resonance spectrometer tuned to 470.7 MHz proton resonant frequency (Magnex Scientific 11.1 Tesla/40 cm horizontal magnet). The MR system was equipped with Resonance Research Inc. spatial encoding gradients (RRI BFG-240/120-S6, maximum gradient strength of 1000 mT/m at 325 Amps and a 200 μs risetime) and controlled by a Bruker AV3 HD console running Paravision 6.0.1 (Billerica, MA). An in house built 2 cm x 2.5 cm quadrature transceive surface radiofrequency (RF) was used for B1 signal generation and detection (RF engineering lab, Advanced Magnetic Resonance Imaging and Spectroscopy Facility, Gainesville, FL).

Mouse imaging setup

Mice were scanned sedated under a continuous paranasal flow of 0.25 % isoflurane gas (delivered at 0.04L/min mixed with medical grade air containing 70% nitrogen and 30% oxygen, Airgas, Inc.) and a continuous subcutaneous infusion of dexmedetomidine. Prior to setup, mice were anesthetized under 2% isoflurane and administered an intraperitoneal (i.p.) injection of 0.1mg/kg dexmedetomidine (at a 1ml/kg volume). Isoflurane was reduced to 0.25% throughout the remaining imaging session. An infusion line was setup for subcutaneous delivery of dexmedetomidine over the course of scanning (0.1mg/kg/ml at an infusion rate of 25 μl/h using a PHD-Ultra microinfusion pump, Harvard Apparatus, Holliston, MA). Functional MRI scans were collected at least 50 minutes after the i.p. injection. Respiratory rates were monitored

continuously, and body temperature was maintained at 36-37°C using a warm water
recirculation system (SA Instruments, Inc., New York).

Functional MRI acquisition protocol

We acquired one T2 weighted anatomical and an fMRI scan per mouse. The T2-weighted
Rapid Acquisition with Relaxation Enhancement (RARE) sequence was acquired with the
following parameters: effective echo time = 41.4 ms, repetition time (TR) = 4 seconds, RARE
factor = 16, number of averages = 12, field of view (FOV) of 15 mm x 15 mm and 0.8 mm thick
slice, and a data matrix of 256 x 256 (0.06 μm^2 in plane) and 12 interleaved ascending coronal
(axial) slices covering the entire brain from the rostral-most extent of the anterior prefrontal
cortical surface, caudally towards the upper brainstem and cerebellum (acquisition time 9
minutes 36 seconds). Functional images were collected using a single-shot spin echo planar
imaging sequence with the following parameters: TE = 16 ms, TR = 1.5 seconds, 600
repetitions, FOV = 15 x 15 mm and 0.9 mm thick slice, and a data matrix of 64 x 48 (0.23 x 0.31
μm in plane) with 14 interleaved ascending coronal slices in the same position as the
anatomical scan. Ten dummy EPI scans were run prior to acquiring data under steady state
conditions. Respiratory rates, isoflurane and dexmedetomidine delivery, body temperature,
lighting, and room conditions were kept constant across subjects.

Image pre-processing

The image processing workflow followed previously published research [27-31]. Resting state
processing was carried out using Analysis of Functional NeuroImages (AFNI) [32], FMRIB
Software Library (FSL) [33], and Advanced Normalization Tools (ANTs) [34]. Binary masks for
anatomical and functional scans were created using ITKSNAP [35]. The brain binary masks were
used for brain extraction prior to registration steps. Times series spikes were removed
(3dDespike, AFNI), image repetition frames aligned to the first time series volume (3dvolreg,
AFNI), and detrended (high pass filter <0.009 Hz using AFNI 3dTproject). Independent
component analysis (ICA) using Multivariate Exploratory Optimized Decomposition into
Independent Components (FSL MELODIC version 3.0) was used to assess structured 'noise' or
artefact components in each subject scan, in their native space. Most, if not all ICA components
in this first stage contained artefact signal voxels along brain edges, in ventricles, and large
vessel regions. These components were suppressed using a soft ('non-aggressive') regression

approach, as implemented in FMRIB Software Library (FSL 6.0.5) using fsl_regfilt [33]. A low-pass

filter (>0.12Hz) and spatial smoothing (0.4mm FWHM) was next applied to the fMRI scans prior

to registration steps. Post-regression ICA was carried out to verify removal of artefact

components and preliminary assessment of putative ICA networks in individual scans.

Atlas registration and resting state signal extraction

Anatomical scans were cropped and bias field corrected (N4BiasFieldCorrection, ANTs).

Functional scans were cropped, and a temporal mean image was registered to the anatomical.

Preprocessed anatomical and fMRI scans were aligned to a parcellated mouse common

coordinate framework (version 3, or CCFv3) template [36]. Bilateral region of interest (ROI)-based

nodes (148 total) were created with the guidance of the annotated CCFv3 parcellation and using

tools in ITKSNAP and FSL, similar to our previous work in rats [37, 38]. Large brain regions, such

as hippocampus, motor, somatosensory, and visual cortices, were assigned multiple nodal

masks. Thus, these are distinguished based on atlas coordinates and a numerical identifier at

the end of the regional code (e.g., hippocampus contains nodes HPC1-HPC5). Anatomical

images were linearly registered to the mouse template using FSL linear registration tool

(FLIRT), using a correlation ratio search cost, full 180-degree search terms, 12 degrees of

freedom and trilinear interpolation. The linear registration output was then nonlinearly warped to

template space using ANTs (antsIntroduction.sh script). Anatomical-to-atlas linear and nonlinear

transformation matrices were applied to fMRI scans at a later stage. Spontaneous signals, each

with 600 data points, were extracted from 148 ROIs and used in cross-correlations and in

calculations of Pearson r coefficients for every pairwise combinations of ROIs (using *corrcoef*

function in MATLAB). The resulting number of pairwise correlations was 10,730 per subject

(after removing 148 self-correlations). Correlation coefficients were Fisher's transform to ensure

normality prior to statistical analyses. The center voxel coordinates for the ICA-based nodes

normalized to the CCFv3 were used in 3D network visualizations using BrainNet viewer [39].

Network analysis

Weighted undirected matrices were analyzed using Brain Connectivity Toolbox [40] in

MATLAB (Mathworks, Natick, MA). Global graph metrics were calculated for edge density

thresholds ranging from 2-40%. Global network measures for this density range were converted

to area under the curve (AUC) values prior to statistical assessments. Node-specific network

measures were converted to AUC values per node. We assessed several graph measures of

network integration and communication efficiency in APP-PS1 and control mice. This included

node strength (sum of edge weights/node) and degree (sum of edges/node) and global

measures, such as transitivity (related to CC; number of triad groups normalized by all possible

triad nodes in a network), and characteristic path length (CPL; the average edges or edge

weights between node pairs). For local node efficiency, a length matrix was used to calculate

inverse values in vicinity of a node, with added weights used to emphasize the highest efficiency

node paths [40, 41]. To corroborate results relative to random networks, all network measures were

calculated on original and randomized versions of the functional connectivity matrices. Positive

and negative edges were randomized by ~5 binary swaps with edge weights re-sorted at each

step. A probabilistic approach for community detection was used to calculate a modularity

statistic (Q), which indexes the rate of intra-group connections versus connections due to

chance [42]. The procedure starts with a random grouping of nodes and iteratively moving nodes

into groups which maximize the value of Q. The final number of modules and node assignments

to each group (e.g., community affiliation assignments) was taken as the median of 100

iterations of the modularity maximization procedure [37]. We analyzed the tendency of assortative

vs dissortative mixing of nodes [43]□. The assortativity index is a correlation coefficient

comparing node strength values between pairs of edge-connected nodes. Positive r values

indicate connectivity between pairs of nodes with similar strengths (e.g., high strength nodes

pairs with high and low with low), while negative r values indicate cross-pairings between low

and high strength nodes. Functional connectivity networks were visualized in BrainNet viewer [39].

Immunohistochemical labeling of amyloid plaques and microglial cells

Mouse brains were harvested and processed for immunohistochemistry. Formalin fixed

brain hemisections were paraffin embedded and sectioned sagittally using microtome at the

following Bregma locations: lateral 0.12mm, 1.56mm, and 3.25mm. Consecutive 4-µm were

collected at each level for staining. The primary antibodies were Iba1 (1:1000, Wako Catalog #

019-19741) and 33.1.1 for staining Aβ plaques (1:13,000; provided by the Chakrabarty

laboratory, Center for Translational Research on Neurodegenerative Diseases, University of

Florida; kind gift of Todd E Golde, Emory University). Whole slides were scanned on a

ScanScope XT Scanner (Leica Biosystems). Images were analyzed with tuned positivity

algorithms for each respective stain using the Aperio ImageScope software. Regions were

hand-drawn on each image using the Paxinos and Franklin's Mouse Brain Atlas in Stereotaxic

Coordinates 4th ed. as a guide. The images had the following regions segmented out: prefrontal
cortex, sensorimotor cortex, and hippocampus (at approx. Bregma coordinates lateral 1.76),
and basolateral amygdala (at approx. Bregma coordinates lateral 2.88). The positivity threshold
(which highlights the dark (stained) tissue beyond a certain threshold) was tuned to highlight the
staining of each respective stain, but not the surrounding tissue. Tuning was done on a single
image for each stain to obtain the threshold which would produce most signal from only the
stained tissue and not surrounding unstained tissue. This threshold for each stain was kept the
same for all regions and all subjects throughout the analysis. The obtained positivity values
were the percentage of pixels positive for staining at the tuned threshold, within the given drawn
region. Each animal has a single positivity value from the specified region drawn on the image
of the stained slide containing that region. 2-way ANOVA with main effects and interaction of
sex and genotype was used to determine p-values for differences between groups within each
specified brain region.

Normalized gene counts and analysis of differentially expressed genes (DEGs)

BAM files were downloaded from Synapse.org as described (doi:
https://doi.org/10.7303/syn2580853). Rsamtools and GenomicAlignments R packages [44, 45] were
used to generate gene counts from the BAM files and were filtered for gene counts to remove
genes with low counts (<10 genes). Subsequent analyses used normalized gene counts
expressed as fragments per kilobase of transcript per million mapped reads (FPKM values).
Analyses of DEGs was carried out using DESeq2 using default setting and a general linear
model fit per gene [46]. Wald tests were used to compare TG vs NTG groups and multiple
comparison adjustments of resulting p-values was done with the Benjamini-Hochberg false
discovery rate (FDR) correction method. DEGs were defined as an absolute log2 fold change >
1 and an adjusted p value < 0.05 (visually evaluated as volcano plots). Gene annotation
symbols were used to group genes according to their known expression in mouse brain cell
types [47]. Within these cell type classifications, the geometric mean of the normalized gene
counts for the genes comprising the cell types was calculated and analyzed between
experimental groups.   The cell type classifications included astrocytes, neurons,
oligodendrocyte precursors, newly formed oligodendrocytes, myelinating oligodendrocytes,
endothelial cells, and microglia/macrophage cells [47].

Weighted Gene Co-Expression Network Analysis (WGCNA)

The WGCNA R package was used to construct gene correlation networks from the normalized expression data after filtering and removing genes with zero variance [48, 49]. Plots of scale-free topology fitting and mean connectivity, both as a function of soft-threshold power values ranging from 1-20, were evaluated to select a gene network threshold using the "pickSoftThreshold" function. Networks were constructed all male and all female samples separately. Adjacency matrices for a signed hybrid network were constructed using expression data and the chosen power threshold. The adjacency matrices were converted to topological overlap maps (TOMs) to be used for module network graphs. Hierarchical clustering was used to identify modules with the cutreeDynamic function. A deepSplit setting of two with a minimum module size of 30 was used for all analyses. Dendrograms were evaluated and compared by identifiable color-coded module statistics. Modules containing highly co-expressed genes were merged and co-expression similarity of entire modules were quantified by first calculating eigengenes and then clustering these based on their Pearson correlations. A cluster height cut was 0.25 (correlations of 0.75) was applied to merge similar modules. To quantify module-trait associations we recalculated module eigengene correlations and p values and displayed these as heat maps. Gene-trait and gene-module relationships were analyzed based on gene significance and module membership statistics. Module hub genes were determined with the 'chooseTopHubInEachModule' function.

For module preservation testing, female data were used as a reference and male data tested against the reference for low ($z < 2$), moderate ($z>2$, $z<10$), high module preservation ($z > 10$). We used sex as a comparison trait since preliminary assessments indicated sex differences in DEGs and cell type compositions. Statistics were run as multiExpr data for signed hybrid networks with 200 permutations and pariwise comparisons are made between common genes. Intramodule degree for eigengenes (kME) is used as a module membership statistic plotted against z score converted summary statistics determining significance. All statistical comparisons for preservation p values were Bonferroni-corrected.

Statistical analysis

Statistical analyses and data visualization were carried out using tools in MATLAB, GraphPad Prism (version 10.2) or ggplot2 in the R statistics package. Unless otherwise stated, statistical analyses used either a two-way full factorial analysis of variance (ANOVA) (genotype

x sex, critical α<0.05) or Mann-Whitney tests. Post hoc tests used Tukey's honest significant

difference (HSD) procedure. Where appropriate, FDR correction (q ≤ 0.05) was used. Further

analysis of fMRI scans was carried out using probabilistic ICA [50] as implemented in MELODIC

Version 3.15, part of FSL [50, 51] and as previously described for mouse fMRI scans [27]. The

resulting components were overlaid on the CCFv3 mouse atlas template and classified

according to the peak z statistic anatomical location. A series of two-stage multiple linear

regressions were used to back-project group ICA components to subject-specific time courses

(using spatial regressions) and spatial components (using temporal regressions). The subject-

specific spatial component maps were then used in statistical comparisons for each component

in the FSL randomise tool. The statistical design matrix was generated using the FSL Glm tool.

We used a one-way ANOVA general linear model design. Randomization tests were carried out

with 500 permutations (corrected p-level for significance is 0.05 and statistical thresholding of

maps by threshold free cluster enhancement).

364   **Results**

Female APP-PS1 mice have a higher amount of amyloid plaques in basolateral amygdala and

sensorimotor cortex than males

Histological analyses confirmed Aβ plaque presence across hippocampal, sensorimotor,

prefrontal and basolateral amygdala regions in TG mice (Figure 1a and 1c). Additional analyses

indicated that female TG mice had a greater amount of Aβ plaques in basolateral amygdala

region (Mann-Whitney p = 0.0087)) and in neocortical areas corresponding to somatomotor

cortex (p = 0.035) (Figure 1b). The hippocampus had a similar trend (p =0.051) and prefrontal

cortex was not significantly different between male and females.

Amyloidosis increased functional connectivity within a posterior striatal/amygdala network

ICA identified 20 components that included previously reported networks [27, 52-54]. These are

summarized in Supplementary Figure 1. The components had peak z-statistic voxels in anterior

and posterior cingulate, ventral and dorsal striatal, motor, somatosensory, visual cortices,

thalamic, midbrain, and cerebellar regions. Statistical analysis using a two factor ANOVA

(genotype x sex) indicated no significant main effects or interactions in established ICA

networks. We analyzed an unclassified mixed network containing peak z score voxels in the left

posterior striatum/basolateral amygdala (Figure 2a). This network included connectivity with

cerebellar, prelimbic, motor, posterior cingulate, ventral hippocampus, entorhinal cortex regions.
Two-way ANOVA revealed a main effect of amyloidosis, with TG mice having greater functional
connectivity across these and other regions compared to NTG mice. No main effect of sex and
sex x genotype interactions were observed for this mixed network (Figure 2b). Figure 2b
summarizes the brain areas with greater functional connectivity in TG vs NTG mice. These
included subregions of the motor cortex, nucleus accumbens, insula, midbrain, piriform cortex,
entorhinal cortex and reticular nuclei ($p < 0.05$, threshold free cluster enhanced corrected).

Amyloidosis is associated with higher eigenvector centrality scores in prefrontal and striatal
subregions in female mice

Global network measures did not vary significantly between NTG and TG groups, regardless
of sex (Supplementary Figure 2a). Figure 3a shows nodal eigenvector centrality scores
calculated at 10% edge density. Eigenvector centrality scores at 10% edge density in several
prefrontal and striatal nodes differed as a function of either sex or genotype (in the case of
female mice) (Figure 3a). A similar difference was not observed in male mice, which had
eigenvector centrality scores similar to TG female mice in these brain areas. TG females had
higher eigenvector scores compared to NTG females in a single node in the left prelimbic cortex
and two in the right prefrontal cortex, left lateral and ventral nodes in the orbital cortex, a node in
the dorsomedial division of the left striatum (group[sex/genotype] x node interaction,
$F_{441,3528}=1.2$, $p=0.005$; multiple between-group comparison t-tests per node with FDR adjusted p
values at $q < 0.05$) (Figure 3a). NTG males had higher eigenvector scores compared to NTG
females in the same nodes and also in left dorsal agranular insular cortex, right anterior
cingulate cortex, right infralimbic cortex, right lateral orbital cortex, right secondary motor cortex,
and right dorsomedial striatum (FDR adjusted $p < 0.05$ for comparisons across all nodes). When
eigenvector scores were analyzed as AUC values per node, we also observed similar regions
with lower eigenvector centrality scores in NTG compared to TG females and NTG females
compared to NTG males (FDR adjusted $p < 0.05$ for comparison across all nodes). Node
eigenvector centrality scores in prelimbic cortex, left lateral orbital cortex, left hippocampus, left
dorsal agranular insula, left dorsolateral striatum were greater in TG compared to NTG female
mice (Figure 3b). Eigenvector centrality in left and right prelimbic cortex, left hippocampus, right
anterior cingulate cortex, and right orbital cortex were greater in NTG males than NTG female
mice (Figure 3b). It should be noted that both left and right hemispheres of these brain regions
had similar differences between male and female NTG and female NTG compared to female TG

(uncorrected p values < 0.05). However, only the mentioned brain regions were significant after

FDR adjustment of p values. The summary 3D functional connectome maps in Figure 3c

illustrate the average AUC eigenvector score brain distribution in each group.

Within-sex effects of amyloidosis on conditioned fear learning and recall

No between-group differences in fear learning were observed (Supplementary Figure 3).

Overall behavioral data was analyzed using 2-way ANOVA of main effects and interactions of

genotype and trial, with Tukey post-hoc analysis for multiple comparisons. Both TG and NTG

mice increased percent time freezing over the course of shock-tone exposure trials on the first

day of testing (Supplementary Figure 3a). Comparison across days also failed to show any

differences between NTG and TG mice on any days (Supplementary Figure 3b). No differences

in motor activity were observed on any of the three test days (Supplementary Figure 3c). Within-

subject analysis of day one of fear conditioning did reveal differences between groups, including

sex differences. First, as expected, we observed an effect of trial on day 1 of testing (main effect

of trial, $F_{3,72}$ = 8.8, p < 0.0001). However, Tukey's multiple comparison test indicated that only

male mice developed fear conditioning when comparing trials 3 and 4 to the first shock-cue

pairing (NTG males: p = 0.0047 trial 1 vs. 3 and p = 0.0017 trial 1 vs. 4; TG males: p = 0.03 trial

1 vs. 3 and p = 0.02 trial 1 vs. 4). Percent time freezing in trials 3 and 4 compared to the first

trial were not significant in NTG and TG female mice (Supplementary Figure 3a). Within-subject

analyses also revealed significant differences in fear conditioning over the 3 test days (main

effect of day, $F_{2,48}$ = 17.4, p < 0.0001; Supplementary Figure 3b). Percent time freezing was

maintained across days in NTG males. Male TG mice showed an increase in percent freezing

time on days 2 and 3 relative to day 1 (p = 0.0035 and p = 0.01, respectively). Female NTG and

TG mice both showed increases in percent time freezing on day 2 compared to day 1 (female

NTG, p = 0.0027 for day 1 vs. 2; female TG, p = 0.026 for day 1 vs. 2). However, percent time

freezing did not differ between day 3 (in a modified context) relative to day 1 in female NTG and

TG mice. No differences in motor activity were observed within-group across days.

Unique neuronal and oligodendrocyte gene signatures associated with APP-PS1 familial AD

mutations in female mice but not in males

Analysis of DEGs revealed a substantially elevated number of gene signatures in TG

relative NTG female mice (Figure 4a). This included 216 upregulated and 24 downregulated

genes (log2FC > 1.0, adjusted p < 0.05). DEGs in male mice included 54 upregulated and 5

downregulated genes (log2FC > 1.0, adjusted p < 0.05) (Figure 4b). We used two-way ANOVA

to assess cell type composition for normalized gene counts, using geometric means as a

composite measure for changes in cell type gene signatures [55] (results for Tukey's post hoc

tests are shown in Figure 4c). We observed significant differences in geometric means for

microglial (genotype $F_{1,22}$ = 93.5, p <0.0001), endothelial (sex $F_{1,22}$ = 12.78, p = 0.0017) and

vascular pericyte (sex $F_{1,22}$ = 5.8, p = 0.02; genotype $F_{1,22}$ = 28.0, p < 0.0001) genes in TG vs

NTG mice (Figure 4c, top row). Both male and female mice showed similar geometric means for

these gene categories. Conversely, we observed reductions in geometric means for neuronal

(genotype x sex $F_{1,22}$ = 8.25, p= 0.0088) and myelinating oligodendrocyte (genotype x sex $F_{1,22}$ =

6.1, p= 0.02; genotype $F_{1,22}$ = 8.9, p= 0.0067) and newly formed oligodendrocyte (genotype x

sex $F_{1,22}$ = 5.1, p= 0.03; genotype $F_{1,22}$ = 13.1, p= 0.0015) cell type categories in TG female mice

relative to NTG females (Figure 4c, bottom row). A similar non-significant trend was observed

for oligodendrocyte precursor category. This was not observed in male mice (Figure 4c, bottom

row).

These results highlighted an interesting relationship between amyloid and neuronal and

white matter related gene signatures, specific to female mice. WGCNA of RNAseq data

revealed 10 modules with significant positively correlated genes and 7 modules with significant

negatively correlated genes between TG vs NTG female mice (Figure 5a). Among the identified

modules, we further analyzed gene signatures within the brown and lightblue4 modules (Figure

5b-c, respectively). The former is comprised of 669 positively correlated genes (r = 0.91, p =

1.87E-05), of which the top ten included genes related to immune activity (macrophage

associated Cd68 was considered the 'hub' gene in this module). The latter module is comprised

of 439 negatively correlated genes (r = -0.81, p = 6.8E-04), with the top10 most significant

genes known for their roles in neuronal and behavioral functions (Gng7 was considered the

'hub' gene in this module). Statistical hierarchical cluster heatmaps in Figure 5d-e summarize

comparisons within these two modules between TG and NTG female mice. This analysis

illustrated a significant segregation of upregulated and downregulated gene clusters between

NTG and TG female mice within each module.

Module preservation analysis indicated that the brown module containing positively

correlated genes was highly preserved between male and female mice (Figure 6a-b). The

lightblue4 module containing negatively correlated genes was moderately preserved in male

relative to female mice (Figure 6a-b). Consistent with this result, analyses of normalized gene

counts for the brown module showed significant increases for the first top 5 genes in both
female and male mice (Figure 6c). These included Cd68 (genotype $F_{1,22}$ = 165, p < 0.0001),
Cd52 (sex x genotype $F_{1,22}$ = 6.6, p = 0.02), Klhl6 (genotype $F_{1,22}$ = 126, p < 0.0001), Irf8
(genotype $F_{1,22}$ = 93.6, p < 0.0001), and Cd14(genotype $F_{1,22}$ = 27.5, p < 0.0001). Analysis of
normalized gene counts for the top ten gene signatures in the lightblue4 module indicated that
these were significantly reduced only in APP-PS1 female mice relative to NTG female controls
(Figure 6d). These included Gng7 (sex x genotype $F_{1,22}$ = 4.98, p = 0.036), Syndig1l (sex x
genotype $F_{1,22}$ = 8.7, p = 0.0075), Pde1b (sex x genotype $F_{1,22}$ = 8.5, p = 0.008), Lrrc10b (sex x
genotype $F_{1,22}$ = 7.8, p = 0.01), Vps26b (genotype $F_{1,22}$ = 15.6, p = 0.0007), and Ppp1r1b (sex x
genotype $F_{1,22}$ = 9.9, p = 0.004).

Amyloidosis is associated with similar microglia abundance in male and female mice

Our histopathological analysis also confirmed the presence of Iba1-labeled microglial cells,
which appeared in greater abundance in TG compared to NTG mice (Supplementary Figure
4a). Iba1 labeling was greater in hippocampus (p = 0.0004), somatomotor (p < 0.0001) and
prefrontal cortex (p = 0.0008) in male and female TG mice relative to their corresponding sex
matched NTG controls (Supplementary Figure 4b). 2-way ANOVA with main effects and
interaction of sex and genotype was used to determine statistical significance.

499   **Discussion**

In this study we examined the effects of AD-amyloidosis on brain wide functional
connectivity and gene expression in middle-aged mice. Our data reveal brain network and gene
network correlates involved in the complex relationship between Aβ, functional connectivity, and
cognitive/affective behavior in the APP-PS1 (line 85) mouse. These mice develop hippocampal
Aβ deposits by 6 mo and abundant diffuse and dense core plaques throughout the cortex and
hippocampus by 9 mo [25]. Based on substantial evidence of Aβ plaque induced changes in
synaptic activity and plasticity in neocortical and hippocampal areas [7, 8, 10, 11, 56-59], we
hypothesized functional connectivity differences as a result of amyloidosis in the APP-PS1
mouse. Surprisingly, functional connectivity in well-known mouse ICA networks [52, 54, 60, 61],
including somatosensory, motor, anterior and posterior cingulate cortices known to develop
abundant extracellular plaques in this AβPP mouse strain [62], did not differ between TG and NTG
mice. The lack of effects of amyloid on functional connectivity is likely due to the mechanism of

Aβ aggregation in the APP-PS1 mouse. ΔE22Aβ mice with intracellular aggregation of Aβ and PSAβPP mice with parenchymal plaque deposits showed either no differences in functional connectivity or exhibited mild differences compared to control mice [23]. On the other hand, Arctic AβPP mice (ArcAβ) with vascular and intracellular Aβ aggregates had significantly reduced cortical functional connectivity in mice [23]. It will be important in future imaging studies to clarify how specific forms of Aβ aggregation affect neuronal circuit activity and large-scale functional connectivity networks.

Despite no observable effect of amyloid on established mouse ICA networks, we did observe differences in functional connectivity between NTG and TG mice within a mixed or unclassified striatal-amygdala network. One objective of this study was to report all networks rather than screening for specific networks from previous literature. This previously unreported network with significant functional connectivity differences between NTG and TG mice was not restricted to previously reported networks or functional systems. Rather it combines regions and structures not thought to be network specific. This is a result of interactions between regions not classically defined in networks which may have been due to sex and transgene specific activity changes. While this network is speculative at this time, ICA decomposes data based on intrinsic connectivity, meaning the statistical separation of this network is not likely to be an artifact of our processing or scan protocol. Peak z-statistic values were observed in voxels located in posterior striatal-basolateral amygdala. The brain areas showing functional connectivity within this network included sensorimotor, cingulate, ventral hippocampal, midbrain and cerebellar regions. The diffuseness of the brain areas included in this ICA component somewhat parallel the similarly diffuse distribution of Aβ plaques in APP-PS1 mice. Within this ICA network, TG mice had higher functional connectivity than NTG mice in motor cortex, orbital cortex, insular cortex, temporal cortex, entorhinal cortex, dorsomedial striatum and midbrain. A shared feature of these brain areas is that they have either first or second order projections to or from the hippocampus, particularly ventral CA1 [63-67]. A possible mechanism underlying the coactivation of structures in this ICA component, and the observed amyloid related differences in functional connectivity, is the compensatory plasticity of local circuits and projection neurons of ventral hippocampus that may occur in response to extracellular Aβ plaques [68]. Such compensatory increases in functional connectivity are likely to fail at much later ages since prior evidence indicates that at 18 months of age interhemispheric fMRI connectivity is reduced in APP-PS1 male mice relative to C57BL6/J wildtype mice [21]. We should note that hypersynchronous activity in parietal and temporal cortical areas were reported in 18-24 mo 3xTg-AD mice [69]. This was correlated with tau but not Aβ burden [69]. This result in 3xTg-AD mice

differs from observations of Aβ associated hyperexcitability in fMRI connectivity in AD, which did

not correlate with tau [70].

DEG analysis suggested that the observed differences in functional connectivity are linked

to changes in neuroimmune activity, particularly microglia activation. Analyses of gene

enrichment modules revealed gene clusters playing roles in the immune response, cell surface

adhesion and receptor-mediated signaling, all elevated in conjunction with elevated human

mutant AβPP in male and female mice. This was consistent with histological data showing

similarly elevated microglial Iba1 immunostaining in cortical and hippocampal tissue in male and

female TG mice. This is an expected result of amyloidosis, which has been reported in AD and

across studies using AβPP mouse strains. It is also consistent with timelines of expression

reported in *in vivo* PET studies of 12-13 mo APP-PS1 using radioligand with affinity for the

translocator protein (TSPO) [71]. Interestingly, temporal cortical TSPO but not Iba1 follows a

staged increase pattern similar to Aβ in immunolabeling studies of postmortem AD cases [72].

Whether the elevated abundance in microglial cells contributes to the reorganization of

functional connectivity is unclear. Microglial cells are known to play a phagocytic role in synaptic

pruning [73], particularly of unused or damaged dendritic spines [74], and this in turn may contribute

to optimization of neural circuit processing [73, 75-77]. Mice deficient of the Crx3cr1 chemokine

receptor have reduced microglia and this is associated with reduced functional coherence

between the prefrontal cortex and hippocampus and reduced functional connectivity [75]. Crx3cr1

knockout mice also exhibit reduced synapses in adult born dentate gyrus cells along with

structural changes in neurons within this region [78]. Iba1 deficiency in the Allograph inflammatory

factor1 (Aif1) knock mouse reduces hippocampal CA1 spines and pyramidal firing frequencies

[79]. Conversely, lifelong depletion of microglia in mice with deletion of the fms-intron regulatory

element (Csfr1$^{\Delta FIRE/\Delta FIRE}$ mice) showed no differences in the number of excitatory and inhibitory

synapses nor spine density in the hippocampal CA1, although neurons within this region did

show deficits in excitability and altered synaptic properties [80].

The observed Aβ related changes in functional connectivity and DEGs were not

accompanied by differences in fear conditioning between TG and NTG mice, which was

unexpected given previous behavioral characterization of the AβPP mouse [26, 81, 82]. We did not

observe differences between NTG and TG mice on any of the test days, although a trend

towards lower freezing responses in male TG mice was observed on day 1. This finding

suggests that the observed brain functional connectivity and gene expression changes might

precede the effects of Aβ on cognitive and emotional behaviors. It is worth noting that changes

in functional connectivity and AB plaque related PET signals have been reported in elderly

individuals without cognitive deficits [83, 84]. Although this seems plausible in the present mouse

imaging study, there are alternative explanations for the lack of differences in fear conditioning

that requires consideration. The combined classical-contextual fear conditioning assay (tone-

context-shock pairing) and the moderate-to-high current level for shock delivery used in the

present study could account for differences relative to previous work using a standard

contextual fear conditioning assay (context-shock pairing) [26]. As discussed in recent work by our

group [27], age- and strain-related declines in sensorimotor function can impact performance on

fear conditioning assays and the outcomes may reflect underlying mechanisms not directly

related to impaired learning and memory. The hybrid classical-contextual fear conditioning

assay using auditory plus contextual cues, and a higher moderate current level, was used here

to induce a robust learning and memory performance in APP-PS1 mice. However, this

approach could have masked milder cognitive declines observed in previous work using

contextual cues only [26].

Our results reveal significant sex dependent effects of amyloidosis on histopathological,

gene expression, functional brain activation and fear conditioning measurements. This is

consistent with previously reported sex differences in this mouse model[85]. We should note that

while the Prnp promoter has been associated with sex dependent patterns in gene expression

in peripheral tissues, this has not been observed for brain[86]. Analysis of the sex dependent gene

Eif2s3x[87] alongside App and Prnp in the present cohort of mice illustrates that the latter genes

were not differentially expressed by sex in NTG mice (Figure 6c and Supplemental Figure 5).

We observed sex differences in eigenvector centrality, a measure of node 'hubness', which was

differentially altered by amyloidosis in female compared to male mice. The brain regions with

highest eigencentrality in both male and females included the anterior thalamus, dorsal striatum

and areas of medial prefrontal cortex and orbital cortex (illustrated in Figure 5c). Functional

connectome eigenvector centrality was lower in several prefrontal areas and striatum of NTG

female mice relative to NTG males. Sex differences in centrality in functional connectivity

networks have been reported in human cortex [88] but in no other rodent fMRI studies to date. In

humans, women reportedly have higher eigenvector centrality scores in hippocampus and

posterior cingulate cortex than men [88]. We tested another centrality measure, betweenness

centrality, which on average resulted in essentially identical distribution of node betweenness

centrality rankings across male and female mice. This suggests that the observed sex

differences in eigencentrality are not generalized to other network centrality measures. It is

unclear if this topological organization has a neurophysiological underpinning relevant to
cognitive or emotional states in humans and other species.

AβPP female mice had higher eigenvector centrality values in prefrontal cortical regions
than controls. Indeed, eigenvector centrality in AβPP female mice were comparable to TG and
NTG male mice. This may have significance to female-specific effects of amyloid in modulating
the functional and behavioral roles of prefrontal brain regions. This is supported by research
showing that PET-based amyloid positivity is related to increased eigenvector centrality in
prefrontal cortex, even in the absence of declines in cognitive performance [89]. The link between
amyloid positivity and eigenvector centrality in cognitively healthy controls appears to be
relatively specific to the default mode network [89, 90]. The increased eigenvector centrality
observed in posterior and mid-cingulate cortical regions correlates with elevated cerebral spinal
fluid (CSF) markers of Aβ and phosphorylated tau (p-tau) [91] and this in turn relates to cognitive
impairment [91]. Others have reported differences in eigenvector centrality linked to stable versus
progressive mild cognitive impairment [92]. Eigenvector centrality in the lateral superior temporal
gyrus changes with age and may inform on cognitive decline in MCI [93]. Overall, there is
evidence from human neuroimaging studies supporting the observed sex differences in
eigenvector centrality and a potentially important link between this network measure and
amyloidosis.

In support of the observed effects of amyloidosis on eigenvector centrality in female mice,
we found that amyloidosis decreased a cluster of genes specifically in female mice. While the
amyloid and immune related genes were highly preserved in modular analyses, the gene
clusters reduced in APP-PS1 females was moderately preserved when comparing to male mice.
This female specific pattern of gene expression changes included Gng7, Syndig1l, Pde1b,
Lrrc10b, Vps26b as the top 5 downregulated genes in AβPP females, and Gad1 (glutamic acid
decarboxylase 1 gene encoding the main enzyme involved in producing GABA in neurons)
within the top 10 downregulated genes. *Gad1* is also known as Gad67 and is found widely
distributed in inhibitory interneuron cell soma [94]. AD associated alterations in GABAergic neuron
function and structure in hippocampal, temporal and parietal regions are well published [6, 8]. Aβ
is linked to hypersynchronous field potential activity in mice via a voltage gated sodium channel
– Nav1.1, mechanism in Gad67-expressing parvalbumin interneurons in parietal cortex of mice
[10]. An intriguing notion is that lightblue4 module genes are linked to functional dysregulation or
loss of GABAergic neurons in response to Aβ, specifically in female mice. *Gng7* reportedly
harbors differentially methylated positions associated with Braak stage pathology [95] and in mice

has been causally linked to ischemic stroke induced motor dysfunction and stressor-induced

anxiety [96] and striatal adenosine-A_{2A} receptor-containing medium spiny neurons' (MSNs) control

over locomotor responses to psychostimulants [97]. Interestingly, *Gng7* is one of the target genes

of microRNAs, miR-881-3p and miR-504, that are downregulated by alcohol treatment in female

but not male rats [98]. *Syndig1l* harbors epigenetic CpG-related single nucleotide polymorphisms

associated with AD in Hispanic populations [99]. *Pde1b*, encoding a phosphodiesterase 1, has

been linked to motor activity and spatial learning [100], identified in motor control in striatum of

transgenic mice of Huntington's [101] and involved in depression-like behaviors in response to

stress in mice [102]. Overexpression of *Pde1b* in dopamine D1 receptor-containing striatal MSNs

attenuated the locomotor response to cocaine in female mice but not males [103]. *Ppp1r1c* also

known as Darpp32, is found in dopamine receptor D1 expressing striatal medium spiny neurons

[104]. Darpp32 follows a sex-dependent pattern of expression [105, 106] and its expression has been

shown to be altered by estradiol [107]. These new data offer useful insight into potential avenues

to investigate sex-specific and hormonally regulated mechanisms of amyloid aggregation that

may differentially impact female brains compared to males.

Finally, we observed a higher number of extracellular plaques in female versus male mice.

This was observed in somatomotor cortex and basolateral amygdala. These findings are

consistent with previous research using the APP-PS1 mouse [108, 109], the Tg2576 [110] and the

triple transgenic AD mouse (3xTg-AD) [111]. The hippocampal concentrations of Aβ40 and Aβ42

and number of plaques at 12 and 17 mo is more than twice the amount in females compared to

male APP-PS1 [108]. The mechanism leading to higher plaque numbers in female versus male

mice may involve effects of estrogen loss on Aβ overproduction. Ovariectomy increases Aβ

levels in female Tg2576 mice (measured in whole brain homogenates by mass spectrometry),

and estrogen replacement reduces Aβ to sham Tg2576 control levels [112]. In that study,

estrogen did not appear to alter full length human AβPP, secreted AβPP nor the PS1 C-terminal

fragment [112]. A similar increase in Aβ load following OVX, and a reduction with estrogen

treatment, is observed in CA1, subiculum and frontal cortex of 3xTg-AD mice [113]. Estrogen

appears to repress BACE1 gene expression, and its activity, via estrogen receptor alpha [114], a

mechanism that may underlie the higher number of Aβ plaques in AβPP female versus male

mice [112]. Results for gene enrichment analysis further substantiated the observed sex

differences in amyloid expression, unexpectedly revealing a diametrically opposed App gene

levels compared to the histological data. App gene counts were higher in males compared to

female mice, whereas the opposite was found with Aβ plaque immunolabeling. This is

consistent with a reduction App gene expression in TG females as a result of putative
estrogenic repression of BACE1 [114].

To conclude, the present study adds to the literature on the functional network changes in
response to increased presence of extracellular plaques. Reducing extracellular plaque
pathology could be one way in which neuronal communication might be safeguarded in
individuals at risk of cognitive decline and Alzheimer disease. Moreover, the identified functional
connectome marker, specific to APP-PS1 females and consistent with the human literature, will
be interesting to further investigate in other AβPP strains. As indicated above, nuanced
mechanisms of Aβ aggregation may be important to understand in terms of how this impacts
broader network activity.

Acknowledgements

The authors would like to thank Marjory Pompilus for her help with portions of the work.
Marcelo Febo and Zachary Simon are thankful for the support provided by the McKnight Brain
Institute of the University of Florida.

Author contributions

Zachary Simon (Data curation; Visualization; Writing - Original Draft Preparation;
Conceptualization; Methodology; Formal Analysis; Investigation); Karen N. McFarland (Writing -
Review & Editing; Conceptualization; Methodology; Formal Analysis); Todd E. Golde (Writing –
Review & Editing; Conceptualization; Methodology); Paramita Chakrabarty (Writing - Review &
Editing; Conceptualization; Methodology; Formal Analysis); Marcelo Febo (Data curation;

Visualization; Writing – Original Draft Preparation; Conceptualization; Methodology; Formal
Analysis).

Statements and declarations

Ethical considerations

All procedures received prior approval from the Institutional Animal Care and Use Committee of
the University of Florida and follow all applicable NIH guidelines.

Consent for publication

All authors approve of the final version of this manuscript.

Declaration of conflicting interests

The authors declare no conflict of interest

Funding statement

This work was funded by NIA R21AG065819 with additional support provided by the McKnight
Brain Institute of the University of Florida. The neuroimaging components of this work were
performed in the National High Magnetic Field Laboratory's AMRIS Facility, which is funded by
National Science Foundation Cooperative Agreement No. DMR-1644779 and the State of
Florida. The original funding for the mouse brain RNA sequencing studies downloaded from AD
Knowledge Portal was provided by the NIH U01 AG046139 to Drs. Todd E Golde, Nilufer Taner
and Nathan Price.

Data availability

The TAUAPPms RNAseq datasets used in this manuscript has been previously deposited in AD
Knowledge Portal (https://adknowledgeportal.org) with the support of NIH/NIA (U01 AG046139
to T.E. Golde, N.D. Price, and N. Ertekin-Taner). The data can be accessed
at https://www.synapse.org/ (Synapse ID:syn2580853). The AD Knowledge Portal is a platform
for accessing data, analyses, and tools generated by the Accelerating Medicines Partnership
(AMP-AD) Target Discovery Program and other National Institute on Aging (NIA)-supported

programs to enable open-science practices and accelerate translational learning. All MRi

datasets are uploaded to openneuro.org.

750   **References**

1.   Alzheimer's Association. 2022 Alzheimer's disease facts and figures. *Alzheimers*
Dement 2022; 18: 700-789. 20220314. DOI: 10.1002/alz.12638.

2.   Hendriks S, Peetoom K, Bakker C, et al. Global Prevalence of Young-Onset Dementia:
A Systematic Review and Meta-analysis. *JAMA Neurol* 2021; 78: 1080-1090. DOI:
10.1001/jamaneurol.2021.2161.

3.   D'Argenio V and Sarnataro D. New Insights into the Molecular Bases of Familial
Alzheimer's Disease. *J Pers Med* 2020; 10 20200419. DOI: 10.3390/jpm10020026.

4.   Guerreiro RJ, Gustafson DR and Hardy J. The genetic architecture of Alzheimer's
disease: beyond APP, PSENs and APOE. *Neurobiol Aging* 2012; 33: 437-456. 20100701. DOI:
10.1016/j.neurobiolaging.2010.03.025.

5.   Mitew S, Kirkcaldie MT, Dickson TC, et al. Altered synapses and gliotransmission in
Alzheimer's disease and AD model mice. *Neurobiol Aging* 2013; 34: 2341-2351. 20130430.
DOI: 10.1016/j.neurobiolaging.2013.04.010.

6.   Fuhrer TE, Palpagama TH, Waldvogel HJ, et al. Impaired expression of GABA
transporters in the human Alzheimer's disease hippocampus, subiculum, entorhinal cortex and
superior temporal gyrus. *Neuroscience* 2017; 351: 108-118. 20170404. DOI:
10.1016/j.neuroscience.2017.03.041.

7.   Dominguez-Alvaro M, Montero-Crespo M, Blazquez-Llorca L, et al. Three-dimensional
analysis of synapses in the transentorhinal cortex of Alzheimer's disease patients. *Acta*
Neuropathol Commun 2018; 6: 20. 20180302. DOI: 10.1186/s40478-018-0520-6.

8.   Kurucu H, Colom-Cadena M, Davies C, et al. Inhibitory synapse loss and accumulation
of amyloid beta in inhibitory presynaptic terminals in Alzheimer's disease. *Eur J Neurol* 2022;
29: 1311-1323. 20210812. DOI: 10.1111/ene.15043.

9.   Garcia-Marin V, Blazquez-Llorca L, Rodriguez JR, et al. Diminished perisomatic
GABAergic terminals on cortical neurons adjacent to amyloid plaques. *Front Neuroanat* 2009; 3:
28. 20091120. DOI: 10.3389/neuro.05.028.2009.

10.   Verret L, Mann EO, Hang GB, et al. Inhibitory interneuron deficit links altered network
activity and cognitive dysfunction in Alzheimer model. *Cell* 2012; 149: 708-721. DOI:
10.1016/j.cell.2012.02.046.

11.   Niraula S, Yan SS and Subramanian J. Amyloid Pathology Impairs Experience-
Dependent Inhibitory Synaptic Plasticity. *J Neurosci* 2024; 44 20240131. DOI:
10.1523/JNEUROSCI.0702-23.2023.

12.   Buckner RL, Sepulcre J, Talukdar T, et al. Cortical hubs revealed by intrinsic functional
connectivity: mapping, assessment of stability, and relation to Alzheimer's disease. *J Neurosci*
2009; 29: 1860-1873. DOI: 10.1523/JNEUROSCI.5062-08.2009.

13.   Adriaanse SM, Binnewijzend MA, Ossenkoppele R, et al. Widespread disruption of
functional brain organization in early-onset Alzheimer's disease. *PLoS One* 2014; 9: e102995.
20140731. DOI: 10.1371/journal.pone.0102995.

14.   Canu E, Agosta F, Mandic-Stojmenovic G, et al. Multiparametric MRI to distinguish early
onset Alzheimer's disease and behavioural variant of frontotemporal dementia. *Neuroimage Clin*
2017; 15: 428-438. 20170525. DOI: 10.1016/j.nicl.2017.05.018.

15.   Filippi M, Basaia S, Canu E, et al. Brain network connectivity differs in early-onset
neurodegenerative dementia. *Neurology* 2017; 89: 1764-1772. 20170927. DOI:
10.1212/WNL.0000000000004577.

16.   Zhou J, Li K, Luo X, et al. Distinct impaired patterns of intrinsic functional network
centrality in patients with early- and late-onset Alzheimer's disease. *Brain Imaging Behav* 2021;
15: 2661-2670. 20210412. DOI: 10.1007/s11682-021-00470-3.

17.   Pini L, Geroldi C, Galluzzi S, et al. Age at onset reveals different functional connectivity
abnormalities in prodromal Alzheimer's disease. *Brain Imaging Behav* 2020; 14: 2594-2605.
DOI: 10.1007/s11682-019-00212-6.

18.   Cacace R, Sleegers K and Van Broeckhoven C. Molecular genetics of early-onset
Alzheimer's disease revisited. *Alzheimers Dement* 2016; 12: 733-748. 20160324. DOI:
10.1016/j.jalz.2016.01.012.

19.   Cochran JN, McKinley EC, Cochran M, et al. Genome sequencing for early-onset or
atypical dementia: high diagnostic yield and frequent observation of multiple contributory alleles.
Cold Spring Harb Mol Case Stud 2019; 5 20191213. DOI: 10.1101/mcs.a003491.

20.   Eissman JM, Dumitrescu L, Mahoney ER, et al. Sex differences in the genetic
architecture of cognitive resilience to Alzheimer's disease. *Brain* 2022; 145: 2541-2554. DOI:
10.1093/brain/awac177.

21.   Shah D, Jonckers E, Praet J, et al. Resting state FMRI reveals diminished functional
connectivity in a mouse model of amyloidosis. *PLoS One* 2013; 8: e84241. 20131217. DOI:
10.1371/journal.pone.0084241.

22.   Parent MJ, Zimmer ER, Shin M, et al. Multimodal Imaging in Rat Model Recapitulates
Alzheimer's Disease Biomarkers Abnormalities. *J Neurosci* 2017; 37: 12263-12271. 20171102.
DOI: 10.1523/JNEUROSCI.1346-17.2017.

23.   Grandjean J, Derungs R, Kulic L, et al. Complex interplay between brain function and
structure during cerebral amyloidosis in APP transgenic mouse strains revealed by multi-
parametric MRI comparison. *Neuroimage* 2016; 134: 1-11. DOI:
10.1016/j.neuroimage.2016.03.042.

24.   Grandjean J, Schroeter A, He P, et al. Early alterations in functional connectivity and
white matter structure in a transgenic mouse model of cerebral amyloidosis. *J Neurosci* 2014;
34: 13780-13789. 2014/10/10. DOI: 10.1523/JNEUROSCI.4762-13.2014.

25.   Jankowsky JL, Fadale DJ, Anderson J, et al. Mutant presenilins specifically elevate the
levels of the 42 residue beta-amyloid peptide in vivo: evidence for augmentation of a 42-specific
gamma secretase. *Hum Mol Genet* 2004; 13: 159-170. 20031125. DOI: 10.1093/hmg/ddh019.

26.   Janus C, Flores AY, Xu G, et al. Behavioral abnormalities in APPSwe/PS1dE9 mouse
model of AD-like pathology: comparative analysis across multiple behavioral domains.
Neurobiol Aging 2015; 36: 2519-2532. 20150521. DOI: 10.1016/j.neurobiolaging.2015.05.010.

27.   Febo M, Mahar R, Rodriguez NA, et al. Age-related differences in affective behaviors in
mice: possible role of prefrontal cortical-hippocampal functional connectivity and metabolomic
profiles. *Front Aging Neurosci* 2024; 16: 1356086. 20240308. DOI:
10.3389/fnagi.2024.1356086.

28.   Colon-Perez LM, Ibanez KR, Suarez M, et al. Neurite orientation dispersion and density
imaging reveals white matter and hippocampal microstructure changes produced by Interleukin-
6 in the TgCRND8 mouse model of amyloidosis. *Neuroimage* 2019; 202: 116138. DOI:
10.1016/j.neuroimage.2019.116138.

29.   Kotlarz P, Nino JC and Febo M. Connectomic analysis of Alzheimer's disease using
percolation theory. *Netw Neurosci* 2022; 6: 213-233. 20220201. DOI: 10.1162/netn_a_00221.

30.   Sakthivel R, Criado-Marrero M, Barroso D, et al. Fixed Time-Point Analysis Reveals
Repetitive Mild Traumatic Brain Injury Effects on Resting State Functional Magnetic Resonance
Imaging Connectivity and Neuro-Spatial Protein Profiles. *J Neurotrauma* 2023 2023/04/14. DOI:
10.1089/neu.2022.0464.

31.   Sadaka AH, Canuel J, Febo M, et al. Effects of inhaled cannabis high in Delta9-THC or
CBD on the aging brain: A translational MRI and behavioral study. *Front Aging Neurosci* 2023;
15: 1055433. 20230201. DOI: 10.3389/fnagi.2023.1055433.

32.   Cox RW. AFNI: software for analysis and visualization of functional magnetic resonance
neuroimages. *Comput Biomed Res* 1996; 29: 162-173.

33.   Jenkinson M, Bannister P, Brady M, et al. Improved optimization for the robust and
accurate linear registration and motion correction of brain images. *Neuroimage* 2002; 17: 825-
841.

34.   Klein A, Andersson J, Ardekani BA, et al. Evaluation of 14 nonlinear deformation
algorithms applied to human brain MRI registration. *Neuroimage* 2009; 46: 786-802. DOI:
10.1016/j.neuroimage.2008.12.037.

35.   Yushkevich PA, Piven J, Hazlett HC, et al. User-guided 3D active contour segmentation
of anatomical structures: significantly improved efficiency and reliability. *Neuroimage* 2006; 31:
1116-1128. DOI: 10.1016/j.neuroimage.2006.01.015.

36.   Wang Q, Ding SL, Li Y, et al. The Allen Mouse Brain Common Coordinate Framework: A
3D Reference Atlas. *Cell* 2020; 181: 936-953 e920. 20200507. DOI: 10.1016/j.cell.2020.04.007.

37.   Pompilus M, Colon-Perez LM, Grudny MM, et al. Contextual experience modifies
functional connectome indices of topological strength and efficiency. *Sci Rep* 2020; 10: 19843.
2020/11/18. DOI: 10.1038/s41598-020-76935-0.

38.   Yang Z, Zhu T, Pompilus M, et al. Compensatory functional connectome changes in a
rat model of traumatic brain injury. *Brain Commun* 2021; 3: fcab244. 2021/11/04. DOI:
10.1093/braincomms/fcab244.

39.   Xia M, Wang J and He Y. BrainNet Viewer: a network visualization tool for human brain
connectomics. *PLoS One* 2013; 8: e68910. 2013/07/19. DOI: 10.1371/journal.pone.0068910.

40.   Rubinov M and Sporns O. Complex network measures of brain connectivity: uses and
interpretations. *Neuroimage* 2010; 52: 1059-1069. DOI: 10.1016/j.neuroimage.2009.10.003.

41.   Wang Y, Ghumare E, Vandenberghe R, et al. Comparison of Different Generalizations of
Clustering Coefficient and Local Efficiency for Weighted Undirected Graphs. *Neural computation*
2017; 29: 313-331. 20161121. DOI: 10.1162/NECO_a_00914.

42.   Blondel VD, Guillaume J-P, Lambiotte R, et al. Fast unfolding of communities in large
networks. *Journal of Statistical Mechanics: Theory and Experiment* 2008; 2008: 2-12. DOI:
10.1088/1742-5468/2008/10/P10008.

43.   Newman ME. Assortative mixing in networks. *Physical review letters* 2002; 89: 208701.
DOI: 10.1103/PhysRevLett.89.208701.

44.   Morgan M, Pages H, Obenchain V, et al. Rsamtools: Binary alignment (BAM), FASTA,
variant call (BCF), and tabix file import. DOI: doi:10.18129/B9.bioc.Rsamtools.

45.   Lawrence M, Huber W, Pages H, et al. Software for computing and annotating genomic
ranges. *PLoS Comput Biol* 2013; 9: e1003118. 20130808. DOI: 10.1371/journal.pcbi.1003118.

46.   Love MI, Huber W and Anders S. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. *Genome Biol* 2014; 15: 550. DOI: 10.1186/s13059-014-0550-8.

883 47. Zhang Y, Chen K, Sloan SA, et al. An RNA-sequencing transcriptome and splicing
884 database of glia, neurons, and vascular cells of the cerebral cortex. *J Neurosci* 2014; 34: 11929-
885 11947. DOI: 10.1523/JNEUROSCI.1860-14.2014.

886 48. Langfelder P and Horvath S. WGCNA: an R package for weighted correlation network
887 analysis. *BMC Bioinformatics* 2008; 9: 559. 20081229. DOI: 10.1186/1471-2105-9-559.

888 49. Langfelder P and Horvath S. Fast R Functions for Robust Correlations and Hierarchical
889 Clustering. *J Stat Softw* 2012; 46.

890 50. Beckmann CF and Smith SM. Probabilistic independent component analysis for
891 functional magnetic resonance imaging. *IEEE Trans Med Imaging* 2004; 23: 137-152. DOI:
892 10.1109/TMI.2003.822821.

893 51. Hyvarinen A. Fast and robust fixed-point algorithms for independent component
894 analysis. *IEEE Trans Neural Netw* 1999; 10: 626-634. DOI: 10.1109/72.761722.

895 52. Bukhari Q, Schroeter A, Cole DM, et al. Resting State fMRI in Mice Reveals Anesthesia
896 Specific Signatures of Brain Functional Networks and Their Interactions. *Front Neural Circuits*
897 2017; 11: 5. 20170203. DOI: 10.3389/fncir.2017.00005.

898 53. Jonckers E, Van Audekerke J, De Visscher G, et al. Functional connectivity fMRI of the
899 rodent brain: comparison of functional connectivity networks in rat and mouse. *PLoS One* 2011;
900 6: e18876. 20110418. DOI: 10.1371/journal.pone.0018876.

901 54. Sforazzini F, Schwarz AJ, Galbusera A, et al. Distributed BOLD and CBV-weighted
902 resting-state networks in the mouse brain. *Neuroimage* 2014; 87: 403-415. DOI:
903 10.1016/j.neuroimage.2013.09.050.

904 55. McFarland KN, Ceballos C, Rosario A, et al. Microglia show differential transcriptomic
905 response to Abeta peptide aggregates ex vivo and in vivo. *Life Sci Alliance* 2021; 4 20210614.
906 DOI: 10.26508/lsa.202101108.

907 56. Busche MA, Wegmann S, Dujardin S, et al. Tau impairs neural circuits, dominating
908 amyloid-beta effects, in Alzheimer models in vivo. *Nat Neurosci* 2019; 22: 57-64. 20181217.
909 DOI: 10.1038/s41593-018-0289-8.

910 57. Calvo-Flores Guzman B, Kim S, Chawdhary B, et al. Amyloid-Beta(1-42) -Induced
911 Increase in GABAergic Tonic Conductance in Mouse Hippocampal CA1 Pyramidal Cells.
912 *Molecules* 2020; 25 20200206. DOI: 10.3390/molecules25030693.

913 58. Lauterborn JC, Scaduto P, Cox CD, et al. Increased excitatory to inhibitory synaptic ratio
914 in parietal cortex samples from individuals with Alzheimer's disease. *Nat Commun* 2021; 12:
915 2603. 20210510. DOI: 10.1038/s41467-021-22742-8.

916 59. Petrache AL, Rajulawalla A, Shi A, et al. Aberrant Excitatory-Inhibitory Synaptic
917 Mechanisms in Entorhinal Cortex Microcircuits During the Pathogenesis of Alzheimer's Disease.
918 *Cereb Cortex* 2019; 29: 1834-1850. DOI: 10.1093/cercor/bhz016.

60.   Shah D, Deleye S, Verhoye M, et al. Resting-state functional MRI and [18F]-FDG PET
demonstrate differences in neuronal activity between commonly used mouse strains.
Neuroimage 2016; 125: 571-577. 20151028. DOI: 10.1016/j.neuroimage.2015.10.073.

61.   Grandjean J, Schroeter A, Batata I, et al. Optimization of anesthesia protocol for resting-
state fMRI in mice based on differential effects of anesthetics on functional connectivity
patterns. *Neuroimage* 2014; 102 Pt 2: 838-847. 20140828. DOI:
10.1016/j.neuroimage.2014.08.043.

62.   Whitesell JD, Buckley AR, Knox JE, et al. Whole brain imaging reveals distinct spatial
patterns of amyloid beta deposition in three mouse models of Alzheimer's disease. *J Comp*
Neurol 2019; 527: 2122-2145. 20181204. DOI: 10.1002/cne.24555.

63.   Arszovszki A, Borhegyi Z and Klausberger T. Three axonal projection routes of individual
pyramidal cells in the ventral CA1 hippocampus. *Front Neuroanat* 2014; 8: 53. 20140625. DOI:
10.3389/fnana.2014.00053.

64.   Orsini CA, Kim JH, Knapska E, et al. Hippocampal and prefrontal projections to the
basal amygdala mediate contextual regulation of fear after extinction. *J Neurosci* 2011; 31:
17269-17277. DOI: 10.1523/JNEUROSCI.4095-11.2011.

65.   Ishikawa A and Nakamura S. Ventral hippocampal neurons project axons
simultaneously to the medial prefrontal cortex and amygdala in the rat. *J Neurophysiol* 2006; 96:
2134-2138. 20060712. DOI: 10.1152/jn.00069.2006.

66.   Swanson LW and Kohler C. Anatomical evidence for direct projections from the
entorhinal area to the entire cortical mantle in the rat. *J Neurosci* 1986; 6: 3010-3023. DOI:
10.1523/JNEUROSCI.06-10-03010.1986.

67.   Thierry AM, Gioanni Y, Degenetais E, et al. Hippocampo-prefrontal cortex pathway:
anatomical and electrophysiological characteristics. *Hippocampus* 2000; 10: 411-419. DOI:
10.1002/1098-1063(2000)10:4<411::AID-HIPO7>3.0.CO;2-A.

68.   Bhembre N, Bonthron C and Opazo P. Synaptic Compensatory Plasticity in Alzheimer's
Disease. *J Neurosci* 2023; 43: 6833-6840. DOI: 10.1523/JNEUROSCI.0379-23.2023.

69.   Liu D, Lu H, Stein E, et al. Brain regional synchronous activity predicts tauopathy in
3xTgAD mice. *Neurobiol Aging* 2018; 70: 160-169. 20180621. DOI:
10.1016/j.neurobiolaging.2018.06.016.

70.   Giorgio J, Adams JN, Maass A, et al. Amyloid induced hyperexcitability in default mode
network drives medial temporal hyperactivity and early tau accumulation. *Neuron* 2024; 112:
676-686 e674. 20231213. DOI: 10.1016/j.neuron.2023.11.014.

71.   Hu W, Pan D, Wang Y, et al. PET Imaging for Dynamically Monitoring
Neuroinflammation in APP/PS1 Mouse Model Using [(18)F]DPA714. *Front Neurosci* 2020; 14:
810. 20200929. DOI: 10.3389/fnins.2020.00810.

72.   Garland EF, Dennett O, Lau LC, et al. The mitochondrial protein TSPO in Alzheimer's
disease: relation to the severity of AD pathology and the neuroinflammatory environment. *J*
Neuroinflammation 2023; 20: 186. 20230814. DOI: 10.1186/s12974-023-02869-9.

73. Giulian D, Li J, Bartel S, et al. Cell surface morphology identifies microglia as a distinct
class of mononuclear phagocyte. *J Neurosci* 1995; 15: 7712-7726. DOI:
10.1523/JNEUROSCI.15-11-07712.1995.

74. Wake H, Moorhouse AJ, Jinno S, et al. Resting microglia directly monitor the functional
state of synapses in vivo and determine the fate of ischemic terminals. *J Neurosci* 2009; 29:
3974-3980. DOI: 10.1523/JNEUROSCI.4363-08.2009.

75. Zhan Y, Paolicelli RC, Sforazzini F, et al. Deficient neuron-microglia signaling results in
impaired functional brain connectivity and social behavior. *Nat Neurosci* 2014; 17: 400-406.
20140202. DOI: 10.1038/nn.3641.

76. Tremblay ME, Lowery RL and Majewska AK. Microglial interactions with synapses are
modulated by visual experience. *PLoS Biol* 2010; 8: e1000527. 20101102. DOI:
10.1371/journal.pbio.1000527.

77. Tremblay ME and Majewska AK. A role for microglia in synaptic plasticity? *Commun*
Integr Biol 2011; 4: 220-222. DOI: 10.4161/cib.4.2.14506.

78. Bolos M, Perea JR, Terreros-Roncal J, et al. Absence of microglial CX3CR1 impairs the
synaptic integration of adult-born hippocampal granule neurons. *Brain Behav Immun* 2018; 68:
76-89. 20171007. DOI: 10.1016/j.bbi.2017.10.002.

79. Lituma PJ, Woo E, O'Hara BF, et al. Altered synaptic connectivity and brain function in
mice lacking microglial adapter protein Iba1. *Proc Natl Acad Sci U S A* 2021; 118. DOI:
10.1073/pnas.2115539118.

80. Surala M, Soso-Zdravkovic L, Munro D, et al. Lifelong absence of microglia alters
hippocampal glutamatergic networks but not synapse and spine density. *EMBO Rep* 2024
20240408. DOI: 10.1038/s44319-024-00130-9.

81. Lalonde R, Kim HD, Maxwell JA, et al. Exploratory activity and spatial learning in 12-
month-old APP(695)SWE/co+PS1/DeltaE9 mice with amyloid plaques. *Neurosci Lett* 2005; 390:
87-92. DOI: 10.1016/j.neulet.2005.08.028.

82. Kilgore M, Miller CA, Fass DM, et al. Inhibitors of class 1 histone deacetylases reverse
contextual memory deficits in a mouse model of Alzheimer's disease.
Neuropsychopharmacology 2010; 35: 870-880. 20091209. DOI: 10.1038/npp.2009.197.

83. Hedden T, Van Dijk KR, Becker JA, et al. Disruption of functional connectivity in clinically
normal older adults harboring amyloid burden. *J Neurosci* 2009; 29: 12686-12694. DOI:
10.1523/JNEUROSCI.3189-09.2009.

84. Lim HK, Nebes R, Snitz B, et al. Regional amyloid burden and intrinsic connectivity
networks in cognitively normal elderly subjects. *Brain* 2014; 137: 3327-3338. 20140929. DOI:
10.1093/brain/awu271.

85. Melnikova T, Park D, Becker L, et al. Sex-related dimorphism in dentate gyrus atrophy
and behavioral phenotypes in an inducible tTa:APPsi transgenic model of Alzheimer's disease.
Neurobiol Dis 2016; 96: 171-185. 20160826. DOI: 10.1016/j.nbd.2016.08.009.

86. Mortberg MA, Zhao HT, Reidenbach AG, et al. Regional variability and genotypic and
pharmacodynamic effects on PrP concentration in the CNS. *JCI Insight* 2022; 7 20220322. DOI:
10.1172/jci.insight.156532.

87. Xu J, Burgoyne PS and Arnold AP. Sex differences in sex chromosome gene expression
in mouse brain. *Hum Mol Genet* 2002; 11: 1409-1419. DOI: 10.1093/hmg/11.12.1409.

88. Zuo XN, Ehmke R, Mennes M, et al. Network centrality in the human functional
connectome. *Cereb Cortex* 2012; 22: 1862-1875. 20111002. DOI: 10.1093/cercor/bhr269.

89. Lorenzini L, Ingala S, Collij LE, et al. Eigenvector centrality dynamics are related to
Alzheimer's disease pathological changes in non-demented individuals. *Brain Commun* 2023; 5:
fcad088. 20230328. DOI: 10.1093/braincomms/fcad088.

90. Ingala S, Tomassen J, Collij LE, et al. Amyloid-driven disruption of default mode network
connectivity in cognitively healthy individuals. *Brain Commun* 2021; 3: fcab201. 20210906. DOI:
10.1093/braincomms/fcab201.

91. Skouras S, Falcon C, Tucholka A, et al. Mechanisms of functional compensation,
delineated by eigenvector centrality mapping, across the pathophysiological continuum of
Alzheimer's disease. *Neuroimage Clin* 2019; 22: 101777. 20190312. DOI:
10.1016/j.nicl.2019.101777.

92. Qiu T, Luo X, Shen Z, et al. Disrupted Brain Network in Progressive Mild Cognitive
Impairment Measured by Eigenvector Centrality Mapping is Linked to Cognition and
Cerebrospinal Fluid Biomarkers. *J Alzheimers Dis* 2016; 54: 1483-1493. DOI: 10.3233/JAD-
160403.

93. Zhang X, Zeng Q, Wang Y, et al. Alteration of functional connectivity network in
population of objectively-defined subtle cognitive decline. *Brain Commun* 2024; 6: fcae033.
20240209. DOI: 10.1093/braincomms/fcae033.

94. Stone DJ, Walsh J and Benes FM. Localization of cells preferentially expressing
GAD(67) with negligible GAD(65) transcripts in the rat hippocampus. A double in situ
hybridization study. *Brain research Molecular brain research* 1999; 71: 201-209. DOI:
10.1016/s0169-328x(99)00185-0.

95. Li QS, Sun Y and Wang T. Epigenome-wide association study of Alzheimer's disease
replicates 22 differentially methylated positions and 30 differentially methylated regions. *Clin
Epigenetics* 2020; 12: 149. 20201017. DOI: 10.1186/s13148-020-00944-z.

96. Halder SK, Sasaki K and Ueda H. Ggamma7-specific prothymosin alpha deletion causes
stress- and age-dependent motor dysfunction and anxiety. *Biochem Biophys Res Commun*
2020; 522: 264-269. 20191120. DOI: 10.1016/j.bbrc.2019.11.103.

97. Brunori G, Pelletier OB, Stauffer AM, et al. Selective Manipulation of G-Protein
gamma(7) Subunit in Mice Provides New Insights into Striatal Control of Motor Behavior. *J
Neurosci* 2021; 41: 9065-9081. 20210920. DOI: 10.1523/JNEUROSCI.1211-21.2021.

98. Choi MR, Han JS, Jin YB, et al. Differential expression of microRNAs in the hippocampi
of male and female rodents after chronic alcohol administration. *Biol Sex Differ* 2020; 11: 65.
20201123. DOI: 10.1186/s13293-020-00342-3.

99. Ma Y, Reyes-Dumeyer D, Piriz A, et al. Multi-omics Characterization of Epigenetic and
Genetic Risk of Alzheimer Disease in Autopsied Brains from two Ethnic Groups. *medRxiv* 2024
20240214. DOI: 10.1101/2024.02.12.24302533.

100. Reed TM, Repaske DR, Snyder GL, et al. Phosphodiesterase 1B knock-out mice exhibit
exaggerated locomotor hyperactivity and DARPP-32 phosphorylation in response to dopamine
agonists and display impaired spatial learning. *J Neurosci* 2002; 22: 5188-5197. DOI:
10.1523/JNEUROSCI.22-12-05188.2002.

101. Hebb AL, Robertson HA and Denovan-Wright EM. Striatal phosphodiesterase mRNA
and protein levels are reduced in Huntington's disease transgenic mice prior to the onset of
motor symptoms. *Neuroscience* 2004; 123: 967-981. DOI: 10.1016/j.neuroscience.2003.11.009.

102. Hufgard JR, Williams MT and Vorhees CV. Phosphodiesterase-1b deletion confers
depression-like behavioral resistance separate from stress-related effects in mice. *Genes, brain,*
and behavior 2017; 16: 756-767. 20170607. DOI: 10.1111/gbb.12391.

103. Teague CD, Markovic T, Zhou X, et al. Circuit-wide gene network analysis reveals sex-
specific roles for phosphodiesterase 1b in cocaine addiction. *J Neurosci* 2024 20240418. DOI:
10.1523/JNEUROSCI.1327-23.2024.

104. Ouimet CC and Greengard P. Distribution of DARPP-32 in the basal ganglia: an electron
microscopic study. *J Neurocytol* 1990; 19: 39-52. DOI: 10.1007/BF01188438.

105. Zhou L, Nazarian A, Sun WL, et al. Basal and cocaine-induced sex differences in the
DARPP-32-mediated signaling pathway. *Psychopharmacology (Berl)* 2009; 203: 175-183.
20081105. DOI: 10.1007/s00213-008-1388-0.

106. Weiner J, Sun WL, Zhou L, et al. PKA-mediated responses in females' estrous cycle
affect cocaine-induced responses in dopamine-mediated intracellular cascades. *Neuroscience*
2009; 161: 865-876. 20090405. DOI: 10.1016/j.neuroscience.2009.03.071.

107. Stroppolo A, Tian C, Guinea B, et al. 17beta-Estradiol promotes striatal medium size
spiny neuronal maturation in vitro. *Neuroendocrinology* 2004; 79: 259-267. 20040624. DOI:
10.1159/000079320.

108. Wang J, Tanila H, Puolivali J, et al. Gender differences in the amount and deposition of
amyloidbeta in APPswe and PS1 double transgenic mice. *Neurobiol Dis* 2003; 14: 318-327.
DOI: 10.1016/j.nbd.2003.08.009.

109. Wu J, Fu B, Lei H, et al. Gender differences of peripheral plasma and liver metabolic
profiling in APP/PS1 transgenic AD mice. *Neuroscience* 2016; 332: 160-169. 20160705. DOI:
10.1016/j.neuroscience.2016.06.049.

110. Lee JY, Cole TB, Palmiter RD, et al. Contribution by synaptic zinc to the gender-
disparate plaque formation in human Swedish mutant APP transgenic mice. *Proc Natl Acad Sci*
U S A 2002; 99: 7705-7710. DOI: 10.1073/pnas.092034699.

111.   Hirata-Fukae C, Li HF, Hoe HS, et al. Females exhibit more extensive amyloid, but not
tau, pathology in an Alzheimer transgenic model. *Brain Res* 2008; 1216: 92-103. 20080410.
DOI: 10.1016/j.brainres.2008.03.079.

112.   Zheng H, Xu H, Uljon SN, et al. Modulation of A(beta) peptides by estrogen in mouse
models. *J Neurochem* 2002; 80: 191-196. DOI: 10.1046/j.0022-3042.2001.00690.x.

113.   Carroll JC, Rosario ER, Chang L, et al. Progesterone and estrogen regulate Alzheimer-
like neuropathology in female 3xTg-AD mice. *J Neurosci* 2007; 27: 13357-13365. DOI:
10.1523/JNEUROSCI.2718-07.2007.

114.   Cui J, Ait-Ghezala G, Sambamurti K, et al. Sex-Specific Regulation of beta-Secretase: A
Novel Estrogen Response Element (ERE)-Dependent Mechanism in Alzheimer's Disease. *J*
Neurosci 2022; 42: 1154-1165. 20211213. DOI: 10.1523/JNEUROSCI.0864-20.2021.

1083

1084

1085

1086

1087

1088   **Figures and figure captions**

1089

1090

1091

1092

1093

1094

1095

1096

1097

1098

1099

1100

1101

1102

1103

1104

Figure 1. Female APP-
PS1 mice have a greater number of amyloid plaques in basolateral amygdala and somatomotor
neocortical areas compared to male APP-PS1 mice. a) Confirmation of amyloid plaque
presence in TG mice using antibody 33.1.1. b) Comparison of amyloid plaque burden between
male (M) and female (F) TG mice. c) Representative histological sections from female and male
NTG and TG mouse basolateral amygdala (BLA), prefrontal cortex (PFC), somatomotor cortex
(SMC), and hippocampus (HPC). Data in a-b are box-whisker showing median, minimum-
maximum values and individual data points. Asterisks are *p < 0.05, **p < 0.01.

1113

1114

Figure 2. Amyloidosis increased functional connectivity within an unclassified component that included nodes in posterior dorsal striatum and basolateral amygdala in both male and female mice. a) Undistinct component measured in a cohort of 28 NTG and TG fMRI scans. Higher intensity/higher functional connectivity voxels in yellow are encircled and indicated by the corresponding brain region. b) Statistical results of full factorial ANOVA assessing genotype x sex interactions and main effects ($p < 0.05$, corrected). Letter codes indicate the brain area according to the Allen Mouse Brain Common Coordinate Reference atlas (CCFv3). Brain area names are matched to letter codes at the bottom of the figure. Significant voxels (main effect genotype) are highlighted in yellow-orange hue indicating TG > NTG.

Figure 3. Sex specific effects of amyloid on functional network 'hub' topology. a) Eigenvector centrality scores for several prefrontal cortical and striatal nodes calculated for a 10% network density. Data in box-whisker plot are median with minimum and maximum values and show individual data points. Top row plots show ROI data for left (L) hemisphere and bottom row plot the right (R) hemisphere. b) Area under the curve (AUC) values calculated for eigenvector centrality across multiple edge densities (2-40%) per ROI. Data in plot is presented as mean standard ± error. Left plot shows ROI data for left (L) hemisphere and right plot the right (R) hemisphere. c) Connectome maps represent mean AUC values for eigenvector centrality per node and edge strength (Pearson correlations) as spherical nodes and lines connecting nodes. Node eigenvector centrality is reflected in sphere size and color intensity. Edges are thresholded at 6% density for visual purposes. Asterisks indicate significantly difference between NTG vs TG females* and NTG females vs males**. Abbreviations: ACG1, anterior cingulate gyrus; PRL, prelimbic cortex; IL, infralimbic cortex; LOC, lateral orbital cortex; VOC, ventral orbital cortex; DMS, dorsomedial striatum.

Figure 4. Diverging patterns in differentially expressed genes (DEGs) in female and male APP-PS1 mice. a) Volcano plot showing significant downregulated (blue numbers) and upregulated (red numbers) DEGs as a function of APP-PS1 genotype in female mice. b) Volcano plot of DEGs in male mice. c) Comparison of gene expression signatures for cell type composition groupings between NTG and TG female (F) and male (M) mice. Data are shown as box-whisker with median, minimum-maximum values and individual data points. Asterisks are *p < 0.05, **p < 0.01, *** p < 0.001.

Figure 5. Weighted gene correlation network analysis identified two distinct modules in female

APP-PS1 brain with significant positively and negatively correlated groups of immune and

neuronal genes. a) Heatmap showing modules with gene networks that are positively (red) and

negatively (blue) correlated with genotype (modules with p-value less than 0.1 are indicated by

color). Modules with nonsignificant correlations are in grey. b) Gene significance by within

module degree (kWithin) for the brown module (evaluated for genotype). Listed are the top 10

most significant genes with hub gene colored red. c) Gene significance by within module degree

(kWithin) for the lightblue4 module (evaluated for genotype). Listed are the top 10 most

significant genes with hub gene colored blue. d) Hierarchical clustering of gene expression in

brown module. e) Hierarchical clustering of gene expression in lightblue4 module.

1195

Figure 6. High module preservation of DEGs involved in immune responses (brown module) and moderate preservation of DEGs involved in neuronal function and structure (lightblue4 module). Female module DEGs used as reference to test male DEGs. A) Module rank. Modules close to 0 are highly preserved. B) Zsummary plot. Moderate preservation Z = 2-10 and high preservation Z > 10. d) Comparison of normalized gene expressions for the top 5 most significant genes in the brown module. App gene included as a comparison. e) Comparison of normalized gene expressions for the top 5 most significant genes in the brown module. Ddx3x gene with sex dependent expression levels is included as a comparison. Data are shown as box-whisker with median, minimum-maximum values and individual data points. Asterisks are *p < 0.05, **p < 0.01, *** p < 0.001, ****p < 0.0001.