E. coli FtsN coordinates synthesis and degradation of septal peptidoglycan by partitioning between a synthesis track and a denuded glycan track

Zhixin Lyu[1,3], Xinxing Yang[1,4], Atsushi Yahashiri[2], Stephen Ha[1], Joshua W. McCausland[1,5], Xinlei Chen[1], Brooke M. Britton[1], David S. Weiss[2*], Jie Xiao[1*]

[1]Department of Biophysics and Biophysical Chemistry, Johns Hopkins School of Medicine, Baltimore, MD 21205, USA.

[2]Department of Microbiology and Immunology, University of Iowa Carver College of Medicine, Iowa City, IA 52242, USA.

[3]Current address: Genetics and Biochemistry Branch, National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health, Bethesda, MD 20892, USA.

[4]Current address: The Chinese Academy of Sciences Key Laboratory of Innate Immunity and Chronic Disease, School of Basic Medical Sciences, Division of Life Sciences and Medicine, University of Science and Technology of China, Hefei, China.

[5]Current address: Department of Biology, Stanford University, Stanford, CA 94305 USA.

*Corresponding authors: david-weiss@uiowa.edu, xiao@jhmi.edu

Abstract

The *E. coli* cell division protein FtsN was proposed to coordinate septal peptidoglycan (sPG) synthesis and degradation to ensure robust cell wall constriction without lethal lesions. Although the precise mechanism remains unclear, previous work highlights the importance of two FtsN domains: the E domain, which interacts with and activates the sPG synthesis complex FtsWIQLB, and the SPOR domain, which binds to denuded glycan (dnG) strands, key intermediates in sPG degradation. Here, we used single-molecule tracking of FtsN and FtsW (a proxy for the sPG synthesis complex FtsWIQLB) to investigate how FtsN coordinates the two opposing processes. We observed dynamic behaviors indicating that FtsN's SPOR domain binds to dnGs cooperatively, which both sequesters the sPG synthesis complex on dnG (termed as the dnG-track) and protects dnGs from degradation by lytic transglycosylases (LTs). The release of the SPOR domain from dnGs leads to activating the sPG synthesis complex on the sPG-track and simultaneously exposing those same dnGs to degradation. Furthermore, FtsN's SPOR domain self-interacts and facilitates the formation of a multimeric sPG synthesis complex on both tracks. The cooperative self-interaction of the SPOR domain creates a sensitive switch to regulate the partitioning of FtsN between the dnG- and sPG-tracks, thereby controlling the balance between sequestered and active populations of the sPG synthesis complex. As such, FtsN coordinates sPG synthesis and degradation in space and time.

Introduction

Gram-negative (diderm) bacteria such as *Escherichia coli* have a rigid cell wall composed of a thin peptidoglycan (PG) layer. The PG is made of repeated N-acetylglucosamine (GlcNAc)-N-acetylmuramic acid (MurNAc) units cross-linked via short peptide bridges[1]. This thin PG layer is constantly remodeled by a large collection of cell wall synthases and hydrolases to allow cell growth and division. In *E. coli*, constriction requires an essential septal PG (sPG) synthase complex formed by the glycosyltransferase FtsW and the transpeptidase FtsI, along with more than ten nonessential PG hydrolases, including three periplasmic amidases (Ami) and eight known and putative lytic transglycosylases (LTs)[2, 3]. Amidases remove peptide crosslinks by cleaving the peptide moiety from MurNAc, which generates denuded glycan (dnG) strands[4, 5]. dnGs are subsequently destroyed by LTs, which cleave the β-1,4-glycosidic bond between MurNAc and GlcNAc[2, 6, 7]. The degradation of PG must be precisely regulated and synchronized with new sPG synthesis in both space and time to ensure robust cell wall constriction without causing lethal lesions. However, the mechanisms underlying this spatiotemporal coordination remain unclear.

FtsN is an essential bitopic membrane protein in *E. coli* and other γ-proteobacteria. It
possesses four functional domains with distinct roles in cell division: (1) a short, N-terminal
cytoplasmic domain (FtsN^{Cyto}) that directly interacts with the early division protein FtsA to trigger
the activation of FtsW[8-10]; (2) an α-helical transmembrane domain (FtsN^{TM}) that anchors FtsN to
the inner membrane[11]; (3) a short, loosely-formed helical essential domain (FtsN^E) in the
periplasm that binds to FtsI and its scaffolding protein FtsL to activate the sPG synthesis complex
formed by FtsWI and FtsQLB[12-15]; and (4) a periplasmic, highly conserved C-terminal SPOR
domain (FtsN^{SPOR}) that binds to dnG[12, 14, 16-19]. A notable feature of FtsN is its long, largely
disordered linker between FtsN^{TM} and FtsN^{SPOR}, which also includes FtsN^E. When fully extended,
this linker can reach ~70 nm in length[14].

In previous single-molecule tracking (SMT) studies in *E. coli*, we observed that FtsW, FtsI,
and FtsB (and hence most likely the full sPG synthesis complex formed by FtsWIQLB) exhibit two
distinct directional moving populations. One population moves fast at ~30 nm s^{-1}, is inactive in
sPG synthesis, and is driven by FtsZ's treadmilling dynamics through a Brownian ratchet
mechanism[15, 20-22]. We term this fast-moving population as on the Z-track. The other population
moves slowly at ~8-9 nm s^{-1}, is driven by active sPG synthesis, and is independent of FtsZ's
treadmilling dynamics[15, 20, 21]. We term this slow-moving population as on the sPG-track.
Additionally, we observed stationary populations of FtsW, FtsI and FtsB, which we identified as
either trapped in the middle of FtsZ polymers or "poised" for synthesis. Thus, SMT provides a
powerful means to link the moving dynamics of cell wall synthesis enzymes and regulators to their
activities and functions in live bacterial cells.

We found that modulating a cell's sPG synthesis activity or FtsZ's treadmilling speed can shift
the partitioning of the synthesis complex between the two tracks, revealing a mechanism to
coordinate sPG synthesis spatiotemporally with septal pole morphogenesis[15, 21, 23]. Such a two-
track behavior was also observed in the diderm bacterium *C. crescentus*[24]. However, in Gram-
positive (monoderm) bacteria such as *S. pneumoniae*[25], *B. subtilis*[26], and *S. aureus*[27], only the
slow-moving, sPG synthesis-driven population of synthases was observed. This difference likely
suggests a less stringent regulation requirement for septation and cell pole morphogenesis due
to the thicker cell walls in these bacteria, and/or a transitory Z-track-dependent behavior of these
enzymes[28].

Recently, we discovered that FtsN exhibits a single, slow-moving population on the sPG-
track and is a required component of the active processive synthesis complex[21, 23]. Structural
modeling shows that FtsN's E domain binds to FtsL and FtsI in the previously identified Activation
of FtsW and FtsI (AWI) domain[29] and Constriction Control Domain (CCD)[29-31], triggering a

conformational change of FtsI's Anchor domain to open up the catalytic pore of FtsW[15]. We also
discovered that the moving dynamics of FtsN depend on its domain composition. For example,
the isolated E domain in the absence of the rest of FtsN exhibits both Z-track and sPG-track
populations and is sufficient to maintain the processivity of the synthesis complex. The
cytoplasmic tail and the transmembrane domain of FtsN (FtsN$^{Cyto-TM}$) exhibits only the Z-track
dependent, fast-moving population[23].

Besides the sPG-track-coupled slow-moving population, we observed that more than half of
septal FtsN molecules are stationary because they are anchored to dnG by the SPOR domain[23].
While the essential E domain is the only region of FtsN strictly required for viability[9, 12], the SPOR
domain improves FtsN's function by concentrating FtsN at the septum through its binding to dnG,
which increases the rate of constriction and sPG synthesis[12, 32, 33]. FtsN also regulates sPG
degradation by recruiting PG amidases that create dnG and mediate daughter cell separation[34].
Conversely, overproduction of FtsN's SPOR domain impairs turnover of dnG by LTs[35]. Thus, the
coexistence within one single protein of a SPOR domain that binds to dnG (a transient
intermediate of sPG degradation) along with an E domain that activates the sPG synthesis
complex, suggests that FtsN could link sPG synthesis and degradation in space and time[19, 36].

In this work, we built on these ideas and dissected the coordination mechanism using SMT
of FtsN and FtsW across various genetic backgrounds and growth conditions. Our studies identify
a third track, termed as the dnG-track, characterized by the stationary population of FtsN bound
to dnG via its SPOR domain. We discovered that FtsN's SPOR domain self-interacts, binds to
dnG cooperatively, and facilitates the formation of a multimeric sPG synthesis complex
FtsWIQLB-FtsN on both the dnG- and sPG-tracks. Our results support a model in which FtsN,
through the cooperative self-interaction of its SPOR domain, sensitively partitions between the
dnG- and sPG-tracks. This partitioning regulates the balance between sequestered and active
sPG synthesis complex, thereby coordinating sPG synthesis and degradation activities both in
space and time.

Results

Altered dnG levels in ∆*amiABC* and ∆6*LT*s cells diminish sPG synthesis activity

To investigate the effect of SPOR-dnG binding on sPG synthesis activities, we monitored the
moving dynamics of FtsN and FtsW using two mutant strains with altered dnG levels. In the
∆*amiABC* strain (strain EC5345, Supplementary Table 1), the dnG level is effectively zero due to
the lack of all three periplasmic amidases[12, 19], which generate dnG[7]. In the ∆6*LT*s strain (strain

JL435, Supplementary Table 1), the dnG level is ~20-40 fold higher than that in the wild type (WT)

background[19]. Cells likely have long stretches of dnG strands due to the deletion of six out of eight

known and putative LTs ($\Delta mltA$, $\Delta mltC$, $\Delta mltD$, $\Delta mltE$, $\Delta slt70$, $\Delta rlpA$), which degrade dnG[6]. Both

the $\Delta amiABC$ and $\Delta 6LT$s mutants exhibited sick chaining phenotypes due to impaired ability to

degrade sPG for proper daughter cell separation[6] (Fig. 1a, b).

To perform SMT of FtsN and FtsW, we ectopically expressed a functional, previously

characterized FtsN-Halo[SW 23] or FtsW-TagRFP-t (abbreviated as FtsW-RFP)[21] fusion protein from

a plasmid in the two strain backgrounds (strains EC5345 and EC3708 respectively,

Supplementary Table 1). As expected, and in accordance with altered dnG levels in the two

strains, septal localization of FtsN-Halo[SW] was reduced in $\Delta amiABC$ cells but enhanced in $\Delta 6LT$s

cells (Fig. 1a, b, top; Supplementary Fig. 1a). FtsW-RFP showed similar localization levels (Fig.

1a, b, bottom; Supplementary Fig. 1b).

SMT of FtsN-Halo[SW] and FtsW-RFP revealed that in $\Delta amiABC$ cells the two fusion proteins

behaved similarly. The active, slow-moving populations on the sPG-track for both fusions were

almost completely abolished (Fig. 1c, red curves). A new, fast-moving population of FtsN-Halo[SW]

appeared ($P_{fm} = 41 \pm 2\%$, $n = 77$ segments for FtsN-Halo[SW]), which was similar to that of FtsW-

RFP ($P_{fm} = 38 \pm 3\%$, $n = 72$ segments, Fig. 1c-e, blue curves and bars, Supplementary Table 4).

The average lifetime of stationary FtsN-Halo[SW] molecules decreased to $T_{stat} = 9.4 \pm 0.6$ s ($P_{stat} =$

$59 \pm 2\%$, $n = 109$ segments, Fig. 1f, Supplementary Table 4) from $T_{stat} = 27 \pm 1$ s we previously

measured under the WT condition[23], comparable to that of stationary FtsW-RFP molecules (T_{stat}

$= 9.1 \pm 0.6$ s, $n = 116$ segments, Fig. 1f, Supplementary Table 4). This shortened stationary

lifetime is similar to the average lifetime of stationary FtsZ molecules (8-12 s) measured in *E. coli*

and other species[23, 25, 37-39]. As there is essentially no dnG in $\Delta amiABC$ cells, these results suggest

that stationary FtsN-Halo[SW] and FtsW-RFP molecules are stuck in the middle of treadmilling FtsZ

polymers, as previously observed both *in vivo* and *in vitro*[21, 23, 40].

In $\Delta 6LT$s cells, FtsN-Halo[SW] and FtsW-RFP also behaved similarly to each other. The majority

of the fusion protein molecules were stationary ($P_{stat} = 91 \pm 3\%$, $n = 546$ segments for FtsN-Halo[SW]

and $P_{stat} = 88 \pm 3\%$, $n = 252$ segments for FtsW-RFP, Fig. 1d, e, black curves and gray bars,

Supplementary Table 4), consistent with the expectation that an elevated dnG level increases the

number of dnG-bound FtsN-Halo[SW] molecules and consequently FtsN-bound FtsW-RFP

molecules. Most strikingly, the stationary lifetime of FtsN-Halo[SW], which reflects the time it takes

for single FtsN-Halo[SW] molecules to dissociate from dnG, increased to $T_{stat} = 59 \pm 2$ s in $\Delta 6LT$s

cells (Fig. 1f, Supplementary Table 4). As we describe further below, the marked increase in both

the percentage and dissociation time of stationary FtsN-Halo[SW] molecules in $\Delta 6LT$s cells is most

consistent with a higher SPOR-dnG affinity arising from cooperative binding of FtsN to long

stretches of dnG. Similar phenomena have been observed on the length-dependent binding of

oligomeric DNA- or mRNA-binding proteins[41-43]. The stationary lifetime of FtsW-RFP molecules

also increased to $T_{stat} = 24 \pm 1$ s (Fig. 1f, Supplementary Table 4), up from ~18 s in the WT

condition we measured previously[21], suggesting FtsW may be bound cooperatively to multiple

molecules of dnG-anchored FtsN (also see below).

The lack of slow-moving population of FtsN-Halo[SW] and FtsW-RFP in the two strains indicates

the lack of active sPG synthesis[21, 23]. To verify this result, we used HADA (HCC-Amino-D-alanine

hydrochloride) labeling to measure the septal cell wall synthesis activity[22]. Both mutants showed

significantly lower HADA incorporation levels than WT cells, which were indistinguishable from

sPG synthesis-inhibited cells treated with cephalexin (Fig. 1g, h). The lack of septal cell wall

synthesis activity in $\Delta amiABC$ cells is consistent with the minimal septal localization of FtsN in the

absence of dnG (Fig.1a). However, in $\Delta 6LT$s cells where septal FtsN and FtsW are abundant, the

lack of HADA incorporation implies that the FtsWIQLB-FtsN complex is sequestered on dnG and

needs to be released from dnG to engage in sPG synthesis.

Truncation of FtsN's SPOR domain leads to the sequestering of FtsW on the Z-

track

To examine the role of FtsN-dnG interaction without modulating dnG levels, we constructed an

$ftsN^{\Delta SPOR}$ strain in which the *spor* sequence is deleted from the full-length *ftsN* gene at its native

chromosomal locus (strain EC3338, Supplementary Table 1). Cells expressing the $ftsN^{\Delta SPOR}$

truncation gene grew slowly and were filamentous, but expression of an additional copy of

$ftsN^{\Delta SPOR}$ from a plasmid restored WT-like growth and similar septum synthesis activity (strain

JL337, termed as $ftsN^{\Delta SPOR+}$ herein for simplicity, Supplementary Table 1, Supplementary Fig. 2).

These results are consistent with the fact that SPOR is the major septum localization determinant

of FtsN, and hence a higher expression level of FtsN$^{\Delta SPOR}$ is needed to support cell division at a

similar level[12].

Next, we performed SMT of FtsW-RFP expressed from a plasmid in the $ftsN^{\Delta SPOR+}$

background (strain JL362, Supplementary Table 1) using the same M9 minimal medium and

imaging conditions as before[23]. We observed that the sPG synthesis processivity of FtsW, defined

as the persistent running time on the sPG-track, remained similar in the absence of SPOR ($T_{sm} =$

21.1 ± 0.5 s, $\mu \pm$ s.e.m. in $ftsN^{\Delta SPOR+}$ *vs.* 20.1 ± 0.1 s in WT, Fig. 2b, top, Supplementary Fig. 3,

Supplementary Table 5). The Z-track processivity of FtsW, similarly defined as the persistent

running time on the Z-track, remained unchanged as well ($T_{fm} = 9.4 \pm 0.2$ s, $\mu \pm$ s.e.m. in $ftsN^{\Delta SPOR+}$

v.s. 8.3 ± 0.3 s in WT, Fig. 2b, bottom, Supplementary Table 5). However, the active, slow-moving

population of FtsW-RFP decreased significantly (from 33 ± 2%, n = 157 segments in WT to 8 ±

3%, n = 38 segments in *ftsN*$^{\Delta SPOR+}$), while the fast-moving population increased from 12 ± 2% to

19 ± 2% (Fig. 2a, c, Supplementary Table 5). Additionally, the stationary population of FtsW-RFP

increased to 73.5 ± 2% (n = 353 segments, Fig. 2c, Supplementary Table 5) from ~55% under

the WT condition as we previously reported[21] and had a short lifetime of T_{stat} = 10.4 ± 0.4 s

(Supplementary Table 5). This stationary lifetime is comparable to that of FtsW-RFP in Δ*amiABC*

cells (Supplementary Table 4), suggesting that they were similarly stuck in the middle of FtsZ

polymers. Taken together, these results suggest that in the absence of FtsN's SPOR domain,

while FtsW's processivities on the sPG- and Z-tracks are largely unaltered, FtsW has a

significantly reduced active, slow-moving population on the sPG-track and is primarily

sequestered on the Z-track.

Truncation of FtsN's SPOR domain leads to the sequestering of FtsN$^{\Delta SPOR}$ on the

Z-track and reduces its sPG-track processivity

To examine how truncation of the SPOR domain affects the moving dynamics of FtsN itself, we

constructed a chromosomal integrated FtsN$^{\Delta SPOR}$-Halo^{SW} fusion protein and overexpressed it in

an *ftsN*-depletion background (strain EC5257, Supplementary Table 1) to restore normal cell

length. We observed that the stationary population of FtsN$^{\Delta SPOR}$-Halo^{SW} at septa decreased to 40%

± 2% with an average T_{stat} = 9.2 ± 0.6 s (n = 85 segments, Fig. 2c, d, Supplementary Table 5),

suggesting that they were trapped in the middle of FtsZ polymers. Most importantly, in contrast to

the single slow-moving population observed for FtsN-Halo^{SW}, FtsN$^{\Delta SPOR}$-Halo^{SW} exhibited two

distinct processively-moving populations, one slow (P_{sm} = 13 ± 3%, V_{sm} = 9.4 ± 0.1 nm s^{-1}, μ ±

s.e.m., n = 28 segments) and one fast (P_{fm} = 46 ± 2%, V_{fm} = 32.7 ± 0.1 nm s^{-1}, μ ± s.e.m., n = 98

segments, Fig. 2c, d, Supplementary Table 5). Inhibiting sPG synthesis activity using aztreonam

completely abolished the slow-moving population of FtsN$^{\Delta SPOR}$-Halo^{SW} but preserved the fast-

moving population (Fig. 2e, Supplementary Table 5), similar to what we previously observed for

FtsW in aztreonam-treated cells[21] or FtsB in fosfomycin-treated cells[15]. However, the persistent

running times, or the processivities, of FtsN$^{\Delta SPOR}$-Halo^{SW} on both the sPG-track and the Z-track

(T_{sm} = 7.5 ± 0.1 s and T_{fm} = 5.5 ± 0.1 s respectively, Fig. 2b, Supplementary Table 5) were shorter

than those of FtsW-RFP in the *ftsN*$^{\Delta SPOR+}$ background and full-length FtsN-Halo^{SW} on the sPG-

track (14.5 ± 0.7 s, Fig. 2b, Supplementary Table 5). These results suggest that similar to FtsW

in the absence of FtsN's SPOR domain, septal FtsN$^{\Delta SPOR}$-Halo^{SW} was mainly sequestered on the

Z-track. Because dnG-bound FtsN molecules are stationary and not running on the sPG-track[23],

the reduced persistent running time of FtsN$^{\Delta SPOR}$-Halo^{SW} on the sPG-track suggests that the

SPOR domain may have a role in supporting the processivity of FtsN in the active sPG synthesis

complex, independent of its dnG-binding ability.

An FtsN^{Q251E} mutant defective in dnG-binding partially restores FtsN's sPG-track

population and processivity

Truncation of FtsN's SPOR domain removes both its dnG-binding ability and potentially other

interactions of the SPOR domain independent of dnG binding. To further investigate the effects

of these two types of interactions, we monitored the moving dynamics of an FtsN^{Q251E}-Halo^{SW}

fusion protein by transforming a plasmid expressing *ftsN^{Q251E}-halo^{SW}* into an *ftsN*-depletion strain

(strain JL378, Supplementary Table 1). Residue Q251 is the most highly conserved amino acid

in the dnG-binding cleft and interacts directly with the glycan chain[16]. The Q251E substitution

reduces septal localization and dnG-binding of the isolated SPOR domain but has not been

studied in the context of full-length FtsN[44].

Consistent with a defect in dnG-binding, FtsN^{Q251E}-Halo^{SW} exhibited a fast-moving population

on the Z-track ($P_{fm} = 33 \pm 2\%$, $T_{fm} = 6.4 \pm 0.2$ s, $n = 64$ segments, Fig. 2b, f, Supplementary Table

5) and a smaller, $42 \pm 2\%$ stationary population with a significantly reduced stationary lifetime of

$T_{stat} = 13.1 \pm 0.9$ s ($n = 81$ segments, Fig. 2c, Supplementary Table 5). These results further

confirm that the dnG-binding ability of SPOR prevents the release of FtsN to the Z-track.

Interestingly, the slow-moving population percentage and processivity of FtsN^{Q251E}-Halo^{SW}

recovered to $P_{sm} = 25 \pm 2\%$ and $T_{sm} = 9.7 \pm 0.2$ s ($n = 49$ segments), in between that of FtsN$^{\Delta SPOR}$-

Halo^{SW} ($P_{sm} = 13\%$ and $T_{sm} = 7.5$ s) and full-length FtsN-Halo^{SW} ($P_{sm} = 45\%$ and $T_{sm} = 14.5$ s)

(Fig. 2b, c, Supplementary Tables 4 and 5). These results suggest that both the dnG-binding

ability and the dnG-binding-independent interaction of SPOR contribute to maintaining the active

sPG synthesis complex on the sPG-track (also see **Discussion**).

Isolated SPOR domain (iSPOR) moves processively with WT FtsN

To further investigate the role of the SPOR domain in supporting FtsN's slow-moving population

on the sPG-track independent of FtsN's other domains, we used a previously characterized fusion

construct FtsN^{WYAA}-Halo^{SW} [23]. This fusion protein contains two critical mutations, W83A and Y85A,

in FtsN's E domain, which abolish the binding of the E domain to FtsI and FtsL and the subsequent

activation of FtsWI[9, 45]. We previously reported that FtsN^{WYAA}-Halo^{SW} molecules were completely

stationary at septa in the superfission variant $ftsB^{E56A}$ $\Delta ftsN$ background[23], despite that FtsW-RFP

still moved processively on the sPG-track in the same strain background[21].

Surprisingly, when we ectopically expressed this FtsN^{WYAA}-Halo^{SW} fusion protein in the

presence of WT FtsN (Fig. 3a, strain EC5259, Supplementary Table 1), it exhibited essentially

the same percentage of the slow-moving population compared to WT FtsN-Halo^{SW} (P_{sm} = 44 ±

4%, n = 79 segments, Fig. 3b, c, Supplementary Table 6) with a modestly reduced processivity

(T_{sm} = 10.7 ± 0.8 s, Fig. 3d, top, Supplementary Table 6). This result suggests that FtsN^{WYAA}-

Halo^{SW} remains part of the processively-moving sPG synthesis complex through the binding of

domains other than the E domain to WT FtsN in the complex, as the direct binding of its E domain

with FtsI and FtsL is abolished by the WYAA mutations.

Our previous studies showed that the cytoplasmic and transmembrane domains of FtsN are

not involved in the formation of the processive sPG complex[23]. Therefore, we reasoned that the

SPOR domain might be responsible for such interaction in the absence of the E domain-mediated

binding to FtsI and FtsL. We constructed a Halo-iSPOR fusion protein, in which a Halo-tag is

fused to the N-terminus of the isolated SPOR domain (hereafter termed as iSPOR to distinguish

from the SPOR domain linked to the rest of FtsN), and expressed it in the presence of WT FtsN

(Fig. 3e, strain EC5275, Supplementary Table 1). Strikingly, while Halo-iSPOR localized to the

septum as expected (Supplementary Fig. 4), these Halo-iSPOR molecules moved directionally

(P_{sm} = 42 ± 3%, v = 9.4 ± 0.5 nm/s, n = 61 segments, Fig. 3c, f, Supplementary Table 6) with a

processivity of T_{sm} = 9.8 ± 0.7 s (Fig. 3d, top, Supplementary Table 6), and became predominately

stationary in cells treated with aztreonam (Supplementary Fig. 5), mimicking what we observed

for full-length FtsN-Halo^{SW} previously[23]. These results strongly suggest that there exists SPOR-

SPOR self-interaction.

iSPOR^{Q251E} defective in dnG-binding moves processively with WT FtsN

An alternative explanation for the observed processive movement of iSPOR with WT FtsN is that

iSPOR and FtsN's SPOR domain may bind to the same dnG fragment disintegrated from the cell

wall without direct protein-protein interactions. To examine this possibility, we constructed an

isolated Halo-iSPOR^{Q251E} fusion, which is defective in dnG-binding, and expressed it in the WT

ftsN background (Fig. 3g, strain JL431, Supplementary Table 1). Septal localization of Halo-

iSPOR^{Q251E} decreased about 2-fold compared to unmutated Halo-iSPOR (Supplementary Fig. 4b).

Surprisingly, Halo-iSPOR^{Q251E} still exhibited a large slow-moving population with a similar

processivity compared to Halo-iSPOR (P_{sm} = 36 ± 4%, T_{sm} = 9.6 ± 0.6 s, n = 33 segments, Fig.

3c, d, h, Supplementary Table 6). The apparent ~2-fold decrease in dnG-binding with little or no

change in the dynamics of the slow-moving population argues the observed iSPOR-FtsN
interaction is independent of dnG-binding. Note that the moving dynamics of Halo-iSPOR^{Q251E} in
the presence of WT FtsN mimics that of WT FtsN but not that of FtsN^{Q251E}-Halo^{SW}. The latter
exhibited a fast-moving population (compare Fig. 2f and Fig. 3h) because it has an N-terminal
cytoplasmic domain that can bind to FtsA, whereas Halo-iSPOR^{Q251E} does not. This comparison
further confirms that there must exist SPOR-SPOR self-interaction in the processive sPG
synthesis complex.

iSPOR becomes stationary when disconnected from the rest of FtsN
To further verify that iSPOR binds to the existing SPOR domain of full-length FtsN in the
processive sPG synthesis complex, we expressed Halo-iSPOR *in trans* with FtsN$^{\Delta SPOR+}$ (Fig. 3i,
strain EC5415, which also contains an extra plasmid copy of *ftsN$^{\Delta SPOR}$*, Supplementary Table 1).
Halo-iSPOR localized to the septum in the *ftsN$^{\Delta SPOR+}$* background similarly as in the WT *ftsN*
background (Supplementary Fig. 4). We reasoned that if the movement of iSPOR were due to its
binding to FtsN domains other than SPOR, we should still observe its processive movement with
FtsN$^{\Delta SPOR+}$. However, Halo-iSPOR molecules became completely stationary (Fig. 3c, j), with a
stationary lifetime of T_{stat} = 29 $\pm$ 1 s (n = 602 segments, Supplementary Table 6) that is
indistinguishable from full-length FtsN-Halo^{SW} (Fig. 1f). Treating cells with aztreonam did not
change the percentage nor the lifetime of these stationary Halo-iSPOR molecules (Fig. 3c, d,
Supplementary Fig. 5). Because we showed that in the same *ftsN$^{\Delta SPOR+}$* background, FtsW-RFP
and FtsN$^{\Delta SPOR}$-Halo^{SW} exhibited both slow- and fast-moving populations (Fig. 2), these results
strongly suggest that SPOR-SPOR self-interaction and the linkage between SPOR and the rest
of FtsN are responsible for the observed processive movement of iSPOR. As such, these results
demonstrate that the active, processive FtsWIQLB-FtsN complex contains multiple FtsN
molecules mediated by SPOR-SPOR self-interaction, and that this interaction is important to
sustain the active population of sPG synthesis complex on the sPG-track.
We also note that the stationary lifetimes (~15-20 s) of the FtsN^{WYAA}-Halo^{SW}, Halo-iSPOR
and Halo-iSPOR^{Q251E} in the presence of WT FtsN were shorter than that of full-length FtsN-Halo^{SW}
(~30 s, Fig. 3d, bottom, Supplementary Table 6), suggesting that FtsN^{WYAA}-Halo^{SW} and iSPOR
domains may not compete effectively with WT FtsN for dnG-binding. Interestingly, in ensemble
time-lapse movies of cells treated with aztreonam, FtsN-Halo^{SW} signal also persisted at the
septum longer than either Halo-iSPOR or FtsN$^{\Delta SPOR}$-Halo^{SW} (Supplementary Fig. 6,
Supplementary Movies 1-3). The longer persistence of full-length FtsN-Halo^{SW} compared to Halo-

iSPOR or FtsN$^{\Delta SPOR}$-Halo^{SW} further suggest that the SPOR domain must be linked to the rest of
FtsN in _cis_ in the same molecule for the full functionality of FtsN.

Stationary and processive sPG synthesis complexes are non-monomeric

In our single-molecule tracking experiments, we used sparse Halo-dye (JF646) labeling to ensure
the tracking of single molecules of interest. We also screened out fluorescent spots with intensities
higher than that of single molecules. In light of the above results, we reexamined some of our
data to assess whether we could directly visualize the presence of more than one molecule in the
same stationary and processive sPG synthesis complex. Indeed, we identified FtsN-Halo^{SW}
molecules in both moving and stationary SMT trajectories where the fluorescent spot intensities
are significantly higher than expected from a single JF646 dye molecule, which also exhibited the
typical quantized photobleaching step or fluorescence jump for single molecules (Fig. 3k;
Supplementary Movies 4-5). We even observed one FtsN-Halo^{SW} molecule that repeatedly
associated and dissociated until it stably associated with a processively moving complex (Fig. 3k,
left; Supplementary Movie 4). Most importantly, we identified similar trajectories showing the
presence of more than one single molecule from our previous SMT of Halo-FtsB[15], FtsI-Halo^{SW} [20]
and FtsW-RFP[21] fusions in _E. coli_ and Halo-FtsW in _C. crescentus_[24] (Supplementary Fig. 7;
Supplementary Movies 6-8). Thus, while the sparse labeling prevents us from further statistically
analyzing the number of molecules in the complex, these observations demonstrate that the
FtsWIQLB-FtsN complex in _E. coli_ exists as a multimer in the form of (FtsWIQLB)$_m$-FtsN$_n$ or
(FtsWIQLB-FtsN)$_n$. The non-monomeric nature of the sPG synthesis complex may be a
conserved feature across different bacterial species, as a recent SMT study in _B. subtilis_ also
reported the presence of multimeric PBP2b, the equivalent of FtsI in _E. coli_, on the sPG-track[26].

Relative sPG synthesis and degradation activities are correlated in WT _ftsN_
background but skewed when SPOR is disconnected

Under normal growth conditions in _E. coli_, sPG synthesis and degradation activities are balanced
to avoid aberrant cell pole morphogenesis or septal cell wall lesions. FtsN has been proposed as
a key player in coordinating these two opposing activities by linking sPG synthesis to the
abundance and location of dnGs[19, 36]. It is known that the binding of FtsN's E domain to the
FtsWIQLB complex activates sPG synthesis activity[15, 23, 46-48]. We also further confirmed that
overexpressing iSPOR in WT cells caused a profound chaining phenotype (Supplementary Fig.
8), suggesting that the binding of FtsN's SPOR domain to dnG protects them from being degraded
by LTs, as also suggested by a previous study[35]. Therefore, we propose that the partitioning of

369 FtsN between the two populations, one slow-moving in complex with active FtsWIQLB and the
370 other stationary bound to dnG, would naturally balance sPG synthesis and degradation activities:
371 release of FtsN and FtsN-bound FtsWIQLB complex from dnG leads to processive sPG synthesis
372 while simultaneously exposing dnGs to degradation by LTs, enhancing both synthesis and
373 degradation; conversely, binding of FtsN to dnG protects dnG and also sequesters FtsN-bound
374 FtsWIQLB complex, diminishing both synthesis and degradation.

375  To analyze the proposed correlation quantitatively, we compared FtsN's occupancies on the
376 processive sPG-track and dnGs (Fig. 4). We calculated FtsN's occupancy on the sPG-track (N_{sPG})
377 using the product of the slow-moving population percentage of FtsN-Halo^{SW} and the
378 corresponding persistent running time and velocity under each condition, normalized to that of
379 the WT condition (**Methods**). Note that FtsN's occupancy on the sPG-track is highly correlated
380 with that calculated using FtsW-RFP data (Supplementary Fig. 9). Additionally, we previously
381 showed that FtsW's occupancy on the processive sPG-track correlated highly to the cell wall
382 constriction rate and HADA incorporation level[21]. Therefore, N_{sPG} can be used as a proxy for sPG
383 synthesis activity.

384  We calculated FtsN's occupancy on dnG (N_{dnG}) using the product of the percentage of the
385 stationary population of FtsN-Halo^{SW} and its corresponding stationary lifetime, normalized to that
386 of the WT condition (**Methods**). As N_{dnG} reflects the level of dnG protected by FtsN, the inverse
387 of N_{dnG} ($1/N_{dnG}$) should reflect LTs' sPG degradation activity (high N_{dnG} indicates low LT activities
388 and low N_{dnG} indicates high LT activities).

389  In Fig. 4a, we plotted $1/N_{dnG}$ (proxy for LTs' sPG degradation activity) against N_{sPG} (proxy for
390 sPG synthesis activity) and observed that $1/N_{dnG}$ is highly correlated with N_{sPG} across a variety of
391 conditions ($p_{pearson}$ = 0.9412). For example, in cells harboring a superfission variant FtsI^{R167S} and
392 grown in rich medium, both N_{sPG} and $1/N_{dnG}$ were high, indicating high activities in both sPG
393 synthesis and degradation (Fig. 4a, conditions j, k). In cells treated with cell wall synthesis
394 inhibitors (MTSES for *ftsW*^{302C}, antibiotics aztreonam or fosfomycin) or deleted of six LTs ($\Delta 6LT$s),
395 both N_{sPG} and $1/N_{dnG}$ were low, indicating low activities in both sPG synthesis and degradation
396 (Fig. 4a, conditions a-d). A modified FtsN variant that lacks the cytoplasmic and transmembrane
397 domains (Halo-FtsN$^{\Delta Cyto-TM}$) also largely followed the same trend (Fig. 4a, condition e), indicating
398 that the correlation is unrelated to the cytoplasmic interactions of FtsN with FtsA.

399  Most strikingly, we found that on average, the ratio between $1/N_{dnG}$ and N_{sPG}, the proxy for
400 the ratio between relative sPG degradation and synthesis activities, was close to one across all
401 conditions that contained a functional, full periplasmic copy of FtsN (Fig. 4b, black bars),
402 irrespective of whether sPG synthesis is enhanced (as in superfission backgrounds, conditions i-

k) or inhibited (as in cells treated with antibiotics, conditions a-d). However, this ratio became

significantly higher (Fig. 4a, b, red) at ~3 when the E domain's binding to FtsWIQLB was disabled

by the WYAA mutations (condition n), or in cells expressing Halo-iSPOR in *trans* with *ftsN*[ΔSPOR+]

(condition m), and further increased to ~5 when cells were treated with aztreonam (condition l).

These results hence suggest that FtsN's partitioning on the sPG- and dnG-tracks maintains

balanced sPG synthesis and degradation activities, and that the linking of the SPOR domain to

the rest of FtsN is required for this role.

Discussion

In *E. coli*, sPG synthesis and turnover must be meticulously coordinated to prevent lysis and

ensure synchronous invagination of the three layers of the envelope. The essential bitopic

membrane protein FtsN was suggested to coordinate sPG synthesis and turnover because its

periplasmic E domain activates the sPG synthesis complex FtsWIQLB and its C-terminal SPOR

domain binds to dnG, a transient intermediate in biogenesis of the division septum. Here we

investigated this role using SMT to monitor the dynamics of septal FtsN and FtsW under various

conditions. Our findings yield several significant insights and lead to a model explaining how FtsN

coordinates sPG synthesis and degradation in space and time, which we discuss below.

A dnG-track anchors stationary FtsWIQLB-FtsN complexes

We previously reported the existence of stationary FtsW-RFP and FtsI-Halo[SW] molecules with an

average stationary lifetime of ~18-20 s, which reflects the contributions of two populations of

FtsWI: one with a short stationary lifetime (~9 s) bound to internal positions in FtsZ polymers, and

another with a longer stationary lifetime whose location was yet unknown[21, 23]. These longer-lived

stationary synthase molecules were inferred as "poised" for sPG synthesis, as their abundance

decreased under conditions stimulating sPG synthesis, such as overproduction of UppS to

provide more Lipid II, and increased under conditions inhibiting sPG synthesis, such as treating

cells with aztreonam or fosfomycin[21, 23]. Our current findings suggest that these "poised" FtsWI

molecules are bound to dnG-anchored FtsN. This suggestion is supported by the disappearance

of the long-lived FtsW-RFP stationary population in Δ*amiABC* cells, which lack dnG (average

stationary lifetime reduced to 9 s, Fig. 1f, Table S4). Conversely, the average stationary lifetime

of FtsW-RFP increased to ~24 s in Δ6*LT*s mutant (Fig. 1f, Table S4), where dnG-anchored FtsN

accumulates to high levels.

We refer to poised sPG synthases bound to dnG-anchored FtsN as being on the "dnG-track".

Along with the previously identified Z-track and sPG-track[15, 21, 23], the dnG-track brings the total

number of notional tracks in the *E. coli* septum to three. What are their purposes? We previously

showed that the dynamic redistribution of FtsWIQLB between the Z-track and sPG-track facilitates

the even distribution of enzymes along the septum, ensuring smooth and symmetric septum

growth[21]. In contrast, the dnG-track likely functions to synchronize sPG synthesis and degradation

both in space and time (see below).

FtsN ensures balanced sPG synthesis and degradation activities

Our results show that under most conditions where FtsN is intact, the normalized activities of sPG

synthesis and degradation are proportional, i.e., balanced. However, this balance is disrupted in

cells where the SPOR domain (iSPOR) is disconnected from the rest of the FtsN (Fig. 4a, b, red).

These findings align with a previous study using fluorescent D-amino acids (FDAAs) to compare

septal cell wall synthesis and degradation rates in WT and an *ftsN*$^{\Delta SPOR}$ mutant[32]. That study found

a clear decrease in sPG synthesis and a marginal increase in sPG degradation in *ftsN*$^{\Delta SPOR}$ cells,

which was not considered significant but nevertheless trended in the expected direction of

increased degradation/synthesis ratio if FtsN's SPOR domain protects dnG from LT activity. The

modest effect of *ftsN*$^{\Delta SPOR}$ on sPG degradation suggests other SPOR proteins may help protect

dnG in the absence of FtsN's SPOR domain. Interestingly, using FDAAs to assay sPG synthesis

and turnover revealed a lower degradation/synthesis ratio in a superfission *ftsL** (*ftsL*^{E88K})

background. This strain still contained a functional full-length FtsN. It would be interesting to

examine whether FtsN's binding to this mutant FtsL is increased, as the mutation FtsL^{E88K} is next

to FtsL^{E87} that is involved in the binding with FtsN's E domain[15].

FtsN coordinates sPG synthesis and degradation both in space and time

Our findings support a model in which FtsN not only coordinates sPG synthesis and degradation

at the activity level (Fig. 4a, b), but also in space and time (Fig. 4c). Specifically, FtsN binds to

dnG generated by amidases, protecting them from degradation by LTs. The sPG synthesis

complex FtsWIQLB becomes anchored to dnG-bound FtsN via its E domain binding to FtsI and

FtsL. When FtsN's SPOR domain dissociates from dnG and when sPG precursors are available,

the FtsWIQLB-FtsN complex initiates processive sPG synthesis in the vicinity of dnG.

Simultaneously, the dissociation of FtsN's SPOR domain from dnG allows LTs to destroy recently-

vacated binding sites for FtsN. Similarly, when the FtsWIQLB-FtsN complex moves processively

to synthesize new sPG, the long, flexible linker of FtsN connecting the SPOR and E domains may

allow SPOR to probe for newly generated dnGs nearby. Once found, SPOR binds to these new

dnGs, potentially bringing over the entire FtsWIQLB-FtsN complex, which will then commence

another round of sPG synthesis nearby when released again. This iterative cycling of sPG

synthesis and degradation ensures that these two activities are closely coordinated both in space

and time.

Self-interaction of SPOR

Our study reveals that FtsN's SPOR domain self-interacts both when bound to dnG and when

bound to the active FtsWIQLB complex. The following lines of evidence support this observation.

First, FtsN is expressed at a level about 3-4-fold higher than FtsW, FtsI, and FtsQLB[23, 49], making

it possible to form at least (FtsWIQLB-FtsN)$_n$ type of complexes. Second, our previous Structured

Illumination Microscopy (SIM) imaging experiments showed large FtsN clusters that either

remained stationary or moved processively together[23]. Third, in our current SMT experiments, we

directly observed more than one FtsN molecule in the stationary or slow-moving trajectories within

our typical imaging resolution limit of ~30-40 nm[23] (Fig. 3k). Fourth, FtsN^{WYAA} moved processively

in the presence of full-length WT FtsN (Fig. 3a), but became completely stationary when *ftsN* was

deleted in the superfission *ftsB*^{E56A} background[23]. Similarly, iSPOR moved processively in the

presence of full-length WT FtsN (Fig. 3e, f), but not in the presence of FtsN$^{\Delta SPOR+}$ (Fig. 3i, j). Finally,

iSPOR^{Q251E}, which is defective in dnG binding, still exhibited similar percentages of both slow-

moving and stationary populations (Fig. 3g, h, Supplementary Table 4). Interestingly, the septal

localization percentage of iSPOR^{Q251E} was half that of Halo-iSPOR, yet the percentage was similar

in the presence and absence of full-length FtsN (Supplementary Fig. 4). This independence

suggests that other SPOR proteins (of which *E. coli* has three[12, 50]) may substitute for FtsN's

SPOR domain. Taken together, these observations strongly support the presence of SPOR-

SPOR self-interaction independent of dnG-binding.

We have tried to characterize the self-interaction of iSPOR *in vitro* but were largely

unsuccessful. Purified iSPOR showed a single monomeric species in size exclusion columns and

static light scattering (Supplementary Fig. 11, Supplementary Table 7). Crosslinking experiments

to trap transient interactions revealed oligomers at high iSPOR concentrations, but it cannot be

excluded that the crosslinked oligomers arose from random collisions (Supplementary Fig. 12).

The published NMR structure of FtsN's periplasmic domain appears to be monomeric[14]. Therefore,

in vitro support for the SPOR-SPOR interaction we observed *in vivo* is lacking. One possible

explanation is that the interaction requires one of the partners contain other domains of FtsN.

Indeed, the Loose group reported that full-length FtsN *in vitro* forms a dimer in sedimentation

experiments[40], and the Vollmer group captured oligomeric FtsN molecules containing the

periplasmic E and SPOR domains using blue native gel electrophoresis[51].

Cooperative binding of SPOR to dnG

Several findings support the idea that FtsN's SPOR domain binds cooperatively to dnG. First, we
observed that not only the stationary population percentages, but also the stationary lifetimes of
single FtsN molecules, changed in response to a variety of altered conditions (Figs.1-3,
Supplementary Fig. 13). The stationary lifetime, which reflects the time it takes for a single
molecule to exit a particular state, directly measures the apparent dnG-dissociation time constant
of FtsN. Changes in these lifetimes are therefore consistent with altered binding affinities between
dnG and SPOR, in addition to shifted binding equilibria where only the bound and unbound
population percentages change.

Second, although we could not directly quantify dnG level or length under different conditions,
FtsN's binding affinity responded to expected dnG levels in cells (Supplementary Fig. 13). For
example, the stationary lifetime of FtsN increased to ~60 s in the $\Delta 6LT$s mutant, where the dnG
level is ~20-40 fold higher than in WT cells (Fig. 1f, Supplementary Fig. 13)[19]. In the $ftsN^{\Delta SPOR+}$
background, iSPOR had a stationary lifetime similar to that of full-length FtsN (~30 s), but in the
presence of WT FtsN, iSPOR's stationary lifetime decreased to ~20 s and further to ~14 s when
cells were treated with aztreonam (Fig. 3d, bottom). In contrast, full-length FtsN's stationary
lifetime increased to ~40 s in aztreonam-treated cells[23]. The decreased binding affinity of iSPOR
to dnG in the presence of WT FtsN can be attributed to the expected low levels of dnG, as most
dnG is occupied by full-length FtsN. The few-fold change in the binding affinities of SPOR to dnG,
together with the *in vivo* evidences for SPOR-SPOR self-interaction, is most consistent with a
cooperative binding mechanism, which has been frequently observed in DNA/mRNA-length-
dependent cooperative binding of oligomeric proteins[41-43]. Notably, the Vollmer group previously
demonstrated that the purified periplasmic domain of FtsN binds exclusively to dnG strands longer
than 25 NAM-NAG units, consistent with a cooperative binding mechanism[18].

Third, we considered alternative explanations for changes in the stationary lifetime of FtsN,
specifically the possibility that these changes result from active "pulling" of SPOR by processive
sPG synthases or "pushing" of SPOR by processive LTs, rather than cooperative binding of SPOR
to dnG. We reasoned that active "pulling" by processive sPG synthases is unlikely because
iSPOR (with FtsN$^{\Delta SPOR+}$), which lacks an E domain to bind to the sPG synthesis complex, had the
same stationary lifetime as full-length WT FtsN (Fig. 3d, bottom). Similarly, active "pushing" of
SPOR away from dnG by processive LTs is also unlikely since overexpression of iSPOR led to a
strong chaining phenotype (Supplementary Fig. 8), suggesting that LTs are unable to "push"
iSPOR off dnG.

Finally, all stationary lifetime distributions of FtsN under these conditions exhibited non-
exponential distributions (Supplementary Fig. 10), indicating that the underlying process cannot
be described as a single rate-limiting step. Many of these distributions showed a rise and decay,
a signature indicative of multiple, consecutive steps[52], which is consistent with a cooperative, non-
monomeric process. These non-monotonic distributions suggest that the affinity of FtsN's SPOR
domain depends on the length of a dnG strand, and the long tails in the stationary lifetime plots
(Supplementary Fig. 10) may arise from longer dnGs.

It remains unclear whether the self-interaction of SPOR on the sPG-track is also cooperative,
but the persistent running time distributions of FtsN exhibited a rise and decay pattern similar to
that of FtsN's stationary lifetime distribution (Supplementary Fig. 3). Notably, FtsW's slow-moving
populations also showed non-exponential distributions (Supplementary Fig. 3). It is likely that
these self-interactions are also cooperative, but further experiments are required to investigate
this possibility.

Biological function of cooperativity of SPOR

What would be the biological function of the cooperative self-interaction of SPOR domains?
Cooperativity is often observed in biology systems when a switch-like behavior in response to a
small change in protein or substrate concentration is required. For example, when LT activities
are low (as mimicked by the $\Delta 6LT$s mutant), tight binding of FtsN to dnG can sequester nearly all
sPG synthesis complexes on the dnG-track, thereby reducing the corresponding sPG synthesis
activity. When sPG synthesis activity is low (as mimicked by aztreonam or fosfomycin treatment),
the diminished slow-moving population of FtsN renders not only more FtsN molecules anchored
to dnG but also tighter binding (stationary lifetime at ~40 s), hence impeding LT activities on dnG
(Fig. 4c, left). Conversely, when sPG synthesis activities are high (such as in the superfission
mutant $ftsI^{R167S}$, in the rich medium EZRDM, or under increased lipid II conditions), more FtsN
molecules bind to the processively-moving sPG synthesis complex, making fewer FtsN molecules
available to bind dnG and also lowering their binding affinities (stationary lifetime at ~14 s).
Consequently, more dnG is exposed to LTs for degradation, leading to higher degradation
activities (Fig. 4c, right).

Furthermore, cooperativity can enhance the binding reaction of SPOR on either the dnG-
track or the sPG-track, facilitating their respective functions. For example, truncation of SPOR led
to a significant reduction in the slow-moving population of FtsW and the processivity of FtsN on
the sPG-rack, suggesting that SPOR's self-interaction on the sPG-track likely enhances the
binding of FtsN to the synthesis complex.

We note that the truncation of SPOR did not impact the processivity of FtsW. The persistent
running time of FtsW on the sPG-track was ~20 s regardless of the presence or absence of FtsN's
SPOR domain (Supplementary Table 5). In contrast, the persistent running time of full-length FtsN
was ~14 s, significantly shorter than those of FtsW under the same conditions (Supplementary
Tables 4 and 5). This discrepancy raises an interesting possibility, that is, multiple FtsN molecules
may cycle within the processive sPG synthesis complex. While each FtsN molecule binds the
complex for ~14 s on average, there is always at least one FtsN molecule in the complex to
maintain FtsWI's processivity for ~20 s. An example trajectory in Fig. 3k may indeed suggest this
process. Furthermore, the slow-moving population percentage and persistent running time
decreased for the dnG-binding defective FtsN^{Q251E}-Halo^{SW} mutant and further for FtsN$^{\Delta SPOR}$-
Halo^{SW} (Fig. 2), suggesting that SPOR's binding to nearby dnG may facilitate the recycling
process by restricting the diffusion of dissociated FtsN molecules.

Lastly, the cooperative binding of SPOR to dnG likely helps prevent the release of dissociated
FtsWIQLB complex to the Z-track. However, the SPOR-dnG binding affinity needs to be in the
right range. If the binding is too tight, as in the Δ6*LT*s background, all FtsWIQLB-FtsN complexes
are sequestered on dnG, reducing sPG synthesis activity. Conversely, in the absence of SPOR-
dnG binding, as in the Δ*amiABC* background, all complexes are sequestered on the Z-track, also
diminishing sPG synthesis activity. We also note that as Δ6*LT*s and Δ*amiABC* cells may have
other significant defects besides different dnG levels, further experiments are required to
investigate these effects.

In summary, we provided strong evidence elucidating a mechanism for how FtsN may
coordinate sPG synthesis and degradation activities in *E. coli*. FtsN controls the populations of
sequestered and active sPG synthesis complexes on the dnG- and sPG-tracks by partitioning
between the two tracks through the cooperative self-interaction of the SPOR domain. While other
protein factors may still be at play, substrate availability and the natural binding affinities of FtsN
to dnG and the synthesis complex can already provide a sensitive switch to regulate the
partitioning of FtsN on the two tracks.

References

1. Vollmer, W. & Bertsche, U. Murein (peptidoglycan) structure, architecture and biosynthesis in *Escherichia coli. Biochim. Biophys. Acta* **1778**, 1714-1734 (2008).
2. van Heijenoort, J. Peptidoglycan hydrolases of *Escherichia coli. Microbiol. Mol. Biol. Rev.* **75**, 636-663 (2011).
3. Jorgenson, M.A., Chen, Y., Yahashiri, A., Popham, D.L. & Weiss, D.S. The bacterial septal ring protein RlpA is a lytic transglycosylase that contributes to rod shape and daughter cell separation in *Pseudomonas aeruginosa. Mol. Microbiol.* **93**, 113-128 (2014).
4. Heidrich, C. *et al.* Involvement of N-acetylmuramyl-L-alanine amidases in cell separation and antibiotic-induced autolysis of *Escherichia coli. Mol. Microbiol.* **41**, 167-178 (2001).
5. Priyadarshini, R., de Pedro, M.A. & Young, K.D. Role of peptidoglycan amidases in the development and morphology of the division septum in *Escherichia coli. J. Bacteriol.* **189**, 5334-5347 (2007).
6. Scheurwater, E., Reid, C.W. & Clarke, A.J. Lytic transglycosylases: bacterial space-making autolysins. *Int. J. Biochem. Cell Biol.* **40**, 586-591 (2008).
7. Vollmer, W., Joris, B., Charlier, P. & Foster, S. Bacterial peptidoglycan (murein) hydrolases. *FEMS Microbiol. Rev.* **32**, 259-286 (2008).
8. Busiek, K.K. & Margolin, W. A role for FtsA in SPOR-independent localization of the essential *Escherichia coli* cell division protein FtsN. *Mol. Microbiol.* **92**, 1212-1226 (2014).
9. Liu, B., Persons, L., Lee, L. & de Boer, P.A. Roles for both FtsA and the FtsBLQ subcomplex in FtsN-stimulated cell constriction in *Escherichia coli. Mol. Microbiol.* **95**, 945-970 (2015).
10. Pichoff, S., Du, S. & Lutkenhaus, J. The bypass of ZipA by overexpression of FtsN requires a previously unknown conserved FtsN motif essential for FtsA-FtsN interaction supporting a model in which FtsA monomers recruit late cell division proteins to the Z ring. *Mol. Microbiol.* **95**, 971-987 (2015).
11. Dai, K., Xu, Y. & Lutkenhaus, J. Topological characterization of the essential *Escherichia coli* cell division protein FtsN. *J. Bacteriol.* **178**, 1328-1334 (1996).
12. Gerding, M.A. *et al.* Self-enhanced accumulation of FtsN at division sites and roles for other proteins with a SPOR domain (DamX, DedD, and RlpA) in *Escherichia coli* cell constriction. *J. Bacteriol.* **191**, 7383-7401 (2009).
13. Park, K.T., Johnson, D.K., Pichoff, S., Du, S. & Lutkenhaus, J. The essential domain of FtsN triggers cell division by promoting interaction between FtsL and FtsI. *bioRxiv* (2023).
14. Yang, J.C., Van Den Ent, F., Neuhaus, D., Brevier, J. & Lowe, J. Solution structure and domain architecture of the divisome protein FtsN. *Mol. Microbiol.* **52**, 651-660 (2004).
15. Britton, B.M. *et al.* Conformational changes in the essential *E. coli* septal cell wall synthesis complex suggest an activation mechanism. *Nat. Commun.* **14**, 4585-4599 (2023).
16. Alcorlo, M. *et al.* Structural basis of denuded glycan recognition by SPOR domains in bacterial cell division. *Nat. Commun.* **10**, 5567-5579 (2019).
17. Moll, A. & Thanbichler, M. FtsN-like proteins are conserved components of the cell division machinery in proteobacteria. *Mol. Microbiol.* **72**, 1037-1053 (2009).
18. Ursinus, A. *et al.* Murein (peptidoglycan) binding property of the essential cell division protein FtsN from *Escherichia coli. J. Bacteriol.* **186**, 6728-6737 (2004).
19. Yahashiri, A., Jorgenson, M.A. & Weiss, D.S. Bacterial SPOR domains are recruited to septal peptidoglycan by binding to glycan strands that lack stem peptides. *Proc. Natl Acad. Sci. USA* **112**, 11347-11352 (2015).
20. McCausland, J.W. *et al.* Treadmilling FtsZ polymers drive the directional movement of sPG-synthesis enzymes via a Brownian ratchet mechanism. *Nat. Commun.* **12**, 609-621 (2021).

21. Yang, X. *et al.* A two-track model for the spatiotemporal coordination of bacterial septal
cell wall synthesis revealed by single-molecule imaging of FtsW. *Nat. Microbiol.* **6**, 584-
593 (2021).
22. Yang, X. *et al.* GTPase activity-coupled treadmilling of the bacterial tubulin FtsZ organizes
septal cell wall synthesis. *Science* **355**, 744-747 (2017).
23. Lyu, Z. *et al.* FtsN maintains active septal cell wall synthesis by forming a processive
complex with the septum-specific peptidoglycan synthases in *E. coli*. *Nat. Commun.* **13**,
5751-5766 (2022).
24. Mahone, C.R. *et al.* Integration of cell wall synthesis and chromosome segregation during
cell division in *Caulobacter*. *J. Cell Biol.* **223**, e20221 (2024).
25. Perez, A.J. *et al.* Movement dynamics of divisome proteins and PBP2x:FtsW in cells of
Streptococcus pneumoniae. *Proc. Natl Acad. Sci. USA* **116**, 3211-3220 (2019).
26. Whitley, K.D. *et al.* Peptidoglycan synthesis drives a single population of septal cell wall
synthases during division in *Bacillus subtilis*. *Nat. Microbiol.* **9**, 1064-1074 (2024).
27. Schaper, S. *et al.* Cell constriction requires processive septal peptidoglycan synthase
movement independent of FtsZ treadmilling in *Staphylococcus aureus*. *Nat. Microbiol.* **9**,
1049-1063 (2024).
28. Perez, A.J. & Xiao, J. Stay on track - revelations of bacterial cell wall synthesis enzymes
and things that go by single-molecule imaging. *Curr. Opin. Microbiol.* **79**, e102490 (2024).
29. Park, K.T., Du, S. & Lutkenhaus, J. Essential role for FtsL in activation of septal
peptidoglycan synthesis. *mBio* **11**, e03012 (2020).
30. Condon, S.G.F. *et al.* The FtsLB subcomplex of the bacterial divisome is a tetramer with
an uninterrupted FtsL helix linking the transmembrane and periplasmic regions. *J. Biol.
Chem.* **293**, 1623-1641 (2018).
31. Marmont, L.S. & Bernhardt, T.G. A conserved subcomplex within the bacterial cytokinetic
ring activates cell wall synthesis by the FtsW-FtsI synthase. *Proc. Natl Acad. Sci. USA*
117, 23879-23885 (2020).
32. Navarro, P.P. *et al.* Cell wall synthesis and remodelling dynamics determine division site
architecture and cell shape in *Escherichia coli*. *Nat. Microbiol.* **7**, 1621-1634 (2022).
33. Yahashiri, A., Kaus, G.M., Popham, D.L., Houtman, J.C.D. & Weiss, D.S. Comparative
study of bacterial SPOR domains identifies functionally important differences in glycan
binding affinity. *J. Bacteriol.* **204**, e0025222 (2022).
34. Peters, N.T., Dinh, T. & Bernhardt, T.G. A fail-safe mechanism in the septal ring assembly
pathway generated by the sequential recruitment of cell separation amidases and their
activators. *J. Bacteriol.* **193**, 4973-4983 (2011).
35. Yakhnina, A.A. & Bernhardt, T.G. The Tol-Pal system is required for peptidoglycan-
cleaving enzymes to complete bacterial cell division. *Proc. Natl Acad. Sci. USA* **117**, 6777-
6783 (2020).
36. Yahashiri, A., Jorgenson, M.A. & Weiss, D.S. The SPOR domain, a widely conserved
peptidoglycan binding domain that targets proteins to the site of cell division. *J. Bacteriol.*
199, e00118 (2017).
37. Squyres, G.R. *et al.* Single-molecule imaging reveals that Z-ring condensation is essential
for cell division in *Bacillus subtilis*. *Nat. Microbiol.* **6**, 553-562 (2021).
38. Buss, J. *et al.* A multi-layered protein network stabilizes the *Escherichia coli* FtsZ-ring and
modulates constriction dynamics. *PLoS Genetics* **11**, e1005128 (2015).
39. Barrows, J.M., Anderson, A.S., Talavera-Figueroa, B.K. & Goley, E.D. Intrinsic and
extrinsic factors regulate FtsZ function in *Caulobacter crescentus*. *bioRxiv* (2023).
40. Baranova, N. *et al.* Diffusion and capture permits dynamic coupling between treadmilling
FtsZ filaments and cell division proteins. *Nat. Microbiol.* **5**, 407-417 (2020).
41. Peisley, A. *et al.* Kinetic mechanism for viral dsRNA length discrimination by MDA5
filaments. *Proc. Natl Acad. Sci. USA* **109**, 3340-3349 (2012).

42.   Morrone, S.R. *et al.* Cooperative assembly of IFI16 filaments on dsDNA provides insights
709   into host defense strategy. *Proc. Natl Acad. Sci. USA* **111**, 62-71 (2014).
710   43.   Hooy, R.M. & Sohn, J. The allosteric activation of cGAS underpins its dynamic signaling
711   landscape. *eLife* **7**, e39984 (2018).
712   44.   Duncan, T.R., Yahashiri, A., Arends, S.J., Popham, D.L. & Weiss, D.S. Identification of
713   SPOR domain amino acids important for septal localization, peptidoglycan binding, and a
714   disulfide bond in the cell division protein FtsN. *J. Bacteriol.* **195**, 5308-5315 (2013).
715   45.   Du, S., Pichoff, S. & Lutkenhaus, J. FtsEX acts on FtsA to regulate divisome assembly
716   and activity. *Proc. Natl Acad. Sci. USA* **113**, 5052-5061 (2016).
717   46.   Mannik, J., Pichoff, S., Lutkenhaus, J. & Mannik, J. Cell cycle-dependent recruitment of
718   FtsN to the divisome in *Escherichia coli*. *mBio* **13**, e0201722 (2022).
719   47.   Kashammer, L. *et al.* Cryo-EM structure of the bacterial divisome core complex and
720   antibiotic target FtsWIQBL. *Nat. Microbiol.* **8**, 1149-1159 (2023).
721   48.   Nguyen, H.T.V. *et al.* Structure of the heterotrimeric membrane protein complex FtsB-
722   FtsL-FtsQ of the bacterial divisome. *Nat. Commun.* **14**, 1903-1915 (2023).
723   49.   Li, G.W., Burkhardt, D., Gross, C. & Weissman, J.S. Quantifying absolute protein
724   synthesis rates reveals principles underlying allocation of cellular resources. *Cell* **157**,
725   624-635 (2014).
726   50.   Arends, S.J. *et al.* Discovery and characterization of three new *Escherichia coli* septal ring
727   proteins that contain a SPOR domain: DamX, DedD, and RlpA. *J. Bacteriol.* **192**, 242-255
728   (2010).
729   51.   Muller, P. *et al.* The essential cell division protein FtsN interacts with the murein
730   (peptidoglycan) synthase PBP1B in *Escherichia coli*. *J. Biol. Chem.* **282**, 36394-36402
731   (2007).
732   52.   Floyd, D.L., Harrison, S.C. & van Oijen, A.M. Analysis of kinetic intermediates in single-
733   particle dwell-time distributions. *Biophys. J.* **99**, 360-366 (2010).

Acknowledgements

We thank all members of the Xiao lab for helpful discussions and feedback on the manuscript, and members of the Weiss lab for help with strain construction. We would like to acknowledge use of resources at the Protein and Crystallography Facility within the Carver College of Medicine at the University of Iowa along with advice and assistance from Devin Reusch and Nicholas Schnicker. Work in the Xiao lab was supported by NIH R01GM086447 and R35GM136436 (to J.X.), GM125656 (subcontract to J.X.), a Hamilton Smith Innovative Research Award (to J.X.). Work in the Weiss lab was supported by NIH R01GM125656 (to D.S.W.).

Author Contributions

Z.L., D.S.W. and J.X. conceived the study. Z.L., A.Y. and D.S.W. constructed the strains and performed genetic and phenotypic experiments. Z.L. and X.Y. performed the imaging experiments and analyzed the data. S.H. performed the HADA labeling and imaging experiments. J.W.M. wrote the custom MATLAB script for speed deconvolution in analyzing the single-molecule tracking data. X.C. performed the crosslinking experiments. X.Y. provided the single-molecule tracking data of Halo-FtsW in *C. crescentus*. B. H. B. provided the single-molecule tracking data of Halo-FtsB in *E. coli*. Z.L., D.S.W. and J.X. wrote the original draft. All authors reviewed and edited the manuscript. D.S.W. and J.X. supervised the study. Funding was acquired by D.S.W. and J.X.

Competing Interests

The authors declare no competing interests.

     Figures and Legends

Fig.1 Altered dnG levels in Δ*amiABC* and Δ6*LT*s cells diminish sPG synthesis activity

a-b, Bright-field and ensemble fluorescent images of FtsN-Halo[SW] (top) and FtsW-RFP (bottom) fusions in Δ*amiABC* (**a**) and Δ6*LT*s (**b**) strain backgrounds. Scale bars, 5 μm. **c-d**, Speed distributions of FtsN-Halo[SW] (top) and FtsW-RFP (bottom) in Δ*amiABC* (**c**) and Δ6*LT*s (**d**) strain

backgrounds overlaid with the fit curves of the stationary (black), slow-moving (red), and fast-moving (blue) populations. Schematic representation of the fusion and strain backgrounds are shown in inset (pink: labeled protein; green: Halo or RFP fluorophore). **e,** Percentages of the stationary (gray), slow-moving (red), and fast-moving (blue) populations of FtsN-Halo^{SW} and FtsW-RFP fusions in WT, $\Delta amiABC$, and $\Delta 6LT$s strain backgrounds. **f,** Box plots of the lifetime of the stationary population under all the conditions in (**e**). Red lines indicate the mean values. **g,** Representative images of HADA-labeled WT, $\Delta amiABC$, and $\Delta 6LT$s cells. Scale bars, 5 µm. **h,** Box plots of septal HADA fluorescence intensity percentage normalized to the whole cell intensity in WT, $\Delta amiABC$, and $\Delta 6LT$s cells treated or untreated with cephalexin to inhibit FtsI.

Fig.2 SPOR domain enhances FtsW and FtsN's slow-moving populations.

a, **d-f**: Speed distributions of FtsW-RFP in $ftsN^{\Delta SPOR+}$ strain background (**a**), FtsN$^{\Delta SPOR}$-Halo^{SW} in $ftsN^{\Delta SPOR+}$ strain background (**d**) and treated with aztreonam (**e**), and FtsN^{Q251E}-Halo^{SW} in WT background (**f**), overlaid with the fit curves of the stationary (black), slow-moving (red), and fast-moving (blue) populations. Schematic representation of the fusion and strain background are shown in insets. **b**, sPG-track processivity (top; defined as the persistent running time of the slow-moving population) and Z-track processivity (bottom; defined as the persistent running time of the fast-moving population) under the conditions in (**a**, **d-f**). **c**, Percentages of the stationary (gray), slow-moving (red), and fast-moving (blue) populations under all the conditions in (**a**, **d-f**). FtsW and FtsN data were replotted from figure (**1e**) for comparison.

Fig.3 SPOR domain exhibits SPOR-SPOR self-interaction.

a, Schematic representation of FtsN^{WYAA}-Halo^{SW} fusion in WT *ftsN* background. **b**, speed distribution of FtsN^{WYAA}-Halo^{SW} overlaid with the fit curves of the stationary (black) and slow-moving (red) populations. **c**, Percentages of the stationary (gray) and slow-moving (red)

populations under all the conditions in **b**, **f**, **h**, **j**, and **S3**. **d**, Top: sPG-track processivity of

FtsN^{WYAA}-Halo^{SW}, Halo-iSPOR, and Halo-iSPOR^{Q251E} fusions in WT strain. Bottom: Stationary

lifetime of all the conditions in (**c**). **e**, Schematic representation of Halo-iSPOR fusion in WT *ftsN*

background. **f**, speed distribution of Halo-iSPOR overlaid with corresponding fit curves. **g**,

Schematic representation of Halo-iSPOR^{Q251E} fusion in WT *ftsN* background. **h**, Speed distribution

of Halo-iSPOR^{Q251E} overlaid with corresponding fit curves. **i**, Schematic representation of Halo-

iSPOR fusion in *ftsN*$^{\Delta SPOR+}$ strain background; **j**, speed distribution of Halo-iSPOR overlaid with

corresponding fit curves. **k**, Representative kymographs (top) and intensity time traces (bottom)

of FtsN-Halo^{SW} multimers at the septum. Left: an FtsN-Halo^{SW} molecule transiently associated

and disassociated (diffusive fluorescence intensity) with a processively moving FtsN-Halo^{SW}

molecule (stable fluorescence intensity) between 20 s and 100 s, and eventually stably associated

(sudden doubling of fluorescence intensity) and moved together at an average speed of 7 nm s^{-1}

; Right: one FtsN-Halo^{SW} molecule was photo-bleached (~155 s) in a stationary dimer. Scale

bars, 1 μm.

Fig.4 FtsN coordinates sPG synthesis and degradation both in space and time.

a, The reverse of normalized FtsN occupancy on dnG ($1/N_{dnG}$) as a proxy for sPG degradation activity correlates with normalized FtsN occupancy on sPG (N_{sPG}) across all conditions (black), except when the SPOR domain is unlinked from the rest of the FtsN (red). All the data points are labeled with alphabetical letters, in the order of elevated sPG occupancy (N_{sPG}) as a proxy for sPG synthesis activity. **b**, The ratio between $1/N_{dnG}$ and N_{sPG}, the proxy for the ratio between relative sPG degradation and synthesis activities is close to one (black), but became significantly higher when the SPOR domain was unlinked from the rest of the FtsN (red). The conditions are labeled with the same letters in (**a**). **c**, A three-track model depicting how FtsN coordinates sPG synthesis and degradation both in space and time. FtsN marks a third track, termed as the dnG-track, through its SPOR domain's binding to dnG. When sPG synthesis activity is low (e.g., under

antibiotic treatment, left), FtsN shifts from the sPG-track to the dnG-track and consequently

sequesters FtsWIQLB on dnG. Cooperative binding of SPOR to dnG inhibits LTs' access to dnG,

thus decreasing sPG degradation activity correspondingly. Conversely, when sPG synthesis

activity is high (e.g., in rich medium, right), FtsN shifts from the dnG-track to the sPG-track,

thereby exposing dnG and facilitating its degradation by LTs, consequently increasing sPG

degradation activity.