Oxidized low-density lipoprotein potentiates angiotensin II-induced Gq activation through the AT1-LOX1 receptor complex：Implications for renal dysfunction

Jittoku Ihara [1]#, Yibin Huang [1]2)#, Yoichi Takami [1]*, Yoichi Nozato [1]*, Toshimasa Takahashi [1]3), Akemi Kakino [4], Cheng Wang [1], Ziwei Wang [1], Yu Guo [1], Weidong Liu [1], Nanxiang Yin [1], Ryoichi Ohara [1], Taku Fujimoto [1], Shino Yoshida [1], Kazuhiro Hongyo [1], Hiroshi Koriyama [1], Hiroshi Akasaka [1], Hikari Takeshita [1], Shinsuke Sakai [5], Kazunori Inoue [5], Yoshitaka Isaka [5], Hiromi Rakugi [1], Tatsuya Sawamura [4], and Koichi Yamamoto [1]

[1] Department of Geriatric and General Medicine, Osaka University Graduate School of Medicine, 2-2 Yamada-oka, Suita, Osaka, 565-0871, Japan

[2] Center for Pulmonary and Vascular Biology, Dept. of Pediatrics, University of Texas Southwestern Medical Center, 5323 Harry Hines Blvd., Dallas, TX, 75390, USA

[3] Department of Medicine, Takahashi clinic, Kobe, Hyogo, 657-0846, Japan

[4] Department of Molecular Pathophysiology, Shinshu University Graduate School of Medicine, Matsumoto, Nagano 390-8621, Japan

[5] Department of Nephrology, Osaka University Graduate School of Medicine, 2-2 Yamada-oka, Suita, Osaka, 565-0871, Japan

#These authors contributed equally to this work.

Correspondence and requests for materials should be addressed to

Yoichi Takami, M.D., Ph.D.*

Department of Geriatric and General Medicine, Osaka University Graduate School of

Medicine, #B6, 2-2 Yamadaoka, Suita, Osaka 565-0871, Japan

Tel.: +81-6-6879-3852, Fax: +81-6-6879-3859, E-mail: takami@geriat.med.osaka-

u.ac.jp

and

Yoichi Nozato, M.D., Ph.D.*

Department of Geriatric and General Medicine, Osaka University Graduate School of

Medicine, #B6, 2-2 Yamadaoka, Suita, Osaka 565-0871, Japan

Tel.: +81-6-6879-3852, Fax: +81-6-6879-3859, E-mail:

yoichi.nozato@geriat.med.osaka-u.ac.jp

Keywords: AT1, LOX-1, oxidized LDL, Angiotensin II, Gq, CKD

Abstract

Chronic kidney disease (CKD) and atherosclerotic heart disease, frequently associated

with dyslipidemia and hypertension, represent significant health concerns. We

investigated the interplay among these conditions, focusing on the role of oxidized low-

density lipoprotein (oxLDL) and angiotensin II (Ang II) in renal injury via G protein αq

subunit (Gq) signaling. We hypothesized that oxLDL enhances Ang II-induced Gq

signaling via the AT1 (Ang II type 1 receptor)-LOX1 (lectin-like oxLDL receptor)

complex. Based on CHO and renal cell model experiments, oxLDL alone did not activate

Gq signaling. However, when combined with Ang II, it significantly potentiated Gq-

mediated inositol phosphate 1 production and calcium influx in cells expressing both

LOX-1 and AT1 but not in AT1-expressing cells. This suggests a critical synergistic

interaction between oxLDL and Ang II in the AT1-LOX1 complex. Conformational

studies using AT1 biosensors have indicated a unique receptor conformational change due

to the oxLDL-Ang II combination. In vivo, wild-type mice fed a high-fat diet with Ang

II infusion presented exacerbated renal dysfunction, whereas LOX-1 knockout mice did

not, underscoring the pathophysiological relevance of the AT1-LOX1 interaction in renal

damage. These findings highlight a novel mechanism of renal dysfunction in CKD driven

by dyslipidemia and hypertension and suggest the therapeutic potential of AT1-LOX1

55   receptor complex in patients with these comorbidities.

Introduction

Dyslipidemia is a major risk factor of atherosclerotic heart disease in patients with chronic kidney disease (CKD). It has been postulated that the association between dyslipidemia and CKD is not solely a result of epidemiological comorbidities but rather a complex interplay of causality, where CKD exacerbates dyslipidemia, while dyslipidemia, in turn, contributes to the onset and progression of CKD[1,2]. Since Moorhead et al. proposed the lipid nephrotoxicity hypothesis in 1982[3], accumulating evidence has suggested that increased plasma lipid levels contribute to the development of renal glomerular and tubular damage, primarily in animal models of dyslipidemia[4]. The etiology of dyslipidemia-induced nephrotoxicity is complex and multifaceted, potentially involving the activation of certain cellular signaling pathways that culminate in renal injury through elevated levels of oxidized LDL (oxLDL)[5-8]. The lectin-like oxLDL receptor, LOX-1, is implicated in organ damage caused by dyslipidemia, and its expression is increased in hypertensive glomerulosclerosis[9]. In murine models, a deficiency of LOX-1 leads to a reduction in renal dysfunction, which was precipitated by a systemic inflammatory state following significant myocardial ischemia and injury[10]. This supports the hypothesis that LOX-1 plays a role in the development of inflammation-induced renal injury. However, to date, no studies have investigated the role of LOX-1 in nephrotoxicity caused by

dyslipidemia.

Hypertension is an established risk factor for CKD, and it has been suggested that

hypertension and dyslipidemia act synergistically to induce renal dysfunction[11]. The

development of renal damage due to hypertension involves direct renal injury by

vasoactive hormones, such as angiotensin II (Ang II), in addition to renal hemodynamic

abnormalities associated with elevated body pressure[12]. We have shown that LOX-1 and

the Ang II type 1 receptor (AT1) of the G protein-coupled receptor (GPCR) are coupled

on the plasma membrane, and that G protein-dependent and β-arrestin-dependent AT1

activation mechanisms are involved in the signaling mechanism by oxLDL and the

intracellular uptake of oxLDL, respectively[13,14].

Interestingly, it was recently demonstrated that AT1 exhibits different modes and degrees

of G protein activation depending on the conformational changes that occur during

activation by various ligands[15]. Specifically, G protein αq subunit (Gq)-biased agonists

induce a more open conformation of AT1 than Ang II, resulting in a more potent activation

of Gq, which is the primary mediator of Ang II-induced hypertension. In contrast, β-

arrestin-biased agonists induce a closer conformational change, leading to the activation

of β-arrestin without the activation of Gq[15]. Indeed, we found that oxLDL selectively

activates G protein αi subunit (Gi) of AT1, without activating Gq[14], similar to that induced

by β-arrestin-biased agonists[16]. However, given that Ang II and oxLDL coexist in

physiological environments, it is plausible to hypothesize that the effect of these ligands

on AT1 in living organisms may differ from their effects when administered individually.

Indeed, we found that the combination treatment of oxLDL and Ang II enhanced pro-

inflammatory NFκB activity compared to each treatment alone in cells overexpressing

both AT1 and LOX-1[14]. Based on our findings and the aforementioned structure-activity

relationship of AT1, we hypothesized that the binding of both oxLDL and Ang II to LOX-

1 and AT1, respectively, may result in a more open AT1 structure, leading to stronger

downstream Gq signaling. Therefore, this study aimed to investigate and clarify this

hypothesis, focusing specifically on renal component cells, and ultimately demonstrate

the in vivo relevance of this phenomenon in the development of renal injury or renal

dysfinction under conditions of Ang II and oxLDL overload.

Results

Oxidized LDL potentiates Ang II-induced Gq signaling in a LOX-1-dependent manner

First, we examined the additive effect of oxLDL on Ang II-stimulated AT-1-Gq signaling in CHO (Chinese hamster ovary) cells that did not endogenously express LOX-1 and AT-1 but were genetically engineered to express these receptors (CHO-LOX-1-AT1) [13].

Fig. 1a shows the dose-response curve of Ang II-induced IP1 production, which serves as a measure of Gq activity, in the presence of varying concentrations of oxLDL in CHO-LOX-1-AT1 cells. Consistent with previous findings, oxLDL alone did not stimulate IP1 production in CHO-LOX-1-AT1 cells [14]. However, when oxLDL was supplemented at concentrations of 10 and 20 μg/mL (but not at 5 μg/mL), it caused a similar leftward shift in the dose-response curve of Ang II, resulting in a more than 80% decrease in the Effective Concentration 50 (EC50) (EC50 values: 0 μg/mL, 9.40×10^{-9} M; 5 μg/mL, 5.21×10^{-9} M; 10 μg/mL, 1.68×10^{-9} M; 20 μg/mL, 1.55×10^{-9} M). The maximum IP1 production induced by Ang II in CHO-LOX-1-AT1 cells was unaffected by the addition of oxLDL.

 Additionally, we observed that oxLDL administration decreased Ang II-induced IP1 production in CHO cells expressing AT1 alone (CHO-AT1), in contrast to CHO cells

expressing both AT1 and LOX-1 (Fig. 1b). This finding suggests that the potentiating

effect of oxLDL on Ang II-AT1-Gq activity is dependent on the presence of LOX-1.

However, the reason for the decrease in IP1 production caused by oxLDL

supplementation was not determined in this study. Native LDL, which does not bind to

LOX-1, did not alter Ang II-induced Gq activity in CHO-LOX-1-AT1 cells (Fig. 1c).

Furthermore, the presence of advanced glycation end products (AGEs) that bind to LOX-

1[17], did not enhance Ang II-induced IP1 production (Fig. 1d).

We found that enhanced IP1 production by co-treatment of oxLDL with Ang II was

similarly observed in CHO-cells expressing LOX-1 and mutated AT1 with impaired

ability to activate β arrestin[14] (Fig 1e). Moreover, the potentiating effect of oxLDL on IP1

production was unaffected by Pertussis Toxin (PTX), a Gi inhibitor, or RKI-1448, a

downstream Rho kinase inhibitor targeting G12/13 signaling in CHO-LOX-1-AT1 cells.

However, this phenomenon was completely inhibited by YM-254890, the Gq inhibitor

(Fig. 1f). These results suggest that the potentiating effect of oxLDL on Ang II-induced

IP1 production is not influenced by β-arrestin, Gi, or G12/13, which are the main effectors

of AT1 signaling aside from Gq[18].

oxLDL potentiates Ang II-induced calcium influx in a LOX-1-dependent manner

Calcium influx is a representative cellular phenomenon that occurs in response to Ang II-

AT1-Gq activation. We found that oxLDL and low concentrations of Ang II (10^{-12} M) did

not induce calcium influx in CHO-LOX-1-AT1 cells when treated alone, but apparently

induced calcium influx when treated together (Fig. 1g, h). Ang II alone at higher

concentrations (10^{-11} M) induced calcium influx, and no further enhancement was

observed with oxLDL supplementation (Fig. 1h). The combined effect of oxLDL and Ang

II on calcium influx was completely blocked by YM254890, suggesting that this

phenomenon was Gq-dependent (Fig. 1i). Importantly, oxLDL supplementation with Ang

II did not affect the calcium influx in CHO cells expressing AT1 alone (Fig. 1j).

Co-treatment of oxLDL with Ang II induces conformational change of AT1 different

from each treatment alone

To gain mechanistic insight into this phenomenon, we initially conducted a live-imaging

analysis of membrane LOX-1 and AT1 to determine whether the combined treatment of

oxLDL and Ang II leads to increased internalization of AT1 upon activation compared to

individual treatments, as described in a previous protocol paper[19] (Fig. S1, Supplemental

Video, and Fig. 2a).

Our findings revealed a decrease in green puncta, which represent AT1-eGFP, upon

treatment with Ang II, oxLDL alone, and their co-treatment compared with the control

group (Fig. 2a). However, the extent of membrane AT1 reduction was similar across all

the treatment groups (Fig. 2a). Consistent with our previous report[14], oxLDL treatment

resulted in a reduction in the red puncta representing LOX-1-mScarlet (Fig. 2a).

Importantly, co-treatment with oxLDL and Ang II did not further enhance the reduction

of red puncta compared to oxLDL treatment alone (Fig. 2a). Based on these findings, it

is conceivable that co-treatment with oxLDL and Ang II does not increase the content of

activated AT1 compared to individual treatments alone.

AT1 intramolecular FlAsH-BRET biosensors were used to detect AT1 conformational

changes in CHO cells expressing LOX-1[20]. Among several biosensors previously tested

[20], we used two sensors with FlAsH insertion at the third intracellular loop (ICL3P3) and

cytoplasmic-terminal tail (C-tailP1) of AT1 that interacts with Renilla luciferase (RlucII)

at the end of the cytoplasmic tail (AT1-ICL3P3 and AT1-C-tailP1, respectively), as these

sensors were shown to enhance BRET signaling induced by Ang II compared to biased

agonists, including SI, SII, DVG, and SBpA[20]. In CHO cells expressing LOX-1 alone

(CHO-LOX-1) transduced with lentivirus encoding AT1-C-tailP1, 10^{-5} M Ang II and the

combination of Ang II and 10 μg/mL oxLDL induced BRET similarly, whereas oxLDL

alone did not alter BRET (Fig. 2b). In contrast, in CHO-LOX-1 cells transduced with

AT1-ICL3P3-encoded lentivirus, the combination of Ang II and oxLDL induced BRET

more prominently than Ang II alone, whereas oxLDL alone did not alter BRET (Fig. 2c).

Furthermore, the difference between oxLDL and the combination treatment was

abolished by a neutralizing antibody against LOX-1 (Fig. 2d). These findings suggested

that the concomitant binding of oxLDL to LOX-1 and Ang II to AT1 induced a

conformational change in AT1 that was distinct from that induced by Ang II or oxLDL

alone.

Oxidized LDL potentiates Ang II-induced Gq-calcium signaling in renal cells

To confirm the pathophysiological significance of this phenomenon observed in the

overexpressing cells, we validated it in cells endogenously expressing LOX-1 and AT1.

We found that oxLDL in combination with Ang II did not increase the cellular IP1 content

in human umbilical vein endothelial cells (HUVECs), bovine vascular endothelial cells

(BACEs), human aortic vascular smooth muscle cells (HAVSMCs), or rat macrophages

(A10) (Fig. S2). In contrast, in normal rat kidney epithelial cells (NRK52E) and

fibroblasts (NRK49F), the combination of oxLDL and Ang II, but not Ang II alone,

increased IP1 accumulation, which was suppressed in the presence of YM-25480, the Gq

inhibitor (Fig. 3a, b). IP1 accumulation induced by co-treatment with Ang II and oxLDL

was abolished by the siRNA-mediated knockdown of either AT1 or LOX-1 (Fig. 3 c, d;

the knockdown efficacy is shown in Fig. S3). Regarding calcium influx, intracellular

calcium concentrations were not increased by the treatment of either 10^{-7} M Ang II or low

concentration of oxLDL (2 μg/ml) alone (Fig. 3e, left and middle, Fig 3f). In contrast, the

combination of Ang II (10^{-7} M) and oxLDL (2 μg/ml) increased intracellular Ca^{2+}

concentration (Fig. 3e, right, Fig. 3f). The combined effect on calcium influx was

attenuated by the siRNA knockdown of either AT1 or LOX-1 (Fig. 3g). The Gq inhibitor

and angiotensin receptor blocker (ARB), Irbesartan, also inhibited this phenomenon (Fig.

3h). Calcium influx was not induced by either combination therapy or monotherapy in

NRK52E cells (Fig. S4).

Co-treatment of oxLDL and Ang II enhanced cellular response upon Gq activation

in renal cells

In NRK49F cells, co-treatment of oxLDL and Ang II, compared to vehicle, increased

mRNA levels of NADPH components, *p67phox* and *p91phox*, fibrosis markers,

fibronectin, *collagen-1*, *collagen-4*, and *TGFβ*, and inflammatory cytokines, *TNFα*, *IL1β*,

IL-6, and *MCP-1* (Fig. 4a). Notably, oxLDL did not alter the expression of the genes of

interest, and Ang II increased the mRNA levels of only *fibronectin* and *MCP-1*, indicating

the apparent synergistic effect of co-treatment with Ang II and oxLDL on specific gene

expression. This amplification effect of co-treatment was also observed in limited genes

including *p67phox*, *TGFβ, IL-6,* and *MCP-1* in NRK52E cells (Fig. 4b). The combined

effects of oxLDL and Ang II on gene expression were completely abolished by the Gq

inhibitor (Fig. 4c, d) and ARB (Fig. 4e, f) in NRK49F and NRK 52E. We also verified

protein expression of αSMA as a molecular marker of epithelial-mesenchymal transition

(EMT) upon the indicated stimulation for 3 days in rat kidney cells. The combination of

Ang II (10^{-7} M) and oxLDL (5 μg/mL) induced αSMA expression in NRK49F or NRK52E

to the same extent or less than TGFβ, a major inducer of EMT, respectively (Fig. 5a, b).

The results of the combination treatment were strikingly different from that of oxLDL or

Ang II treatment alone, which did not affect αSMA expression in both types of cells (Fig.

5c, d). The induction of αSMA by the combined treatment was suppressed in the presence

of a Gq inhibitor or ARB in both types of cells, suggesting the AT1-Gq-dependent

pathway in this phenomenon (Fig. 5e, f and Fig. 5g, h). As a final in vitro assay, the

proliferative activity of NRK49F cells after 24 h of treatment with Ang II and oxLDL,

either alone or in combination, was measured using the BrdU assay (Fig. 6a). When

administered alone, Ang II and oxLDL increased and reduced BrdU incorporation,

respectively. The reducing effect of oxLDL on proliferation was blocked by Irbesartan,

but not by a Gq inhibitor, consistent with our findings that oxLDL induces biased

activation of AT1, which favors Gi but not Gq signaling[14]. Oxidized LDL potentiated the

Ang II-induced increase in BrdU incorporation, which was blocked in the presence of a

Gq inhibitor or an ARB (Fig. 6a). The siRNA-mediated knockdown of AT1 completely

abolished the effects of Ang II and oxLDL, either alone or in combination. Knockdown

of LOX-1 did not affect the pro-proliferative effect of Ang II itself but blocked the

enhanced proliferation induced by co-treatment with oxLDL (Fig. 6b).

Oxidized LDL-inducible diet exacerbates Ang II-induced renal dysfuntion in

wildtype mice, but not in LOX-1 knockout mice.

To examine the relevance of this phenomenon in renal injury, we replicated the in vivo

conditions that occurred during simultaneous stimulation with oxLDL and Ang II in

wildtype (WT) and LOX-1 knockout (LOX-1 KO) mice. This was achieved by inducing

oxLDL via a high-fat diet (HFD) [21] and Ang II via a subcutaneously implanted osmotic

mini-pump (Fig. 7a). Two different doses of Ang II were used in the experiment: a pressor

dose of 0.7 γ, which was demonstrated to increase blood pressure (BP) and induce renal

dysfunction[22,23], and a subpressor dose of 0.1 γ, which does not affect BP. For the intended

analysis, the results are presented separately for mice exposed to subpressor and pressor

doses of Ang II, utilizing the same outcomes of control animals infused with saline for

comparison. Consistent with a previous report, the HFD used in the study prominently

increased the plasma LOX-1 ligand concentration, which was undetectable under a

normal diet (ND) after 6 weeks of feeding (Fig. S5). Notably, the HFD did not intensify

but rather attenuated the body weight increase during the experimental period compared

to the ND (Fig. S6a-c). The pressor dose of Ang II similarly increased BP in HFD-fed and

ND-fed WT mice (Fig. 7b, Fig. S6d). However, notably, there was a modest trend of

attenuated BP elevation by 0.7 γ Ang II infusion in LOX-1 KO mice, in line with the

previous report showing reduced Ang II-induced BP elevation by LOX-1 deficiency in

mice (Fig. 7b, Fig. S6d, f)[24]. As expected, a subpressor dose of Ang II did not alter BP in

the corresponding mouse group (Fig. 7b, Fig. S6e). Regarding biofluid analysis, HFD

with saline infusion did not alter the urinary 8-OHdG concentration as a marker of

oxidative stress and urinary albumin excretion (UAE) compared to ND with saline

infusion in either WT or LOX-1 KO mice (Fig. 7c, d). The pressor dose of Ang II with

ND significantly increased the urinary 8-OHdG concentration and UAE in WT mice (Fig.

7c, d). Notably, HFD feeding in WT mice with a pressor dose of Ang II resulted in a

prominent increase in urinary 8-OHdG concentration and UAE compared to mice fed

with ND (Fig. 7c, d). In WT mice administered with a subpressor dose of Ang II, a

significant increase in UAE was observed when comparing the effects of HFD to ND (Fig.

7d). There were no significant differences in urinary 8-OHdG levels between the two

dietary conditions (Fig. 7c). Interestingly, when LOX-1 KO mice were fed either HFD or

ND and then administered the corresponding dose of Ang II, no differences were observed

in the measured parameters (Fig. 7c, d). These findings indicate that the combination of

HFD and Ang II administration appears to have a more pronounced effect on certain

biofluid markers of renal injury in WT mice than in LOX-1 KO mice. The presence or

absence of LOX-1 appears to influence the interaction between HFD and Ang II, affecting

these specific parameters in mice. We did not find a difference in plasma aldosterone

concentration between ND- and HFD-fed WT mice with a pressor dose of Ang II (Fig.

S7).

We then conducted quantitative real-time PCR analysis on genes within kidney sections,

encompassing NADPH components (*p40phox, p47phox, p67phox, and p91phox*),

inflammatory cytokines (*IL-6, TNFα, IL1β, MCP-1, and COX-2*), fibrosis markers (*TGFβ,*

fibronectin, collagen-1a, collagen-4a, αSMA, and vimentin), epithelial markers (*E-*

cadherin and cadherin-16), and tubular marker (*NAGL*) (Fig. 8, Fig S8). In mice treated

with a pressor dose of Ang II, 15 out of 18 genes examined showed enhanced alterations

in gene expression, indicative of heightened NADPH oxidase components, inflammatory

cytokines, fibrosis, decreased epithelial markers, and exacerbated tubular injury in HFD-

fed WT mice compared to ND-fed mice (Fig. 8a, Fig. S8a). For many of these genes, a

synergistic effect of the combination of a pressor dose of Ang II and HFD was not

observed in LOX-1 KO mice (Fig. 8a, Fig. S8a). Seven genes displayed discernible

differences between HFD- and ND-fed WT mice treated with a subpressor dose of Ang

II (Fig. 8b Fig. S8b). Other than *fibronectin*, no differences were observed between ND-

and HFD-fed LOX-1 mice exposed to a subpressor dose of Ang II (Fig. 8b Fig. S8b). In

relation to AT1 and LOX-1 expression, no variations emerged between the ND and HFD

groups when subjected to the corresponding dose of Ang II treatment in both WT and

LOX-1 KO mice (Fig. S8a, b).

In the histological analysis of kidney samples, in contrast to the gene expression data, the

administration of either a subpressor or pressor dose of Ang II, both in isolation and in

combination with HFD for 4 weeks, did not reveal any notable increase in fibrosis, as

evaluated by Masson-Trichrome staining (Fig. S9a). Similarly, there was no significant

change in the degree of mesangial expansion or glomerular area, as assessed by PAS

staining. (Fig. S9b).

Finally, immunofluorescence staining of mouse kidney specimens was performed to

detect the colocalization sites of LOX-1 and AT1 in the kidney. As shown in Fig. 9a and

Fig. 9b, LOX-1 and AT1a were predominantly colocalized in the renal tubules, but not in

the glomeruli.

Discussion

We conducted a series of experiments to provide empirical support for our hypotheses.

We propose that the simultaneous binding of Ang II to AT1 and oxLDL to LOX-1 triggers distinct and more pronounced structural modifications in AT1 than the individual modifications induced by each ligand. This structural alteration in AT1 leads to the enhanced activation of Gq signaling.

Our experiments showed that oxLDL enhanced the effects of Ang II-induced production of IP1, a downstream signaling molecule associated with Gq activation and subsequent calcium influx. However, this effect was observed only in the presence of both LOX-1 and AT1. Considering that oxLDL selectively activates Gi but not Gq through the LOX-1-AT1 dependent pathway[14], it is evident that the observed phenomenon cannot be attributed solely to the additive effect of oxLDL on AT1 activation. Rather, the simultaneous binding of oxLDL and Ang II to their respective receptors, LOX-1 and AT1, which form a single complex, underlies this phenomenon. Indeed, our findings from live imaging of the membrane receptors revealed that the combination of Ang II and oxLDL did not induce any additional influence on the internalization of both receptors upon AT1-β-arrestin activation compared to each ligand alone, suggesting that the quantity of activated receptors is not affected by the combination treatments. Considering the

conformation-activation relationship in AT1 activation[20], this supports the hypothesis that

the simultaneous binding of Ang II and oxLDL in a single AT1-LOX-1 complex induces

a greater conformational change, resulting in a more open conformation of AT1 compared

with each ligand alone. However, it is technically challenging to directly identify

structural modifications of the receptor complex using techniques such as crystal structure

analysis. Instead, we utilized AT1 conformational sensors capable of differentiating

between the conformational changes induced by Ang II and biased AT1[20]. Interestingly,

we observed that one of the two conformational sensors employed in this study detected

an augmented response in the presence of the combination treatment compared with Ang

II alone. Importantly, this enhanced response was significantly inhibited when an LOX-1

antibody was introduced, indicating the dependency of LOX-1 on this phenomenon.

These results indicate that the combined treatment with Ang II and oxLDL in the presence

of LOX-1 induces a unique conformational change in individual AT1 molecules, which

differs from the conformational changes induced by each single treatment alone.

It is important to note that the ability of LOX-1 ligands to enhance Ang II-AT1 signaling

is not commonly observed. This was corroborated by the observation that BSA-

conjugated AGE, a recognized ligand of LOX-1[17], failed to augment the production of

IP1 by Ang II compared to control BSA (Fig. 1d). While the exact reason for this

discrepancy is not yet understood, it is noteworthy that the predicted particle size of

oxLDL[25] is significantly larger at 250 Å compared to the maximum particle size of AGE-

BSA[26], which is 120 Å. This suggests a potentially greater impact of oxLDL on the

structural modifications within the AT1-LOX-1 complex. However, further structural

analysis is required to validate this hypothesis.

Notably, the Gq bias resulting from combination treatment varied across the mammalian

cells examined. Specifically, we observed a combinatorial effect exclusively in renal

epithelial and fibroblasts, whereas vascular endothelial and smooth muscle cells did not

display the same response. In these renal cells, we observed increased Gq signaling, along

with other cellular phenomena such as calcium influx, changes in gene expression, and

alterations in cellular characteristics, including EMT and cell proliferation. Specifically,

EMT is an important cellular phenomenon in renal epithelial cells to acquire the

phenotype of myofibroblasts and induce fibrosis by the excessive production and

deposition of extracellular matrix (ECM) proteins, although there is a debate regarding

the precise role of EMT in vivo[27]. Numerous studies have suggested that the EMT plays

a critical role in Ang II-mediated renal fibrosis. EMT is also involved in the pathological

transition of normal fibroblasts to myofibroblasts, which promotes fibrosis. We found that

αSMA expression, a well-known indicator of EMT, was dramatically enhanced when

renal epithelial cells and fibroblast cells were subjected to the combination treatment of

Ang II and oxLDL, in comparison to individual treatments. Additionally, fibroblast

proliferation, as assessed by the BrdU assay, was notably enhanced by the combined

treatment, as opposed to each treatment administered separately. Interestingly, the use of

5 μg/mL oxLDL in our study showed a tendency to decrease proliferation, which was

counteracted by the administration of an ARB but not a Gq inhibitor. These findings

suggest that the presence of Ang II leads to a significant transformation in the function of

oxLDL, primarily due to its altered influence within the AT1-LOX-1 complex. Taken

together, these results suggest that the simultaneous binding of oxLDL to LOX-1 and Ang

II to AT1 results in a Gq-biased shift in AT1 activation, leading to a cellular phenomenon

that could potentially contribute to renal damage.

In an animal study, we introduced a biological environment in which both circulating Ang

II and oxLDL (an LOX-1 ligand) were increased in mice. Previous studies using mice fed

an HFD have consistently reported the onset of renal injury, as determined by various

measurements[28-32]. In contrast, 6 weeks of HFD feeding without Ang II treatment did not

alter renal function in our mice. This can be attributed to the lack of obesity induced by

the high-fat diet in this study. We used this diet based on a previous study that confirmed

increased circulating LOX-1 ligand levels without body weight gain in mice[33], and this

diet decreased body weight in our experiments. Obesity is a well-established and

clinically proven risk factor of renal dysfunction. The mechanisms underlying this

association are complex and involve various factors other than lipid abnormalities, such

as hemodynamic changes that affect kidney circulation and the impact of adipose tissue

on the production of adipokines and other inflammatory mediators[34,35]. Consequently, we

observed the influence of elevated lipid particle levels on renal function, independent of

obesity. We found that the effect of an HFD became obvious with an increase in the Ang

II load in WT mice. In particular, in WT mice treated with high-dose Ang II, which

elevated systolic BP by approximately 30 mm Hg, an HFD induced notable increases in

urinary reactive oxygen species and urinary albumin. In contrast, an HFD had no impact

on Ang II-infused LOX-1 KO mice, as evidenced by equivalent urinalysis measurements

for renal injury between the diets. Correspondingly, simultaneous administration of an

HFD and Ang II resulted in a consistent alteration in the expression of genes related to

renal injury, including fibrosis, inflammation, and oxidative stress, except for some genes

in WT mice, but not in LOX-1 KO mice. This strongly suggests that the combined effect

of Ang II and HFD on renal function is LOX-1 dependent. Nevertheless, it should be

noted that the effect of high-dose Ang II infusion on BP tended to be less pronounced in

LOX-1 KO mice compared to WT mice, although there were no differences in BP

elevation between the diets in each group of mice. These findings align with those of

previous studies indicating that LOX-1 knockout mice show resistance to Ang II-induced

elevation of BP[24,36,37]. Specifically, when mice were infused with 2 γ Ang II (equivalent

to 25 g mice) for a duration of 28 days, wildtype mice experienced a BP increase

exceeding 180 mmHg, while LOX-1 knockout mice demonstrated a reduction of

approximately 40 mmHg in this elevation[24]. Furthermore, the same research group

reported that Ang II infusion led to less severe renal injury in LOX-1KO mice compared

to WT mice[36]. Importantly, these findings were observed in mice fed with an ND,

suggesting that the protective effect of LOX-1 loss-of-function against Ang II-induced

elevated BP occurs through the LOX-1-AT1 complex, independent of the presence of

oxLDL. Taken together, the effects of LOX-1 on AT1 signaling are complex, involving

both ligand-dependent and -independent mechanisms, and further investigation is

required for a comprehensive understanding of the LOX-1-AT1 interaction.

Finally, the current findings unequivocally demonstrated the molecular interactions

between key molecules associated with dyslipidemia and hypertension in the kidneys.

Moreover, this interaction can be effectively inhibited by ARBs, suggesting an additive

effect in preventing the development of CKD, particularly in patients with hypertension

and dyslipidemia. Interestingly, Ang II-dependent hypertensive animal models, including

constriction of the renal artery and infusion of Ang II in rats and mice, have revealed a

progressive increase in intrarenal Ang II levels, surpassing what can be accounted for by

circulating Ang II levels alone[38]. This is due to Ang II-dependent renal activation of the

renin-angiotensin system, as indicated by increased urinary angiotensinogen (AGT)

secretion[38,39]. Importantly, the elevation of urinary AGT is also evident in patients with

various pathologies, including hypertension and CKD, implying that renal Ang II levels

increase even in individuals who do not exhibit elevated levels of circulating Ang II[21,39,40].

Taken together, the current finding of the synergistic effect of Ang II and oxLDL on AT1

activation in renal tissue is highly relevant for the development of kidney disease. RA

system inhibitors, ARB, and ACE inhibitors are prioritized therapies to prevent the

development of CKD with proteinuria in patients with hypertension[41]. In addition to the

well-established inhibitory effects of Ang II on the contraction of efferent arterioles[42], a

novel renal protective action of RA system inhibitors has been proposed. Particularly in

hypertension accompanied by dyslipidemia, RA system inhibitors may exhibit anti-

inflammatory, antifibrotic, and antioxidant effects in the kidneys by inhibiting the Gq

signaling pathway through the AT1-LOX-1 complex in renal tubular cells and fibroblasts.

Collectively, the current findings suggest that RA inhibitors can concomitantly reduce the

effect of increased renal Ang II and oxLDL levels on the development of CKD in patients

with hypertension and dyslipidemia, although direct evidence in clinical studies to

support this remains to be elucidated.

This study has several limitations as follows: 1) The kidney, a complex organ vital for

maintaining homeostasis, comprises a myriad of distinct cell types working in concert to

execute its multifaceted functions[43]. The experiments conducted in mice raised questions

regarding the specific cell types implicated in the synergistic effect of Ang II and oxLDL

within the LOX-1-AT1 complex in the kidney. Addressing this issue would ideally require

single-cell analysis, which is a challenge for future research. 2) The study employed

systemic LOX-1 knockout mice. For a more detailed analysis, a phenotypic investigation

using renal tissue-specific LOX-1 and AT1 knockout mice is required. Of particular

importance is the analysis using tubule-specific knockout mice, in which the localization

of these components has been verified through immunohistochemical staining. 3) The

administration of Ang II (both pressor and subpressor doses) and its combination with

HFD did not result in any histological changes in terms of fibrosis and mesangial

expansion, despite the observed alterations in the associated gene expression and

urinalysis for renal injury. Alterations in gene expression and urinalysis for renal injury

may be sensitive and relatively early phenomena, and this discrepancy could potentially

be attributed to the relatively short intervention duration of 4 weeks. Therefore,

concurrent pathohistological alterations in the renal tissue might become evident with a

more extended intervention period. 4) This study was limited by its inability to detect the

amplification of Gq signaling in mouse renal tissue due to the concurrent administration

of Ang II and HFD. Overcoming this limitation is a challenge for future studies.

In conclusion, the current findings suggest that the simultaneous binding of oxLDL and

Ang II to their respective receptors within the complex induces a distinct conformational

change compared with the effect of each ligand alone. This unique conformational change

results in the heightened activation of G protein signaling and subsequent unfavorable

cellular reactions in renal component cells. The relevance of this phenomenon was

confirmed in mouse models, in which renal dysfunction was prominently exacerbated

when there was a concomitant increase in Ang II and oxLDL levels. Notably, this effect

was abolished by the deletion of LOX-1, indicating LOX-1 dependency of this in vivo

phenomenon (Fig. 10). These findings indicate the clinical relevance of the direct

interaction between hypertension and dyslipidemia and further support the clinical

significance of RA inhibition in treating patients with CKD.

Materials and methods

Cell culture and materials

HUVECs and BAECs were cultured in EGM-2 (Lonza, Basel, Switherlad). Cells with

fewer than five passages were used in the experiments. Transgenic CHO cells were

maintained in an F-12 Nutrient Mixture with GlutamaxTM-I (Thermo Fisher Scientific,

MA, USA), 10% fetal bovine serum (FBS; Gibco, USA), and appropriate selection

reagents, as described below. CHO-K1 cells were maintained in F-12 Nutrient Mixture

with GlutamaxTM-I and 10% FBS. HAVSMCs were cultured in Dulbecco's modified

Eagle's medium/F12 (DMEM/F12) (Nacalai Tesque, Japan) supplemented with 1%

penicillin-streptomycin (Fujifilm, Japan) and 10% FBS. A10 cells were grown in DMEM

(Wako, Osaka, Japan) supplemented with 10% FBS and 1% penicillin-streptomycin.

NRK52E and NRK49F cells (ECACC, UK) were cultured in DMEM (Wako, Japan)

supplemented with 5% FBS and the appropriate selection reagents. Gene transcription in

CHO cells was induced by adding 100 ng/mL doxycycline (Merck KGaA, Darmstadt,

Germany). Cells were incubated at 37°C in 5% CO_2 and 95% air.

Construction of plasmid vectors

For stable transformants, pTRE2hyg vector (Clontech, USA) encoding mutated hAT1

with impaired ability to activate β-arrestin (pTRE2hyg-HA-FLAG-hAT1mg) were

created using site direct mutagenesis as previously described[14]. For real-time imaging,

LOX-1 tagged with V5-6×His at the C-terminus (V5-LOX-1) was subcloned into

pmScarlet_C1 (plasmid #85042; Addgene) (mScarlet-LOX-1). HA-FLAG-hAT1 was

subcloned into pcDNA3-EGFP (plasmid #85042; Addgene) (AT1-eGFP)[14].

Stable transformants

We constructed CHO-K1 cells expressing tetracycline-inducible human LOX-1 tagged

with V5-6×His at the C-terminus (CHO-LOX-1), human HA-FLAG-hAT1 (CHO-AT1),

or cells expressing both human LOX-1 and AT1 (CHO-LOX-1-AT1), as previously

described[13,14] To establish cells expressing both LOX-1 and mutated AT1 (pTRE2hyg-

HA-FLAG-hAT1mg), they were co-transfected with the pSV2bsr vector (Funakoshi,

Japan) into CHO-LOX-1 using the Lipofectamine 2000 transfection reagent (Thermo

Fisher Scientific, USA). The stable transformants were selected with 400 μg/mL of

hygromycin B (Wako, Osaka, Japan) and 10 μg/mL of blasticidin S (Funakoshi, Japan).

The resistant clone expressing LOX-1 and mutated AT1 in response to doxycycline

(Calbiochem, USA) was selected for use in the experiments (CHO-LOX-1-AT1mg) [14].

Small interfering RNA

NRK52 and NRK47 cells were plated at 50% confluence on the day of transfection.

Silencer Select small interfering RNA (siRNAs) for *LOX-1* and *AT1a* (Thermo Fisher

Scientific, MA, USA) were transfected into cells in medium without serum or antibiotics

using Lipofectamine RNAiMAX (Thermo Fisher Scientific, MA, USA), according to the

manufacturer's instructions.

Preparation of oxLDL

Human plasma LDL (1.019-1.063 g/mL), isolated by sequential ultracentrifugation,

was oxidized using 20 μM $CuSO_4$ in PBS at 37 °C for 24 h. Oxidation was terminated

by adding excess EDTA. LDL oxidation was analyzed by agarose gel

electrophoresis for migration versus LDL[13].

Quantification of cellular IP1 accumulation

Gq-dependent activation of phospholipase C was quantified by measuring IP1 using the

IP-One assay kit (Cisbio, France) as previously described[19]. Briefly, cells were seeded at

80,000 cells/well in 96 well transparent cell culture plates and incubated under serum-

free conditions for 24 h. Thereafter, cells were treated for 1 h with IP1 stimulation buffer,

including vehicle, native LDL, oxLDL, Ang II, AGE, PTX

(Merck KGaA, Darmstadt, Germany), YM-254890 (Fujifilm Wako, Osaka, Japan), and

RKI-1448 (Selleck, USA), as described in the text. Cell lysates with Triton X at a final

concentration of 1% were transferred to a 384-well white plate, and IP1 levels were

measured by incubating the cell lysates with FRET reagents (cryptate-labeled anti-IP1

antibody and d2-labeled IP1 analog).

Quantitative real-time PCR

RNA samples were purified using RNeasy Mini Kit (Qiagen, Germantown, MD, USA).

One microgram of RNA was converted into cDNA using a ReverTra Ace qPCR RT kit

(TOYOBO, Osaka, Japan) according to the manufacturer's instructions. All genes were

evaluated using the ViiA7 Real-Time PCR System (Applied Biosystems, Thermo Fisher

Scientific, Waltham, MA, USA). The data were analyzed using the ΔΔCt method with

normalization against the GAPDH RNA expression in each sample. The primer sequences

are listed in the Supplemental Table.

Western blotting

Proteins were separated using sodium dodecyl sulfate-polyacrylamide gel electrophoresis

and electrophoretically transferred onto polyvinylidene fluoride membranes. The

membranes were blocked with 5% nonfat dried milk and incubated with primary

antibodies overnight at 4 ºC. The primary antibodies used in this study were as follows:

anti-SMA antibody (1:1000), anti-α-Tubulin antibody (1:1000) (Cell Signaling

Technology, Inc, Danvers, MA, USA). The bands were visualized with a

chemiluminescence detection system (LAS-4000 mini; GE Healthcare Life Sciences,

Buckinghamshire, UK) using Chemi-Lumi One Super (Nacalai Tesque, Kyoto, Japan).

Calcium influx assay

Calcium influx was measured using Fura 2-AM (Dojindo, Kumamoto, Japan) with slight

modifications to the manufacturer's protocol. In brief, cells plated in 96 wells were

incubated with 5 μM Fura 2-AM in HEPES buffer saline (20 mM HEPES, 115 mM NaCl,

5.4 mM KCl, 0.8 mM MgCl2, 1.8 mM $CaCl_2$, 13.8 mM glucose, pH 7.4) for 1 h at 37°C,

followed by replacement with recording medium without Fura 2-AM. Cells were treated

with oxLDL, Ang II, or a combination of both at the indicated concentrations. Changes

in F340/F380, an index of intracellular calcium concentration, were measured by dual-

excitation microfluorometry using a digital image analyzer (Aquacosmos; Hamamatsu

Photonics, Hamamatsu, Japan).

BrdU assay

Proliferative activity was assessed using a BrdU Cell Proliferation ELISA Kit (Funakoshi,

Japan). NRK49F cells were seeded in 96-well tissue culture plates and incubated with the

test reagents for 24 h. After incubating the cells with BrdU, we fixed the cells and

denatured their DNA using a Fixing Solution. The plate was washed thrice with Wash

Buffer before adding the Detector Antibody. Next, 100 μL/well of anti-BrdU monoclonal

Detector Antibody was added, and the plate was incubated for 1 h at room temperature.

Subsequently, 100 μL/well of Goat Anti-Mouse IgG Conjugate was pipetted and

incubated for 30 minutes at room temperature. After five washes, the reaction was stopped

by adding Stop Solution to each well. The color of the positive wells changed from blue

to bright yellow. Finally, the plate was read at a wavelength of 450/550 nm using a

spectrophotometric microtiter plate reader.

Creation of lentivirus encoding AT1 conformational sensors

For lentivirus encoding AT1 conformational sensors, rat AT1 and RlucII were subcloned

into the pLVSIN-CMV Neo vector (Takara Bio, Japan). Next, the FlAsH binding

sequence (CCPGCC) was inserted between residues K135 and S136 in the third

intracellular loop (AT1-ICL3P3), as well as between residues K333 and M334 in the

cytoplasmic-terminal tail (AT1-Ctail), utilizing the KOD-Plus Mutagenesis Kit (Toyobo,

Japan), at the same site as previously documented[14].

Intramolecular FlAsH Bioluminescence resonance energy transfer assay to detect

AT1 conformational change

CHO-LOX-1 cells were initially plated onto a white clear-bottom 96-well culture plate at

a density of 1×10^5 cells/well. The following day, the cells were transduced with lenti-

virus encoding AT1-Ctail or AT1-ICL3P3 [20] in 10% FBS. After 24 h of transduction, the

cultures were transferred to serum-free conditions and incubated for an additional 24 h.

Add 1.5 µM FlAsH-EDT$_2$ labeling reagent of TC-FlAsH™ II In-Cell Tetracysteine Tag

Detection Kit (Thermo Fisher Scientific, MA, USA), wash twice with 250 µM BAL buffer,

and assays were promptly conducted on a Spark® microplate reader (TECAN,

Switzerland). The BRET ratio (emission mVenus/emission Rluc) was calculated as

follows: Following a 3-minute baseline reading (with the final baseline reading presented

at 0), cells were exposed to vehicle, oxLDL alone, AII alone, a combination of AII and

oxLDL, or a combination of AngII, oxLDL, and LOX-1 antibodies. The BRET ratios

were calculated every 16 seconds for a total of 320 seconds and the relative change in

intramolecular BRET ratio was calculated by subtracting the average BRET ratio

measured for cells stimulated with vehicle at each time point.

Analysis of LOX-1 and AT1 dynamics by real-time imaging

Live imaging was performed using previously reported methods[19]. Briefly, twenty-four

hours prior to imaging experiments, CHO-K1 cells were transfected with LOX-1-

mScarlet and AT1-eGFP by electroporation. Subsequently, the cells were seeded in a 35-

mm glass base dish (Iwaki, Japan) that had been pre-coated with a 1000X diluted solution

of 10 mg/mL poly-L-lysine (ScienCell, USA) one hour before seeding. The growth

medium was substituted with imaging buffer (pH 7.4), which consisted of 125 mM NaCl,

5 mM KCl, 1.2 mM $MgCl_2$, 1.3 mM $CaCl_2$, 25 mM HEPES, and 3 mM D-glucose, with

the pH adjusted to 7.4 using NaOH. Dynamic images of the cells were acquired at 25 °C

using a SpinSR10 inverted spinning disk-type confocal super-resolution microscope

(Olympus, Japan). The microscope was equipped with a 100x NA1.49 objective lens

(UAPON100XOTIRF, Olympus, Japan) and an ORCA-Flash 4.0 V2 scientific CMOS

camera (Hamamatsu Photonics KK, Japan) at 5-second intervals. The imaging

experiment was conducted using CellSens Dimension 1.11 software, employing a 3D

deconvolution algorithm (Olympus, Japan), and the number of puncta was determined

using ImageJ1.53K[19].

Animals and diets

Male WT mice (C57BL/6J) and LOX-1 KO mice with a C57BL/6 background were used

in this study. LOX-1 KO mice were generated as described previously[44]. Mice were

housed in a temperature-controlled (20-22°C) room on a 12 hours light/dark cycle and

fed an ND (MF; Oriental Yeast, Osaka, Japan) or an HFD (High Fat Diet without DL-α-

tocopherol, CLEA Japan Inc, Tokyo, Japan), which reported to increase plasma LOX-1

ligand in ApoE KO mice[21]. All study protocols were approved by the Animal Care and

Use Committee of Osaka University and were conducted according to the guidelines of

the NIH for the Care and Use of Laboratory Animals.

Blood pressure measurement in mice

642 The blood pressure of the mice was measured using the tail- cuff method with BP-98A

643 (Softron, Japan). The measurements were performed after restraining the mice. The blood

644 pressure was calculated as the average of 6 readings for each animal at each time point.

Urine tests in mice

Urine tests in mice included the measurement of urine 8-OHdG, creatinine, and albumin

concentrations. The DNA Damage (8-OHdG) ELISA Kit (StressMarq Bioscience,

Canada), Creatinine Kit L type Wako (Fujifilm, Japan), and Mouse Albumin ELISA Kit

(Bethyl Laboratories, Inc., TX, USA) were utilized for these measurements, following

their respective instructions.

Plasma LOX-1 ligand concentration

Measurement of LOX-1 ligands containing apoB (LAB) in mouse plasma was performed

using a modified protocol based on a previously reported method [33]. Briefly, recombinant

human LOX-1 (0.25 μg/well) was immobilized on 384 well plate (greiner, Frickenhausen,

Germany) by incubating overnight at 4°C in 50 μl of PBS. After three washes with PBS,

80 μl of 20% (v/v) ImmunoBlock (KAC, Kyoto, Japan) was added, and the plates were

incubated for 2 h at 25°C. After three washes with PBS, the plates were incubated for 2 h

at 25°C with 40 μl of standard oxidized LDL or samples. Samples were prepared by 4-

fold dilution of plasma with HEPES-EDTA buffer (10 mM HEPES, 150 mM NaCl, 2 mM

EDTA, pH 7.4), and standards were prepared by dilution of oxidized LDL with HEPES-

EDTA buffer. Following three washes with PBS, the plates were incubated for 1 h at 25°C

with chicken monoclonal anti-apoB antibody (HUC20, 0.5 μg/mL) in HEPES-EDTA

containing 1% (w/v) BSA. After three washes with PBS, the plates were incubated for 1

h at 25°C with peroxidase-conjugated donkey anti-chicken IgY (Merck, NJ, USA) diluted

5000 times with HEPES-EDTA containing 1% (w/v) BSA. After five washes with PBS,

the substrate solution containing 3,3',5,5'-tetramethylbenzidine (TMB solution, Bio-Rad

Laboratories, CA, USA) was added to the plates and incubated them for 30 min at room

temperature. The reaction was terminated with 2 M sulfuric acid. Peroxidase activity was

determined by measuring absorbance at 450 nm using a SpectraMax 340PC384

Microplate Reader (Molecular Devices, CA, USA).

Tissue preperation

Kidneys were perfused with cold PBS before removal. Kidney samples were rapidly

excised. A quarter of samples were stored at 4°C in RNAlater (Thermo Fisher Scientific,

MA, USA) for RNA extraction. The remaining quarters were fixed in 4%

paraformaldehyde overnight at 4°C for histological evaluation. The remaining half was

snap-frozen in liquid nitrogen and stored at -80°C for further analysis.

Periodic acid-Schiff and Masson-Trichrome staining

The degree of glomerular mesangial expansion and glomerular area (representing the

structural integrity of the glomeruli) were assessed in a blinded manner using periodic

acid-Schiff (PAS) staining. Collagen accumulation was determined by Masson-Trichrome

(MTC) staining. For MTC staining, the area displaying fibrosis was quantitatively

evaluated in a blinded manner by measuring the blue staining in six strongly magnified

fields of view using the ImageJ software, and the average was calculated after

determining the ratio of the total area.

Fluorescent immunostaining

For fluorescent immunostaining of LOX-1 and AT-1, the mice were perfused with cold

saline before tissue removal. After 3 days of zinc fixation, the tissue was replaced with

70% ethanol. The 3-μm-thick kidney tissue sections were immunohistochemically stained

with antibodies against ATGR (1:200, Cosmo Bio, Japan) and OLR-1 (1:200, TS58 from

the laboratory of T.S., Shinshu University School of Medicine, Nagano, Japan).

Following deparaffinization (using Lemosol and gradient ethanol) and rehydration, the

slices were subjected to antigen retrieval by autoclaving in citrate buffer (0.01 M; pH 6.0).

Subsequently, the slices were washed thrice with PBS and blocked with 5% bovine serum

albumin for 30 min at room temperature. The slides were then incubated with primary

antibodies for 2 h at room temperature. Goat anti-Rabbit IgG (H+L) High Cross-

Adsorbed Secondary Antibody, Alexa Fluor Plus 488 and 594 (Thermo Fisher Scientific,

MA, USA) were used as secondary antibodies for ATGR and OLR-1, respectively. After

incubation with secondary antibodies for 1 h at room temperature, the slices were washed

with PBS. Finally, slides were sealed and photographed. Visual analyses were performed

using a BZ-800L microscope (Keyence, Japan).

Statistical analyses

All data are presented as the mean ± SEM. Differences between two treatments or among

multiple treatments were determined using the Student's t-test or one-way ANOVA

followed by Tukey's multiple comparison test.

Data availability

All data supporting the findings of this study are available within the paper and its

Supplementary Information. Source data are provided in this paper.

Acknowledgements

This work was partially supported by JSPS KAKENHI Grant Numbers 21K07389 (Y.T.),

22K08181 (Y.N.), 20H03576 (H.R.), and 18H02732 (K.Y.). We are grateful to Tomoko

Sato, Yoshinori Koishi, and Chika Takana for technical assistance. We would like to thank

Editage (www.editage.com) for the English language editing.

Author contributions

Competing interests

734 The authors declare no competing interests.

References

References

1   Harper, C. R. & Jacobson, T. A. Managing dyslipidemia in chronic kidney disease.

J Am Coll Cardiol **51**, 2375-2384 (2008).

https://doi.org/10.1016/j.jacc.2008.03.025

2   Cases, A. & Coll, E. Dyslipidemia and the progression of renal disease in chronic

renal failure patients. *Kidney Int Suppl*, S87-93 (2005).

https://doi.org/10.1111/j.1523-1755.2005.09916.x

3   Moorhead, J. F., Chan, M. K., El-Nahas, M. & Varghese, Z. Lipid nephrotoxicity

in chronic progressive glomerular and tubulo-interstitial disease. *Lancet* **2**, 1309-

1311 (1982). https://doi.org/10.1016/s0140-6736(82)91513-6

4   Ruan, X. Z., Varghese, Z. & Moorhead, J. F. An update on the lipid nephrotoxicity

hypothesis. *Nat Rev Nephrol* **5**, 713-721 (2009).

https://doi.org/10.1038/nrneph.2009.184

5   Bussolati, B. *et al.* Statins prevent oxidized LDL-induced injury of glomerular

podocytes by activating the phosphatidylinositol 3-kinase/AKT-signaling

pathway. *J Am Soc Nephrol* **16**, 1936-1947 (2005).

https://doi.org/10.1681/ASN.2004080629

6   Dai, Y. *et al.* High fat diet causes renal fibrosis in LDLr-null mice through MAPK-

NF-kappaB pathway mediated by Ox-LDL. *J Cardiovasc Pharmacol* **63**, 158-166

(2014). https://doi.org/10.1097/FJC.000000000000035

7 Deng, S., Jin, T., Zhang, L., Bu, H. & Zhang, P. Mechanism of tacrolimus-induced

chronic renal fibrosis following transplantation is regulated by ox-LDL and its

receptor, LOX-1. *Mol Med Rep* **14**, 4124-4134 (2016).

https://doi.org/10.3892/mmr.2016.5735

8 Gai, Z. *et al.* Lipid Accumulation and Chronic Kidney Disease. *Nutrients* **11**

(2019). https://doi.org/10.3390/nu11040722

9 Nagase, M. *et al.* Expression of LOX-1, an oxidized low-density lipoprotein

receptor, in experimental hypertensive glomerulosclerosis. *J Am Soc Nephrol* **11**,

1826-1836 (2000). https://doi.org/10.1681/ASN.V11101826

10 Lu, J. *et al.* Abrogation of lectin-like oxidized LDL receptor-1 attenuates acute

myocardial ischemia-induced renal dysfunction by modulating systemic and local

inflammation. *Kidney Int* **82**, 436-444 (2012). https://doi.org/10.1038/ki.2012.186

11 Manttari, M., Tiula, E., Alikoski, T. & Manninen, V. Effects of hypertension and

dyslipidemia on the decline in renal function. *Hypertension* **26**, 670-675 (1995).

https://doi.org/10.1161/01.hyp.26.4.670

12 Brewster, U. C. & Perazella, M. A. The renin-angiotensin-aldosterone system and

the kidney: effects on kidney disease. *Am J Med* **116**, 263-272 (2004).

https://doi.org/10.1016/j.amjmed.2003.09.034

13   Yamamoto, K. *et al.* Oxidized LDL (oxLDL) activates the angiotensin II type 1

receptor by binding to the lectin-like oxLDL receptor. *FASEB J* **29**, 3342-3356

(2015). https://doi.org/10.1096/fj.15-271627

14   Takahashi, T. *et al.* The endocytosis of oxidized LDL via the activation of the

angiotensin II type 1 receptor. *iScience* **24**, 102076 (2021).

https://doi.org/10.1016/j.isci.2021.102076

15   Wingler, L. M. *et al.* Angiotensin Analogs with Divergent Bias Stabilize Distinct

Receptor Conformations. *Cell* **176**, 468-478 e411 (2019).

https://doi.org/10.1016/j.cell.2018.12.005

16   Namkung, Y. *et al.* Functional selectivity profiling of the angiotensin II type 1

receptor using pathway-wide BRET signaling sensors. *Sci Signal* **11** (2018).

https://doi.org/10.1126/scisignal.aat1631

17   Jono, T. *et al.* Lectin-like oxidized low density lipoprotein receptor-1 (LOX-1)

serves as an endothelial receptor for advanced glycation end products (AGE).

FEBS Lett **511**, 170-174 (2002). https://doi.org/10.1016/s0014-5793(01)03325-7

18   Lymperopoulos, A., Borges, J. I., Carbone, A. M., Cora, N. & Sizova, A.

791  Cardiovascular angiotensin II type 1 receptor biased signaling: Focus on non-Gq-,

792  non-betaarrestin-dependent signaling. *Pharmacol Res* **174**, 105943 (2021).

793  https://doi.org/10.1016/j.phrs.2021.105943

794 19 Huang, Y., Takahashi, T., Gaisano, H., Rakugi, H. & Yamamoto, K. A live-

795  imaging protocol for tracking receptor dynamics in single cells. *STAR Protoc* **3**,

796  101347 (2022). https://doi.org/10.1016/j.xpro.2022.101347

797 20 Devost, D. *et al.* Conformational Profiling of the AT1 Angiotensin II Receptor

798  Reflects Biased Agonism, G Protein Coupling, and Cellular Context. *J Biol Chem*

799  **292**, 5443-5456 (2017). https://doi.org/10.1074/jbc.M116.763854

800 21 Kobori, H. *et al.* Urinary angiotensinogen as a novel biomarker of the intrarenal

801  renin-angiotensin system status in hypertensive patients. *Hypertension* **53**, 344-

802  350 (2009). https://doi.org/10.1161/HYPERTENSIONAHA.108.123802

803 22 Jennings, B. L. *et al.* Cytochrome P450 1B1 contributes to renal dysfunction and

804  damage caused by angiotensin II in mice. *Hypertension* **59**, 348-354 (2012).

805  https://doi.org/10.1161/HYPERTENSIONAHA.111.183301

806 23 Wolak, T. *et al.* Osteopontin modulates angiotensin II-induced inflammation,

807  oxidative stress, and fibrosis of the kidney. *Kidney Int* **76**, 32-43 (2009).

808  https://doi.org/10.1038/ki.2009.90

24   Hu, C. *et al.* Modulation of angiotensin II-mediated hypertension and cardiac

remodeling by lectin-like oxidized low-density lipoprotein receptor-1 deletion.

Hypertension **52**, 556-562 (2008).

https://doi.org/10.1161/HYPERTENSIONAHA.108.115287

25   Ohki, I. *et al.* Crystal structure of human lectin-like, oxidized low-density

lipoprotein receptor 1 ligand binding domain and its ligand recognition mode to

OxLDL. *Structure* **13**, 905-917 (2005). https://doi.org/10.1016/j.str.2005.03.016

26   Wright, A. K. & Thompson, M. R. Hydrodynamic structure of bovine serum

albumin determined by transient electric birefringence. *Biophys J* **15**, 137-141

(1975). https://doi.org/10.1016/s0006-3495(75)85797-3

27   Kriz, W., Kaissling, B. & Le Hir, M. Epithelial-mesenchymal transition (EMT) in

kidney fibrosis: fact or fantasy? *J Clin Invest* **121**, 468-474 (2011).

https://doi.org/10.1172/jci44595

28   Jiang, T. *et al.* Diet-induced obesity in C57BL/6J mice causes increased renal lipid

accumulation and glomerulosclerosis via a sterol regulatory element-binding

protein-1c-dependent pathway. *J Biol Chem* **280**, 32317-32325 (2005).

https://doi.org/10.1074/jbc.M500801200

29   Kume, S. *et al.* Role of altered renal lipid metabolism in the development of renal

injury induced by a high-fat diet. *J Am Soc Nephrol* **18**, 2715-2723 (2007).

https://doi.org/10.1681/ASN.2007010089

30   Yamamoto, T. *et al.* High-Fat Diet-Induced Lysosomal Dysfunction and Impaired

Autophagic Flux Contribute to Lipotoxicity in the Kidney. *J Am Soc Nephrol* **28**,

1534-1551 (2017). https://doi.org/10.1681/ASN.2016070731

31   Sun, Y. *et al.* High-fat diet promotes renal injury by inducing oxidative stress and

mitochondrial   dysfunction.   *Cell   Death   Dis*   **11**,   914   (2020).

https://doi.org/10.1038/s41419-020-03122-4

32   Yu, Y., Mo, H., Zhuo, H., Yu, C. & Liu, Y. High Fat Diet Induces Kidney Injury

via Stimulating Wnt/beta-Catenin Signaling. *Front Med (Lausanne)* **9**, 851618

(2022). https://doi.org/10.3389/fmed.2022.851618

33   Sato, Y. *et al.* Determination of LOX-1-ligand activity in mouse plasma with a

chicken monoclonal antibody for ApoB. *Atherosclerosis* **200**, 303-309 (2008).

https://doi.org/10.1016/j.atherosclerosis.2008.02.001

34   Garcia-Carro, C. *et al.* A Nephrologist Perspective on Obesity: From Kidney

Injury to Clinical Management. *Front Med (Lausanne)* **8**, 655871 (2021).

https://doi.org/10.3389/fmed.2021.655871

35   Tsuboi, N., Okabayashi, Y., Shimizu, A. & Yokoo, T. The Renal Pathology of

Obesity. *Kidney Int Rep* **2**, 251-260 (2017).

https://doi.org/10.1016/j.ekir.2017.01.007

36   Hu, C. *et al.* Deletion of LOX-1 attenuates renal injury following angiotensin II

infusion. *Kidney Int* **76**, 521-527 (2009). https://doi.org/10.1038/ki.2009.234

37   Li, X. *et al.* LOX-1 Deletion Attenuates Myocardial Fibrosis in the Aged Mice,

Particularly Those With Hypertension. *Front Cardiovasc Med* **8**, 736215 (2021).

https://doi.org/10.3389/fcvm.2021.736215

38   Navar, L. G., Kobori, H., Prieto, M. C. & Gonzalez-Villalobos, R. A. Intratubular

renin-angiotensin system in hypertension. *Hypertension* **57**, 355-362 (2011).

https://doi.org/10.1161/HYPERTENSIONAHA.110.163519

39   Navar, L. G. Translational studies on augmentation of intratubular renin-

angiotensin system in hypertension. *Kidney Int Suppl (2011)* **3**, 321-325 (2013).

https://doi.org/10.1038/kisup.2013.67

40   Mills, K. T. *et al.* Increased urinary excretion of angiotensinogen is associated

with risk of chronic kidney disease. *Nephrol Dial Transplant* **27**, 3176-3181

(2012). https://doi.org/10.1093/ndt/gfs011

41   Kidney Disease: Improving Global Outcomes Blood Pressure Work, G. KDIGO

2021 Clinical Practice Guideline for the Management of Blood Pressure in

863   Chronic Kidney Disease. *Kidney Int* **99**, S1-S87 (2021).

864   https://doi.org/10.1016/j.kint.2020.11.003

865   42   Yang, R., Smolders, I. & Dupont, A. G. Blood pressure and renal hemodynamic

866   effects of angiotensin fragments. *Hypertens Res* **34**, 674-683 (2011).

867   https://doi.org/10.1038/hr.2011.24

868   43   Balzer, M. S., Rohacs, T. & Susztak, K. How Many Cell Types Are in the Kidney

869   and What Do They Do? *Annu Rev Physiol* **84**, 507-531 (2022).

870   https://doi.org/10.1146/annurev-physiol-052521-121841

871   44   Mehta, J. L. *et al.* Deletion of LOX-1 reduces atherogenesis in LDLR knockout

872   mice fed high cholesterol diet. *Circ Res* **100**, 1634-1642 (2007).

873   https://doi.org/10.1161/CIRCRESAHA.107.149724

 Figure Legends

Fig. 1| Oxidized LDL potentiates Ang II-induced Gq signaling and calcium influx in

a LOX-1-dependent manner

a Dose-dependent response of IP1 concentration by the activation of Gq signaling in

response to oxLDL and Ang II in CHO-LOX-1-AT1 cells. Cells were treated with oxLDL

and Ang II at the concentrations described in the figure. (n=4 for each oxLDL

concentration). $^{*}P$=0.0004 for 0 μg/mL vs 20 μg/mL, $^{\dagger}P$=0.0003 for 5 g/mL vs 20 μg/mL,

$^{\ddagger}P$=0.0020 for 0 μg/mL vs 10 μg/mL, and $^{\S}P$=0.0015 for 5 μg/mL vs 10 μg/mL at 10^{-9} M

Ang II; $^{\parallel}P$=0.0051 for 0 μg/mL vs 10 μg/mL, $^{\P}P$=0.0004 for 0 μg/mL vs 20 μg/mL at 10^{-}

^{8} M Ang II.

b IP1 concentration in response to vehicle, native LDL (nLDL 10 μg/mL), and oxLDL

(10 μg/mL) in the combination of Ang II (10^{-8} M) in CHO-LOX-1-AT1 cells (n=5 for

each group)

c IP1 concentration in response to vehicle, oxLDL (10 μg/mL) in the combination of Ang

II (10^{-8}M) in CHO-LOX-1-AT1 and CHO-AT1 cells (n=5 for each group).

d IP1 concentration in response to vehicle, oxLDL (10 μg/mL), BSA (10 or 100 μg/mL),

BSA-conjugated AGE (10 or 100 μg/mL) in the combination of Ang ll (10^{-8} M) in CHO-

LOX-1-AT1 cells.

e IP1 concentration in response to vehicle, oxLDL (10 μg/mL) in the combination of Ang

II (10^{-8} M) in genetically engineered CHO cells with or without intact β-arrestin binding

domain (n=5 for each group). AT1mg indicates AT1 a mutant AT1 lacking a functional β-

arrestin binding domain but retaining G-protein-biased signaling capability.

f IP1 concentration in response to oxLDL (10 μg/mL) in the combination of Ang II (10^{-8}

M) and additional effect of PTX, a Gi inhibitor, YM-254890, a Gq inhibitor, and RKI-

1448, a downstream Rho kinase inhibitor targeting G12/13 signaling, in CHO-LOX-1-

AT1 cells (n=5 for each group).

g Intracellular calcium dynamics measured using Fura 2-AM by the ratio of the mission

signals at excitation wavelength 340 nm and 380 nm in response to oxLDL (5 μg/mL),

Ang II (10^{-12} M), and their combination (for each agonist, 4-7 regions of interest were

selected). Addition of these agonists is marked with arrows on the timeline of the assay.

h Percentage changes from baseline in the ratio of emission signals (F340/F380)

measured by Fura 2-AM were quantified following treatment with oxLDL and Ang II at

specified concentrations in CHO-LOX-1-AT1 cells as detailed in the figure (n=4-8).

i Percentage change from baseline in the ratio of emission signals (F340/F380) measured

by Fura 2-AM after stimulation with oxLDL (5 μg/mL), Ang II (10^{-12} M), and YM-

254890, a Gq inhibitor, in CHO-LOX-1-AT1 cells.

j Percentage changes from baseline in the ratio of emission signals (F340/F380) measured

by Fura 2-AM were quantified following treatment with oxLDL (5 μg/mL) and Ang II at

specified concentrations in CHO-AT1 cells, as detailed in the figure.

Data are represented as mean ± SEM. Differences were determined using one-way

ANOVA, followed by Tukey's multiple comparison test for (a-f) and (h-j).

Fig. 2| Co-treatment of oxLDL with Ang II induces conformational change of AT1

different from each treatment alone

a Change in green puncta (AT1-eGFP) and red puncta (LOX-1-mScarlet) by the treatment

with vehicle, oxLDL (10μg/mL), Ang II (10^{-5}M), and the combination of oxLDL and Ang

II in CHO cell overexpressing these fluorescent protein-conjugated receptors (n=10-11

for each group). The puncta were manually counted by a blinded observer and the number

of puncta at 0 and 3 min was determined (N0 and N3, respectively). The change in puncta

was calculated as (N0-N3)/N0.

b-d Changes in BRET signals were monitored in CHO-LOX-1 cells expressing the

following conformational biosensors: AT1-C-tailP1 (b) and AT1-ICL3P3 (c, d) bearing

FlAsH insertion at the cytoplasmic-terminal tail (C-tailP1) and the third intracellular loop

(ICL3P3) of AT1, respectively, that interact with Renilla luciferase at the end of the

cytoplasmic tail. Cells were subjected to treatments with vehicle, oxLDL (10 μg/mL),

Ang II (10^{-5}M), the combination of Ang II and oxLDL, and the combination

of AngII, oxLDL and LOX-1 antibody. The BRET ratios were calculated every 16

seconds for a total of 320 seconds and the relative change in intramolecular BRET ratio

(ΔBRET) was calculated by subtracting the average BRET ratio measured for cells

stimulated with vehicle at each time point. Lower panels indicate average ΔBRET of all

the time points during measurement.

Data are represented as mean ± SEM. Differences were determined by one-way ANOVA,

followed by Tukey's multiple comparison test (a-d).

Fig. 3| Oxidized LDL potentiates Ang II-induced Gq-calcium signaling in renal cells

a, b IP1 concentration in response to Ang II (10^{-7} M), oxLDL (10 μg/mL), and the

combination of both with or without YM-254890, Gq inhibitor, in NRK52E (a) and

NRK49F cells (b) (n=5 for each group).

c, d IP1 concentration in response to Ang II (10^{-7} M) and oxLDL (10 μg/mL) in the

combination of Ang II (10^{-7} M) and the additional effect of siRNA-mediated knockdown

of *AT1a* or *LOX-1* in NRK52E (c) and NRK49F cells (d) (n=5 for each group).

e Intracellular calcium concentration in NRK49F cells using Fura 2-AM and dual-

excitation microfluorometry. Changes in the fluorescence intensity ratio (F340/F380)

served as an index of the calcium dynamics. Cells were exposed to Ang II (10^{-7} M),

oxLDL (2 μg/mL), and a combination of both agents. Addition of these agonists is marked

with arrows on the timeline of the assay. Data acquisition and analysis were performed

using a digital image analyzer to monitor real-time calcium flux (n=4-9).

f Percentage changes from baseline in the ratio of emission signals (F340/F380) measured

by Fura 2-AM were quantified following treatment with Ang II (10^{-7} M) and oxLDL at

the concentrations detailed in the figure in NRK49F cells (n=5 for each group).

g Impact of siRNA-mediated knockdown *AT1* or *LOX-1* on the percentage changes from

baseline in the ratio of emission signals (F340/F380) measured by Fura 2-AM were

quantified following treatment with Ang II (10^{-7} M) and oxLDL (2 μg/mL) in NRK49F

cells (n=3-7).

g Impact of co-treatment of YM-254890, Gq inhibitor, or Irbesartan, an angiotensin

receptor blocker (ARB), on the the percentage changes from baseline in the ratio of

emission signals (F340/F380) measured by Fura 2-AM were quantified following

treatment with Ang II (10^{-7} M) and oxLDL (2 μg/mL) in NRK49F cells (n=4 for each

group).

Data are represented as mean ± SEM. Differences were determined using one-way

ANOVA, followed by Tukey's multiple comparison test for (a-d) and (f-h).

Fig. 4| Co-treatment of oxLDL and AII enhanced cellular response upon Gq

activation in renal cells

a,b Quantitative real-time PCR analysis was performed to measure the gene expression

of NADPH oxidase subunits (*p67phox* and *p91phox*), fibrosis markers (*fibronectin*,

collagen-1, *collagen-4*, and *TGFβ*), and inflammatory cytokines (*TNFα*, *IL1β*, *IL-6*, and

MCP-1) in NRK49F cells (a) and NRK 52E cells (b). Gene expression levels were

normalized to those of GAPDH. Cells were stimulated by oxLDL (5 μg/mL), Ang II (10^{-}

^{7} M) or their combination (n=4 for each group).

c, d Cells were pre-treated with vehicle or Gq inhibitor (YM-254890, Gqi), followed by

treatment with vehicle or the combination of oxLDL (5 μg/mL) and Ang II (10^{-7} M) in

NRK49F cells (c) and NRK 52E cells (d) (n=4 for each group).

e, f Cells were pre-treated with vehicle or ARB (Irbesartan, Arb), followed by treatment

with vehicle or the combination of oxLDL (5 μg/mL) and Ang II (10^{-7} M) in NRK49F

cells (e) and NRK 52E cells (f) (n=4 for each group).

Data are represented as mean ± SEM. Differences were determined by one-way ANOVA,

followed by Tukey's multiple comparison test (a-f).

Fig. 5| Oxidized LDL enhanced Ang II-induced epithelial mesenchymal transition in

NRK52E and NRK49F cells

a, b Left: Western blot analysis of α-smooth muscle actin (α-SMA), a marker of

epithelial–mesenchymal transition (EMT), in NRK49F (a) and NRK52E (b) cells. Cells

were stimulated with oxLDL (5 μg/mL), Ang II (10^{-7} M), and TGF-β (10 ng/mL), with

TGF-β serving as a well-known EMT inducer. Right: Densitometric analysis of α-SMA

protein expression normalized to α-Tubulin (n=3 for each group).

c, d Left: Western blot analysis of α-SMA in NRK49F (c) or NRK52E (d) after stimulation

with oxLDL (5 μg/mL), Ang II (10^{-7} M), and their combination. Right: Densitometric

analysis of α-SMA protein expression normalized to α-tubulin (n=3 for each group).

e, f Left: Western blot analysis of α-SMA in NRK49F(e) or NRK52E (f) after treatment

with a combination of oxLDL (5 μg/mL) and Ang II (10^{-7} M). Prior to this treatment, the

cells were pre-treated with either a vehicle or a Gq inhibitor (YM-254890, Gqi). Right:

Densitometric analysis of α-SMA protein expression normalized to α-Tubulin (n=3 for

each group).

g, h Left: Western blot analysis of α-SMA in NRK49F (e) or NRK52E (f) after treatment

with a combination of oxLDL (5 μg/mL) and Ang II (10^{-7} M). Prior to treatment, cells

1008   were pre-treated with either vehicle or ARB (Irbesartan, Arb). Right: Densitometric

1009   analysis of α-SMA protein expression normalized to α-tubulin (n=3 for each group).

1010   Data are represented as mean ± SEM. Differences were determined by one-way ANOVA,

1011   followed by Tukey's multiple comparison test (a-f).

1012

Fig. 6| Oxidized LDL enhanced Ang II-induced renal fibroblast proliferation via

AT1-Gq signaling and LOX-1-dependent manner

a Proliferative activity assessed by BrdU incorporation into NRK49F cells. Cells were

pretreated with vehicle, YM-254890, or ARB, Irbesartan, followed by the treatment with

oxLDL (5 μg/mL), Ang II (10^{-7} M), or their combination. (n=5 for each group).

b NRK49F cells were subjected to siRNA-mediated knockdown using specific siRNAs

for *AT1a* (siAT1) or *LOX-1* (siLOX-1). Following knockdown, cells were treated with

either vehicle, oxLDL (5 μg/mL), Ang II (10^{-7} M), or their combination. Proliferative

activity was assessed by measuring the BrdU levels (n=5 for each group).

Data are represented as mean ± SEM. Differences were determined using one-way

ANOVA, followed by Tukey's multiple comparison test for (a) and (b).

1024

Fig. 7| Oxidized LDL-inducible diet exacerbates Ang II-induced renal dysfuntion in

wildtype mice, but not in LOX-1 knockout mice

a A Schematic protocol for the animal experiments. Eight-week-old male WT mice and

male LOX-1 KO mice were fed either an ND or an HFD for 6 weeks. After 10 weeks of

age, the mice were treated over a 4-week period with infusions of either saline or Ang II.

Ang II was administered at two dosage levels: a subpressor dose of 0.1 γ and a pressor

dose of 0.7 γ, delivered via subcutaneously implanted osmotic pumps. At the end of the

infusion period, urine was collected, the animals were sacrificed, and comprehensive

tissue analysis was conducted to evaluate the renal effects of the treatments.

b Average systolic blood pressure (SBP) measured at half-week intervals in WT and

LOX-1 KO mice during the 4-week infusion period.

c, d Urine 8-OHDG concentrations (mg/g creatinine [Cr]) (c) and urine albumin

concentrations (mg/g creatinine [Cr]) (d) in WT and LOX-1 KO mice at the conclusion

of the 4-week infusion period.

Data are represented as mean ± SEM. Differences were determined by one-way ANOVA,

followed by Tukey's multiple comparison test (a-d).

1041

1042

1043 **Fig. 8| A high-fat diet enhanced Ang II-induced renal injury-related gene expression**

in the kidney in a LOX-1-dependent manner.

Quantitative real-time PCR analysis for gene expression of NADPH components

(*p67phox and p91phox*), inflammatory cytokines (*IL-6, TNFα, IL1β, and MCP-1*), and

fibrosis markers (*TGFβ, fibronectin, collagen-1a, and collagen-4a*) in the kidney

harvested from WT and LOX-1 KO mice.

The experimental procedures, including the dietary regimen and Ang II administration,

are detailed in Fig. 7a.

Data are represented as mean ± SEM. Differences were determined using one-way

ANOVA, followed by Tukey's multiple comparison test for (a) and (b).

Fig. 9| LOX-1 and AT1a were predominantly co-localized in renal tubules

a, b Representative images depicting staining for LOX-1 (a) and AT1 (b) in the renal

cortex tissues from wildtype mice (WT) and LOX-1 knockout mice (LOX-1 KO). Nuclei

are stained blue with DAPI. Green and red signals indicates AT1, while the red signal

indicates LOX-1. Overlay images demonstrate the merged visualization of AT1 or LOX-

1 with DAPI, highlighting the predominant colocalization of LOX-1 and AT1 in renal

tubules as opposed to the glomerulus.

Fig. 10| Schematic overview of the AT1 and LOX-1 Interaction dynamics in renal

cells

This schematic summary illustrates the predicted structure-activation relationship of the

AT1 receptor within the LOX-1-AT1 complex in renal component cells. This highlights

how the simultaneous binding of Ang II to AT1 and oxLDL to LOX-1 induces

conformational changes in AT1. These changes were more pronounced than those

triggered by the individual ligands. Such structural alterations have been proposed to

amplify Gq signaling pathway activation, subsequently leading to renal damage.

1070

1071

1072

1073

1074

1075

1076

1077

1078

1079

Supplemental Figure Legends

Fig. S1| Live-imaging analysis of membrane LOX-1 and AT1 in response to the co-treatment of oxLDL with AngII.

Real-time membrane imaging of CHO cells co-transfected with LOX-1-mScarlet and AT1-eGFP in response to oxLDL (10μg/ml) in the combination of Ang II (10^{-7}M) (Supplemental Video). A count of puncta was performed using separate images visualizing LOX-1-mScarlet (red puncta) and AT1-eGFP (green puncta) immediately before and 3 min after ligand application.

Fig. S2| Oxidized LDL in combination with Ang II do not increase cellular IP1 content in human umbilical vein endothelial cells and bovine vascular endothelial cells, human aortic vascular smooth muscle cells, and rat macrophages

IP1 concentration in response to oxLDL (10μg/ml) in the combination of Ang II (10^{-7}M) in HUVECs (human umbilical vein endothelial cell), BAECs (bovine aortic endothelial cell), HAVSMCs (human aortic vascular smooth muscle cell), and A10 cells (rat macrophages).

Data are represented as mean ± SEM. Differences were determined using one-way ANOVA, followed by Tukey's multiple comparison test (n=5 for each group).

1098

Fig. S3| Efficiency of siRNA-mediated knockdown for *AT1a* and *LOX-1* in NRK52E

and NRK49F cells

NRK52E and NRK49F cells were transfected with siRNAs against scrambled siRNA,

AT1a or *LOX-1*. The efficiency of siRNA-mediated gene silencing was quantified by

assessing *AT1a* and *LOX-1* expression levels using quantitative real-time PCR.

Data are represented as mean ± SEM. Differences were determined using one-way

ANOVA, followed by Tukey's multiple comparison test.

1106

Fig. S4| Calcium influx was not induced by either the combination treatment of Ang

II or oxLDL or each treatment alone in NRK52E cells

Percentage changes from baseline in the ratio of emission signals (F340/F380) measured

by Fura 2-AM were quantified following treatment with Ang II (10^{-7}M) and oxLDL at

the concentrations detailed in the figure for NRK49E cells (n=5-7 for each group).

Data are represented as mean ± SEM. Differences were determined using one-way

ANOVA, followed by Tukey's multiple comparison test.

1114

1115 **Fig. S5| High Fat Diet used in the study prominently increased plasma LOX-1 ligand**

concentration

Plasma LOX-1 ligand concentration of 14-week-old wildtype mice upon ND and HFD

for 6 weeks. From 10 weeks of age, these mice also received concurrent 4-week infusions

of Ang II at a pressor dose of 0.7 γ, administered through subcutaneously implanted

osmotic pumps.

Data are represented as mean ± SEM. Differences were determined using Student's t-test

(n=7 for each group).

Fig. S6 | Impact of Diet and Ang II Infusion on Body Weight and Systolic Blood Pressure in Mice

a Final Body weight: This figure shows the body weights of WT and LOX-1 KO mice at

the end of the 4-week infusion period, highlighting the effects of diet and pharmacological

treatment.

b, c Serial Body Weight Changes: These graphs depict the progression of body weight

over time in WT and LOX-1 KO mice, illustrating the impact of the dietary regimen and

Ang II infusion on weight dynamics.

d, e Systolic Blood Pressure Trajectory: Serial measurements of systolic blood pressure

(SBP) obtained using the tail-cuff method are presented for WT and LOX-1 KO mice.

The figures show the changes in SBP over the course of the study, corresponding to the

administration of a pressor dose of 0.7 γ (d) and a subpressor dose of 0.1 γ (e) of Ang II,

respectively.

f Final SBP: This figure shows SBP of WT and LOX-1 KO mice at the end of the 4-week

infusion period, highlighting the effects of diet and pharmacological treatments.

Beginning at 8 weeks of age, mice were fed either an ND or an HFD for 6 weeks. From

10 weeks of age, coinciding with the 2-week time point in the figure, the rats underwent

a 4-week period of infusion with either vehicle or Ang II. The infusion was delivered at

specific dosage levels through subcutaneously implanted osmotic pumps.

Data are represented as mean ± SEM. Differences were determined using one-way

ANOVA, followed by Tukey's multiple comparison test for (a) and (b).

1145

Fig. S7| No significant difference was found in plasma aldosterone concentration

between a normal diet and a high fat diet-fed wildtype mice with a pressor dose of

Ang II

Plasma aldosterone concentration in 14-week-old wild-type mice fed an ND or an HFD

for 6 weeks. From 10 weeks of age, these mice also received concurrent 4-week infusions

of Ang II at a pressor dose of 0.7 γ, administered through subcutaneously implanted

osmotic pumps.

Data are represented as mean ± SEM. Differences were determined using Student's t-test

(n=7 for each group).

1155

Fig. S8| A high-fat diet enhanced Ang II-induced renal injury-related gene

expression in the kidney in a LOX-1-dependent manner.

Quantitative real-time PCR analysis for gene expression of NADPH components

(*p40phox and p47phox*), inflammatory gene (*COX-2*), fibrosis markers (*αSMA and*

vimentin), epithelial markers (*E-cadherin and cadherin-16*), tubular marker (*NAGL*),

AT1a, *AT1b*, and *LOX-1* in the kidney harvested from WT and LOX-1 KO mice.

The experimental procedures, including the dietary regimen and Ang II administration,

are detailed in Fig. 7a.

Data are represented as mean ± SEM. Differences were determined using one-way

ANOVA, followed by Tukey's multiple comparison test for (a) and (b).

1166

Fig. S9| The treatment with Ang II, a High Fat Diet, or their combination for 4 weeks

did not induce any histological changes indicative of renal injury

a Left: Representative histological images of Masson-Trichrome staining used to detect

fibrosis in renal tissues harvested from WT and LOX-1 KO mice. Right: Quantitative

analysis by Masson-Trichrome staining.

Data are represented as mean ± SEM. Differences were determined using one-way

ANOVA, followed by Tukey's multiple comparison test.

b Representative histological images of renal tissues harvested from WT and LOX-1 KO

mice stained for PAS to assess mesangial expansion and glomerular area, providing

insight into the structural integrity of the glomeruli.

Eight-week-old mice were fed an ND or an HFD for 6 weeks. From 10 weeks of age,

these mice were concurrently treated for 4 weeks with infusions of vehicle or Ang II (a

subpressor dose of 0.1 γ or a pressor dose of 0.7 γ) through subcutaneously implanted

osmotic pomps.

1181

1182

1183

1184

1185

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6

6a

6b

Fig. 7

7a

WT
- saline
- Ang II 0.1γ
- Ang II 0.7γ

ND or HFD

LOX-1 KO
- saline
- Ang II 0.1γ
- Ang II 0.7γ

ND or HFD

8 weeks of age

Ang II infusion by Alzet® 2004

Urine analysis
Sacrifice
Tissue analysis

0   2   6 (weeks)

7b

Average SBP during Ang II infusion (mmHg)

Ang II 0.7γ

WT: P<0.0001, P<0.0001
LOX-1 KO: P=0.0431

Ang II 0.1γ

WT: P=0.0260

ND+saline, HFD+saline, ND+Ang II, HFD+Ang II

7c

Urine 8-ohdg concentration (mg/gCr)

Ang II 0.7γ

WT: P<0.0001, P<0.0001, P<0.0001
LOX-1 KO: P=0.0193, P=0.0124

Ang II 0.1γ

LOX-1 KO: P=0.0158, P=0.0119

ND+saline, HFD+saline, ND+Ang II, HFD+Ang II

7d

Urine albumin (mg/gCr)

Ang II 0.7γ

WT: P<0.0001, P=0.0005, P=0.0082

Ang II 0.1γ

WT: P<0.0001, P<0.0001, P=0.0120
LOX-1 KO: P<0.0001, P<0001

ND+saline, HFD+saline, ND+Ang II, HFD+Ang II

Fig. 8

Fig. 9

9a

9b

Fig. 10

Fig. S1

AT1-eGFP LOX-1-mScarlet

N=283 N=185

5μm

N=283 N=185

1min

N=167 N=125

N=167 N=125

4min

00:00

Fig. S2

Fig. S3

Fig. S4

Fig. S5

Fig. S6

Fig. S7

Fig. S8

Fig. S9

S9a
Masson-Trichrome staining

S9b
PAS staining