**Oxidized low-density lipoprotein potentiates angiotensin II-induced Gq activation**

**through the AT1-LOX1 receptor complex：Implications for renal dysfunction**

Jittoku Ihara [1]#, Yibin Huang [1)2)#], Yoichi Takami [1)*], Yoichi Nozato [1)*], Toshimasa

Takahashi [1)3)*], Akemi Kakino [4], Cheng Wang [1], Ziwei Wang [1], Yu Guo [1], Weidong

Liu [1], Nanxiang Yin [1], Ryoichi Ohara [1], Taku Fujimoto [1], Shino Yoshida [1], Kazuhiro

Hongyo [1], Hiroshi Koriyama [1], Hiroshi Akasaka [1], Hikari Takeshita [1], Shinsuke Sakai

[5], Kazunori Inoue [5], Yoshitaka Isaka [5], Hiromi Rakugi [1], Tatsuya Sawamura [4], and

Koichi Yamamoto [1]

[1] Department of Geriatric and General Medicine, Osaka University Graduate School of

Medicine, 2-2 Yamada-oka, Suita, Osaka, 565-0871, Japan

[2] Center for Pulmonary and Vascular Biology, Dept. of Pediatrics, University of Texas

Southwestern Medical Center, 5323 Harry Hines Blvd., Dallas, TX, 75390, USA

[3] Department of Medicine, Takahashi clinic, Kobe, Hyogo, 657-0846, Japan

[4] Department of Molecular Pathophysiology, Shinshu University Graduate School of

Medicine, Matsumoto, Nagano 390-8621, Japan

[5] Department of Nephrology, Osaka University Graduate School of Medicine, 2-2

Yamada-oka, Suita, Osaka, 565-0871, Japan

#These authors contributed equally to this work.

Correspondence and requests for materials should be addressed to

Yoichi Takami, M.D., Ph.D.*

Department of Geriatric and General Medicine, Osaka University Graduate School of

Medicine, #B6, 2-2 Yamadaoka, Suita, Osaka 565-0871, Japan

Tel.: +81-6-6879-3852, Fax: +81-6-6879-3859, E-mail: takami@geriat.med.osaka-

u.ac.jp

and

Yoichi Nozato, M.D., Ph.D.*

Department of Geriatric and General Medicine, Osaka University Graduate School of

Medicine, #B6, 2-2 Yamadaoka, Suita, Osaka 565-0871, Japan

Tel.: +81-6-6879-3852, Fax: +81-6-6879-3859, E-mail:

yoichi.nozato@geriat.med.osaka-u.ac.jp

and

Toshimasa Takahashi, M.D., Ph.D.*

Department of Geriatric and General Medicine, Osaka University Graduate School of

Medicine, #B6, 2-2 Yamadaoka, Suita, Osaka 565-0871, Japan

1    Tel.: +81-6-6879-3852, Fax: +81-6-6879-3859, E-mail:

2    toshimasa.takahashi@geriat.med.osaka-u.ac.jp

4    Keywords: AT1, LOX-1, oxidized LDL, Angiotensin II, Gq, CKD

**Abstract**

Chronic kidney disease (CKD) and atherosclerotic heart disease, frequently associated

with dyslipidemia and hypertension, represent significant health concerns. We

investigated the interplay among these conditions, focusing on the role of oxidized low-

density lipoprotein (oxLDL) and angiotensin II (Ang II) in renal injury via G protein $\alpha q$

subunit (Gq) signaling. We hypothesized that oxLDL enhances Ang II-induced Gq

signaling via the AT1 (Ang II type 1 receptor)-LOX1 (lectin-like oxLDL receptor)

complex. Based on CHO and renal cell model experiments, oxLDL alone did not activate

Gq signaling. However, when combined with Ang II, it significantly potentiated Gq-

mediated inositol phosphate 1 production and calcium influx in cells expressing both

LOX-1 and AT1 but not in AT1-expressing cells. This suggests a critical synergistic

interaction between oxLDL and Ang II in the AT1-LOX1 complex. Conformational

studies using AT1 biosensors have indicated a unique receptor conformational change due

to the oxLDL-Ang II combination. In vivo, wild-type mice fed a high-fat diet with Ang

II infusion presented exacerbated renal dysfunction, whereas LOX-1 knockout mice did

not, underscoring the pathophysiological relevance of the AT1-LOX1 interaction in renal

damage. These findings highlight a novel mechanism of renal dysfunction in CKD driven

by dyslipidemia and hypertension and suggest the therapeutic potential of AT1-LOX1

1    receptor complex in patients with these comorbidities.

**Introduction**

Dyslipidemia is a major risk factor of atherosclerotic heart disease in patients with chronic

kidney disease (CKD). It has been postulated that the association between dyslipidemia

and CKD is not solely a result of epidemiological comorbidities but rather a complex

interplay of causality, where CKD exacerbates dyslipidemia, while dyslipidemia, in turn,

contributes to the onset and progression of CKD[1,2]. Since Moorhead et al. proposed the

lipid nephrotoxicity hypothesis in 1982[3], accumulating evidence has suggested that

increased plasma lipid levels contribute to the development of renal glomerular and

tubular damage, primarily in animal models of dyslipidemia[4]. The etiology of

dyslipidemia-induced nephrotoxicity is complex and multifaceted, potentially involving

the activation of certain cellular signaling pathways that culminate in renal injury through

elevated levels of oxidized LDL (oxLDL)[5-8]. The lectin-like oxLDL receptor, LOX-1, is

implicated in organ damage caused by dyslipidemia, and its expression is increased in

hypertensive glomerulosclerosis[9]. In murine models, a deficiency of LOX-1 leads to a

reduction in renal dysfunction, which was precipitated by a systemic inflammatory state

following significant myocardial ischemia and injury[10]. This supports the hypothesis that

LOX-1 plays a role in the development of inflammation-induced renal injury. However,

to date, no studies have investigated the role of LOX-1 in nephrotoxicity caused by

dyslipidemia.

Hypertension is an established risk factor for CKD, and it has been suggested that

hypertension and dyslipidemia act synergistically to induce renal dysfunction[11]. The

development of renal damage due to hypertension involves direct renal injury by

vasoactive hormones, such as angiotensin II (Ang II), in addition to renal hemodynamic

abnormalities associated with elevated body pressure[12]. We have shown that LOX-1 and

the Ang II type 1 receptor (AT1) of the G protein-coupled receptor (GPCR) are coupled

on the plasma membrane, and that G protein-dependent and β-arrestin-dependent AT1

activation mechanisms are involved in the signaling mechanism by oxLDL and the

intracellular uptake of oxLDL, respectively[13,14].

Interestingly, it was recently demonstrated that AT1 exhibits different modes and degrees

of G protein activation depending on the conformational changes that occur during

activation by various ligands[15]. Specifically, G protein αq subunit (Gq)-biased agonists

induce a more open conformation of AT1 than Ang II, resulting in a more potent activation

of Gq, which is the primary mediator of Ang II-induced hypertension. In contrast, β-

arrestin-biased agonists induce a closer conformational change, leading to the activation

of β-arrestin without the activation of Gq[15]. Indeed, we found that oxLDL selectively

activates G protein αi subunit (Gi) of AT1, without activating Gq[14], similar to that induced

by β-arrestin-biased agonists[16]. However, given that Ang II and oxLDL coexist in physiological environments, it is plausible to hypothesize that the effect of these ligands on AT1 in living organisms may differ from their effects when administered individually. Indeed, we found that the combination treatment of oxLDL and Ang II enhanced pro-inflammatory NFκB activity compared to each treatment alone in cells overexpressing both AT1 and LOX-1[14]. Based on our findings and the aforementioned structure-activity relationship of AT1, we hypothesized that the binding of both oxLDL and Ang II to LOX-1 and AT1, respectively, may result in a more open AT1 structure, leading to stronger downstream Gq signaling. Therefore, this study aimed to investigate and clarify this hypothesis, focusing specifically on renal component cells, and ultimately demonstrate the in vivo relevance of this phenomenon in the development of renal injury or renal dysfinction under conditions of Ang II and oxLDL overload.

**Results**

**Oxidized LDL potentiates Ang II-induced Gq signaling in a LOX-1-dependent manner**

First, we examined the additive effect of oxLDL on Ang II-stimulated AT-1-Gq signaling in CHO (Chinese hamster ovary) cells that did not endogenously express LOX-1 and AT-1 but were genetically engineered to express these receptors (CHO-LOX-1-AT1) [13].

Fig. 1a shows the dose-response curve of Ang II-induced IP1 production, which serves as a measure of Gq activity, in the presence of varying concentrations of oxLDL in CHO-LOX-1-AT1 cells. Consistent with previous findings, oxLDL alone did not stimulate IP1 production in CHO-LOX-1-AT1 cells [14]. However, when oxLDL was supplemented at concentrations of 10 and 20 μg/mL (but not at 5 μg/mL), it caused a similar leftward shift in the dose-response curve of Ang II, resulting in a more than 80% decrease in the Effective Concentration 50 (EC50) (EC50 values: 0 μg/mL, $9.40 \times 10^{-9}$ M; 5 μg/mL, $5.21 \times 10^{-9}$ M; 10 μg/mL, $1.68 \times 10^{-9}$ M; 20 μg/mL, $1.55 \times 10^{-9}$ M). The maximum IP1 production induced by Ang II in CHO-LOX-1-AT1 cells was unaffected by the addition of oxLDL.

Additionally, we observed that oxLDL administration decreased Ang II-induced IP1 production in CHO cells expressing AT1 alone (CHO-AT1), in contrast to CHO cells

expressing both AT1 and LOX-1 (Fig. 1b). This finding suggests that the potentiating

effect of oxLDL on Ang II-AT1-Gq activity is dependent on the presence of LOX-1.

However, the reason for the decrease in IP1 production caused by oxLDL

supplementation was not determined in this study. Native LDL, which does not bind to

LOX-1, did not alter Ang II-induced Gq activity in CHO-LOX-1-AT1 cells (Fig. 1c).

Furthermore, the presence of advanced glycation end products (AGEs) that bind to LOX-

1[17], did not enhance Ang II-induced IP1 production (Fig. 1d).

We found that enhanced IP1 production by co-treatment of oxLDL with Ang II was

similarly observed in CHO-cells expressing LOX-1 and mutated AT1 with impaired

ability to activate β arrestin[14] (Fig 1e). Moreover, the potentiating effect of oxLDL on IP1

production was unaffected by Pertussis Toxin (PTX), a Gi inhibitor, or RKI-1448, a

downstream Rho kinase inhibitor targeting G12/13 signaling in CHO-LOX-1-AT1 cells.

However, this phenomenon was completely inhibited by YM-254890, the Gq inhibitor

(Fig. 1f). These results suggest that the potentiating effect of oxLDL on Ang II-induced

IP1 production is not influenced by β-arrestin, Gi, or G12/13, which are the main effectors

of AT1 signaling aside from Gq[18].

**oxLDL potentiates Ang II-induced calcium influx in a LOX-1-dependent manner**

Calcium influx is a representative cellular phenomenon that occurs in response to Ang II-AT1-Gq activation. We found that oxLDL and low concentrations of Ang II ($10^{-12}$ M) did not induce calcium influx in CHO-LOX-1-AT1 cells when treated alone, but apparently induced calcium influx when treated together (Fig. 1g, h). Ang II alone at higher concentrations ($10^{-11}$ M) induced calcium influx, and no further enhancement was observed with oxLDL supplementation (Fig. 1h). The combined effect of oxLDL and Ang II on calcium influx was completely blocked by YM254890, suggesting that this phenomenon was Gq-dependent (Fig. 1i). Importantly, oxLDL supplementation with Ang II did not affect the calcium influx in CHO cells expressing AT1 alone (Fig. 1j).

**Co-treatment of oxLDL with Ang II induces conformational change of AT1 different from each treatment alone**

To gain mechanistic insight into this phenomenon, we initially conducted a live-imaging analysis of membrane LOX-1 and AT1 to determine whether the combined treatment of oxLDL and Ang II leads to increased internalization of AT1 upon activation compared to individual treatments, as described in a previous protocol paper[19] (Fig. S1, Supplemental Video, and Fig. 2a).

Our findings revealed a decrease in green puncta, which represent AT1-eGFP, upon

treatment with Ang II, oxLDL alone, and their co-treatment compared with the control

group (Fig. 2a). However, the extent of membrane AT1 reduction was similar across all

the treatment groups (Fig. 2a). Consistent with our previous report[14], oxLDL treatment

resulted in a reduction in the red puncta representing LOX-1-mScarlet (Fig. 2a).

Importantly, co-treatment with oxLDL and Ang II did not further enhance the reduction

of red puncta compared to oxLDL treatment alone (Fig. 2a). Based on these findings, it

is conceivable that co-treatment with oxLDL and Ang II does not increase the content of

activated AT1 compared to individual treatments alone.

AT1 intramolecular FlAsH-BRET biosensors were used to detect AT1 conformational

changes in CHO cells expressing LOX-1[20]. Among several biosensors previously tested

[20], we used two sensors with FlAsH insertion at the third intracellular loop (ICL3P3) and

cytoplasmic-terminal tail (C-tailP1) of AT1 that interacts with Renilla luciferase (RlucII)

at the end of the cytoplasmic tail (AT1-ICL3P3 and AT1-C-tailP1, respectively), as these

sensors were shown to enhance BRET signaling induced by Ang II compared to biased

agonists, including SI, SII, DVG, and SBpA[20]. In CHO cells expressing LOX-1 alone

(CHO-LOX-1) transduced with lentivirus encoding AT1-C-tailP1, $10^{-5}$ M Ang II and the

combination of Ang II and 10 μg/mL oxLDL induced BRET similarly, whereas oxLDL

alone did not alter BRET (Fig. 2b). In contrast, in CHO-LOX-1 cells transduced with

AT1-ICL3P3-encoded lentivirus, the combination of Ang II and oxLDL induced BRET

more prominently than Ang II alone, whereas oxLDL alone did not alter BRET (Fig. 2c).

The difference in sensitivity between the CHO-LOX-1-AT1-3p3 and CHO-LOX-1-AT1-

C-tail P1 sensors likely explains why only the former showed a significant response to

the combination of Ang II and oxLDL, underscoring the importance of FlAsH insertion

site selection in these assays. Furthermore, the difference between oxLDL and the

combination treatment was abolished by a neutralizing antibody against LOX-1 (Fig. 2d).

These findings suggested that the concomitant binding of oxLDL to LOX-1 and Ang II to

AT1 induced a conformational change in AT1 that was distinct from that induced by Ang

II or oxLDL alone.

**Oxidized LDL potentiates Ang II-induced Gq-calcium signaling in renal cells**

To confirm the pathophysiological significance of this phenomenon observed in the

overexpressing cells, we validated it in cells endogenously expressing LOX-1 and AT1.

We found that oxLDL in combination with Ang II did not increase the cellular IP1 content

in human umbilical vein endothelial cells (HUVECs), bovine vascular endothelial cells

(BACEs), human aortic vascular smooth muscle cells (HAVSMCs), or rat macrophages

(A10) (Fig. S2). In contrast, in normal rat kidney epithelial cells (NRK52E) and

fibroblasts (NRK49F), the combination of oxLDL and Ang II, but not Ang II alone,

increased IP1 accumulation, which was suppressed in the presence of YM-25480, the Gq

inhibitor (Fig. 3a, b). IP1 accumulation induced by co-treatment with Ang II and oxLDL

was abolished by the siRNA-mediated knockdown of either AT1 or LOX-1 (Fig. 3 c, d;

the knockdown efficacy is shown in Fig. S3). Regarding calcium influx, intracellular

calcium concentrations were not increased by the treatment of either $10^{-7}$ M Ang II or low

concentration of oxLDL (2 μg/ml) alone (Fig. 3e, left and middle, Fig 3f). In contrast, the

combination of Ang II ($10^{-7}$ M) and oxLDL (2 μg/ml) increased intracellular $Ca^{2+}$

concentration (Fig. 3e, right, Fig. 3f). The combined effect on calcium influx was

attenuated by the siRNA knockdown of either AT1 or LOX-1 (Fig. 3g). The Gq inhibitor

and angiotensin receptor blocker (ARB), Irbesartan, also inhibited this phenomenon (Fig.

3h). Calcium influx was not induced by either combination therapy or monotherapy in

NRK52E cells (Fig. S4).

**Co-treatment of oxLDL and Ang II enhanced cellular response upon Gq activation**

**in renal cells**

In NRK49F cells, co-treatment of oxLDL and Ang II, compared to vehicle, increased

mRNA levels of NADPH components, *p67phox* and *p91phox*, fibrosis markers,

*fibronectin*, *collagen-1*, *collagen-4*, and *TGFβ*, and inflammatory cytokines, *TNFα*, *IL1β*,

*IL-6,* and *MCP-1* (Fig. 4a). Notably, oxLDL did not alter the expression of the genes of

interest, and Ang II increased the mRNA levels of only *fibronectin* and *MCP-1*, indicating

the apparent synergistic effect of co-treatment with Ang II and oxLDL on specific gene

expression. This amplification effect of co-treatment was also observed in limited genes

including *p67phox*, *TGFβ, IL-6,* and *MCP-1* in NRK52E cells (Fig. 4b). The combined

effects of oxLDL and Ang II on gene expression were completely abolished by the Gq

inhibitor (Fig. 4c, d) and ARB (Fig. 4e, f) in NRK49F and NRK 52E. We also verified

protein expression of αSMA as a molecular marker of epithelial-mesenchymal transition

(EMT) upon the indicated stimulation for 3 days in rat kidney cells. The combination of

Ang II ($10^{-7}$ M) and oxLDL (5 μg/mL) induced αSMA expression in NRK49F or NRK52E

to the same extent or less than TGFβ, a major inducer of EMT, respectively (Fig. 5a, b).

The results of the combination treatment were strikingly different from that of oxLDL or

Ang II treatment alone, which did not affect αSMA expression in both types of cells (Fig.

5c, d). The induction of αSMA by the combined treatment was suppressed in the presence

of a Gq inhibitor or ARB in both types of cells, suggesting the AT1-Gq-dependent

pathway in this phenomenon (Fig. 5e, f and Fig. 5g, h). As a final in vitro assay, the

proliferative activity of NRK49F cells after 24 h of treatment with Ang II and oxLDL,

either alone or in combination, was measured using the BrdU assay (Fig. 6a). When

administered alone, Ang II and oxLDL increased and reduced BrdU incorporation,

respectively. The reducing effect of oxLDL on proliferation was blocked by Irbesartan,

but not by a Gq inhibitor, consistent with our findings that oxLDL induces biased

activation of AT1, which favors Gi but not Gq signaling[14]. Oxidized LDL potentiated the

Ang II-induced increase in BrdU incorporation, which was blocked in the presence of a

Gq inhibitor or an ARB (Fig. 6a). The siRNA-mediated knockdown of AT1 completely

abolished the effects of Ang II and oxLDL, either alone or in combination. Knockdown

of LOX-1 did not affect the pro-proliferative effect of Ang II itself but blocked the

enhanced proliferation induced by co-treatment with oxLDL (Fig. 6b).

**Oxidized LDL-inducible diet exacerbates Ang II-induced renal dysfuntion in**

**wildtype mice, but not in LOX-1 knockout mice.**

To examine the relevance of this phenomenon in renal injury, we replicated the in vivo

conditions that occurred during simultaneous stimulation with oxLDL and Ang II in

wildtype (WT) and LOX-1 knockout (LOX-1 KO) mice. This was achieved by inducing

oxLDL via a high-fat diet (HFD) [21] and Ang II via a subcutaneously implanted osmotic

mini-pump (Fig. 7a). Two different doses of Ang II were used in the experiment: a pressor

dose of 0.7 γ, which was demonstrated to increase blood pressure (BP) and induce renal

dysfunction[22,23], and a subpressor dose of 0.1 γ, which does not affect BP. For the intended

analysis, the results are presented separately for mice exposed to subpressor and pressor

doses of Ang II, utilizing the same outcomes of control animals infused with saline for

comparison. Consistent with a previous report [24], the HFD used in the study prominently

increased the plasma LOX-1 ligand concentration, which was undetectable under a

normal diet (ND) after 6 weeks of feeding (Fig. S5). Notably, the HFD did not intensify

but rather attenuated the body weight increase during the experimental period compared

to the ND (Fig. S6a-c), likely due to reduced food intake in mice fed the HFD used in this

study (Fig. S6d, e). The pressor dose of Ang II similarly increased BP in HFD-fed and

ND-fed WT mice (Fig. 7b, Fig. S6f). However, notably, there was a modest trend of

attenuated BP elevation by 0.7 γ Ang II infusion in LOX-1 KO mice, in line with the

previous report showing reduced Ang II-induced BP elevation by LOX-1 deficiency in

mice (Fig. 7b, Fig. S6f) [25]. As expected, a subpressor dose of Ang II did not alter BP in

the corresponding mouse group (Fig. 7b, Fig. S6g). Regarding biofluid analysis, HFD

with saline infusion did not alter the urinary 8-OHdG concentration as a marker of

oxidative stress and urinary albumin excretion (UAE) compared to ND with saline

infusion in either WT or LOX-1 KO mice (Fig. 7c, d). The pressor dose of Ang II with

ND significantly increased the urinary 8-OHdG concentration and UAE in WT mice (Fig.

7c, d). Notably, HFD feeding in WT mice with a pressor dose of Ang II resulted in a

prominent increase in urinary 8-OHdG concentration and UAE compared to mice fed

with ND (Fig. 7c, d). In WT mice administered with a subpressor dose of Ang II, a

significant increase in UAE was observed when comparing the effects of HFD to ND (Fig.

7d). There were no significant differences in urinary 8-OHdG levels between the two

dietary conditions (Fig. 7c). Interestingly, when LOX-1 KO mice were fed either HFD or

ND and then administered the corresponding dose of Ang II, no differences were observed

in the measured parameters (Fig. 7c, d). These findings indicate that the combination of

HFD and Ang II administration appears to have a more pronounced effect on certain

biofluid markers of renal injury in WT mice than in LOX-1 KO mice. The presence or

absence of LOX-1 appears to influence the interaction between HFD and Ang II, affecting

these specific parameters in mice. We did not find a difference in plasma aldosterone

concentration between ND- and HFD-fed WT mice with a pressor dose of Ang II (Fig.

S7).

We then conducted quantitative real-time PCR analysis on genes within kidney sections,

encompassing NADPH components (*p40phox, p47phox, p67phox, and p91phox*),

inflammatory cytokines (*IL-6, TNFα, IL1β, MCP-1, and COX-2*), fibrosis markers (*TGFβ,*

*fibronectin, collagen-1a, collagen-4a, αSMA, and vimentin*), epithelial markers (*E-cadherin and cadherin-16*), and tubular marker (*NAGL*) (Fig. 8, Fig S8). In mice treated with a pressor dose of Ang II, 15 out of 18 genes examined showed enhanced alterations in gene expression, indicative of heightened NADPH oxidase components, inflammatory cytokines, fibrosis, decreased epithelial markers, and exacerbated tubular injury in HFD-fed WT mice compared to ND-fed mice (Fig. 8a, Fig. S8a). For many of these genes, a synergistic effect of the combination of a pressor dose of Ang II and HFD was not observed in LOX-1 KO mice (Fig. 8a, Fig. S8a). Seven genes displayed discernible differences between HFD- and ND-fed WT mice treated with a subpressor dose of Ang II (Fig. 8b Fig. S8b). Other than *fibronectin*, no differences were observed between ND- and HFD-fed LOX-1 mice exposed to a subpressor dose of Ang II (Fig. 8b Fig. S8b). In relation to AT1 and LOX-1 expression, no variations emerged between the ND and HFD groups when subjected to the corresponding dose of Ang II treatment in both WT and LOX-1 KO mice (Fig. S8a, b).

In the histological analysis of kidney samples, in contrast to the gene expression data, the administration of either a subpressor or pressor dose of Ang II, both in isolation and in combination with HFD for 4 weeks, did not reveal any notable increase in fibrosis, as evaluated by Masson-Trichrome staining (Fig. S9a). Similarly, there was no significant

change in the degree of mesangial expansion or glomerular area, as assessed by PAS staining. (Fig. S9b).

Finally, immunofluorescence staining of mouse kidney specimens was performed to detect the colocalization sites of LOX-1 and AT1 in the kidney. As shown in Fig. 9a and Fig. 9b, LOX-1 and AT1a were predominantly colocalized in the renal tubules, but not in the glomeruli. We also performed co-immunofluorescence staining with megalin, a well-established marker of proximal renal tubules, as shown in Fig. S10. Both AT1 and LOX-1 were observed to colocalize with megalin, especially at the brush borders, indicating their presence within the same renal compartments involved in AT1/LOX-1 signaling.

**Discussion**

We conducted a series of experiments to provide empirical support for our hypotheses.

We propose that the simultaneous binding of Ang II to AT1 and oxLDL to LOX-1 triggers distinct and more pronounced structural modifications in AT1 than the individual modifications induced by each ligand. This structural alteration in AT1 leads to the enhanced activation of Gq signaling.

To confirm AT1 and LOX-1 colocalization and interaction at the cell membrane, our previous study used in situ proximity ligation assays (PLA) and membrane protein immunoprecipitation assays in CHO cells expressing tagged AT1 and LOX-1, successfully demonstrating AT1/LOX-1 complex formation [13]. Additionally, we provided histological evidence of their co-localization with megalin in proximal renal tubules to support these findings, although the limitation remains regarding the lack of direct in vivo evidence for membrane co-localization of LOX-1 and AT1. While additional co-staining with other markers to identify specific cell types was not conducted, the prominent localization of AT1R with megalin in our study provides strong evidence of its expression in proximal renal tubules, consistent with established findings regarding AT1R presence in this nephron segment. Previous studies have documented AT1R expression in various renal cells, including mesangial, interstitial, and juxtaglomerular (JG) cells, as well as

proximal tubules. In our immunofluorescence analysis, however, we did not observe

significant AT1R expression in the glomerulus or mesangium. The pronounced

expression of AT1R in proximal tubules aligns with previous reports [26], though limitations

in immunofluorescence sensitivity do not exclude AT1R presence in other compartments.

Notably, our focus on proximal tubules enabled clear observation of AT1/LOX-1 co-

localization, especially under oxLDL and AngII stimulation. This interaction underscores

a potential focal point for AT1R/LOX-1 signaling in kidney disease pathogenesis within

the renal system.

At the cellular level, live imaging analysis (Fig. S1) displayed a limited number of

overlapping LOX-1 and AT1R puncta, which could be attributed to the dynamic and

transient nature of the LOX-1 and AT1 receptor interaction. As described in our previous

study [13], this interaction is sensitive to buffer conditions, with complex formation

occurring under non-reducing but not reducing conditions. This indicates that the LOX-1

and AT1 interaction is mediated by non-covalent rather than stable covalent bonds,

leading these receptors to form and dissociate rapidly. Consequently, only a small fraction

of LOX-1 and AT1 receptors may be co-localized at any given time point, explaining the

limited number of overlapping puncta observed in live imaging. Importantly, despite this

limited spatial overlap, our findings indicate that co-treatment with oxLDL and Ang II

significantly enhances Gq signaling. This highlights that the functional impact of the LOX-1/AT1 interaction, especially in response to specific stimuli like oxLDL and Ang II, is more crucial to downstream signaling outcomes than the extent of stable receptor co-localization.

The current experiments showed that oxLDL enhanced the effects of Ang II-induced production of IP1, a downstream signaling molecule associated with Gq activation and subsequent calcium influx, further supporting functional activation of the AT1/LOX-1 complex. However, this effect was observed only in the presence of both LOX-1 and AT1. The differences in activation levels observed between the IP1 assay and the calcium influx assay, both indicators of Gq activity, likely arise due to variations in assay sensitivity. While the absolute differences in the IP1 assay between treatment groups may appear modest, the critical comparison between Ang II alone and Ang II with oxLDL consistently demonstrated significant differences, in alignment with the calcium influx results. Notably, co-treatment with oxLDL reduced the EC50 for IP1 production by 80% compared to Ang II alone, underscoring a robust enhancement of Gq signaling, even though the IP1 assay differences were relatively small in absolute terms.

Considering that oxLDL selectively activates Gi but not Gq through the LOX-1-AT1 dependent pathway[14], it is evident that the observed phenomenon cannot be attributed

solely to the additive effect of oxLDL on AT1 activation. Rather, the simultaneous binding

of oxLDL and Ang II to their respective receptors, LOX-1 and AT1, which form a single

complex, underlies this phenomenon. Indeed, our findings from live imaging of the

membrane receptors revealed that the combination of Ang II and oxLDL did not induce

any additional influence on the internalization of both receptors upon AT1-β-arrestin

activation compared to each ligand alone, suggesting that the quantity of activated

receptors is not affected by the combination treatments. Considering the conformation-

activation relationship in AT1 activation[20], this supports the hypothesis that the

simultaneous binding of Ang II and oxLDL in a single AT1-LOX-1 complex induces a

greater conformational change, resulting in a more open conformation of AT1 compared

with each ligand alone. However, it is technically challenging to directly identify

structural modifications of the receptor complex using techniques such as crystal structure

analysis. Instead, we utilized AT1 conformational sensors capable of differentiating

between the conformational changes induced by Ang II and biased AT1[20]. Interestingly,

we observed that one of the two conformational sensors employed in this study detected

an augmented response in the presence of the combination treatment compared with Ang

II alone. Importantly, this enhanced response was significantly inhibited when an LOX-1

antibody was introduced, indicating the dependency of LOX-1 on this phenomenon.

These results indicate that the combined treatment with Ang II and oxLDL in the presence

of LOX-1 induces a unique conformational change in individual AT1 molecules, which

differs from the conformational changes induced by each single treatment alone.

It is important to note that the ability of LOX-1 ligands to enhance Ang II-AT1 signaling

is not commonly observed. This was corroborated by the observation that BSA-

conjugated AGE, a recognized ligand of LOX-1[17], failed to augment the production of

IP1 by Ang II compared to control BSA (Fig. 1d). While the exact reason for this

discrepancy is not yet understood, it is noteworthy that the predicted particle size of

oxLDL[27] is significantly larger at 250 Å compared to the maximum particle size of AGE-

BSA[28], which is 120 Å. This suggests a potentially greater impact of oxLDL on the

structural modifications within the AT1-LOX-1 complex. However, further structural

analysis is required to validate this hypothesis.

Notably, the Gq bias resulting from combination treatment varied across the mammalian

cells examined. Specifically, we observed a combinatorial effect exclusively in renal

epithelial and fibroblasts, whereas vascular endothelial and smooth muscle cells did not

display the same response. In these renal cells, we observed increased Gq signaling, along

with other cellular phenomena such as calcium influx, changes in gene expression, and

alterations in cellular characteristics, including myofibroblast activation and cell

proliferation. While our study demonstrates a significant upregulation of α-SMA, a well-established marker of myofibroblast, in renal epithelial and fibroblast cells exposed to combined Ang II and oxLDL treatment, we acknowledge the evolving understanding of EMT, particularly the role of partial epithelial-mesenchymal transition (pEMT) in CKD. Notably, pEMT has garnered attention under conditions of inflammation, oxidative stress, and elevated TGF-β, which are also relevant to our Ang II and HFD model. Unlike full EMT, where epithelial cells completely transition into mesenchymal cells, pEMT represents a state in which epithelial cells partially acquire mesenchymal characteristics, such as increased α-SMA expression and the secretion of pro-fibrotic cytokines, while remaining attached to the basement membrane without fully transitioning into fibroblasts [31]. Importantly, previous studies using the unilateral ureteral obstruction (UUO) model suggest that full EMT is unlikely to play a significant role in renal fibrosis and that most kidney fibroblasts are thought to originate from interstitial cells rather than via EMT [29]. The observed increase in α-SMA in our model may therefore indicate a pEMT-like state, indirectly contributing to fibrosis by promoting cytokine and growth factor release rather than directly driving fibroblast generation. This interpretation aligns with findings from other kidney fibrosis models, including the UUO model, which shares pathophysiological features such as inflammation and oxidative stress with our model. Given these

considerations, the increased α-SMA expression observed in our study may be indicative

of pEMT rather than definitive evidence of EMT directly contributing to fibroblast

differentiation. Additionally, extrapolating in vitro pEMT findings to in vivo models

presents inherent challenges, as detecting these subtle phenotypic changes remains

complex [31]. Further mechanistic investigations are required to clarify the contribution of

pEMT to renal fibrosis. Nevertheless, our findings support the possibility that pEMT may

contribute to fibrosis within the specific pathophysiological context of Ang II and HFD

co-administration.

Additionally, fibroblast proliferation, as assessed by the BrdU assay, was notably

enhanced by the combined treatment, as opposed to each treatment administered

separately. Interestingly, the use of 5 μg/mL oxLDL in our study showed a tendency to

decrease proliferation, which was counteracted by the administration of an ARB but not

a Gq inhibitor. These findings suggest that the presence of Ang II leads to a significant

transformation in the function of oxLDL, primarily due to its altered influence within the

AT1-LOX-1 complex. Taken together, these results suggest that the simultaneous binding

of oxLDL to LOX-1 and Ang II to AT1 results in a Gq-biased shift in AT1 activation,

leading to a cellular phenomenon that could potentially contribute to renal fibrosis.

In an animal study, we introduced a biological environment in which both circulating Ang

II and oxLDL (an LOX-1 ligand) were increased in mice. Previous studies using mice fed an HFD have consistently reported the onset of renal injury, as determined by various measurements[32-36]. In contrast, 6 weeks of HFD feeding without Ang II treatment did not alter renal function in our mice. This can be attributed to the lack of obesity induced by the HFD in this study. We used this diet based on a previous study that confirmed increased circulating LOX-1 ligand levels without body weight gain in mice [37]. In our experiments, this diet led to a decrease in body weight, likely due to reduced food intake in HFD-fed mice (Fig. S6d, e). Although modest, this weight reduction may influence renal function. Obesity is a well-established and clinically proven risk factor of renal dysfunction. The mechanisms underlying this association are complex and involve various factors other than lipid abnormalities, such as hemodynamic changes that affect kidney circulation and the impact of adipose tissue on the production of adipokines and other inflammatory mediators[38,39]. Consequently, we observed the influence of elevated lipid particle levels on renal function, independent of obesity. We found that the effect of an HFD became obvious with an increase in the Ang II load in WT mice. In particular, in WT mice treated with high-dose Ang II, which elevated systolic BP by approximately 30 mm Hg, an HFD induced notable increases in urinary reactive oxygen species and urinary albumin. In contrast, an HFD had no impact on Ang II-infused LOX-1 KO mice, as

evidenced by equivalent urinalysis measurements for renal injury between the diets.

Correspondingly, simultaneous administration of an HFD and Ang II resulted in a consistent alteration in the expression of genes related to renal injury, including fibrosis, inflammation, and oxidative stress, except for some genes in WT mice, but not in LOX-1 KO mice. This strongly suggests that the combined effect of Ang II and HFD on renal function is LOX-1 dependent. Nevertheless, it should be noted that the effect of high-dose Ang II infusion on BP tended to be less pronounced in LOX-1 KO mice compared to WT mice, although there were no differences in BP elevation between the diets in each group of mice. This reduction in BP may contribute to the decreased renal injury observed in LOX-1 KO mice, independent of the AT1/LOX-1 interaction. These findings align with those of previous studies indicating that LOX-1 knockout mice show resistance to Ang II-induced elevation of BP[25,40,41]. Specifically, when mice were infused with 2 γ Ang II (equivalent to 25 g mice) for a duration of 28 days, wildtype mice experienced a BP increase exceeding 180 mmHg, while LOX-1 knockout mice demonstrated a reduction of approximately 40 mmHg in this elevation[25]. Furthermore, the same research group reported that Ang II infusion led to less severe renal injury in LOX-1KO mice compared to WT mice[40]. Importantly, these findings were observed in mice fed with an ND, suggesting that the protective effect of LOX-1 loss-of-function against Ang II-induced

elevated BP occurs through the LOX-1-AT1 complex, independent of the presence of oxLDL. Additionally, under subpressor doses of Ang II, where no significant differences in BP were observed, HFD-fed WT mice still exhibited increased renal injury compared to ND-fed mice, an effect that was reduced in LOX-1 KO mice. These findings suggest that the protective effects of loss-of-function of LOX-1 are partly independent of BP changes, underscoring the role of the AT1/LOX-1 interaction in renal injury with Ang II and HFD co-treatment. We recognize that our single time-point analysis (1.5 months post-treatment) also may limit these observations, as the effects of AT1/LOX-1 interaction on renal injury could vary with treatment duration. Taken together, the effects of LOX-1 on AT1 signaling are complex, involving both ligand-dependent and -independent mechanisms, and further investigation is required for a comprehensive understanding of the LOX-1-AT1 interaction.

Regarding the effects of AT1 and LOX-1 interaction on the renin-angiotensin system (RAS) and blood pressure, oxLDL binding to LOX-1 enhances AT1 receptor-mediated Gq signaling, thereby promoting Ang II effects such as vasoconstriction, oxidative stress, and inflammation, all of which contribute to elevated blood pressure. However, in HFD-fed mice treated with a pressor dose of Ang II, plasma aldosterone levels showed an increasing tendency but remained statistically non-significant compared with ND-fed

mice (ND: 102.8 ± 11.6 pg/mL vs. HFD: 141.8 ± 15.0 pg/mL, P = 0.081), as shown in

Fig S7, indicating a limited response in aldosterone production under these conditions.

Additionally, BP did not significantly change (Fig S6f, g), potentially due to

heterogeneous cellular responses across cell types, as indicated by the lack of reaction in

vascular endothelial cells, vascular smooth muscle cells, and macrophages (Fig S2),

and/or possibly due to aldosterone saturation from the high Ang II dose.

Finally, the current findings unequivocally demonstrated the molecular interactions

between key molecules associated with dyslipidemia and hypertension in the kidneys.

Moreover, this interaction can be effectively inhibited by ARBs, suggesting an additive

effect in preventing the development of CKD, particularly in patients with hypertension

and dyslipidemia. Interestingly, Ang II-dependent hypertensive animal models, including

constriction of the renal artery and infusion of Ang II in rats and mice, have revealed a

progressive increase in intrarenal Ang II levels, surpassing what can be accounted for by

circulating Ang II levels alone[42]. This is due to Ang II-dependent renal activation of the

RAS, as indicated by increased urinary angiotensinogen (AGT) secretion[42,43]. Importantly,

the elevation of urinary AGT is also evident in patients with various pathologies,

including hypertension and CKD, implying that renal Ang II levels increase even in

individuals who do not exhibit elevated levels of circulating Ang II[21,43,44]. Taken together,

the current finding of the synergistic effect of Ang II and oxLDL on AT1 activation in

renal tissue is highly relevant for the development of kidney disease. RAS inhibitors,

ARB, and ACE inhibitors are prioritized therapies to prevent the development of CKD

with proteinuria in patients with hypertension[45]. In addition to the well-established

inhibitory effects of Ang II on the contraction of efferent arterioles[46], a novel renal

protective action of RAS inhibitors has been proposed. Particularly in hypertension

accompanied by dyslipidemia, RAS inhibitors may exhibit anti-inflammatory, antifibrotic,

and antioxidant effects in the kidneys by inhibiting the Gq signaling pathway through the

AT1-LOX-1 complex in renal tubular cells and fibroblasts. In terms of ARBs'

effectiveness in inhibiting AT1/LOX-1 receptor conformational changes, all ARBs

generally block the downstream signaling from AT1-LOX-1 interaction by preventing

Ang II binding to AT1. Our previous study also showed that ARBs, including olmesartan,

telmisartan, valsartan, and losartan, could inhibit AT1 activation by oxLDL in the absence

of Ang II [13]. However, certain ARBs—olmesartan, telmisartan, and valsartan—also act

as inverse agonists, reducing baseline AT1 activity by preventing conformational changes

even without Ang II [13]. This inverse agonist property may offer additional therapeutic

benefits by reducing receptor activation beyond what is achieved by simple antagonism

in pathological states where AT1 activation occurs independently of Ang II, such as in

oxLDL presence. Collectively, the current findings suggest that RAS inhibitors, some of which possess inverse agonist properties, can concomitantly mitigate the effect of increased renal Ang II and oxLDL levels on the development of CKD in patients with hypertension and dyslipidemia, although direct evidence in clinical studies to support this remains to be elucidated.

This study has several limitations as follows: 1) The kidney, a complex organ vital for maintaining homeostasis, comprises a myriad of distinct cell types working in concert to execute its multifaceted functions [47]. The experiments conducted in mice raised questions regarding the specific cell types implicated in the synergistic effect of Ang II and oxLDL within the LOX-1-AT1 complex in the kidney. Addressing this issue would ideally require single-cell analysis, which is a challenge for future research. 2) The study employed systemic LOX-1 knockout mice. For a more detailed analysis, a phenotypic investigation using renal tissue-specific LOX-1 and AT1 knockout mice is required. Of particular importance is the analysis using tubule-specific knockout mice, in which the localization of these components has been verified through immunohistochemical staining. 3) The administration of Ang II (both pressor and subpressor doses) and its combination with HFD did not result in any histological changes in terms of fibrosis and mesangial expansion, despite the observed alterations in the associated gene expression and

urinalysis for renal injury. Alterations in gene expression and urinalysis for renal injury

may be sensitive and relatively early phenomena, and this discrepancy could potentially

be attributed to the relatively short intervention duration of 4 weeks for combined HFD

and Ang II administration. Therefore, concurrent pathohistological alterations in the renal

tissue might become evident with a more extended intervention period. 4) This study was

limited by its inability to detect the amplification of Gq signaling in mouse renal tissue

due to the concurrent administration of Ang II and HFD. Overcoming this limitation is a

challenge for future studies.

In conclusion, the current findings suggest that the simultaneous binding of oxLDL and

Ang II to their respective receptors within the complex induces a distinct conformational

change compared with the effect of each ligand alone. This unique conformational change

results in the heightened activation of G protein signaling and subsequent unfavorable

cellular reactions in renal component cells. The relevance of this phenomenon was

confirmed in mouse models, in which renal dysfunction was prominently exacerbated

when there was a concomitant increase in Ang II and oxLDL levels. Notably, this effect

was abolished by the deletion of LOX-1, indicating LOX-1 dependency of this in vivo

phenomenon (Fig. 10). These findings indicate the clinical relevance of the direct

interaction between hypertension and dyslipidemia and further support the clinical

1    significance of RA inhibition in treating patients with CKD.

**Materials and methods**

**Cell culture and materials**

HUVECs and BAECs were cultured in EGM-2 (Lonza, Basel, Switherlad). Cells with

fewer than five passages were used in the experiments. Transgenic CHO cells were

maintained in an F-12 Nutrient Mixture with GlutamaxTM-I (Thermo Fisher Scientific,

MA, USA), 10% fetal bovine serum (FBS; Gibco, USA), and appropriate selection

reagents, as described below. CHO-K1 cells were maintained in F-12 Nutrient Mixture

with GlutamaxTM-I and 10% FBS. HAVSMCs were cultured in Dulbecco's modified

Eagle's medium/F12 (DMEM/F12) (Nacalai Tesque, Japan) supplemented with 1%

penicillin-streptomycin (Fujifilm, Japan) and 10% FBS. A10 cells were grown in DMEM

(Wako, Osaka, Japan) supplemented with 10% FBS and 1% penicillin-streptomycin.

NRK52E and NRK49F cells (ECACC, UK) were cultured in DMEM (Wako, Japan)

supplemented with 5% FBS and the appropriate selection reagents. Gene transcription in

CHO cells was induced by adding 100 ng/mL doxycycline (Merck KGaA, Darmstadt,

Germany). Cells were incubated at 37°C in 5% $CO_2$ and 95% air.

**Construction of plasmid vectors**

For stable transformants, pTRE2hyg vector (Clontech, USA) encoding mutated hAT1

with impaired ability to activate β-arrestin (pTRE2hyg-HA-FLAG-hAT1mg) were

created using site direct mutagenesis as previously described[14]. For real-time imaging,

LOX-1 tagged with V5-6×His at the C-terminus (V5-LOX-1) was subcloned into

pmScarlet_C1 (plasmid #85042; Addgene) (mScarlet-LOX-1). HA-FLAG-hAT1 was

subcloned into pcDNA3-EGFP (plasmid #85042; Addgene) (AT1-eGFP)[14].

**Stable transformants**

We constructed CHO-K1 cells expressing tetracycline-inducible human LOX-1 tagged

with V5-6×His at the C-terminus (CHO-LOX-1), human HA-FLAG-hAT1 (CHO-AT1),

or cells expressing both human LOX-1 and AT1 (CHO-LOX-1-AT1), as previously

described[13,14] To establish cells expressing both LOX-1 and mutated AT1 (pTRE2hyg-

HA-FLAG-hAT1mg), they were co-transfected with the pSV2bsr vector (Funakoshi,

Japan) into CHO-LOX-1 using the Lipofectamine 2000 transfection reagent (Thermo

Fisher Scientific, USA). The stable transformants were selected with 400 μg/mL of

hygromycin B (Wako, Osaka, Japan) and 10 μg/mL of blasticidin S (Funakoshi, Japan).

The resistant clone expressing LOX-1 and mutated AT1 in response to doxycycline

(Calbiochem, USA) was selected for use in the experiments (CHO-LOX-1-AT1mg) [14].

**Small interfering RNA**

NRK52 and NRK47 cells were plated at 50% confluence on the day of transfection.

Silencer Select small interfering RNA (siRNAs) for *LOX-1* and *AT1a* (Thermo Fisher

Scientific, MA, USA ) were transfected into cells in medium without serum or antibiotics

using Lipofectamine RNAiMAX (Thermo Fisher Scientific, MA, USA ), according to the

manufacturer's instructions.

**Preparation of oxLDL**

Human plasma LDL (1.019-1.063 g/mL), isolated by sequential ultracentrifugation, was

oxidized using 20 μM $CuSO_4$ in PBS at 37 °C for 24 h. Oxidation was terminated by

adding excess EDTA. LDL oxidation was analyzed by agarose gel electrophoresis for

migration versus LDL[13].

**Quantification of cellular IP1 accumulation**

Gq-dependent activation of phospholipase C was quantified by measuring IP1 using the

IP-One assay kit (Cisbio, France) as previously described[19]. Briefly, cells were seeded at

80,000 cells/well in 96 well transparent cell culture plates and incubated under serum-

free conditions for 24 h. Thereafter, cells were treated for 1 h with IP1 stimulation buffer,

including vehicle, native LDL, oxLDL, Ang II, AGE, PTX (Merck KGaA, Darmstadt, Germany), YM-254890 (Fujifilm Wako, Osaka, Japan), and RKI-1448 (Selleck, USA), as described in the text. Cell lysates with Triton X at a final concentration of 1% were transferred to a 384-well white plate, and IP1 levels were measured by incubating the cell lysates with FRET reagents (cryptate-labeled anti-IP1 antibody and d2-labeled IP1 analog).

**Quantitative real-time PCR**

RNA samples were purified using RNeasy Mini Kit (Qiagen, Germantown, MD, USA). One microgram of RNA was converted into cDNA using a ReverTra Ace qPCR RT kit (TOYOBO, Osaka, Japan) according to the manufacturer's instructions. All genes were evaluated using the ViiA7 Real-Time PCR System (Applied Biosystems, Thermo Fisher Scientific, Waltham, MA, USA). The data were analyzed using the $\Delta\Delta$Ct method with normalization against the GAPDH RNA expression in each sample. The primer sequences are listed in the Supplemental Table.

**Western blotting**

Proteins were separated using sodium dodecyl sulfate-polyacrylamide gel electrophoresis

and electrophoretically transferred onto polyvinylidene fluoride membranes. The

membranes were blocked with 5% nonfat dried milk and incubated with primary

antibodies overnight at 4 ºC. The primary antibodies used in this study were as follows:

anti-SMA antibody (1:1000), anti-$\alpha$-Tubulin antibody (1:1000) (Cell Signaling

Technology, Inc, Danvers, MA, USA). The bands were visualized with a

chemiluminescence detection system (LAS-4000 mini; GE Healthcare Life Sciences,

Buckinghamshire, UK) using Chemi-Lumi One Super (Nacalai Tesque, Kyoto, Japan).

**Calcium influx assay**

Calcium influx was measured using Fura 2-AM (Dojindo, Kumamoto, Japan) with slight

modifications to the manufacturer's protocol. In brief, cells plated in 96 wells were

incubated with 5 $\mu$M Fura 2-AM in HEPES buffer saline (20 mM HEPES, 115 mM NaCl,

5.4 mM KCl, 0.8 mM MgCl2, 1.8 mM $CaCl_2$, 13.8 mM glucose, pH 7.4) for 1 h at 37°C,

followed by replacement with recording medium without Fura 2-AM. Cells were treated

with oxLDL, Ang II, or a combination of both at the indicated concentrations. Changes

in F340/F380, an index of intracellular calcium concentration, were measured by dual-

excitation microfluorometry using a digital image analyzer (Aquacosmos; Hamamatsu

Photonics, Hamamatsu, Japan).

**BrdU assay**

Proliferative activity was assessed using a BrdU Cell Proliferation ELISA Kit (Funakoshi,

Japan). NRK49F cells were seeded in 96-well tissue culture plates and incubated with the

test reagents for 24 h. After incubating the cells with BrdU, we fixed the cells and

denatured their DNA using a Fixing Solution. The plate was washed thrice with Wash

Buffer before adding the Detector Antibody. Next, 100 μL/well of anti-BrdU monoclonal

Detector Antibody was added, and the plate was incubated for 1 h at room temperature.

Subsequently, 100 μL/well of Goat Anti-Mouse IgG Conjugate was pipetted and

incubated for 30 minutes at room temperature. After five washes, the reaction was stopped

by adding Stop Solution to each well. The color of the positive wells changed from blue

to bright yellow. Finally, the plate was read at a wavelength of 450/550 nm using a

spectrophotometric microtiter plate reader.

**Creation of lentivirus encoding AT1 conformational sensors**

For lentivirus encoding AT1 conformational sensors, rat AT1 and RlucII were subcloned

into the pLVSIN-CMV Neo vector (Takara Bio, Japan). Next, the FlAsH binding

sequence (CCPGCC) was inserted between residues K135 and S136 in the third

intracellular loop (AT1-ICL3P3), as well as between residues K333 and M334 in the

cytoplasmic-terminal tail (AT1-Ctail), utilizing the KOD-Plus Mutagenesis Kit (Toyobo,

Japan), at the same site as previously documented[14].

**Intramolecular FlAsH Bioluminescence resonance energy transfer assay to detect**

**AT1 conformational change**

CHO-LOX-1 cells were initially plated onto a white clear-bottom 96-well culture plate at

a density of $1 \times 10^5$ cells/well. The following day, the cells were transduced with lenti-

virus encoding AT1-Ctail or AT1-ICL3P3 [20] in 10% FBS. After 24 h of transduction, the

cultures were transferred to serum-free conditions and incubated for an additional 24 h.

Add 1.5 µM FlAsH-EDT$_2$ labeling reagent of TC-FlAsH™ II In-Cell Tetracysteine Tag

Detection Kit (Thermo Fisher Scientific, MA, USA), wash twice with 250 µM BAL buffer,

and assays were promptly conducted on a Spark® microplate reader (TECAN,

Switzerland). The BRET ratio (emission mVenus/emission Rluc) was calculated as

follows: Following a 3-minute baseline reading (with the final baseline reading presented

at 0), cells were exposed to vehicle, oxLDL alone, AII alone, a combination of AII and

oxLDL, or a combination of AngII, oxLDL, and LOX-1 antibodies. The BRET ratios

were calculated every 16 seconds for a total of 320 seconds and the relative change in

intramolecular BRET ratio was calculated by subtracting the average BRET ratio

measured for cells stimulated with vehicle at each time point.

**Analysis of LOX-1 and AT1 dynamics by real-time imaging**

Live imaging was performed using previously reported methods[19]. Briefly, twenty-four

hours prior to imaging experiments, CHO-K1 cells were transfected with LOX-1-

mScarlet and AT1-eGFP by electroporation. Subsequently, the cells were seeded in a 35-

mm glass base dish (Iwaki, Japan) that had been pre-coated with a 1000X diluted solution

of 10 mg/mL poly-L-lysine (ScienCell, USA) one hour before seeding. The growth

medium was substituted with imaging buffer (pH 7.4), which consisted of 125 mM NaCl,

5 mM KCl, 1.2 mM $MgCl_2$, 1.3 mM $CaCl_2$, 25 mM HEPES, and 3 mM D-glucose, with

the pH adjusted to 7.4 using NaOH. Dynamic images of the cells were acquired at 25 °C

using a SpinSR10 inverted spinning disk-type confocal super-resolution microscope

(Olympus, Japan). The microscope was equipped with a 100x NA1.49 objective lens

(UAPON100XOTIRF, Olympus, Japan) and an ORCA-Flash 4.0 V2 scientific CMOS

camera (Hamamatsu Photonics KK, Japan) at 5-second intervals. The imaging

experiment was conducted using CellSens Dimension 1.11 software, employing a 3D

deconvolution algorithm (Olympus, Japan), and the number of puncta was determined

using ImageJ1.53K[19].

**Animals and diets**

Male WT mice (C57BL/6J) and LOX-1 KO mice with a C57BL/6 background were used

in this study. LOX-1 KO mice were generated as described previously[48]. Mice were

housed in a temperature-controlled (20-22°C) room on a 12 hours light/dark cycle and

fed an ND (MF; Oriental Yeast, Osaka, Japan) or an HFD (High Fat Diet without DL-α-

tocopherol, CLEA Japan Inc, Tokyo, Japan), which reported to increase plasma LOX-1

ligand in ApoE KO mice[21]. All study protocols were approved by the Animal Care and

Use Committee of Osaka University and were conducted according to the guidelines of

the NIH for the Care and Use of Laboratory Animals.

**Blood pressure measurement in mice**

The blood pressure of the mice was measured using the tail- cuff method with BP-98A

(Softron, Japan). The measurements were performed after restraining the mice. The blood

pressure was calculated as the average of 6 readings for each animal at each time point.

**Urine tests in mice**

Urine tests in mice included the measurement of urine 8-OHdG, creatinine, and albumin

concentrations. The DNA Damage (8-OHdG) ELISA Kit (StressMarq Bioscience,

Canada), Creatinine Kit L type Wako (Fujifilm, Japan), and Mouse Albumin ELISA Kit

(Bethyl Laboratories, Inc., TX, USA) were utilized for these measurements, following

their respective instructions.

**Plasma LOX-1 ligand concentration**

Measurement of LOX-1 ligands containing apoB (LAB) in mouse plasma was performed

using a modified protocol based on a previously reported method [37]. Briefly, recombinant

human LOX-1 (0.25 μg/well) was immobilized on 384 well plate (greiner, Frickenhausen,

Germany) by incubating overnight at 4°C in 50 μl of PBS. After three washes with PBS,

80 μl of 20% (v/v) ImmunoBlock (KAC, Kyoto, Japan) was added, and the plates were

incubated for 2 h at 25°C. After three washes with PBS, the plates were incubated for 2 h

at 25°C with 40 μl of standard oxidized LDL or samples. Samples were prepared by 4-

fold dilution of plasma with HEPES-EDTA buffer (10 mM HEPES, 150 mM NaCl, 2 mM

EDTA, pH 7.4), and standards were prepared by dilution of oxidized LDL with HEPES-

EDTA buffer. Following three washes with PBS, the plates were incubated for 1 h at 25°C

with chicken monoclonal anti-apoB antibody (HUC20, 0.5 μg/mL) in HEPES-EDTA

containing 1% (w/v) BSA. After three washes with PBS, the plates were incubated for 1

h at 25°C with peroxidase-conjugated donkey anti-chicken IgY (Merck, NJ, USA) diluted

5000 times with HEPES-EDTA containing 1% (w/v) BSA. After five washes with PBS,

the substrate solution containing 3,3',5,5'-tetramethylbenzidine (TMB solution, Bio-Rad

Laboratories, CA, USA) was added to the plates and incubated them for 30 min at room

temperature. The reaction was terminated with 2 M sulfuric acid. Peroxidase activity was

determined by measuring absorbance at 450 nm using a SpectraMax 340PC384

Microplate Reader (Molecular Devices, CA, USA).

**Tissue preperation**

Kidneys were perfused with cold PBS before removal. Kidney samples were rapidly

excised. A quarter of samples were stored at 4°C in RNAlater (Thermo Fisher Scientific,

MA, USA) for RNA extraction. The remaining quarters were fixed in 4%

paraformaldehyde overnight at 4°C for histological evaluation. The remaining half was

snap-frozen in liquid nitrogen and stored at -80°C for further analysis.

**Periodic acid-Schiff and Masson-Trichrome staining**

The degree of glomerular mesangial expansion and glomerular area (representing the

structural integrity of the glomeruli) were assessed in a blinded manner using periodic

acid-Schiff (PAS) staining. Collagen accumulation was determined by Masson-Trichrome

(MTC) staining. For MTC staining, the area displaying fibrosis was quantitatively

evaluated in a blinded manner by measuring the blue staining in six strongly magnified

fields of view using the ImageJ software, and the average was calculated after

determining the ratio of the total area.

**Fluorescent immunostaining**

For fluorescent immunostaining of LOX-1, AT-1, and meglin, the mice were perfused

with cold saline before tissue removal. After 3 days of zinc fixation, the tissue was

replaced with 70% ethanol. The 3-μm-thick kidney tissue sections were

immunohistochemically stained with antibodies against ATGR (1:200, Cosmo Bio,

Japan), OLR-1 (1:200, TS58 from the laboratory of T.S., Shinshu University

School of Medicine, Nagano, Japan), and megalin (1:200, BiCell Scientific, MO,

USA). Following deparaffinization (using Lemosol and gradient ethanol) and rehydration,

the slices were subjected to antigen retrieval by autoclaving in citrate buffer (0.01 M; pH

6.0). Subsequently, the slices were washed thrice with PBS and blocked with 5% bovine

serum albumin for 30 min at room temperature. The slides were then incubated with

primary antibodies for 2 h at room temperature. Goat anti-Rabbit IgG (H+L) High Cross-

Adsorbed Secondary Antibody, Alexa Fluor Plus 488 and 594 (Thermo Fisher Scientific,

MA, USA) were used as secondary antibodies for ATGR and OLR-1, respectively. After

incubation with secondary antibodies for 1 h at room temperature, the slices were washed

with PBS. Finally, slides were sealed and photographed. Visual analyses were performed

using a BZ-800L microscope (Keyence, Japan).

**Statistical analyses**

All data are presented as the mean ± SEM. Differences between two treatments or among

multiple treatments were determined using the Student's t-test or one-way ANOVA

followed by Tukey's multiple comparison test.

**Data availability**

All data supporting the findings of this study are available within the paper and its

Supplementary Information. Source data are provided in this paper.

**Acknowledgements**

This work was partially supported by JSPS KAKENHI Grant Numbers 21K07389 (Y.T.),

22K08181 (Y.N.), 20H03576 (H.R.), and 18H02732 (K.Y.). We are grateful to Tomoko

Sato, Yoshinori Koishi, and Chika Takana for technical assistance. We would like to thank

Editage (www.editage.com) for the English language editing.

**Author contributions**

**Competing interests**

The authors declare no competing interests.

**References**

1     Harper, C. R. & Jacobson, T. A. Managing dyslipidemia in chronic kidney

disease. *J Am Coll Cardiol* **51**, 2375-2384 (2008).

https://doi.org/10.1016/j.jacc.2008.03.025

2     Cases, A. & Coll, E. Dyslipidemia and the progression of renal disease in

chronic renal failure patients. *Kidney Int Suppl*, S87-93 (2005).

https://doi.org/10.1111/j.1523-1755.2005.09916.x

3     Moorhead, J. F., Chan, M. K., El-Nahas, M. & Varghese, Z. Lipid nephrotoxicity

in chronic progressive glomerular and tubulo-interstitial disease. *Lancet* **2**, 1309-

1311 (1982). https://doi.org/10.1016/s0140-6736(82)91513-6

4     Ruan, X. Z., Varghese, Z. & Moorhead, J. F. An update on the lipid

nephrotoxicity hypothesis. *Nat Rev Nephrol* **5**, 713-721 (2009).

https://doi.org/10.1038/nrneph.2009.184

5     Bussolati, B. *et al.* Statins prevent oxidized LDL-induced injury of glomerular

podocytes by activating the phosphatidylinositol 3-kinase/AKT-signaling

pathway. *J Am Soc Nephrol* **16**, 1936-1947 (2005).

https://doi.org/10.1681/ASN.2004080629

6     Dai, Y. *et al.* High fat diet causes renal fibrosis in LDLr-null mice through

MAPK-NF-kappaB pathway mediated by Ox-LDL. *J Cardiovasc Pharmacol* **63**, 158-166 (2014). https://doi.org/10.1097/FJC.0000000000000035

Deng, S., Jin, T., Zhang, L., Bu, H. & Zhang, P. Mechanism of tacrolimus-induced chronic renal fibrosis following transplantation is regulated by ox-LDL and its receptor, LOX-1. *Mol Med Rep* **14**, 4124-4134 (2016). https://doi.org/10.3892/mmr.2016.5735

Gai, Z. *et al.* Lipid Accumulation and Chronic Kidney Disease. *Nutrients* **11** (2019). https://doi.org/10.3390/nu11040722

Nagase, M. *et al.* Expression of LOX-1, an oxidized low-density lipoprotein receptor, in experimental hypertensive glomerulosclerosis. *J Am Soc Nephrol* **11**, 1826-1836 (2000). https://doi.org/10.1681/ASN.V11101826

Lu, J. *et al.* Abrogation of lectin-like oxidized LDL receptor-1 attenuates acute myocardial ischemia-induced renal dysfunction by modulating systemic and local inflammation. *Kidney Int* **82**, 436-444 (2012). https://doi.org/10.1038/ki.2012.186

Manttari, M., Tiula, E., Alikoski, T. & Manninen, V. Effects of hypertension and dyslipidemia on the decline in renal function. *Hypertension* **26**, 670-675 (1995). https://doi.org/10.1161/01.hyp.26.4.670

Brewster, U. C. & Perazella, M. A. The renin-angiotensin-aldosterone system and the kidney: effects on kidney disease. *Am J Med* **116**, 263-272 (2004). https://doi.org/10.1016/j.amjmed.2003.09.034

Yamamoto, K. *et al.* Oxidized LDL (oxLDL) activates the angiotensin II type 1 receptor by binding to the lectin-like oxLDL receptor. *FASEB J* **29**, 3342-3356 (2015). https://doi.org/10.1096/fj.15-271627

Takahashi, T. *et al.* The endocytosis of oxidized LDL via the activation of the angiotensin II type 1 receptor. *iScience* **24**, 102076 (2021). https://doi.org/10.1016/j.isci.2021.102076

Wingler, L. M. *et al.* Angiotensin Analogs with Divergent Bias Stabilize Distinct Receptor Conformations. *Cell* **176**, 468-478 e411 (2019). https://doi.org/10.1016/j.cell.2018.12.005

Namkung, Y. *et al.* Functional selectivity profiling of the angiotensin II type 1 receptor using pathway-wide BRET signaling sensors. *Sci Signal* **11** (2018). https://doi.org/10.1126/scisignal.aat1631

Jono, T. *et al.* Lectin-like oxidized low density lipoprotein receptor-1 (LOX-1) serves as an endothelial receptor for advanced glycation end products (AGE). *FEBS Lett* **511**, 170-174 (2002). https://doi.org/10.1016/s0014-5793(01)03325-7

Lymperopoulos, A., Borges, J. I., Carbone, A. M., Cora, N. & Sizova, A.

Cardiovascular angiotensin II type 1 receptor biased signaling: Focus on non-

Gq-, non-betaarrestin-dependent signaling. *Pharmacol Res* **174**, 105943 (2021).

https://doi.org/10.1016/j.phrs.2021.105943

19    Huang, Y., Takahashi, T., Gaisano, H., Rakugi, H. & Yamamoto, K. A live-

imaging protocol for tracking receptor dynamics in single cells. *STAR Protoc* **3**,

101347 (2022). https://doi.org/10.1016/j.xpro.2022.101347

20    Devost, D. *et al.* Conformational Profiling of the AT1 Angiotensin II Receptor

Reflects Biased Agonism, G Protein Coupling, and Cellular Context. *J Biol*

*Chem* **292**, 5443-5456 (2017). https://doi.org/10.1074/jbc.M116.763854

21    Kobori, H. *et al.* Urinary angiotensinogen as a novel biomarker of the intrarenal

renin-angiotensin system status in hypertensive patients. *Hypertension* **53**, 344-

350 (2009). https://doi.org/10.1161/HYPERTENSIONAHA.108.123802

22    Jennings, B. L. *et al.* Cytochrome P450 1B1 contributes to renal dysfunction and

damage caused by angiotensin II in mice. *Hypertension* **59**, 348-354 (2012).

https://doi.org/10.1161/HYPERTENSIONAHA.111.183301

23    Wolak, T. *et al.* Osteopontin modulates angiotensin II-induced inflammation,

oxidative stress, and fibrosis of the kidney. *Kidney Int* **76**, 32-43 (2009).

https://doi.org/10.1038/ki.2009.90

24    Fujita, Y. *et al.* Oxidized LDL receptor LOX-1 binds to C-reactive protein and

mediates its vascular effects. *Clin Chem* **55**, 285-294 (2009).

https://doi.org/10.1373/clinchem.2008.119750

25    Hu, C. *et al.* Modulation of angiotensin II-mediated hypertension and cardiac

remodeling by lectin-like oxidized low-density lipoprotein receptor-1 deletion.

*Hypertension* **52**, 556-562 (2008).

https://doi.org/10.1161/HYPERTENSIONAHA.108.115287

26    Arthur, G., Osborn, J. L. & Yiannikouris, F. B. (Pro)renin receptor in the kidney:

function and significance. *Am J Physiol Regul Integr Comp Physiol* **320**, R377-

R383 (2021). https://doi.org/10.1152/ajpregu.00259.2020

27    Ohki, I. *et al.* Crystal structure of human lectin-like, oxidized low-density

lipoprotein receptor 1 ligand binding domain and its ligand recognition mode to

OxLDL. *Structure* **13**, 905-917 (2005). https://doi.org/10.1016/j.str.2005.03.016

28    Wright, A. K. & Thompson, M. R. Hydrodynamic structure of bovine serum

albumin determined by transient electric birefringence. *Biophys J* **15**, 137-141

(1975). https://doi.org/10.1016/s0006-3495(75)85797-3

29    Kriz, W., Kaissling, B. & Le Hir, M. Epithelial-mesenchymal transition (EMT)

in kidney fibrosis: fact or fantasy? *J Clin Invest* **121**, 468-474 (2011).

https://doi.org/10.1172/jci44595

30    Hadpech, S. & Thongboonkerd, V. Epithelial-mesenchymal plasticity in kidney

fibrosis. *Genesis* **62**, e23529 (2024). https://doi.org/10.1002/dvg.23529

31    Sheng, L. & Zhuang, S. New Insights Into the Role and Mechanism of Partial

Epithelial-Mesenchymal Transition in Kidney Fibrosis. *Front Physiol* **11**,

569322 (2020). https://doi.org/10.3389/fphys.2020.569322

32    Jiang, T. *et al.* Diet-induced obesity in C57BL/6J mice causes increased renal

lipid accumulation and glomerulosclerosis via a sterol regulatory element-

binding protein-1c-dependent pathway. *J Biol Chem* **280**, 32317-32325 (2005).

https://doi.org/10.1074/jbc.M500801200

33    Kume, S. *et al.* Role of altered renal lipid metabolism in the development of

renal injury induced by a high-fat diet. *J Am Soc Nephrol* **18**, 2715-2723 (2007).

https://doi.org/10.1681/ASN.2007010089

34    Yamamoto, T. *et al.* High-Fat Diet-Induced Lysosomal Dysfunction and

Impaired Autophagic Flux Contribute to Lipotoxicity in the Kidney. *J Am Soc*

*Nephrol* **28**, 1534-1551 (2017). https://doi.org/10.1681/ASN.2016070731

35    Sun, Y. *et al.* High-fat diet promotes renal injury by inducing oxidative stress

and mitochondrial dysfunction. *Cell Death Dis* **11**, 914 (2020).

https://doi.org/10.1038/s41419-020-03122-4

36    Yu, Y., Mo, H., Zhuo, H., Yu, C. & Liu, Y. High Fat Diet Induces Kidney Injury

via Stimulating Wnt/beta-Catenin Signaling. *Front Med (Lausanne)* **9**, 851618

(2022). https://doi.org/10.3389/fmed.2022.851618

37    Sato, Y. *et al.* Determination of LOX-1-ligand activity in mouse plasma with a

chicken monoclonal antibody for ApoB. *Atherosclerosis* **200**, 303-309 (2008).

https://doi.org/10.1016/j.atherosclerosis.2008.02.001

38    Garcia-Carro, C. *et al.* A Nephrologist Perspective on Obesity: From Kidney

Injury to Clinical Management. *Front Med (Lausanne)* **8**, 655871 (2021).

https://doi.org/10.3389/fmed.2021.655871

39    Tsuboi, N., Okabayashi, Y., Shimizu, A. & Yokoo, T. The Renal Pathology of

Obesity. *Kidney Int Rep* **2**, 251-260 (2017).

https://doi.org/10.1016/j.ekir.2017.01.007

40    Hu, C. *et al.* Deletion of LOX-1 attenuates renal injury following angiotensin II

infusion. *Kidney Int* **76**, 521-527 (2009). https://doi.org/10.1038/ki.2009.234

41    Li, X. *et al.* LOX-1 Deletion Attenuates Myocardial Fibrosis in the Aged Mice,

Particularly Those With Hypertension. *Front Cardiovasc Med* **8**, 736215 (2021).

https://doi.org/10.3389/fcvm.2021.736215

42   Navar, L. G., Kobori, H., Prieto, M. C. & Gonzalez-Villalobos, R. A.

Intratubular renin-angiotensin system in hypertension. *Hypertension* **57**, 355-362

(2011). https://doi.org/10.1161/HYPERTENSIONAHA.110.163519

43   Navar, L. G. Translational studies on augmentation of intratubular renin-

angiotensin system in hypertension. *Kidney Int Suppl (2011)* **3**, 321-325 (2013).

https://doi.org/10.1038/kisup.2013.67

44   Mills, K. T. *et al.* Increased urinary excretion of angiotensinogen is associated

with risk of chronic kidney disease. *Nephrol Dial Transplant* **27**, 3176-3181

(2012). https://doi.org/10.1093/ndt/gfs011

45   Kidney Disease: Improving Global Outcomes Blood Pressure Work, G. KDIGO

2021 Clinical Practice Guideline for the Management of Blood Pressure in

Chronic Kidney Disease. *Kidney Int* **99**, S1-S87 (2021).

https://doi.org/10.1016/j.kint.2020.11.003

46   Yang, R., Smolders, I. & Dupont, A. G. Blood pressure and renal hemodynamic

effects of angiotensin fragments. *Hypertens Res* **34**, 674-683 (2011).

https://doi.org/10.1038/hr.2011.24

47   Balzer, M. S., Rohacs, T. & Susztak, K. How Many Cell Types Are in the

1          Kidney and What Do They Do? *Annu Rev Physiol* **84**, 507-531 (2022).

2          https://doi.org/10.1146/annurev-physiol-052521-121841

3     48    Mehta, J. L. *et al.* Deletion of LOX-1 reduces atherogenesis in LDLR knockout

4          mice fed high cholesterol diet. *Circ Res* **100**, 1634-1642 (2007).

5          https://doi.org/10.1161/CIRCRESAHA.107.149724

**Figure Legends**

**Fig. 1| Oxidized LDL potentiates Ang II-induced Gq signaling and calcium influx in**

**a LOX-1-dependent manner**

**a** Dose-dependent response of IP1 concentration by the activation of Gq signaling in

response to oxLDL and Ang II in CHO-LOX-1-AT1 cells. Cells were treated with oxLDL

and Ang II at the concentrations described in the figure. (n=4 for each oxLDL

concentration). $^{*}P$=0.0004 for 0 μg/mL vs 20 μg/mL, $^{†}P$=0.0003 for 5 g/mL vs 20 μg/mL,

$^{‡}P$=0.0020 for 0 μg/mL vs 10 μg/mL, and $^{§}P$=0.0015 for 5 μg/mL vs 10 μg/mL at $10^{-9}$ M

Ang II; $^{\|}P$=0.0051 for 0 μg/mL vs 10 μg/mL, $^{¶}P$=0.0004 for 0 μg/mL vs 20 μg/mL at $10^{-}$

$^{8}$ M Ang II.

**b** IP1 concentration in response to vehicle, native LDL (nLDL 10 μg/mL), and oxLDL

(10 μg/mL) in the combination of Ang II ($10^{-8}$ M) in CHO-LOX-1-AT1 cells (n=5 for

each group)

**c** IP1 concentration in response to vehicle, oxLDL (10 μg/mL) in the combination of Ang

II ($10^{-8}$M) in CHO-LOX-1-AT1 and CHO-AT1 cells (n=5 for each group).

**d** IP1 concentration in response to vehicle, oxLDL (10 μg/mL), BSA (10 or 100 μg/mL),

BSA-conjugated AGE (10 or 100 μg/mL) in the combination of Ang ll ($10^{-8}$ M) in CHO-

LOX-1-AT1 cells.

**e** IP1 concentration in response to vehicle, oxLDL (10 μg/mL) in the combination of Ang

II ($10^{-8}$ M) in genetically engineered CHO cells with or without intact β-arrestin binding

domain (n=5 for each group). AT1mg indicates AT1 a mutant AT1 lacking a functional β-

arrestin binding domain but retaining G-protein-biased signaling capability.

**f** IP1 concentration in response to oxLDL (10 μg/mL) in the combination of Ang II ($10^{-8}$

M) and additional effect of PTX, a Gi inhibitor, YM-254890, a Gq inhibitor, and RKI-

1448, a downstream Rho kinase inhibitor targeting G12/13 signaling, in CHO-LOX-1-

AT1 cells (n=5 for each group).

**g** Intracellular calcium dynamics measured using Fura 2-AM by the ratio of the mission

signals at excitation wavelength 340 nm and 380 nm in response to oxLDL (5 μg/mL),

Ang II ($10^{-12}$ M), and their combination (for each agonist, 4-7 regions of interest were

selected). Addition of these agonists is marked with arrows on the timeline of the assay.

**h** Percentage changes from baseline in the ratio of emission signals (F340/F380)

measured by Fura 2-AM were quantified following treatment with oxLDL and Ang II at

specified concentrations in CHO-LOX-1-AT1 cells as detailed in the figure (n=4-8).

**i** Percentage change from baseline in the ratio of emission signals (F340/F380) measured

by Fura 2-AM after stimulation with oxLDL (5 μg/mL), Ang II ($10^{-12}$ M), and YM-

254890, a Gq inhibitor, in CHO-LOX-1-AT1 cells.

**j** Percentage changes from baseline in the ratio of emission signals (F340/F380) measured

by Fura 2-AM were quantified following treatment with oxLDL (5 μg/mL) and Ang II at

specified concentrations in CHO-AT1 cells, as detailed in the figure.

Data are represented as mean ± SEM. Differences were determined using one-way

ANOVA, followed by Tukey's multiple comparison test for (a-f) and (h-j).

**Fig. 2| Co-treatment of oxLDL with Ang II induces conformational change of AT1**

**different from each treatment alone**

**a** Change in green puncta (AT1-eGFP) and red puncta (LOX-1-mScarlet) by the treatment

with vehicle, oxLDL (10μg/mL), Ang II ($10^{-5}$M), and the combination of oxLDL and Ang

II in CHO cell overexpressing these fluorescent protein-conjugated receptors (n=10-11

for each group). The puncta were manually counted by a blinded observer and the number

of puncta at 0 and 3 min was determined (N0 and N3, respectively). The change in puncta

was calculated as (N0-N3)/N0.

**b-d** Changes in BRET signals were monitored in CHO-LOX-1 cells expressing the

following conformational biosensors: AT1-C-tailP1 (b) and AT1-ICL3P3 (c, d) bearing

FlAsH insertion at the cytoplasmic-terminal tail (C-tailP1) and the third intracellular loop

(ICL3P3) of AT1, respectively, that interact with Renilla luciferase at the end of the

cytoplasmic tail. Cells were subjected to treatments with vehicle, oxLDL (10 µg/mL),

Ang II ($10^{-5}$M), the combination of Ang II and oxLDL, and the combination

of AngII, oxLDL and LOX-1 antibody. The BRET ratios were calculated every 16

seconds for a total of 320 seconds and the relative change in intramolecular BRET ratio

(ΔBRET) was calculated by subtracting the average BRET ratio measured for cells

stimulated with vehicle at each time point. Lower panels indicate average ΔBRET of all

the time points during measurement.

Data are represented as mean ± SEM. Differences were determined by one-way ANOVA,

followed by Tukey's multiple comparison test (a-d).

**Fig. 3| Oxidized LDL potentiates Ang II-induced Gq-calcium signaling in renal cells**

**a, b** IP1 concentration in response to Ang II ($10^{-7}$ M), oxLDL (10 µg/mL), and the

combination of both with or without YM-254890, Gq inhibitor, in NRK52E (a) and

NRK49F cells (b) (n=5 for each group).

**c, d** IP1 concentration in response to Ang II ($10^{-7}$ M) and oxLDL (10 µg/mL) in the

combination of Ang II ($10^{-7}$ M) and the additional effect of siRNA-mediated knockdown

of *AT1a* or *LOX-1* in NRK52E (c) and NRK49F cells (d) (n=5 for each group).

**e** Intracellular calcium concentration in NRK49F cells using Fura 2-AM and dual-

excitation microfluorometry. Changes in the fluorescence intensity ratio (F340/F380)

served as an index of the calcium dynamics. Cells were exposed to Ang II ($10^{-7}$ M),

oxLDL (2 μg/mL), and a combination of both agents. Addition of these agonists is marked

with arrows on the timeline of the assay. Data acquisition and analysis were performed

using a digital image analyzer to monitor real-time calcium flux (n=4-9).

**f** Percentage changes from baseline in the ratio of emission signals (F340/F380) measured

by Fura 2-AM were quantified following treatment with Ang II ($10^{-7}$ M) and oxLDL at

the concentrations detailed in the figure in NRK49F cells (n=5 for each group).

**g** Impact of siRNA-mediated knockdown *AT1* or *LOX-1* on the percentage changes from

baseline in the ratio of emission signals (F340/F380) measured by Fura 2-AM were

quantified following treatment with Ang II ($10^{-7}$ M) and oxLDL (2 μg/mL) in NRK49F

cells (n=3-7).

**g** Impact of co-treatment of YM-254890, Gq inhibitor, or Irbesartan, an angiotensin

receptor blocker (ARB), on the the percentage changes from baseline in the ratio of

emission signals (F340/F380) measured by Fura 2-AM were quantified following

treatment with Ang II ($10^{-7}$ M) and oxLDL (2 μg/mL) in NRK49F cells (n=4 for each

group).

Data are represented as mean ± SEM. Differences were determined using one-way

ANOVA, followed by Tukey's multiple comparison test for (a-d) and (f-h).

**Fig. 4| Co-treatment of oxLDL and AII enhanced cellular response upon Gq**

**activation in renal cells**

**a,b** Quantitative real-time PCR analysis was performed to measure the gene expression

of NADPH oxidase subunits (*p67phox* and *p91phox*), fibrosis markers (*fibronectin*,

*collagen-1*, *collagen-4*, and *TGFβ*), and inflammatory cytokines (*TNFα*, *IL1β*, *IL-6*, and

*MCP-1*) in NRK49F cells (a) and NRK 52E cells (b). Gene expression levels were

normalized to those of GAPDH. Cells were stimulated by oxLDL (5 μg/mL), Ang II ($10^-$

$^7$ M) or their combination (n=4 for each group).

**c,** d Cells were pre-treated with vehicle or Gq inhibitor (YM-254890, Gqi), followed by

treatment with vehicle or the combination of oxLDL (5 μg/mL) and Ang II ($10^{-7}$ M) in

NRK49F cells (c) and NRK 52E cells (d) (n=4 for each group).

**e, f** Cells were pre-treated with vehicle or ARB (Irbesartan, Arb), followed by treatment

with vehicle or the combination of oxLDL (5 μg/mL) and Ang II ($10^{-7}$ M) in NRK49F

cells (e) and NRK 52E cells (f) (n=4 for each group).

Data are represented as mean ± SEM. Differences were determined by one-way ANOVA,

followed by Tukey's multiple comparison test (a-f).

**Fig. 5| Oxidized LDL enhanced Ang II-induced epithelial mesenchymal transition in**

**NRK52E and NRK49F cells**

**a, b** Left: Western blot analysis of α-smooth muscle actin (α-SMA), a marker of

epithelial–mesenchymal transition (EMT), in NRK49F (a) and NRK52E (b) cells. Cells

were stimulated with oxLDL (5 μg/mL), Ang II ($10^{-7}$ M), and TGF-β (10 ng/mL), with

TGF-β serving as a well-known EMT inducer. Right: Densitometric analysis of α-SMA

protein expression normalized to α-Tubulin (n=3 for each group).

**c, d** Left: Western blot analysis of α-SMA in NRK49F (c) or NRK52E (d) after stimulation

with oxLDL (5 μg/mL), Ang II ($10^{-7}$ M), and their combination. Right: Densitometric

analysis of α-SMA protein expression normalized to α-tubulin (n=3 for each group).

**e, f** Left: Western blot analysis of α-SMA in NRK49F(e) or NRK52E (f) after treatment

with a combination of oxLDL (5 μg/mL) and Ang II ($10^{-7}$ M). Prior to this treatment, the

cells were pre-treated with either a vehicle or a Gq inhibitor (YM-254890, Gqi). Right:

Densitometric analysis of α-SMA protein expression normalized to α-Tubulin (n=3 for

each group).

**g, h** Left: Western blot analysis of α-SMA in NRK49F (e) or NRK52E (f) after treatment

with a combination of oxLDL (5 μg/mL) and Ang II ($10^{-7}$ M). Prior to treatment, cells

were pre-treated with either vehicle or ARB (Irbesartan, Arb). Right: Densitometric

analysis of α-SMA protein expression normalized to α-tubulin (n=3 for each group).

Data are represented as mean ± SEM. Differences were determined by one-way ANOVA,

followed by Tukey's multiple comparison test (a-f).

**Fig. 6| Oxidized LDL enhanced Ang II-induced renal fibroblast proliferation via**

**AT1-Gq signaling and LOX-1-dependent manner**

**a** Proliferative activity assessed by BrdU incorporation into NRK49F cells. Cells were

pretreated with vehicle, YM-254890, or ARB, Irbesartan, followed by the treatment with

oxLDL (5 μg/mL), Ang II ($10^{-7}$ M), or their combination. (n=5 for each group).

**b** NRK49F cells were subjected to siRNA-mediated knockdown using specific siRNAs

for *AT1a* (siAT1) or *LOX-1* (siLOX-1). Following knockdown, cells were treated with

either vehicle, oxLDL (5 μg/mL), Ang II ($10^{-7}$ M), or their combination. Proliferative

activity was assessed by measuring the BrdU levels (n=5 for each group).

Data are represented as mean ± SEM. Differences were determined using one-way

ANOVA, followed by Tukey's multiple comparison test for (a) and (b).

**Fig. 7| Oxidized LDL-inducible diet exacerbates Ang II-induced renal dysfuntion in**

**wildtype mice, but not in LOX-1 knockout mice**

**a** A Schematic protocol for the animal experiments. Eight-week-old male WT mice and male LOX-1 KO mice were fed either an ND or an HFD for 6 weeks. After 10 weeks of age, the mice were treated over a 4-week period with infusions of either saline or Ang II. Ang II was administered at two dosage levels: a subpressor dose of 0.1 γ and a pressor dose of 0.7 γ, delivered via subcutaneously implanted osmotic pumps. At the end of the infusion period, urine was collected, the animals were sacrificed, and comprehensive tissue analysis was conducted to evaluate the renal effects of the treatments.

**b** Average systolic blood pressure (SBP) measured at half-week intervals in WT and LOX-1 KO mice during the 4-week infusion period.

**c, d** Urine 8-OHDG concentrations (mg/g creatinine [Cr]) (c) and urine albumin concentrations (mg/g creatinine [Cr]) (d) in WT and LOX-1 KO mice at the conclusion of the 4-week infusion period.

Data are represented as mean ± SEM. Differences were determined by one-way ANOVA, followed by Tukey's multiple comparison test (a-d).

**Fig. 8| A high-fat diet enhanced Ang II-induced renal injury-related gene expression in the kidney in a LOX-1-dependent manner.**

Quantitative real-time PCR analysis for gene expression of NADPH components

(*p67phox and p91phox*), inflammatory cytokines (*IL-6, TNFα, IL1β, and MCP-1*), and

fibrosis markers (*TGFβ, fibronectin, collagen-1a, and collagen-4a*) in the kidney

harvested from WT and LOX-1 KO mice.

The experimental procedures, including the dietary regimen and Ang II administration,

are detailed in Fig. 7a. Mice were administered either a pressor dose of 0.7 γ Ang II (a)

or a subpressor dose of 0.1 γ Ang II (b).

Data are represented as mean ± SEM. Differences were determined using one-way

ANOVA, followed by Tukey's multiple comparison test for (a) and (b).

**Fig. 9| LOX-1 and AT1a were predominantly co-localized in renal tubules**

**a, b** Representative images depicting staining for LOX-1 (a) and AT1 (b) in the renal

cortex tissues from wildtype mice (WT) and LOX-1 knockout mice (LOX-1 KO). Nuclei

are stained blue with DAPI. Green and red signals indicates AT1, while the red signal

indicates LOX-1. Overlay images demonstrate the merged visualization of AT1 or LOX-

1 with DAPI, highlighting the predominant colocalization of LOX-1 and AT1 in renal

tubules as opposed to the glomerulus.

**Fig. 10| Schematic overview of the AT1 and LOX-1 Interaction dynamics in renal**

**cells**

This schematic summary illustrates the predicted structure-activation relationship of the

AT1 receptor within the LOX-1-AT1 complex in renal component cells. This highlights

how the simultaneous binding of Ang II to AT1 and oxLDL to LOX-1 induces

conformational changes in AT1. These changes were more pronounced than those

triggered by the individual ligands. Such structural alterations have been proposed to

amplify Gq signaling pathway activation, subsequently leading to renal damage.

**Supplemental Figure Legends**

**Fig. S1| Live-imaging analysis of membrane LOX-1 and AT1 in response to the co-treatment of oxLDL with AngII.**

Real-time membrane imaging of CHO cells co-transfected with LOX-1-mScarlet and AT1-eGFP in response to oxLDL (10μg/ml) in the combination of Ang II ($10^{-7}$M) (Supplemental Video). A count of puncta was performed using separate images visualizing LOX-1-mScarlet (red puncta) and AT1-eGFP (green puncta) immediately before and 3 min after ligand application.

**Fig. S2| Oxidized LDL in combination with Ang II do not increase cellular IP1 content in human umbilical vein endothelial cells and bovine vascular endothelial cells, human aortic vascular smooth muscle cells, and rat macrophages**

IP1 concentration in response to oxLDL (10μg/ml) in the combination of Ang II ($10^{-7}$M) in HUVECs (human umbilical vein endothelial cell), BAECs (bovine aortic endothelial cell), HAVSMCs (human aortic vascular smooth muscle cell), and A10 cells (rat macrophages).

Data are represented as mean ± SEM. Differences were determined using one-way ANOVA, followed by Tukey's multiple comparison test (n=5 for each group).

**Fig. S3| Efficiency of siRNA-mediated knockdown for *AT1a* and *LOX-1* in NRK52E**

**and NRK49F cells**

NRK52E and NRK49F cells were transfected with siRNAs against scrambled siRNA,

*AT1a* or *LOX-1*. The efficiency of siRNA-mediated gene silencing was quantified by

assessing *AT1a* and *LOX-1* expression levels using quantitative real-time PCR.

Data are represented as mean ± SEM. Differences were determined using one-way

ANOVA, followed by Tukey's multiple comparison test.

**Fig. S4| Calcium influx was not induced by either the combination treatment of Ang**

**II or oxLDL or each treatment alone in NRK52E cells**

Percentage changes from baseline in the ratio of emission signals (F340/F380) measured

by Fura 2-AM were quantified following treatment with Ang II ($10^{-7}$M) and oxLDL at

the concentrations detailed in the figure for NRK49E cells (n=5-7 for each group).

Data are represented as mean ± SEM. Differences were determined using one-way

ANOVA, followed by Tukey's multiple comparison test.

**Fig. S5| High Fat Diet used in the study prominently increased plasma LOX-1 ligand**

**concentration**

Plasma LOX-1 ligand concentration of 14-week-old wildtype mice upon ND and HFD

for 6 weeks. From 10 weeks of age, these mice also received concurrent 4-week infusions

of Ang II at a pressor dose of 0.7 γ, administered through subcutaneously implanted

osmotic pumps.

Data are represented as mean ± SEM. Differences were determined using Student's t-test

(n=7 for each group).

**Fig. S6 | Impact of Diet and Ang II Infusion on Body Weight and Systolic Blood**

**Pressure in Mice**

**a** Final Body weight: This figure shows the body weights of WT and LOX-1 KO mice at

the end of the 4-week infusion period, highlighting the effects of diet and pharmacological

treatment.

**b, c** Serial Body Weight Changes: These graphs depict the progression of body weight

over time in WT and LOX-1 KO mice, illustrating the impact of the dietary regimen and

Ang II infusion on weight dynamics.

**d, e** Food Intake: Food intake Trajectory: Serial measurements of weekly food intake

(g/week per mouse) are presented for WT and LOX-1 KO mice. The figures show the

changes in food intake over the course of the study, corresponding to the administration

of a pressor dose of 0.7 γ (d) and a subpressor dose of 0.1 γ (e) of Ang II, respectively.

**f, g** Systolic Blood Pressure Trajectory: Serial measurements of systolic blood pressure

(SBP) obtained using the tail-cuff method are presented for WT and LOX-1 KO mice.

The figures show the changes in SBP over the course of the study, corresponding to the

administration of a pressor dose of 0.7 γ (d) and a subpressor dose of 0.1 γ (e) of Ang II,

respectively.

Beginning at 8 weeks of age, mice were fed either an ND or an HFD for 6 weeks. From

10 weeks of age, coinciding with the 2-week time point in the figure, the mice underwent

a 4-week period of infusion with either vehicle or Ang II. The infusion was delivered at

specific dosage levels through subcutaneously implanted osmotic pumps.

Data are represented as mean ± SEM. Differences were determined using one-way

ANOVA, followed by Tukey's multiple comparison test for (a).

**Fig. S7| No significant difference was found in plasma aldosterone concentration**

**between a normal diet and a high fat diet-fed wildtype mice with a pressor dose of**

**Ang II**

Plasma aldosterone concentration in 14-week-old wild-type mice fed an ND or an HFD

for 6 weeks. From 10 weeks of age, these mice also received concurrent 4-week infusions

of Ang II at a pressor dose of 0.7 γ, administered through subcutaneously implanted

osmotic pumps.

Data are represented as mean ± SEM. Differences were determined using Student's t-test

(n=7 for each group).

**Fig. S8| A high-fat diet enhanced Ang II-induced renal injury-related gene**

**expression in the kidney in a LOX-1-dependent manner.**

Quantitative real-time PCR analysis for gene expression of NADPH components

(*p40phox and p47phox*), inflammatory gene (*COX-2*), fibrosis markers (*αSMA and*

*vimentin*), epithelial markers (*E-cadherin and cadherin-16*), tubular marker (*NAGL*),

*AT1a*, *AT1b*, and *LOX-1* in the kidney harvested from WT and LOX-1 KO mice.

The experimental procedures, including the dietary regimen and Ang II administration,

are detailed in Fig. 7a. Mice were administered either a pressor dose of 0.7 γ Ang II (a)

or a subpressor dose of 0.1 γ Ang II (b).

Data are represented as mean ± SEM. Differences were determined using one-way

ANOVA, followed by Tukey's multiple comparison test for (a) and (b).

**Fig. S9| The treatment with Ang II, a High Fat Diet, or their combination for 4 weeks**

**did not induce any histological changes indicative of renal injury**

**a** Left: Representative histological images of Masson-Trichrome staining used to detect

fibrosis in renal tissues harvested from WT and LOX-1 KO mice. Right: Quantitative

analysis by Masson-Trichrome staining.

Data are represented as mean ± SEM. Differences were determined using one-way

ANOVA, followed by Tukey's multiple comparison test.

**b** Representative histological images of renal tissues harvested from WT and LOX-1 KO

mice stained for PAS to assess mesangial expansion and glomerular area, providing

insight into the structural integrity of the glomeruli.

Eight-week-old mice were fed an ND or an HFD for 6 weeks. From 10 weeks of age,

these mice were concurrently treated for 4 weeks with infusions of vehicle or Ang II (a

subpressor dose of 0.1 γ or a pressor dose of 0.7 γ) through subcutaneously implanted

osmotic pomps.

**Fig. S10| LOX-1 and AT1a were co-localized with megalin**

Representative images showing co-staining of LOX-1 (upper panels) and AT1 (lower

panels) with megalin in renal cortex tissues from WT mice. Nuclei are stained blue with

1   DAPI. Green signals indicate LOX-1 or AT1, while red signals indicate megalin. The

2   overlay images show the merged visualization of AT1 or LOX-1 with megalin and DAPI,

3   highlighting the predominant co-localization of LOX-1 and AT1 at the brush borders.

# Fig. 2

**Fig. 3**

# Fig. 4

**Fig. 5**

# Fig. 6

6a

6b

# Fig. 7

7a

7b

7c

7d

# Fig. 8

# Fig. 9

9a

9b

**Fig. 10**

**Fig. S1**

AT1-eGFP    LOX-1-mScarlet

N=283    N=185

5μm

N=283    N=185

N=167    N=125

N=167    N=125

1min

4min

00:00

# Fig. S2

# Fig. S3

**Fig. S4**

# Fig. S5

# Fig. S6

**Fig. S7**

# Fig. S8

# Fig. S9

## S9a
Masson-Trichrome staining

## S9b
PAS staining

# Fig. S10