Diurnal butterfly diversity in a human-modified landscape of the subtropical montane

forest of NW Argentina

Andrea del V. Guanuco

Universidad Nacional de Jujuy, San Salvador de Jujuy, Argentina. E-mail::

andruguanuco@gmail.com

Mariano Ordano

Fundación Miguel Lillo, Tucumán, Argentina; and Instituto de Ecología Regional (IER),

CONICET - Universidad Nacional de Tucumán, Yerba Buena, Tucumán, Argentina. E-mail:

maordano@lillo.org.ar

Laura C. Pereyra

Instituto de Ecorregiones Andinas (INECOA), CONICET – Universidad Nacional de Jujuy,

San Salvador de Jujuy, Argentina.E-mail: laureech@gmail.com

María José Barrionuevo

Instituto de Ecorregiones Andinas (INECOA), CONICET – Universidad Nacional de Jujuy,

San Salvador de Jujuy, Argentina. Correo electrónico: mariajosebarrionuevo@gmail.com

Noelia V. Gonzalez Baffa Trasci, Universidad Nacional de Jujuy, San Salvador de Jujuy,

Argentina.E-mail: novi.arg31@gmail.com

Marcos Vaira

Instituto de Ecorregiones Andinas (INECOA), CONICET – Universidad Nacional de Jujuy,

San Salvador de Jujuy, Argentina.E-mail: marcos.vaira@conicet.gov.ar

Running title: Butterfly diversity on a modified forest landscape

Abstract.

The change and intensification in land-use is currently among the main causes of

species declines and local extinctions around the world, therefore, forecasting changes in

species diversity concerning habitat conditions may be crucial for conservation strategies. We

explored the diurnal lepidopteran diversity in a modified landscape of subtropical montane

forests of Jujuy, NW Argentina. We considered that degradation of the natural forest habitat

would likely impact on resources crucial for butterflies, consequently altering both species

richness and composition within these forests. We assessed and compared alpha diversity

through Hill diversity profiles and beta diversity through the beta-diversity partitioning

method. Additionally, we employed a multivariate permutational analysis of variance, and

rank-abundance curves of butterfly species at different habitat types. Our results suggest that

land-use changes diminish the number of forest-dependent species and increase species more

tolerant to modified habitats and open areas. While alpha diversity did not decrease as land

use changed, beta diversity showed significant changes in the butterfly species composition

with a worrying reduction of forest-related species in altered habitats. Species composition

became increasingly dominated by open area butterfly species resulting in biota

homogenization, with possible consequences for ecosystem functioning and services of these

forests. Further research on the mechanisms underlying the effects of human-induced habitat

changes on forest butterfly diversity could help clarify which mitigation efforts are most

likely to be successful for the conservation of butterflies of the subtropical montane forests.

Keywords. Lepidoptera, butterflies, species richness, land use-change, abundance, Van

Someren Rydon trap, sweep net, diurnal, Nymphalidae, Jujuy.

Introduction

The transformation of large areas of native forests into mosaics of human-modified

habitats with different land uses is among the main drivers of biodiversity loss and currently

represents an important threat to species extinctions globally (Newbold *et al.* 2015; Picanco

et al. 2017). Therefore, forecasting changes in species diversity concerning habitat

deterioration may be crucial for conservation strategies. Several studies have found that

human-made disturbances on natural landscapes such as habitat loss and fragmentation are

among the most critical causes of species loss and declines of butterfly communities (Steffan-

Dewenter & Tscharntke 2000; Uehara-Prado *et al.* 2007; Vasconcelos *et al.* 2019).

Consequently, critical ecosystem processes may be affected, altering the ecosystem

functioning and also the provision of goods and services for humans (Klein *et al.* 2007;
Steffan-Dewenter & Westphal 2008; Potts *et al.* 2010). To these conditions, species
communities react differently, being able to persist or become extinct when the habitats are
smaller or very isolated (Barlow *et al*. 2009; Thomas 2016; Vasconcelos *et al*. 2019).
63   Among insect groups, butterflies have been pointed out as good biological models to
detect responses to land-use changes by several authors (Barlow *et al*. 2007ab; Lazzeri *et al*.
2011; Sagwe *et al*. 2015) mainly because they react sensitively and quickly to changes in
habitat conditions (Thomas *et al*. 2004; Thomas 2005; Bonebrake *et al*. 2010). In this way,
changes in ambient temperature, microclimate, humidity, and lighting (Checa *et al*. 2014), as
well as the composition and structure of vegetation can drive variation in species diversity
(Barlow *et al*. 2007b; Uehara-Prado & Freitas 2007). Unlike most other insect groups, its
systematics and taxonomy are relatively well documented (Thomas 2005; Lucci Freitas *et al*.
2014).

72   Several studies measuring the impacts of land-use changes have shown distinct effects
73   on species richness and composition such as local extinctions or lower persistence of certain
74   threatened or rare species and increases in the number of more non-threatened or widespread
75   species (Barlow *et al*. 2007ab; Bubová *et al.*, 2015; Thomas, 2016; Vasconcelos *et al*. 2019).
76   These responses have also been found not only at the level of the family taxa but also within
77   the same subfamily of butterflies (Veddeler *et al*. 2005).

78   The Southern Andean Yungas ecoregion in Argentina, defined as an outstanding
biodiverse region, is considered a critical area for biodiversity conservation because of the
levels of species richness and endemism that are among the highest in the country, but also
increasing land-use changes that threaten several species (Brown *et al.*, 2006). This ecoregion
covers only 2% of the country's land and is the second most diverse ecoregion (Pacheco &
Brown 2006; Nanni *et al*. 2020) that extends as a narrow strip of montane forest in the NW
region of the country, corresponding to the southernmost extension of Neotropical cloud
forests (Kapelle & Brown, 2001) that stands out for its high heterogeneity of environments
with important extensions of subtropical montane forests, small plots with subsistence
agriculture, logging sectors, large areas with commercial crops, and areas with different
degrees of urbanization (Malizia *et al.*, 2012). In such human-intervened landscapes, it will
be particularly relevant to investigate the response of local biodiversity to human pressures
and, thus, to estimate biodiversity changes at local scales to determine conservation priorities

(Reynolds *et al.*, 2011). However, the lack of information on how several groups of species

respond to such heterogeneity and land-use changes in these montane forests makes a great

challenge to the reversion of species declines or disappearances and the preservation and

restoration of suitable habitats to maintain the species diversity (Izquierdo & Grau, 2009).

Successful long-term conservation plans will require an understanding of the relationships

between land-use intensification and species composition to identify variables driving species

diversity in the region.

In this study, we describe diurnal lepidopteran diversity in a portion of the subtropical

montane forests of Jujuy, NW Argentina where human intervention has transformed the

entire area into a mosaic formed by habitat patches of different land-uses, such as remnants of

native forest, *Eucalyptus* plantations, and secondary grasslands. We expect that butterfly

assemblage of the area should be particularly sensitive to human-induced changes showing

the highest richness and densities at patches of native forests. Meanwhile, the species

assemblage will show a reduction in diversity at more altered patches.

We expect that butterfly assemblage of the area should be particularly sensitive to

human-induced changes, showing the highest richness and densities at patches of native

forests. Meanwhile, the species assemblage will show a reduction in diversity at more altered

patches.

Material and methods

Study area

The study was conducted at Finca "Las Capillas", (24° 05☐ 34'' S; 65° 09☐ 32'' W,

1160 m asl), located in the province of Jujuy, NW Argentina (Fig. 1). The average annual

temperature varies between 18 and 20 °C (Bianchi *et al*. 2005). This area is composed of a

mixed-use landscape of secondary forests and plantations of exotic *Eucalyptus* sp. (Entrocassi

2015).

We selected habitat patches defined by three dominant land-uses types: (1) Secondary

native forest (FO) area, covering approximately 0.08 km^2. This has undergone wood

extraction through selective logging and has low-intensity livestock grazing. This was

considered as a relatively preserved natural forest with minimal land-use change (2)

Eucalyptus plantation (EU) area, covering approximately 0.065 km², with a low-intensity

livestock grazing, and (3) Secondary grassland (GR) area, located at a mixed-use flat area

covering approximately 0.10 km^2, this habitat type was originated following the clearance

and burning of the primary forest. Poaceas dominate it and have moderate livestock grazing

(Fig. 1).

Data collection

Butterfly surveys were carried out between September and May 2016-2018, in

coincidence with the period of higher adult butterfly activity in the region and favourable

weather conditions. To ensure a comprehensive assessment of diversity, we employed a

combination of hand nets and bait trap survey techniques, as both methods are considered

complementary (Sparrow *et al.* 1994; Checa *et al.* 2019). In each habitat patch, we placed

twenty 50 mts-long transects. We set two Van Someren Rydon bait traps (VSR) (DeVries

1987, 1988; Villareal *et al.* 2004) located at 10 and 40 meters from the start of each transect,

accounting for a total of 40 traps per habitat type. The VSRs, manufactured according to the

specifications of DeVries (1987, 1988), were slightly modified (i.e. replacing the metal plate

of the base with a plastic container). The bait traps consisting of decaying fruit (banana,

apple, peach, mango, pear, based on seasonal availability in the local market) were suspended

1.5-2.0 m above the ground. The traps were operational for three consecutive days (72 hours)

and visited every 24 hours. Additionally, we conducted entomological net sampling (Villareal

et al. 2004) along the transects on the second day after activating the VSR traps. Two

researchers sampled flying butterflies using a sweep net for 15 minutes along a standard line

transect of 2 m wide (Pollard & Yates 1993), between 11:00 am and 4:00 pm, in coincidence

with the peak of butterfly activity (Villareal *et al.* 2004).

Trapped butterflies were carefully placed in paper envelopes. Subsequently,

specimens were kept in a humid chamber for 24 to 48 hours for later assembly (Villareal *et*

al. 2004). All collected material was examined and identified at the species levels following

Canals (2000, 2003), Lamas (2004), Volkmann & Núñez-Bustos (2010) and Klimaitis *et al.*

(2018ab). In addition, we reviewed the database of butterflies of America (Warren *et al.*

2023). We followed the taxonomy proposed by Mielke (2008) for the Hesperiidae family,

and Lamas (2004) for the remaining Papilionoidea. Approximately three hundred specimens

were prepared and deposited at the "Entomological Collection of the Instituto de Biología de

la Altura Dra. Liliana Estela Neder (CEINBIAL)", Universidad Nacional de Jujuy, Jujuy,

Argentina.

Statistical analyses

To evaluate the adequacy of sampling effort at each habitat patch, we employed

measures of sample coverage following Chao & Jost (2012). Coverage represents the

percentage of individuals in an assemblage that belong to a species represented in the sample;

it ranges from 0 to 100% (Chao & Jost 2012). As this value is a measure of inventory

completeness, when coverage is close to 100%, the sample is near complete and diversity

values (^{q}D) can be compared directly (Chao & Jost 2012). We constructed diversity profiles

based on Hill numbers (^{q}D) to assess the changes in cumulative species diversity among the

three habitat patches. We considered q-values from 0 to 2 at increments of 0.25 and

computed the 95% confidence intervals. Significant differences in species diversity between

two assemblages were inferred when the confidence intervals of the diversity profile curves

did not overlap.

We compared species composition among the three habitat patches with the beta-

diversity partitioning method following Baselga (2010). This method partitions the total beta

diversity into its components of nestedness and species turnover. Nestedness refers to the

overlap of species occurrence among habitat patches, indicating that species composition of a

given assemblage is a subset of the species composition of a larger assemblage (Ulrich &

Gotelli 2007). Conversely, species turnover refers to the replacement of species in one habitat

patch by different ones in another habitat, leading to less frequent or even segregated species

occurrence, with many species never co-occurring (Ulrich & Gotelli 2007). To further

examine the assemblage structure between habitat patches, we applied a multivariate

permutational analysis of variance on Bray-Curtis distances (PERMANOVA; Anderson

2001). Lastly, we assessed changes in community structure with rank-abundance curves

based on the log-transformed relative abundances of butterfly species in each habitat patch.

These curves visually compare species richness, relative abundance, dominance, and the

sequence of each species without losing their identity (Feinsinger 2004). We statistically

tested for differences in rank☐abundance distribution between habitat patches using

Kolmogorov–Smirnov two☐sample tests under the null hypothesis that two samples can be

generated from the same distribution.

All statistical analyses were conducted using R 3.6.1 (R Development Core Team,

2019). The database, R script, and packages are available as supplementary material

(Supplemental information, Tables S1 and S2).

Results

The combined diurnal butterfly collection comprised 3914 individuals, representing

an overall diversity of 78 species distributed in six families (Table 1), including Hesperiidae

(with two subfamilies: Eudaminae and Pyrginae), Nymphalidae (with nine subfamilies:

Apaturinae, Biblidinae, Charaxinae, Danainae, Heliconiinae, Libytheinae, Limenitidinae,

Satyrinae, and Nymphalinae), Pieridae (with two subfamilies: Coliadinae and

Dismorphiinae), and Lycaenidae, Riodinidae, and Papilionidae, with only one subfamily

each (Theclinae, Riodininae, and Papilioninae, respectively).

Sample coverage estimates exceeded (90%) for each habitat patch. Diversity profiles

based on Hill numbers (CI: 95%) showed that the GR exhibited the highest diversity with 74

species, followed by the EU with 41 species, and FO with 18 species (Table 1, Fig. 2).

Butterfly species composition differed significantly among the three habitat patches

(PERMANOVA, F= 30.69, P <0.001). The FO habitat presented a significant dissimilarity in

species composition with both, GR and EU habitats, slightly above 60% (PERMANOVA F=

59.82, P <0.001; F= 30.44, P= 0.001, respectively). Observed differences between FO and

GR were given mainly by a nestedness process, indicating that the species composition of

forest assemblage is a subset of the one found in the grassland (Fig 3.)(Fig. 5). Conversely,

beta diversity observed between FO and EU was predominantly due to species turnover, with

few species co-occurring in both habitat patches (Fig. 3). Finally, GR and EU presented a

relatively lower difference in species composition (Fig. 3), with a statistically significant

dissimilarity value of 32% (PERMANOVA F= 10.45, P < 0.001). This dissimilarity was

highly influenced by a nestedness process, indicating that the species composition of the

Eucalyptus sp. assemblage constitutes a subset of the grassland community (Fig. 3).

The rank-abundance distribution was more even in the grassland assemblage, with

three species accounting for over 50% of all individuals registered in this habitat (Fig. 4). In

GR, *Hermeuptychia* sp. Forster, 1964 (Nymphalidae: Satyrinae) was the most abundant

species (21.7%), followed by *Biblis hyperia nectanabis* (Fruhstorfer, 1909) (Nymphalidae:

Biblidinae) (19.6%), and *Cybdelis petronita* Burmeister, 1861 (Nymphalidae: Biblidinae)

(11.5%). The EU also presented three species summing more than 50% of the individuals

registered in this habitat (Fig. 4): *H.* sp., was again the most abundant species in the

assemblage (28.7%), followed by *C. petronita* (12.8%), and *B. hyperia nectanabis* (11.2%).

In contrast, FO presented two dominant species, which accounted for 90% of the individuals

registered for this habitat: *Pedaliodes palaepolis,* (Hewitson, 1878) (Nymphalidae: Satyrinae)

(53.2%), and *Eunica tatila bellaria* Fruhstorfer, 1908 (Nymphalidae: Biblidinae) (36.8%)

(Fig. 4). The Kolmogorov-Smirnov test showed a statistically significant shift in species

rank-abundance distribution between the three habitat patches (FO vs. GR: D= 0.72, P

<0.001, FO vs. EU: D= 0.22, P= 0.002, and GR vs. EU D= 0.42, P <0.001).

Discussion

In this study, we documented the diversity of a diurnal butterfly community across habitat

patches with different land-use within a sector of the subtropical montane forest of NW

Argentina (Yungas ecoregion). Our results suggest that the impacts of land-use change have a

strong effect on the diurnal butterfly assemblage of the area. Contrary to our initial

expectations, the butterflies assemblage did not show a reduction in species richness with

increased land-use change. Instead, we observed the opposite trend, revealing clear

differences in alpha diversity among the three habitat patches, where GR emerged as the most

diverse habitat, followed by EU and, lastly, FO. The highest diversity registered in GR can be

attributed to the structure of the vegetation found in this patch, which includes a small

remnant of native forest and open areas with a variety of herbaceous plants. This habitat

heterogeneity likely provides a large amount of resources and niches for several species.

238   Previous studies established that species belonging to the Satyrinae and

Biblidinae subfamily can display diverse responses to habitat disturbances, with some

species being highly sensitive to such changes, while others became more abundant

following alterations (Shahabuddin & Terborgh 1999; Fermon et al. 2005; Barlow et

al. 2007ab; Barlow et al. 2009; Uehara-Prado et al. 2007; Ribeiro et al. 2012;

Vasconcelos et al. 2019). This aligns with our results, where *Hermeuptychia* sp.

emerged as the principal satirine species successfully occupying patches of altered

habitat compared to the more conserved secondary native forest. Modified

environments appeared to be suitable for opportunistic species tolerant to structural

changes in the vegetation. The genus *Hermeuptychia*, has been geographically

expanding in recent years due to its great adaptability, the capacity of larvae to feed on

a variety of pioneer plant species belonging to grasses, such as *Poa*, *Paspalum*, and

Stenotaphrum (Nuñez-Bustos et al. 2010). Therefore, these modified habitats may not

only host more satyrins species, but also support a higher abundance per species

(Brown 1992). This is consistent with previous studies that identified the genus,

Hermeuptychia and *Biblis* as typical invasive species of open areas (Ramos 2000). In

addition to *B. hyperia nectanabis*, *C. petronita* was also abundant in GR and EU

habitat patches, probably due to the association with the type of host plant used during

larval development. These butterfly species feed mainly on herbaceous plants of the

genus Tragia (Freitas et al. 1997; Bauza 2000; Nuñez-Bustos 2010), which are

perennials and typically inhabit open areas, such as cultivated fields or stream banks

(Romero & Sanguinetti 1989).

Species from the Pieridae family, such as *Abaeis albula albula* (Cramer, 1775),

Teriocolias deva deva (E. Doubleday, 1847), *Phoebis sennae marcelina* (Linnaeus,

1758), *Pyrisitia nise floscula* (Cramer 1775), *Anteos clorinde* (Godart 1824), *Abaeis*

arbela arbela (Geyer 1832), and *Pseudopieris nehemia prasina*, Hayward, 1949

(Nuñez-Bustos 2010; Klimaitis et al. 2018a; Klimaitis et al. 2018b, Warren et al.

2023), recorded in the habitat patches with the most remarkable land-use

intensification, suggest adaptations that enable them to colonize different

environments. Their rapid flight, the use of a variety of nutritional plants, and

preference for open or edge areas contribute to their capability to occupy extensive

modified areas (Ospina López et al. 2010). Other species tolerant to modified

environments, such as "jumping" hesperids *Burnsius orcus* (Stoll, 1780), *Burnsius*

orcynoides (Giacomelli, 1928), *Urbanus proteus proteus* (Linnaeus, 1758), and certain

nymphalids of the subfamily Nymphalinae, represented by heliophilous species of

open areas such as *Junonia genoveva hilaris* C. Felder & R. Felder, 1867, *Chlosyne*

lacinia saundersi (E. Doubleday, [1847]), *Tegosa claudina* (Eschscholtz, 1821),

Telenassa berenice drusinilla (Röber, 1913), and *Vanessa carye* (Hübner, 1812), may

colonize different habitats due to their lack of specific habitat requirements (Pastrana et

al. 2004; Nuñez-Bustos 2010; Volkmann & Nuñez-Bustos 2010; Klimaitis et al. 2018a;

Klimaitis et al. 2018b, Warren et al. 2023). We also recorded species from the

Charaxinae subfamily in the modified habitats patches (GR and EU). These

lepidopterans, with their large bodies and high mobility, can move long distances

(Barlow et al. 2009; Freire et al. 2021), allowing them to occupy various habitat types.

The absence or reduced abundance of *P. palaepolis* and *E. tatila bellaria* in the GR

and EU habitats suggests that these diurnal frugivorous lepidopterans, can respond sensitively

to habitat disturbances. *P. palaepolis*, was recognized as a typical species of montane forest

of the Yungas region (F. Moschioni, pers. comm.). Consequently, the heightened species

richness in GR could be attributed to an increase in species tolerant to modified habitats and

open areas, resulting in a mechanism of compensatory dynamics, which likely masks the

negative effect of land-use change on diversity (Supp & Ernest 2014; Filgueiras *et al*. 2019).

Similarly, we considered EU a heterogeneous habitat despite exhibiting a simplified

vegetation structure, including small portions of native forest and open areas, the latter used

as cattle pasture. Consequently, species richness in this habitat could also be compensated by

common and abundant species, such as *H. sp.*, *Yphthimoides celmis*, Godart, (1824), and

Cissia phronius Godart (1824). These findings were consistent with studies carried out in

different regions of Brazil, such as the Amazon Region and the Brazilian Atlantic Forest in

Southeast Brazil (Ramos 2000; Ribeiro *et al.* 2012; Vasconcelos *et al.* 2019), where the same

genera were registered as abundant and common.

Despite the structural simplification observed in EU the presence of tree canopy

coverage may enable certain species associated with canopy habitats to tolerate land-use

changes, explaining the occurrence of species linked to the forest canopy (DeVries 1987).

However, this structural simplification could lead to biotic homogenization, often acting as a

filter for those species with specific habitat requirements (Steffan-Dewenter & Tscharntke

2000), particularly understory butterflies associated with the vertical stratification of a forest,

and favoring species capable of using a new spectrum of plants (Ekroos *et al.* 2010; Borschig

et al. 2013). Furthermore, the absence or decline of certain dominant forest species in EU

suggests that some land uses may not adequately preserve local diversity.

Studying species composition through beta diversity measures revealed

significant changes in butterfly assemblages, with a concerning reduction of forest-

related species. GR and EU, representing higher land-use intensification, exhibited

greater similarities to each other, and both differed significantly in community

composition from the FO. A nestedness process mainly determined the observed

difference in species composition of butterfly assemblages between GR and FO. This

suggests that while GR maintains a set of butterfly species relatively similar to the FO,

new structural conditions might facilitate the occurrence of a new set of species

capable of exploiting highly modified habitats. This may result in a butterfly

community increasingly dominated by species adapted to modified environments and

open areas, leading to biota homogenization with significant consequences for

ecosystem functioning and services of the Yungas forest. On the other hand, the beta

diversity observed between FO and EU was mainly driven by a species turnover

process.

Comparison of the community structure among the three land-use areas showed

not only a significant difference in species composition but, a variation in the dominant

species within each habitat type. Species that were abundant in GR and EU, such as *H.*

sp. (Satyrinae), *B. hyperia nectanabis* (Biblidinae), and *C. petronita* (Biblidinae), were

practically absent in the secondary native forest. Conversely, dominant species in FO,

such as *P. palaepolis* (Satyrinae) and *E. tatila bellaria* (Biblidinae) were less abundant or even absent in more modified habitats. *E. tatila bellaria*, is characteristic of humid forests associated with habitats of a complex vegetation structure. Its larvae feed on a native tree from the Euphorbiaceae family, specifically the genus *Sebastiania* (Nuñez-Bustos 2010), dominant in the Montane forest strata of the Yungas (Cabrera 1994). Cavanzón-Medrano et al. (2003), showed that *E. tatila* is a neotropical species identified as a bioindicator with significant relevance in undisturbed areas of the Yucatan Peninsula in Mexico. The species predominantly associates with the arboreal stratum, providing essential food resources for the adults (Salazar 2009). Consequently, the dominance of *P. palaepolis* and *E. tatila bellaria* in FO could be considered indicative of low levels of habitat alteration in forest habitats. Montane forest likely serves as a source of host plants, fleshy fruits, and organic matter for larvae and adult stages of these frugivorous species. Our results suggest that, alongside these two species, other forest-dependent butterflies present in the area like *Taguaiba ypthima* (Hübner, 1821), may also be affected by land-use changes. Although its classification as a forest species (DeVries 1987), coincides with our results given that it was only recorded in FO.

To our knowledge, our work represents the first study on the response of diurnal butterflies to land-use changes in the Southern AndeanYungas ecoregion of Argentina. This research constitutes a first step for future regional-scale studies in this field. Additional studies are needed to assess the impacts of habitat composition and configuration on species diversity in the studied region. Further research into the mechanisms driving the effects of land-use changes on forest-dependent butterfly diversity could provide insights into which mitigation strategies are most likely to succeed in conserving butterfly diversity in theYungas of Argentina. To establish the amount of forest vital to protect even a fraction of the original biodiversity holds value for local butterfly conservation. The dispersal abilities of most species and the distances they travel, crucial for species to cope with alterations in habitat conditions, are still poorly known in the region. Additionally, specific habitat requirements and functional traits of the species, such as body size or mobility, may determine whether a particular species benefits from or is adversely affected by intense land-use changes.

Acknowledgements

We thank the owners and Juan José Correa, in charge of the "Las Capillas" farm,

359 Sergio José Flores for giving us a place to stay, my friends, family and parents for

360 accompanying me in all the campaigns carried out. Fundación Miguel Lillo partially

361 supported the research of MO (Projects Z-0113-1 and Z-048-1). This project was partially

362 supported by a PIO Grant Conicet #14020140100094CO and a PUE Grant Conicet

363 (#22920170100027CO).

References

Anderson M. J. 2001. A new method for non-parametric multivariate analysis of
variance. *Austral Ecology* 26(1): 32-46. https://doi.org/10.1111/j.1442-
9993.2001.01070.pp.x

Bauza J. A. T. 2000. Ciclo de vida de *Biblis hyperia* (Cramer) en Puerto Rico
(Lepidoptera:Nymphalidae). *Caribbean Journal of Science* 36(1/2): 87-93.

Barlow J. O. S., Overal W. L., Araujo I. S., Gardner T. A. & Peres C. A. 2007a. The
value of primary, secondary and plantation forests for fruit□feeding butterflies in
the Brazilian Amazon. *Journal of Applied Ecology* 44(5): 1001-1012.

Barlow J., Gardner T. A., Araujo I. S., Ávila-Pires T. C., Bonaldo A. B., Costa J. E.,
Esposito M. C., Ferreira L. V., Hawes J., Hernandez M. I. M., Hoogmoed M. S.,
Leite R. N., Lo-Man-Hung N. F., Malcolm J. R., Martins M. B., Mestre L. A. M.,
Miranda-Santos R., Nunes-Gutjahr A. L., Overal W. L., Parry L., Peters S. L.,
Ribeiro-Junior M. A., da Silva M. N. F., da Silva Motta C. & Peres C. A. 2007b.
Quantifying the biodiversity value of tropical primary, secondary, and plantation
forests. *Proceedings of the National Academy of Sciences* 104(47): 18555-18560.

Barlow J., Araujo I. S., Overal W. L., Gardner T. A., da Silva Mendes F., Lake I. R.
& Peres C. A. 2009. Diversity and composition of fruit-feeding butterflies in
tropical *Eucalyptus* plantations. *Plantation Forests and Biodiversity: Oxymoron*
or Opportunity? 9: 165-180.

Baselga A. 2010. Partitioning the turnover and nestedness components of beta
diversity. *Global ecology and biogeography* 19(1): 134-143.
https://doi.org/10.1111/j.1466-8238.2009.00490.x

Bianchi A., Yañez C., Acuña L. 2005. *Base de datos mensuales de precipitaciones*
del Noroeste Argentino. Editorial INTA, EEA. Salta-Jujuy.

Bonebrake T. C., Ponisio L. C., Boggs C. L. & Ehrlich P. R. 2010. More than just
indicators: a review of tropical butterfly ecology and conservation. *Biological*
conservation 143(8): 1831-1841.

Börschig C., Klein A. M., von Wehrden H. & Krauss J. 2013. Traits of butterfly
communities change from specialist to generalist characteristics with increasing
land-use intensity. *Basic and Applied Ecology* 14(7): 547-554.

Brown Jr. K. S. & Brown G. G. 1992. Habitat alteration and species loss in Brazilian
forests. *Tropical Deforestation and Species Extinction* 10: 119-142.

Brown A. D., Pacheco S., Lomáscolo T. & Malizia L. 2006. Situación ambiental en

los bosques andinos yungueños. In La situación ambiental Argentina 2005: 52–72. Brown A. D., Ortíz U. M., Acerbi M. & Corcuera J. (Eds). Buenos Aires: Fundación Vida Silvestre Argentina.

Bubová T., Vrabec V., Kulma M. & Nowicki P. 2015. Land management impacts on European butterflies of conservation concern: A review. *Journal of Insect Conservation* 19: 805-821.

Cabrera A. L. 1994. *Regiones Fitogeográficas Argentinas*. Editorial Acme S.A.C.I., Buenos Aires, Argentina. Reimpresión. 85 pp.

Canals G.R. 2000. *Mariposas bonaerenses*. LOLA, Buenos Aires, 347pp.

Canals G.R. 2003. *Mariposas de Misione*s. LOLA, Buenos Aires, 325pp.

Cavanzón-Medrano L. E., Machkour-M'Rabet S., Chablé-Iuit, L. R., Pozo C., Hénaut Y. & Legal L. 2018. Effect of Climatic Conditions and Land Cover on Genetic Structure and Diversity of *Eunica tatila* (Lepidoptera) in the Yucatan Peninsula, Mexico. *Diversity* 10(3): 1-79.

Chao A. & Jost L. 2012. Coverage□based rarefaction and extrapolation: standardizing samples by completeness rather than size. *Ecology* 93(12): 2533-2547.

Checa M. F., Rodriguez J., Willmott K. R., & Liger B. 2014. Microclimate variability significantly affects the composition, abundance and phenology of butterfly communities in a highly threatened neotropical dry forest. *Florida Entomologist* 97(1): 1-13.

Checa M. F., Donoso D. A., Rodriguez J., Levy E., Warren A. & Willmott K. 2019. Combining sampling techniques aids monitoring of tropical butterflies. *Insect Conservation and Diversity* 12(4): 362-372.

DeVries P. J. 1987. *Butterflies of Costa Rica and their Natural History: Papilionidae, Pieridae, Nymphalidae*. Vol. 1. New Jersey: Princeton. University Press, 97pp.

DeVries P. 1988. Stratification of fruit-feeding nymphalid butterflies in a Costa Rica rainforest. *Journal of Research Lepidoptera* 26(1-4): 98-108.

Ekroos J., Heliölä J. & Kuussaari M. 2010. Homogenization of lepidopteran communities in intensively cultivated agricultural landscapes. *Journal of Applied Ecology* 47(2): 459-467.

Entrocassi F.G. 2015. Estudio de los bosques subtropicales de montaña de la Reserva Ecológica de Uso Múltiple Serranías de Zapla (Jujuy, Argentina): Composición florística, distribución de la vegetación y caracterización bioclimática. Tesis Doctoral. Universidad Complutense de Madrid, Fac. Farmacia, Dept. Biología Vegetal II.

Feinsinger P. 2004. *El Diseño de Estudios de Campo para la Conservación de la Biodiversidad*. Editorial FAN, Santa Cruz de la Sierra, Bolivia. 454 pp.

Fermon H., Waltert M., Vane-Wright R. I. & Mühlenberg M. 2005. Forest use and vertical stratification in fruit-feeding butterflies of Sulawesi, Indonesia: impacts for conservation. *Biodiversity & Conservation* 14: 333-350.

Filgueiras B. K., Melo D. H., Uehara-Prado M., Freitas A.V. L., Leal I. R. & Tabarelli M. 2019. Compensatory dynamics on the community structure of fruit-

feeding butterflies across hyper-fragmented Atlantic forest habitats. *Ecological Indicators* 98: 276-284.

Freitas A. V. L., Brown Jr. K. S. & Otero L. D. 1997. Juvenile stages of *Cybdelis*, a key genus uniting the diverse branches of the *Eurytelinae* (Lepidoptera: Nymphalidae). *Tropical Lepidoptera Research* 8(1): 29-34.

Freire Jr G. B., Silva T., Oliveira H., Collier C., Rodrigues H. P., Dias J. P., Diniz I. R., Santos J. P., Marini-Filho O. J., Freitas A. V. L., Smilanich A. M. & Dyer L. A. 2021. Good things come in larger packages: size matters for adult fruit-feeding butterfly dispersal and larval diet breadth. *Diversity* 13(12): 664.

Izquierdo A. E. & Grau H. R. 2009. Agriculture adjustment, land-use transition and protected areas in Northwestern Argentina. *Journal of Environmental Management* 90(2): 858-865.

Kappelle M. & Brown A. D. 2001. *Bosques nublados del Neotrópico*. Editorial IMBIO, Costa Rica. 397-442pp.

Klein A. M., Vaissiere B.E., Cane J. H., Steffan-Dewenter I., Cunningham S.A., Kremen, C. & Tscharntke T. 2007. Importance of pollinators in changing landscapes for world crops. *Proceedings of the Royal Society B: Biological Sciences* 274(1608): 303-313.

Klimaitis J. F., Moschione F. N. & Klimaitis C. L. 2018a. *Maravillas aladas de Argentina: Mariposas de la puna a la Patagonia*. Ecoval, Córdoba, Argentina.454 pp.

Klimaitis J. F., Nuñez Bustos E. O., Klimaitis C. L. & Guller, R. M. 2018b. *Mariposas de Argentina: Guía de identificación.* Vazquez Mazzini. Buenos Aires, Argentina. 327 pp.

Lamas G., 2004. Checklist: Part 4 A. Hesperioidea-Papilionoidea. *Atlas of Neotropical Lepidoptera: XXXVI* (ed. by Heppner, J.B.), Association for Tropical Lepidoptera, Gainesville, USA. 1- 439.

Lazzeri M. G., Bar M. E. & Pieri Damborsky M. 2011. Diversidad del orden Lepidoptera (Hesperioidea y Papilionoidea) de la ciudad Corrientes, Argentina. *Revista de Biología Tropical* 59(1): 299-308.

Lucci Freitas A. V., Agra Iserhard C., Pereira Santos J., Oliveira Carreira I. J. Y. Bandini Ribeiro D., Alves Melo D. H. A., Rosa-Batista H. A. Marini-Fhilo J. O. Accacio G. M. & Uehara-Prado M. 2014. Studies with butterfly bait traps: an overview. *Revista Colombiana de Entomología* 40(2): 203-212.

Malizia L., Pacheco S., Blundo C. & Brown A.D. 2012. Caracterización altitudinal, uso y conservación de las Yungas Subtropicales de Argentina. *Ecosistema*s 21(1-2): 53-73.

Mielke O. H. H., Emery E. D. O. & Pinheiro C. E. G. 2008. As borboletas Hesperiidae (Lepidoptera, Hesperioidea) do Distrito Federal, Brasil. *Revista Brasileira de Entomologia* 52(2): 283-288.

Nanni A. S., Piquer Rodríguez M., Rodriguez M. D., Núñez Regueiro M. M., Periago M. E., Aguiar S., Ballari S A., Blundo C. M., Derlindati E. J., Di Blanco Y. E., Eljall A., Grau H. R., Herrera L. P. Huertas-Herrera A., Izquierdo A. E., Lescano J. N., Machi L., Mazzini F., Milkovic M., Montti L. F. Paviolo A. J.,

Pereyra M., Quintana R. D., Quiroga V. A., Renison D., Beade S. M., Schaaf A. A. & Gasparri N. I. 2020. Presiones sobre la conservación asociadas al uso de la tierra en las ecorregiones terrestres de la Argentina. *Ecología Austral* 30:304-320.

Newbold T., Hudson L. N., Hill S. L. L., Contu S., Gray C. L., Scharlemann J. P. W., Börger L., Phillips H. R. P., Sheil D., Lysenko I. & Purvis A. 2016. Global patterns of terrestrial assemblage turnover within and among land uses. Ecography 39: 1151–1163.

Núñez Bustos E. 2010. *Mariposas de la Ciudad de Buenos Aires y alrededores*. Buenos Aires, Argentina, Vázquez Mazzini. Buenos Aires, Argentina. 264 pp.

Ospina-López L. A., García-Pérez J. F., Villa-Navarro F. A. & Reinoso-Flórez G. 2010. Mariposas Pieridae (Lepidóptera: Papilionoidea) de la Cuenca del Río Coello (Tolima), Colombia. *Actualidades Biológicas* 32(93): 173-188.

Pacheco S. y Brown A. 2006. Ecología y producción de cedro (Género Cedrela) en las Yungas australes. Ediciones del Subtrópico. Tucumán. Argentina. 9-18pp.

Pastrana J. A. 2004. *Los lepidópteros argentinos. Sus plantas hospedadoras y otros sustratos alimenticios*. Sociedad Entomológica. Buenos Aires, Argentina. 334 pp.

Picanco A., Rigal F., Matthews T. J., Cardoso P. & Borges P. A. 2017. Impact of land use change on flowers□visiting insect communities on an oceanic island. *Insect Conservation and Diversity* 10(3): 211-223.

Pollard E. & Yates T. J. 1993. *Monitoring butterflies for ecology and conservation: The British butterfly monitoring scheme*. Springer Science & Business Media. Londres. 1-61pp.

Pott, S. G., Biesmeijer J. C., Kremen C., Neumann P., Schweiger O. & Kunin, W. E. 2010. Global pollinator declines: trends, impacts and drivers. *Trends in Ecology & Evolution* 25(6): 345-353.

Ramos F. A. 2000. Nymphalid butterfly communities in an amazonian forest fragment. *Journal of Research on the Lepidoptera* 35(1): 29-41.

Reynolds C., Fletcher R. J., Carneiro C. M., Jennings N., Ke A., LaScaleia M. C., Lukhele M. B., Mamba M. L., Sibiya M. D., Austin J. D., Magagula C. N., Mahlaba T., Monadjem A., Wisely S. M. & McCleery R. A. 2018. Inconsistent effects of landscape heterogeneity and land-use on animal diversity in an agricultural mosaic: a multi-scale and multi-taxon investigation. *Landscape ecology* 33: 241-255.

Ribeiro D. B., Batista R., Prado P. I. Brown K. S. & Freitas A. V. 2012. The importance of small scales to the fruit-feeding butterfly assemblages in a fragmented landscape. *Biodiversity and Conservation* 21: 811-827.

De Romero M. E. M. & De Sanguinetti M. M. G. 1989. Actualizacion taxonomica de Tragia (Euphorbiaceae) para Argentina Y regiones limítrofes. *Darwiniana* 29(1/4): 77-138.

Salazar J. A. 2009. *Catálogo de Las Plantas Huésped de Las Mariposas Neotropicales*. Boletín Científico Centro De Museos De Historia Natural, 13, 230-232.

Sagwe R. N., Muya S. M. & Maranga R. 2015. Effects of land use patterns on the diversity and conservation status of butterflies in Kisii highlands, Kenya. *Journal

of insect conservation 19: 1119-1127.

Shahabuddin G. & Terborgh J. W. 1999. Frugivorous butterflies in Venezuelan
forest fragments: abundance, diversity and the effects of isolation. *Journal of*
Tropical Ecology 15(6): 703-722.

Sparrow H. R., Sisk T. D., Ehrlich P. R. & Murphy D. D. 1994. Techniques and
guidelines for monitoring neotropical butterflies. *Conservation Biology* 8(3): 800-
809.

Steffan□Dewenter I. & Tscharntke T. 2000. Butterfly community structure in
fragmented habitats. *Ecology Letters* 3(5): 449-456.

Steffan□Dewenter I. & Westphal C. 2008. The interplay of pollinator diversity,
pollination services and landscape change. *Journal of Applied Ecology* 45(3):
737-741.

Supp S. R. & Ernest S. M. 2014. Species□level and community□level responses to
disturbance: a cross□community analysis. *Ecology* 95(7): 1717-1723.

Thomas J. A., Telfer M. G., Roy D. B., Preston C. D., Greenwood J. J. D., Asher J.
& Lawton J. H. 2004. Comparative losses of British butterflies, birds, and plants
and the global extinction crisis. *Science* 303(5665): 1879-1881.

Thomas J. A. 2005. Monitoring change in the abundance and distribution of insects
using butterflies and other indicator groups. *Philosophical Transactions of the*
Royal Society B: Biological Sciences 360(1454): 339-357.

Thomas J. A. 2016. Butterfly communities under threat. *Science* 353(6296): 216-
218.

Uehara□Prado M., Brown Jr. K. S. & Freitas A. V. L. 2007. Species richness,
composition and abundance of fruit□feeding butterflies in the Brazilian Atlantic
Forest: comparison between a fragmented and a continuous landscape. *Global*
Ecology and Biogeography 16(1): 43-54.

Ulrich W. & Gotelli J. N. 2007. Disentangling community patterns of nestedness
and species co□occurrence. *Oikos* 116(12): 2053-2061.

Vasconcelos R. N., Cambui E. C. B., Mariano-Neto E., da Rocha P. L. B. & Cardoso
M. Z. 2019. The role of *Eucalyptus* planted forests for fruit-feeding butterflies'
conservation in fragmented areas of the Brazilian Atlantic forest. *Forest Ecology*
and Management 432: 115-120.

Veddeler D., Schulze C. H., Steffan-Dewenter I., Buchori D. & Tscharntke T. 2005.
The contribution of tropical secondary forest fragments to the conservation of
fruit-feeding butterflies: effects of isolation and age. Biodiversity & Conservation
14: 3577-3592.

Villareal H. M., Álvarez M., Córdoba-Córdoba S., Escobar F., Fagua G., Gast F. &
Umaña A. M. 2004. *Manual de métodos para el desarrollo de inventarios de*
biodiversidad. Instituto de Investigación de Recursos Biológicas Alexander von
Humboldt, Bogotá, Colombia. 454 pp.

Volkmann L. & Núñez Bustos E. 2010. M*ariposas Serranas de Argentina Central:*
guía de especies más comunes halladas en sierras, valles y salinas del centro
oeste argentino (Córdoba, San Luis, La Rioja, Catamarca y Santiago del Estero).
Equipo gráfico, Córdoba, Argentina. 454 pp.

Warren A. D., Davis K. J., Stangeland E. M., Pelham J. P. & Grishin N. V. 2023.

Illustrated Lists of American Butterflies. [12-X-2023]. < http://www.

butterfliesofameric

Table 1. List of butterfly species recorded at three habitat patches of Finca "Las Capillas",

(24° 05 34'' S; 65° 09 32'' W, 1160 m asl), Jujuy, Argentina. **FO=** Secondary native

forest; **EU=** *Eucalyptus* plantation; GR= Secondary grassland.

Family	Subfamily	Species	FO	EU	GR
Hesperiidae	Eudaminae	*Urbanus proteus proteus* (Linnaeus, 1758)		x	x
	Eudaminae	*Spicauda teleus* (Hübner, 1821)			x
	Pyrginae	*Burnsius orcus (Stoll, 1780)*		x	x
	Pyrginae	*Burnsius orcynoides* (Giacomelli, 1928)			x
Lycaenidae	NA	*sp1*			x
	Theclinae	*Chlorostrymon simaethis* (Drury, 1773)			x
	Theclinae	*Micandra sylvana* (Jörgensen, 1934)			x
	Theclinae	*Pseudolycaena marsyas* ((Linnaeus, 1758)			x
Nymphalidae	Apaturinae	*Doxocopa cyane burmeisteri* (Godman & Salvin, 1884)	x	x	x
	Biblidinae	*Hamadryas epinome* (C. Felder & R. Felder, 1867)		x	x
	Biblidinae	*Hamadryas fornax fornax* (Hübner, [1823])		x	
	Biblidinae	*Biblis hyperia nectanabis* (Fruhstorfer, 1909)	x	x	x
	Biblidinae	*Mestra hersilia apicalis* (Staudinger, 1886)			x
	Biblidinae	*Callicore sorana sorana* (Godart, [1824])			x
	Biblidinae	*Diaethria neglecta neglecta* (Salvin, 1869)		x	x
	Biblidinae	*Haematera pyrame pyrame* (Hübner, [1819])		x	x
	Biblidinae	*Cybdelis petronita* Burmeister, 1861	x	x	x
	Biblidinae	*Eunica tatila bellaria* Fruhstorfer, 1908	x	x	x
	Biblidinae	*Epiphile orea negrina* C. Felder & R. Felder, 1862	x		x
	Charaxinae	*Fountainea glycerium cratais* (Hewitson, 1874)	x		x
	Charaxinae	*Fountainea nessus* (Latreille, [1813])		x	x
	Charaxinae	*Memphis moruus morpheus* (Staudinger, [1886])		x	x
	Charaxinae	*Archaeoprepona chromus chromus* (Guérin-Méneville, [1844])			x
	Charaxinae	*Archaeoprepona demophoon thalpius* (Hübner, [1814])			x
	Danainae	*Danaus eresimus plexaure* (Godart, 1819)			x
	Danainae	*Danaus erippus* (Cramer, 1775)			x
	Danainae	*Lycorea halia pales* C. Felder & R. Felder, 1862			x
	Danainae	*Episcada hymenaea hymenaea* (Prittwitz, 1865)		x	x
	Danainae	*Mechanitis lysimnia elisa* (Guérin-Méneville, [1844])		x	x
	Danainae	*Pteronymia ozia tanampaya* Haensch, 1909		x	x
	Danainae	*Thyridia psidii ino* C. Felder & R. Felder, 1862			x
	Heliconiinae	*Actinote pellenea pellenea* Hübner, [1821]			x
	Heliconiinae	*Euptoieta hortensia* (Blanchard, 1852)		x	x
	Heliconiinae	*Dione vanillae* (Linnaeus, 1758)			x
	Heliconiinae	*Dione moneta moneta* Hübner, [1825]		x	x
	Heliconiinae	*Dryas iulia* (Cramer, 1779)			x
	Heliconiinae	*Heliconius erato phyllis* (Fabricius, 1775)			x
	Libytheinae	*Libytheana carinenta carinenta* (Cramer, 1777)		x	x

Family	Subfamily	Species			
	Limenitidinae	*Adelpha alala negra* (C. Felder & R. Felder, 1862)			x
	Limenitidinae	*Adelpha thessalia cesilas* Fruhstorfer, 1915		x	x
	Nymphalinae	*Junonia genoveva hilaris* C. Felder & R. Felder, 1867		x	x
	Nymphalinae	*Chlosyne lacinia saundersi* (E. Doubleday, [1847])			x
	Nymphalinae	*Dagon catula* (Hopffer, 1874)			x
	Nymphalinae	*Ortilia gentina* Higgins, 1981		x	x
	Nymphalinae	*Ithra ithra* (W. F. Kirby, 1900)		x	x
	Nymphalinae	*Tegosa claudina* (Eschscholtz, 1821)	x	x	x
	Nymphalinae	*Telenassa berenice* (Röber, 1913)			x
	Nymphalinae	*Hypanartia bella* (Fabricius, 1793)		x	x
	Nymphalinae	*Hypanartia lethe lethe* (Fabricius, 1793)		x	x
	Nymphalinae	*Smyrna blomfildia blomfildia* (Fabricius, 1781)			x
	Nymphalinae	*Vanessa braziliensis* (Moore, 1883)	x	x	x
	Nymphalinae	*Vanessa carye* (Hübner, [1812])	x		x
	Nymphalinae	*Anartia jatrophae jatrophae* (Linnaeus, 1763)		x	x
	Satyrinae	*Corades iduna procellaria* Thieme, 1907		x	x
	Satyrinae	*Hermeuptychia sp.* Forster, 1964	x	x	x
	Satyrinae	*Lasiophila orbifera pura* Thieme, 1907			x
	Satyrinae	*Cissia phronius* (Godart, [1824])		x	x
	Satyrinae	*Paryphthimoides poltys poltys* (Prittwitz, 1865)	x	x	x
	Satyrinae	*Pedaliodes palaepolis* (Hewitson, 1878)	x		x
	Satyrinae	*Pharneuptychia phares* (Godart, [1824])	x		x
	Satyrinae	*Physcopedaliodes physcoa* (Hewitson, 1862)	x		
	Satyrinae	*Taguaiba ypthima Hübner, [1821]*	x		
	Satyrinae	*Yphthimoides celmis* (Godart, [1824])	x	x	x
Papilionidae	Papilioninae	*Heraclides lamarchei* (Staudinger, 1892)			x
	Papilioninae	*Battus polydamas polydamas* (Linnaeus, 1758)		x	x
Pieridae	Coliadinae	*Anteos clorinde* (Godart, [1824])			x
	Coliadinae	*Abaeis albula* (Cramer, 1775)	x	x	x
	Coliadinae	*Abaeis arbela arbela* Geyer, 1832		x	x
	Coliadinae	*Abaeis salome salome* (C. Felder & R. Felder, 1861)		x	
	Coliadinae	*Teriocolias deva deva* (E. Doubleday, 1847)		x	x
	Coliadinae	*Phoebis neocypris neocypris* (Hübner, [1823])	x	x	x
	Coliadinae	*Phoebis sennae marcelina* (Cramer, 1777)		x	x
	Coliadinae	*Pyrisitia nise flóscula* (Weeks, 1901)		x	x
	Coliadinae	*Zerene cesonia cesonides* (Staudinger, 1894)		x	x
	Dismorphiinae	*Pseudopieris nehemia prasina* Hayward, 1949			x
Riodinidae	Riodininae	*Barbicornis basilis tucumana* Thieme, 1907			x
	Riodininae	*sp1*			x
	Riodininae	*Riodina lysippoides* Berg, 1882			x

1. **Figure 1.** Study area and surveyed patches location at Finca "Las Capillas", (24° 05⬜ 34'' S; 65° 09⬜ 32'' W, 1160 m asl), Jujuy, NW Argentina. **FO=** Secondary native forest; **EU**= *Eucalyptus* plantation;GR= Secondary grassland.

2. **Figure 2.** Diversity profiles based on Hill numbers (^{q}D) for diurnal butterfly assemblages present in three surveyed habitat patches at Finca "Las Capillas", Jujuy, NW Argentina: **EU**: *Eucalyptus* plantation, **FO**: secondary native forest, and **GR**: Secondary grassland. The x axis represents the order *q*. Colored ribbons represent the 95% Confidence intervals.

3. **Figure 3.** Beta diversity partition into nestedness and species turnover components among the three surveyed habitat patches at Finca "Las Capillas", Jujuy, NW Argentina: **FO**: secondary native forest, **EU**: *Eucalyptus* plantation, and **GR**: Secondary grassland. Bars correspond to total beta diversity.

4. **Figure 4.** Species rank abundance curves based on the log-transformed relative abundances of butterfly species at each surveyed habitat patch at Finca "Las Capillas", Jujuy, NW Argentina. **FO**: secondary native forest; **EU**: *Eucalyptus* plantation, and **GR**: Secondary grassland. Only dominant species were identified by a name code. *Pedaliodes palaepolis* (**Pedpala**), *Eunica tatila bellaria* (**Euntati**), *Hermeuptychia sp.* (**Hersp**), *Cybdelis petronita* (**Cybpetr**), *Biblis hyperia nectanabis* (**Bibhype**).