Longitudinal Proteomic Profiling of T Cell Differentiation In Vivo Reveals Biochemical Remodeling Underlying Exhaustion

Christian M. Beusch[1,2,3,4,10]*, Abdelhameed S. Dawood[1,2]*, Ahmet Ozdilek[1,2], Sarah Welbourn[1,2,3,4], Youssef M. Zohdy[2], Christopher M. Monaco[1,2,3,4,5], Alexandra S. Flegle[1,2,6], Ahmed Mahgoub[11], Sakshi Malik[2,7,9], Christina Niavi[2,7,8], Akil Akhtar[2,7], Carly Roman[2,7,8], Autumn A. Gavora[2,7], David E. Gordon[1,2,3,4,5] #, and Mohamed S. Abdel-Hakeem[1,2,3,6,8,12] #

[1] Pathology Advanced Translational Research Unit (PATRU), Department of Pathology and Laboratory Medicine, Emory University School of Medicine, Atlanta, GA, USA
[2] Emory Vaccine Center, Emory University School of Medicine, Atlanta, GA, USA
[3] Winship Cancer Institute, Emory University and Georgia Institute of Technology
[4] Wallace H. Coulter Department of Biomedical Engineering, Emory University and Georgia Institute of Technology
[5] Molecular and Systems Pharmacology Graduate Program, Emory University, Atlanta, GA, USA
[6] Cancer Biology Graduate Program, Emory University, Atlanta, GA, USA
[7] Department of Microbiology and Immunology, Emory University School of Medicine, Atlanta, GA
[8] Immunology and Molecular Pathogenesis Graduate Program, Emory University, Atlanta, GA, USA
[9] Currently: Department of Pediatrics, Emory University School of Medicine, Atlanta, GA, USA
[10] Department of Surgical Sciences, Uppsala University, Uppsala, Sweden
[11] Jeham AI, Alpharetta, GA, USA
[12] Department of Microbiology and Immunology, Faculty of Pharmacy, Cairo University, Cairo, Egypt

*Equal contribution, # Corresponding authors

Author contributions

MSA and DEG conceived the experiments. ASD and AO performed all immunology experiments, including LCMV infections, sorting, and flow cytometry, with help from ASF, SM, AA, and CR. SM and CN provided technical help with flow cytometry panels. ASD was responsible for mouse breeding and colony maintenance with help from AAG. CMB optimized global and phosphoproteomics workflows and performed proteomics sample preparation, phosphopeptide enrichment, mass spectrometry acquisition, and data analysis with input from DEG. CMB and SLW conducted protein abundance mass spectrometry, supported by CMM. CMB performed bioinformatic analyses of proteomics, pan-cancer single-cell RNAseq, and conservation analysis with input from DEG. CMB generated figures with input from DEG and MSA. YMZ and CMB performed bioinformatic analyses for RNAseq data with help from AM. ASD performed flow cytometry analyses and prepared the immunology figures. CMB, MSA, and DEG wrote the manuscript.

Acknowledgments

This work was supported by an Emory MP3 Seed Grant (121861), a Winship Cancer Center Invest$ Pilot Award (147150), and Department of Pathology and Laboratory Medicine Start-up funds to MSA and DEG. CMB was supported by a Swedish Research Council grant (2023-00510). ASF was supported by the National Institutes of Health through the Initiative for Maximizing Student Development (IMSD) T32 Fellowship (T32GM148391-01). The authors would like to thank members of the Hakeem and Gordon labs, as well as members of the Pathology Advanced Translational Research Unit (PATRU) and Emory Vaccine Center (EVC) for their valuable discussions and comments. We would also like to thank the Emory Flow Cytometry Core (EFCC) and the Emory Integrated Genomics Core (EIGC) for their support. This study was supported in part by the Emory Flow Cytometry Core (EFCC), one of the Emory Integrated Core Facilities (EICF), and is subsidized by the Emory University School of Medicine. Additional support was provided by the National Center for Georgia Clinical & Translational Science Alliance of the National Institutes of Health under Award Number UL1TR002378. Research reported in this publication was supported in part by the

Emory Integrated Genomics Core (EIGC) shared resource of Winship Cancer Institute of Emory
University and NIH/NCI under award number P30CA138292. We thank the NIH Tetramer Core
Facility (contract number 75N93020D00005) for providing tetramers used in this study (H2-Db
LCMV-GP33-41, KAVYNFATC). The NIH Tetramer Core Facility is supported through National
Institute of Allergy and Infectious Diseases (NIAID) and co-funding from the National Cancer
Institute (NCI). The content is solely the responsibility of the authors and does not necessarily
represent the official views of the National Institutes of Health.

Abstract

CD8 T cell exhaustion impedes immune responses to cancer and chronic infections, and a biochemical understanding of exhaustion is essential to improving immunotherapy. Here, we present the first longitudinal protein abundance and phosphoproteomic analysis of antigen-specific CD8 T cells undergoing differentiation in vivo during acute (LCMV-Armstrong) and chronic (LCMV-Clone 13) infection. Comparing protein abundance across the two infection conditions identified over 180 known and novel exhaustion-associated proteins, including proteins missed by transcriptional analyses. Phosphoproteomic analysis identified >900 differentially regulated phosphosites on >400 proteins, including known inhibitory phosphosites on PD1, PAG1, SHP-1/PTPN6, SLAMF1/CD150. We also calculated phosphosite conservation across mammals, to direct follow-up studies towards sites with likely essential function. Lastly, our analysis uncovers exhaustion-associated kinases with clinical-stage inhibitors, underscoring the translational utility of our dataset to guide immunotherapy development. Together, our datasets define a biochemical atlas of T cell exhaustion in vivo, shedding light on the molecular mechanisms of T cell dysfunction.

Introduction

CD8 T cells are central in combating tumors and intracellular pathogens. During acute infection, naïve T cells are activated, giving rise to effector cells (T_{EFF}) which differentiate into memory T cells (T_{MEM}) upon clearing the pathogen. This is accompanied by extensive changes in the phenotype and cellular profiles of T cells[1, 2, 3]. However, under chronic infections and cancer, antigen persistence leads to alternative T cell differentiation through the "exhaustion" pathway, resulting in exhausted T cells (T_{EX}) that progressively lose function and exhibit distinct phenotypic, transcriptional, and epigenetic profiles compared to T_{EFF}/T_{MEM} cells [4, 5, 6, 7, 8, 9]. The biochemical landscape of antigen-specific T cell differentiation *ex vivo* is understudied, limiting our ability to therapeutically modulate T_{EX} cells.

While transcriptomic and epigenetic approaches offer valuable insights into gene expression and regulation, they do not directly measure the functional molecules driving cellular processes – the proteome. Several studies have reported discrepancies between RNA and protein levels, highlighting the importance of direct protein measurements[10, 11, 12]. More importantly, analysis of post-translational modifications (PTMs) of proteins, e.g., phosphorylation, offers biochemical insights that are not accessible to transcriptional readouts. Notably, the central importance of phosphorylation for T cell activation, and its antagonism by inhibitory signaling pathways, are central to the regulation of T cell response to antigen stimulus[13, 14, 15]. Therefore, analyses of protein abundance and phosphorylation in the context of T cell differentiation afford an enhanced perspective on the mechanisms underlying T cell differentiation and exhaustion. Previous proteomic studies of T cell differentiation have primarily been conducted *in vitro*, often utilizing non-specific stimulation and focusing on the early stages of T cell activation[16, 17, 18]. Other investigations have profiled immune cells' proteome in the context of viral infections or cancer, demonstrating that proteomic analyses can uncover key proteins overlooked in transcriptomic datasets—some of which hold promise as immunotherapeutic targets[19, 20]. However, comprehensive proteomic analysis of *ex vivo* CD8 T cells in an antigen-specific model across T cell differentiation states has not been performed.

Here, we profiled the dynamics of proteomic alterations during the differentiation of antigen-specific CD8 T cells *in vivo* in the context of acute versus chronic infection with lymphocytic choriomeningitis virus (LCMV) Armstrong (Arm) and Clone-13 (CL13) strains, respectively. Overall, our comprehensive analysis of global proteomics aligns with established knowledge of T cell biology, highlighting key canonical inhibitory receptors, metabolic pathways, and transcription factors associated with the different states of T cell differentiation. Crucially, our analysis reveals inconsistencies between mRNA and protein abundance for several differentially expressed proteins, underscoring the indispensable role of proteomics in capturing the true biochemical landscape of T cell differentiation. Additionally, our phosphoproteomic analysis highlights known phosphorylation sites mediating inhibitory signaling, and also identifies a multitude of novel phosphorylation sites associated with T cell differentiation that merit further study. Using kinase substrate enrichment analysis, we identified state-specific kinases, e.g., T_{EX}-specific kinases, as targets for kinase inhibitor interventions. Notably, we applied a sequence conservation-based prioritization approach to focus on phosphorylation sites of potential functional importance and highlight several understudied conserved phosphorylation sites identified in our dataset. Given the widespread adoption and translational relevance of the LCMV mouse model, our dataset provides a powerful resource for deeper insights into T cell differentiation and serves as a valuable complement to other omics approaches.

Results

Longitudinal proteomic profiling of CD8 T cells reveals dynamic proteome modulations

For decades, the LCMV mouse model has been instrumental in the phenotypic, transcriptional, and epigenetic characterization of T cell differentiation. A central strength of this system is the ability to compare mice infected with the Clone 13 strain (LCMV-CL13), which induces chronic infection, to those infected with the Armstrong strain (LCMV-Arm), which induces acute infection. This comparison specifically reveals exhaustion-associated molecular programs in T cells during chronic infection, offering valuable insights into the underlying mechanisms of exhaustion that revealed successful immunotherapeutic approaches[1, 5, 8, 21, 22, 23].

Leveraging the power of the LCMV model, we employed liquid chromatography-tandem mass spectrometry (LC-MS/MS) to profile proteomic alterations of transgenic LCMV-GP33-specific P14 CD8 T cells at different stages of chronic and acute infection (Fig.1A). P14 cells were adoptively transferred into congenically distinct recipient mice, which were either infected with chronic LCMV-CL13 or acute LCMV-Arm, respectively generating exhausted T cells (T_{EX}) or effector/memory T cells (T_{EFF}/T_{MEM}). Mice were sacrificed, and P14 cells were isolated and sort-purified across a time course at 8, 15, and 30 days post-infection (hereafter d8, d15, and d30), representing early, intermediate, and late stages of infection[1, 5, 8, 21, 23]. Naïve P14 cells were sorted as a benchmark for protein expression in the naïve state. Our flow cytometry analyses reproduced established phenotypes of T_{EX} and T_{EFF}/T_{MEM} (Fig.1B-C, Extended Data Fig.1). Sorted P14 cells were processed for global protein abundance analysis by LC-MS/MS. We quantified over 6,600 proteins across the entire dataset with a robust correlation in reproducibility across all biological replicates, and quantified proteins within a broad dynamic expression range (Extended Data Fig.2A-E). We confirmed concordance between canonical T cell differentiation markers' expression measured by flow cytometry and mass spectrometry-based proteomics (Fig. 1D, Extended Data Fig.2F-G), including markers associated with activation and effector phenotype (KLRG1, CX3CR1, CD44, and granzyme B) as well as exhaustion-associated markers (PD1, Tox, and CD39/*Entpd1*).

Overall, differentiation of CD8 T cells during acute and chronic infection led to significant alterations in their proteome, delineated by two distinct trajectories (Fig.1E), confirming unique differentiation of T_{EX} compared to the T_{EFF}/T_{MEM} on the proteome level as previously shown phenotypically[22, 23], transcriptionally[8, 21], and epigenetically[1, 5]. Quantitatively, activation from the naïve T cell state to either T_{EFF} or early-T_{EX} (naïve vs. d8) was accompanied by changes in the abundance of ~1,000 proteins, which is significantly higher than modulations occurring at further stages of differentiation (Fig. 1F). For acute infection, significant changes continued as the cells transitioned from early T_{EFF}-dominated response (d8) towards a T_{MEM}-dominated response (d15), but to a much lesser extent from intermediate to late T_{MEM}. Crucially, in the context of chronic/CL13 infection, proteome modulation persisted throughout the early-to-intermediate exhaustion transition and the intermediate-to-late exhaustion transition (Fig.1F). Overall, global protein alterations quantified across our time course analysis of acute and chronic LCMV infections align with previously reported transcriptional and epigenetic studies[5, 8, 21].

Delineating T cell proteome alterations during acute versus chronic infection identifies exhaustion-specific proteins

T cell exhaustion is a hallmark of cancer and chronic viral infection, representing a divergent differentiation path for T_{EX} distinct from T_{EFF}/T_{MEM}[1, 6, 7, 8, 21, 23, 24, 25, 26, 27, 28, 29, 30, 31]. We harnessed the full potential of the LCMV mouse model, which enables comparison of antigen-specific CD8 T cells with the same specificity during both acute and chronic infection, to dissect the proteome dynamics accompanying differentiation of T_{EFF}/T_{MEM} and T_{EX}.

We applied unsupervised hierarchical clustering to group proteome alterations of T_{EFF}/T_{MEM} and T_{EX} T cells into distinct clusters based on their longitudinal dynamics (Fig.2A, Extended Data Fig.3A-B), followed by gene ontology (GO) enrichment analysis to functionally categorize proteins within each cluster (Extended Data Fig.3C). For example, T_{EX}-enriched modules reflected in cluster 4 included Eomes and Lag3, cluster 6 included PD1, cluster 8 included Tox, and cluster 11 included Tigit and CD244/2B4. These clusters were associated with GO terms including negative regulation

of cell activation and defense response (cluster 4 and cluster 6); inflammatory response and
negative regulation of virus replication (cluster 8); and regulation of cell killing and positive
regulation of cytokine production (cluster 11). Focusing on specific protein families highlighted
several biological processes, including downregulation of respiratory chain complex 1 and
mitochondrial translation-associated proteins during chronic infection (Fig.2B and Extended Data
Fig.3D), features of T_{EX} associated with metabolic dysfunction and impaired oxidative
phosphorylation[32, 33]. We also observe a spike in protein expression of MCM complex associated
with cell proliferation in T_{EFF} cells, which returns to baseline in T_{MEM}, but is sustained in T_{EX} (Fig.
2B, bottom panel) – this is consistent with previous literature[23].
By leveraging time-matched samples of T_{EFF}/T_{MEM} from acutely infected (LCMV-Arm) mice
and T_{EX} from chronically infected (LCMV-CL13) mice, we can directly compare the distinct
differentiation trajectories, analogous to previous transcriptomic and epigenetic studies. Indeed,
analyzing LCMV-Arm and LCMV-CL13 in parallel distinguishes the core program of T cell
differentiation at the intersection between Arm and CL13, versus context-specific programs unique
to either acute or chronic conditions at the early, intermediate, and late stages of infection,
respectively (Fig.2C-E, Supplementary Table 1, Extended Data Fig.3E-I). These comparisons
confirmed elevated expression of T_{EX} inhibitory surface proteins: PD1/*Pdcd1*, TIM3/*Havcr2*,
2B4/*Cd244*, CD39/*Entpd1*, Lag3, and TIGIT; in addition to the main transcription factor
orchestrating the exhaustion program, Tox (Fig.2D). Conversely, several proteins consistently
exhibited higher abundance in T_{EFF}/T_{MEM}, including the activation marker KLRG1, the memory-
associated markers CD127/*Il7r* and TCF1/*Tcf7* at the intermediate and late stages, and SATB1
which recruits the NuRD deacetylase complex to repress PD1/*Pdcd1* expression[34]. To identify
proteins specifically regulated during T cell exhaustion, we used trinary plot analysis comparing
naïve, exhausted (CL13 d30), and memory (Arm d30) T cells. Naïve markers like Sell (CD62L) and
Tcf7 were enriched in naïve cells, while exhaustion markers such as Pdcd1, Lag3, and Tox were
specific to exhausted T cells (Fig.2E).
While the exhaustion program in T_{EX} is best understood when contrasted with T_{EFF}/T_{MEM},
there is also value in analyzing the two trajectories on their own. Our longitudinal profiling of protein
abundance in T_{EFF}/T_{MEM} during acute viral infection revealed dynamic changes throughout
transitions from naïve, to effector, to memory (Extended Data Fig.4), and corroborates
transcriptional and epigenetic trends previously reported[2, 3, 8, 21]. Studies of *in vitro* T cell stimulation
demonstrated extensive proteome remodeling upon activation[12, 16, 18], however, our study details a
comprehensive profiling of the *in vivo* trajectory of T_{EFF}/T_{MEM}. Our analysis of the T_{EX} proteomic
trajectory during chronic viral infection also represents the first insight into the longitudinal *in-vivo*
protein abundance dynamics of T_{EX} (Extended Data Fig.5).
212
Tumor-associated transcriptomics highlight commonalities of T_{EX} biology across chronic
diseases
To validate that our proteomic results are not LCMV-context-dependent, we compared our
protein abundance results to the pan-cancer atlas single-cell transcriptomics dataset of human
tumor-infiltrating lymphocytes[43]. Across several T_{EX}-specific proteins identified in our dataset, we
quantified the proportion of cells expressing each transcript across all CD8 T cell clusters (Fig.2G
circle diameter), and the average RNA expression (Fig.2G color bar, yellow = highest). Across both
canonical markers (*) and less characterized proteins, we consistently observed higher percent
expression and greater RNA expression in T_{EX}-clusters, demonstrating that T_{EX}-specific proteins
identified by our proteomic analyses could be extended to the T cell behavior present in human
cancer. In several cases, we noted T_{EX}-upregulated proteins in our dataset were near/below the
limit of detection in the pan-cancer atlas T cell scRNAseq dataset (Fig.2G).
225
Integrating our protein abundance and transcriptomics data identified additional proteins
implicated in T cell exhaustion
Transcriptional analyses are pivotal to our understanding of T cell differentiation[1, 8, 21].
However, previous studies underscored the partial correlation between mRNA and protein

abundance upon T cell activation *in vitro*, highlighting the significance of post-transcriptional
mechanisms governing protein expression[10, 11, 12, 17, 44].

To compare transcriptome and proteome changes during T cell differentiation, we
performed RNA-seq on the same samples used for proteomics, and compared RNA and protein
abundance (Fig.3, Extended Data Fig.6, Supplementary Table 3). This analysis revealed a Pearson
correlation coefficient of 0.51 at 30 dpi and 0.32-0.33 at earlier time points (Fig.3A-C), aligning with
other published comparisons of RNA vs protein abundance in diverse contexts[10, 19, 45]. Gene
ontology analysis of genes with concordant changes in both RNA and protein levels reveals
enrichment for functions related to lymphocyte activation and regulation, and cell-cell adhesion. In
contrast, genes with discordant RNA and protein changes are enriched for functions such as cell
migration, organelle disassembly, RNA capping, and signal peptide processing (Fig.3D–F).

Generally, we observe concordance between RNA and protein abundance, notably with
canonical exhaustion markers including PD1/*Pdcd1*, Lag3, and Tox, which were initially identified
via transcriptomics (Fig.3A-C). Together, these analyses validate the use of global proteomics to
identify T_{EX}-associated markers and uncover novel proteins involved in exhaustion biology,
highlighting the critical importance of integrating proteomic and transcriptomic data for a more
comprehensive and unbiased understanding of T cell exhaustion.

Phosphoproteomic analyses reveal exhaustion-specific signaling modularity
Post-translational modifications (PTMs) such as phosphorylation regulate biochemical
interactions and signaling pathways underpinning cellular functions[13, 49, 50, 51]. Previous studies have
explored phosphoproteomic analysis of T cell activation *in vitro*[52, 53], however, comprehensive
phosphoproteomic analysis of antigen-specific T cells directly *ex vivo* has not yet been performed.
We performed phosphopeptide enrichment on the same samples as used for our global
proteomics analysis (Fig.4A), identifying over 10,000 localized phosphosites with high quantitative
overlap and reproducibility between biological replicates (Extended Data Fig.7A-B). Importantly, to
derive unbiased biochemical insights, we corrected for protein abundance differences of our
phosphoproteomics quantities (Extended Data Fig.7C). Principle component analysis shows clear
separation by infection condition and time point post-infection indicating distinct phosphorylation
states between sample groups (Fig.4B). Phosphoproteomic analysis uncovered extensive
remodeling of signaling networks during the transition from naïve T cells to either T_{EFF} or early-T_{EX}
states (Fig. 4C), and dynamic changes persisted throughout T_{EX} differentiation, whereas under
acute infection phosphoproteomic changes are reduced as cells transition to T_{MEM} throughout d15
and d30. These findings reinforce the notion that T cell exhaustion is not merely a consequence of
phenotypic, epigenetic, transcriptional, or global proteomic changes but is also shaped by distinct
post-translational modifications and their impacts on cellular biochemical networks and signaling
pathways.

Phosphoproteomics reveals kinase signatures of T cell exhaustion

Kinase-substrate enrichment analysis (KSEA) enables inference of upstream kinases
responsible for specific phosphorylation events, and thus identifies potential kinase targets for
therapeutic intervention[54, 55] (Fig.4D). Comparing phosphosite changes in the different cell states
throughout the differentiation trajectory revealed a consistent trend across cyclin-dependent
kinases (CDKs). CDK activity surged at d8 in both conditions, persisting in T_{EX} while returning to
quiescent levels in T_{MEM} (Fig.4E, Extended Data Fig.7D, Supplementary Table 4). In late-$T_{EX,}$ CDK
activity ultimately diminished, aligning with reduced proliferative capacity in late-T_{EX}[23].
Reassuringly, this temporal pattern mirrored the protein abundance trend of MCM complex proteins
(Fig. 2B, bottom panel), which are also associated with cell proliferation.

Thus, comprehensive KSEA of our phosphoproteomic data might revealed functionally
relevant kinases that orchestrate T cell regulation, pinpointing key regulatory nodes that could serve
as therapeutic targets.

281
282 **Comparative Phosphoproteomics of Exhausted and Memory T Cells Uncovers Regulatory**
283 **Mechanisms**

284
Phosphorylation at specific protein residues acts as a molecular switch, regulating protein
activity, stability, and localization – and ultimately influencing cell fate and function[52, 63]. In some
cases, specific phosphorylation events can initiate or alter entire signaling pathways[64, 65]. With this
in mind, we focused on differentially regulated phosphorylation sites between T_{EX} (CL13-d30) and
T_{MEM} (Arm-d30), as these sites could provide biochemical mechanistic and therapeutic insights
(Fig.4G, Supplementary Table 5). Among the most significantly modulated sites on T_{EX}, we
identified several located on key inhibitory receptors and transcription factors. These include PD1
Tyrosine 225 (phosphosite notion Y225) and Y248; Lag3 S476; CD244 S287 and S306; and Eomes
T520, S617, S652 and S654. Additionally, we observed dysregulated phosphorylation of proteins
involved in TCR cell signaling, such as Ptpn6 Y564, Ptpn12 S673, Lck S166, Grap2 T254, Lat T62
and Y67, and Map3k8 S141. Several of these phosphorylation sites have already been studied in
the context of T cell exhaustion. Notably, Grap2 T254 (T262 in humans) is increased in T_{EX},
consistent with its function negatively regulating T cell activation. As an adaptor protein, Grap2
plays a pivotal role in linking the TCR to downstream signaling cascades, thereby mediating T cell
activation and effector function[66, 67, 68].

300
Leveraging the longitudinal nature of our data, we examined PTM regulation on key proteins
(Fig.4H). PD1 Y225 and Y248[15, 69] – located within the ITIM and ITSM motifs, respectively –
exhibited contrasting phosphorylation patterns in T_{EX} vs T_{EFF}/T_{MEM}, with Y225 downregulated and
Y248 upregulated in T_{EX} (Fig.4G, H). This is consistent with functions of these PD1 phosphosites:
p-Y248 is known to mediate inhibitory signaling via SHP-2 and SHIP recruitment, dampening T cell
activation[70], while Y225 (Y223 in humans) is dispensable for SHP-2 binding and downstream PD1
signaling in T cells[15]. On SHP-1/Ptpn6, a negative regulator of TCR signaling, we also observed a
significant increase in Y564 phosphorylation in T_{EX}. This is consistent with published work,
indicating phosphorylation of Y536 or Y564 on SHP-1/Ptpn6 relieves auto-inhibition, allowing its
phosphatase activity to dampen signaling[71, 72, 73, 74]. We also observed upregulation of Lag3 S476,
embedded within its FSALE motif (conserved between mouse 475-479 and human 481-487), a
sequence essential for LAG3's suppressive activity. Lag3 S476 (S484 in humans) is dispensable
for Lag3 inhibitory activity, however considering its conservation (see Fig.5D) in mouse and
humans, it may have another uncharacterized regulatory function[75, 76].

To prioritize functionally relevant phosphorylation sites for potential pharmacological
interventions, we assessed sequence conservation of phosphosites quantified in our mouse
dataset across five other mammalian species (human, gorilla, macaque, rabbit, and rat), based on
the premise that conserved sites are more likely to have functional significance (Fig.5A-B). We
quantified conservation of phosphosites on a scale of 0-100, with 100 representing perfect
conservation across all six mammalian proteomes (Fig.5B, Supplementary Table 6). As an
example, we studied conservation of phosphorylation sites on Eomes, a transcription factor
associated with T cell exhaustion (Fig.5B-C). We observe a conservation score range of 71-98
across all phosphorylation sites in Eomes, and a range of 76-98 for four significantly altered sites
(T_{EX} vs T_{MEM} on d30). Among these four significant sites, T520 and S617 are upregulated, and S652
and S654 are downregulated in T_{EX} (Fig.5C). The functions of these phosphosites are unknown,
but three of the four were located within the transactivation domain which is responsible for
transcriptional activity[77], and two (S652 and S654) are located adjacent to a putative nuclear
localization signal located at residues 662-665[78], with potential relevance to the nuclear trafficking
of Eomes.

330   We also examined other phosphosites of interest with high conservation (Fig.5D,
Supplementary Tables 6 and 7). For example, we observe increased phosphorylation of Pag1 at
Y296 in T_{EX}. Phosphorylated Pag1 (also known as Cbp) regulates T cell signaling by recruiting Csk
to the membrane, where Csk attenuates TCR signaling by negatively regulating Src kinases[79, 80].
Importantly, mutational studies demonstrate that Y296 phosphorylation is crucial for Csk binding:
substitution of Pag1 Y296 with phenylalanine weakens the Csk interaction, while combined
mutation of Y296 and Y314 abolishes it entirely. This evidence links elevated Pag1 Y296
phosphorylation in T_{EX} to Csk inhibition of TCR signaling[81]. We also observed increased Slamf1 p-
Y288 (Y281 in humans) in T_{EX}. This residue is located within the ITSM motif of Slamf1/CD150, a T-
cell co-stimulatory molecule, and Slamf1 Y288 phosphorylation mediates binding to SHP-2 and
SHIP, two key negative regulators of T cell signaling[70]. Finally, we also highlight high conservation
of PD1 Y225 and Lag3 S476 (discussed earlier).

342   Our analysis identifies known and novel phosphorylation modifications in exhausted T
343   cells, suggesting potential mechanisms of immune regulation meriting further investigation.

344
345   **Discussion**
346
347   This study presents the first in vivo longitudinal proteomic and phosphoproteomic profiling
348   of antigen-specific CD8 T cell differentiation during both chronic and acute viral infections. By
349   contrasting T cells from chronic (T_{EX}) and acute (T_{EFF}/T_{MEM}) infection contexts, we identified
350   differentially abundant proteins and phosphosites that distinguish these states.
351
352   Most systems-level T cell exhaustion analyses have been performed using NGS-based
technologies. While transcriptomics captures a momentary view of RNA expression and
epigenetics sheds light on gene regulation, these techniques do not directly measure the global
proteome, which comprises the molecular machines and functional components that orchestrate
cellular functions. Reassuringly, we observe a correlation between transcriptomic and protein
abundance data, and thus many of the top exhaustion-associated hits in our protein abundance
dataset have previously been identified and validated[8, 21]. These include classical inhibitory
receptors: PD1/*Pdcd1*, Lag3, TIM3/*Havcr2*, 2B4/*Cd244*, and TIGIT as well as transcription factors
implicated in exhaustion: Tox and Eomes. In addition, our protein abundance analysis identified
several proteins elevated in T_{EX} that were not highlighted by sequencing-based analyses. This
discrepancy is explained by limited correlation between protein abundance and mRNA levels, also
observed in diverse contexts[10, 19, 45], as well as limited read depth in scRNAseq analyses, which
we've observed for some exhaustion-associated transcripts (see Fig.2G). Based on current
knowledge of these gene functions, they warrant further analysis towards their role in T cell function.

366   Our phosphorylation analysis highlighted state-specific PTMs modulating immune signaling,
representing opportunities for novel insights into T cell regulation and immunotherpeutic strategies.
While a few phosphosites in our dataset have been implicated in key T cell signaling pathways,
e.g., PD1 Y248[15, 69] and Grap2 T254[66, 67, 68], the majority of phosphosites identified from our
phosphoproteomics dataset are not implicated in T cell regulation, and some are entirely novel.
Interestingly, our data indicate that Eomes is rich in phosphosites that are state-dependent,
reflecting a previously unrecognized layer of regulation of the T cell differentiation program. Eomes
is implicated in both the memory state and the late exhaustion state, and displays differentially
regulated phosphosites that are enriched in either state, suggesting a potential involvement of these
PTMs in regulating the differential role of the same transcription factor under various conditions.
Thus, beyond known regulatory sites, our dataset highlights novel phosphosites with potential
functional relevance, providing new insights into T cell differentiation. More broadly, biochemical
properties such as post-translational modifications (PTMs) are not accessible by NGS-based

readouts, and these represent new opportunities for studying immune regulatory biochemistry and
identifying therapeutic strategies.

Moving forward, a key challenge in PTM analysis is the functional annotation of modification
sites. For example, large-scale MS-based studies have reported over 200,000 phosphorylation
sites in human proteins, yet their functional significance remains largely unknown[85]. Recent
advances in CRISPR/Cas9 have yielded efficient base editors applicable to unbiased high-
throughput mutational analyses of protein residues[86, 87], and new innovations continue to expand
the precision and efficiency of gene editing[88]. Utilizing the phosphoproteomic data in this study to
design targeted mutational libraries, future work will apply precision gene editing to identify the
phosphosites and protein surfaces mediating T cell differentiation and exhaustion. Phosphorylation
is only one of hundreds of PTMs, and further proteomic analyses will explore other biochemical
modifications mediating T cell differentiation and exhaustion.

In addition to identifying individual phosphosites, a higher-order interpretation of our
phosphoproteomics data is provided by kinase substrate enrichment analysis (KSEA), which infers
state-specific kinase activity based on phosphorylation sites. We validated our approach by
analyzing the activity of CDKs associated with cell proliferation, which reflected expected activity in
both acute and chronic settings[23]. Additionally, KSEA highlighted several exhaustion-associated
kinases, several of which have been functionally validated in previous work. Crucially, there are
over 80 FDA-approved kinase inhibitors and approximately 180 orally effective kinase inhibitors in
clinical trials[89], providing opportunities to leverage our data towards clinical translation.
399
Finally, our analyses of the evolutionary conservation of phosphosite motifs across
mammalian species further reinforce the value of our dataset. Mouse models, especially LCMV,
have been invaluable in dissecting the T_{EX} biology that was translated into human settings. Since
the inception of T cell exhaustion studies, a plethora of work has confirmed the conserved T_{EX} core
program across species (mice/macaques/humans) and variable chronic settings[90, 91, 92, 93, 94]. Given
the conservation of T cell regulatory mechanisms across vertebrates, our analysis highlights high
priority phosphosites to guide the field's focus.
407
A key strength of the LCMV system is the differential analysis of chronic versus acute
infection, permitting isolation of specific molecular features differentiating T_{EX} from T_{EFF}/T_{MEM}. For
brevity, we chose to focus largely on this analysis to highlight the most impactful and translatable
features in our dataset. Isolated trajectories of naïve to T_{EX}, or naïve to T_{EFF}/T_{MEM}, are also provided
in our extended data, and can be leveraged to identify and study novel biochemical mechanisms
underlying T cell differentiation.
414
The fields of immunology and cancer research are on a quest for novel interventions to
enhance the efficacy and durability of immunotherapy[95]. One understudied aspect of T cell biology
is profiling their biochemical dynamics. Here, our study provides the first-of-its-kind dataset
dissecting the kinetics of protein abundance and phosphorylation in antigen-specific T cells. Our
protein abundance profiling of T_{EX} revealed novel proteins with potential roles in exhaustion biology
not previously highlighted by transcriptional readouts, highlighting the importance of proteomic
analysis to identify the 'ground truth' of protein expression. Furthermore, investigation of PTMs
provides a biochemical perspective that is largely inaccessible through sequencing-based
approaches. This comprehensive biochemical data is essential for dissecting the regulatory
networks that govern T cell exhaustion programs and fate decisions, and ultimately developing
therapeutic strategies to modulate immunity and combat T cell dysfunction.
426

NOTE: In-depth validation of key findings from our proteomics-guided analysis is currently underway. This includes confirming the roles of newly identified proteins and phosphorylation sites in T cell signaling and exhaustion. These follow-up studies will be essential for translating our large-scale proteomic observations into mechanistic insights and potential therapeutic strategies.

References

1.   Scott-Browne, J.P. *et al.* Dynamic Changes in Chromatin Accessibility Occur in CD8(+) T Cells Responding to Viral Infection. *Immunity* **45**, 1327-1340 (2016).

2.   Kaech, S.M. *et al.* Selective expression of the interleukin 7 receptor identifies effector CD8 T cells that give rise to long-lived memory cells. *Nat Immunol* **4**, 1191-1198 (2003).

3.   Joshi, N.S. *et al.* Inflammation directs memory precursor and short-lived effector CD8(+) T cell fates via the graded expression of T-bet transcription factor. *Immunity* **27**, 281-295 (2007).

4.   Blackburn, S.D. *et al.* Coregulation of CD8+ T cell exhaustion by multiple inhibitory receptors during chronic viral infection. *Nat Immunol* **10**, 29-37 (2009).

5.   Ghoneim, H.E. *et al.* De Novo Epigenetic Programs Inhibit PD-1 Blockade-Mediated T Cell Rejuvenation. *Cell* **170**, 142-157 e119 (2017).

6.   McLane, L.M., Abdel-Hakeem, M.S. & Wherry, E.J. CD8 T Cell Exhaustion During Chronic Viral Infection and Cancer. *Annu Rev Immunol* **37**, 457-495 (2019).

7.   Pauken, K.E. *et al.* Epigenetic stability of exhausted T cells limits durability of reinvigoration by PD-1 blockade. *Science* **354**, 1160-1165 (2016).

8.   Wherry, E.J. *et al.* Molecular signature of CD8+ T cell exhaustion during chronic viral infection. *Immunity* **27**, 670-684 (2007).

9.   Wherry, E.J. & Kurachi, M. Molecular and cellular insights into T cell exhaustion. *Nat Rev Immunol* **15**, 486-499 (2015).

10. Wang, D. *et al.* A deep proteome and transcriptome abundance atlas of 29 healthy human tissues. *Mol Syst Biol* **15**, e8503 (2019).

11. Edfors, F. *et al.* Gene-specific correlation of RNA and protein levels in human cells and tissues. *Mol Syst Biol* **12**, 883 (2016).

12. Hukelmann, J.L. *et al.* The cytotoxic T cell proteome and its shaping by the kinase mTOR. *Nat Immunol* **17**, 104-112 (2016).

13. Locard-Paulet, M. *et al.* LymphoAtlas: a dynamic and integrated phosphoproteomic resource of TCR signaling in primary T cells reveals ITSN2 as a regulator of effector functions. *Molecular Systems Biology* **16**, e9524 (2020).

476 14. Jiang, Y. *et al.* Ligand-induced ubiquitination unleashes LAG3 immune checkpoint
477   function by hindering membrane sequestration of signaling motifs. *Cell* (2025).
478
479 15. Patsoukis, N. *et al.* Interaction of SHP-2 SH2 domains with PD-1 ITSM induces PD-
480   1 dimerization and SHP-2 activation. *Commun Biol* **3**, 128 (2020).
481
482 16. Rieckmann, J.C. *et al.* Social network architecture of human immune cells unveiled
483   by quantitative proteomics. *Nat Immunol* **18**, 583-593 (2017).
484
485 17. Wolf, T. *et al.* Dynamics in protein translation sustaining T cell preparedness. *Nat*
486   *Immunol* **21**, 927-937 (2020).
487
488 18. Joshi, R.N. *et al.* TcellSubC: An Atlas of the Subcellular Proteome of Human T Cells.
489   *Front Immunol* **10**, 2708 (2019).
490
491 19. Canale, F.P. *et al.* Proteomics of immune cells from liver tumors reveals
492   immunotherapy targets. *Cell Genom* **3**, 100331 (2023).
493
494 20. Jiang, Y. *et al.* Proteomics identifies new therapeutic targets of early-stage
495   hepatocellular carcinoma. *Nature* **567**, 257-261 (2019).
496
497 21. Doering, T.A. *et al.* Network analysis reveals centrally connected genes and
498   pathways involved in CD8+ T cell exhaustion versus memory. *Immunity* **37**, 1130-
499   1144 (2012).
500
501 22. Fuller, M.J. & Zajac, A.J. Ablation of CD8 and CD4 T cell responses by high viral
502   loads. *J Immunol* **170**, 477-486 (2003).
503
504 23. Wherry, E.J., Blattman, J.N., Murali-Krishna, K., van der Most, R. & Ahmed, R. Viral
505   persistence alters CD8 T-cell immunodominance and tissue distribution and results
506   in distinct stages of functional impairment. *J Virol* **77**, 4911-4927 (2003).
507
508 24. Sen, D.R. *et al.* The epigenetic landscape of T cell exhaustion. *Science* **354**, 1165-
509   1169 (2016).
510
511 25. Bengsch, B. *et al.* Bioenergetic Insufficiencies Due to Metabolic Alterations
512   Regulated by the Inhibitory Receptor PD-1 Are an Early Driver of CD8(+) T Cell
513   Exhaustion. *Immunity* **45**, 358-373 (2016).
514
515 26. Paley, M.A. *et al.* Progenitor and terminal subsets of CD8+ T cells cooperate to
516   contain chronic viral infection. *Science* **338**, 1220-1225 (2012).
517
518 27. Shin, H. *et al.* A role for the transcriptional repressor Blimp-1 in CD8(+) T cell
519   exhaustion during chronic viral infection. *Immunity* **31**, 309-320 (2009).
520
521 28. Alfei, F. *et al.* TOX reinforces the phenotype and longevity of exhausted T cells in
522   chronic viral infection. *Nature* **571**, 265-269 (2019).
523

524   29.   Khan, O. *et al.* TOX transcriptionally and epigenetically programs CD8(+) T cell
525   exhaustion. *Nature* **571**, 211-218 (2019).

527   30.   Scott, A.C. *et al.* TOX is a critical regulator of tumour-specific T cell differentiation.
528   *Nature* **571**, 270-274 (2019).

530   31.   Yao, C. *et al.* Single-cell RNA-seq reveals TOX as a key regulator of CD8(+) T cell
531   persistence in chronic infection. *Nat Immunol* **20**, 890-901 (2019).

533   32.   Wu, H. *et al.* Mitochondrial dysfunction promotes the transition of precursor to
534   terminally   exhausted   T   cells   through   HIF-1alpha-mediated   glycolytic
535   reprogramming. *Nat Commun* **14**, 6858 (2023).

537   33.   Vardhana, S.A. *et al.* Impaired mitochondrial oxidative phosphorylation limits the
538   self-renewal of T cells exposed to persistent antigen. *Nat Immunol* **21**, 1022-1033
539   (2020).

541   34.   Stephen, T.L. *et al.* SATB1 Expression Governs Epigenetic Repression of PD-1 in
542   Tumor-Reactive T Cells. *Immunity* **46**, 51-64 (2017).

544   35.   Chang, G.W. *et al.* The Adhesion G Protein-Coupled Receptor GPR56/ADGRG1 Is
545   an Inhibitory Receptor on Human NK Cells. *Cell Rep* **15**, 1757-1770 (2016).

547   36.   Palacios, D. *et al.* The G Protein-Coupled Receptor GPR56 Is an Inhibitory
548   Checkpoint for NK Cell Migration. *J Immunol* **213**, 1349-1357 (2024).

550   37.   Bilemjian, V. *et al.* The Novel Immune Checkpoint GPR56 Is Expressed on Tumor-
551   Infiltrating Lymphocytes and Selectively Upregulated upon TCR Signaling. *Cancers*
552   *(Basel)* **14** (2022).

554   38.   Carpino, N. *et al.* Regulation of ZAP-70 activation and TCR signaling by two related
555   proteins, Sts-1 and Sts-2. *Immunity* **20**, 37-46 (2004).

557   39.   Gorelik, A., Randriamihaja, A., Illes, K. & Nagar, B. A key tyrosine substitution
558   restricts nucleotide hydrolysis by the ectoenzyme NPP5. *FEBS J* **284**, 3718-3726
559   (2017).

561   40.   Horenstein, A.L. *et al.* A CD38/CD203a/CD73 ectoenzymatic pathway independent
562   of   CD39   drives   a   novel   adenosinergic   loop   in   human   T   lymphocytes.
563   *Oncoimmunology* **2**, e26246 (2013).

565   41.   Ruiz-Fernandez de Cordoba, B., Martinez-Monge, R. & Lecanda, F. ENPP1
566   Immunobiology as a Therapeutic Target. *Clin Cancer Res* **29**, 2184-2193 (2023).

568   42.   Linnemann, C. *et al.* Adenosine regulates CD8 T-cell priming by inhibition of
569   membrane-proximal T-cell receptor signalling. *Immunology* **128**, e728-737 (2009).

571   43.   Chu, Y. *et al.* Pan-cancer T cell atlas links a cellular stress response state to
572   immunotherapy resistance. *Nature Medicine* **29**, 1550-1562 (2023).

44. Nicolet, B.P. & Wolkers, M.C. The relationship of mRNA with protein expression in CD8+ T cells associates with gene class and gene characteristics. *PLoS One* **17**, e0276294 (2022).

45. Qie, J. *et al.* Integrated proteomic and transcriptomic landscape of macrophages in mouse tissues. *Nat Commun* **13**, 7389 (2022).

46. Wan, J. *et al.* De novo NAD(+) synthesis contributes to CD8(+) T cell metabolic fitness and antitumor function. *Cell Rep* **42**, 113518 (2023).

47. Wu, D. & Zhu, Y. Role of kynurenine in promoting the generation of exhausted CD8(+) T cells in colorectal cancer. *Am J Transl Res* **13**, 1535-1547 (2021).

48. Ando, S. *et al.* mTOR regulates T cell exhaustion and PD-1-targeted immunotherapy response during chronic viral infection. *J Clin Invest* **133** (2023).

49. Tan, H. *et al.* Integrative Proteomics and Phosphoproteomics Profiling Reveals Dynamic Signaling Networks and Bioenergetics Pathways Underlying T Cell Activation. *Immunity* **46**, 488-503 (2017).

50. Okazaki, T., Maeda, A., Nishimura, H., Kurosaki, T. & Honjo, T. PD-1 immunoreceptor inhibits B cell receptor-mediated signaling by recruiting src homology 2-domain-containing tyrosine phosphatase 2 to phosphotyrosine. *Proc Natl Acad Sci U S A* **98**, 13866-13871 (2001).

51. Wang, H. *et al.* Tonic ubiquitylation controls T-cell receptor:CD3 complex expression during T-cell development. *EMBO J* **29**, 1285-1298 (2010).

52. Ross, S.H. *et al.* Phosphoproteomic Analyses of Interleukin 2 Signaling Reveal Integrated JAK Kinase-Dependent and -Independent Networks in CD8(+) T Cells. *Immunity* **45**, 685-700 (2016).

53. Navarro, M.N., Goebel, J., Hukelmann, J.L. & Cantrell, D.A. Quantitative phosphoproteomics of cytotoxic T cells to reveal protein kinase d 2 regulated networks. *Mol Cell Proteomics* **13**, 3544-3557 (2014).

54. Johnson, J.L. *et al.* An atlas of substrate specificities for the human serine/threonine kinome. *Nature 2023 613:7945* **613**, 759-766 (2023).

55. Yaron-Barir, T.M. *et al.* The intrinsic substrate specificity of the human tyrosine kinome. *Nature* **629**, 1174-1181 (2024).

56. Huang, T. *et al.* MST4 Phosphorylation of ATG4B Regulates Autophagic Activity, Tumorigenicity, and Radioresistance in Glioblastoma. *Cancer Cell* **32**, 840-855 e848 (2017).

57. Xu, X. *et al.* Autophagy is essential for effector CD8(+) T cell survival and memory formation. *Nat Immunol* **15**, 1152-1161 (2014).

58. Breuer, M.L., Cuypers, H.T. & Berns, A. Evidence for the involvement of pim-2, a new common proviral insertion site, in progression of lymphomas. *EMBO J* **8**, 743-748 (1989).

59. Allen, J.D., Verhoeven, E., Domen, J., van der Valk, M. & Berns, A. Pim-2 transgene induces lymphoid tumors, exhibiting potent synergy with c-myc. *Oncogene* **15**, 1133-1141 (1997).

60. Daenthanasanmak, A. *et al.* PIM-2 protein kinase negatively regulates T cell responses in transplantation and tumor immunity. *J Clin Invest* **128**, 2787-2801 (2018).

61. Sharma, A. *et al.* PIM kinase inhibitors: an updated patent review (2016-present). *Expert Opin Ther Pat* **34**, 365-382 (2024).

62. Pettersen, I.K.N. *et al.* Upregulated PDK4 expression is a sensitive marker of increased fatty acid oxidation. *Mitochondrion* **49**, 97-110 (2019).

63. Ubersax, J.A. & Ferrell, J.E., Jr. Mechanisms of specificity in protein phosphorylation. *Nat Rev Mol Cell Biol* **8**, 530-541 (2007).

64. Tong, J., Taylor, P., Peterman, S.M., Prakash, A. & Moran, M.F. Epidermal growth factor receptor phosphorylation sites Ser991 and Tyr998 are implicated in the regulation of receptor endocytosis and phosphorylations at Ser1039 and Thr1041. *Mol Cell Proteomics* **8**, 2131-2144 (2009).

65. Zhuang, C. *et al.* A novel gain-of-function phosphorylation site modulates PTPN22 inhibition of TCR signaling. *J Biol Chem* **300**, 107393 (2024).

66. Yoder, J. *et al.* Requirement for the SLP-76 adaptor GADS in T cell development. *Science* **291**, 1987-1991 (2001).

67. Lugassy, J. *et al.* Modulation of TCR responsiveness by the Grb2-family adaptor, Gads. *Cell Signal* **27**, 125-134 (2015).

68. Lasserre, R. *et al.* Release of serine/threonine-phosphorylated adaptors from signaling microclusters down-regulates T cell activation. *J Cell Biol* **195**, 839-853 (2011).

69. Bardhan, K. *et al.* Phosphorylation of PD-1-Y248 is a marker of PD-1-mediated inhibitory function in human T cells. *Sci Rep* **9**, 17252 (2019).

70. Shlapatska, L.M. *et al.* CD150 association with either the SH2-containing inositol phosphatase or the SH2-containing protein tyrosine phosphatase is regulated by the adaptor protein SH2D1A. *J Immunol* **166**, 5480-5487 (2001).

71. Yang, J. *et al.* Crystal structure of human protein-tyrosine phosphatase SHP-1. *J Biol Chem* **278**, 6516-6520 (2003).

72.   Wang, W. *et al.* Crystal structure of human protein tyrosine phosphatase SHP-1 in
the open conformation. *J Cell Biochem* **112**, 2062-2071 (2011).

73.   Zhang, Z., Shen, K., Lu, W. & Cole, P.A. The role of C-terminal tyrosine
phosphorylation in the regulation of SHP-1 explored via expressed protein ligation.
J Biol Chem **278**, 4668-4674 (2003).

74.   Xiao, W., Ando, T., Wang, H.Y., Kawakami, Y. & Kawakami, T. Lyn- and PLC-beta3-
dependent regulation of SHP-1 phosphorylation controls Stat5 activity and
myelomonocytic leukemia-like disease. *Blood* **116**, 6003-6013 (2010).

75.   Maeda, T.K., Sugiura, D., Okazaki, I.M., Maruhashi, T. & Okazaki, T. Atypical motifs
in the cytoplasmic region of the inhibitory immune co-receptor LAG-3 inhibit T cell
activation. *J Biol Chem* **294**, 6017-6026 (2019).

76.   Aigner-Radakovics, K. *et al.* The ligand-dependent suppression of TCR signaling by
the immune checkpoint receptor LAG3 depends on the cytoplasmic RRFSALE motif.
Sci Signal **16**, eadg2610 (2023).

77.   Intlekofer, A.M. *et al.* Effector and memory CD8+ T cell fate coupled by T-bet and
eomesodermin. *Nat Immunol* **6**, 1236-1244 (2005).

78.   Brameier, M., Krings, A. & MacCallum, R.M. NucPred--predicting nuclear
localization of proteins. *Bioinformatics* **23**, 1159-1160 (2007).

79.   Brdicka, T. *et al.* Phosphoprotein associated with glycosphingolipid-enriched
microdomains (PAG), a novel ubiquitously expressed transmembrane adaptor
protein, binds the protein tyrosine kinase csk and is involved in regulation of T cell
activation. *J Exp Med* **191**, 1591-1604 (2000).

80.   Davidson, D., Bakinowski, M., Thomas, M.L., Horejsi, V. & Veillette, A.
Phosphorylation-dependent regulation of T-cell activation by PAG/Cbp, a lipid raft-
associated transmembrane adaptor. *Mol Cell Biol* **23**, 2017-2028 (2003).

81.   Tanaka, H. *et al.* Identification of a new interaction mode between the Src homology
2 domain of C-terminal Src kinase (Csk) and Csk-binding protein/phosphoprotein
associated with glycosphingolipid microdomains. *J Biol Chem* **288**, 15240-15254
(2013).

82.   Kim, J. *et al.* Polycomb- and Methylation-Independent Roles of EZH2 as a
Transcription Activator. *Cell Rep* **25**, 2808-2820 e2804 (2018).

83.   Wei, Y. *et al.* CDK1-dependent phosphorylation of EZH2 suppresses methylation of
H3K27 and promotes osteogenic differentiation of human mesenchymal stem cells.
Nat Cell Biol **13**, 87-94 (2011).

84. Isshiki, Y. *et al.* EZH2 inhibition enhances T cell immunotherapies by inducing lymphoma immunogenicity and improving T cell function. *Cancer Cell* **43**, 49-68 e49 (2025).

85. Needham, E.J., Parker, B.L., Burykin, T., James, D.E. & Humphrey, S.J. Illuminating the dark phosphoproteome. *Sci Signal* **12** (2019).

86. Richter, M.F. *et al.* Phage-assisted evolution of an adenine base editor with improved Cas domain compatibility and activity. *Nat Biotechnol* **38**, 883-891 (2020).

87. Komor, A.C., Kim, Y.B., Packer, M.S., Zuris, J.A. & Liu, D.R. Programmable editing of a target base in genomic DNA without double-stranded DNA cleavage. *Nature* **533**, 420-424 (2016).

88. Anzalone, A.V. *et al.* Search-and-replace genome editing without double-strand breaks or donor DNA. *Nature* **576**, 149-157 (2019).

89. Roskoski, R. Properties of FDA-approved small molecule protein kinase inhibitors: A 2024 update. *Pharmacological Research* **200**, 107059 (2024).

90. Barber, D.L. *et al.* Restoring function in exhausted CD8 T cells during chronic viral infection. *Nature* **439**, 682-687 (2006).

91. Day, C.L. *et al.* PD-1 expression on HIV-specific T cells is associated with T-cell exhaustion and disease progression. *Nature* **443**, 350-354 (2006).

92. Pauken, K.E., Torchia, J.A., Chaudhri, A., Sharpe, A.H. & Freeman, G.J. Emerging concepts in PD-1 checkpoint biology. *Semin Immunol* **52**, 101480 (2021).

93. Yates, K.B. *et al.* Epigenetic scars of CD8(+) T cell exhaustion persist after cure of chronic infection in humans. *Nat Immunol* **22**, 1020-1029 (2021).

94. Abdel-Hakeem, M.S. *et al.* Epigenetic scarring of exhausted T cells hinders memory differentiation upon eliminating chronic antigenic stimulation. *Nat Immunol* **22**, 1008-1019 (2021).

95. Saez-Ibanez, A.R., Sommers, E., Upadhaya, S. & Campbell, J. PD1/PDL1 clinical trials adapt to a growing landscape of patients resistant to PDx. *Nat Rev Drug Discov* **22**, 944-945 (2023).

96. Welsh, R.M. & Seedhom, M.O. Lymphocytic choriomeningitis virus (LCMV): propagation, quantitation, and storage. *Curr Protoc Microbiol* **Chapter 15**, Unit 15A 11 (2008).

97. Meier, F. *et al.* diaPASEF: parallel accumulation-serial fragmentation combined with data-independent acquisition. *Nat Methods* **17**, 1229-1236 (2020).

98. Skowronek, P. *et al.* Rapid and In-Depth Coverage of the (Phospho-)Proteome With Deep Libraries and Optimal Window Design for dia-PASEF. *Mol Cell Proteomics* **21**, 100279 (2022).

99. Bruderer, R. *et al.* Optimization of Experimental Parameters in Data-Independent Mass Spectrometry Significantly Increases Depth and Reproducibility of Results. *Mol Cell Proteomics* **16**, 2296-2309 (2017).

100. Jin, L. *et al.* A comparative study of evaluating missing value imputation methods in label-free proteomics. *Sci Rep* **11**, 1760 (2021).

101. Lazar, C., Gatto, L., Ferro, M., Bruley, C. & Burger, T. Accounting for the Multiple Natures of Missing Values in Label-Free Quantitative Proteomics Data Sets to Compare Imputation Strategies. *J Proteome Res* **15**, 1116-1125 (2016).

102. Zhou, Y. *et al.* Metascape provides a biologist-oriented resource for the analysis of systems-level datasets. *Nat Commun* **10**, 1523 (2019).

103. Hu, C. *et al.* CellMarker 2.0: an updated database of manually curated cell markers in human/mouse and web tools based on scRNA-seq data. *Nucleic Acids Res* **51**, D870-D876 (2023).

104. Chen, E.Y. *et al.* Enrichr: interactive and collaborative HTML5 gene list enrichment analysis tool. *BMC Bioinformatics* **14**, 128 (2013).

105. Martinez-Val, A. *et al.* Spatial-proteomics reveals phospho-signaling dynamics at subcellular resolution. *Nat Commun* **12**, 7113 (2021).

106. Ewels, P.A. *et al.* The nf-core framework for community-curated bioinformatics pipelines. *Nat Biotechnol* **38**, 276-278 (2020).

107. S., A. FastQC: a quality control tool for high throughput sequence data. Available online at: http://www.bioinformatics.babraham.ac.uk/projects/fastqc. (2010).

108. Martin, M. CUTADAPT removes adapter sequences from high-throughput sequencing reads. *EMBnet.journal* **17** (2011).

109. Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21 (2013).

110. Patro, R., Duggal, G., Love, M.I., Irizarry, R.A. & Kingsford, C. Salmon provides fast and bias-aware quantification of transcript expression. *Nature Methods* **14**, 417-419 (2017).

111. Love, M.I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biology* **15**, 550 (2014).

112. Aboyoun P, G.R. pwalign: Perform pairwise sequence alignments. Available online at: https://bioconductor.org/packages/pwalign. (2025).

814
815   113.   Vizcaino, J.A. *et al.* ProteomeXchange provides globally coordinated proteomics
816   data submission and dissemination. *Nat Biotechnol* **32**, 223-226 (2014).
817
818

Figure 1

Figure 1: Longitudinal proteomic profiling of antigen-specific CD8 T cells distinguishes proteome alterations under acute and chronic infection conditions. (**A**) Overview of the experimental design to analyze proteome-wide changes in CD8 T cells differentiating during acute (Armstrong) and chronic (CL13) LCMV infections. (**B, C**) Representative flow cytometry dot plots showing expression levels of the exhaustion marker PD-1 and activation marker KLRG1 on transgenic GP33-specific P14 CD8 T cells. (**D**) Comparative analysis of protein expression per-cell (MFI) assessed by flow cytometry (top) and mass spectrometry-based proteomics (bottom). N = 4, with error bars representing mean ± standard deviation. Naïve B6 mice were used as flow cytometry staining controls. The absence of a measurement bar indicates proteins not quantified by mass spectrometry. (**E**) Principal component analysis (PCA) of proteomics samples, highlighting distinct clustering of acute versus chronic infection conditions. (**F**) Quantification of protein abundance changes across the T cell differentiation trajectory in acute (top) and chronic (bottom) LCMV infection. Dark wedges indicate upregulated proteins (log$_2$FC > 1), while grey wedges represent downregulated proteins (log$_2$FC < -1), p < 0.05, one-sided ANOVA. Curved lines denote the number of differentially expressed proteins comparing acute/Armstrong versus chronic/CL13 at each post-infection time point.

Figure 2

Figure 2: Exhaustion-specific dynamic changes of the T cell proteome in the context of chronic LCMV-CL13 infection. (**A**) Unsupervised hierarchical clustering of differentially expressed proteins across acute (Arm) and chronic (CL13) infections, highlighting protein clusters with distinct temporal expression patterns. (**B**) Distribution of protein expression changes for components of the respiratory chain complex I and minichromosome maintenance (MCM) complex across all samples, illustrating metabolic and cell cycle regulation dynamics. (**C**) Scatter plot comparing protein expression in memory and exhausted T cells at day 30 post-infection relative to naïve T cells. The Venn diagram quantifies the overlap of significantly regulated proteins between the two conditions. Sample correlation was determined using Pearson correlation, while the Jaccard index assessed the similarity of significantly regulated proteins. (**D**) Volcano plot displaying differential protein abundance in P14 CD8 T cells from Arm- versus CL13-infected mice at day 30 post-infection. (**E**) Ternary plot illustrating protein-specific expression patterns across naïve, memory (Arm day 30), and exhausted (CL13 day 30) T cells, with labeled proteins serving as reference points. (**F**) Boxplots of selected proteins from (D), depicting individual proteomics measurements along the T cell differentiation trajectory. (**G**) Bubble plot showing gene expression of selected proteins from (F) across tumor-infiltrating CD8 T lymphocyte (TIL) subsets, based on data from the Pan-Cancer T Cell Atlas[1]. Asterisks denote canonical exhaustion-associated genes. N = 4 for all proteomics samples.

Figure 3

Figure 3: Integration of T cell transcriptional and proteomic profiles reveals variability in transcript-to-protein correlation. (**A–C**) Scatter plots comparing protein expression (proteomics) to mRNA levels (bulk RNAseq) at different time points post-infection. Venn diagrams highlight the overlap of significantly regulated gene products between the two methods. Pearson correlation was used to assess overall sample correlation, while the Jaccard index was used to quantify the similarity of significantly regulated proteins. (**D**) Histogram showing the distribution of Pearson correlation coefficients between protein abundance and corresponding mRNA levels, illustrating the variability in transcript-to-protein relationships. (**E, F**) Gene Ontology (GO) enrichment analysis of the top 250 positively correlated (concordant) and negatively correlated (discordant) gene products. N = 4 for all proteomics samples, N = 3–4 for RNA sequencing.

Figure 4

Figure 4: Phosphoproteomic profiling reveals unique kinase activity and phosphorylation sites in the chronic LCMV-CL13 infection context. (**A**) Schematic representation of the phosphopeptide enrichment workflow used for phosphoproteomic data acquisition. (**B**) Principal component analysis (PCA) of phosphoproteomics samples, showing distinct clustering between acute and chronic infection conditions. (**C**) Quantification of phosphosite abundance changes across the T cell differentiation trajectory in acute and chronic viral infections. Dark wedges indicate upregulated phosphosites (log$_2$FC > 0.75), while grey wedges denote downregulated phosphosites (log$_2$FC < -0.75), p < 0.01, one-sided ANOVA. (**D**) Overview of Kinase-Substrate Enrichment Analysis (KSEA), a computational method to identify kinases potentially responsible for phosphorylating specific phosphosites. (**E**) Longitudinal kinase activity profiling of selected Cyclin-Dependent Kinases (CDKs) in acute and chronic infection. (**F**) Volcano plot depicting differentially regulated phosphosites in CD8 T cells from Arm- vs. CL13-infected mice at day 30 post-infection. (**G**) Schematic representation of PD-1/Pdcd1 (left), SHP-1/Ptpn6 (middle), and Lag3 (right), highlighting all quantified phosphorylation sites. N = 4 for all proteomics samples.

Figure 5

Figure 5: Phosphoproteomic profiling of T cells identifies regulatory exhaustion-specific modification sites conserved across mammalian species. (**A**) Comparative analysis of phosphorylation site conservation across multiple mammalian species. (**B**) Histogram quantifying phosphosite conservation scores across mammalian species compared with mouse. Sites on Eomes are highlighted in purple; asterisks (*) denote Eomes sites with significant phosphorylation difference between T_{EX} and T_{MEM} at day 30. (**C**) Schematic indicating quantified Eomes phosphorylation sites compared between T_{EX} and T_{MEM} (Arm) at day 30. (**D**) Boxplots display phosphorylation levels at individual phosphosites along the T cell differentiation trajectory. Amino acid sequence alignments show protein conservation across six mammalian species, with the phosphorylated residue centered and marked by a tick. N = 4 for all proteomics samples.

Extended Data Figure 1

Extended Data Figure 1: Flow cytometry validation of the phenotype and sorting purity of antigen-specific CD8 T cells used for proteomic profiling by mass spectrometry-based proteomics. (**A**) Representative dot plots reflecting the sorting purity (post-sorting, right panels) compared to the original frequencies pre-sorting. (**B**) Gating strategy for flow cytometry analysis. Representative dot plots of the phenotypic analysis of for the markers KLRG1, CD127, and PD-1 for naïve CD8 T cells (**C**), and for the markers KLRG1 and CD127 for P14 cells longitudinally during LCMV-Arm and LCMV-CL13 infection (**D**).

Extended Data Figure 2

Extended Data Figure 2: Assessment of protein abundance profiling of antigen-specific CD8 T cells in the context of acute and chronic infection. (**A**) Number of quantified proteins in each proteomics sample. (**B**) Correlation between the different samples from all four biological replicates. (**C-E**) Dynamic range for the protein abundance quantification for the naive (**C**), acute (**D**), and chronic (**E**) samples. (**F**) Comparison of protein expression assessed through flow cytometry (top) and proteomics (bottom). The absence of a measurement bar indicates no quantification. (**G**) Pearson correlation between protein expression levels assessed through flow cytometry (Y-axis) and proteomics (X-axis). Naive B6 mice were used as staining controls in (**F-G**).

Extended Data Figure 3

Extended Data Figure 3: Protein alterations in the context of T cell exhaustion compared to differentiation in an acute context. (**A**) Schematic representation of analyzed samples used in an unsupervised clustering approach to identify protein expression patterns specific to exhausted T cells. (**B**) Longitudinal protein expression dynamics across all clusters from Figure 3A, with bold lines indicating the average protein trajectory over time. (**C**) Gene Ontology (GO) enrichment analysis of protein clusters identified in Figure 3A, highlighting functional pathways associated with exhaustion. (**D**) Distribution of protein expression changes for mitochondrial translation components across all samples, illustrating metabolic and cell cycle regulatory shifts. (**E, F**) Scatter plots comparing protein expression changes in acute versus chronically infected T cells relative to naïve cells at days 8 and 15 post-infection. The Venn diagram quantifies the overlap of significantly regulated proteins between the two conditions. Pearson correlation was used to assess sample similarity, while the Jaccard index evaluated the overlap of significantly regulated proteins. (**G**) Signatures of differentially regulated proteins from Figure 3D, identified using the CellMarker database[2]. (**H, I**) Volcano plots illustrating differential protein abundance in CD8$^+$ T cells from Armstrong- versus Clone 13-infected mice at days 8 and 15 post-infection. P-values were calculated using a two-sided Student's t-test with equal variance. N = 4 for all proteomics samples.

Extended Data Figure 4

Extended Data Figure 4: Expression profiles of proteins defining T cell differentiation during acute LCMV-Arm infection. (**A**) Unsupervised hierarchical clustering of differentially expressed proteins throughout the LCMV Armstrong (acute) infection time course. (**B**) Longitudinal expression dynamics of proteins within selected clusters. The bold blue line represents the average protein change. (**C**) Gene Ontology (GO) enrichment analysis of selected protein clusters, highlighting biological processes associated with different phases of T cell differentiation. (**D-F**) Volcano plot depicting differentially expressed proteins in naïve CD8+ T cells compared to antigen-experienced effector CD8+ T cells at day 8 post-infection, d8 (left), d15 (middle, and d30 (right). P-values were calculated using a two-sided Student's t-test with equal variance. (**G**) Longitudinal expression trends of selected proteins across the differentiation trajectory of CD8+ T cells, tracking key regulators and markers from naïve to effector and memory states. (**H**) Explanatory ternary plot. (**I**) Ternary plot depicting protein-specific expression patterns across the transition trajectory from naïve, to effector (Arm d8), then memory (Arm d30) T cells, illustrating key regulators of T cell differentiation. Named proteins act as reference points. N = 4 for all proteomics samples

Extended Data Figure 5

Extended Data Figure 5: Expression profiles of proteins defining T cell exhaustion. (**A**) Unsupervised hierarchical clustering of differentially expressed proteins throughout the LCMV-CL13 (chronic) infection time course. (**B-D**) Volcano plots depicting differentially expressed proteins in naïve versus early-Tex d8 (left), d8 versus d15 (middle), and d15 versus d30 (right). P-values were calculated using a two-sided Student's t-test with equal variance. N = 4 for all proteomics samples.

Extended Data Figure 6

Extended Data Figure 6: Transcriptomic analysis and correlation of changes in transcript abundance compared with protein expression. (**A**) Principal component analysis (PCA) of RNA sequencing samples. (**B**) Differentially expressed genes in the different samples, reflecting previously reported state-specific modules[3, 4]. Each column represents a biological replicate. (**C**) Pearson correlation analysis comparing protein expression levels measured by flow cytometry (Y-axis) with RNA sequencing counts (X-axis). Naïve B6 mice were used as staining controls. N = 3-4 for RNA sequencing.

Extended Data Figure 7

Extended Data Figure 7: Phosphoproteomic analysis and kinase activity in T cell differentiation. (**A**) Quantification of phosphosites in each sample, showing the number of identified sites per condition. (**B**) Overlap of phosphosite quantification across all samples, highlighting shared and unique phosphosites. (**C**) Correlation analysis between protein expression measurements and phosphosite changes, both before and after adjusting for intrinsic protein expression changes during T cell differentiation. Sample correlation was determined using Pearson correlation, with only one replicate shown per sample. (**D**) Kinase-substrate enrichment analysis (KSEA) for all cyclin-dependent kinases (CDKs), comparing each time point to naïve T cells.

References

1. Chu, Y. *et al.* Pan-cancer T cell atlas links a cellular stress response state to immunotherapy resistance. *Nature Medicine* **29**, 1550-1562 (2023).

2. Hu, C. *et al.* CellMarker 2.0: an updated database of manually curated cell markers in human/mouse and web tools based on scRNA-seq data. *Nucleic Acids Res* **51**, D870-D876 (2023).

3. Wherry, E.J. *et al.* Molecular signature of CD8+ T cell exhaustion during chronic viral infection. *Immunity* **27**, 670-684 (2007).

4. Doering, T.A. *et al.* Network analysis reveals centrally connected genes and pathways involved in CD8+ T cell exhaustion versus memory. *Immunity* **37**, 1130-1144 (2012).