Title: A new avian feeding mechanism: Nectar suction by sunbirds

Authors: David Cuban[1,2], Cynthia Wang-Claypool[3,4], Yohanna Dalimunthe[5], Colleen T. Downs[6],

Rauri Bowie[3,4], Fabian Brau[7] Steve Johnson[6], Alejandro Rico-Guevara[1,2]

Affiliations:

[1]*Department of Biology, University of Washington, Life Sciences Building, Box 351800, Seattle,*

WA 98105, USA

[2]*The Burke Museum of Natural History and Culture, Ornithology Department, 4300 15th Avenue*

NE, Seattle, WA 98105, USA

[3]*Department of Integrative Biology, University of California, Berkeley 3040 Valley Life Sciences,*

Building # 3140, Berkeley, CA 94720-3200

[4]*The Museum of Vertebrate Zoology, University of California, Berkeley, 3101 Valley Life*

Sciences Bldg | Berkeley, CA 94720-3160

[5]*Museum Zoologicum Bogoriense, Research Center for Biosystematics and Evolution, National*

Research and Innovation Agency of Indonesia (BRIN) | Jalan Raya Jakarta-Bogor KM 46,

Cibinong, West Java, Indonesia 16911

[6]*Centre for Functional Biodiversity, School of Life Sciences, University of KwaZulu-Natal,*

Private Bag X01, Scottsville, Pietermaritzburg, 3209, South Africa

[7] *Nonlinear Physical Chemistry Unit, Université Libre de Bruxelles, Av. Franklin Roosevelt*

50, 1050 Bruxelles, Belgium

* Corresponding author: David Cuban

Email: dcuban@uw.edu; ORCID: https://orcid.org/0000-0003-4661-7180

Other emails and ORCIDs of others:

Cynthia Wang-Claypool Email: cynthia.yh.wang@berkeley.edu; ORCID: https://orcid.org/0000-
0002-7725-5787

Yohanna Dalimunthe Email: yoha008@brin.go.id; ORCID: https://orcid.org/0000-0001-6051-
3199

Colleen T. Downs Email: downs@ukzn.ac.za; ORCID: http://orcid.org/0000-0001-8334-1510

Rauri Bowie Email: bowie@berkeley.edu; ORCID: https://orcid.org/0000-0001-8328-6021

Fabian Brau Email: Fabian.Brau@ulb.be; ORCID: https://orcid.org/0000-0002-8715-5464

Steve Johnson Email: JohnsonSD@ukzn.ac.za; ORCID:https://orcid.org/0000-0002-5114-5862

Alejandro Rico-Guevara Email: colibri@uw.edu; ORCID: https://orcid.org/0000-0003-4067-5312

Abstract

Nectarivory has independently evolved about 30 times within birds, yet little is known about the

nectar-feeding mechanisms used by these very different types of birds. Multiple groups have

relatively long bills and tube-like tongues hypothesized to be adaptations for nectar extraction.

Sunbirds are the group that exhibits the largest bill and tongue convergence with the well-

studied hummingbirds. Using microCT scans we characterize the details of the sunbird feeding

apparatus and using high-speed video experiments we describe their nectar-feeding

mechanism. Sunbirds collect nectar through their bi-cylindrical tongues reciprocating them

around 9 times per second. Multiple lines of kinematic evidence and further fluid dynamics

modeling indicate that sunbirds use a unique drinking mechanism not found in any other bird:

suction feeding through the tongue. These findings are consistent across a wide sampling of

sunbird species from the geographical extremes of their range, implying that these mechanics

are shared by the entire Nectariniidae family. Our findings describe a heretofore undescribed

feeding mechanism in vertebrates, suction through the inside of the tongue, and sets the stage

for investigating the physical limitations imposed by flowers and fluid dynamical effects of

different nectar types in the flowers that sunbirds visit in nature.

Introduction

Suction is one of the most common feeding mechanisms for vertebrates that capture

prey underwater (Wainwright et al., 2015). Although most terrestrial vertebrates use lapping or

licking to drink, a few groups can suck from pools of liquid by creating a small oral opening

connected to an expandable oral cavity (e.g., some mammals; Williams, 2019). The only

vertebrate that has been reported to exhibit lingual pumping is a species of bat with lateral

tongue grooves, where suction is not possible because these groove edges do not overlap

resulting in a lack of seal (Tschapka et al., 2015). For vertebrates without lips or cheeks that

allow suction, drinking or surface filter-feeding is a challenge. In birds, suction feeding is

restricted to a few groups (e.g., filter feeders) and it is always accomplished by the tongue

pushing liquid inside the oral cavity (Rico-Guevara et al., 2019). There are no known examples

of vertebrates able to move liquids using active suction along the entire length of the tongue

cavities, as in some nectar-feeding insects. Nectarivorous vertebrates would then be prime

candidates for such a mechanism to evolve.

Nectarivory allows animals to consume high-caloric and easily digestible meals to fuel

their energy-thirsty metabolisms. In return, these nectar-feeders provide pollination services to

flowering plants that offer nectar as a reward. Nectar-feeding vertebrates exhibit a variety of

sizes, ecologies, and evolutionary histories (Fleming and Muchhala, 2005). Within birds,

specialized nectar feeding has evolved independently in approximately 30 clades (Pauw, 2019;

Hewes et al. 2022). These groups have separately evolved to fill this specialized ecological

niche and, consequently, exhibit behavioral, functional, and morphological convergence (Stiles,

1981; Fleischer et al., 2008; Janeček et al., 2015; Cuban et al., 2022; Hewes et al. 2022). Some

of the most notorious convergent traits evolved in their feeding apparati, such as bill and tongue

similarities among taxa in distant geographic regions, for example hummingbirds (Trochilidae) in

the Americas, sunbirds (Nectariniidae) in Africa and Asia, honeyeaters (Meliphagidae) in

Australia, and honeycreepers (Thraupidae) in Hawaii (Skead, 1967; Paton and Collins, 1989;

Downs, 2004; Hewes et al., 2022). Of the many nectar-feeding birds across the world, the

feeding mechanisms used to ingest nectar are best known for hummingbirds (Rico-Guevara and

Rubega, 2011; Rico-Guevara et al., 2015). Among nectarivorous birds, the taxon most

morphologically similar to hummingbirds are sunbirds.

Sunbirds and hummingbirds fill the same ecological niches in different continents,

providing pollination services to flowering plants in return for energy-packed rewards (Nicolson

and Fleming, 2014; Cheml et al., 2021; Rico-Guevara et al., 2021). These two phylogenetically

distinct lineages of birds are visually very similar, with overall small body sizes, relative bill

length and shape, and even iridescent coloration converging between them (Stiles, 1981).

Given that both sunbirds and hummingbirds have greatly elongated and narrow bills with ample

oral cavities, housing equally long and distally bi-cylindrical-hollow-tongues, we expected that

these groups would share similar feeding mechanisms, as have been described for

hummingbirds and that do not include active suction throughout the entire tongue.

The drinking mechanics have been explored across a diversity of hummingbird species;

we studied sunbirds from the geographical ends of their distribution, different parts of their

phylogeny, and with different body and bill sizes. There are 143 species of sunbirds with their

ranges spanning Africa, southern Asia, and the Malay Archipelago (Cheke and Mann, 2001;

Winkler et al., 2020). Although every species within this family includes nectar in its diet (except

Deleornis fraseri; Cheke and Mann, 2020), we know nearly nothing about how they drink nectar.

Hypotheses were made about their feeding mechanics beginning over a century ago, yet these

ideas haven't been tested (Gadow, 1883; Liversidge, 1967; Downs, 2004; Cuban et al., 2022).

Using micro computed tomography (μCT) scans of species from both groups, high-speed videos

of sunbirds drinking from transparent artificial flowers, kinematic analyses and fluid dynamics

models, we unveil the unique feeding mechanism that sunbirds use to feed on nectar.

Results

Tongue and bill morphology

Sunbird bills are long relative to their body size and generally decurved, and their

tongues have approximately the same length as the bill. The sunbird tongue can be described

along three distinct sections: (1) distally, the tongue is bifurcated, and made of medially curling,

keratinous, and very thin walls that form two hollow cylinders (Fig. S1); (2) in the middle section,

the walls are less curled inward with their ends opening up to form a single elliptical cylinder,

also hollow across its entire length; (3) in this proximal section, the tongue opens up entirely to

form a "V" shaped channel with its edges flared out laterally (Fig 1.

Sunbird and hummingbird tongues are more similar to each other than any of the

tongues found among all other nectarivores. Both species' tongues are long, thin, keratinous

structures with longitudinal grooves used to transport nectar. At the tip (Fig 1, cross-section 5),

both are bifurcated tubes with fringed ends. Although the middle section of the hummingbird

tongue has two separate grooves and sunbirds have one, both species can carry nectar in the

middle section of their tongue (Fig. 1, cross-section 4). At the proximal portion of the tongue the

two groups notably differ. Hummingbird tongues become a solid keratin structure with no

internal space to transport nectar within the tongue whereas sunbird tongues unfurl laterally at

the base, increasing the volume the tongue could carry (Fig 1, cross-sections 1 and 2). The

hummingbird tongue morphology aids in their feeding as they need a strong, rigid structure to

keep the tongue oriented properly when it pushes it between its compressed bill tips. This

allows the tongue to store elastic energy before rapidly expanding and filling its hollow portions

with nectar, known as elastic-filling. Sunbird tongue morphology consists of a continuous

groove, opening the possibility of continuous flow through the entire length of the tongue.

Sunbird and hummingbird bills also show significant convergence. Both are long and

narrow, often decurved, and have internal structures that interact with the tongue to aid in

nectar-feeding. When closed, hummingbird bills have minimal empty space outside of the

tongue grooves, with that volume decreasing towards the distal tip. Protrusions on the inner

surfaces of the upper and lower bills guide the tongue while it is protruded and help compress

the tongue at the bill tips to wring the tongue of the nectar it held from the previous lick and to

elastically load the tongue for the next expansion. Sunbird bills also have some protrusions on

the internal surfaces. However, if the bill is closed the tongue is not compressed at any point

along its length, indicating that these structures are likely for guiding the tongue only and not for

compressing it.

Tongue kinematics and nectar flow

Nectar was offered to birds in this study inside artificial flowers consisting of clear

rectangular cross-section corollas and red plastic disks to guide the birds. High-speed videos

showed that all sampled sunbirds (15 individuals from 7 species, Table S1) fed in similar ways.

First, they protrude their tongue past the slightly open bill tips until the distal portion of the

lingual cylinders go into the nectar. Second, they keep their tongue inserted, pausing

reciprocation when the tongue is at the point of maximum protrusion, while nectar flows inside

the tongue towards the mouth filling the internal tongue capacity. Third, they retract the tongue

only partially to swallow, leaving the distal cylinders partly protruded and full of nectar, but not

contacting the nectar pool. And fourth, they re-extend the tongue for the next lick cycle stage at

which the nectar in the tongue cylinders moves mouthwards (Video S1). The licking cycle period

is approximately 0.11 seconds from one tongue extension to the next with variable tongue

protrusion lengths between individual feeding bouts. An example of tongue kinematics and

nectar flow during feeding are plotted in Figure 2. To visualize the nectar flow and tongue

movement of sunbirds feeding on artificial nectar we fed birds *ad libitum* with red dyed sucrose

solutions of 20% w/w. The tongue, nectar reservoir meniscus, and nectar meniscus seen in the

tongue are measured relative to an arbitrary point at the edge of the video (Fig. 2A). As the bird

fed, it extended its tongue into the nectar reservoir and pause, while nectar could be observed

flowing inside the tongue towards the mouth before the tongue retracted partially and began the

cycle over again at a frequency of ~9 s^{-1} (Fig. 2B). By subtracting the position of the reservoir

meniscus from the tongue position we find the immersion depth to vary between 1.3 and 2.1

mm (Fig. 2C). We subtract the reservoir nectar meniscus position from the tongue nectar

meniscus position and find that the average licking period is ~ 0.11 s (Fig. 2D). Each of the licks

are overlaid with each other relative to the tongue insertion time comparing the position, h, of

the nectar meniscus in the tongue (Fig. 2E). Averaging all the tongue meniscus positions for

each lick at the same relative time in the lick cycle shows strong evidence for a linear

relationship between meniscus position and time (Fig. 2F). The first lick was not included in the

average as in each trial it differs from the subsequent licks.

Fluid dynamic modeling

Figure 3 shows the nectar meniscus rise in the tongue averaged across multiple licks of

a *Nectarinia famosa* (linear fit) in comparison with three feeding mechanics fluid models of

increasing complexity. The capillarity model has been proposed before as the mechanism used

by nectar-feeding birds. The model accounts for the surface tension of the sucrose solution at

the tongue-air-nectar interface and the viscosity of the nectar. As the model shows significant

error relative to the measured meniscus position capillarity on its own does not explain our

observations. Including the inertial effects of accelerating the fluid from a static position brings

the initial meniscus position closer to that of the observed values. However, when we include a

time-varying pressure differential in the model the predicted values almost perfectly match our

experimental data. The pressure differential itself is found to increase linearly with time, implying

that whatever internal motion the bird is doing to generate suction must be happening

continuously during the tongue's immersion period in the nectar. Derivations of the fluid models

are available in the supplementary materials.

Additional mechanistic observations

In many videos bubbles can be seen moving in the tongue when a bird does not make

complete contact between its tongue tips and the nectar reservoir (Video S2). If capillarity was

the sole mechanism used by sunbirds then no bubbles could be present as the surface tension

at the distal end of the tongue would not allow for bubbles to pass into the tongue. Moreover,

the nectar columns inside the tongue cylinders from the previous lick move mouthwards before

the tongue tips contact the nectar reservoir (see trailing menisci in Video S1, S3). When the

tongue is being extended the distal end is not in contact with a nectar reservoir, therefore there

is a surface tension force applied pulling the nectar in the tongue towards the tongue tip,

opposing any surface tension forces applied at the unseen proximal end of the tongue cavity.

Since the nectar does, in fact, move toward the mouth at this time, an additional force, must be

applied to affect this motion. Additionally, a *Chalcomitra amethystina* was recorded while its

tongue suddenly dropped down from contact with its upper bill resulting in nectar flowing out of

its mouth. We hypothesize this results from the failure of a hermetic seal between the tongue

and upper bill as the bird moved its tongue too far down while feeding (Video S4). In all feeding

events, synchronized side views evidence a downward motion of the throat while nectar is

flowing inside the tongue cylinders, as observed in the top view (e.g., Videos S1). This

downward motion of the throat corresponds to the area where the hyoid bones that support the

tongue base move through, thus consequently corresponding to the V section of the tongue

being depressed within the oral cavity. Lastly, at the end of each lick cycle a stereotypic

swallowing motion was observed in the throat when the tongue is partially retracted before re-

extension.

Discussion

For hundreds of years it was believed that capillarity was the main mechanism for nectar

feeding in birds. Using fluid dynamic models, Kim et al (2011) showed correspondence between

the optimal concentration for caloric ingestion, from previous studies, and the predicted energy

intake peak based on the feeding mechanism. In such study, three bird groups were assumed

to use capillarity, called by them capillary suction, as their drinking mechanism: hummingbirds,

honeyeaters, and sunbirds (Fig. 1, Kim et al. 2011). Since then, for hummingbirds, the capillary

filling mechanism has been disproved (Rico-Guevara and Rubega, 2011), for honeyeaters, only

one out of five species exhibited tongue kinematics consistent with capillarity (Hewes et al.,

2023), and here we tested if capillary rise alone could explain the recorded flow patterns.

Hummingbirds and most honeyeaters employ an expansive filling mechanism (Rico-Guevara et

al., 2015; Cuban et al., 2022; Hewes et al., 2023)., so our initial hypothesis was that sunbirds –

with convergent traits in their feeding apparatus morphology– would also use a similar

technique. However, our study shows that for sunbirds, the main nectar intake mechanism is

neither capillarity nor expansive filling. Instead, we found that sunbirds move nectar through

their tongues over relatively longer immersion times in the floral nectar reservoirs and do not

use of their bill tips to compress their tongues to squeeze the nectar and move it inside the bill,

which also stores the elastic energy needed for the expansive filling mechanism to work (Rico-

Guevara et al., 2015; Cuban et al., 2022; Hewes et al., 2023). In fact, sunbirds keep their bills

slightly open during the entire feeding process with multiple licks occurring before the bill and

tongue are removed from the flower corolla. The kinematic measurements of the tongue and

nectar movements demonstrate that a simple capillary filling mechanism is unable to explain the

nectar uptake patterns reported here. In the end, the only mechanistic model that matches the

collected data is the one that includes a pressure differential within the tongue cylinders, strong

evidence for the use of active suction.

Coupling the inferences from the top view videos of the flower corolla and the side view

videos of the birds bill and upper body, we propound the mechanism used by sunbirds to

generate an increasing pressure gradient when drinking nectar. Figure 4 shows a cut-away

diagram of the feeding cycle with the tongue motions that can generate the pressure differential.

The tongue lies centered inside the slightly open bill; a ridge in the roof of the oral cavity

matches the dorsal tongue shape (cross-section 1 in Fig. 4A) and would help to maintain the

tongue centered as it slides along the upper bill. When the tongue is protruded to reach the

nectar reservoir, the proximal and middle sections are pressed against the maxilla's ventral

surface (Fig. 4B). The "U" shape and flexion of the proximal tongue section allows for a hermetic

seal against the maxilla (cross-sections 2-3 in Fig. 4B), which allows for the combined action of

capillary forces and active suction that explain the intralingual menisci advance rate (Fig. 3).

The tongue is then depressed in the oral cavity resulting in the slight visible movement of the

throat. The increased volume between the tongue and maxilla creates a region of lower

pressure relative to the atmosphere which draws nectar through the tubular shaped distal and

middle sections of the tongue (cross-sections 3-4 in Fig. 4C). Finally, the tongue is pulled

proximally towards the throat and unseals from the maxilla, allowing for the bird to swallow the

nectar now in the oral cavity (Fig. 4D) before re-extending the tongue for the next lick cycle (Fig.

4E and 4F).

The mechanistically unique feeding style of sunbirds adds another interesting physical

solution to the evolutionary challenge of feeding on small quantities of liquid food concealed in

hard-to-reach spaces. The flowers that sunbirds and hummingbirds feed from often have

relatively long, narrow, and tubular corollas with nectar located at the base. However, the two

bird lineages must have had slightly different morphological traits or environmental pressures

that led to the different adaptations of their feeding apparatus to be able to extract energy

efficiently enough from the local flowers they co-evolved with. As natural selection trends

towards relatively advantageous traits even a small difference between each sunbird and

hummingbird ancestors could result in evolutionary patterns that separately selected for

different behaviors and morphologies that improved their nectar extraction rates. Further studies

of how efficient nectar-feeding birds are at drinking nectar of different sugar concentrations

would allow for deeper understanding of the plant-pollinator evolution as well as the interspecies

dependencies of birds and flowers in the world today.

In our work, we unveil a previously unknown method to generate suction through a

straw-like tongue in vertebrates without cheeks or lips. Additionally, the tongue itself is not a

closed tube and is not made of soft tissue that typically would be seen in pressure-generating

organs. This discovery raises new mechanistic questions for other vertebrate and invertebrate

groups that achieve similarly unique results with the many diverse materials, structures, and

physics phenomena of which the biological world is comprised.

Materials and Methods

Morphological analysis of bill and tongue

We used micro computed tomography (microCT) scans of the sunbird *Cyanomitra olivacea* and

the hummingbird *Calypte anna* at ~20 micron resolution to compare morphology and measured

internal tongue and oral cavity volumes. The CT scanner x-ray source voltage and current was

42 kV and 560 microAmps, respectively. We analyzed the scans using the 3D Slicer software

and the SlicerMorph package (Rolfe et al., 2021). Transverse section images were taken along

the bill and tongue at 5 points that showed relevant anatomical features of interest. We

measured the tongue structure dimensions and internal volume by adding landmark points and

segmenting out the internal volume of the tongue.

Experimental Design

We used high-speed cameras to capture the nectar, tongue, and bill movement XX individuals

from 7 sunbird species (*Nectarinia famosa* [n = 3]*, Cinnyris afer* [n = 2], *C. talatala* [n = 1], *C.*

jugularis [n = 1]*, Chalcomitra amethystina* [n = 3], *Cyanomitra olivacea* [n = 2]*, and Anthreptes*

malacensis [n = 1])(Table S1) while they fed on nectar from artificial flowers (described below).

South African sunbird species (5, Table S1) were captured in Umgungundlovu district of

KwaZulu Natal in grassland areas with aloes and foothill areas with proteas between June and

August of 2022 and 2023. Experiments with these species were conducted at the University of

KwaZulu-Natal Pietermaritzburg in the outdoor Animal House aviaries. The Indonesian sunbird

species (2, Table S1) were captured on Gunung Ambang and Gunung Klabat in North Sulawesi,

Indonesia in July 2023. Experiments with these species were conducted in field camps

established at each site. The artificial flower (Fig. S2) consisted of a red plastic ring attached to

the end of a clear rectangular prism made of thin glass (5.9 mm width x 1 mm height) oriented

horizontally that was filled with artificial nectar (20% w/w sugar concentration). When birds fed

from the artificial flower, they partially inserted their bills through the red disk into the tube and

extended their tongues to feed. Three high-speed cameras (Chronos 1.4, Krontech) were used

to film, one oriented to film the clear tube from above, and a second filming the side of the bird's

bill and head. Experiments with the Indonesian species used two cameras (Chronos 1.4,

Krontech) filming the clear tube from above and the side of the bill and head. The cameras used

Nikkor 105mm macro lenses (Nikon) and a film rate of 1000 to 4000 frames/second.

Video Analysis

We analyzed the video data using dltdv8a (Hedrick, 2008), tracking the tongue tip, bill tip,

corolla nectar meniscus in the prismatic corolla of the artificial flower, and nectar menisci inside

the tongue for multiple licks across multiple feeding bouts for each individual sunbird recorded in

the experiment. Corolla nectar meniscus is defined as the edge of the nectar in the clear

prismatic chamber at roughly the midpoint of the tube's width. The tongue nectar menisci are

defined as the leading surface of nectar as it is drawn into the tongue and the trailing surface of

the column of nectar as it moves through the tongue (these are found at different times in the

feeding process, Video S1, S2, S3). The video data was calibrated using the known width of the

nectar reservoir (5.9 mm) and the recording frame rate. The volume and volumetric rate of

nectar consumption was calculated by measuring the position change of the corolla nectar

meniscus multiplied by the nectar reservoir's cross-sectional area and the duration of time

passed between the beginning and end of the time that the tongue was in contact with the

nectar. Detailed derivations of the fluid models are available in the supplementary materials.

References:

Cheke, R. and Mann, C. (2020). Fraser's Sunbird (Deleornis fraseri), version 1.0. *Birds of*

the World.

Chmel, K., Ewome, F. L., Gómez, G. U., Klomberg, Y., Mertens, J. E. J., Tropek, R. and

Janeček, Š. (2021). Bird pollination syndrome is the plant's adaptation to ornithophily,

but nectarivorous birds are not so selective. *Oikos* **130**, 1411–1424.

Cuban, D., Hewes, A. E., Sargent, A. J., Groom, D. J. E. and Rico-Guevara, A. (2022).

On the feeding biomechanics of nectarivorous birds. *Journal of Experimental Biology*

225, jeb243096.

Eastham, L. E. S., Eassa, Y. E. E. and Wigglesworth, V. B. (1997). The feeding

mechanism of the butterfly Pieris Brassicae L. *Philosophical Transactions of the Royal*

Society of London. Series B, Biological Sciences **239**, 1–43.

Erdoğan, S. and Iwasaki, S. (2014). Function-related morphological characteristics and

specialized structures of the avian tongue. *Annals of Anatomy - Anatomischer*

Anzeiger **196**, 75–87.

Fleischer, R. C., James, H. F. and Olson, S. L. (2008). Convergent Evolution of Hawaiian

and Australo-Pacific Honeyeaters from Distant Songbird Ancestors. *Current Biology*

18, 1927–1931.

Fleming, T. H. and Muchhala, N. (2008). Nectar-feeding bird and bat niches in two worlds:

pantropical comparisons of vertebrate pollination systems. *Journal of Biogeography* **35**,

764–780.

Frost, S. K. and Frost, P. G. H. (1981). Sunbird Pollination of Strelitzia nicolai. *Oecologia*

49, 379–384.

Gill, F. B. and Wolf, L. L. (1975). Economics of Feeding Territoriality in the Golden-Winged

Sunbird. *Ecology* **56**, 333–345.

Hedrick, T. L. (2008). Software techniques for two- and three-dimensional kinematic

measurements of biological and biomimetic systems. Bioinspir. Biomim. 3, 034001.

Hewes, A. E., Cuban, D., Groom, D. J. E., Sargent, A. J., Beltrán, D. F. and Rico-

Guevara, A. (2022). Variable evidence for convergence in morphology and function

across avian nectarivores. *Journal of Morphology* **283**, 1483–1504.

Hewes, A. E., Baldwin, M. W., Buttemer, W. A. and Rico-Guevara, A. (2023). How do

honeyeaters drink nectar? *Integrative and Comparative Biology* **63**, 48–58.

Köhler, A., Leseigneur, C. D. C., Verburgt, L. and Nicolson, S. W. (2010). Dilute bird

nectars: viscosity constrains food intake by licking in a sunbird. *American Journal of*

Physiology-Regulatory, Integrative and Comparative Physiology **299**, R1068–R1074.

Krenn, H. W. (2010). Feeding Mechanisms of Adult Lepidoptera: Structure, Function, and

Evolution of the Mouthparts. *Annual Review of Entomology* **55**, 307–327.

Lotz, C. N. and Nicolson, S. W. (1999). Energy and water balance in the lesser double-

collared sunbird (Nectarinia chalybea) feeding on different nectar concentrations. *J*

Comp Physiol B **169**, 200–206.

Mbatha, K., Downs, C. T., Penning, M. and Dean, W. R. J. (2002). Nectar passage and

gut morphology in the Malachite Sunbird and the Black-capped Lory: implications for

feeding in nectarivores. *Ostrich* **73**, 138–142.

Nicolson, S. W. (2002). Pollination by passerine birds: why are the nectars so dilute?

Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology

131, 645–652.

Nicolson, S. W. and Fleming, P. A. (2003). Energy Balance in the Whitebellied Sunbird

Nectarinia talatala: Constraints on Compensatory Feeding, and Consumption of

Supplementary Water. *Functional Ecology* **17**, 3–9.

Nicolson, S. W. and Fleming, P. A. (2014). Drinking problems on a 'simple' diet:

physiological convergence in nectar-feeding birds. *Journal of Experimental Biology*

217, 1015–1023.

Nicolson, S. W. and Nepi, M. (2005). Dilute Nectar in Dry Atmospheres: Nectar Secretion

Patterns in Aloe castanea (Asphodelaceae). *International Journal of Plant Sciences*

166, 227–233.

Padyšáková, E. and Janeček, Š. (2016). Sunbird hovering behavior is determined by both

the forager and resource plant. *Biotropica* **48**, 687–693.

Pyke, G. H., Pulliam, H. R. and Charnov, E. L. (1977). Optimal Foraging: A Selective

Review of Theory and Tests. *The Quarterly Review of Biology* **52**, 137–154.

Rico-Guevara, A. and Rubega, M. A. (2011). The hummingbird tongue is a fluid trap, not

a capillary tube. *Proc Natl Acad Sci USA* **108**, 9356.

Rico-Guevara, A. and Rubega, M. A. (2017). Functional morphology of hummingbird bill

tips: their function as tongue wringers. *Zoology* **123**, 1–10.

Rico-Guevara, A., Fan, T.-H. and Rubega, M. A. (2015). Hummingbird tongues are elastic

micropumps. *Proceedings of the Royal Society B: Biological Sciences* **282**, 20151014.

Rico-Guevara, A., Sustaita, D., Gussekloo, S., Olsen, A., Bright, J., Corbin, C. and

Dudley, R. (2019). Feeding in Birds: Thriving in Terrestrial, Aquatic, and Aerial Niches.

In *Feeding in Vertebrates: Evolution, Morphology, Behavior, Biomechanics* (ed. Bels,

V.) and Whishaw, I. Q.), pp. 643–693. Cham: Springer International Publishing.

Rico-Guevara, A., Hurme, K. J., Elting, R. and Russell, A. L. (2021). Bene"fit"

Assessment in Pollination Coevolution: Mechanistic Perspectives on Hummingbird

Bill–Flower Matching. *Integrative and Comparative Biology* **61**, 681–695.

Rico-Guevara, A., Hurme, K. J., Rubega, M. A. and Cuban, D. (2023). Nectar feeding

beyond the tongue: hummingbirds drink using phase-shifted bill opening, flexible

tongue flaps and wringing at the tips. *Journal of Experimental Biology* **226**, jeb245074.

Rolfe, S., Pieper, S., Porto, A., Diamond, K., Winchester, J., Shan, S., Kirveslahti, H.,

Boyer, D., Summers, A. and Maga, A. M. (2021). SlicerMorph: An open and

extensible platform to retrieve, visualize and analyse 3D morphology. Methods in

Ecology and Evolution 12, 1816–1825.

Schlamowitz, R., Hainsworth, F. R. and Wolf, L. L. (1976). On the Tongues of Sunbirds.

The Condor **78**, 104–107.

Schwenk, K. and Rubega, M. (2005). Diversity of Vertebrate Feeding Systems.pp. 1–41.

Stiles, F. G. (1981). Geographical Aspects of Bird-Flower Coevolution, with Particular

Reference to Central America. *Annals of the Missouri Botanical Garden* **68**, 323–351.

Tschapka, M., Gonzalez-Terrazas, T. P. and Knörnschild, M. (2015). Nectar uptake in

bats using a pumping-tongue mechanism. *Science Advances* **1**, e1500525.

Wainwright, P. C., McGee, M. D., Longo, S. J. and Patricia Hernandez, L. (2015).

Origins, Innovations, and Diversification of Suction Feeding in Vertebrates. *Integrative*

and Comparative Biology **55**, 134–145.

Wei, J., Rico-Guevara, A., Nicolson, S. W., Brau, F., Damman, P., Gorb, S. N., Wu, Z.

and Wu, J. (2023). Honey bees switch mechanisms to drink deep nectar efficiently.

Proceedings of the National Academy of Sciences **120**, e2305436120.

Williams, S. H. (2019). Feeding in Mammals: Comparative, Experimental, and Evolutionary

Insights on Form and Function. In *Feeding in Vertebrates: Evolution, Morphology,*

Behavior, Biomechanics (ed. Bels, V.) and Whishaw, I. Q.), pp. 695–742. Cham:

Springer International Publishing.

Winkler, D. W., Billerman, S. M. and Lovette, I. J. (2020). Sunbirds and Spiderhunters

(Nectariniidae), version 1.0. *Birds of the World*.

Figures:

Figure 1: Cross-sections of a sunbird (above, *Cyanomitra olivacea*) and hummingbird

(below, *Calypte anna*) from microCT images. The gray structures at the top and bottom are

lateral views of the bills. Dashed red lines crossing them correspond to the numbered cross-

sections. Cross-sections were taken at proportionally similar points along the longitudinal axis of

the bills, showing structures of the bill and tongue relevant to the feeding mechanics of each.

The cross-sections show the keratin structure (gray), the underlying bone (light-gray), and the

tongues (orange and yellow) of each bird. Both species have internal morphology of the bill that

guides the tongue along the longitudinal axis when it is extended. The sunbird has notably more

empty volume in the bill outside of the tongue grooves at the dorsal tip.

Figure 2: Kinematic measurements of a *Nectarinia famosa* tongue and red colored nectar

during multiple lick cycles. (A) Frame of a video showing the nectar reservoir (W = 5.9 mm)

and the bi-cylindrical distal portion of the tongue where only the tips are immersed in the nectar

(dyed magenta to improve visualization). Arrows indicate the position of the nectar, x_N, of the

tongue tip, x_T, and of the meniscus inside the tongue, x_M, with respect to an arbitrary fixed

point. Bill tip (not visible) towards the top of the frame. **(B)** Evolution of x_N, x_T and $d_{MT} = x_M -$

x_T as a function of time. **Inset**: Evolution of x_N as a function of time together with a linear fit. **(C)**

Evolution of the position of the tip of the tongue relative to the position of the nectar, $d_{TN} = x_T -$

x_N, as a function of time. The shaded area corresponds to the intervals of time where the

tongue is in the nectar. Average tongue dipping time, $T_D \simeq 0.07$ s. The maximal immersion

depth of the tongue in the nectar varies between 1.3 and 2.1 mm. **(D)** Evolution of the distance

h between the liquid meniscus inside the tongue and the nectar reservoir as a function of time.

t_0 corresponds to the time at which the tip of the tongue touches the nectar in the extension

phase for each capture process. Average licking period, $T_L \simeq 0.11$ s. **(E)** Evolution of h as a

function of $t - t_0$. **(F)** Average of the data shown in panel E (without the first lick). The region

where the average is performed is shown in panel E and corresponds to the time interval where

there is data for all the licks averaged.

490   **Figure 3:** Evolution of ⟨h⟩ as a function of time (see Fig. Yf) together with the evolution

491   predicted by capillary rise only (blue dashed curve), by capillary rise and inertial effects (green

492   dashed curve) and by capillary rise, inertial, effects, and a difference of pressure across

493   tongue's length varying linearly in time (red dashed dotted curve).

Figure 4: Biomechanical method of sunbird suction feeding. Sagittal cross sections of the

bill and tongue of a sunbird while feeding **(left)**. Transverse cross sections of the bill and tongue

while feeding **(right)**. (A) Initial position, (B) Tongue extension and pressing into upper palate,

(C) Tongue depression while tongue edges sealed against upper bill, (D) Partial retraction and

swallowing to remove nectar from proximal trough of tongue, (E) Re-extension for next lick,

some nectar often remains in the tubular portion of the tongue from the previous lick cycle, (F)

During re-extension, nectar left over from the previous lick will be drawn back in the tongue via

suction as the bird begins to generate a pressure differential across the length of the tongue

before the tongue tips make contact with the nectar reservoir.