# EZH2-driven immune evasion defines high-risk pediatric AML with t(16;21) *FUS::ERG* gene fusion.

Nathaniel J Buteyn,[1] Connor G Burke,[1] Vincent J Sartori,[1] Eve Deering-Gardner,[1] Zachary J DeBruine,[1] Dahlya Kamarudin,[1] Darrell P Chandler,[1] Alexander C Monovich,[2] Monika W Perez,[3] Joanna S Yi,[4] Rhonda E Ries,[5] Todd A Alonzo,[6,7] Russell JH Ryan,[2] Soheil Meshinchi,[5,6,8] and Timothy J Triche Jr.[1,9,10]

[1]Department of Epigenetics, Van Andel Research Institute, Grand Rapids, MI;

[2]Department of Pathology, University of Michigan, Ann Arbor, MI;

[3]Department of Genome Sciences, University of Washington, Seattle, WA;

[4]Department of Pediatrics, Texas Children's Cancer and Hematology Centers, Baylor College of Medicine, Houston, TX;

[5]Clinical Research Division, Fred Hutchinson Cancer Research Center, Seattle, WA;

[6]Children's Oncology Group, Monrovia, CA;

[7]Department of Translational Genomics, University of Southern California, Los Angeles, CA;

[8]Department of Pediatrics, University of Washington, Seattle, WA;

[9]Department of Translational Genomics, Keck School of Medicine, University of Southern California, Los Angeles, CA;

[10]Department of Pediatrics, College of Human Medicine, Michigan State University, East Lansing, MI.

**ABSTRACT**

Minimal improvement in outcomes for high-risk pediatric acute myeloid leukemia (pAML) patients has been made in the past decades. Nowhere is this more evident than in patients carrying a t(16;21)(p11;q22) *FUS::ERG* translocation; quick time to relapse and universal failure of hematopoietic stem cell transplant contribute to one of the lowest survival rates in childhood leukemia. Here, we have identified a unique, defining immune-evasion phenotype in *FUS::ERG* pAML driven by EZH2 and characterized by loss of MHC class I and II molecules and immune co-stimulatory receptors. This loss of immune engagement, present at diagnosis, allows pervasiveness of blasts that prove resistant to standard treatment. We demonstrate that treatment with the FDA-approved EZH2 inhibitor tazemetostat, in combination with IFN-γ, reverses the phenotype, re-expresses MHC receptor expression, and reduces blast viability. EZH2 inhibitors provide a novel therapeutic option for this high-risk population and may prove a beneficial supplemental treatment for *FUS::ERG* pAML.

**STATEMENT OF SIGNIFICANCE**

*FUS::ERG* pAML patients have dismal outcomes. Here we show a ubiquitous immune-evasive phenotype, defined by elevated EZH2 levels and loss of MHC class I and II receptors, is present in these patients at diagnosis. Treatment with the EZH2 inhibitor Tazemetostat successfully reverses the phenotype in a patient-derived cell line model.

**INTRODUCTION**

The E26 transformation-specific (ETS) family of transcription factors regulate a host of cellular functions including differentiation, proliferation, and apoptosis. When aberrantly expressed, often through chromosomal translocations resulting in chimeric fusion proteins, ETS factors drive a variety of cancers[1–5] including acute leukemias[6,7]. Translocations involving ETS factors have historically been considered high-risk in pediatric acute myeloid leukemia (pAML). In particular, the ETS family member ERG, when fused with the RNA-binding protein FUS in a t(16;21)(p11;q22) translocation, drives particularly poor prognosis in patients. Standard therapy has proven ineffective for *FUS::ERG* pAML; median time to relapse was <10 months, overall five-year survival was 33%, and 100% of patients who received stem cell transplant (SCT) died [8,9]. There is a desperate need for novel therapeutic options that target this high-risk subset.

Children's Oncology Group (COG) pAML trials AAML0531 and AAML1031[8,9] include one of the largest collections of pediatric AML patients with ETS fusions ever assembled. Analysis of this cohort allowed us, for the first time, to identify a pervasive immune-evasion phenotype ubiquitously present across *FUS::ERG* pAML. Through aberrant ERG activity, *FUS::ERG* pAML significantly overexpresses the histone-lysine methyltransferase EZH2 resulting in silencing of CIITA and class I and II major histocompatibility complex (MHC) receptors; loss of MHC expression has been shown to strongly correlate with relapse in leukemic disease[10–13]. We demonstrate here that treatment with the FDA-approved EZH2 inhibitor tazemetostat, when in combination with IFN-γ, successfully reverses the immune-evasion

phenotype and decreases blast viability in *FUS::ERG* AML cells.  Immune evasion often develops in a post-relapse setting[14], but in *FUS::ERG* pAML it presents at full strength at diagnosis, presenting a unique therapeutic problem.  Targeted EZH2 inhibition from the onset of treatment may prove vital in restoring immunogenicity to patients and extending time to relapse.

## METHODS

### AML Cohort

Patient data from the COG trials AAML0531[8] and AAML1031[9] were used for this study and contained RNA-sequencing data for 1,410 individuals.  Consent was obtained from all study participants in accordance with the Declaration of Helsinki.  The Fred Hutchinson Cancer Research Center Institutional Review Board and the COG Myeloid Biology Committee approved and oversaw the use of patient data.

### Non-negative Matrix Factorization and Gene Set Enrichment Analysis

Non-negative Matrix Factorization was run using the Rcpp Machine Learning Library (github.com/zdebruine/RcppML), with monte carlo cross-validation with 3 random replicates using Wold's method to identify an optimally robust rank at around k = 100.  The sample loadings in the "H" matrix were used for UMAP projection and for enrichment analysis of factors within *FUS::ERG* positive samples.  Gene set enrichment analysis was performed using the R-package msigdbr (Molecular Signatures Database) on the gene expression matrix resulting from the non-negative matrix factorization (NMF) of TARGET AML initiative patient data by way of the R-package RcppML[15,16]. NMF factors of interest were chosen and positively enriched gene sets were ranked by adjusted p-value. The top 20 MSigDB GO:BP gene sets were visualized using the R-package ggplot2[17].

### H3K27ac chromatin immunoprecipitation–sequencing processing

Data provided from pAML patients at Texas Children's Hospital.  Methods previously published.[18]

## Cell culture

TSU-1621MT cells were generously provided by Dr. Olaf Heidenreich of the Princess Máxima Center, Utrecht, The Netherlands . Cells were cultured at 37°C in 5% $CO_2$ in RPMI Medium 1640 (Life Technologies, Grand Island, NY) supplemented with 10% heat-inactivated FBS (Life Technologies), 56 U/ml/56 µg/ml penicillin/streptomycin (Invitrogen, Waltham, MA) and 10 ng/ml human recombinant G-CSF (Stemcell Technologies, Vancouver, Canada). Tazemetostat (EPZ-6438) was purchased from Selleck Chemicals (Houston, TX).  Recombinant human IFN-γ was purchased from R&D Systems (Minneapolis, MN).

## Flow cytometry

Cells were incubated with anti-human HLA-A, B, C antibody conjugated to Pacific Blue™ (Biolegend, San Diego, CA, USA) and anti-human HLA-DR, DP, DQ antibody conjugated to Alexa Fluor® 647 (Biolegend, San Diego, CA).  Dead cells were labeled with Propidium Iodide Ready Flow™ Reagent (Invitrogen).  Samples were analyzed for HLA expression using the Beckman Coulter CytoFLEX S flow cytometer and analyzed using FlowJo software (BD Life Sciences).

## Statistical analysis

For all analyses, a $p$ value of ≤ 0.05 was considered statistically significant.  Mantel-Cox Log-rank tests were used to analyze survival curves in Figure 1.  An unpaired t test was performed for analyses between two groups; for analyses involving multiple groups, data was analyzed by One-way ANOVA followed by Šídák's multiple comparisons tests.  Pearson correlation coefficients were calculated for scatterplots in Figures 3 and 5.

**RESULTS**

**Cohort characteristics**

The COG AAML0531 and AAML1031 cohorts comprise 2399 pAML patients, for which 66 patients have an ETS fusion.  Of those 66 patients, 59 fusions contained either the ETV6 or ERG transcription factor, with *FUS::ERG* accounting for 18.2% of all ETS fusions (**Figure 1A**).  The *FUS::ERG* fusions occurred in older patients (median 11.2 years) than the *ETV6::MNX1* fusions (median 9 months).  66.7% (8/12) of *FUS::ERG* patients were female and 41.7% were Hispanic, proportions that are higher than what is observed across all pAML patients in AAML0531/1031 (47.9% female and 16.1% Hispanic).  Of the 6 patients enrolled in AAML0531, four received gemtuzumab ozogamicin (Mylotarg) in addition to standard therapy.  Five out of six AAML1031 patients were enrolled in Arm B and received bortezomib in addition to standard therapy, with one receiving standard therapy alone (Arm A)[8,9].  Taken together, 8/12 *FUS::ERG* patients achieved complete remission (CR) after the first round of chemotherapy, and 11/12 were in remission after the second round.

**FUS::ERG patients are non-responsive to current therapies**

Whereas 51.2% of all AAML0531/1031 pAML patients had an event-free survival (EFS) event (classified as induction failure, relapse, or death), 83% (10/12) of *FUS::ERG* patients registered EFS events, the majority occurring after initial remission (7 relapses, 2 deaths, and 1 induction failure).  Median EFS for *FUS::ERG* patients was 299 days (compared to  629 days for all pAML).  Site of relapse/induction failure was reported for patients, excluding the two patients where primary EFS event was death, with 6/8 recorded in the bone marrow, one unknown, and one patient reporting relapse at a chloroma.  Five year overall survival rates for *FUS::ERG* patients was 33.3% (4/12), lower than the rest of pAML and the other major ETS fusions, with a median time of death after diagnosis of 376 days (**Figure 1B,C**).  25% of *FUS::ERG* patients (4/12) received SCT in first CR; of note, 100% of patients receiving SCT died (compared to 40.9% for ETS

fusion pAML and 37.4% for all pAML) (**Figure 1D**). Given the relatively low occurrence of ETS fusion patients, it has been difficult to properly categorize the population – risk group classification from the AAML1031 protocol initially labeled 33% of *FUS::ERG* patients "low-risk".  Only in its most recent edition of "Classification of Haematolymphoid Tumors (2022)" did the World Health Organization include *FUS::ERG* as a defining genetic alteration[19].  From **Figure 1**, however, it is clear that pAML patients harboring *FUS::ERG* fusions comprise an "ultra high-risk" population.

**EZH2-driven immune evasion**

The uniquely poor prognosis for *FUS::ERG* pAML – patients often achieve a CR before rapidly relapsing and are ubiquitously nonresponsive to SCT – suggests a distinct mechanism of resistance may be prevalent from the onset of disease.  As an initial test of this hypothesis, we analyzed RNA-sequencing data obtained from 1,410 AAML0531/1031 patients at diagnosis, ten of whom had a *FUS::ERG* fusion. UMAP dimensional reduction showed that *FUS::ERG* patients presented with a distinct transcriptional profile from all other pAML patients, including those harboring other ETS fusions (e.g., *ETV6::MNX1*, *ETV6::X*, and *HNRNPH1::ERG*; **Figure 2A**; two-color plot in **Supp Figure 1**).  We next used non-negative matrix factorization (NMF) on RNA-Seq data to characterize factors for which *FUS::ERG* samples were enriched. We identified one factor enriched for up-regulated genes, and three factors enriched for down-regulated genes (**Figure 2B**).  Gene set enrichment analysis (GSEA) provided the gene ontology of each NMF factor and showed that some of the most down-regulated genes were immune-related, especially in NMF factor 90 (**Figure 2C**).  Relative to all other patients harboring an ETS fusion, *FUS::ERG* patients were also characterized by significant up-regulation of the histone-lysine methyltransferase *EZH2* (**Figure 2D**).

We then looked more closely at immune-related gene expression in *FUS::ERG* patients relative to all other pAML patients, and quantified transcript levels from the most prevalent fusion groups in AAML0531/1031 (including the four major ETS fusion subsets *FUS::ERG*, *ETV6::MNX1*, *ETV6::X*, and

*HNRNPH1::ERG*).  Elevated levels of *EZH2* correlated with loss of MHC class I and II expression alongside suppression of the master regulator of MHC II, *CIITA* (**Figure 3A**).  Core-binding factor AML, those patients with a t(8;21)(q22;q22) *RUNX1::RUNX1T1* or inv(16)(p13q22)/t(16;16)(p13;q22) *CBFB::MYH11* fusion, as well as those with a *RUNX1::CBFA2T3* fusion, populated the opposite side of the heatmap.  These fusions displayed heightened MHC expression and immunogenic tumors characterized by low *EZH2*, high *CIITA*, and successful response to therapy in AAML0531/1031, consistent with other published cohorts.[20,21] **Figure 3B** shows positive correlation between expression of *CIITA* and mean MHC II score amongst all patients in AAML0531/1031 for which transcriptional data was available (n=1410).

Christopher, *et al.* showed that adult AML patients who relapse post-SCT demonstrate global loss of MHC receptors compared to levels at diagnosis (Adult Dx vs. Adult Relapse in **Figure 3C**)[14].  Importantly, when compared to the pAML diagnosis mean, *FUS::ERG* patients display the same, if not more exacerbated, MHC silenced phenotype as post-SCT relapse patients, but at diagnosis (pAML Dx vs. *FUS::ERG* Dx in **Figure 3C**).  Indeed, instead of acquiring immune deficiency via the selective pressures of standard induction and SCT, *FUS::ERG* patients enter treatment overwhelmingly defined by a treatment-resistant phenotype.  It is no surprise then that time-to-event is one of the shortest and response to SCT, nonexistent, compared to the rest of pAML.

AML is generally considered an immunogenic malignancy, where allogeneic SCT is effective at promoting graft-versus-leukemia (GvL) in lieu of graft-versus-host disease (GvHD)[22,23]. Effector immune cells engage their targets via MHC receptors, but also rely on a host of co-stimulatory receptors to initiate activation and facilitate clearance (**Figure 4A**). We therefore analyzed RNA-seq data for expression of ligands for natural killer cell and T cell co-stimulatory immune receptors that are typically expressed on monocytic cells.  As shown in **Figure 4B**, *FUS::ERG* pAML patients showed the lowest mean expression of these ligands compared to the other major pAML fusions (average expression [TPM] of each ligand across pAML shown in **Figure 4C**).  Of note, the ligands for the T cell receptor CD28 were almost completely absent in *FUS::ERG* patients.  Moreover, *FUS::ERG* patients showed a significant increase in transcripts for

TGF-β (*TGFB1*) and the TIM-3 ligand Galectin-9 (*LGALS9*), both inducers of immunosuppressive T regulatory (Treg) cells (**Figure 4D**)[24–26]. The resistance of *FUS::ERG* patients to standard therapy might therefore be explained by both the existence of an immune-desensitized leukemic blast population and a stunted immune effector cell population.

**ERG drives EZH2 over-expression**

All ETS transcription factors contain a winged helix-turn-helix DNA binding domain that interacts with the GGA(A/T) motif known as an ETS-binding sites (EBS). Numerous variants of the *FUS::ERG* fusion have been reported in pediatric leukemias, yet all retain their DNA binding domain[27]. Additionally, repeats of the GGAA sequence have been shown to recruit ETS proteins and facilitate initiation of transcription[28]. Chromatin immunoprecipitation sequencing (ChIP-seq) for ERG in the patient-derived *FUS::ERG* cell line TSU-1621MT shows binding of ERG to one such GGAA repeat site in the promoter region of *EZH2* (**Figure 5A**)[Data from Sotoca, *et al.*[29]]. This exaggerated ERG binding at EZH2 may be a unique feature of *FUS::ERG* leukemia. In primary pAML patient H3K27ac ChIP-seq data provided from Texas Children's Hospital (data from Perez, *et al.*[18]) the GGAA repeat site in EZH2 was heavily acetylated, indicative of higher transcriptional activation, in only the *FUS::ERG* patient. Other patients, including those with MLL fusions, core-binding factor AML, and even another ETS fusion – *ETV6::MNX1*, showed no H3K27ac at the locus (**Figure 5B**). Predictably, and matching the RNA-seq data from AAML0531/1031, the *FUS::ERG* patient additionally showed a lack of accessible chromatin marks at HLA loci – including HLA-DRA show in **Figure 5C**. Further analysis of H3K27 acetylation at EBS GGAA repeats across pediatric malignancies revealed that *FUS::ERG* expressed a global repression of acetylation, unique amongst the other pediatric leukemias and similar only to the ETS-driven Ewing's Sarcoma (**Figure 5D; Supp Figure 2**).

EZH2 itself operates canonically as the catalytic subunit of the polycomb repressive complex 2 (PRC2) to carry out histone methyl transferase activity[30,31]. Expression of other members of the complex,

both core (EED, SUZ12) and auxiliary subunits (RbAp46, RbAp48, JARID2, AEBP2), are not significantly altered in *FUS::ERG* compared the rest of pAML (**Figure 5E**). ERG has been previously shown to regulate *EZH2* in prostate cancer, where up to fifty percent of patients harbor the *TMPRSS2::ERG* fusion, localizing to the *EZH2* promoter and inducing transcription[32]. Analysis of RNA-seq from AAML0531/1031 supported these findings in pAML – a correlation is seen between expression of *ERG* and *EZH2* (Pearson = 0.6045)(**Figure 5F**).

Interestingly, there was a strong correlation observed between the expression of *FUS* and *EZH2* (Pearson = 0.9255). *FUS*, or Fused in Sarcoma, is part of the FET family of RNA-binding proteins; other members include *TAF15* and *EWSR1*. Like *FUS*, a strong correlation was observed between transcript levels of *EWSR1* and *EZH2* (Pearson = 0.9677); *TAF15* did not correlate (Pearson = -0.2095) (**Figure 5F**). As both wild-type *ERG* and *FUS* correlate with *EZH2* expression, it is unsurprising that their fusion, contributing aberrant activity of both members, significantly upregulates *EZH2* above the general pAML population as well as above patients with a non-ERG FET-ETS fusion (e.g. *FUS::FEV*, *FUS:FLI1*, *EWSR1::ELF5*, *EWSR1::FEV*) or another -ERG fusion (e.g. *HNRNPH1::ERG*)(**Figure 5G**). Furthermore, only *FUS::ERG* showed elevated wild-type ERG expression, potentially via feedback mechanisms with EZH2[33] or direct transcriptional control (an EBS exists within the *ERG* gene). Wild-type *FUS* was not differentially expressed between all pAML, other FET-ETS, other -ERG fusions, and *FUS::ERG* (**Figure 5G**). The lack of heightened *EZH2* observed in other FET-ETS fusions when compared to *FUS::ERG* suggests that aberrant FUS activity from the N-terminus chimeric protein fragment may not be sufficient to over-induce *EZH2*. Taken together with the heightened epigenetic-mediated accessibility to EZH2, *FUS::ERG* pAML seems uniquely primed towards EZH2-driven leukemogenesis.

**Tazemetostat restores MHC expression**

To test if inhibition of EZH2 could reverse the immune-evasion phenotype, we treated a patient-derived *FUS::ERG* cell line (TSU-1621MT) with the FDA-approved EZH2 inhibitor tazemetostat. The reversal of epigenetic suppressive H3K27me3 marks by tazemetostat allows access to *CIITA,* which can be transcriptionally activated via treatment with IFN-γ; it has been long established that IFN-γ potently induces *CIITA* via its promoter IV and drives expression of MHC II molecules[34,35]. As predicted, combined tazemetostat and IFN-γ treatment robustly synergized to significantly upregulate MHC II receptor surface expression after 7 days of treatment *in vitro* (33.5 fold increase over Ctrl). MHC I molecules were also upregulated significantly (49.5 fold increase over Ctrl) – CIITA has shown to moderately control this class as well (**Figure 6A,B**).[36,37] Additionally, tazemetostat significantly reduced cell viability (48.9% of Ctrl)(**Figure 6C**). The robust reversal of the *FUS::ERG* immune-evasion phenotype in patient derived cells suggests that tazemetostat might be a novel therapeutic option for *FUS::ERG* pAML patients.

Given our data, the mechanism we propose (**Figure 6D**) suggests that the aberrant activity from the chimeric fusion protein *FUS::ERG* directly induces *EZH2* transcription. *EZH2* upregulation is further exacerbated by heightened expression of wildtype ERG. Elevated EZH2 carries out its canonical deposition of H3K27me3 which leads to silencing of *CIITA* and suppression of immune receptor expression. Unchecked by host immunity, or donor immunity in the scenario of SCT, leukemic blasts proliferate unfettered, leading to rapid relapse, SCT failure, and one of the poorest survival rates in childhood leukemia.

**DISCUSSION**

The breadth of AAML0531/1031 allowed for analysis of a cohort of ETS patients the size of which was not previously available. Here, we show that *FUS::ERG* fusions drive EZH2 overexpression, a novel immune-evasion phenotype, and adverse outcomes in pAML patients. Our data and conceptual model (**Figure 6D**)

are consistent with known roles for EZH2 in silencing *CIITA*[38], an essential component of the enhanceosome that positively regulates MHC II molecules. The absence of MHC receptors on leukemic blasts (**Figure 3**) and the resulting loss of immunogenicity is explicitly tied to relapse in AML[10–14,39]. Monocytic leukemic blasts also express activating and inhibitory ligands on their cell surface, which interact with effector immune cells receptors to mediate immunosurveillance and target elimination. For example, NKG2D ligands increase susceptibility to allorecognition and NK cell elimination[40]; activity of NKp30 with its ligand NCR3LG1 (B7-H6) is correlated with improved overall survival in AML[41,42]; patients with high expression of DNAM-1 and its ligands have significantly longer remission and overall survival[43]; and expression of CD80 and CD86 is associated with prolonged remission[44]. AML patients with an activating NK ligand phenotype also see a significant improvement in 2-year overall survival (59.6% vs 24.4%)[45]. In *FUS::ERG* patients, these ligands were down-regulated while inducers of Treg were upregulated (**Figure 4**). Thus, it appears *FUS::ERG* patients enter treatment at diagnosis with an immune-desensitized blast population that is not susceptible to standard induction or SCT, resulting in an overall prognosis that is one of the worst in pediatric leukemia (**Figure 1**).

Induction and intensification chemotherapies release tumor neoantigens which, when loaded on MHC receptors, elicit target elimination via effector immune cells[46]; renewal of this anti-leukemic mechanism could prove significantly beneficial in pediatric *FUS::ERG* patients. Immediate intervention at time of diagnosis could successfully bridge patients to transplant where further maintenance could elicit the desired GvL. Moreover, treatment with tazemetostat may be particularly effective in conjunction with allo-SCT to elicit potent GvL effects. There is a window immediately following SCT where endogenous inflammatory cytokines such as IFN-γ are elevated[47,48]. Treating these patients with tazemetostat should make *CIITA* transcriptionally accessible to the inevitable interferon cascade, thus up-regulating MHC receptors and enabling donor lymphocytes to engage with any remaining leukemic blasts.

Modeling ETS fusions, and *FUS::ERG* in particular, brings certain challenges, both due to the relatively limited patient population as well as the inherent biology that defines the fusions. The locations

of ETS-binding GGAA tandem repeat sites has been shown to be poorly conserved between humans and rodents[49], and prevents accurate modeling of the ETS-driven mechanisms of leukemogenesis explored here (**Figure 5A**).  Moreover, giving the vital role immunity plays in *FUS::ERG* biology, xenograft modeling in immune-incompetent mice is also ineffective.  For this study we utilized a patient-derived cell line that matched the MHC absent phenotype observed in pAML *FUS::ERG* patients (confirmed by flow cytometry).  Given the ubiquitousness of the phenotype amongst patients – all AAML0531/1031 *FUS::ERG* patients expressed EZH2 at least 1 standard deviation above the pAML population mean; mean standard score was 6.15 (**Figure 2D**) – and the robust reversal of the observed phenotype after tazemetostat treatment, consideration of EZH2 inhibition for this high-risk subset is warranted.

ETS fusions make up a relatively small percentage of pAML cases (in AAML0531/1031 = 4-5%), and *FUS::ERG* an even smaller percentage.  However, the EZH2-driven immune-evasion phenotype described here may be more widely operative – both in other pAML subtypes and other cancers.  For example, patients in the AAML0531/1031 cohort who fall within the top 10% of EZH2 expression have significantly lower HLA-DRA expression compared to those with the lowest 10% of EZH2 expression (P-val = 0.0073).  The top 10% EZH2 group includes the *FUS::ERG* patients, but also other ETS fused leukemias (*ETV6::INO80D*, *ETV6::FOXO1*, and *ERG::GATA2)*; other high-risk fusions (*CBFA2T3::GLIS2* and *KMT2A::MLLT10)*; as well as patients with traditionally low-risk fusions (*CBFB::MYH11* and *RUNX1::RUNX1T1)*.  In fact, within the top 10% of EZH2 group (n =141), there are 37 unique fusions, and 32 patients who do not have an identified driver fusion.  It is possible that augmenting standard treatment with an EZH2 inhibitor may benefit the 61.7% of these patients who experience an adverse event during standard treatment (higher than the rate for all of AAML0531/1031 = 51.2%).

Given the role EZH2 plays in driving oncogenesis in a number of malignancies[50,51], there has been considerable effort focused on developing EZH2 small molecule inhibitors.  Tazemetostat (Tazverik) is an orally available, S-adenosyl methionine (SAM) competitive inhibitor of EZH2 that has been approved by the FDA for adults (16+) with metastatic or locally advanced epithelioid sarcoma and adults with relapsed

or refractory follicular lymphoma[52]. Tazemetostat was reported to be well tolerated with a favorable safety profile[53,54]. It is also being used to suppress oncogenic transformation and sensitize Ewing's sarcoma patients to CAR T cell therapy[50,55], for treatment of pediatric rhabdoid tumors harboring SMARCB1/SMARCA4 or EZH2 alterations[56], and is under consideration for prostate cancer patients with a TMPRSS2::ERG fusion and aberrant EZH2 activity[57].

In summary, there is currently no effective treatment for *FUS::ERG* pAML, but we have identified for the first time a distinct immune evasion phenotype present at diagnosis and readily targetable by the EZH2 inhibitor tazemetostat. Reinvigoration of the patient's immune response has the potential to thoroughly eliminate residual disease after standard induction as well as elicit potent graft-versus-leukemic effects upon introduction of donor immune effector cells (via SCT or donor lymphocyte infusions). Success here could radically alter clinical outcomes for *FUS::ERG* pAML.

**ACKNOWLEDGEMENTS**

Research was supported by Van Andel Institute's T32 Cancer Epigenetics Training Program (5T32CA251066-03), the Children's Oncology Group, the National Cancer Institute of the National Institutes of Health (NCTN Operations Center Grant U10CA180886, NCTN Statistics & Data Center Grant U10CA180899), and St. Baldrick's Foundation.

Disclaimer: The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health

# FIGURE 1

# FIGURE 2

# FIGURE 3

# FIGURE 4

# FIGURE 5

# FIGURE 6

# SUPPLEMENTAL FIGURE 1

**RNA-seq by Primary Fusion**

**FUSION**
- 🔴 FUS::ERG
- ⚫ pAML

# SUPPLEMENTAL FIGURE 2

**Figure 1: Characteristics of ETS fused pAML patients in trials AAML0531/1031.** A) Demographic and clinical characteristics of 2399 patients enrolled in Childrens Oncology Group trials AAML0531[8] and AAML1031;[9] major ETS fusions further expanded. B) Event-free and overall survival curves of ETS fused patients compared to all pAML. C) Survival curves for *FUS::ERG* patients compared to assigned risk categories from AAML0531/1031. D) Breakdown of frequency and success rate of hematopoietic stem cell transplant after first CR for ETS fused patients compared to all pAML.

**Figure 2: Loss of immune function characterizes *FUS::ERG* pAML.** A) UMAP projection of gene expression from 1410 AAML0531/1031 patients who had RNA-sequencing at diagnosis. Sorted by major pAML fusion. B) Non-negative matrix factorization (90-factor) of RNA-sequencing data showing factors enriched for *FUS::ERG* signatures. C) Gene set enrichment analysis for Gene Ontology: Biological Processes of each enriched factor. Down-regulated factors were heavily enriched for immune related processes. D) Expression of *EZH2* (TPM) amongst major pAML fusions (AAML0531/1031; n = 1410). ****p-val ≤0.0001.

**Figure 3: EZH2 overexpression correlates with global loss of MHC receptors.** A) Heatmap of *EZH2*, *CIITA*, and MHC receptor expression among the major pAML fusions (TPM z-score; n = 1410). B) Scatterplot of MHC II expression (mean z-score) verses *CIITA* in AAML0531/1031 (n = 1410; Pearson = 0.6480); *FUS::ERG* patients highlighted in red. C) Expression of MHC I and II genes in adult AML at diagnosis (black) and relapse post-SCT (grey) from Christopher, *et al*.[14] was compared to AAML0531/1031 pAML at diagnosis (white) and *FUS::ERG* pAML at diagnosis (red).

**Figure 4: Immune co-stimulatory receptors are downregulated in *FUS::ERG*.** A) Natural killer cell and T cell immune receptor ligands found on the surface of pAML blasts. Graphic created with biorender.com. B) Expression of ligands in the major pAML fusions (TPM z-score; n = 1410). C) Average expression of ligands in TPM across pAML. D) Expression of pro-Treg stimulants *TGFB1* and *LGALS9* in pAML versus *FUS::ERG* (n = 1410). ****p-val ≤0.0001; *p-val ≤0.05.

**Figure 5: Aberrant ERG activity correlates with overexpression of *EZH2*.** A) ChIP-seq for ERG in TSU-1621MT *FUS::ERG* cells at the GGAA repeat sequence in the promoter region of *EZH2*.[29] B,C) H3K27ac ChIP-seq in primary pAML patients[18] at the EZH2 (B) and HLA-DRA loci (C). D) H3K27 acetylation analysis

at 6x GGAA motifs conducted on primary pAML[18], T-lymphoid/myeloid mixed-phenotype acute leukemia (TMPAL)[58], Desmoplastic small round cell tumors (DSRCT)[59], and Ewing's Sarcoma[60] patients with CD34 hematopoietic progenitor cells (HPC) and mesenchymal stem cell (MSC) controls[61]; pAML *FUS::ERG*[18] cases highlighted in blue. Controls shown in Supplemental figure 2.  E) Expression of PRC2 members in pAML compared to *FUS::ERG* (n = 1410).  F) Scatter plots of expression of *EZH2* versus *ERG* (Pearson = 0.6045), *FUS* (Pearson = 0.9255), and *EWSR1* (Pearson = 0.9677) (n = 1410).  G) Expression of *EZH2*, *ERG*, and *FUS* among pAML, other FET-ETS, other *ERG*, and *FUS::ERG* (n = 1410).  ****p-val ≤0.0001; *p-val ≤0.05.

**Figure 6: Tazemetostat restores MHC receptor expression in *FUS::ERG* cells.** A,B) TSU-1621MT cells were treated with 10 µM tazemetostat and/or 10 ng/ml IFN-γ on days 0, 2, 4, and 6.  Cells were collected on day 7 and surface expression of MHC I and II receptors was assessed via flow cytometry. Representative histogram shown in (A) and mean florescent intensity (MFI) graphed in (B) (n = 3).  C) Cell viability in TSU-1621MT cells measured by trypan blue exclusion assay on day 6 of treatment (schedule identical to that in A; n = 3). D) Proposed mechanism for immune suppression in *FUS::ERG* pAML. ****p-val ≤0.0001; ***p-val ≤0.001; **p-val ≤0.01; *p-val ≤0.05.

**Supplemental Figure 1:** UMAP projection of gene expression from 1410 AAML0531/1031 patients who had RNA-sequencing at diagnosis.  *FUS::ERG* patients highlighted in red.

**Supplemental Figure 2:** H3K27 acetylation analysis at 6x GGAA motifs conducted on primary pAML[18], T-lymphoid/myeloid mixed-phenotype acute leukemia (TMPAL)[58], Desmoplastic small round cell tumors (DSRCT)[59], and Ewing's Sarcoma[60] patients with CD34 hematopoietic progenitor cells (HPC) and mesenchymal stem cell (MSC) controls[61]; pAML *FUS::ERG*[18] cases highlighted in blue..  Analysis was conducted on 3x repeats with one mismatch (A), 3x repeats with no mismatch (B), and 6x repeats with no mismatch (C).  Random genomic regions and housekeeping promoters shown in (D) and (E) as controls.

1.	Chi, P. *et al.* ETV1 is a lineage survival factor that cooperates with KIT in gastrointestinal stromal tumours. *Nature* **467**, 849–853 (2010).

2.	Tognon, C. *et al.* Expression of the ETV6-NTRK3 gene fusion as a primary event in human secretory breast carcinoma. *Cancer Cell* **2**, 367–376 (2002).

3.	Tomlins, S. A. *et al.* Recurrent Fusion of TMPRSS2 and ETS Transcription Factor Genes in Prostate Cancer. *Science* **310**, 644–648 (2005).

4.	de Alava, E. *et al.* EWS-FLI1 fusion transcript structure is an independent determinant of prognosis in Ewing's sarcoma. *J. Clin. Oncol.* **16**, 1248–1255 (1998).

5.	Jané-Valbuena, J. *et al.* An Oncogenic Role for ETV1 in Melanoma. *Cancer Res.* **70**, 2075–2084 (2010).

6.	Sa, S. *et al.* TEL/AML1 fusion resulting from a cryptic t(12;21) is the most common genetic lesion in pediatric ALL and defines a subgroup of patients with an excellent prognosis. *Leukemia* **9**, 1985–1989 (1995).

7.	Shimizu, K. *et al.* An ets-related gene, ERG, is rearranged in human myeloid leukemia with t(16;21) chromosomal translocation. *Proc. Natl. Acad. Sci.* **90**, 10280–10284 (1993).

8.	Children's Oncology Group. *A Phase III Randomized Trial of Gemtuzumab Ozogamicin (Mylotarg) Combined With Conventional Chemotherapy for De Novo Acute Myeloid Leukemia (AML) in Children, Adolescents, and Young Adults*. https://clinicaltrials.gov/study/NCT00372593 (2021).

9.	National Cancer Institute (NCI). *A Phase III Randomized Trial for Patients With De Novo AML Using Bortezomib and Sorafenib (NSC# 681239, NSC# 724772) for Patients With High Allelic Ratio FLT3/ITD*. https://clinicaltrials.gov/study/NCT01371981 (2023).

10.	Vago, L. *et al.* Loss of Mismatched HLA in Leukemia after Stem-Cell Transplantation. *N. Engl. J. Med.* **361**, 478–488 (2009).

11.	Stölzel, F. *et al.* Clonal evolution including partial loss of human leukocyte antigen genes favoring extramedullary acute myeloid leukemia relapse after matched related allogeneic hematopoietic stem cell transplantation. *Transplantation* **93**, 744–749 (2012).

12.	Masuda, K. *et al.* Loss or down-regulation of HLA class I expression at the allelic level in freshly isolated leukemic blasts. *Cancer Sci.* **98**, 102–108 (2007).

13.	Dermime, S. *et al.* Immune escape from a graft-versus-leukemia effect may play a role in the relapse of myeloid leukemias following allogeneic bone marrow transplantation. *Bone Marrow Transplant.* **19**, 989–999 (1997).

14.	Christopher, M. J. *et al.* Immune Escape of Relapsed AML Cells after Allogeneic Transplantation. *N. Engl. J. Med.* **379**, 2330–2341 (2018).

15.	DeBruine, Z. J., Melcher, K. & Triche, T. J. Fast and robust non-negative matrix factorization for single-cell experiments. 2021.09.01.458620 Preprint at https://doi.org/10.1101/2021.09.01.458620 (2021).

16.	igor. msigdbr: MSigDB Gene Sets for Multiple Organisms in a Tidy Data Format. (2023).

17.	ggplot2: Elegant Graphics for Data Analysis (3e). https://ggplot2-book.org/.

18.     Perez, M. W. *et al.* Defining the transcriptional control of pediatric AML highlights RARA as a superenhancer-regulated druggable dependency. *Blood Adv.* **5**, 4864–4876 (2021).

19.     Khoury, J. D. *et al.* The 5th edition of the World Health Organization Classification of Haematolymphoid Tumours: Myeloid and Histiocytic/Dendritic Neoplasms. *Leukemia* **36**, 1703–1719 (2022).

20.     Schlenk, R. f. *et al.* Individual Patient Data–Based Meta-Analysis of Patients Aged 16 to 60 Years With Core Binding Factor Acute Myeloid Leukemia: A Survey of the German Acute Myeloid Leukemia Intergroup. *J. Clin. Oncol.* **22**, 3741–3750 (2004).

21.     Marcucci, G. *et al.* Prognostic Factors and Outcome of Core Binding Factor Acute Myeloid Leukemia Patients With t(8;21) Differ From Those of Patients With inv(16): A Cancer and Leukemia Group B Study. *J. Clin. Oncol. Off. J. Am. Soc. Clin. Oncol.* **23**, 5705–17 (2005).

22.     Passweg, J. R. *et al.* The EBMT activity survey report 2017: a focus on allogeneic HCT for nonmalignant indications and on the use of non-HCT cell therapies. *Bone Marrow Transplant.* **54**, 1575–1585 (2019).

23.     Stern, M. *et al.* Sensitivity of hematological malignancies to graft-versus-host effects: an EBMT megafile analysis. *Leukemia* **28**, 2235–2240 (2014).

24.     Pang, N. *et al.* Activated Galectin-9/Tim3 promotes Treg and suppresses Th1 effector function in chronic lymphocytic leukemia. *FASEB J. Off. Publ. Fed. Am. Soc. Exp. Biol.* **35**, e21556 (2021).

25.     Zhu, C. *et al.* The Tim-3 ligand galectin-9 negatively regulates T helper type 1 immunity. *Nat. Immunol.* **6**, 1245–1252 (2005).

26.     Oh, S. A. & Li, M. O. TGF-β: Guardian of T Cell Function. *J. Immunol. Baltim. Md 1950* **191**, 3973–3979 (2013).

27.     PANAGOPOULOS, I., GORUNOVA, L., ZELLER, B., TIERENS, A. & HEIM, S. Cryptic FUS-ERG fusion identified by RNA-sequencing in childhood acute myeloid leukemia. *Oncol. Rep.* **30**, 2587–2592 (2013).

28.     Uchiumi, F., Miyazaki, S. & Tanuma, S. The possible functions of duplicated ets (GGAA) motifs located near transcription start sites of various human genes. *Cell. Mol. Life Sci.* **68**, 2039–2051 (2011).

29.     Sotoca, A. M. *et al.* The oncofusion protein FUS–ERG targets key hematopoietic regulators and modulates the all-trans retinoic acid signaling pathway in t(16;21) acute myeloid leukemia. *Oncogene* **35**, 1965–1976 (2016).

30.     Viré, E. *et al.* The Polycomb group protein EZH2 directly controls DNA methylation. *Nature* **439**, 871–874 (2006).

31.     Cao, R. *et al.* Role of histone H3 lysine 27 methylation in Polycomb-group silencing. *Science* **298**, 1039–1043 (2002).

32.     Kunderfranco, P. *et al.* ETS Transcription Factors Control Transcription of EZH2 and Epigenetic Silencing of the Tumor Suppressor Gene Nkx3.1 in Prostate Cancer. *PLOS ONE* **5**, e10547 (2010).

33.     Zoma, M. *et al.* EZH2-induced lysine K362 methylation enhances TMPRSS2-ERG oncogenic activity in prostate cancer. *Nat. Commun.* **12**, 4147 (2021).

34.      Steimle, V., Siegrist, C. A., Mottet, A., Lisowska-Grospierre, B. & Mach, B. Regulation of MHC class II expression by interferon-gamma mediated by the transactivator gene CIITA. *Science* **265**, 106–109 (1994).

35.      Muhlethaler-Mottet, A., Di Berardino, W., Otten, L. A. & Mach, B. Activation of the MHC class II transactivator CIITA by interferon-gamma requires cooperative interaction between Stat1 and USF-1. *Immunity* **8**, 157–166 (1998).

36.      Gobin, S. J., Peijnenburg, A., Keijsers, V. & van den Elsen, P. J. Site alpha is crucial for two routes of IFN gamma-induced MHC class I transactivation: the ISRE-mediated route and a novel pathway involving CIITA. *Immunity* **6**, 601–611 (1997).

37.      Martin, B. K. *et al.* Induction of MHC class I expression by the MHC class II transactivator CIITA. *Immunity* **6**, 591–600 (1997).

38.      Truax, A. D., Thakkar, M. & Greer, S. F. Dysregulated Recruitment of the Histone Methyltransferase EZH2 to the Class II Transactivator (CIITA) Promoter IV in Breast Cancer Cells. *PLoS ONE* **7**, e36013 (2012).

39.      Romero, D. Reduced MHCII levels in relapsed AML. *Nat. Rev. Clin. Oncol.* **16**, 7–7 (2019).

40.      Diermayr, S. *et al.* NKG2D ligand expression in AML increases in response to HDAC inhibitor valproic acid and contributes to allorecognition by NK-cell lines with single KIR-HLA class I specificities. *Blood* **111**, 1428–1436 (2008).

41.      Chretien, A.-S. *et al.* NKp30 expression is a prognostic immune biomarker for stratification of patients with intermediate-risk acute myeloid leukemia. *Oncotarget* **8**, 49548–49563 (2017).

42.      Fiegler, N. *et al.* Downregulation of the activating NKp30 ligand B7-H6 by HDAC inhibitors impairs tumor cell recognition by NK cells. *Blood* **122**, 684–693 (2013).

43.      Chashchina, A. *et al.* DNAM-1/CD226 is functionally expressed on acute myeloid leukemia (AML) cells and is associated with favorable prognosis. *Sci. Rep.* **11**, 18012 (2021).

44.      Whiteway, A., Corbett, T., Anderson, R., Macdonald, I. & Prentice, H. G. Expression of co-stimulatory molecules on acute myeloid leukaemia blasts may effect duration of first remission. *Br. J. Haematol.* **120**, 442–451 (2003).

45.      Mastaglio, S. *et al.* Natural killer receptor ligand expression on acute myeloid leukemia impacts survival and relapse after chemotherapy. *Blood Adv.* **2**, 335–346 (2018).

46.      Xie, N. *et al.* Neoantigens: promising targets for cancer therapy. *Signal Transduct. Target. Ther.* **8**, 1–38 (2023).

47.      Rybicka-Ramos, M. *et al.* Profiles of interferon-gamma and interleukin-2 in patients after allogeneic hematopoietic stem cell transplantation. *World J. Biol. Chem.* **13**, 72–82 (2022).

48.      Melenhorst, J. J. *et al.* Cytopenia and leukocyte recovery shape cytokine fluctuations after myeloablative allogeneic hematopoietic stem cell transplantation. *Haematologica* **97**, 867–873 (2012).

49.      Kodgule, R. *et al.* ETV6 Deficiency Unlocks ERG-Dependent Microsatellite Enhancers to Drive Aberrant Gene Activation in B-Lymphoblastic Leukemia. *Blood Cancer Discov.* **4**, 34–53 (2023).

50.     Richter, G. H. S. *et al.* EZH2 is a mediator of EWS/FLI1 driven tumor growth and metastasis blocking endothelial and neuro-ectodermal differentiation. *Proc. Natl. Acad. Sci.* **106**, 5324–5329 (2009).

51.     Melling, N. *et al.* Overexpression of enhancer of zeste homolog 2 (EZH2) characterizes an aggressive subset of prostate cancers and predicts patient prognosis independently from pre- and postoperatively assessed clinicopathological parameters. *Carcinogenesis* **36**, 1333–1340 (2015).

52.     TAZVERIK (tazemetostat) tablets, for oral use. Initial U.S. Approval: 2020. (2020).

53.     Proudman, D. *et al.* A Matching-Adjusted Indirect Comparison of Single-Arm Trials in Patients with Relapsed or Refractory Follicular Lymphoma Who Received at Least Two Prior Systemic Treatments: Tazemetostat was Associated with a Lower Risk for Safety Outcomes Versus the PI3-Kinase Inhibitors Idelalisib, Duvelisib, Copanlisib, and Umbralisib. *Adv. Ther.* **39**, 1678–1696 (2022).

54.     Stacchiotti, S. *et al.* Safety and efficacy of tazemetostat, a first-in-class EZH2 inhibitor, in patients (pts) with epithelioid sarcoma (ES) (NCT02601950). *J. Clin. Oncol.* **37**, 11003–11003 (2019).

55.     Kailayangiri, S. *et al.* EZH2 Inhibition in Ewing Sarcoma Upregulates GD2 Expression for Targeting with Gene-Modified T Cells. *Mol. Ther.* **27**, 933–946 (2019).

56.     Chi, S. N. *et al.* Tazemetostat for tumors harboring SMARCB1/SMARCA4 or EZH2 alterations: results from NCI-COG pediatric MATCH APEC1621C. *J. Natl. Cancer Inst.* **115**, 1355–1363 (2023).

57.     Choudhury, A. D. *et al.* A phase Ia/Ib study of talazoparib in combination with tazemetostat in metastatic castration-resistant prostate cancer (mCRPC). *J. Clin. Oncol.* **40**, TPS195–TPS195 (2022).

58.     Botten, G. A. *et al.* Structural variation cooperates with permissive chromatin to control enhancer hijacking–mediated oncogenic transcription. *Blood* **142**, 336–351 (2023).

59.     Lamhamedi-Cherradi, S.-E. *et al.* The androgen receptor is a therapeutic target in desmoplastic small round cell sarcoma. *Nat. Commun.* **13**, 3057 (2022).

60.     Riggi, N. *et al.* EWS-FLI1 utilizes divergent chromatin remodeling mechanisms to directly activate or repress enhancer elements in Ewing sarcoma. *Cancer Cell* **26**, 668–681 (2014).

61.     Subramanian, S. *et al.* Genome-wide transcription factor-binding maps reveal cell-specific changes in the regulatory architecture of human HSPCs. *Blood* **142**, 1448–1462 (2023).