Title: Development of RT-RPA-based point-of-care tests for epidemic arthritogenic

alphaviruses

Authors: Sainetra Sridhar[1,2], Prince Baffour Tonto[1,2], Lily Lumkong[1,2], Bobby Brooke

Herrera[1,2*]

Affiliations: [1]Rutgers Global Health Institute, Rutgers University, New Brunswick, NJ,

USA

[2]Department of Medicine, Division of Allergy, Immunology, and Infectious Diseases, and

Child Health Institute of New Jersey, Rutgers Robert Wood Johnson Medical School,

Rutgers University, New Brunswick, NJ, USA

***Corresponding Author**: BBH, bherrera@globalhealth.rutgers.edu

Keywords: chikungunya virus, o'nyong-nyong virus, Mayaro virus, recombinase

polymerase amplification, rapid tests, amplicon sequencing

Abstract:

Chikungunya (CHIKV), o'nyong-nyong (ONNV), and Mayaro (MAYV) viruses are

transmitted by mosquitoes and known to cause a debilitating arthritogenic syndrome.

These alphaviruses have emerged and re-emerged, leading to outbreaks in tropical and

subtropical regions of Asia, South America, and Africa. Despite their prevalence, there

persists a critical gap in the availability of sensitive and virus-specific point-of-care

(POC) diagnostics. Traditional immunoglobulin-based tests such as enzyme-linked

immunosorbent assay (ELISAs) often yield cross-reactive results due to the close

genetic relationship between these viruses. Molecular diagnostics such as quantitative

polymerase chain reaction (qPCR) offer high sensitivity but are limited by the need for

specialized laboratory equipment. Recombinase polymerase amplification (RPA), an

isothermal amplification method, presents a promising alternative to qPCR, providing

rapid results with minimal equipment requirements. Here, we report the development

and validation of three virus-specific RPA-based rapid tests for CHIKV, ONNV, and

MAYV. These tests demonstrated both speed and sensitivity, capable of detecting 10

viral copies within 20 minutes of amplification, without exhibiting cross-reactivity to

related alphaviruses and other arboviruses. Furthermore, we evaluated the clinical

potential of these tests using serum and tissue samples from CHIKV, ONNV, and

MAYV-infected mice. Ultimately, findings highlight the significance of these rapid and

specific POC diagnostics in improving the early detection and management of these

arboviral infections.

Introduction

Chikungunya (CHIKV), o'nyong-nyong (ONNV) and Mayaro (MAYV) viruses belong to the family *Togaviridae* and are closely related mosquito-borne alphaviruses[1]. Their genomes consist of approximately 12 kilobases of positive-sense RNA encased within a nucleocapsid, encoding the four nonstructural proteins (nsp1-4) and five structural proteins including three envelope proteins (E1-3), the capsid, and 6K across two open reading frames[2]. While evidence of alphavirus-induced human diseases dates back centuries, CHIKV, ONNV, and MAYV were formally isolated in the 1950s: CHIKV in Tanzania (1952), ONNV in Uganda (1959), and MAYV in Trinidad (1954)[3-5].

CHIKV is primarily transmitted by *Aedes* mosquitoes[1,3]. Following its isolation in 1952 during an outbreak in Tanzania, CHIKV caused several intermittent outbreaks from the 1960s-1980s in the Sub-Saharan Africa[6,7]. Meanwhile, the first record of CHIKV in Asia, was in Thailand in 1958, where it was found to be associated with dengue fever in a significant urban outbreak[6-8]. Several outbreaks in Southeast and South Asia were reported throughout the 1960s [6-8]. Only sporadic cases were reported in the years that followed, until its reemergence in 2004, on Lamu Island, in Kenya[6-9]. This particular outbreak caused a seroprevalence in 75% of the island's population had anti-CHIKV Immunoglobulins (IgM and IgG)I[9]. It subsequently spread to several nearby Islands in the Indian ocean [9]. Of these, the La reunion epidemic of 2005-2006, was particularly severe, with 266,000 cases and over 200 deaths resulting from the more severe disease manifestations, particularly among the older patient population[6,7,9]. Also part of this series of outbreaks was a major CHIKV outbreak in India, from 2005-2006, that resulted in over 1.4 million cases[6-8]. Around this time, outbreaks were also reported

in parts of Southeast Asia and also in some pacific islands[6-8]. Then, in 2011, CHIKV

reached the western hemisphere, and rapidly diffused through the Caribbean and finally

reached the South America in 2014, affecting hundreds of thousands of people along

the way [6-8]. Since then, there have been periodic CHIKV outbreaks in all of these

regions, as well as some autochthonous cases in parts of Europe, Middle East, and

North America[3,6,7]. Overall, according to the European Center for Disease Prevention

and Control, CHIKV lays claim to over 100 countries, and in 2024 alone, as of March

31, 160,000 new cases and 50 deaths have been reported[10].

ONNV, closely related to CHIKV, is limited in its endemicity within Africa,

although, reports of the virus reaching countries outside of Africa via international

travelers has been previously reported[11-13]. Transmitted by *Anopheles* mosquitoes,

ONNV has been implicated in two major outbreaks, that impacted millions of people[1].

This first outbreak, spanning 1959-1962, involved countries in both East and West

Africa, infecting over 2 million people[12,14]. The second outbreak between 1996-1997,

originated in Uganda and then spread to Kenya and Tanzania, with estimated infection

rates at 45-68%[1,12]. Following these, isolated cases and smaller outbreaks have been

reported across East and West Africa, as well as in travelers returning from those

regions[12].

Like ONNV, MAYV, is also geographically limited in its spread, causing sporadic

outbreaks in Central and South America, since it was first isolated in 1954 in

Trinidad[15,16]. Outbreaks have been recorded in Brazil, Bolivia, Venezuela, and Haiti[15,16].

Outside of these, wide ranging seroprevalence (21% to 72%) has been reported all

across the region, and sporadic cases occur both within the region and abroad, in

Europe and the USA through travelers[15,17]. Although primarily transmitted by

Hemagogous mosquitoes, certain strains of MAYV have the ability to adapt to replicate

in and be transmitted by different, more anthropophilic mosquito vectors[15,18,19]. This is

noteworthy because this ability had enabled the broader dissemination of related

viruses like CHIKV and suggests a possibility of global spread of MAYV[15,18,19].

Infection with these viruses typically results in a febrile illness characterized by

fever, mild to severe arthralgia, occasionally accompanied by edema, and/or a rash, but

can also lead to a debilitating, long-term arthritogenic syndrome[3]. In the case of CHIKV,

approximately 40% of infections progress to chronic arthritis, persisting for up to 3

months. Long-term manifestations have not been reported for ONNV, whereas, rare

chronic arthritis and encephalitic disease has been reported for MAYV[3,20]. Although

seldom fatal, the illnesses induced by these viruses substantially impact quality of life.

Given the millions of reported cases worldwide, CHIKV, ONNV, and MAYV pose a

significant global health risk[1,21].

The prevalence of these viruses is anticipated to escalate due to the influences

of climate change, urbanization, and international travel[21], necessitating improved

monitoring to prevent major outbreaks through effective, point-of-care (POC) diagnostic

tools. Diagnosis of CHIKV, ONNV, and MAYV infections primarily rely on laboratory

testing owing to clinical overlaps among these viruses with other arboviral diseases like

dengue and Zika fever [1,3,21]. Additionally, the geographical co-circulation of CHIKV and

ONNV in Africa and CHIKV and MAYV in South America, further complicates accurate

diagnosis[1,3,21]. Current diagnostic tests are predominantly either serological or

molecular techniques, based on the detection of patient antiviral immunoglobulins (IgM

and IgG) or viral RNA, respectively[1].

Commercially available serological tests, such as enzyme-linked immunosorbent

assays (ELISAs) detect anti-viral IgM and IgG[1]. IgM antibodies can usually be detected

in patient serum or plasma within 6 days after infection and persist for 3-4 months after

infection[1]. IgG antibodies emerge subsequent to IgM antibodies and can be detected

decades after the initial infection[1,22,23]. ELISAs, however, are reported to have low

sensitivity and lack specificity since antibodies tend to cross-react between the three

viruses due to their genetic similarity[1]. Plaque reduction neutralization tests (PRNTs)

are typically considered gold standard assays to distinguish viral infections based on

neutralizing antibodies but these tests often require enhanced biosafety laboratories

and require expert handling[24]. Moreover, there is evidence suggesting potential cross-

reactivity between CHIKV and ONNV in PRNTs as well[1,25-27].

Molecular diagnostics, on the other hand, are more sensitive and specific. Of

these, reverse transcription (RT)-PCRs or qRT-PCRs are the most popular[1,21]. There

are several commercial and/or in-house RT-PCR tests for CHIKV, ONNV, and MAYV.

These assays are reported to be sensitive and specific and can reliably detect viruses

particularly during the acute phase of the infection (the first 3-4 days), given the high

viremia (10^4-10^8 RNA copies/mL in the blood) during that time[1]. However, these assays

fall short of being ideal POC diagnostics, especially in more resource-limited settings,

due to their reliance on expensive and high precision equipment such as thermal

cyclers.

Therefore, rapid, inexpensive, sensitive, and specific POC diagnostics are still an

unmet need with these three viruses. One possible solution is using isothermal

amplification methods to detect viral RNA. These techniques have emerged as faster

and less expensive alternatives to PCR, particularly because they do not require precise

thermal cycling, and have shorter reaction times[28]. Furthermore, they possess

comparable to potentially enhanced sensitivity and specificity, and there are numerous

methods to detect their final products, several of which can be accomplished without

scanners or fluorescence readers[21,28]. Of these, recombinase polymerase amplification

(RPA) is of interest due to its non-complicated assay design, shorter reaction running

times, and its potential to be multiplexed[28].

RPA was first described in 2006 and it works by using a single pair of single-

stranded DNA primers, recombinase proteins (T4 Uvs X and UvsY), single-strand

binding proteins (Gp32), and a strand displacing polymerase (B. *subtilis* polymerase

large fragment), to isothermally amplify a double-stranded DNA template in the

presence of ATP and magnesium acetate[29]. Reactions are typically run at 37°C-42°C

for 5-20 minutes[28]. It can be used in the detection of RNA with the addition of a reverse

transcription step, either as a separate step or in a one-step reaction[28,30-32].

Furthermore, the assay can be configured to comply with several detection modalities,

including agarose gel electrophoresis, colorimetric readouts, and rapid test/lateral flow

assay (LFA) strips [28-33].

There currently are not many RPA tests for the detection of these three

alphaviruses[21], with the only example being a CHIKV RPA test designed by Patel et al.,

which was sensitive but cross-reactive with ONNV[34]. In this study, we report the

development of three virus-specific and sensitive RPA-based diagnostic tests for

CHIKV, ONNV, and MAYV. We initially performed multiple sequence alignments

(MSAs) to identify genomic targets for primer design, followed by an extensive primer

screen, and then finally re-configuring the successful primer pairs to make the assay

compatible with LFA-based detection. The assays target E2 for CHIKV, and NSP2 for

both MAYV and ONNV and all three tests can detect down to 10 viral copies on LFA

strips. The RPA-based rapid tests were further evaluated on serum and tissue samples

from clinically relevant mouse infection models for each virus. Ultimately, our results

highlight the sensitive and specific nature of RT-RPA-based diagnostics and allude to

their value as POC diagnostic tool better characterize these viruses and monitor

outbreaks.

Methods

Selecting regions for primer/probe design

MSAs were constructed using NCBI BLAST by first querying a complete genome

for each virus (NC_004162.2 for CHIKV, NC_075006.1 for ONNV, NC_003417.1 for

MAYV). The search results were filtered down to obtain a set of complete genomes with

over 90% similarity to the query sequence. Those sequences were aligned using

BLAST's MSA tool and viewed using the NCBI MSA viewer 1.25. The MSA viewer was

used to generate a consensus sequence with a 70% threshold, i.e., if >=70% of the

aligned sequences have the same nucleotide at the same position in the alignment, the

consensus sequence will reflect that, and, if there is less than 70% consensus at a

particular position, an ambiguous IUPAC symbol is used instead. The positions with a

less than 70% consensus (i.e., with an ambiguous IUPAC symbol in the consensus

sequence) were considered "mismatches" while calculating the % non-homology for

each gene of interest. Finally, the % mismatch values for the various genes were

compared and the top 4 most conserved, i.e., with the lowest % mismatch values were

selected for primer design. A list of all the aligned accession numbers is provided in

Supplementary Table 1.

Primer and Probe design

Genomes, NC_004162.2, NC_075006.1 and NC_003417.1 for CHIKV, ONNV

and MAYV, were respectively used to design primers and probes, within the genomic

regions of interest identified through the MSAs, for each virus. Primers were designed

manually, ensuring rules: a length of 30-35 nt, a GC content of 40-60% and absence of

palindromes or tandem nucleotide repeats[30,33,35]. The target amplicons were all between

120-200 bp in size and also had a GC content of 40%-60%. Finally, the designed

primers were mapped onto another MSA of the complete genomes of the three viruses

to ensure that sufficient mismatch was present to prevent cross reactivity[35]. After the

primer screen, probes were designed corresponding to the primer pair identified in the

screen for each virus. Nfo probe for the ONNV assay was designed as described by

Piepenburg et al., ensuring a complete length of about 45 nt, with at least 30 nt

preceding the abasic site, and 15 nt following it[29]. The hybridization probes for the

CHIKV and MAYV assays were designed as described by Qian et al., to be around 15-

25 nt in size, and binding to a region within the amplicon, avoiding overlap with the

primer binding sites[31]. The designed primers were synthesized by Integrated DNA

Technologies. A list of the RPA primers and probes designed and used in this study are

provided in supplementary Table 2.

RNA extraction

The QiaAmp viral Mini kit (Qiagen) or the PureLink Virl RNA/DNA Mini kit

(ThermoFisher) were used for extraction of RNA from serum or culture supernatant for

ONNV (NR-51661, BEI Resources Repository), following the manufacturers'

instructions. RNA from mouse tissues was extracted using the RNeasy Lipid tissue Mini

Kit (Qiagen) following manufacturer instructions. The concentration of the RNA extracts

was determined using a NanoDrop One device (ThermoFisher).

cDNA synthesis

Viral gRNA for CHIKV (181/25, BEI Resources Repository) and for MAYV (NR-

50081, BEI Resources Repository) and extracted RNA from ONNV (NR-51661, BEI

Resources Repository) propagated on Vero cells (CCL-81, ATCC) was used. Viral RNA

was converted to cDNA using the SuperScript IV Vilo master mix (ThermoFisher),

following the manufacturer's instructions. RNA input of $2x10^7$ viral copies was used in a

20 μL cDNA synthesis reaction to obtain cDNA with a concentration of 10^6 copies/μL.

For the primer screen, the cDNA stock was diluted to 10^5 copies/μL and 1 μL was used

for the RPA reactions. For the sensitivity assays, the cDNA stock was serially diluted to

obtain a range of concentrations, to enable inputs ranging from 10^6 copies to 1. For the

serum and tissue samples, a fixed volume of RNA (2 μL) was used as input into cDNA

synthesis, and 1μL of cDNA was used as input into RPA.

RT-RPA reactions using unlabeled primers

RPA reactions were performed using the TwistAmp Basic Lyophilized kit (TwistDx), following the manufacturer's instructions, but scaling measurements down to 10 µL reactions. Briefly, lyophilized mix was rehydrated using 29.5 µL of the provided rehydration buffer and 10 µL of nuclease free water, and 7.9 µL of the rehydrated mix was used per 10µL reaction, along with 0.3 µM of each primer, 280 mM magnesium acetate, and the requisite amount of cDNA. The reactions were incubated for 20 minutes at either 37°C, 39°C or 41°C, as indicated in the text. Diluted reaction products (1:5) were run on a 2.5% agarose (BioRad) gel in 1x TAE buffer (ThermoScientific) and stained with Gelgreen (Biotium). For the sensitivity assay gels, RPA products were purified using the PureLink PCR purification kit (ThermoFisher), following the manufacturer's instructions.

RT-RPA reactions with labelled primers

For the hybridization method, reactions were set up as described above with one modification—the replacement of the reverse primer with a 5'biotin- or 5'digitonin-labelled version, keeping the concentration the same. The hybridization step was performed as described by Qian et al.[31]. Briefly, after the RPA reaction run, 1 µL of 5 µM hybridization probe (final concentration of 0.167 µM) and 19 µL of hybridization buffer (10 mM Tris HCl at pH 8.5) was added to each reaction. This was heated to 95°C for 3 minutes and then allowed to gradually cool to room temperature to enable hybridization. The nfo-probe reactions were performed as described by Piepenburg et al[29]. Briefly, the basic master mix was modified, replacing a portion of the water with nfo or Endonuclease IV (New England Biosciences) at a final concentration of 0.2 U/µL and

nfo-probe to a final conc of 0.1 µM. The reverse primer was also replaced with a

5'digitonin-labelled version, keeping the concentration the same.

Detection of RPA products on LFA strips

The products from the hybridization and nfo-probe reactions were processed

using the HybriDetect 2T LFA strips (Milenia Biosciences), following the manufacturer's

instructions. Briefly, unpurified reaction products were directly diluted in the provided

assay buffer (1:20 for the nfo-probe reactions and 1:15 for hybridization reactions) and

applied to the strip. The strip was then immersed in 70 µL of the provided assay buffer

and was allowed to react for 5-30 minutes. The strips were imaged using a smartphone

camera.

qRT-PCR

For the CHIKV, ONNV and MAYV qPCRs conducted for this study, the

PrimeTime one-step qRT-PCR kit (Integrated DNA Technologies), along with primers

and TaqMan probes were used. CT values were recorded using a QuantStudio3 device

(ThermoFisher). The primer and probe sequences and the publications they are derived

from are as follows:

1) CHIKV- 5'AAGCTCCGCGTCCTTTACCAAG 3', 5'CCAAATTGTCCGGGTCCTCCT

3', 5'/FAM/CCATGTCTTCAGCCTGGACACCTTT/TAMRA/3'[36].

2) ONNV- 5'CGCAGCTTACGGGTTTCATA3', 5'GCAACGCCTTCAGAAACGC3',

5'/FAM/TGCTCTACTCTGCATTGCAAGA/BHQ1/3'[37].

3) MAYV- 5'AAGCTCTTCCTCTGCATTGC3', 5'TGCTGGAAACGCTCTCTGTA3', 5'/

CalFluorOrange560/GCCGAGAGCCCGTTTTTAAAATCAC/ BHQ-1/3'[38].

Preparation of Mouse Samples

Adult AG129 mice were inoculated with 1000 PFU of either CHIKV (181/25

strain, BEI Resources), ONNV (NR-50081, BEI Resources), or MAYV (NR-51661, BEI

Resources) via subcutaneous injection on the ventral side of the left footpads.

Submandibular blood was collected was collected at day 1 and/or day 3 post infection,

and into heparin coated tubes and centrifuged at 2000 x g for 10 minutes, to obtain

serum. The mice succumbed to the infections at day 3 (CHIKV and ONNV) and at day 5

(MAYV) at which point, the liver, spleen, kidneys, and cardiac blood were collected.

These samples were subjected to RNA extraction and RT-RPA as described above.

Results

In order to obtain primer pairs and probes capable of detecting RNA from all

strains of a given virus, we wanted to target them to the most conserved regions of that

virus' genome. To identify such targets for primer design, we used BLAST to construct

MSAs of a set of complete genomes for each virus. We then analyzed the consensus

sequence generated from the MSA to identify the most conserved genomic regions,

focusing our analysis on the 4 NSPs, capsid, E1 and E2. The two remaining structural

proteins—6K, E3—were overlooked due to their smaller size and therefore less area

within which to design a set of primers. The top 4 most conserved genomic regions

identified for each virus were as follows: NSP2, NSP4, E2 and E1 for CHIKV; NSP1,

NSP2, NSP4, E2 for ONNV; NSP1, NSP2, NSP4 and E1 for MAYV (Fig. 1A-C). Primers

were designed within the identified genes for each virus, with one exception: for ONNV,

we opted to design primers within E1 rather than NSP4. This was due to its high degree

of homology, with less than 40% sequence divergence within the Semliki Forest

Serocomplex, the antigenic complex to which the CHIKV, ONNV, and MAYV belong

and also due to E1's popularity as a target in existing molecular assays for these

viruses[1,39].

16 primer pairs were designed for ONNV and 17 for CHIKV and MAYV, spanning

the genomic regions identified through the MSAs. It was ensured that the primers

followed the stipulations for RPA primer design: a length of 30-35 nt, a GC content of

40-60% and absence of palindromes or tandem nucleotide repeats[30,33,35]. Additionally,

all designed primers were mapped onto the genomes of the other two viruses, to ensure

sufficient mismatches were present to minimize chances of cross-reactivity[35]. To screen

the designed primers, viral genomic RNA (gRNA) was first reverse transcribed and then

an equivalent of 10^5 viral copies in complementary DNA (cDNA) was used as input into

each RPA reaction. The reactions were run for 20 minutes, at three temperatures 37°C,

39°C and 41°C, which span the typical RPA temperature range[35]. They were evaluated

by running the products out on an agarose gel (Fig. 2A-C). Each primer pair varied in its

ability to amplify cDNA, and this activity sometimes influenced by temperature, for

example: at 37°C, CHIKV primers #7, 9, 11 fared better than #3, 6, 10, and #7 fared

better at 37°C than at 39°C (Fig. 2A). At the end of the screen, several successful

primer-pair/reaction-temperature combinations were identified, and the following

combinations were selected: #9 at 37°C was identified for CHIKV, #2 at 41°C for ONNV

and #1 at 39°C for MAYV. Their respective targets are within the E2 gene for CHIKV

and within NSP2 for both ONNV and MAYV. The analytical sensitivity of these three

primer-pairs was evaluated by using a serial dilution of cDNA, ranging from 10^6 viral

copies to 1 viral copy. Also included were reactions using 10^5 viral copies of the other

two viruses to evaluate cross-reactivity of the primers. The limits of detection (LODs) for

these three primers, processed via agarose gel electrophoresis, was found to be

approximately 10^2 viral copies for CHIKV and ONNV and between 10^3-10^2 viral copies

for MAYV, and no cross-reactivity to cDNA from the other two viruses was seen (Fig. 3).

Since we sought to design rapid tests for these three viruses, our next step was

to re-configure our assays to enable the detection of the reaction products on LFA

strips. For this, two approaches were used: hybridization and nfo-probes. At the outset,

both approaches involve replacing one of the reaction primers with a 5'labelled version

to generate a singly labelled amplification product. The first method involves a separate

hybridization step, where the singly-labelled amplification product, is denatured and

then hybridized with a labelled probe[32,40]. The second approach involves introducing an

nfo-probe, which are labelled probes with an internal abasic site, into the reaction, along

with an endonuclease (nfo or Endo IV) that cleaves at that abasic site upon probe

hybridization[29]. Ultimately, both approaches result in an amplification product that is

doubly labelled—one label with affinity for test line on the strip, and the other, with

affinity for the antibody conjugated to the gold nanoparticles (Fig. 4A). The LFA strips

we used had two test-bands, one coated with streptavidin and the other with an anti-

digitonin antibody. The gold nanoparticles in the strip are conjugated to an anti-FITC

antibody. We therefore opted to label the reverse primers using either 5'biotin tag (for

CHIKV) or 5'Digitonin tag (for ONNV and MAYV) and labelled all the probes with a

5'FAM tag.

We chose the hybridization option for the CHIKV and MAYV tests, and the nfo-

probe option for ONNV. The reaction temperatures were maintained at the same values

that were identified in the agarose gel-based primer screen for the ONNV and MAYV

tests. However, for CHIKV, running the reaction at 39°C instead of 37°C resulted in

darker coloration of the test-band on the LFA strip, prompting us to modify the running

conditions accordingly. Sensitivity of the rapid test was determined in the same manner

as described for Figure 3, using a serial dilution of viral cDNA. All three tests were

highly sensitive, with the ability to detect down to 10 viral copies, and similar to their un-

labeled counterparts (Fig. 3), they did not cross-react with 10^5 cDNA copies from the

other two viruses (Fig. 4B-D). Furthermore, we conducted a broader cross-reactivity

screen for these three rapid tests using cDNA from other arboviruses and saw no cross

reactivity, enabling us to confirm that the designed assays were highly specific to their

intended targets (Supplementary Table 3).

Having designed sensitive and specific RT-RPA-based rapid tests for CHIKV,

ONNV, and MAYV, we then sought to evaluate their performance on clinically relevant

samples. For this, adult AG129 mice, which are homozygous interferon

alpha/beta/gamma receptor (IFNα/β/γR) knockouts, were inoculated with either CHIKV,

ONNV, or MAYV at a dose of 1000 PFU via subcutaneous footpad injection. RNA was

then extracted from serum collected 1- and 3-days post-infection (dpi) and from the

spleens, livers, and kidneys. The extracted RNA was subjected to the designed RT-

RPA tests and also to RT-PCR for comparison (Fig. 5A-C). The CHIKV test was able to

detect virus in 2 out of 3 of the 1-dpi serum samples, 3-dpi serum samples, and livers.

But all three kidney and spleen samples produced a positive test. The CHIKV RT-PCR

test in comparison only detected viral RNA in two samples: one of the 3-dpi serum

samples and in one of the livers (Fig. 5A). The ONNV rapid test detected viral RNA in 1

out of 2 of the 3-dpi serum samples, spleens, and kidneys and both liver samples (Fig.

5B). Several of the RT-RPA-negative ONNV samples did however test positive with RT-

PCR. All but one of the MAYV samples tested positive on both RT-RPA and RT-PCR,

with the only negative result coming from one of the 1-dpi serum samples (Fig. 5C).

Discussion

In this study we describe three RT-RPA based rapid tests for the sensitive and

specific detection of CHIKV, ONNV and MAYV. The three viruses are closely related

alphaviruses, belonging to the same serocomplex, with ONNV and CHIKV being the

closest related viruses in that group[39]. There is currently a lack of sensitive and specific

POC diagnostic tools for these three viruses, given the cross reactivity of common

serological tests, and the expensive cost of running common molecular diagnostic tests

like RT-PCRs[1,21]. We sought to bridge that gap by designing RT-RPA based rapid tests,

that utilize the isothermal amplification technique, RPA, and are detected on rapid test

strips, making them ideal for use at the POC. We show that our three tests were

sensitive, with the ability to detect down to 10 viral copies, and specific, with no cross

reactivity to related alphaviruses or other arboviruses.

We further demonstrated that our tests are capable of detecting viral RNA from

more clinically relevant samples, starting with those obtained from mouse infection

models. Mice deficient in interferon signaling, like AG129 mice, are commonly used to

model CHIKV, ONNV, and MAYV infections because wildtype mice are not as

susceptible and are able to rapidly clear infections from these viruses[41,42]. In our mouse

experiments, we were able to detect viral RNA in the day 3 sera more reliably than in

the day 1 sera, consistent with the trend of viremia peaking around day 2-3 for arboviral

infections in this mouse model[42]. We were also able to detect viral RNA in the spleens

and livers, which, along with joints and muscles, are considered common replication

sites for alphaviruses in their hosts, allowing for their efficient dissemination

thereafter[41,43-45]. We were also able to detect viral RNA in the kidneys of infected mice,

and while they not a common target or replication site for alphavirus infections, viral

loads have been detected in the kidneys of AG129 mice infected with100 PFU of

CHIKV[46]. Additionally, we were only able to produce faint positive test bands on our

rapid-test strips for most of these samples. One possible explanation for this could be

that these mice succumbed to the infections quite rapidly, and possibly before the

viruses had a chance to replicate to produce robust viral loads. It is worth noting that the

3 CHIKV infected mice and the 2 ONNV infected mice experienced mortality at day 3,

and the 2 MAYV infected mice succumbed to the infection at day 5. This is consistent

with what with what is known about the behavior of CHIKV, ONNV, and MAYV

infections in AG129, A129, or similar mouse models[36,41,42,45,47-51].

One improvement that would further bolster these assays would be optimizing

them to function as one-step RT-RPA reactions. Secondly, we had designed the MAYV

and ONNV assays to produce products that have a different (digitonin) tag than the

CHIKV assay products (which have a biotin tag) because we had initially aimed at

multiplexing their detection on our double test-band LFA strips. Further optimization to

enable this would allow for the creation of two alphavirus tests, one for the South

American viruses-CHIKV and MAYV, and the other for the African viruses-CHIKV and

ONNV. Lastly, being able to detect RNA directly from crude biological samples such as

plasma/serum, urine, or saliva, without an RNA extraction step, would truly enable field

deployment. This is possible and has been attempted before, because compared to

PCR, RPA is less susceptible to inhibitors present in crude samples[33,52].

Despite their spread and disease severity, approved vaccines and antivirals for

these three alphaviruses are limited, with the first approved CHIKV vaccine being as

recent as November 2023[53]. As these efforts ramp up with the increasing recognition of

the threats posed by these viruses, effective POC diagnostic tools will be critical for the

efficient monitoring of outbreaks, and for the dissemination of prophylactics and

therapeutics. Given their potential as sensitive, specific, POC diagnostics, we think that

the RT-RPA assays described here are a promising step forward towards that goal.

Data Availability

The data that support the findings of this study are available within the

manuscript or upon request from the corresponding authors.

Acknowledgments

425   We would like to thank Rutgers Global Health Institute, Rutgers Robert Wood

426 Johnson Medical School, and the Child Health Institute of New Jersey for their support.

Conflicts of Interest

429   BBH is a co-founder of Mir Biosciences, Inc., a biotechnology company focused

430 on T cell-based diagnostics and vaccines for infectious diseases, cancer, and

431 autoimmunity.

References

1.   Pezzi L, Reusken C, Weaver S, C., et al. GloPID-R report on Chikungunya, O'nyong-

nyong and Mayaro virus, part I: Biological diagnostics. *Antiviral Research*.

2019;doi:10.1016/j.antiviral.2019.03.009

2.   Kim A, S. , Diamond M, S. . A molecular understanding of alphavirus entry and

antibody protection. *Nature Reviews Microbiology*. 2022;doi:10.1038/s41579-022-00825-7

3.   Zaid A, Burt F, J., Liu X, et al. Arthritogenic alphaviruses: epidemiological and

clinical perspective on emerging arboviruses. *Lancet Infectious Diseases*.

2020;doi:10.1016/s1473-3099(20)30491-6

4.   Diagne C, Tidiane, Bengue M, Choumet V, Hamel R, Pompon J, Misse D. Mayaro

Virus Pathogenesis and Transmission Mechanisms. *Pathogenetics*.

2020;doi:10.3390/pathogens9090738

5.   Rezza G, Chen R, Weaver S, C. O'nyong-nyong fever: a neglected mosquito-borne

viral disease. *Pathogens and Global Health*. 2017;doi:10.1080/20477724.2017.1355431

6.   Wahid B, Ali A, Rafique S, Idrees M. Global expansion of chikungunya virus:

mapping the 64-year history. *International Journal of Infectious Diseases*. 2017/05/01/

2017;58:69-76. doi:https://doi.org/10.1016/j.ijid.2017.03.006

7.   Zeller H, Van Bortel W, Sudre B. Chikungunya: Its History in Africa and Asia and Its

Spread to New Regions in 2013–2014. *The Journal of Infectious Diseases*.

2016;214(suppl_5):S436-S440. doi:10.1093/infdis/jiw391

8.   Manzoor KN, Javed F, Ejaz M, et al. The global emergence of Chikungunya infection:

An integrated view. *Reviews in medical virology*. 2022;(1099-1654 (Electronic))

9.   Renault P, Balleydier E, D'Ortenzio E, Bâville M, Filleul L. Epidemiology of

chikungunya infection on Reunion Island, Mayotte, and neighboring countries. *Médecine et*

Maladies Infectieuses. 2012/03/01/ 2012;42(3):93-101.

doi:https://doi.org/10.1016/j.medmal.2011.12.002

10.   Chikungunya worldwide overview. online. European Centre for Disease Prevention

and Control. Accessed 5/14/2024, 2024. https://www.ecdc.europa.eu/en/chikungunya-

monthly

11.   Pezzi L, LaBeaud A, D., Reusken C, B., et al. GloPID-R report on chikungunya,

o'nyong-nyong and Mayaro virus, part 2: Epidemiological distribution of o'nyong-nyong

virus. *Antiviral Research*. 2019/12/01/ 2019;172:104611.

doi:https://doi.org/10.1016/j.antiviral.2019.104611

12.   Rezza G, Chen R, Weaver SC. O'nyong-nyong fever: a neglected mosquito-borne

viral disease. *Pathogens and global health*. 2017;(2047-7732 (Electronic))

13.   Tappe D, Kapaun A, Emmerich P, et al. O'nyong-nyong virus infection imported to

Europe from Kenya by a traveler. *Emerg Infect Dis*. Oct 2014;20(10):1766-7.

doi:10.3201/eid2010.140823

14.   Powers A, M., Brault A, C., Tesh R, B., Weaver S, C. Re-emergence of Chikungunya

and O'nyong-nyong viruses: evidence for distinct geographical lineages and distant

evolutionary relationships. *The Journal of general virology*. 2000;(0022-1317 (Print))

15.   Caicedo E-Y, Charniga K, Rueda A, et al. The epidemiology of Mayaro virus in the

Americas: A systematic review and key parameter estimates for outbreak modelling. *PLoS*

Neglected Tropical Diseases. 2021;(1935-2735 (Electronic))

16.   Pezzi L, Rodriguez-Morales A, J., Reusken C, et al. GloPID-R report on chikungunya,

o'nyong-nyong and Mayaro virus, part 3: Epidemiological distribution of Mayaro virus.

Antiviral Research. 2019;doi:10.1016/j.antiviral.2019.104611

17.   Silva-Ramos C, Ramiro, Mejorano-Fonseca J, A., Hidalgo M, Rodríguez-Morales A,

J., Faccini-Martínez Á, A. Clinical, Epidemiological, and Laboratory Features of Mayaro

Virus Infection: a Systematic Review. *Current Tropical Medicine Reports*. 2023/12/01

2023;10(4):309-319. doi:10.1007/s40475-023-00308-6

18.   Santos F, Martins, Dias R, Sousa, Fernandes L, de Souza, da Silva C, Canedo, de

Paula S, Oliveira. Mayaro Virus Infection: Clinical Features and Global Threat. *Current*

Treatment Options in Infectious Diseases. 2020/12/01 2020;12(4):387-397.

doi:10.1007/s40506-020-00240-7

19.   Mota M, Tasso de Oliveira, Avilla C, M., S., Nogueira M, Lacerda. Mayaro virus: a

neglected threat could cause the next worldwide viral epidemic. *Future Virology*.

2019;doi:10.2217/fvl-2019-0051

20.   Amaral JK, Bilsborrow J, B. , Schoen R, T. Chronic Chikungunya Arthritis and

Rheumatoid Arthritis: What They Have in Common. *The American Journal of Medicine*.

2020;doi:10.1016/j.amjmed.2019.10.005

21.   Mori A, Pomari E, Deiana M, et al. Molecular techniques for the genomic viral RNA

detection of West Nile, Dengue, Zika and Chikungunya arboviruses: a narrative review.

Expert Review of Molecular Diagnostics. 2021;doi:10.1080/14737159.2021.1924059

22.   Chusri S, Siripaitoon P, Silpapojakul K, et al. Kinetics of Chikungunya Infections

during an Outbreak in Southern Thailand, 2008-2009. *American Journal of Tropical*

Medicine and Hygiene. 2014;doi:10.4269/ajtmh.12-0681

23.   Nitatpattana N, Kanjanopas K, Yoksan S, et al. Long-term persistence of

Chikungunya virus neutralizing antibodies in human populations of North Eastern Thailand.

Virology Journal. 2014;doi:10.1186/1743-422x-11-183

24.   Natrajan M, S. , Rojas A, Waggoner J, J. . Beyond Fever and Pain: Diagnostic

Methods for Chikungunya Virus. *Journal of Clinical Microbiology*.

2019;doi:10.1128/jcm.00350-19

25.   Lim J, Kyungah, Ridde V, Agnandji S, T., et al. Seroepidemiological reconstruction of

long-term chikungunya virus circulation in Burkina Faso and Gabon. *The Journal of*

Infectious Diseases. // 2023;doi:10.1093/infdis/jiac246

26.   Hozé N, Diarra I, Sangaré AK, et al. Model-based assessment of Chikungunya and

O'nyong-nyong virus circulation in Mali in a serological cross-reactivity context. *Nature*

Communications. 2021;(2041-1723 (Electronic))

27.   LaBeaud AD, Banda T, Brichard J, et al. High rates of o'nyong nyong and

Chikungunya virus transmission in coastal Kenya. *PLOS Neglected Tropical Diseases*.

2015;(1935-2735 (Electronic))

28.   Tan M, Liao C, Liang L, et al. Recent advances in recombinase polymerase

amplification: Principle, advantages, disadvantages and applications. *Frontiers in Cellular*

and Infection Microbiology. 2022;doi:10.3389/fcimb.2022.1019071

29.   Piepenburg O, Williams C, H., Stemple D, L., Armes N, A. DNA Detection Using

Recombination Proteins. *PLOS Biology*. 2006;doi:10.1371/journal.pbio.0040204

30.   Shanshan Z, Mingyuan D, Shuang L, et al. Current status of recombinase

polymerase amplification technologies for the detection of pathogenic microorganisms.

Diagnostic microbiology and infectious disease.

2023;doi:10.1016/j.diagmicrobio.2023.116097

31.   Qian J, Boswell SA, Chidley C, et al. An enhanced isothermal amplification assay for

viral detection. *Nature Communications*. 2020;doi:10.1038/s41467-020-19258-y

32.   Salcedo N, Sena B, F., Qu X, Herrera BB. Comparative Evaluation of Rapid

Isothermal Amplification and Antigen Assays for Screening Testing of SARS-CoV-2. *Viruses*.

2022;doi:10.3390/v14030468

33.   Daher RK, Stewart G, Boissinot M, Bergeron MG. Recombinase Polymerase

Amplification for Diagnostic Applications. *Clinical Chemistry*.

2016;doi:10.1373/clinchem.2015.245829

34.   Patel P, Abd El Wahed A, Faye O, et al. A Field-Deployable Reverse Transcription

Recombinase Polymerase Amplification Assay for Rapid Detection of the Chikungunya

Virus. *PLOS Neglected Tropical Diseases*. 2016;doi:10.1371/journal.pntd.0004953

35.   Daher R, K. , Stewart G, Boissinot M, Boudreau DK, Bergeron MG. Influence of

sequence mismatches on the specificity of recombinase polymerase amplification

technology. *Molecular and Cellular Probes*. 2015;doi:10.1016/j.mcp.2014.11.005

36.   Schilte C, Couderc T, Chretien F, et al. Type I IFN controls chikungunya virus via its

action on nonhematopoietic cells. *The Journal of experimental medicine*. 2010;(1540-9538

(Electronic))

37.   Waggoner J, J., Heath C, J., Ndenga B, et al. Development of a Real-Time Reverse

Transcription Polymerase Chain Reaction for O'nyong-nyong Virus and Evaluation with

Clinical and Mosquito Specimens from Kenya. *The American journal of tropical medicine*

and hygiene. 2017;(1476-1645 (Electronic))

38.   Waggoner J, J., Rojas A, Mohamed-Hadley A, de Guillén Y, Arévalo, Pinsky B, A. Real-

time RT-PCR for Mayaro virus detection in plasma and urine. *Journal of Clinical Virology*.

2018/01/01/ 2018;98:1-4. doi:https://doi.org/10.1016/j.jcv.2017.11.006

39.   Powers A, M. , Brault A, C. , Shirako Y, et al. Evolutionary Relationships and

Systematics of the Alphaviruses. *Journal of Virology*. 2001;doi:10.1128/jvi.75.21.10118-

10131.2001

40.   Mancuso C, P., Lu Z, Qian Z, Boswell SA, Springer M. A Semi-Quantitative

Isothermal Diagnostic Assay Utilizing Competitive Amplification. *Analytical Chemistry*.

2021;doi:10.1021/acs.analchem.1c01576

41.   Couderc T, Chrétien F, Schilte C, et al. A Mouse Model for Chikungunya: Young Age

and Inefficient Type-I Interferon Signaling Are Risk Factors for Severe Disease. *PLOS*

Pathogens. 2008;4(2):e29. doi:10.1371/journal.ppat.0040029

42.   Baldon LVR, de Mendonça SF, Ferreira FV, et al. AG129 Mice as a Comprehensive

Model for the Experimental Assessment of Mosquito Vector Competence for Arboviruses.

LID - 10.3390/pathogens11080879 [doi] LID - 879. *Pathogens*. 2022;(2076-0817 (Print))

43.   Assunção-Miranda I, Cruz-Oliveira C, Da Poian A, T. Molecular mechanisms

involved in the pathogenesis of alphavirus-induced arthritis. *BioMed research*

international. 2013;(2314-6141 (Electronic))

44.   Kafai N, M., Diamond M, S. , Fox J, M. Distinct Cellular Tropism and Immune

Responses to Alphavirus Infection. *Annual Review of Immunology*.

2022;doi:10.1146/annurev-immunol-101220-014952

45.   Marín-López A, Calvo-Pinilla E, Moreno S, et al. Modeling Arboviral Infection in Mice

Lacking the Interferon Alpha/Beta Receptor. *Viruses*. 2019;doi:10.3390/v11010035

46.   Ngwe Tun MM, Muthugala R, Kyaw Kyaw A, Shimada S, Morita K, Hayasaka D.

Pathogenetic Potential Relating to Metabolic Activity in a Mouse Model of Infection with the

Chikungunya Virus East/Central/South African Genotype. *Viruses*. 2020;12(2).

doi:10.3390/v12020169

47.   Gary W, Xiangguo Q. Type I interferon receptor knockout mice as models for

infection of highly pathogenic viruses with outbreak potential. *Zoological Research*.

2018;doi:10.24272/j.issn.2095-8137.2017.052

48.   Seymour R, L. , Rossi S, L., Bergren N, A. , Plante K, S. , Weaver S, C. . The Role of

Innate versus Adaptive Immune Responses in a Mouse Model of O'Nyong-Nyong Virus

Infection. *American Journal of Tropical Medicine and Hygiene*. 2013;doi:10.4269/ajtmh.12-

0674

49.   Webb E, M., Azar S, R., Haller S, L., et al. Effects of Chikungunya virus immunity on

Mayaro virus disease and epidemic potential. *Scientific Reports*. 2019/12/31

2019;9(1):20399. doi:10.1038/s41598-019-56551-3

50.   Figueiredo CM, Neris R, Gavino-Leopoldino D, et al. Mayaro Virus Replication

Restriction and Induction of Muscular Inflammation in Mice Are Dependent on Age, Type-I

Interferon Response, and Adaptive Immunity. *Frontiers in microbiology*. 2019;(1664-302X

(Print))

51.   Partidos C, D., Paykel J, Weger J, et al. Cross-protective immunity against o'nyong-

nyong virus afforded by a novel recombinant chikungunya vaccine. *Vaccine*. 2012;(1873-

2518 (Electronic))

52.   Vasileva Wand N, I., Bonney L, C., Watson R, J., Graham V, Hewson R. Point-of-care

diagnostic assay for the detection of Zika virus using the recombinase polymerase

amplification method. *Journal of General Virology*. 2018;(1465-2099 (Electronic))

53.   Chikungunya vaccine approved. *Nature Biotechnology*. 2023/12/01

2023;41(12):1667-1667. doi:10.1038/s41587-023-02071-5

Figures

Figure 1: Schematic depicting the regions selected for primer design. Multiple

sequence alignments (MSAs) were assembled for (A) CHIKV, (B) ONNV and (C) MAYV

using the indicated number of sequences following which the consensus sequence was

analyzed focusing on the four NSPs, the Capsid, E2 and E1 to identify the most

conserved regions for each virus. The heatmaps show the percent non-homology

calculated for each region and the top four most homologous sites (with the darkest

shading), highlighted in a red border, were selected for primer design. For ONNV, the

analysis identified NSP4, highlighted with a discontinuous border, was identified as the

fourth target for primer design, however, we chose to prioritize designing primers in E1

instead.

Figure 2: Primer screens conducted by agarose gel electrophoresis of RPA products. All 17 primers for CHIKV, 16 primers for ONNV and 17 primes for MAYV were screened in 20-minute RPA reactions using 10^5 viral copies with cDNA as input, The reactions were run at three temperatures as indicted. Depicted here is a representative subset of those screened primers, A) corresponding to CHIKV, (B) to ONNV, and (C) to MAYV.

Figure 3: Analytical sensitivities of the designed primers on an agarose gel.

Candidate (A) CHIKV, (B) ONNV, and (C) MAYV primer pairs identified in the screen were evaluated on a serial dilution of cDNA inputs ranging from 10^6 viral copes to 1 viral copy. Cross reactivity of these primers was also tested using 10^5 copies of cDNA from the other two viruses. The reaction inputs for each well are denoted in the image. The sensitivity was determined to be 10^2 for CHIKV and ONNV, and between 10^3 and 10^2 for MAYV, with not cross-reactivity.

Figure 4: Detecting RPA reactions on LFA strips. A) Schematic representation of the

Millenia Biosciences 2T test strips utilized in this study. The strips have two test bands

"T1" and "T2," one coated with streptavidin and the other with an anti-Digitonin antibody.

The gold nanoparticles (AuNPs) are conjugated to an anti-FITC antibody. Accordingly,

all our reverse primers were either tagged with a 5'Biotin tag (CHIKV) or a 5' Digitonin

tag (ONNV and MAYV), and their probes- hybridization probe for CHIKV and MAYV or

nfo-probe for ONNV- were all 5'FAM tagged. Shown in the figure are depictions of the

reaction product from the CHIKV assay, that uses a hybridization probe, binding at T1

and the ONNV assay, that uses a nfo-probe, binding at T2. The MAYV RPA product

would appear like the CHIKV product since it also uses hybridization probes, however, it

would bind at T2, since its reverse primer has a 5'digitonin tag. (B-D) show the

analytical sensitivity of the rapid-test versions of the CHIKV (B), ONNV, (C) and MAYV

(D) assays. The reactions used a serial dilution of cDNA as their input. Also included

are reactions with 10^5 copies of cDNA from the other two viruses to assess cross-

reactivity. RPA reaction products were diluted and applied to the strips and then

immersed in the manufacturer-provided assay buffer for up to 30 minutes. The strips

were photographed using a smartphone camera. All three assays were able to detect

down to 10 copies of viral cDNA and were not cross-reactive.

Figure 5: Detecting alphaviruses in serum and tissues from mouse infection models using RT-RPA rapid tests. AG129 mice were infected with 1000 PFU of (A) CHIKV (181/25 strain, n=3), (B) ONNV (NR-50081 strain, n=2) and (C) MAYV (NR-51661 strain, n=2) via subcutaneous footpad injection. Serum was collected 1- and 3-days post infection, and organs were harvested at day 3 for CHIKV and ONNV and at day 5 for MAYV, when the mice experienced mortality. RNA was extracted, reverse-transcribed and subjected to RPA using labelled primers. The reaction products were applied to the test strips and were photographed using a smartphone camera. The extracted RNA from each sample was also subjected to RT-PCR for comparison. The

678   results of both assays for each sample are indicated with a + (or ++) and –, as

679   observed.