# Recombinant inbred line panels inform the genetic architecture and interactions of adaptive traits in *Drosophila melanogaster*

**Authors**: Tiago da Silva Ribeiro[1,2,a,@], Matthew J. Lollar[1], Quentin D. Sprengelmeyer[1b], Yuheng Huang[1c], Derek M. Benson[1], Megan S. Orr[1], Zachary C. Johnson[1], Russell B. Corbett-Detig[3,4], John E. Pool[1,2,@]

**Mailing address:**

[1] Laboratory of Genetics, University of Wisconsin-Madison, Madison, WI, 53706, USA

[2] Department of Integrative Biology, University of Wisconsin-Madison, Madison, WI, 53706, USA

[3] Genomics Institute, University of California, Santa Cruz, Santa Cruz, CA, 95064, USA

[4] Department of Biomolecular Engineering, University of California, Santa Cruz, Santa Cruz, CA, 95064, USA

[a] Current address: Instituto de Biociências, Universidade de São Paulo, São Paulo, Brazil, 05508-090.

[b] Current address: Department of Equity and Inclusion, University of Wisconsin-Platteville, Platteville, WI, 53818

[c] Current address: Department of Ecology and Evolutionary Biology, University of California, Irvine, Irvine, CA, 92697.

[@] Corresponding authors:

Tiago da Silva Ribeiro, tribeiro@wisc.edu

John E. Pool, jpool@wisc.edu

**Keywords**: epistasis, Quantitative Trait Loci (QTL), adaptation, quantitative genetics, *Drosophila melanogaster.*

## Abstract

The distribution of allelic effects on traits, along with their gene-by-gene and gene-by-environment interactions, contributes to the phenotypes available for selection and the trajectories of adaptive variants. Nonetheless, uncertainty persists regarding the effect sizes underlying adaptations and the importance of genetic interactions. Herein, we aimed to investigate the genetic architecture and the epistatic and environmental interactions involving loci that contribute to multiple adaptive traits using two new panels of *Drosophila melanogaster* recombinant inbred lines (RILs). To better fit our data, we re-implemented functions from R/qtl (Broman *et al.* 2003) using additive genetic models. We found 14 quantitative trait loci (QTL) underlying melanism, wing size, song pattern, and ethanol resistance. By combining our mapping results with population genetic statistics, we identified potential new genes related to these traits. None of the detected QTLs showed clear evidence of epistasis, and our power analysis indicated that we should have seen at least one significant interaction if sign epistasis or strong positive epistasis played a pervasive role in trait evolution. In contrast, we did find roles for gene-by-environment interactions involving pigmentation traits. Overall, our data suggest that the genetic architecture of adaptive traits often involves alleles of detectable effect, that strong epistasis does not always play a role in adaptation, and that environmental interactions can modulate the effect size of adaptive alleles.

## Introduction

Adaptive evolution is the result of natural selection, which acts by increasing the frequency of beneficial traits in a population. A number of important uncertainties persist regarding the genetic architectures of adaptive trait changes in nature, including the number and effect sizes of contributing variants, and their frequencies before and after the action of positive selection (Pritchard *et al.* 2010, Savoleinen *et al.* 2013, Barghi *et al.* 2020). Furthermore, research on the genetic basis of adaptation has largely been centered on the individual additive effects of each gene underlying the adaptive trait, whereas the role of gene-by-gene and gene-by-environment interactions in adaptation has remained more elusive (Whitlock *et al.* 1995, Malmberg & Mauricio 2005, Martin & Lenormand 2006, Bank 2022).

Theoretical and empirical studies have yielded varied findings concerning the number and strength of loci contributing to adaptive trait change. Fisher (1930) argued that traits were governed by numerous genetic variants of very small magnitude, and that natural selection proceeded via minute incremental changes in allele frequency at these loci. Extending Fisher's geometric model, Orr (1998) found that in a sequential model of genetic adaptation, a few larger effect sizes should tend to be followed by many smaller ones, yielding an exponential distribution of effects. However, adaptation in natural populations may sometimes depart from the assumptions of such models in multiple respects. First, the

role of standing variation (as opposed to new mutations) in the adaptive

process continues to be investigated (Fuhrmann *et al.* 2023, Schlötterer

2023), and these two sources of genetic variation may differ in their effect

sizes, dominance, and other properties (e.g. Orr & Betancourt 2001, Ségurel

& Bon 2017, Bomblies & Peichel 2022).  Second, the genetic architecture of

local adaptation, in which populations in a heterogeneous landscape adapt

to local conditions in the presence of migration, has a stronger bias towards

alleles of large effect sizes (Griswold 2006, Yeaman & Whitlock 2011,

Yeaman & Otto 2011).  Third, interactions among genes impacting

phenotypes (*i.e.* epistasis) may alter the dynamics of adaptive variants, as

further discussed below.

In regards to the role of epistasis in adaptation, a discussion has

existed since the onset of the Modern Synthesis, with some arguing that it is

largely irrelevant to the trajectory of Darwinian selection (Fisher 1930,

Cohan *et al.* 1989, Hill *et al.* 2008, Crow 2010) and others proposing that it

plays an important role (Fenster *et al.* 1997, Hansen 2013, Wright 1931).  The

above literature suggests that on the one hand, epistasis might not be

expected to strongly affect adaptation because alleles at unlinked loci can

segregate independently and selection would act on the combined effect of

each allele against different genetic backgrounds; and yet, epistasis has the

potential to alter the rate at which adaptive trait changes can occur by

producing larger or smaller effects in some of these backgrounds, and in

certain contexts, it may wield a stronger influence on the course of

adaptation (Cohan *et al.* 1989, Crow 2010, Fenster *et al.* 1997, Fisher 1930,

Hansen 2013, Hill *et al.* 2008, Wright 1931).

Epistasis has been shown to play a role in many evolutionary

processes, including the evolution of sex and recombination, speciation,

canalization, maintenance of genetic variation, and the evolution of co-

adapted gene complexes (Hansen 2013).  How epistatic interactions alter the

response to directional selection is determined by the nature of the epistatic

interaction (Hansen 2013).   Some kinds of epistasis affect the magnitude of

the phenotypic effects, such as positive and negative epistasis, and another

kind, called sign epistasis, changes the direction of the effect.  Multiple

theoretical and simulation studies have investigated how each of these

epistatic scenarios might influence the adaptive process (Carter *et al.* 2005;

Hansen *et al.* 2006; Yukilevich *et al.* 2008; Barton 2017).  For positive

(synergistic) epistasis, in which the combined effect of both alleles is greater

than the sum of the individual effects, additive genetic variance would

increase, and the mean fitness in an adapting population is expected to

change faster and reach a higher plateau than without epistasis.  For

negative epistasis, in which the combined effect is lower than the sum of the

individual effects but still qualitatively consistent with the additive effects,

additive genetic variance decreases and the mean fitness during adaptation

changes more slowly and reaches a lower plateau than without

epistasis.  Particularly in the context of adaptation, negative epistasis is also

referred to as diminishing returns epistasis.  For sign epistasis, whether the

allele will contribute to a larger or smaller phenotype will depend on the

allele at another locus.  In this case, the evolutionary dynamics can be more

complex, and the fate of one allele is attached to the other, but while both

variants persist, their interaction may impede adaptation. Furthermore, to

the extent that epistasis masks the phenotypic influence of some variants, it

may facilitate the accumulation of cryptic genetic variation for a given trait,

which may subsequently be exposed to selection when elements of the

genetic background (or the environment) change (Gibson & Dworkin 2004,

Steiner *et al.* 2007).

Despite the pervasiveness of epistasis at the genome-wide scale

(Huang *et al.* 2012), as well as between adaptive variants within proteins

(Lunzer *et al.* 2010, Gong & Bloom 2014, Starr & Thorthon 2016), the

prevalence of epistatic interactions among adaptive variants that contribute

to the evolution of more complex adaptive traits remains incompletely

understood.  Studies of crop domestication have reported examples of

epistasis among artificially selected alleles, including grapevine varieties

(Duchêne *et al.* 2012) and many examples in maize domestication (Stitzer &

Ross-Ibarra 2018).  Still, compared to wild alleles, domesticated alleles were

usually less sensitive to the genetic background (reviewed in Doust *et al.*

2014).  Epistasis was also found among adaptive quantitative trait loci

(QTLs) underlying flowering time in *Arabidopsis thaliana* (Juenger *et al.*

2002). In a recent genome-wide screen in yeast, using genes known to be

involved in a high number of interactions, 24% of the adaptive variants were

strain-specific and indicative of epistasis (Ang *et al.* 2023). Positive epistasis

has been shown to underlie the evolution of bacteria with multiple

antibiotic resistance alleles (Trindade *et al.* 2009). Examples in animals

involve epistasis between alleles underlying adaptive coloration in

butterflies (Papa *et al.* 2013), negative epistasis for traits underlying fitness in

populations of *Drosophila melanogaster* experimentally selected on a

chronic larval malnutrition regimen (Vijendravarma & Kawecki 2013), and

epistasis between genes underlying coat color in oldfield mice, in which the

effect of one allele is only expressed in the presence of another allele

(Steiner *et al.* 2007).

      The genetic architecture of adaptive traits can also be affected by

gene-by-environment interactions because the effect of a given allele can

change depending on its environment (Zeng *et al.* 1999, Mackay

2001). Transcriptome analysis of *D. melanogaster* reared at 20 different

environments have shown that expression of ~15% of genes changes

detectably with environment (Zhou et al. 2012). In some cases, gene-by-

environment interactions can be seen at the gene expression level but not

the metabolic level (Reed et al. 2014), highlighting the complexity of

interpreting its role in adaptation. Examples of interactions between the

environment and adaptive alleles include *D. melanogaster* reproductive

performance at different temperatures (Fry *et al.* 1998), drought stress

adaptation in wheat (Mathews *et al.* 2008), and local adaptation reflected in

biomass for switchgrass (Lowry *et al.* 2019). Similarly to epistatic

interaction, gene-by-environment interaction can also uncover cryptic

genetic variation (Gibson & Dworkin 2004). The chaperone protein Hsp90

offers an example of how genetic and environmental interactions can

uncover cryptic variation underlying discrete and continuous traits

(Rutherford & Lindquist 1998, McGuigan & Sgro 2009, Flynn *et al.* 2020).

Here, we contribute to the understanding of the genetic architecture

of adaptive traits and the emerging knowledge about genetic and

environmental interactions by identifying adaptive quantitative trait loci

(QTLs) underlying recent local adaptation in natural populations of *D.*

*melanogaster* and asking whether there is evidence of epistasis among them

and, for a subset, whether they interact with the environment. Using two

newly-described panels of recombinant inbred lines (RILs), we investigate

several traits that appear to have evolved directly or indirectly under

adaptive differentiation between an ancestral range population (Zambia)

and a population from either France or highland Ethiopia, regions

colonized by *D. melanogaster* approximately 2 kya (Sprengelmeyer *et al.*

2020).

The traits examined here (cold tolerance, ethanol resistance,

pigmentation, song pattern, and wing length) show evidence of being either

direct targets of local adaptation or else pleiotropic readouts of local

adaptation targeting correlated traits.  The pigmentation traits show strong

correlations with environmental variables, especially ultraviolet radiation

(Bastide *et al.* 2014), which reaches particularly high levels in the Ethiopian

highlands.  Larger wings may help Ethiopian flies navigate in cooler,

thinner air, and wing size was found to show unusually strong quantitative

trait differentiation ($Q_{ST}$ = 0.985; Lande 1992; Spitze 1993) between the

Ethiopia and Zambia populations, compared to genome-wide genetic

differentiation (Lack *et al.* 2016), indicating a role for selection in this trait's

differentiation.  France and Zambia have markedly different ethanol

resistance, also exceeding patterns of genome-wide genetic differentiation

(Sprengelmeyer *et al.* 2021).  Cold developmental survival shows a similarly

consistent differentiation between France and Zambia in particular (Huang

*et al.* 2021), with clear presumptive adaptive value in light of climate

differences between these regions.  A male song trait – the proportion of

pulses classified as slow (Clemens *et al.* 2018) – was included based on strong

population differentiation, with a $Q_{ST}$ of 0.797 exceeding the $F_{ST}$ of >96% of

non-singleton SNPs (Lollar *et al.* 2025).  Although directional selection has

not been previously reported to have acted directly on song traits within *D.*

*melanogaster*, it is also possible that this trait difference reflects selection on

a pleiotropically connected trait, such as synaptic function in a novel

thermal environment (Pool *et al.* 2017).

We also note that for the pigmentation, size, and ethanol resistance

traits examined here, bulk mapping studies found that all of these

population trait changes had very different genetic architectures from one

mapping cross to the next (Bastide *et al.* 2016; Sprengelmeyer *et al.* 2021;

Sprengelmeyer *et al.* 2022).  Given that the differences included QTLs with

effect size high enough to be detected across all the mapping crosses (Pool

2016), these results imply considerable segregating variation in genetic basis

of each trait difference within one or both of the populations studied.

However, it is possible (though as yet untested) that epistasis may also play a

role in yielding distinct QTLs in different mapping crosses for these traits.

For each of the above traits, we first identify QTLs and candidate

genes that may underlie each of these traits.  Then, we quantify the strength

of evidence for epistasis impacting each detected QTL, and then assess the

overall signal of epistasis across traits.  Lastly, we investigate the reaction

norm of the QTLs underlying a subset of the traits, related to pigmentation,

that were studied in two different temperature treatments.  These

investigations provide new insights into both the genetic basis of adaptive

trait changes and the dependence of adaptive alleles on genetic interactions.

226
227   **Methods**

## Mapping cross design

We report two new Recombinant Inbred Line (RIL) mapping panels

of *Drosophila melanogaster*.  Each RIL set was derived from a cross

between two inversion-free inbred lines from distinct geographic

populations, in each case pairing a strain from the southern-central African

ancestral range of the species (Sprengelmeyer *et al.* 2020) with a strain from

a cooler derived environment.  One cross was between a Zambian inbred

line (ZI215N, BioProject PRJNA861300) and a highland Ethiopian inbred

line (EF43N, sequenced in Lack *et al.* 2016); the other was between a

different Zambian inbred line (ZI418N, sequenced in Sprengelmeyer & Pool

2021) and a French inbred line (FR320N, sequenced in Lack *et al.*

2016).  Each cross was allowed to interbreed in an intercross design for 13

and 12 non-overlapping generations, in the Ethiopian and French cross,

respectively.  Then, the offspring of individual females were inbred for 5

generations to create each RIL.  We obtained 293 and 328 RILs for the

Ethiopian and French panels, respectively (BioProject PRJNA861300).  The

Zambian lines were collected in Siavonga, the Ethiopian line was collected

in Fiche, and the French line was collected in Lyon (Lack *et al.* 2015).  The

food used to rear the flies during the experiment and during the

phenotyping assays was prepared in batches of 4.5 L of water, 500 mL of

cornmeal, 500 mL of molasses, 200 mL of yeast, 54 g of agar,  45 mL of 10%

tegosept (in 95% ethanol), and 20 mL of propionic acid.

*Phenotypes*

Ethiopian flies have increased wing area, even after correcting size

relative to their larger body mass (Lack *et al.* 2016). We measured the wing

length of female flies that were at least 3-days old. 10 female flies per RIL

were photographed using an Amscope SM-4TZZ-144A dissection

microscope under $CO_2$ anesthesia. The pictures were scored with ImageJ

(Abramoff *et al.* 2004). Wing length was measured as the distance from the

intersection of the L4 longitudinal vein and the anterior cross vein to the L3

longitudinal vein intersection with the wing margin.

Ethiopian flies are darker than Zambian flies (Bastide *et al.* 2014). We

photographed and measured two to five females per RIL using a dissecting

scope. Resolution (3584 x 2748 pixels), gain (3.0), exposure time (99.84 ms),

magnification (20x), and illumination level were kept constant using an

Amscope adaptor for LED lamp at maximum lighting. White balance was

set at the same resolution using ColorChecker white balance. We measured

three common traits for pigmentation: the pigmentation of the

mesopleuron in greyscale proportion ranging from 0 for white to 1 for

black (herein, mesopleuron, a thorax trait), the pigmentation of the

background of the fourth abdominal segment in the same greyscale

proportion (herein, A4 Background), and the proportion of the fourth

abdominal segment that was covered by the black stripe (herein, stripe

ratio).  Relatively low correlations (computed via regression) among these

three traits were observed among independent isofemale strains within the

Ethiopia and Zambia populations ($r < 0.35$), except for a higher correlation

between mesopleural and A4 Background for Ethiopia specifically ($r =$

0.646; Bastide *et al.* 2014).

Pigmentation is also plastically influenced by temperature, and the

Ethiopian population is located in a colder region, on a high plateau 3 km

above sea level.  Therefore, we measured the three pigmentation traits in

flies raised at two temperatures (25 °C and 15 °C), to uncover potential

cryptic variation that could be present in the warmer Zambian (midpoint

temperature: 25 °C) but is only expressed in the colder Ethiopian (midpoint

temperature: 11 °C) population.  We also report plasticity for these three

pigmentation traits, as the difference in the trait measured at 25 °C and

15 °C.

French flies have higher resistance to ethanol (Cohan & Graf 1985,

Sprengelmeyer & Pool 2021) and cold (Pool *et al.* 2017, Huang *et al.* 2021)

than Zambian flies.  Ethanol resistance was measured as the average time

300 female flies, 3- to 5-day old, from each RIL needed to become

immobile when exposed to 18% ethanol.  The flies were placed in 50 ml

falcon tubes with ethanol-saturated tissue placed on the bottom, and the

number of mobile flies was scored every 15 minutes for six hours.  Cold

tolerance was measured as egg to pupa viability when raised at 15 °C.  Given

that very few dead pupa were observed across RILs, this measure was taken

as a reasonable time-saving proxy for the egg to adult viability difference

previously demonstrated between the parental strains (Huang *et al.*

2021).   To obtain this metric, six females and six males (3-5-day old) were

collected from each RIL and allowed to lay eggs for 15 hours.  The egg to

pupa viability was estimated as the number of emerged adults plus left pupa

in a vial at Day 34-35 divided by the number of eggs present in the vial after

the parents were removed (Day 1).

The song pattern trait was the ratio of slow pulses over all pulses (slow

and fast) in the song made by males during courtship.  Although not

expected to be under local environmental selection *a priori*, male courtship

is often under strong sexual selection.  In addition, a large number of

nervous system genes appear to have evolved between warm- and cold-

adapted *D. melanogaster* populations (Pool *et al.* 2017), representing one

potential source of pleiotropic effects on song traits.  We found that

Zambian male flies display a greater fraction of slow to fast pulses than

French flies (Lollar *et al.* 2025), a novel pulse mode classification recently

discovered within the *D. melanogaster* species complex (Clemens *et al.*

2018).  The song data was recorded in the Stern Lab at Janelia Farms

Research Campus, using materials and methods described by Arthur and

colleagues (Arthur *et al.* 2013).  Song was annotated without human

intervention with SongExplorer (Arthur *et al.* 2021).  Annotations were

classified with locally modified versions of BatchSongAnalysis

(https://github.com/dstern/BatchSongAnalysis), using previously trained *D.*

*melanogaster* models provided by Ben Arthur (Lollar *et al.* 2025).

We expected that there might be some level of correlation among the

traits of any given cross purely due to either linkage among causative

variants or else pleiotropy.  We calculated the pairwise correlation between

all the traits measured for each cross to investigate the degree to which they

are correlated.

324
*DNA Extraction and Sequencing*

For each RIL, we extracted DNA from five to ten female flies from each

line.  The flies were crushed and homogenized using pipette tips in strip

tubes, in a solution of 100 µl of lysis buffer (50 mM Tris-HCl with pH = 8,

100 mM EDTA, 100 mM NaCl, and 0.5% SDS) and 2 µl of proteinase K (IBI

Scientific).  The samples were incubated to inactivate the proteinase K (55 °C

for 5 minutes, 95 °C for 10 minutes, and 25 °C hold).  The supernatant of the

fly lysate was transferred to new tubes, and we added SPRI beads (Sera-mag

SpeedBeads SPRI beads, [GE Healthcare 24152105050250]) in a 1.8 to 1

proportion of beads to lysate.  The beads were used in a solution of PEG

8000 (9 g), Tween 20 (27.5 µl), 5M NaCl, and 3 mL TE Buffer (composed of

100 µl of 1 mM EDTA, 500 µL of 10 mM Tris HCl).  1 mL of the beads were

added to the buffer mix and filled with water until the final volume was

mL.  The tubes were incubated for at least three minutes at room

temperature for DNA to bind to the beads and then placed on a magnetic

plate for five minutes to remove the beads from the solution.  The

supernatant was discarded, and the beads were washed with two rounds of

80% ethanol (90 µL per well).  DNA was eluted beads with 30 µL of $H_2O$ and

then used the magnetic plate to transfer only the resuspended DNA to new

tubes.

Genomic libraries were prepared following Adams *et al.* (2020), with

size selection, cutoff at 300 bp, and cleanup performed using the Zymo

Select-a-Size DNA clean & Concentrator [catalog No. D4080].  Libraries

were sequenced at the UW-Madison Biotechnology Center on the Illumina

NovaSeq 6000 platform, with 150 bp paired-end reads.

350
*DNA Alignment and Ancestry Calling*

The reads were mapped to the *D. melanogaster* (v5.57) reference

genome using BWA-MEM, version 0.7.17 (Li 2013).  We used Ancestry

HMM (Corbett-Detig & Nielsen 2017) to estimate ancestry along the

genome of each RIL, allowing for diploid ancestry calls (*i.e.* homozygous for

either parent, or heterozygous).  Ancestries were summarized in non-

overlapping windows defined to contain 1,000 SNPs in the Zambian

population (19 kb average size), using the majority ancestry call in the case

of an ancestry breakpoint within the window.  Window size was chosen to

ensure ample variation for ancestry calling within each window, and to

moderate the number of QTL and epistasis tests performed in order to

ensure computational tractability, while still allowing a genomic resolution

much narrower than the expected megabase scale ancestry tracts. When

two adjacent windows had the same ancestry genotype across all RILs, the

two windows were merged. We used a recombination map based on

Comeron *et al.* (2012) but modeled only one generation of recombination

when using Ancestry HMM, in order to focus on conservatively-defined

ancestry switches (pipeline available on

https://github.com/ribeirots/RILepi).

370
*QTL Mapping*

To test whether adaptive loci showed evidence of epistatic

interactions we first used QTL mapping to identify the adaptive loci and

then performed epistasis tests between the focal loci and all other loci in the

genome. Each locus is a genomic window of ~19kb, defined and genotyped

as described above.

To identify loci underlying trait changes, we employed a similar

approach to the R/qtl (Broman *et al.* 2003) *scanone()* function, but with an

additive framework, modifying the genotypes to be coded as numeric

variables instead of categorical (https://github.com/ribeirots/RILepi), using

Python3 (Van Rossum & Drake 2009). The additive modification was

preferred in light of the small counts of specific genotypes due to the low

heterozygosity and the variable genomic ancestry skew present in our RIL

panels.  We fit a model of the phenotype given the ancestry genotype of

each window (Trait ~ Marker).  In our model, we coded the genotypes at Q

as numeric values: 0, 1, and 2 for the homozygous Zambian ancestry,

heterozygous ancestry, and Ethiopian or French homozygous ancestry,

respectively.

The initial QTL mapping is based on the comparison of a model with

a single QTL *versus* a model without any QTL, performed repeatedly one

locus at a time.  For each locus, we obtained a logarithm of the odds (LOD)

score as the ratio between the log-likelihoods of the model with one QTL

over the model without a QTL.  Because both RIL sets had average ancestral

frequencies skewed against the Zambian ancestry (68% Ethiopia ancestry

and 75% France ancestry), we excluded the windows with an ancestry bias of

90% or greater for either ancestry to avoid the effects of extremely uneven

sample sizes on the model.  The genome-wide significance of a QTL was

tested using 10,000 permutations, in which trait values were randomly

shuffled.  From each permutation, we saved the highest LOD score

obtained for any window (excluding windows with ancestry skew of 90% or

greater) and empirical LOD scores from each window were compared to

the null, permutation LOD score distribution.  We considered the windows

whose LOD score's p-value was lower than 0.10 as putative QTLs (*i.e.* a less

than 10% chance of having a QTL signal of that strength anywhere in the

genome). A p-value threshold of 0.10 was chosen aiming to include a

sufficient number of QTLs to be tested for epistasis. We emphasize that

these QTL p-values are trait-specific, and do not attempt to quantify the

likelihood of any QTL peak for any analyzed trait having a given LOD

score.

Each putative QTL surrounded by windows with a lower LOD score

was considered a putative QTL peak. If two putative QTL peaks were not

separated by a window with a LOD score lower than the minor peak's LOD

score minus 1.5, the minor peak was removed. Finally, the remaining

putative QTL peaks were filtered to remove any peak within 10 cM of a

QTL peak with a higher LOD score, in order to increase the independence

of QTLs used in epistasis testing. The confidence interval for these QTLs

was defined as all contiguous windows with a LOD higher than the QTL's

peak LOD score minus 1.5. Heritability-based QTL effect ($h^2$) was

calculated as described in R/qtl for an intercrossing mapping design

(Broman *et al.* 2003).

The power of the QTL mapping analysis was calculated based on

simulated data for each RIL panel. We simulated a normally distributed

phenotype (mean = 0, standard deviation = 0.75) with additive effect of the

derived allele varying from 0.05 to 0.5 in 0.05 increments and without

dominance. We randomly sampled genotype distributions for each

simulation from the empirical RIL panels, to a total of 10,000 simulations

for each additive effect and RIL panel.  We calculated the LOD score and

heritability-based QTL effect ($h^2$) for each simulation as described above,

and calculate its p-value based on the permutations of the empirical data for

each RIL panel.  QTL mapping power was determined for each additive

effect and RIL panel as the proportion of simulations with p-value equal to

or lower than 0.1–the same threshold used in our analysis.

433
*Candidate Genes and Population Genetics*

To identify genes potentially underlying the adaptive traits, we

scanned our QTL regions for signatures of local adaptation based on

population genetics.  We sequenced genomes from inbred lines from each

of the studied populations to calculate a haplotype-based statistics: $\chi_{MD}$

(Lange & Pool 2016) and two $F_{ST}$ statistics (using Reynolds *et al.* 1983), $F_{ST\_FullWin}$

(window-wide $F_{ST}$ calculated using all the SNPs in the window) and $F_{ST\_MaxSNP}$

(the highest $F_{ST}$ value from a single SNP within the window), that have

power to distinct power to detect distinct kinds of selective events (da Silva

Ribeiro *et al.* 2022).  We defined as outlier windows those that fell within the

top 1% of windows on the same chromosome arm for any of the above three

statistics.  We focused on regions with recombination rates generally above

0.5 cM/Mb (Comeron *et al.* 2012) due to more localized signatures of

selection: 2.3–21.4 Mb of the chromosome X, 0.5–17.5 Mb of arm 2L, 5.2–

20.8 Mb of arm 2R, 0.6–17.7 Mb of arm 3L, and 6.9–26.6 Mb of arm 3R.  We

selected as candidates all genes that overlapped with the outlier windows as

well as the first gene up- and down-stream of the outlier window, to

account for instances in which the target of selection is in a regulatory

region outside the gene region.  We also identified other genes within these

QTLs with a known or potential association with the phenotype based on

FlyBase (release FB2023_05) annotation.

455
*Epistasis Testing*

Epistasis tests were performed between each focal QTL and all of the

remaining genomic windows outside its confidence interval.  The epistasis

test for each window pair consisted of an interaction LOD score obtained as

the ratio between the log-likelihoods of the model with two QTLs and their

interaction (Trait ~ Marker1 + Marker2 + Marker1:Marker2) *versus* a model

including both QTLs but no interaction (Trait ~ Marker1 + Marker2).  The

interaction LOD score for each pair was compared against

permutations.  Permutations were specific for each QTL analyzed.  We

shuffled the phenotype, fixed the focal QTL (Marker1), and calculated the

interaction LOD score of Marker1 against all other windows in the

genome.  The highest genome-wide interaction LOD score from each

permutation was kept and the empirical interaction LOD scores were

compared against this null, permuted interaction LOD distribution.

*Epistasis Meta-analysis*

We combined the results from all the epistasis tests across mapping

crosses and phenotypes to test the hypothesis of whether adaptive QTLs

have epistatic interactions. We collected the set of epistasis p-values from

each QTL and its most likely interactor, quantifying the probability of

obtaining any interaction term between that QTL and any partner locus in

the genome with an interaction LOD score as high as the one observed. We

used Fisher's combined probability test to ask if the p-values from each

QTL and their most likely interacting locus reject the null hypothesis that

there is no epistasis. The p-values should follow a uniform distribution if

the null hypothesis is true. Therefore, we performed an additional step to

investigate how many of the lowest p-values would need to be removed

from the data set to obtain a median value near 0.50, in order to ask

whether our cumulative data set showed any hints of epistasis across all

analyzed QTLs.

*Epistasis Power Analysis*

488    We calculated the power of our approach and our data sets to detect

489    an interaction by simulating phenotypes with different interaction

490    strengths. We chose one empirical QTL with intermediate sample size and

491    estimated the additive QTL effect among the detected QTL to serve as the

492    basis for the simulations. The QTL chosen was detected for the abdominal

background pigmentation at 25 °C, it had an estimated effect size equal to
0.035, mean trait value equal to 0.3485, and a standard deviation equal to
0.0697. The observed genotype distribution of the QTL peak window was
used for the simulations. Our null simulations generated phenotypes based
on a normal distribution with the observed trait mean and standard
deviation, and the simulated values were modified based on the trait effect
and individual genotypes (1x effect for heterozygous and 2x effect for non-
Zambian homozygous). We then calculated the interaction LOD score of
the focal peak window against all other genome windows outside the
original QTL's confidence interval and kept the highest genome-wide
interaction LOD score obtained. We repeated this process 10,000 times to
obtain a null (no interaction) distribution of interaction LOD scores.

   To simulate epistasis, we used the genotype distribution of the
window most likely to be interacting with our focal QTL: the window with
the highest empirical interaction LOD score. We modified the empirical
effect of the focal QTL on the simulated phenotype based on the genotype
of the interacting genomic window. If the genotype of the interacting
window was Zambia homozygous, the original effect was not modified, if it
was non-Zambia homozygous, the original effect was multiplied by the
interacting factor $I$, and if it was heterozygous, the original effect was
intermediate (*i.e.* multiplied by $(1+I)/2$). The values of $I$ we simulated varied
from -2 to 4, depending on the nature of the epistatic interaction being

simulated. We simulated positive epistasis effects (original QTL effect

increased by the presence of the epistatic allele) with $I$ ranging from 1.125 to

4. For diminishing returns negative epistasis (original QTL effect decreased

by the presence of the epistatic allele), $I$ ranged from 0.889 (1/1.125) to

0.5. We simulated masking epistasis (the original QTL effect nullified in the

presence of the epistatic allele) with $I = 0$. Lastly, for sign epistasis (the

original effect changes in sign, *e.g.* instead of the non-Zambia QTL allele

producing larger trait it produces smaller traits), we examined effects from -

0.5 to -2.

Each interaction simulation was compared against the null

distribution and a p-value was calculated as the proportion of null

simulations with an interaction LOD score higher than the simulated

interacting LOD score. Power was calculated for each interaction effect $I$ as

the proportion of simulations that had p-value lower than 0.05.

We also calculated the power of our meta analysis based on 14 QTLs'

interaction p-values. For each interaction effect, we sampled 14 interaction

LOD score p-values and calculated (1) how often the fisher combined p-

value was as extreme or greater than the one we observed and (2) how often

at least one p-value was lower than the lowest p-value we obtained for our

empirical data.

*Reaction Norms*

To investigate genotype-by-environment interactions for

pigmentation traits, we combined phenotypic data from RILs raised at the

two temperature treatments (15 °C and 25 °C) with the genotype of each

non-overlapping QTL.  To test whether the genotype-by-environment

interaction was significant, we used a linear model with the phenotype as

the response variable and the genotype, environment, and genotype-by-

environment interaction as the explanatory variables.

544
545
**Results**

*Genotype and phenotype data*

Whole genome sequences for the RILs had mean depth of coverage of

3.3X (S.D. 1.96 among RIL mean depths, Table S1) and 7.58X (S.D. 9.09,

Table S2) per site for EF and FR RILs, respectively, which should be more

than sufficient to call population ancestry in large chromosomal blocks

(Corbett-Detig & Nielsen 2017).  Ancestries for both RIL sets were skewed

toward lower Zambian ancestry, with averages of 67.84% Ethiopian ancestry

and 77.88% French ancestry (full ancestry genotypes in Tables S3, S4 ).  After

five generations of inbreeding, a low level of parental strain heterozygosity

was present in the final panel: 8.71% and 5.3% on the Ethiopian and French

panels, respectively (at the level of genomic windows).  The sample size

varied for each trait based on the number of RILs successfully phenotyped

(Table 1, S1, S2).

**Table 1.** Summary of mapped traits. Listing the mapping cross used for each

trait (EF = Ethiopia, FR = France), the number of phenotyped lines for each

trait, the number of QTLs, and the number of QTLs with epistatic

interaction for each phenotypic trait.

| RIL panel | Trait | Sample size | # QTLs |
|---|---|---|---|
| EF | A4 Background 25 °C | 278 | 1 |
| EF | Stripe Ratio 25 °C | 278 | 1 |
| EF | Mesopleuron 25 °C | 278 | 1 |
| EF | A4 Background 15 °C | 190 | 0 |
| EF | Stripe Ratio 15 °C | 190 | 1 |
| EF | Mesopleuron 15 °C | 190 | 1 |
| EF | A4 Background (Plast.) | 178 | 0 |
| EF | Stripe Ratio (Plast.) | 178 | 0 |
| EF | Mesopleuron (Plast.) | 178 | 0 |
| EF | Wing Length | 268 | 5 |
| FR | Cold Tolerance | 304 | 0 |
| FR | Ethanol Resistance | 277 | 3 |
| FR | Song Pattern | 294 | 1 |

The distribution of the phenotypes were unimodal for all traits with

varying degrees of variance and low to moderate skew (Figures S1, S2).

Correlations among the analyzed traits were examined for each RIL panel,

which may indicate pleiotropy or else linkage of causative alleles. Among

pigmentation traits, the thoracic mesopleuron trait was strongly correlated

with abdominal background: r = 0.687 at 15 °C, r = 0.800 at 25 °C, r = 0.782

572 for the plasticity of these traits (Table S5). Stripe width showed moderate,

573 significant correlations with both of those traits at 25 °C ($r$ = 0.460, and $r$ =

574 0.253), but weaker correlations at 15 °C ($r$ = 0.169, and $r$ = -

575 0.085). Abdominal background showed a non-significant correlation

576 between the two temperatures, while the other two traits showed moderate,

577 significant correlations between temperatures. In light of the correlations

578 observed among many pigmentation traits, overlapping QTLs for

579 pigmentation traits were not treated as independent QTLs, and only the

580 QTL with the highest LOD score was used in downstream analyses. No

581 pigmentation trait was significantly correlated with wing length (Table S5),

582 the only other trait scored among the Ethiopian RIL panel. Among traits

583 scored from the French RIL panel, the song pattern was weakly correlated

584 with ethanol resistance, $r_2$ = 0.13 (Table S6).

585
586 *Single locus QTL mapping*

587  We identified potential QTLs for eight out of the thirteen investigated

588 traits (Figure 1; Table 1). We list the genome-wide distribution of LOD

589 scores and p-values for each trait in the Table S7. We also present the

590 phenotypic distributions observed for each genotype at the QTL peak

591 window (Figures S3, S4) including the pulse song usage trait that is

592 discussed in more detail by Lollar *et al.* (2025). The traits with no detected

593 QTLs were A4 Background at 25 °C, all the three pigmentation plasticity

traits, and cold tolerance (genome-wide LOD scores shown in Figure S5).

The number of QTLs per trait ranged up to five, for wing length, with other

traits yielding one or three QTLs (Figure 1, Table 1).  Twelve of the fourteen

QTLs were significant at $p < 0.05$, while two marginally significant QTLs -

one for A4 Background at 25 °C ($p = 0.0591$) and one QTL for ethanol

resistance ($p = 0.0693$) – were also considered in downstream analyses

(Table 2).  The effect of each QTL, based on estimated $h^2$, varied from 0.033

to 0.285 and we had high power (>0.8) to detect QTLs of effect size 0.1

(Figure S6).

**Figure 1.** The number of QTLs for each trait ranged from 0 to 5. In this figure, only the traits with at least one identified QTL are shown (phenotype distributions for each genotype at each QTL peak window are shown in Figures S3, S4); the traits without significant QTLs (Table 1) can be seen in Figure S5. Each panel shows the -log of the (trait-specific) p-value for the LOD score of the genomic windows. A4 Background (Figure 1E) is shown here with no QTL above the threshold, but one marginally significant QTL (p-value = 0.053, Table 2). Windows filtered out for ancestry skew are given a value of -0.25. The red dashed line represents the 0.05 p-value cutoff based on 10,000 permutations. The color of the dots represents the chromosome arm of each genomic window. Note that for wing length, a single QTL spans a broad low recombination centromeric region between 2L and 2R.

**Table 2.** Characteristics of detected QTLs for all analyzed phenotypes, and possible underlying genes. The single QTL coefficient shows the magnitude of phenotypic change by substituting one allele of the derived population ancestry at that locus, which is shown on the scale of the measured phenotype. The heritability due to the QTL ($h^2$) shows how much of the phenotypic variance is explained by the QTL (Broman *et al.* 2003). QTL p-value and epistasis p-value were obtained based on 10,000 permutations. * an estimated p-value of zero means the empirical value was

more extreme than any permutation.  Q4 was excluded from subsequent

meta-analysis due to a stronger overlapping QTL for a correlated trait.

Further details of each QTL are given in Table S8.  Candidate genes are

given in light of a priori functional knowledge based on FlyBase annotation

or as population genetic outliers for local adaptation statistics (n/a indicates

that QTL was in a region of low recombination and was not analyzed for

signatures of selection).  Genes that may underlie the song trait are instead

described by Lollar *et al.* (2025).

| RIL | Trait | QTL ID | LOD score | Trait Coeff | $h^2$ | QTL p-value | Epist. p-value | QTL Confidence Interval | Functional Candidate Genes | Genes with Signatures of Selection |
|---|---|---|---|---|---|---|---|---|---|---|
| EF | A4Back 25C | Q1 | 3.274 | 0.012 | 0.033 | 0.0591 | 0.736 | Chr3R:21614478 . 22512754 | | *TwdlQ, beat-vii, scrib* |
| EF | Mesopl 15C | Q2 | 5.819 | 0.012 | 0.112 | 0.0006 | 0.699 | Chr3R:16750589 . 16945416 | | |
| EF | Mesopl 25C | Q3 | 5.89 | 0.014 | 0.076 | 0.0003 | 0.714 | Chr3R:17549430 . 17666763 | *burs* | |
| EF | Stripe 25C | Q4 | 6.335 | 0.035 | 0.285 | 0.0002 | 0.643 | ChrX:1057169 . 2420139 | *Hr4, y* | n/a |
| EF | Stripe 15C | Q5 | 9.924 | 0.071 | 0.176 | 0* | 0.244 | ChrX:0 . 2141596 | *Hr4* | n/a |
| EF | Wing | Q6 | 9.581 | 0.026 | 0.109 | 0* | 0.322 | Chr2L:12902781 . 13046286 | | *A16, Vha68-2, ACXE, CG16800, Tor, Ube4B* |
| EF | Wing | Q7 | 8.628 | 0.024 | 0.098 | 0* | 0.672 | Chr2L:22740849 . 23011543 | *Slmap* | n/a |
| EF | Wing | Q8 | 3.867 | 0.022 | 0.087 | 0.0183 | 0.444 | Chr3L:10353468 . 11407946 | *Cpr67Fa2, FoxK, Rbfox1, scyl, wls* | *Fad2, CG32079, klu* |

| | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|
| EF | Wing | Q9 | 3.526 | 0.019 | 0.059 | 0.0369 | 0.941 | Chr3R:18722896 . 19232530 | *Cow, Gbp3, hh, lrk1, p53, pnt, Rassf* | *Pnt* |
| EF | Wing | Q10 | 4.286 | 0.016 | 0.053 | 0.0073 | 0.183 | ChrX:20419847 . 20893609 | | *CG15450, Cyp6v1, hydra, shakB, CG1835, CG11227* |
| FR | EtOH | Q11 | 3.565 | 22.516 | 0.051 | 0.0368 | 0.208 | Chr3L:10829907 . 11166414 | *Aps* | *CG42831, tna* |
| FR | EtOH | Q12 | 3.253 | 27.506 | 0.127 | 0.0693 | 0.853 | Chr3R:3268049 . 3791969 | | n/a |
| FR | EtOH | Q13 | 3.74 | 19.166 | 0.038 | 0.0246 | 0.476 | Chr3R:23770233 . 24580087 | *fkh* | *trv, htt, CG34354, CG43125, snu* |
| FR | Song | Q14 | 9.111 | 0.019 | 0.079 | 0* | 0.273 | Chr3R:21997113 . 22171358 | | |

Of the five pigmentation QTLs, the two identified for stripe ratio (Q4

and Q5, at 25 °C and at 15 °C respectively) overlapped.  We mapped the

residuals values of the correlation between these two traits to investigate

whether there was a signal beyond their correlation and did not obtain any

significant QTLs (Figure S7).  Therefore, only the overlapping QTL with the

higher LOD score (15 °C) was retained for meta-analysis (Table

S8).  Confidence intervals for the two QTLs detected for mesopleuron (one

at 25 °C and one at 15 °C) were separated by just over 600 kb, so they were

kept as two independent QTLs for the meta-analysis (Table 2).

The direction of all QTLs (Table 2) was in the expected direction:

substituting alleles of the derived population resulted in phenotypic

changes in the direction of the derived populations.  This pattern is

congruent with a scenario where these QTLs are contributing to locally

adaptive trait changes (involving either the measured traits or else

pleiotropically correlated traits under selection), given that if the

phenotypic difference were neutral, we would expect some QTLs to show

an effect in the opposite direction (Orr 1998).

*Outlier and Candidate Genes within QTLs*

To generate hypotheses for possible causative genes underlying trait

changes, we used population genetic summary statistics to identify genes

within our QTLs that are potentially under local adaptation.  Specifically,

we flagged genes associated with windows that fell within the chromosome

arm's top 1% of windows for window $F_{ST}$, maximum SNP $F_{ST}$, or the haplotype

statistic $\chi_{MD}$ (Table 2).  These local adaptation scans were not conducted in low

recombination QTL regions due to the expected lack of gene-scale

resolution of selection signals in such genomic intervals.  Since a striking

signal of local adaptation is not guaranteed at loci underlying these strong

population trait differences (*e.g.* if they involved soft partial sweeps), we also

noted the presence of other functionally relevant candidate genes within

each QTL.

We found a total of 26 candidate genes with functionally relevant

annotations on FlyBase related to their specific measured phenotype. Q3

(mesopleuron 25 °C) included the gene *burs* (Dewey *et al.* 2004), implicated

in cuticle tanning and hardening.  Q4 and Q5 (the two stripe ratio

QTLs)both included the candidate gene *Hr4*, a gene that when suppressed

results in reduced pigmentation (Rogers *et al.* 2014).  One also included the

gene *yellow*, canonically implicated in pigmentation variation and

evolution (e.g. Massey & Wittkopp 2015).  This pair of QTLs overlapped

others previously detected for the same trait in distinct Ethiopia-Zambia

crosses (Bastide *et al.* 2016).  The gene *ebony*, at which a soft sweep signal

associated with a pigmentation QTL was previously characterized for

Ethiopian *D. melanogaster* (Bastide *et al.* 2016), fell just between the closely-

located Q2 and Q3 (mesopleuron 15 °C and 25 °C).

At least one gene related to wing development was found within each

of the five wing length QTL regions.  Within Q6 (on arm 2L), our

population genetic outliers included the *mechanistic Target of rapamycin*

gene (*mTor*), which plays a key role in insulin signaling and growth

regulation, and has been tied to wing size and development specifically

(Parker & Struhl 2015). Outliers inside this QTL also included the potential

wing regulators *Vha68-2* and *Ube4B* (Krupp *et al.* 2005, Blanco *et al.* 2010,

Okada *et al.* 2016, López-Varea *et al.* 2021).  Within Q7 (low recombination

regions of 2L), among the few genes present was *Slmap*, a hippo signaling

gene shown to alter wing size (Zheng *et al.* 2017).  Within Q8 (3L),

population genetic outliers included *klu*, which regulates cell size and

proliferation, and is associated with wing size defects (Schertel *et al.*

2015).  Within Q9 (3R), outliers included *pnt*, related to wing morphogenesis

(Dworkin & Gibson, 2006, Bejarano *et al.* 2008, Paul *et al.* 2013).  The X-

linked wing QTL (Q10) included an outlier gene possibly involved in wing

development, *shakB* (Krishnan *et al.* 1993).  Q7, Q8, and Q10 overlap with

previously detected wing length QTLs, each from a different Ethiopia-

Zambia cross out of four previously analyzed mapping crosses

(Sprengelmeyer *et al.* 2022).

For the ethanol resistance QTLs, we identified two population genetic

outlier genes (*htt* and *trv*) previously implicated in response to ethanol

(Fochler *et al.* 2017) within Q13 (3R). Outlier genes detected within the other

two QTLs had no annotated connection to ethanol. These QTLs did not

overlap with those previously detected for the same trait from distinct

France-Zambia crosses (Sprengelmeyer *et al.* 2021), in line with the

genetically variable basis of this trait identified by that study.

Overall, we identified both novel candidate genes associated with the

detected QTLs and genes that had already been implicated with their

respective trait. The identification of genes already known to underlie these

traits (although in many cases not their evolution) provides support for the

approach we used here.

711
*Epistasis*

To investigate whether there is evidence of epistasis among adaptive

loci, we performed an epistasis scan for each single QTL and a meta-

analysis combining the results across non-correlated traits. We did not

identify any individually significant epistatic QTL pairs. The lowest

genome-wide interaction LOD score p-value we obtained was 0.183, for a

wing length QTL (Table 2).

719  Our meta-analysis of the combined non-overlapping single QTLs also

720 supports the hypothesis that there is no strong epistasis involving adaptive

721 loci. By combining the interaction p-values for all non-overlapping QTLs

722 (where each p-value denotes the probability of that QTL having any

723 interaction LOD effect that strong in the genome, based on 10,000

724 permutations), we obtained a p-value of 0.763 using Fisher's combined

725 probability test. This result indicated that the observed p-values do not

726 deviate from the null expectation of uniformly distributed p-values

727 expected without epistasis involving adaptive loci.

728  Because our sample sizes of RILs may not be suitable for detecting all

729 magnitudes of epistasis, we conducted a simulation-driven analysis to

730 indicate the strength of our negative results, and to indicate if there was a

731 parameter space of epistasis that we could confidently rule out based on

732 these results. Based on this analysis, our method had a high power (over

733 80%) to detect an epistatic QTL pair in scenarios of sign epistasis, in which

734 the interaction effect changed the direction of the main effect, as well as

735 high power to detect strong positive epistasis in which the interaction

736 effected increased the main effect at least 2.5-fold (Figure 2, Table S9). We

737 also tested the power of our meta-analysis by re-sampling 13 p-values from

738 simulated data sets of different interaction effects (Figure 2). This analysis

739 indicated that we should have had power to detect similar scenarios of

740 strong epistasis as indicated for the lowest p-value analysis above (*i.e.* sign

epistasis and positive epistasis at least doubling the main effect, and that we

would likely have detected masking epistasis as well, in which the main

effect was erased by the modifier). Therefore, although the power to detect

moderate and weak epistasis is reduced rapidly, the lack of detected

epistatic interactions from our empirical data supports the hypothesis that

strong forms of epistasis did not play an important role in these instances of

evolution.

**Figure 2.** Simulation analysis indicating that strong positive epistasis or sign

epistasis should have been detectable for the empirical data. Simulations

based on randomly permuted individual genotypes and simulated epistatic

effects (see Materials and Methods) were conducted to assess our ability to

detect varying models of epistasis based on either the lowest empirical

epistasis p-value (red series) or the Fisher-combined epistasis p-value across

13 analyzed QTLs (blue series). Here, the epistasis factor (x-axis) represents

the multiplier that a modifier locus exerts on the primary QTL's

effect. Thus, negative values represent sign epistasis (pink shaded area),

zero represents masking epistasis, values between 0 and 1 indicate negative

epistasis (yellow shaded area), and values above 1 indicate positive epistasis

(green shaded area). The y-axis shows the proportion of occurrences out of

the 1,000 resampled instances in which a given model of epistasis yielded a

lowest p-value lower than the observed 0.183 (red), or else how often the

combined Fisher p-value was lower than the observed 0.763 (blue).

**Figure 3.** The reaction norms of the four non-overlapping pigmentation
QTL show darker phenotypes for flies raised at 15 °C than at 25 °C and for
flies with Ethiopia ancestry alleles, corroborating expectations. (A) Q1: A4
Background at 25 °C, with peak centered near 21.6 Mb on 3R. (B) Q2:
Mesopleuron at 15 °C, near 16.8 Mb on 3R. (C) Q3: Mesopleuron at 25 °C,
near 17.6 Mb on 3R. (D) Q5: Stripe ratio at 15 °C, near 1.3 Mb on the X. The

y-axis shows the mean phenotype for each genotype and temperature

treatment; a higher number is a darker phenotype.  ZZ = Zambia

homozygous, EZ = heterozygous, EE = Ethiopia homozygous.

*Phenotype plasticity for pigmentation QTLs*

Fly pigmentation is known for its temperature-based plasticity; flies

reared at lower temperatures develop darker phenotypes than those raised

at warm temperatures (David *et al.* 1990).   Our results recapitulated this

behavior, in that average pigmentation traits were higher (*i.e.* darker) at

15 °C (Figure 3).  We identified one overlapping QTL for stripe ratio at both

temperatures (Table 2).  Based on the trait correlation and the shared QTL,

at least part of the genetic architecture underlying adaptive melanism

appears to be shared between temperatures.

The reaction norm of four pigmentation QTLs also showed that flies

raised at 15 °C showed darker phenotypes for all traits and even darker

phenotypes for the Ethiopia homozygotes, as expected (Figure 3).  We also

found significant genotype-by-environment interactions for two of these

QTLs – A4 Background (p = 0.003) and stripe ratio (p = 0.007) (Figure

3A,D).  The change in stripe ratio (Figure 3D) was greater for flies with the

homozygous Ethiopia genotype.  Here, the more derived-like environment

appears to enhance the phenotypic consequences of the Ethiopian

pigmentation variants.  In contrast, for the A4 Background QTL, the

phenotype means at 15 °C only show a small difference between

homozygous genotypes, compared to a larger difference at 25 °C, which is

congruent with this QTL being found only at 25 °C (whereas differences in

the heterozygote means may reflect small sample size effects).

798
799

**Discussion**

We report the creation of two new *Drosophila melanogaster* RIL

panels established from crosses between single inbred lines from Ethiopia

and Zambia, and from France and Zambia.  We used these new data sets,

together with an updated QTL mapping approach focusing on additive

effects, to investigate the genetic architectures of multiple traits that have

evolved to become differentiated between *D. melanogaster*

populations.  We did not find evidence for strong epistasis underlying their

evolution, but genotype-by-environment interactions were supported for

two of the three traits investigated.

810
*Genetic architectures underlying trait evolution*

Our QTL mapping results indicate genetic architectures for most

traits involving detectable loci with non-trivial effect sizes (5.3% to 21.4%;

Table 2), consistent with past bulk mapping studies for some of these traits

(Bastide *et al.* 2016, Sprengelmeyer *et al.* 2020, Sprengelmeyer *et al.*

2021).  In combination with those studies, our results reinforce previous

findings that evolved populations maintain persistent variability in the

genetic basis of pigmentation traits, wing length, and ethanol resistance.  We

did not find significant QTLs for cold tolerance or thermal plasticity of

pigmentation, which could reflect either a more complex genetic basis of

those traits or else greater non-genetic variance in those assays.

We have also identified genes within the detected QTLs that might be

underlying the relevant traits, including genes previously known to be

related to the phenotype and, with the use of population genetic signatures

of selection, novel candidate genes (Table 2).  The identified genes

represent viable hypotheses for contributors to the genetic architecture of

traits that appear to have evolved due to local adaptation targeting either

these or pleiotropically correlated traits but deserve further

investigation.  Additional studies using functional approaches such as

genome editing will be needed to confirm the roles of these genes in the

evolution of the respective traits.

832
*Lack of evidence for epistasis*

Despite the pervasiveness of epistasis underlying the genetic basis of

complex traits (Huang *et al.* 2012, Mackay 2014), we did not find evidence of

epistasis involving any of the detected adaptive loci (Table 2).  In light of

our power analysis (Figure 2), our empirical results primarily argue against

the presence of modifier variants within our RIL panels that trigger sign

epistasis, complete masking epistasis, or strong positive epistasis with regard

to the main effect loci. In contrast, our study does not speak to the presence

of quantitatively negative epistasis (*e.g.* diminishing returns epistasis) or

more moderate positive synergistic epistasis.

Previous studies have also found mixed results when investigating

epistatic interaction among adaptive loci (Malmberg & Mauricio 2005). In

some cases, the varied outcome for adaptive traits might reflect the

transient nature of epistatic interactions in a population, given that the

epistatic effects of two loci on a trait depend on the allele frequency of the

interacting alleles. Interestingly, studies mapping QTLs for fitness-related

traits (more directly related to reproduction and survival) have a more

consistent result, often showing more epistatic QTLs than additive QTLs

(Malmberg & Mauricio 2005).

One of the difficulties in extrapolating the results of experimental

crosses to natural populations is that experimental crosses start with a

limited amount of genetic variation (Mackay 2014). In our case, the crosses

were made between one inbred line from each parental population, which

will exclude some variants present in the source populations while elevating

some naturally rare variants to moderate frequency. Previous genetic

mapping studies of traits such as pigmentation (Bastide *et al.* 2016), ethanol

resistance (Sprengelmeyer *et al.* 2021), and wing size (Sprengelmeyer *et al.*

2022) have shown that single crosses made with different inbred lines from

the same population do not contain most of the genetic variation

underlying population differentiation in each trait.  However, our crosses

have the advantage of focusing on a simplified genetic architecture, which

may improve power to detect interactions among the variants present, in

addition to clear inference of the parental origin of alleles genomewide.

We highlight that our study likely underestimates the epistatic

interactions affecting *D. melanogaster* adaptive traits, since we only had

power to detect relative strong interactions (Figure 2, Table S9).  Based on

the limited numbers of RILs available, we could only detect strong additive

QTLs, and we needed to focus our epistasis scan on interactions with those

few strong QTLs.  Future research with a larger sample size (e.g. Torgeman

& Zamir 2023) would be helpful to achieve more extensive insights into the

scope of epistatic interactions affecting traits such as these.  Studies using a

more diverse set of starting lineages could mitigate some of the issues

related to the limited genetic variation in our study.  Additionally, drawing

information from diverse sources, such as genome-wide association studies,

bulk segregating approaches, and population genetics signatures of selection

could be useful in designing mapping studies capable of not only

identifying QTLs but candidate loci within QTLs.  Multi-trait QTL mapping

could also be a fruitful future direction to investigate the role of epistasis on

adaptations as it can increase mapping power (Pitchers *et al.* 2019).  Multi-

trait approaches can also infer pleiotropy and epistasis (Zhang *et al.* 2016),

adding a layer of complexity to the interpretation of results that could be

especially challenging when some of the measured traits are not expected to

have similar developmental and genetic bases.

*Genetic Architectures at Two Temperatures*

Although the three body pigmentation traits we studied were

somewhat correlated (Table S5), no shared QTL was noted between them

(Table 2).  Instead, we found two instances of overlapping or neighboring

QTLs for the same trait measured at different temperatures, with 25 °C and

15 °C being more similar to the mean temperatures in the Zambian and

Ethiopian populations, respectively.  These results are compatible with the

same or nearby genes being responsible for pigmentation differences at

both temperatures.

*Drosophila* pigmentation is known to show thermal plasticity, and

Ethiopian flies would have been expected to become somewhat darken on

this basis alone, and yet they have also adapted genetically to become

darker (Bastide *et al.* 2014).  It is unclear whether this pigmentation plasticity

is adaptive, which in itself can facilitate adaptive evolution (Ghalambor *et al.*

2007).  Unlike larger ectotherms, it has been estimated that the solar gain in

temperature from having dark pigmentation is at most a fraction of a

degree for an organism with the small size and high surface to volume ratio

of *Drosophila* (Willmer & Unwin 1981, Hirai & Kimura 1997).  Across African

populations of *D. melanogaster*, levels of ultraviolet radiation have been

found to be stronger predictors of pigmentation than temperature (Bastide

*et al.* 2014).

Gene-by-environment loci have recently been shown to be enriched

for signatures of positive selection (Lea *et al.* 2022), supporting the

hypothesis that environment-specific loci play an important role in local

adaptation. Here, we did find that two QTLs underlying pigmentation

evolution showed significant genotype-by-environment interactions

(Figure 3), suggesting that the rate of their response to selective events

might have changed upon colonization of colder environments. However,

we did not find consistent patterns of either gene-by-gene or gene-by-

environment interactions that met the predictions of cryptic variation. For

example, the absence of epistatic interactions means that the Ethiopian

alleles did not have their effect completely masked or reversed by another

gene on the ancestral background. And in the case of gene-environment

interactions, there was an enhanced QTL effect in the derived Ethiopia-like

cool environment in just one of two significant cases (*i.e.* the stripe

QTL). While cryptic variation may have played a role in the evolution of

some or all of these traits, our results suggest that it may not have been

pervasive with respect to the strongest effect changes.

**Conclusion**

While the genetic architecture underlying adaptive evolution may

vary somewhat depending upon the trait, there remains a key interest in

assessing potentially general patterns.  In combination with past studies, our

results suggest that most adaptive trait changes involve variants of non-

trivial effect size, and that these variants often do not initially reach

fixation.  Epistasis is an important component underlying complex traits,

including in flies (Huang *et al.* 2012), but we did not find evidence that

strong epistasis was relevant for the adaptive evolution of the studied

loci.  In light of the existence of other examples in which epistasis among

adaptive loci have been shown (Malmberg & Mauricio 2005), our study

highlights that the answer to the classic debate on whether epistasis is

important to natural selection might vary case by case.  Lastly, we found

evidence of gene-by-environment interactions underlying pigmentation,

stressing that the variability of genetic architectures can also be

environment-dependent, which is particularly relevant in light of rapidly

changing global environments.

**Data availability**: Source code used in the analyses is available on GitHub

(https://github.com/ribeirots/RILepi).  Raw sequence read data for the

Recombinant Inbred Lines are available on the Sequence Read Archive

(BioProject: PRJNA861300).

**Acknowledgments**: We thank members of the Pool lab for helpful

comments on this manuscript.

**Conflict of interest**: The authors declare no conflict of interest.

**Funding:** This research was funded by NIH grants R01 GM127480 and R35

GM136306.

## References

Abramoff, M. D., Magalhães, P. J., & Ram, S. J. (2004). Image processing with ImageJ. Biophotonics International, 11(7), 36–42.

Adams, M., McBroome, J., Maurer, N., Pepper-Tunick, E., Saremi, N. F., Green, R. E., Vollmers, C., & Corbett-Detig, R. B. (2020). One fly–one genome: chromosome-scale genome assembly of a single outbred *Drosophila melanogaster*. Nucleic Acids Research, 48(13), e75-e75. doi:10.1093/nar/gkaa450.

Ang, R. M. L., Chen, S. A. A., Kern, A. F., Xie, Y., & Fraser, H. B. (2023). Widespread epistasis among beneficial genetic variants revealed by high-throughput genome editing. Cell Genomics, 3(4). doi:10.1016/j.xgen.2023.100260.

Arthur, B. J., Ding, Y., Sosale, M., Khalif, F., Kim, E., Waddell, P., Turaga, S. C., & Stern, D. L. (2021). SongExplorer: A deep learning workflow for discovery and segmentation of animal acoustic communication signals (p. 2021.03.26.437280). bioRxiv. doi:10.1101/2021.03.26.437280.

Arthur, B. J., Sunayama-Morita, T., Coen, P., Murthy, M., & Stern, D. L. (2013). Multi-channel acoustic recording and automated analysis of *Drosophila* courtship songs. BMC Biology, 11(1), 11. doi:10.1186/1741-7007-11-11.

Bank, C. (2022). Epistasis and adaptation on fitness landscapes. Annual Review of Ecology, Evolution, and Systematics, 53, 457-479. doi:10.1146/annurev-ecolsys-102320-112153.

Barghi, N., Hermisson, J., & Schlötterer, C. (2020). Polygenic adaptation: a unifying framework to understand positive selection. Nature Reviews Genetics 21(12), 769-781. doi:10.1038/s41576-020-0250-z.

Barton, N. H. (2017). How does epistasis influence the response to selection? Heredity, 118(1), 96-109. doi:10.1038/hdy.2016.109.

Bastide, H., Lange, J. D., Lack, J. B., Yassin, A., & Pool, J. E. (2016). A variable genetic architecture of melanic evolution in *Drosophila melanogaster*. Genetics, 204(3), 1307–1319. doi:10.1534/genetics.116.192492.

Bastide, H., Yassin, A., Johanning, E. J., & Pool, J. E. (2014). Pigmentation in *Drosophila melanogaster* reaches its maximum in Ethiopia and correlates most strongly with ultra-violet radiation in sub-Saharan

Africa. BMC Evolutionary Biology, 14(1), 179. doi:10.1186/s12862-014-
0179-y.

Bejarano, F., Luque, C. M., Herranz, H., Sorrosal, G., Rafel, N., Pham, T. T.,
& Milán, M. (2008). A gain-of-function suppressor screen for genes
involved in dorsal–ventral boundary formation in the *Drosophila* wing.
Genetics, 178(1), 307-323.. http://dx.doi.org/10.1534/genetics.107.081869.

Blanco, E., Ruiz-Romero, M., Beltran, S., Bosch, M., Punset, A., Serras, F., &
Corominas, M. (2010). Gene expression following induction of
regeneration in *Drosophila* wing imaginal discs. Expression profile of
regenerating wing discs. BMC Developmental Biology, 10(1), 1-14.
http://dx.doi.org/10.1186/1471-213X-10-94.

Bomblies, K., & Peichel, C. L. (2022). Genetics of adaptation. Proceedings of
the National Academy of Sciences, 119(30), e2122152119.
doi:10.1073/pnas.2122152119.

Broman, K. W., Wu, H., Sen, Ś., & Churchill, G. A. (2003). R/qtl: QTL
mapping in experimental crosses. Bioinformatics, 19(7), 889–890.
doi:10.1093/bioinformatics/btg112.

Carter, A. J., Hermisson, J., & Hansen, T. F. (2005). The role of epistatic gene
interactions in the response to selection and the evolution of evolvability.
Theoretical Population Biology, 68(3), 179-196.
doi:10.1016/j.tpb.2005.05.002.

Clemens, J., Coen, P., Roemschied, F. A., Pereira, T. D., Mazumder, D.,
Aldarondo, D. E., Pacheco, D. A., & Murthy, M. (2018). Discovery of a
new song mode in *Drosophila* reveals hidden structure in the sensory
and neural drivers of behavior. Current Biology, 28(15), 2400-2412.e6.
doi:10.1016/j.cub.2018.06.011.

Cohan, F. M., & Graf, J.-D. (1985). Latitudinal cline in *Drosophila*
*melanogaster* for knockdown resistance to ethanol fumes and for rates of
response to selection for further resistance. Evolution, 39(2), 278–293.
doi:10.2307/2408362.

Cohan, F. M., Hoffmann, A. A., & Gayley, T. W. (1989). A test of the role of
epistasis in divergence under uniform selection. Evolution, 43(4), 766-
774. doi:10.2307/2409305.

Comeron, J. M., Ratnappan, R., & Bailin, S. (2012). The many landscapes of recombination in *Drosophila melanogaster*. PLOS Genetics, 8(10), e1002905. doi:10.1371/journal.pgen.1002905.

Corbett-Detig, R., & Nielsen, R. (2017). A hidden Markov model approach for simultaneously estimating local ancestry and admixture time using next generation sequence data in samples of arbitrary ploidy. PLoS Genetics, 13(1), e1006529. doi:10.1371/journal.pgen.1006529.

Crow, J. F. (2010). On epistasis: why it is unimportant in polygenic directional selection. Philosophical Transactions of the Royal Society B: Biological Sciences, 365(1544), 1241–1244. doi:10.1098/rstb.2009.0275.

da Silva Ribeiro, T., Galván, J. A., & Pool, J. E. (2022). Maximum SNP $F_{ST}$ outperforms full-window statistics for detecting soft sweeps in local adaptation. Genome Biology and Evolution, 14(10), evac143. doi:10.1093/gbe/evac143.

David, J. R., Capy, P., & Gauthier, J.-P. (1990). Abdominal pigmentation and growth temperature in *Drosophila melanogaster*: similarities and differences in the norms of reaction of successive segments. Journal of Evolutionary Biology, 3(5–6), 429–445. doi:10.1046/j.1420-9101.1990.3050429.x.

Dewey, E. M., McNabb, S. L., Ewer, J., Kuo, G. R., Takanishi, C. L., Truman, J. W., & Honegger, H. W. (2004). Identification of the gene encoding bursicon, an insect neuropeptide responsible for cuticle sclerotization and wing spreading. Current Biology, 14(13), 1208-1213.. http://dx.doi.org/10.1016/j.cub.2004.06.051.

Doust, A. N., Lukens, L., Olsen, K. M., Mauro-Herrera, M., Meyer, A., & Rogers, K. (2014). Beyond the single gene: How epistasis and gene-by-environment effects influence crop domestication. Proceedings of the National Academy of Sciences, 111(17), 6178-6183. doi:10.1073/pnas.1308940110.

Duchêne, E., Butterlin, G., Dumas, V., & Merdinoglu, D. (2012). Towards the adaptation of grapevine varieties to climate change: QTLs and candidate genes for developmental stages. Theoretical and Applied Genetics, 124(4), 623-635. doi:10.1007/s00122-011-1734-1.

Dworkin, I., & Gibson, G. (2006). Epidermal growth factor receptor and transforming growth factor-β signaling contributes to variation for wing shape in *Drosophila*

*melanogaster*. Genetics, 173(3), 1417-1431.
http://dx.doi.org/10.1534/genetics.105.053868.

Fenster, C. B., Galloway, L. F., & Chao, L. (1997). Epistasis and its
consequences for the evolution of natural populations. Trends in
Ecology & Evolution, 12(7), 282–286. doi:10.1016/S0169-5347(97)81027-0.

Fisher, R. A. (1930). The genetical theory of natural selection (pp. xiv, 272).
Clarendon Press. doi:10.5962/bhl.title.27468.

Flynn, J. M., Rossouw, A., Cote-Hammarlof, P., Fragata, I., Mavor, D.,
Hollins III, C., Bank, C., & Bolon, D. N. (2020). Comprehensive fitness
maps of Hsp90 show widespread environmental dependence. Elife, 9,
e53810. doi:10.7554/eLife.53810.

Fochler, S., Morozova, T. V., Davis, M. R., Gearhart, A. W., Huang, W.,
Mackay, T. F., & Anholt, R. R. (2017). Genetics of alcohol consumption in
*Drosophila melanogaster*. Genes, Brain and Behavior, 16(7), 675-685.
doi:10.1111/gbb.12399.

Fry, J. D., Nuzhdin, S. V., Pasyukova, E. G., & Mackay, T. F. (1998). QTL
mapping of genotype–environment interaction for fitness in *Drosophila*
*melanogaster*. Genetics Research, 71(2), 133-141.
doi:10.1017/S0016672398003176.

Fuhrmann, N., Prakash, C., & Kaiser, T. S. (2023). Polygenic adaptation from
standing genetic variation allows rapid ecotype formation. Elife, 12,
e82824. doi:10.7554/eLife.82824.

Ghalambor, C. K., McKay, J. K., Carroll, S. P., & Reznick, D. N. (2007).
Adaptive versus non-adaptive phenotypic plasticity and the potential for
contemporary adaptation in new environments. Functional Ecology,
21(3), 394–407. doi:10.1111/j.1365-2435.2007.01283.x.

Gibson, G., & Dworkin, I. (2004). Uncovering cryptic genetic variation.
Nature Reviews Genetics 5, 681–690. doi:10.1038/nrg1426.

Gong, L. I., & Bloom, J. D. (2014). Epistatically interacting substitutions are
enriched during adaptive protein evolution. PLoS Genetics, 10(5).
doi:10.1371/journal.pgen.1004328.

Griswold, C. (2006). Gene flow's effect on the genetic architecture of a local
adaptation and its consequences for QTL analyses. Heredity, 96, 445-453.
doi:10.1038/sj.hdy.6800822.

Hansen, T. F., Álvarez-Castro, J. M., Carter, A. J., Hermisson, J., & Wagner, G. P. (2006). Evolution of genetic architecture under directional selection. Evolution, 60(8), 1523-1536. doi:10.1111/j.0014-3820.2006.tb00498.x.

Hansen, T. F. (2013). Why epistasis is important for selection and adaptation. Evolution, 67(12), 3501–3511. doi:10.1111/evo.12214.

Hill, W. G., Goddard, M. E., & Visscher, P. M. (2008). Data and theory point to mainly additive genetic variance for complex traits. PLOS Genetics, 4(2), e1000008. doi:10.1371/journal.pgen.1000008.

Hirai, Y., & Kimura, M. T. (1997). Incipient reproductive isolation between two morphs of *Drosophila elegans* (Diptera: Drosophilidae). Biological Journal of the Linnean Society, 61(4), 501-513. doi:10.1111/j.1095-8312.1997.tb01804.x.

Huang, Y., Lack, J. B., Hoppel, G. T., & Pool, J. E. (2021). Parallel and population-specific gene regulatory evolution in cold-adapted fly populations. Genetics, 218(3). doi:10.1093/genetics/iyab077.

Huang, W., Richards, S., Carbone, M. A., Zhu, D., Anholt, R. R. H., Ayroles, J. F., Duncan, L., Jordan, K. W., Lawrence, F., Magwire, M. M., Warner, C. B., Blankenburg, K., Han, Y., Javaid, M., Jayaseelan, J., Jhangiani, S. N., Muzny, D., Ongeri, F., Perales, L., … Mackay, T. F. C. (2012). Epistasis dominates the genetic architecture of *Drosophila* quantitative traits. Proceedings of the National Academy of Sciences, 109(39), 15553–15559. doi:10.1073/pnas.1213423109.

Juenger, T. E., Sen, S., Stowe, K. A., & Simms, E. L. (2005). Epistasis and genotype-environment interaction for quantitative trait loci affecting flowering time in *Arabidopsis thaliana*. Genetica, 123, 87-105. doi:10.1007/s10709-003-2717-1.

Krishnan, S. N., Frei, E., Swain, G. P., & Wyman, R. J. (1993). Passover: a gene required for synaptic connectivity in the giant fiber system of *Drosophila*. Cell, 73(5), 967-977. doi:10.1016/0092-8674(93)90274-T.

Krupp, J. J., Yaich, L. E., Wessells, R. J., & Bodmer, R. (2005). Identification of genetic loci that interact with cut during *Drosophila* wing-margin development. Genetics, 170(4), 1775-1795. http://dx.doi.org/10.1534/genetics.105.043125.

Lack, J. B., Cardeno, C. M., Crepeau, M. W., Taylor, W., Corbett-Detig, R. B., Stevens, K. A., Langley, C. H., & Pool, J. E. (2015). The *Drosophila*

genome nexus: a population genomic resource of 623 *Drosophila melanogaster* genomes, including 197 from a single ancestral range population. Genetics, 199(4), 1229–1241. doi:10.1534/genetics.115.174664.

Lack, J. B., Monette, M. J., Johanning, E. J., Sprengelmeyer, Q. D., & Pool, J. E. (2016). Decanalization of wing development accompanied the evolution of large wings in high-altitude *Drosophila*. Proceedings of the National Academy of Sciences, 113(4), 1014–1019. doi:10.1073/pnas.1515964113.

Lande, R. 1992. Neutral theory of quantitative genetic variance in an island model with local extinction and colonization. Evolution, 46:381–389. doi:10.1111/j.1558-5646.1992.tb02046.x.

Lange, J. D., & Pool, J. E. (2016). A haplotype method detects diverse scenarios of local adaptation from genomic sequence variation. Molecular Ecology, 25(13), 3081-3100. doi:10.1111/mec.13671.

Lea, A. J., Peng, J., & Ayroles, J. F. (2022). Diverse environmental perturbations reveal the evolution and context-dependency of genetic effects on gene expression levels. Genome Research, 32(10), 1826-1839.https://www.genome.org/cgi/doi/10.1101/gr.276430.121.

Li, H. (2013). Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. arXiv:1303.3997 [q-Bio]. http://arxiv.org/abs/1303.3997.

Lollar, M. J., da Silva Ribeiro, T., Stern. D., & Pool, J. E. (2025). Quantitative trait mapping of courtship song variation between male *Drosophila melanogaster* from two allopatric populations. biorxiv, 2025-02.

López-Varea, A., Vega-Cuesta, P., Ruiz-Gómez, A., Ostalé, C. M., Molnar, C., Hevia, C. F., Martín, M., Organista, M. F., de Celis, J., Culí, J., Esteban, N., & de Celis, J. F. (2021). Genome-wide phenotypic RNAi screen in the *Drosophila* wing: Phenotypic description of functional classes. G3, 11(12), jkab349. doi:10.1093/g3journal/jkab349.

Lowry, D. B., Lovell, J. T., Zhang, L., Bonnette, J., Fay, P. A., Mitchell, R. B., Lloyd-Reilley, J., Boe, A. R., Wu, Y., Rouquette Jr, F. M., Wynia, R. L., Weng, X., Behrman, K. D., Healey, A., Barry, K., Lipzen, A., Bauer, D., Sharma, A., Jenkins, J., Schmutz, J., Fritschi, F. B., & Juenger, T. E. (2019). QTL × environment interactions underlie adaptive divergence in switchgrass across a large latitudinal gradient. Proceedings of the

National Academy of Sciences, 116(26), 12933-12941. doi:10.1073/pnas.1821543116.

Lunzer, M., Golding, G. B., & Dean, A. M. (2010). Pervasive cryptic epistasis in molecular evolution. PLoS Genetics, 6(10). doi:10.1371/journal.pgen.1001162.

Mackay, T. F. (2001). The genetic architecture of quantitative traits. Annual Review of Genetics, 35(1), 303-339. doi:10.1146/annurev.genet.35.102401.090633.

Mackay, T. F. C. (2014). Epistasis and quantitative traits: Using model organisms to study gene–gene interactions. Nature Reviews Genetics, 15(1), 22–33. doi:10.1038/nrg3627.

Malmberg, R. L., & Mauricio, R. (2005). QTL-based evidence for the role of epistasis in evolution. Genetics Research, 86(2), 89–95. doi:10.1017/S0016672305007780.

Martin, G., & Lenormand, T. (2006). The fitness effect of mutations across environments: A survey in light of fitness landscape models. Evolution, 60(12), 2413-2427. doi:10.1111/j.0014-3820.2006.tb01878.x.

Massey, J. H., & Wittkopp, P. J. (2016). The genetic basis of pigmentation differences within and between *Drosophila* species. Current Topics in developmental Biology, 119, 27-61. doi:10.1016/bs.ctdb.2016.03.004.

Mathews, K. L., Malosetti, M., Chapman, S., McIntyre, L., Reynolds, M., Shorter, R., & Van Eeuwijk, F. (2008). Multi-environment QTL mixed models for drought stress adaptation in wheat. Theoretical and Applied Genetics, 117, 1077-1091. doi:10.1007/s00122-008-0846-8.

McGuigan, K., & Sgro, C. M. (2009). Evolutionary consequences of cryptic genetic variation. Trends in Ecology & Evolution, 24(6), 305-311. doi:10.1016/j.tree.2009.02.001.

Okada, H., Ebhardt, H. A., Vonesch, S. C., Aebersold, R., & Hafen, E. (2016). Proteome-wide association studies identify biochemical modules associated with a wing-size phenotype in *Drosophila melanogaster*. Nature Communications, 7(1), 12649. http://dx.doi.org/10.1038/ncomms12649.

Orr, H. A. (1998). The population genetics of adaptation: the distribution of factors fixed during adaptive evolution. Evolution, 52(4), 935–949. doi:10.1111/j.1558-5646.1998.tb01823.x.

Orr, H. A., & Betancourt, A. J. (2001). Haldane's sieve and adaptation from the standing genetic variation. Genetics, 157(2), 875-884. doi:10.1093/genetics/157.2.875.

Papa, R., Kapan, D. D., Counterman, B. A., Maldonado, K., Lindstrom, D. P., Reed, R. D., Nijhout, H. F., Hrbek, T., & McMillan, W. O. (2013). Multi-allelic major effect genes interact with minor effect QTLs to control adaptive color pattern variation in *Heliconius erato*. PLoS One, 8(3), e57033. doi:10.1371/journal.pone.0057033.

Parker, J., & Struhl, G. (2015). Scaling the *Drosophila* wing: TOR-dependent target gene access by the Hippo pathway transducer Yorkie. PLoS Biology, 13(10), e1002274.. http://dx.doi.org/10.1371/journal.pbio.1002274.

Paul, L., Wang, S. H., Manivannan, S. N., Bonanno, L., Lewis, S., Austin, C. L., & Simcox, A. (2013). Dpp-induced Egfr signaling triggers postembryonic wing development in *Drosophila*. Proceedings of the National Academy of Sciences, 110(13), 5058-5063. http://dx.doi.org/10.1073/pnas.1217538110.

Pitchers, W., Nye, J., Márquez, E. J., Kowalski, A., Dworkin, I., & Houle, D. (2019). A multivariate genome-wide association study of wing shape in Drosophila melanogaster. Genetics, 211(4), 1429-1447. doi:10.1534/genetics.118.301342.

Pool, J. E. (2016). Genetic mapping by bulk segregant analysis in Drosophila: experimental design and simulation-based inference. Genetics, 204(3), 1295-1306. https://doi.org/10.1534/genetics.116.192484.

Pool, J. E., Braun, D. T., & Lack, J. B. (2017). Parallel evolution of cold tolerance within *Drosophila melanogaster*. Molecular Biology and Evolution, 34(2), 349–360. doi:10.1093/molbev/msw232.

Pritchard, J. K., Pickrell, J. K., & Coop, G. (2010). The genetics of human adaptation: hard sweeps, soft sweeps, and polygenic adaptation. Current Biology, 20(4), R208-R215. doi:10.1016/j.cub.2009.11.055.

Reed, L. K., Lee, K., Zhang, Z., Rashid, L., Poe, A., Hsieh, B., Deighton, N., Glassbrook, N., Bodmer, R. & Gibson, G. (2014). Systems genomics of metabolic phenotypes in wild-type *Drosophila melanogaster*. Genetics, 197(2), 781-793. doi:10.1534/genetics.114.163857.

Reynolds, J., Weir, B. S., & Cockerham, C. C. (1983). Estimation of the
coancestry coefficient: basis for a short-term genetic distance. Genetics,
105(3), 767-779. doi:10.1093/genetics/105.3.767.

Rogers, W. A., Grover, S., Stringer, S. J., Parks, J., Rebeiz, M., & Williams, T.
M. (2014). A survey of the trans-regulatory landscape for *Drosophila*
*melanogaster* abdominal pigmentation. Developmental Biology, 385(2),
417-432. doi:10.1016/j.ydbio.2013.11.013.

Rutherford, S. L., & Lindquist, S. (1998). Hsp90 as a capacitor for
morphological evolution. Nature, 396(6709), 336-342. doi:10.1038/24550.

Savolainen, O., Lascoux, M., & Merilä, J. (2013). Ecological genomics of local
adaptation. Nature Reviews Genetics, 14, 807-820. doi:10.1038/nrg3522.

Schertel, C., Albarca, M., Rockel-Bauer, C., Kelley, N. W., Bischof, J., Hens,
K., van Nimwegen, E., Basier, K., & Deplancke, B. (2015). A large-scale, in
vivo transcription factor screen defines bivalent chromatin as a key
property of regulatory factors mediating *Drosophila* wing development.
Genome Research, 25(4), 514-523. doi:10.1101/gr.181305.114.

Schlötterer, C. (2023). How predictable is adaptation from standing genetic
variation? Experimental evolution in *Drosophila* highlights the central
role of redundancy and linkage disequilibrium. Philosophical
Transactions of the Royal Society B, 378(1877), 20220046.
doi:10.1098/rstb.2022.0046.

Ségurel, L., & Bon, C. (2017). On the evolution of lactase persistence in
humans. Annual review of genomics and human genetics, 18(1), 297-319.
doi:10.1146/annurev-genom-091416-035340.

Spitze, K. (1993). Population structure in *Daphnia obtusa*: quantitative
genetic and allozymic variation. Genetics. 135:367–374.
doi:10.1093/genetics/135.2.367

Sprengelmeyer, Q. D., Lack, J. B., Braun, D. T., Monette, M. J., & Pool, J. E.
(2022). The evolution of larger size in high-altitude Drosophila
melanogaster has a variable genetic architecture. G3, 12(3), jkab454.
doi:10.1093/g3journal/jkab454.

Sprengelmeyer, Q. D., Mansourian, S., Lange, J. D., Matute, D. R., Cooper, B.
S., Jirle, E. V., Stensmyr, M. C., & Pool, J. E. (2020). Recurrent collection
of *Drosophila melanogaster* from wild African environments and

genomic insights into species history. Molecular Biology and Evolution, 37(3), 627–638. doi:10.1093/molbev/msz271.

Sprengelmeyer, Q. D., & Pool, J. E. (2021). Ethanol resistance in *Drosophila melanogaster* has increased in parallel cold-adapted populations and shows a variable genetic architecture within and between populations. Ecology and Evolution, 11(21), 15364–15376. doi:10.1002/ece3.8228.

Starr, T. N., & Thornton, J. W. (2016). Epistasis in protein evolution. Protein Science, 25(7), 1204-1218. doi:10.1002/pro.2897.

Steiner, C. C., Weber, J. N., & Hoekstra, H. E. (2007). Adaptive variation in beach mice produced by two interacting pigmentation genes. PLoS Biology, 5(9), e219. doi:10.1371/journal.pbio.0050219.

Stitzer, M. C., & Ross-Ibarra, J. (2018). Maize domestication and gene interaction. New Phytologist, 220(2), 395-408. doi:10.1111/nph.15350.

Torgeman, S., & Zamir, D. (2023). Epistatic QTLs for yield heterosis in tomato. Proceedings of the National Academy of Sciences, 120(14), e2205787119. doi:10.1073/pnas.2205787119.

Trindade, S., Sousa, A., Xavier, K. B., Dionisio, F., Ferreira, M. G., & Gordo, I. (2009). Positive epistasis drives the acquisition of multidrug resistance. PLOS Genetics, 5(7), e1000578. doi:10.1371/journal.pgen.1000578.

Van Rossum, G., & Drake, F. L. (2009). Python 3 Reference Manual. Scotts Valley, CA: CreateSpace.

Vijendravarma, R. K., & Kawecki, T. J. (2013). Epistasis and maternal effects in experimental adaptation to chronic nutritional stress in *Drosophila*. Journal of Evolutionary Biology, 26(12), 2566-2580. doi:10.1111/jeb.12248.

Whitlock, M. C., Phillips, P. C., Moore, F. B.-G., & Tonsor, S. J. (1995). Multiple fitness peaks and epistasis. Annual Review of Ecology and Systematics, 26(1), 601–629. doi:10.1146/annurev.es.26.110195.003125.

Willmer, P. G., & Unwin, D. M. (1981). Field analyses of insect heat budgets: reflectance, size and heating rates. Oecologia, 50, 250-255. doi:10.1007/BF00348047.

Wright, S. (1931). Evolution in Mendelian populations. Genetics, 16(2), 97–159. doi:10.1093%2Fgenetics%2F16.2.97.

Yeaman, S. & Otto, S. P. (2011). Establishment and maintenance of adaptive
genetic divergence under migration, selection, and drift. Evolution, 65(7),
2123–2129. doi:10.1111/j.1558-5646.2011.01277.x.

Yeaman, S., & Whitlock, M. C. (2011). The genetic architecture of adaptation
under migration-selection balance. Evolution, 65(7), 1897–1911.
doi:10.1111/j.1558-5646.2011.01269.x.

Yukilevich, R., Lachance, J., Aoki, F., & True, J. R. (2008). Long-term
adaptation of epistatic genetic networks. Evolution, 62(9), 2215-2235.
doi:10.1111/j.1558-5646.2008.00445.x.

Zeng, Z. B., Kao, C. H., & Basten, C. J. (1999). Estimating the genetic
architecture of quantitative traits. Genetics Research, 74(3), 279-289.
doi:10.1017/S0016672399004255.

Zhang, F., Xie, D., Liang, M., & Xiong, M. (2016). Functional regression
models for epistasis analysis of multiple quantitative traits. PLoS
genetics, 12(4), e1005965. doi:10.1371/journal.pgen.1005965.

Zheng, Y., Liu, B., Wang, L., Lei, H., Prieto, K. D. P., & Pan, D. (2017).
Homeostatic control of Hpo/MST kinase activity through
autophosphorylation-dependent recruitment of the STRIPAK PP2A
phosphatase complex. Cell Reports, 21(12), 3612-3623.
doi:10.1016/j.celrep.2017.11.076.

Zhou, S., Campbell, T. G., Stone, E. A., Mackay, T. F., & Anholt, R. R. (2012).
Phenotypic plasticity of the *Drosophila* transcriptome. PLoS genetics,
8(3), e1002593. doi:10.1371/journal.pgen.1002593.