Courtship song differs between African and European populations of *Drosophila*

***melanogaster* and involves a strong effect locus**

Matthew J. Lollar[1,@], Elizabeth Kim[2], David L. Stern[2], John E. Pool[1,@]

[1]Laboratory of Genetics, University of Wisconsin – Madison, Madison, WI, 53705, USA

[2]Janelia Research Campus, Howard Hughes Medical Institute, Ashburn, VA, 20147

USA

Corresponding author contact: mlollar@wisc.edu; jpool@wisc.edu

Running head: Genetics of song evolution within *D. melanogaster*

Keywords: courtship song, behavior, QTL mapping, variation, evolution

ABSTRACT

The courtship song of *Drosophila melanogaster* has long served as excellent model

system for studies of animal communication and differences in courtship song have

been demonstrated among populations and between species. Here, we report that flies

of African and European origin, which diverged approximately 13,000 years ago, show

significant genetic differentiation in the use of slow versus fast pulse song. Using a

combination of quantitative trait mapping and population genetic analysis we detected a

single strong QTL underlying this trait and we identified candidate genes that may

contribute to the evolution of this trait. Song trait variation between parental strains of

our recombinant inbred panel enabled detection of genomic intervals associated with six

additional song traits, some of which include known courtship-related genes. These

findings improve the prospects for further genetic insights into the evolution of

reproductive behavior and the biology underlying courtship song.

29 INTRODUCTION

Communication, the process of information transfer through signal production and

recognition, is abundant across the animal kingdom. There are many forms of

communication in nature, from chemical interactions between bacteria (Keller & Surette

2006), to intricate mating displays of suboscine passerines (Prum 1990), or the

complexity of human language (Lieberman 1971). Accompanying this diversity of form

is a diversity of purpose: communication is used by animals in mate choice and

conspecific recognition (Anholt *et al.* 2020), antagonistic displays (Maynard Smith

1979), and cooperative behavior such as territory defense and foraging (Fitzgerald &

Peterson 1988). Often, the proper transmission of information from signaler to receiver

has important consequences for the fitness and reproductive success of an individual

(Maynard Smith 1979; Anholt *et al.* 2020), underpinning its importance in both standing

diversity and the evolutionary history of populations. Acoustic communication has

evolved independently over 20 times across insects (Greenfield 2016) and is an

important component of reproductive behavior (Leonhardt *et al.* 2016).

Male *Drosophila melanogaster* "sing" to females by vibrating their wings, typically as

unilateral extension of a single outstretched wing (Ewing & Bennet-Clark 1968). In *D.*

melanogaster, song consists of two primary modes; a series of continuous sinusoidal

waves or hums (sine song) and trains of more punctuated wave clusters (pulse song)

with amplitude and pitch typically exceeding those produced by sine song (Swain &

Philipsborn 2021). Furthermore, two modes of pulse song have been identified based

on the pitch of the pulse song: fast and slow pulse modes (Clemens *et al.* 2018).

Female *Drosophila* typically become more receptive to mating upon hearing song (von

Schilcher 1976) and may distinguish between subtle differences in song components.

The active female responses to courtship that encourage or discourage copulation

underpin the role of song in the reproductive success of male flies (Cook 1973;

Riabinina *et al*. 2011; Laturney & Moehring 2012).

Male courtship song has been characterized in many *Drosophila* species, (O'Grady &

DeSalle 2018), providing deep historical knowledge in a system benefiting from

excellent genetic tractability (St. Johnston 2002; *Drosophila* 12 Genomes Consortium

2007). The acoustic properties of pulse song, such as the interpulse interval (IPI)

typically receive the greatest attention (Ritchie *et al.* 1994; Talyn & Dowse 2004; Stern

2014), although components of sine song also vary among closely related species (Ding

et al. 2016). Pulse song is highly variable across species and has been implicated in

mate and conspecific recognition (Spieth 1952; Bennet-Clark & Ewing 1969; Watanabe

& Kawanishi 1979; Cowling & Burnet 1981; Talyn & Dowse 2004; Ding *et al.* 2019),

although song does may not contribute to mate choice in every species (Richie *et al.*

1999; Iglesias & Hasson 2017).

The genetic architectures and underpinnings of variation in courtship song and other

premating isolating mechanisms are poorly understood. Early studies supported the

view that variation in these behaviors was caused by a small number of loci of large

effect (Mackay 2001; Mackay *et al.* 2005). Arbuthnott (2009) found, for instance, that

among 36 studies in insects (predominantly *Drosophila)* on courtship traits contributing

to premating isolating barriers, 25 studies considered traits to be influenced by a few

loci of large effect. However, quantitative trait locus (QTL) studies are likely biased

toward detecting large effect loci (Rockman 2012). More recent studies have revealed

that the genetic architecture of premating barriers is likely more complex (Mackay 2009;

Arbuthnott 2009; Oh *et al.* 2012; Shahandeh & Turner 2020). The number of loci

contributing to variation in a trait involved in premating isolation can also have

implications for the rate of speciation between species (Gavrilets & Vose 2007),

providing additional incentive to study the evolution of such phenotypes.

Differences between well-isolated species in reproductive behavior may or may not

have contributed to the intial reproductive isolation of those species (Price 2002). On

the other hand, within-population variation in such traits may be distinct in its genetic

architecture and basis from evolutionary trait changes (Arbuthnott 2009). Whereas,

evolved differences in courtship traits between interfertile taxa, including recently

diverged populations of the same species, may hold particular interest for the study of

reproductive isolation.

D. melanogaster originated in sub-Saharan Africa and has spread nearly worldwide

(Lachaise & Silvain 2004, Lachaise *et al.* 1988, Keller 2007). The demographic history

of the species is well-studied (e.g. Sprengelmeyer *et al.* 2020), and striking genetic and

phenotypic changes have occurred on a relatively short evolutionary time-scale. Genetic

differentiation is in part due to at least one moderate population bottleneck during the

initial migration out of sub-Saharan Africa (Begun & Aquadro 1993, Sprengelmeyer *et*

al. 2020), as well as locally adaptive changes in response to novel environmental

pressures (David & Capy 1988, Fabian *et al.* 2015; Lack *et al.* 2016a). Genes under

selection as a consequence of adaptation often have neuronal or morphological and

developmental functions (Fabian *et al.* 2015; Lack *et al.* 2016a; Pool *et al.* 2017). Such

changes could have either specific or pleiotropic effects on courtship traits such as

song, in light of the multifaceted roles of many genes involved in courtship behavior

(Gleason 2005; Greenspan & Ferveur 2000).

While some components of courtship song variation within *D. melanogaster* have been

studied (Colegrave *et al.* 1999; Gleason *et al.* 2002), there have been no prior

comprehensive intraspecific studies of many courtship song components. Here, we

utilize panels of naturally-derived *D. melanogaster* from two geographic origins, Zambia

and France, and high-throughput methods of recording and analyzing courtship song

(Arthur *et al.* 2013; Arthur *et al.* 2021) to study the genetic basis of song differences

between populations. We find that the proportion of slow and fast pulses ("pulse mode

preference") differs significantly between populations. We use a panel of recombinant

inbred lines (RILs) derived from reciprocal crosses between France and Zambia strains

to map pulse mode preference as a quantitative trait, as well as mapping eight

additional song traits that differ significantly between RIL parental strains. Finally we

complement our mapping analysis with population genetic screens across QTL

intervals, highlighting the most genetically differentiated loci between populations to

identify additional genes that may underlie song variation and evolution. Collectively,

our results reveal that naturally evolved differences in male song components exist

between populations within the *D. melanogaster* species and include moderately strong

effect loci, opening the door to targeted genetic and neurophysiological studies of their

molecular basis.

MATERIALS AND METHODS

Drosophila husbandry and stocks

Two distinct types of strains were used in this study. Strains used to measure trait

differences between population samples were (8 generation) inbred lines from wild-

derived isofemale stocks, collected at locations described in Pool *et al.* (2012). Strains

used to map quantitative traits represented a recombinant inbred line (RIL) panel

generated in the lab and derived from a reciprocal cross between two inbred strains,

one each from our France (FR) and Zambia (ZI) collections (FR320N and ZI418N).

Following 12 generations of intercrossing, isofemale lines were established from these

interbred flies and inbred for 5 generations to establish the final panel, from which 305

recombinant strains were included in the present study. Females used in mating assays

were derived from a single France strain, FR54N. We chose this strain beacuse

European females show less preference across populations compared to some

southern African strains (Grillet *et al*. 2012; Hollocher & Wu, 1996; Wu *et al*. 1995), who

tend to disfavor non-African males, and female response can influence the expression

of song (Riabinina *et al*. 2011; Laturney & Moehring 2012).

Flies were maintained as stocks in laboratory conditions on food prepared in batches of

4.5 L water, 500 mL cornmeal, 500 mL molasses, 200 mL yeast, 54 g agar, 20 mL

propionic acid, and 45 mL tegosept 10% (in 95% ethanol). Flies used in assays were

collected under carbon dioxide anesthesia and aged in one of two 12 hour light/dark

controlled incubators, separated in cycle by two real time hours to allow additional

assays each day. Incubator temperatures were not strictly controlled but measured daily

and ranged from 23-25C. Males were collected and aged individually in glass fly food

vials. Females were collected and aged in plastic fly food bottles containing

approximately 50 females per bottle.

Song recording

Male song was recorded on a multi-channel acoustic recording chamber (Arthur et al.

2013), at Janelia Research Campus from June-July of 2022 (Table S1). Males used in

assays were collected 0-24 hours post-eclosion, isolated into individual vials, and aged

for 3-5 days. Males were recorded in the presence of a single unmated female,

collected 0-6 hours post-eclosion, and aged an additional 12-24 hours before use in

assays. This age was selected as an optimal range to allow aging of females past a

"young nymph" stage, without allowing them to become old enough to be receptive to

mating (Manning 1967), which would diminish the duration of captured song before

copulation (observationally, males that initiated song typically sang throughout the entire

duration of recording if they did not mount successfully). Flies were mouth aspirated into

recording chambers, and assays were performed within the first two hours from lights

on. Male flies were recorded for a consecutive duration of 30 minutes. Replicate assays

were performed such that strains were assayed in sets of 6-8 males across 3-5 unique

days, for a total target sample size of 24 for each RIL, and 50-60 for each inbred strain

(Table S1).

Song analysis

Song data was automatically segmented and analyzed without human intervention

using SongExplorer (doi:10.1101/2021.03.26.437280) to annotate acoustic components

of song. A previously trained *D.melanogaster* model detected four principle sounds

(sine song, pulse song, ambient noise, other), trained using audio collected on the

recording instrumentation used in this study. Samples were recorded at a sampling rate

twice that of the provided model, so raw audio files were downsampled at a rate of one

half using the "signal" function in the python open-source package SciPy (version 1.13)

for proper segmentation (Virtanen *et al.* 2020).

Segmented   song   parameters   were   quantified   using   BatchSongAnalysis

(https://github.com/dstern/BatchSongAnalysis)   with   small   modifications   for

individualized trait parameterization (https://github.com/mjlollar/song_work), MATLAB

version 9.13 (R2022b) (The MathWorks Inc. 2022). The numbers of song bouts, sine

and pulse song durations, interpulse interval characteristics, pulse/sine carrier

frequencies and pulse/sine train lengths were measured as described in Arthur *et al.*

(2013). Pulse and sine song amplitude were measured by calculating the square root of

the mean of the squares of all pulses in a song (output as arbitrary units). Transition

probabilities were calculated using the Matlab "transProb()" function for pairwise

measurements of events (The MathWorks Inc. 2022). Slow and fast pulse

characterization was available thanks to previous classification training in the model

provided and used methods similar to those described in Clemens *et al.* (2018) to

differentiate and classify pulse modes. Interpulse intervals were estimated

independently of pulse carrier frequency by fitting an envelope to each pulse to estimate

pulse duration and calculating the interpulse interval as the time from the end of one

pulse to the center of the next pulse. Here, pulse event carrier frequency was measured

as the modal frequency detected by SongExplorer. Further discussion on the

characterization of carrier frequency can be found in Ding *et al.* (2019).

Replicates with fewer than one second of song per minute, or those with an aberrantly

high sine song rate per minute (recordings with >55 seconds of characterized sine song

appeared to represent a failure of recording equipment) were excluded from analysis.

Additionally, replicates with zero values for a trait were not considered when calculating

strain mean for that trait, as a zero value from BatchSongAnalysis was typically the

result of an inability to properly measure trait values (due to minimal sampling

throughout the assay, etc.). Trait values for strains were represented as the mean value

of the total remaining replicates.

Differences in the population mean trait values were quantified using Mann-Whitney

(MW) tests using the two-sample wilcox.test() function in R (R Core Team

2023)(https://github.com/mjlollar/song_work). To statistically control for the testing of

multiple traits, we used a permutation-based approach to correct for multiple testing that

accounted for non-independence among some traits (Figure S1). 10000 permutations

were performed in the following design. For each replicate, the identities of each strain

were randomly shuffled and drawn without replacement to represent 10 FR strains and

7 ZI strains (matching sample sizes of the empirical data). MW tests were calculated

between these groups and recorded. We then determined the false positive rate (the

likelihood the empirical MW p-value occurs by chance) as the proportion of permutation

replicates that gave a p-value as low or lower than the empirical value. Correlation

statistics in Supplementary Figure 1 were generated in R using Pearson's test (R Core

Team 2023).

For a trait showing evidence of population differentiation from the above methodology,

we also calculated Q_{ST} as a measure of quantitative trait differentiation (Lande 1992;

Spitze 1993). To indicate whether this trait might fall outside the range of neutral

population differentiation, we compared the estimated Q_{ST} to the distribution of F_{ST} at

individual SNPs (which conservatively maximizes the variance of F_{ST}), calculated from

allelic sample sizes identical to the number of strains from which phenotypic data was

obtained.

QTL mapping based on RIL ancestries

Genomic sequencing of the RIL panel is described in da Silva Ribeiro *et al.* (2024),

which also documents the calling of parental strain ancestry. Briefly, for each RIL,

homozygous and heterozygous parental strain ancestry calls were assigned using

Ancestry HMM (Corbett-Detig & Nielsen 2017) on windows defined by intervals

containing 2,500 SNP genotype differences between the two parental strains (which

were used as the parental panels).

QTL mapping was performed using parental strain ancestry calls as marker genotypes.

We used a modified version of r/QTL (Broman *et al.* 2003) that allowed us to use an

additive QTL model, rather than allowing each genotypic class to have an independent

mean trait value (da Silva Ribeiro *et al.* 2024), in light of the smaller counts of some

ancestry genotypes observed in many genomic windows. The primary model tests the

alternative hypothesis of a single QTL against the null hypothesis of no QTL, while

using phenotypic permutations to control for genome-wide multiple testing. More explicit

details of the pipeline can be found in da Silva Ribeiro *et al.* (2024), as well as source

code at https://github.com/ribeirots/RILepi.

The above analysis yields a genomic LOD score landscape for any given song trait.

However, when this landscape is complex or ragged in its topography, in practice it can

be difficult to assess which local peaks to consider as distinct QTLs. We therefore

implemented a series of criteria to determine which local peaks to integrate into the

same QTL region, and also to denote which regions within a broad/complex QTL may

have the greatest likelihood of containing a causative locus. Preliminary QTL peak

regions were defined based on contiguous runs of windows above the genome-wide

LOD significance threshold. A pair of initially separate peaks separated by a valley with

a minimum LOD value of greater than the shorter peak's maximum LOD score minus

1.5 were merged into the same peak region - since a simple confidence interval (CI) for

the shorter peak, based on a 1.5 LOD drop, would end up including the taller peak. After

defining and integrating peaks in this manner, we accounted for peaks with complex

LOD landscapes by differentiating between "inclusive" and "restrictive" CIs. An inclusive

CI was defined as the full span of the windows within a contiguous run of windows

above the LOD significance threshold in which the local LOD score was greater than the

region's maximum LOD score minus 1.5. A QTL region's inclusive CI was therefore

contiguous by definition, and could encompass some windows with LOD scores less

than the maximum LOD minus 1.5. In contrast, restrictive CIs were defined as only the

subset of windows within the inclusive CI that had LOD scores greater than the

maximum LOD minus 1.5 (https://github.com/mjlollar/song_work). A QTL region's

restrictive CI could therefore encompass multiple non-contiguous runs of windows

representing different LOD sub-peaks within the same QTL region as defined above.

Based on the LOD scores of their constituent windows, restrictive CIs were considered

to be the most likely intervals to contain a causative locus, and were therefore the focus

of downstream analysis as described below. Windows yielding a RIL panel-wide

ancestry proportion above 90% for either parental strain were excluded from the

mapping analysis, and confidence intervals (CIs) encountering such regions were

extended until encountering an included window that ended the CI based on the criteria

detailed above.

Population Genetic and Candidate Gene Analysis of QTL regions

Previous population genomic data available from the *Drosophila* Genome Nexus (Lack

et al. 2016a) was used to generate a population genetics database. We used all

available Zambia and France genomes for a total of 197 and 96 strains, respectively.

Under the hypothesis that the two population-differentiated song traits may have been

influenced by population-specific selection, we assessed three window-based

population genetic statistics across the restrictive CIs of each QTL region for those

traits: 1) $F_{ST\_FullWin,}$ which represents the estimated F_{ST} value of a window using all

variable sites within the window; 2) F_{ST_MaxSNP}, the maximum F_{ST} value of any variable

site within a window (da Silva Ribeiro *et al.* 2022); and 3) Comparative Haplotype

Identity (χ_{MD}), which tests for evidence of local adaptation by comparing the summed

lengths of pairwise identical haplotypes within one population versus another (Lange &

Pool 2016).

We then identified "candidate genes" within QTL CIs (using the restrictive CI definition

above, including only the windows within an extended CI that had LOD scores greater

than the peak value minus 1.5). We associated genes with windows if any of their exons

(coding or non-coding) overlapped a given window, or if an exon of this gene was the

first exon to the left or right of this window's boundaries (in order to account for potential

regulatory regions of a given gene). We included as candidates any genes that had

been previously implicated in courtship behavior. A list of 92 such genes was generated

using genes sourced from the Gene Ontology database category "male mating

behavior" (ID GO:0060179) as well as additional genes identified from cursory literature

searches (Table S2). For traits with evidence of population differentiation, we also

included as candidates (1) genes with a documented role in wing development or male-

biased neuronal expression, and (2) genes with evidence for local adaptation, in that

they were associated with at least one window that fell within the top 1% quantile of

values for either $F_{ST\_FullWin}$, F_{ST_MaxSNP}, or X_{MD}. A gene was considered to be associated

with an outlier window if at least one of its coding exons overlapped the window

boundaries, or represented the nearest neighboring exon (in either direction).

RESULTS

AFRICAN AND EUROPEAN POPULATIONS OF D. MELANOGASTER *MAY DIFFER*

IN MULTIPLE COMPONENTS OF PULSE SONG

We first sought to determine whether any quantative aspects of male courtship song

varied between two *D. melanogaster* populations: one from within the species' inferred

ancestral range (Siavonga, Zambia) and one from Europe (Lyon, France). We divided

components of male song into a range of acoustic properties, each considered as a

single quantitative trait. To first identify population-level differences in specific song

traits, we selected 10 France-derived and 7 Zambia-derived inversion-free inbred

strains from previous collection efforts (Pool *et al.* 2012), and recorded male songs in 30

minute intervals using 40-60 males per strain (raw individual recording measurements

available in Tables S3 and S4). Strains were assigned single trait values for each song

behavior representing the mean of individual replicates (Table S5), and population

mean trait values were compared statistically between France and Zambia strain

cohorts to test for significant trait divergence in each examined song characteristic

(Figure 1; Table S5).

In total, 37 traits were analyzed, some of which were found to have significant positive

or negative correlation, often between traits with an *a priori* expectation of possible

correlation (e.g. the amount of song per minute and the number of bouts per minute)

(Figure S1). Two related traits, described by Clemens *et al.* (2020), had significant raw

Mann-Whitney (MW) p-values for the population comparison: the number of "fast"

pulses ($p = 0.0185$) and fraction of "slow" pulses to total pulses ($p = 0.0068$; Figure 1A;

Table S5). Hereafter, we focus on the latter of these correlated traits (Figure S1) in light

of its clearer signal of differentiation. Aside from those two characteristics of pulse type,

the only trait with even a marginal raw p-value (p = 0.087) was the median amplitude of

all pulses (Figure 1B; Table S5), with France strains associated with somewhat higher

(louder) pulse amplitudes than Zambia on average.

We applied a permutation-based multiple test correction to MW p-values (Table S5).

Here, we found that the ratio of "slow" pulses to total pulses had the lowest least

likelihood of false positive discovery (~10%) and represented the greatest difference in

population trait differentiation (Figure 1; Table S5). Thus, the rate of slow to total pulses

is likely to represent a genuinely differentiated trait between the Zambia and France

populations. Whereas, evidence for population differentiation in pulse amplitude, which

showed little correlation with the slow/fast pulse traits (Figure S1), was more tentative.

Figure 1 demonstrates that both of these traits had somewhat overlapping interquartile

ranges, with more distinct distributions for pulse mode.

The above findings suggested that throughout the course of multiple courtship bouts in

an uninterrupted mating setting, males from France strains were relatively more likely to

utilize a fast pulse mode over a slow pulse mode. This trait's population differentiation

was also quantified by the Q_{ST} statistic, which yielded an elevated value of 0.797, which

exceeds the F_{ST} values of 96.44% of examined SNPs genome-wide (see Materials and

Methods). In general, Q_{ST} values that exceed the bulk of the genomic F_{ST} distribution

are often considered potential targets of geographically variable directional selection,

and our comparison conservatively used F_{ST} values from individual SNPs. Although this

result should also be interpreted in the context of the number of traits initially examined,

it suggests the possibility that the usage of fast versus slow pulses may have been

influenced by differential selection between populations, whether acting on this or a

pleiotropically correlated trait. Furthermore, the relatively consistent population

differentiation observed for this trait makes it an appropriate target for QTL analyses

involving phenotypically contrasting strains.

SONG VARIATION BETWEEN RIL PARENTAL STRAINS REVEALS ADDITIONAL

TRAITS FOR MAPPING

Next, we compared mean trait values between the two strains used in the establishment

of our RIL panel, FR320N and ZI418N ("parental" strains) (Figure 1 - dotted rectangles;

Table S5). Between these strains we found a total of 10 out of 37 traits with significant

raw MW p-values (p<0.05) (Table S5). The two song traits mentioned above with regard

to population-level differences, pulse mode preference and median pulse amplitude,

were represented among this set of traits (Figure 1A & 1B, dotted rectangles). The

similarity in magnitude and direction of parental differences in these traits has important

consequences for our ability to interpret mapping results as representing likely

contributors to population differentiation, particularly for the pulse mode trait.

Seven additional traits displayed quantitatively significant differences between parental

strains: the number of bouts per minute, the transition probability of 'null' (no) song to

sine song, the median sine and pulse train length, median sine amplitude, and mode

carrier frequencies of sines and pulses (Table S4). Quantitative differences in these

traits between parental strains imply that mappable variation could exist in the RIL

panel, and we therefore elected to map these traits in addition to the two previously

mentioned. While mapping efforts for most of these traits will not reveal loci that

contribute to significant population differentiation between populations, they could reveal

loci that make a meaningful biological contribution to the production of courtship

behavior.

SOME QTLS FOR TRAITS DIFFERENTIATED BETWEEN PARENTAL STRAINS

INCLUDE KNOWN COURTSHIP GENES

We used a modified version of r/QTL (Broman *et al.* 2003; da Silva Ribeiro *et al.* 2024)

to first map the song traits associated with parental strain but not population-level

differences, under a single-locus QTL model. QTL mapping results for these traits are

depicted in Figure 2 for the amount and type of song and in Figure 3 for the properties

of pulse or sine song. For two traits, mode sine carrier frequency and median sine

amplitude, no genome-wide significant QTLs were obtained. Conversely, a total of 21

significant QTL peaks were identified for the remaining six traits, as described

individually below (Table S6). Within each of these QTLs, we highlight below any known

genes associated with male mating behavior (Table S2). However, not all QTLs include

genes known to contribute to courtship behavior, and even where they are present it is

possible that other genes with currently unknown roles in courtship behaviors may

underlie some QTLs (Mackay 2009).

The average number of bouts per minute differed between the RIL parental strains

(Table S5) [FR320N (6.24+/-2.832) vs. ZI418N (7.831+/-2.294)], and mapped to two

primary QTLs on the X chromosome (Figure 2A; Table S6). The second interval

(ChrX:10.6-11.4 Mb; note that all positions given here refer to *D. melanogaster*

reference release 5.9) was found to include two courtship genes from our annotated list,

discs large 1 (*dlg1*, FBgn0001624) and *gustatory receptor 10b* (*Gr10b*, FBgn0030297).

Intuitively, a second parental-differentiated trait, the transition probability from null sound

to sine song was found to be correlated with the above-discussed bouts per minute trait.

As expected, this "Null to Sine" trait value was higher in the parental strain with greater

bouts per minute [ZI418N (0.005+/-0.004) vs. FR320N (0.003+/-0.002)] (Figure 1; Table

S5). In spite of the above-mentioned correlation with bouts per minute, the identified

QTLs for these traits were distinct, although some non-significant peaks coincided

(Figure 2B). The null to sine probability mapped to a region near the centromere on

chromosome arm 2L starting at ~12.1 Mb (Figure 2B). This QTL's CI spans the low

recombination centromeric region and extends until chromosome arm 2R at ~7.04 Mb.

Ten courtship-associated genes were found in this interval, 4 of which encode gustatory

receptors (*Gr33a*, *Gr39a*, *Gr43a*, and *Gr47a*) (Table S7). Three genes that have been

previously established to play roles in male courtship initiation were also found in this

interval (Table S7). The gene *Chemosensory protein B 42a* (*CheB42a,* FBgn0053348)

encodes a pheromone-binding protein that plays a role in the rate and speed of

courtship initiation in males (Park *et al.* 2016). A neighboring gene, *pickpocket 25*

(*ppk25*, FBgn0053349), was similarly implicated in the initiation of courtship (Liu *et al.*

2012). Although the simplest hypothesis would be that only a single gene is responsible

for this QTL, the presence of genes relevant to courtship initiation is notable with regard

to the transition probability to sine song.

The median lengths of sine and pulse trains were both significantly different between

parental strains, and shared positive correlations with one another, which was

statistically significant in the France cohort (Figure S1). In both cases, train lengths were

longer in the Zambian [sine: 0.094+/-0.032, pulse: 1.158+/-0.136] cohort and shorter in

the French cohort [sine: 0.078+/-0.025, pulse: 1.069+/-0.128] (Figure 1; Table S5).

Median pulse train length mapped to a single QTL with restrictive peak CIs (Figure 2C;

Table S6) between Chr2L:3.21-3.39 Mb that does not harbor annotated male courtship-

related genes. Whereas, we found that median sine train length mapped to several

separate QTLs on chromosome 3: three peaks on the left arm and two peaks on the

right arm (Figure 2D) (Table S6). A single QTL at Chr3R:12.1-12.6 Mb overlaps a

known courtship gene, *spineless* (*ss*, FBgn0003513) (Table S7).

We also investigated trait variation in the mode carrier frequency of pulse and sine

songs, which were higher in the France strain [211.7+/-40.7 versus 199.3+10.9 for

pulses, 297.6+/-11.3 versus 291.9+/-5.5 for sines] (Figure 1; Table S5). Pulse carrier

frequency mapped to a single primary QTL on chromosome 3L (Figure 3A & 3B; Table

S6) and was not found to harbor known courtship genes. There was no significant QTL

associated with sine carrier frequency, and underlying LOD landscapes suggest that, as

with train lengths, sine and pulse carrier frequencies may be governed by unique loci

(Figure 3A & 3B, Table S6). This trend is somewhat recapitulated in mapping the

amplitude of pulse (Figure 3C) and sine song (Figure 3D), although no QTLs were

detected for the latter trait.

Relative to the above traits, the median amplitude of pulses gave a relatively more

polygenic signal, mapping to 11 unique QTLs that form roughly three clusters across

chromosome 3 (Figure 3C; Table S6). The two most statistically significant peaks

mapped to chromosome 3L (inclusive CIs 3L:5.98-6.71 Mb and 3L:6.80-7.24 Mb) and

were flanked by two 'minor' peaks (Figure 3A; Figure S2; Figure S3; Table S6). The

more significant of these two (peak 3) contains the courtship-associated gene *pale* (*ple,*

FBgn0005626). *pale* encodes a rate-limiting enzyme involved in catecholamine

synthesis, such as the production of dopamine precursors, and has been demonstrated

to play roles in a variety of behaviors, including courtship (Liu *et al.* 2009).

A second cluster on chromosome 3L harbored 5 additional QTLs, ranging between

3L:11 Mb and 3L:16.5 Mb and spaced roughly 1 Mb apart (Figure 3C). Peak 5 (Figure

S3, Table S7) contains a second identified courtship-associated gene, the gustatory

receptor *Gr68a* (FBgn0041231). The third cluster on chromosome 3R contained the

final two QTLs, which each have confidence intervals within the 3R:10.5-13.4 Mb

region. Distal portions of this interval (approximately 3R:12.0-13.0 Mb) constituted the

only instance of an overlapping QTL CI between any traits mapped in the present study,

in this instance overlapping with a QTL for median sine train length (Figure 3; Table S6).

Because this trait showed possible evidence of population differentiation, for

completeness we report a scan for local adaptation statistic outliers (Materials and

Methods; Figures S2 & S3; Table S7, Table S8). However, we emphasize that the

evolutionary forces underlying variation in this trait remain uncertain.

A SONG TRAIT DIFFERENTIATED BETWEEN FRANCE AND ZAMBIA

POPULATIONS YIELDS A SINGLE STRONG QTL

We next mapped the song trait that represented the most geographically differentiated

trait among those studied (Figure 1; Table S5). The ratio of slow pulses to the total

number of pulses, which was significantly higher in the Zambian strains relative to

France strains (0.833+/-0.067 vs. 0.725+/-0.125, respectively) (Figure 1A; Table S5).

One major QTL peak on chromosome 3R was identified with a broad CI interval at

approximately 3R:20.0..22.6 Mb (Figure 4A; Table S6), which was estimated to explain

42% of the parental strain trait difference.

Above, this trait was estimated to have approximately a 90% probability of reflecting

genuine population differentiation, and presented a Q_{ST} value consistent with the

differential action of directional selection (on either this or a pleiotropically correlated

trait) between populations. Under the hypothesis that a gene underlying a trait with

elevated population differentiation might display elevated genetic differentiation, we

chose to supplement our QTL mapping with a population genetic analysis of the

detected QTL region, in which we identified genomic windows within the QTL's

restrictive CIs that were associated with values of either window F_{ST}, maximum SNP F_{ST}

(da Silva Ribeiro *et al.* 2022) or the Comparative Haplotype Identity statistic (Lange &

Pool 2016) within the top 1% on the chromosome arm (Materials and Methods; Table

S8). We also made note of an expanded range of candidate genes on functional

grounds, including male-biased neuronal genes and those related to wing development.

We identified a cluster of outlier statistics within the major QTL restrictive CI (Figure 4B,

orange intervals; Materials and Methods; Table; S6) around 3R:21.75-22 Mb,

encompassing a gene-dense region that contains encouraging putative genetic

candidates. Here, we note the presence of the *enhancer of split* complex and flanking

genes, which have been previously implicated in the quantitative trait mapping of

courtship behavior (Moehring & Mackay 2004). Additionally, the transcriptional

corepressor *groucho* (*gro*, FBgn0001139) falls within the region, and has been

implicated in the *retained*-mediated repression of transcriptional targets involved in male

courtship behavior (Shirangi *et al.* 2006) (Figure 4B).

Genes previously implicated in male courtship were additionally found within restrictive

CIs of this interval (Figure 4B; Table S7). The gene *takeout*, a lipid binding protein that

is a target of the sex determination pathway and implicated in male courtship

(Dauwalder *et al.* 2002), is flanked by highly differentiated windows (Figure 4B).

Additionally, the *slowpoke* gene, which has a demonstrated role in male courtship

variation in the *D. simulans* clade (Ding *et al.* 2016) is present within the CI. Both genes

do not directly overlap population genetic outlier windows flanking them to the left,

which themselves harbor a few additional genetic candidates (Figure 4B), but this

pattern does not exclude the possibility of elevated differentiation at regulatory regions

of these genes.

Three additional regions of this QTL interval were found to contain population genetic

outlier windows. The first window contains the gene *jaguar* (*jar*, FBgn0011225). The

second window overlaps three genes: *BRWD3*, *Rrp5*, and *Dis3* (FBgn0011785,

FBgn0039182, FBgn00339183) (Figure 4). The third window does not contain any

genes, but sits upstream of the first exon of the closest proximal gene, *Furin-1* (*Fur1,*

FBgn0004509). This gene has known roles in synaptic transmission at neuromuscular

junctions (Kim *et al.* 2015), and its disruption via RNA-interference is associated with

abnormal locomotion and flight (Schnorrer *et al.* 2010). Although we highlight multiple

potential genes that may underlie this QTL, we note that it may be driven by any one of

them, by more than one of them, or by none of those listed above.

DISCUSSION

Here we describe natural variation in male courtship song components between two

natural populations of *D. melanogaster* from contrasting environments. Comparison of

population cohorts revealed significant phenotypic differences in pulse mode

preference, and relatedly the total number of fast pulses performed in the assay. We

found that the slow to fast preference is partially explained by a QTL with moderate

effect size. Conversely, the median amplitude of pulses, which had limited support for

population differentiation, was found to be under the control of multiple QTL intervals of

relatively small effects. Other traits differentiated between the RIL parental strains had

anywhere from zero to six significant QTLs detected. However, in light of the total effect

sizes estimated across QTLs for each trait (Table S6), along with the finite statistical

power expected for detecting QTLs from a few hundred RILs (*e.g.* King *et al.* 2012), we

would predict that each of these traits is also influenced by an unknown number of

smaller effect loci that did not achieve genome-wide significance in our mapping study.

Still, the distinctness of our QTL results across traits suggests variable genetic

architectures among these song components.

Phenotypic variability among RIL parental strains reveals unique QTL for multiple

courtship traits

Mapped traits that did not have significant differences at the population level but did

display significant parental strain differences were mapped in an attempt to identify

genes that may regulate courtship behavior. Components of courtship behavior are

responsive to artificial selection in lab settings using wild-derived populations (Ritchie &

Kyriacou 1994; Gaertner *et al*. 2015), reinforcing the view that fitness-relevant variation

in courtship behaviors may be present within natural populations. The generation of a

RIL panel from two inbred strains will capture a limited quantity of naturally segregating

variation, but the origins of our parental strain from two distinct natural populations may

have enhanced the extent of trait variation in behavior that is amenable to QTL

mapping.

Aside from pulse mode preference, we found that four of eight parental-differentiated

traits yielded significant QTLs overlapping known courtship-related genes (Table S2,

Table S7), and our results provide a preliminary association of these male song

components with previously implicated genes. Most of the significant QTL were found

on autosomes (Figures 2 & 3), with only two neighboring QTLs for the same trait

identified on the X chromosome. (Y chromosome effects were not considered in the

present study, due to the absence of available RIL sequencing data from males) While

the total number of QTLs and the linkage that exists among some of them preclude

strong conclusions regarding X-linkage, a reduced occurrence of X-linked QTLs might

be predicted if genes involved in male song production are sometimes subject to

sexually antagonistic selective pressures (Snook *et al.* 2005), in light of the X

chromosome spending two thirds of its time in females.

In general, we found that even when corresponding pulse and sine song traits both

differed between RIL parental strains, they mapped to distinct genomic intervals. This

pattern held for pulse versus sine train length (Figure 2C & 2D; Table S6) and for pulse

versus sine carrier frequency (Figure 3A & 3B; Table S6), in agreement with the lack of

strong correlations between these traits (Figure S1). This evidence for distinct genetic

architectures, in spite of co-directional differences in these pulse and sine song traits

between our France and Zambia parental strains, supports previous findings that

components of sine and pulse song can have unique genetic (and neuronal) bases

(Shirangi *et al.* 2013; Ding *et al.* 2016; Hiroshi *et al.* 2022; Shiozaki *et al.* 2023; Lillvis *et*

al. 2024). In contrast, we found that the nearly genome-wide significant (p = 0.06) peak

for sine amplitude (Figure 3D) falls within the inclusive CI of the most significant QTL for

pulse amplitude, at roughly 3L:6.30 Mb - 6.32 Mb (Figure 3C; Table S6, peak 3). The

potentially overlapping architecture of the two amplitude traits may indicate a pleiotropic

basis and aligns with the stronger correlation between these traits within both

populations (Figure S1).

The rate of slow to fast pulses is the most phenotypically differentiated song trait

between Zambia and France populations and involves a single detectable QTL

Among the 37 song traits examined here, two traits related to pulse mode usage,

particularly the preference for slow versus fast pulses, received the strongest statistical

support for differentiation between the France and Zambia populations examined. The

distinction between fast and slow pulse modes is a relatively recent discovery and its

possible role in sexual selection is currently unknown, although the trait does seem to

affect female movement responses to song (Clemens *et al.* 2018). This same study

showed that males use sensory input (primarily visual) to bias which pulse mode is used

(Clemens *et al.* 2018).

Assay conditions, such as chamber size and female behavioral plasticity are thus an

important consideration when interpreting results regarding this trait. Recording

chambers used in the study were small (Arthur *et al.* 2013), constraining flies to a

minimal space. We did not track the positions of flies during courtship, so we cannot

rule out that male song varied in response to female signals. The difference in pulse

mode preference in the present study may either represent a cumulative difference in

male response to females, or simply represent a difference in a probabilistic preference

of pulse choice when other environmental parameters are held constant. Still, QTL

signals for this trait were very strong (Figure 4A) and suggest that population genetic

differentiation underlies trait differences.

Our analysis provides evidence that pulse usage may have been influenced either

directly or indirectly by population-specific directional selection. In a conservatively

designed comparison between Q_{ST} and F_{ST}, the pulse type ratio showed greater

differentiation than all but around 3.5% of genome-wide SNPs. If pulse type usage has

in fact been influenced by recent directional selection, the reasons remain unclear.

Although female preference varies between some southern African and European

strains, there is no documented role of courtship song in M/Z-type mating preference

(Colegrave *et al.* 1999). Hence, it is unclear to what degree any directional selection has

resulted in song differentiation that impacts male reproductive fitness. However, it is

worth emphasizing that all experiments in this study were conducted at room

temperature. In contrast, the far more variable temperatures encountered by natural

populations, especially in the temperate zone, may exert plastic influences on song

components (Stanley & Kyriacou 2021), and it is possible that genetic adaptation may

allow for the production of universally favorable song characteristics across divergent

environments, even if some specific song differences persist at a given temperature.

Alternatively, it is possible that directional selection, if present here, may not have been

acting upon song at all. Population genomic scans for selection in this species have

reported evidence for local adaptation between warm- and cold-adapted natural

populations for multiple categories of genes that modulate synaptic function, including

neuropeptide receptors, ion channels, and structural components of the synapse (*e.g.*

Pool *et al.* 2017). It is possible that the adaptive evolution of one or more aspects of

synaptic function, even if driven by selection to maintain nervous system function in a

novel environment, might have pleiotropic consequences for behaviors such as those

involved in courtship song. Adaptive responses to other selective pressures, such as

insecticides, some of which target insect nervous systems, also may trigger pleiotropic

effects on traits that are not the direct target of selection. Hence, further study will be

needed to assess whether pulse mode usage has been targeted or pleiotropically

influenced by directional selection between African and European *D. melanogaster*

populations.

The pleiotropy hypothesis could potentially also explain variation in amplitude (*i.e.*

volume), which showed weaker evidence of differentiation between populations. Pulse

amplitude was found to be somewhat higher in France than Zambia males (Figure 1;

Table S5), sine amplitude showed a lesser trend in the same direction, and both traits

were significantly higher in the France parental strain than in the Zambia parent (Table

S5). Flies from these populations tend to be slightly larger (Lack *et al.* 2016b). Based on

measurements of females from that study, the France RIL parent has 8.1% greater

thorax length and 5.5% greater wing width than the Zambia RIL parent. If larger flies

tend to sing with greater volume, then differences in amplitude could be downstream of

differences in body or wing size. Empirically, this relationship is unclear, although there

is indirect evidence that physical size of wing components, such as the musculature,

can play roles in the sexually dimorphic expression of male song (Shirangi *et al.* 2013).

Further research will be needed to determine the degree to which song variation and

differentiation within *D. melanogaster* is shaped by directional selection to modify or

maintain sexually-selected traits in populations experiencing different environments,

directional selection on pleiotropically correlated traits, stabilizing selection, genetic drift,

or other forces.

We idenified a single large-effect QTL for pulse mode usage on chromosome arm 3R.

Curiously, this QTL for the trait with the strongest evidence for population differentiation

had a stronger mapping signal than any of the 21 QTLs identified for any of the other

traits that differed between the RIL parent strains. This QTL had a somewhat bimodal

shape (Figure 4A) that could reflect either stochastic factors in QTL mapping or else the

contribution of more than one locus.

In *Drosophila*, an average QTL may encompass hundreds of genes (Mackay 2001).

Functional approaches, such as complementation tests or gene disruption experiments,

have traditionally been employed to identify candidate genes from QTL intervals, but

can become a laborious task when attempting to test all possible candidates within a

QTL CI. We thus supplemented our QTL mapping with a population genetic analysis of

QTL intervals, leveraging available population genomic data (Lack *et al.* 2016a) to

identify specific loci with elevated genetic differentiation between populations that might

contribute to elevated trait differentiation. The candidates identified for pulse model

preference, such as the region surrounding the *enhancer of split* complex, *groucho*,

takeout, and other genes (Figure 4B; Table S7), provide promising candidates to be

tested in future studies. Subsequent analyses such as fine mapping and functional

testing such as via reciprocal hemizygosity tests (Stern 2014) may ultimately confirm

specific genes that contribute to population differences in courtship song.

Collectively, our results provide evidence of variation in pulse song usage between

naturally occurring populations of *D. melanogaster*. By combining RIL-based QTL

mapping with population genetic inference, we identified a set of genes that may be

tested in future studies. We also note that all 37 traits examined showed meaningful

variation among the studied strains, further reinforcing the vast potential of natural

variation for investigating the genetic mechanisms underlying behavior.

ACKNOWLEDGMENTS

We thank Ben Arthur and Joshua Lillvis for assistance with song analysis and members

of the Pool Lab for helpful comments on this manuscript. This work was funded by NIH

grant R35 GM13630 to J.E.P. and by the HHMI Janelia Research Campus Visitor

Project Program.

DATA AVAILABILITY STATEMENT

All raw trait data is provided in this article's Supplemental Tables. No new genomic

sequencing data was generated for this study. RIL sequence data reported by da Silva

Ribeiro *et al.* (2024) is available from the NIH Sequence Read Archive (BioProject:

PRJNA861300).

CONFLICT OF INTEREST

The authors declare that they have no conflicts of interest.

CITATIONS

Anholt RRH, O'Grady P, Wolfner MF, Harbison ST. 2020. Evolution of Reproductive

Behavior. *Genetics.* 214:49-73.

Arbuthnott D. 2009. The genetic architecture of insect courtship behavior and premating

isolation. *Heredity.* 103:15-22.

Arthur BJ, Sunayama-Morita T, Coen P, Murthy M, Stern DL. 2013. Muli-channel

acoustic recording and automated analysis of *Drosophila* courtship songs. *BMC Biol.*

11:11.

Arthur BJ, Ding Y, Sosale M, Khalif F, Kim E, Waddell P, Turaga SC, Stern DL. 2021.

Songexplorer: a deep learning workflow for discovery and segmentation of animal

acoustic communication signals. *bioRxiv.* 2021-03.

Begun DJ, Aquadro, CF. 1993. African and North American populations of *Drosophila*

melanogaster are very different at the DNA level. *Nature,* 365:548–550.

Bennet-Clark HC, Ewing AW. 1969. Pulse interval as a critical parameter in the

courtship song of *Drosophila melanogaster. Anim Behav.* 17:755-759.

Broman KW, Wu H, Sen Ś, Churchill GA. 2003. R/qtl: QTL mapping in experimental

crosses. *Bioinformatics.* 19:889-890.

Clemens J, Coen P, Remschied FA, Pereira TD, Mazumder D, Aldarondo DE, Pacheco

DA, Murthy M. 2018. Discovery of a New Song Mode in *Drosophila* Reveals Hidden

Structure in the Sensory and Neural Drivers of Behavior. *Curr Biol.* 28:2400-2412.

Colegrave, Hollocher, Hinton, Ritchie. 1999. The courtship song of African *Drosophila*

melanogaster. J Evol Biol. 13:143-150.

Cook RM. 1973. Courtship processing in *Drosophila melanogaster*. II. An adaptation to

selection for receptivity to wingless males. *Anim Behav*. 21:349-358.

Corbett-Detig R & Nielsen R. 2017. A Hidden Markov Model Approach for

Simultaneously Estimating Local Ancestry and Admixture Time Using Next Generation

Sequence Data in Samples of Arbitrary Ploidy. PLOS Genet. 13(1): e1006529.

Cowling DE, Burnet B. 1981. Courtship songs and genetic control of their acoustic

characteristics in sibling species of the *Drosophila melanogaster subgroup. Anim*

Behav. 29:924-935

da Silva Ribeiro T, Galván JA, Pool JE. 2023. Maximum SNP F_{ST} Outperforms Full-

Window Statistics for Detecting Soft Sweeps in Local Adaptation. *Genome Biol Evol*.

7:evac143.

da Silva Ribeiro T, Lollar MJ, Sprengelmeyer QD, Huang Y, Benson DM, Orr MS,

Johnson ZC, Corbett-Detig RB, Pool JE. Recombinant inbred line panels inform the

genetic architecture and interactions of adaptive traits in *Drosophila melanogaster*.

bioRxiv 2024-05.

Dauwalder B, Tsujimoto S, Moss J, Mattox W. 2002. The *Drosophila takeout* gene is

regulated by the somatic sex-determination pathway and affects male courtship

behavior. *Genes Dev*. 16:2879-2892

David JR, Capy P. 1988. Genetic variation of *Drosophila melanogaster* natural

populations. *Trends in Genetics*. 4:106–111.

Ding Y, Berrocal A, Morita T, Longdon KD, Stern DL. 2016. Natural courtship song

variation caused by an intronic retroelement in an ion channel gene. *Nature.* 536:329-

332.

Ding Y, Lillvis JL, Cande J, Berman GJ, Arthur BJ, Long X, Xu M, Dickson BJ, Stern DL.

2019. Neural Evolution of Context-Dependent Fly Song. *Curr Biol.* 29:1089-1099.

Drosophila 12 Genomes Consortium; Clark AG, Eisen MB, Smith DR, *et. al.* 2007.

Evolution of genes and genomes on the Drosophila phylogeny. *Nature*. 450: 203-218.

Ewing AW, Bennet-Clark HC. 1968. The courtship songs of *Drosophila*. *Behaviour*.

31:288-301.

Fabian DK, Lack JB, Vinayak M, Schlötterer C, Schmidt PS, Pool JE, Flatt, T. 2015.

Spatially varying selection shapes life history clines among populations of *Drosophila*

melanogaster from sub-Saharan Africa. *J Evol Biol*. 28:826–840.

Fitzgerald TD, Peterson SC. 1988. Cooperative Foraging and Communication in

Caterpillars. *Bioscience*. 38:20-25.

Gaertner BE, Ruedi EA, McCoy LJ, Moore JM, Wolfner MF, Mackay TFC. Heritable

variation in courtship patterns in *Drosophila*. *G3*. 5:531-539.

Gavrilets S, Vose A. 2007. Case studies and mathematical models of ecological

speciation. 2. Palms on an oceanic island. *Mol. Ecol.* 16:2910-2921.

Gleason JM, Nuzhdin SV, Ritchie MG. 2002. Quantitative trait loci affecting a courtship

signal in *Drosophila melanogaster*. *Heredity.* 89:1-6.

Gleason JM. 2005. Mutations and Natural Genetic Variation in the Courtship Song of

Drosophila. Behav Genet. 35:265-277.

Greenfield MD. 2016. Evolution of Acoustic Communication in Insects. *Insect Hearing.*

Springer. Manhattan (NY).

Greenspan, RJ, Ferveur, JF. 2000. Courtship in Drosophila. *Annu Rev Genet.* 34:205-

232.

Grillet M, Everaerts C, Houot B, Ritchie MG, Cobb M, Ferveur JF. 2012. Incipient

speciation in *Drosophila melanogaster* involves chemical signals. *Sci Rep.* 2:224.

Hollocher H, Wu C-I. 1996. The genetics of reproductive isolation in the *Drosophila*

simulans clade: X vs. autosomal effects and male vs. female effects. *Genetics.*

143:1243-1255.

Hope KA, Flatten D, Cavitch P, May B, Sutcliffe JS, O'Donnell J, Reiter LT. 2019. The

Drosophila Gene *Sulfateless* Modulates Autism-Like Behaviors. *Front Genet.* 10:574.

Iglesias PP, Hasson E. 2017. The role of courtship song in female mate choice in South

American Cactophilic *Drosophila. PLoS ONE.* 12:e0176119.

Keller A. 2007. *Drosophila melanogaster*'s history as a human commensal. *Curr Biol.*

17:R77-R81.

Keller L, Surette MG. 2006. Communication in bacteria: an ecological and evolutionary

perspective. *Nat Rev Microbiol.* 4:249-58.

Kim YJ, Igiesuorobo O, Ramos CI, Bao H, Zhang B, Serpe M. 2015. Prodomain

removal enables Neto to stablize glutamate receptors at the *Drosophila* neuromuscular

junction. *PLoS Genet.* 11:e1004988

King EG, Merkes CM, McNeil CL, Hoofer SR, Sen S, Broman KW, Long AD, Macdonald

SJ. 2012. Genetic dissection of a model complex trait using the *Drosophila* Synthetic

Population Resource. *Genome Res.* 22:1558-1566.

Lachaise D, Carious ML, David JR, Lemeunier F, Tsacas L, Ashburner M. 1988.

Historical biogeography of the *Drosophila melanogaster* species subgroup. *Evol Biol.*

22:159-225.

Lachaise D, Silvain JF. 2004. How two Afrotropical endemics made two cosmopolitan

human commensals: the *Drosophila melanogaster-D. simulans* palaeogeographic

riddle. *Genetica.* 120:17-39.

Lack JB, Lange JD, Tang AD, Corbett-Detig RB, Pool JE. 2016a. A thousand fly

genomes: an expanded *Drosophila* Genome Nexus. *Mol Biol Evol.* 33:3308-3313.

Lack JB, Monette MJ, Johanning EJ, Sprengelmeyer QD, Pool JE. 2016b.

Decanalization of wing development accompanied the evolution of large wings in high

altitude *Drosophila*. *Proc Natl Acad Sci USA*. 113: 1014–1019.

Lande R. 1992. Neutral theory of quantitative genetic variance in an island model with

local extinction and recolonization. *Evolution*. 46:381-389

Lange JD, Pool JE. 2016. A haplotype method detects diverse scenarios of local

adaptation from genomic sequence variation. *Mol Ecol.* 25:3081-3100

Laturney M, Moehring AJ. 2012. The Genetic Basis of Female Mate Preference and

Species Isolation in *Drosophila. Intl J Evol Biol.* 2012:328392

Leonhardt SD, Menzel F, Nehring V. 2016. Ecology and Evolution of Communication in

Social Insects. *Cell.* 164:1277-1287.

Lieberman P. 1971. On the Evolution of Human Language. *Speech Res.* 113-131.

Lillvis JL, Wang K, Shiozaki HM, Stern DL, Dickson BJ. Nested neural circuits generate

distinct signals during *Drosophila* courtship. *Curr Biol.* 34:808-824.

Liu T, Dartevelle L, Yuan C, Wei H, Wang Y, Ferveur JF, Guo A. 2009. Reduction of

dopamine level enhances the attractiveness of male *Drosophila* to other males. *PLoS*

ONE. 4:e4574.

Liu T, Starostina E, Vijayan V, Pikielny CW. 2012. Two *Drosophila* DEG/ENaC Channel

Subunits Have Distinct Functions in Gustatory Neurons That Activate Male Courtship. *J*

Neurosci. 32:11879-11889.

Mackay TFC. 2001. The Genetic Architecture of Quantitative Traits. *Annu Rev Genet.*

35:303-339.

Mackay TFC, Heinsohn SL, Lyman RF, Moehring AJ, Morgan TJ, Rollmann SM. 2005.

Genetics and genomics of *Drosophila* mating behavior. *Proc Natl Acad Sci USA.*

102:6622-6629.

Mackay TFC. 2009. The genetic architecture of complex behaviors: lessons from

Drosophila. Genetica. 136:295-302.

Manning A. 1967. The control of sexual receptivity in female *Drosophila. Anim Behav.*

15:239-250.

Maynard Smith J. 1979. Game theory and the evolution of behaviour. *Proc R Soc Lond*

B. 205:475-488.

Moehring AJ, Mackay TFC. 2004. The quantitative genetic basis of male mating

behavior in *Drosophila melanogaster.* Genetics. 167:1249-1263.

O'Grady P, DeSalle R. 2018. Phylogeny of the Genus *Drosophila. Genetics.* 209:1-25.

Oh KP, Fergus DJ, Grace JL, Shaw KL. Interspecific genetics of speciation phenotypes:

song and preference coevolution in Hawaiian crickets. *J Evol Biol.* 25:1500-1512.

Park SK, Mann KJ, Lin H, Starostina E, Kolski-Andreaco A. 2006. A Drosophila Protein

Specific to Pheromone-Sensing Gustatory Hairs Delays Males' Copulation Attempts.

Curr Biol. 16:P1154-1159,

Pool JE, Aquadro CF. 2007. The genetics basis of adaptive pigmentation variation in

Drosophila melanogaster. Mol Ecol. 16:2844-2851.

Pool JE, Corbett-Detig RB, Sugino RP, Stevens KA, Cardeno CM, Crepeau MW,

Duchen P, Emerson JJ, Saelao P, Begun, DJ, Langley CH. 2012. Population genomics

of sub-Saharan *Drosophila melanogaster*: African diversity and non-African admixture.

PLoS Genet. 8:e1003080.

Pool JE, Braun DT, Lack JB. 2017. Parallel evolution of cold tolerance within *Drosophila*

melanogaster. Mol Biol Evol. 34:349-360.

Price TD. 2002. Domesticated birds as a model for the genetics of speciation by sexual

selection. *Genetica*. 116:311-327.

Prum RO. 1990. Phylogenetic analysis of the evolution of display behavior in the

Neotropical manakins (Aves, Pipridae). *Ethology*. 84:202-231.

R Core Team. 2023. R: A Language and Environment for Statistical Computing. R

Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>

Riabinina O, Dai M, Duke T, Albert JT. 2011. Active Process Mediates Species-Specific

Tuning of Drosophila Ears. *Curr Biol.* 21:658-664.

Ritchie MG, Yate VH, Kyriacou CP. 1994. Genetic variability of the interpulse interval of

courtship song among some European populations of *Drosophila melanogaster*.

Heredity. 72:459-464.

Ritchie MG, Halsey EJ, Gleason JM. 1999. *Drosophila* song as a species-specific

mating signal and the behavioural importance of Kyriacou & Hall cycles in *D.*

melanogaster song. *Anim Behav.* 58:649-657.

Rockman MV. 2012. The QTN program and the alleles that matter for evolution: All

that's gold does not glitter. *Evolution*. 66:1-17.

Schnorrer F, Schönbauer C, Langer CHC, Dietzl G, Novatchkova M, Schernhuber M,

Fellner M, Azaryan A, Radolf M, Stark A, Keleman K, Dickson BJ. 2010. Systematic

genetic analysis of muscle morphogenesis and function in *Drosophila*. *Nature*. 464:287-

291.

Shahandeh MP, Turner TL. 2020. The complex architecture of male mate choice

evolution between *Drosophila* species. *Heredity*. 124:737-750.

Shirangi TR, Taylor BJ, McKeown M. 2006. A double-switch system regulates male

courtship behavior in male and female Drosophila melanogaster. *Nat Genet*. 38:1435-

1439.

Shirangi TR, Stern DL, Truman JW. 2013. Motor control of *Drosophila* courtship song.

Cell Rep. 5:678-86.

Shiozaki HM, Wang K, Lillvis JL, Xu M, Dickson BJ, Stern DL. 2023. Combinatorial

dynamics orchestrate flexible motor patterns in *Drosophila* courtship. *bioRxiv*.

doi:10.1101/2022.12.14.520499.

Snook RR, Robertson A, Crudgington HS, Ritchie MG. 2005. Experimental manipulation

of sexual selection and the evolution of courtship song in *Drosophila pseudoobscura*.

Behav Genet. 35:245-255.

Speith H. 1952. Mating behavior within the genus *Drosophila* (Diptera). *Bull Am Mus*

Nat Hist. 99:399-474.

Spitze K. 1993. Population structure in *Daphnia obtusa*: Quantitative genetic and

allozyme variation. *Genetics.* 135:67-374.

Sprengelmeyer QD, Mansourian S, Lange JD, Matute DR, Cooper, BS, Jirle, EV,

Stensmyr MC, Pool JE. 2020. Recurrent collection of *Drosophila melanogaster* from

wild African environments and genomic insights into species history. *Mol Biol Evol*.

11:15354-15376.

St. Johnston D. 2002. The art and design of genetic screens: Drosophila melanogaster.

Nat Rev Genet. 3:176-188

Stanley RA, Kyriacou CP. 2021. A latitudinal cline in a courtship song character of

Drosophila melanogaster. J Neurogenet. 35:333-340

Stern DL. 2014. Reported *Drosophila* courtship song rhythms are artifacts of data

analysis. *BMC Biol.* 12:38.

Sturtevant AH 1915. Experiments on sex recognition and the problem of sexual

selection in Drosophila. *J Anim Behav*. 5:351-366.

Swain B, von Philipsborn AC. 2021. Chapter Three - Sound production in *Drosophila*

melanogaster: Behavior and neurobiology. Amsterdam (Netherlands): Elsevier.

Talyn BC, Dowse HB. 2004. The role of courtship song in sexual selection and species

recognition by female *Drosophila melanogaster. Anim Behav.* 68:1165-1180

The MathWorks Inc. 2022. MATLAB version: 9.13.0 (R2022b), Natick, Massachusetts:

The MathWorks Inc. <https://www.mathworks.com>

Turissini, DA, Mcgirr, JA, Patel, SS, David, JR, Matute, DR. 2018. The Rate of Evolution

of Postmating-Prezygotic Reproductive Isolation in *Drosophila. Mol Biol Evol.* 35:312-

334.

Virtanen P, Gommers R, Oliphant TE, Haberland M, Reddy T, Cournapeau D *et al.*

2020. SciPy 1.0: Fundamental Algorithms for Scientific Computing in Python. *Nat*

Methods. 17(3), 261-272.

von Schilcher F. 1976. The role of auditory stimuli in the courtship of *Drosophila*

melanogaster. Anim Behav. 24:18-26.

Watanabe TK, Kawanishi, M. 1979. Mating preference and the direction of evolution in

the *Drosophila. Science*. 205:906-907.

Wu C-I, Hollocher H, Begun DJ, Aquadro CF, Xu X, Wu M-L 1995. Sexual isolation in

Drosophila melanogaster: A possible case of incipient speciation. *Proc Natl Acad Sci*

USA. 92:2519-2523.

FIGURES

Figure 1: *Zambia and France D. melanogaster males display differences in pulse mode*

preference (slow to total pulses) and in pulse amplitude. Violin and boxplots for France

(Blue) and Zambia (Red) males are displayed, with the middle line in boxplots

representing median values. Parental strains used in the establishment of the RIL panel

are highlighted by dotted boxes. The pairwise differences in both traits between parental

strains is of similar magnitude and direction to that of the population trend, suggesting

that some of the genetic variation contributing to phenotypic differences between

populations may be represented among the RIL panel.

Figure 2: *Traits associated with the amount and type of song map to unique QTLs*. The

number of bouts per minute (A) mapped to a pair of nearby QTLs on the X chromosome

- the only X-linked QTLs detected in this study. The probability of initiating sine song

("Null to Sine") (B) and the median length of pulse trains (C) mapped to single QTLs. In

contrast, the median length of sine trains displayed a more polygenic basis, with

multiple QTLs across both arms of chromosome 3, and distinct peaks from the pulse

song counterpart of this trait (D). LOD values are plotted against *D. melanogaster*

genome (release 5.9) positions, where color denotes chromosome arm. Genome-wide

significance thresholds of LOD values corresponding to $p = 0.05$ for each scan are

displayed as dashed lines. Significant peaks are denoted with a red star, with p-values

noted.   Further   characteristics   of   each   QTL   are   given   in   Table   S6.

1017

Figure 3: *Traits associated with sine and pulse song qualities map to unique autosomal*

QTLs. Both pulse (A) and sine (B) song amplitude and carrier frequencies were distinct

between RIL parental strains, and map to unique QTLs. The only instance of partially

overlapping QTL peaks was a region on chromosome 3R, overlapping QTLs for pulse

amplitude (C) and sine train length (D), while the QTL for pulse carrier frequency was

nearby, as was the nearly significant peak for sine amplitude. LOD values are plotted

against *D. melanogaster* genome (rel 5.9) positions, where color denotes chromosome

arm. Genome-wide significance thresholds of LOD values corresponding to p=0.05 are

displayed as dashed lines. Significant peaks are denoted with a red star, with p-values

noted.   Further   characteristics   of   each   QTL   are   given   in   Table   S6.

Figure 4: *Pulse mode preference maps to a single strong effect QTL, with potentially*

relevant genes indicated by functional annotation and population genetic analysis. (A)

QTL mapping of the quantitative trait "slow to total pulses" in song reveals a single

strong QTL on chromosome 3R with a complex LOD landscape. All features of this plot

are as described for Figures 2 and 3. (B) Negative log quantile values for $F_{ST\_FullWin}$

(Red), F_{ST_MaxSNP} (yellow), and χ_{MD} (blue) reveal genomic regions with elevated

population differentiation quantile values, sometimes overlapping peak LOD windows. -

$\log_{10}$ quantile values on the left *y*-axis begin at 1.00 to better visualize outlier windows.

Window LOD scores are represented as a black line (right *y*-axis). Restrictive

confidence intervals are represented as orange boxes under the *x*-axis. We find an

aggregation of more differentiated genomic sites within restrictive QTLs around

3R:21.75-22.25 Mb. This genomic region has been previously implicated in courtship

behavior (Moehring & Mackay 2004). Further gene annotation details are available in

Table S7**.**