POSTERIOR-SUPERIOR INSULA REPETITIVE TRANSCRANIAL MAGNETIC STIMULATION REDUCES EXPERIMENTAL TONIC PAIN AND PAIN-RELATED CORTICAL INHIBITION IN HUMANS

Nahian S Chowdhury[1,2,3], Samantha K Millard[1], Enrico de Martino[1], Dennis Boye Larsen[1], David A Seminowicz[4], Siobhan M Schabrun[5,6], Daniel Ciampi de Andrade[1], Thomas Graven-Nielsen[1]

[1] Center for Neuroplasticity and Pain (CNAP), Department of Health Science and Technology, Aalborg University, Aalborg, Denmark

[2] Center for Pain IMPACT, Neuroscience Research Australia, Sydney, New South Wales, Australia

[3] University of New South Wales, Sydney, New South Wales, Australia

[4] Department of Medical Biophysics, Schulich School of Medicine & Dentistry, University of Western Ontario, London, Canada

[5] The Gray Centre for Mobility and Activity, Parkwood Institute, London, Canada

[6] School of Physical Therapy, University of Western Ontario, London, Canada

Running title: Brain Stimulation, Experimental Pain and Cortical Inhibition

Keywords: rTMS, Posterior Insular Cortex, Transcranial-Magnetic Stimulation, Electroencephalography, Combined TMS-EEG, Experimental Pain

Number of Text Pages: 20

Number of Figures: 8

Number of Tables: 1

Corresponding author:

Professor Thomas Graven-Nielsen, DMSc, Ph.D.

Center for Neuroplasticity and Pain (CNAP),

Department of Health Science and Technology,

Faculty of Medicine, Aalborg University.

Selma Lagerløfs Vej 249, 9260 Gistrup, Denmark.

Tel.: +45 9940 9832.

E-mail: tgn@hst.aau.dk

ABSTRACT

High frequency repetitive transcranial magnetic stimulation (rTMS) to the posterosuperior insula (PSI) may produce analgesic effects. However, the neuroplastic changes behind PSI-rTMS analgesia remain poorly understood. The present study aimed to determine whether tonic capsaicin-induced pain and cortical inhibition (indexed using TMS-electroencephalography) are modulated by PSI-rTMS. Twenty healthy volunteers (10 females) attended two sessions randomized to active or sham rTMS. Experimental pain was induced by capsaicin administered to the forearm for 90 minutes, with pain ratings collected every 5 minutes. Left PSI-rTMS was delivered (10Hz, 100 pulses per train, 15 trains) ~50 minutes post-capsaicin administration. TMS-evoked potentials (TEPs) and thermal sensitivity were assessed at baseline, during capsaicin pain prior to rTMS and after rTMS. Bayesian evidence of reduced pain scores and increased heat pain thresholds were found following active rTMS, with no changes occurring after sham rTMS. Pain (prior to active rTMS) led to an increase in the frontal negative peak ~45 ms (N45) TEP relative to baseline. Following active rTMS, there was a decrease in the N45 peak back to baseline levels. In contrast, following sham rTMS, the N45 peak was increased relative to baseline. We also found that the reduction in pain NRS scores following active vs. sham rTMS was partially mediated by decreases in the N45 peak. These findings provide evidence of the analgesic effects of PSI-rTMS and suggest that the TEP N45 peak is a potential marker and mediator of both pain and analgesia.

INTRODUCTION

High frequency repetitive transcranial magnetic stimulation (rTMS) involves the delivery of magnetic pulses over the brain and has been shown to be a promising, safe, and non-invasive treatment for pain [38; 68; 71]. A common target in rTMS treatments for pain is the primary motor cortex (M1) [38]. However, M1 rTMS shows a 25-50% reduction in pain intensity in only ~50% of chronic pain patients [5; 46; 47]. Moreover, the effects of M1 rTMS on pain are believed to be mediated by functional alterations in extra motor areas [2; 40; 42]. As such, one strategy to improve the effects of rTMS on pain has been to explore non-M1 rTMS targets, with one of these being the posterior insular cortex (PSI) [17; 21; 27; 30; 37; 49], which plays a critical role in pain processing [39; 53]. In 10 healthy individuals, the effect of a single session of cTBS of the operculo-insular cortex demonstrated a decrease in heat pain sensitivity [49]. Likewise, another study reported decreased heat pain sensitivity and reduced clinical pain intensities after 5 repeated sessions of 10 Hz rTMS to PSI in 31 patients with chronic neuropathic pain [30]. This suggests PSI-rTMS can reduce pain severity and decrease heat pain sensitivity. While this is encouraging, the neuroplastic changes that occur during PSI-rTMS analgesia remain unknown.

One method of assessing pain and analgesia-related neuroplasticity has been to use single pulse TMS to measure changes in corticomotor excitability (CME) in response to painful stimuli and following rTMS [11; 12; 15; 24; 51]. However, this methodology limits the investigation of neuroplasticity to the motor system and is an indirect measure of cortical plasticity given it is confounded by activity of spinal and subcortical processes [13]. A novel method combining TMS with electroencephalography (EEG) allows for neuroplastic changes to be directly measured from multiple cortical regions [33] with excellent temporal resolution. When TMS is applied to M1 during concurrent EEG, several commonly observed peaks are detected in the TMS-evoked potential (TEP), with larger peaks occurring at 45 ms (N45) and 100 ms (N100) linked to stronger inhibitory (GABAergic) neurotransmission [9; 19; 20; 60; 61].

Recent studies using TMS-EEG suggest cortical inhibitory processes may be implicated in rTMS-induced analgesia [13; 23]. One study showed the amplitude of the frontocentral N45 and N100 TEP peak was increased in response to acute heat pain, with a larger increase in the frontocentral N45 associated with higher pain ratings [13]. Furthermore, 10Hz rTMS over dorsolateral prefrontal cortex (dlPFC) led to a decrease in TEP indices of GABAergic activity (N45 and N100) in people with major depression [74], while another study in healthy individuals showed a decrease in the amplitude of the frontocentral TEP negative peak indexing GABA (N100) following 10Hz dlPFC rTMS, with this decrease associated with an increase in pain thresholds [77]. Thus, it is possible that changes in pain perception following PSI-rTMS are mediated by changes in TEP peaks that index GABAergic activity.

The present study aimed to determine whether tonic experimental pain and cortical

inhibition are modulated by PSI-rTMS. The study was conducted on healthy human participants,

with active or sham PSI-rTMS applied during tonic capsaicin-induced pain, and TEPs and thermal

pain sensitivity measures assessed before pain, during pain before rTMS, and during pain after

rTMS. It was hypothesized that active PSI-rTMS would induce i) a reduction in capsaicin-induced

pain intensity, ii) a decrease in pain sensitivity, and iii) a reduction in TEPs that index GABAergic

activity (N45 and N100).

METHODS

Participants

This study was conducted at the Center for Neuroplasticity and Pain (CNAP), Aalborg University,

Aalborg, Denmark. All procedures adhered to the Declaration of Helsinki, with written, informed

consent obtained prior to study commencement. The study was approved by the local ethics

committee (Videnskabsetiske Komite for Region Nordjylland: N-20210047). A sample size

calculation was conducted (G*Power 3.1.9.7) based on available means/SDs reported in a previous

study exploring the effects of 10 Hz rTMS on frontal TEPs that index GABA [74] and another study

exploring the effects of PSI-rTMS on pain thresholds/pain intensity in healthy individuals [49]. We

also used our previous work which reported correlation values between repeated measurements of

TEP peaks before and after pain [13]. For the effect of rTMS on pain intensity ($\alpha = 0.05$, $\beta = 0.8$, d

$= 1.82$), a sample size of at least 5 individuals was required, while for TEPs ($\alpha = 0.05$, $\beta = 0.8$, $r =$

.9, d $= 0.92$ for N45 and d $= 0.97$ for N100) a sample size of at least 12 individuals was required.

We opted for a higher sample size of 20 participants to improve statistical power.

Twenty healthy participants (10 females; aged 26.5 ± 4.6 years (mean $\pm$ SD) were recruited

through online advertisement. Participants were excluded if they presented with any acute pain, had

a history or presence of chronic pain, neurological, musculoskeletal, psychiatric or other major

medical condition, were pregnant and/or lactating, or were contraindicated for TMS (e.g., metal

implants in the head) as assessed using the Transcranial Magnetic Stimulation Adult Safety Screen

questionnaire [64]. To further characterize the mental health profile and degree of catastrophic

thinking related to pain, participants completed the following questionnaires: Beck-Depression

Inventory [8], State-Trait Anxiety Inventory Pain Catastrophizing Scale [70] and Positive and

Negative Affective Schedule [76].

Experimental Protocol

This study used a cross-over, randomised, sham-controlled design. Participants attended two

sessions (spaced ~2-3 weeks apart) of either active or sham PSI-rTMS in a randomized sequence.

Each session involved the administration of tonic pain induced by capsaicin applied to the right volar forearm for 90 minutes (Fig. 1). Left PSI-rTMS was delivered ~50 minutes after capsaicin administration for 7.5 minutes. TEPs using combined TMS-EEG to M1 followed by thermal pain sensitivity were assessed three times: at baseline (pre-pain), during capsaicin pain prior to rTMS (pain pre-rTMS: between ~30-45 minutes after capsaicin administration), and after rTMS (pain post-rTMS: between ~65-80 minutes after capsaicin administration).

Transcranial Magnetic Stimulation Evoked Electroencephalography

Participants sat in a comfortable chair with their eyes fixated on a cross placed on the wall in front of them. Single, biphasic transcranial magnetic stimuli were delivered using a Magstim unit (Magstim Ltd., UK) and 70 mm figure-of-eight flat coil. EEG was recorded using a TMS-compatible amplifier (g.HIamp EEG amplifier, g.tec-medical engineering GmbH, Schiedlberg, Austria) at a sampling rate of 4800 Hz. Signals were recorded from 63 passive electrodes, embedded in an elastic cap (EASYCAP GmbH, Etterschlag, Germany), in line with the 10-5 system. Recordings were referenced online to 'right mastoid' and the ground electrode placed on right cheekbone. This was to reduce artifacts produced by the TMS, which was applied on the left side of the head. Electrolyte gel was used to reduce electrode impedances below ~5 kΩ. To maintain low impedances throughout the experiment, we used two net caps (GVB-geliMED GmbH, Ginsterweg Bad Segeberg, Germany) and a plastic stretch wrap handle film over the EEG cap [23]. In order to minimize the effect of the auditory response generated by the TMS coil click sound, a masking toolbox [66] was used with the participants wearing noise-cancelling headphones (Shure SE215-CL-E Sound Isolating, Shure Incorporated, United States).

Neuronavigation (Brainsight TMS Neuronavigation, Rogue Research Inc., Montréal, Canada) was used with a template MRI (MNI ICBM 152 average brain) from Brainsight software to track and calibrate each participant's head position and TMS coil position in 3D space. Surface disposable silver/silver chloride adhesive electrodes (Ambu Neuroline 720, Ballerup, Denmark) were applied over the right first dorsal interosseous (FDI) muscle parallel to muscle fibres, with the ground electrode placed on the right ulnar styloid process. The coil was oriented at 45° to the midline, inducing a current in the posterior-anterior direction. To identify the left M1 target, the scalp site ('hotspot') that evoked the largest motor evoked potential (MEP) measured at the FDI was determined and marked. The rest motor threshold (RMT) was determined using the ML-PEST (maximum likelihood strategy using parametric estimation by sequential testing) algorithm to estimate the TMS intensity required to induce an MEP of 50 microvolts with a 50% probability [3]. This method has been shown to achieve the accuracy of methods such as the Rossini-Rothwell

method [65] but with fewer pulses [69]. The test stimulus intensity was set at 90% RMT to

minimize contamination of EEG signal from re-afferent muscle activation [13].

The real-time TEP visualization tool was used [10] to confirm that artefacts (muscle,

auditory) in the signal were minimal, and that, given the coil orientation and 90% RMT stimulus

intensity, that the early peaks (<100ms) at the stimulation site were evident (P30-N100) [10; 23;

43]. The neuronavigation system and real-time TEP visualization tool were used throughout each

session to monitor coil positioning and TEP data quality across measurements within and between

sessions. For each TEP measurement (baseline, pain pre-rTMS, pain post-rTMS), ~150 TMS pulses

(~7 minutes total) were delivered with a jitter of 2.6-3.4 s [23; 43].

Thermal Pain Sensitivity Assessment

Cold and heat pain thresholds were assessed at each timepoint (immediately after TEP

measurement), in line with a previous study [13]. A 27 mm diameter thermode (Medoc Pathway

ATS device; Medoc Advanced Medical Systems Ltd) was applied over the right thenar eminence.

With the baseline temperature set at a neutral skin temperature of 32°C, participants completed two

threshold tests in the following order: to report when a decreasing temperature first became painful

(cold pain threshold, CPT) and to report when an increasing temperature first became painful (heat

pain threshold, HPT). A total of three trials were conducted for each test to obtain an average, with

an interstimulus interval of six seconds. Participants provided their threshold for each trial by

pressing a button (with their left hand) on a hand-held device connected to the Medoc Pathway.

Temperatures were applied with a rise/decrease rate of 1°C/s and return rate of 2°C/s (initiated by

the button click).

Heat-Evoked Pain

At each of the three timepoints, an additional test for heat-evoked pain was conducted. The heat

thermode (Medoc Pathway ATS device; Medoc Advanced Medical Systems Ltd) was applied at the

capsaicin administration site, and the temperature was increased towards a target temperature for 5 s

(increase rate of 1°C/s, return rate of 2°C/s). The target temperature was determined at the baseline

timepoint prior to capsaicin administration, by measuring the HPT (across 3 trials) over the

capsaicin administration site, and then adding 2 degrees above this pain threshold. Participants

provided a pain NRS rating to each 5 s stimulus, with this process repeated 3 times.

Capsaicin-induced Tonic Pain

After the baseline TEP, heat-evoked pain and thermal sensitivity assessment, an 8% topical

capsaicin patch (Transdermal patch, 'Qutenza', Astellas, 4× 4 cm) was applied to induce cutaneous

pain over the volar part of the right forearm (5 cm from the wrist) [1]. A numerical rating scale

(NRS) scale between 0 (no pain) to 10 (worst imaginable pain) was used to assess pain intensity

every 5 minutes following patch application.

Repetitive Transcranial Magnetic Stimulation

Active or sham rTMS over the orthogonal projection of the PSI was delivered with a double cone

coil (D110, Magstim Ltd., UK). The location and intensity of stimulation was determined in

between capsaicin administration and the first TEP measurement (i.e., within the first 30 minutes

after capsaicin administration). The stimulation intensity was determined by delivering single pulse

TMS using the double cone coil over the left motor representation of the tibialis anterior (TA)

muscle, which has a similar depth within the cortex as the PSI [21]. The TA hotspot and RMT were

determined by visually inspecting responses in the leg to the TMS pulse, with the RMT determined

using the ML-PEST procedure [3].

The fast PSI method was used to identify the PSI target without the need for MRI-guided

neuronavigation [18]. A recommended rTMS protocol for inducing analgesic effects was used:

1500 pulses (10 Hz, 15 trains of 10 s each, inter-train interval of 20 s, 7.5 minutes total) [16], with

the intensity of stimulation set to 80% of the TA RMT [30], and the coil oriented so that the main

phase of the biphasic waveform induced a current in the posterior-anterior direction [48]. In both

active and sham conditions, an D70 figure-of-eight TMS coil (Magstim Ltd., UK) was placed in

contact with the double cone coil but faced orthogonally using an adjustable mechanical arm.

During active rTMS, the double cone coil was activated, whereas during sham rTMS, the second

coil was activated [30].

Data Processing

Pre-processing of the TEPs was completed using EEGLAB [25] and TESA [63] in MATLAB

(R2021b, The Math works, USA), and based on previously described methods [13; 14; 55; 56; 63].

First, the data was epoched 1000 ms before and after the TMS pulse, and baseline corrected

between -1000 ms and -5 ms before the TMS pulse. Bad channels which showed large decay

artefacts from the TMS pulse were removed. The period between -5 ms and 12 ms after the TMS

pulse was removed and interpolated by fitting a cubic function. Noisy epochs were identified via the

EEGLAB auto-trial rejection function [26] and then visually confirmed. The fastICA algorithm

with auto-component rejection was used to remove eyeblink and muscle artefacts [63]. The source-

estimation noise-discarding (SOUND) algorithm was applied [55; 56], which estimates and

supresses noise at each channel based on the most likely cortical current distribution given the

recording of other channels. This signal was then re-referenced (to average). A band-pass (1-100

Hz) and band-stop (48-52 Hz) Butterworth filter was then applied. Any previously removed bad

channels were then interpolated.

The grand-averaged TEPs (across participants) for the baseline, pain pre-rTMS, and pain

post-rTMS were obtained. In line with previous studies investigating the effects of pain on TEPs

[13], and rTMS on TEPs during pain [77], the mean TEP was extracted from a frontocentral region

of interest (F1, F2, F3, F4, Fz, FC1, FC2, FC3, FC4, FCz). Peaks of the TEP from this ROI (e.g.

N15, P30, N45, P60, N100, P180) were identified for each participant using the TESA peak

function [63], with predetermined windows of interest (N15: 12-20 ms, P30: 25-40 ms, N45: 40-60

ms, P60: 55-70 ms, N100: 70-110 ms, P180: 150-200 ms) chosen to account for variation between

participants in the latency of the peaks.

Statistical Analysis

For CPT and HPTs, evoked pain NRS scores and TEP peak amplitudes, we computed the change

(Δ) scores by subtracting the pain pre-rTMS timepoint and the pain post-rTMS timepoint from the

baseline value of each respective session. Data are presented as mean ± standard deviations unless

otherwise stated. Where relevant, Cohen's *d* was reported to quantify effect sizes. Where violations

of normality occurred according to Shapiro-Wilk tests, log-transformations of the data were

conducted.

Bayesian inference was used to analyse the data, which considers the strength of the

evidence for the alternative vs. null hypothesis, using JASP software (Version 0.12.2.0, JASP

Team, 2020). Bayes factors were expressed as BF_{10} values, where BF_{10}'s of 1–3, 3–10, 10–30, 30-

100 and >100 indicated 'weak', 'moderate', 'strong', 'very strong' and 'extreme' evidence for the

alternative hypothesis, while BF_{10}'s of 1/3–1, 1/10-1/3, 1/30-1/10 and 1/100-1/30 indicated

'anecdotal', 'moderate', 'strong' ,'very strong' and 'extreme' evidence in favour of the null

hypothesis [73]. Given the novelty of the study (no prior studies on PSI rTMS on TEPs), default

priors in JASP were used to provide a balance between informed and non-informed hypotheses.

We first ran Bayesian paired t-tests to determine evidence for a difference between active

and sham sessions in pain thresholds, evoked pain NRS scores and TEP peak amplitudes at the

baseline timepoint. For capsaicin pain NRS Ratings, a 2 (session: active vs. sham) x 10 (timepoint:

0-45 minutes) Bayesian repeated measures ANOVA was conducted to determine the evidence for a

difference in pain ratings between active and sham rTMS sessions prior to stimulation. Then, a 2

(session: active vs. sham) x 9 (timepoint: 50-90 minutes) Bayesian repeated measures ANOVA was

conducted to assess pain NRS ratings following rTMS. The main effect of stimulation determined

evidence for a difference in pain ratings between active and sham rTMS, while the interaction effect

determined evidence for whether this difference changed across time. Follow-up Bayesian paired t-

tests were conducted to compare pain ratings between the end of the session (90 minutes) and onset of rTMS (50 minutes) for each group separately.

For ΔHPT, ΔCPT, Δ evoked pain NRS scores and ΔTEP peak amplitudes, A 2 (session: active vs. sham) x 2 (timepoint: pain pre-RTMS, pain post-rTMS) Bayesian repeated measures ANOVA was conducted. The interaction between session and timepoint determined evidence of modulation of the outcomes as a result of active vs. sham rTMS. Follow-up Bayesian paired t-tests were conducted to determine evidence for a change in Δ scores between pain pre and pain post-rTMS for active and sham sessions separately. Note that the difference in the baseline-normalized Δ scores between pain pre and pain post-rTMS was identical to the difference in the non-normalized raw scores. Thus, increases and decreases in the Δ scores were directly interpreted as increases and decreases in the raw outcomes. Finally, a follow-up Bayesian one sample t-test was also conducted at each timepoint and session separately to determine overall change in outcomes relative to baseline.

For any TEP peaks and pain outcomes that demonstrated at least moderate evidence of a change between pre and post active rTMS, we further explored the link between these peak changes and both pain and analgesia. We conducted a Bayesian correlation analysis to determine whether, across the whole sample, ΔTEP peak amplitudes were associated with Δ pain outcomes following capsaicin administration (pain pre-rTMS – baseline) or following rTMS (pain post-rTMS – pain pre-rTMS). We also determined whether Δ pain outcomes following rTMS were mediated by ΔTEP peak amplitudes. We used the bmlm package in R [75] to conduct a mediation analysis for repeated measures designs. This package uses a Bayesian framework to compute the mean and 95% credibility interval of plausible posterior parameter values for the total effect of the mediation model, direct effects between each variable, and the indirect effect (i.e. mediation effect). For each model, the outcome variable was Δ pain outcome following rTMS (pain post-rTMS– pain pre-rTMS), the predictor variable was PSI-rTMS session (active vs sham) and the mediating variable was ΔTEP peak amplitude (pain post-rTMS – pain pre-rTMS).

RESULTS

All participants completed the active and sham sessions, with no missing data. The mean interval between sessions was 21.5 ± 11.4 days. Eight out of 20 participants were able to correctly identify the sequence of the active and sham sessions, suggesting blinding was successful. All participants tolerated the rTMS without side effects. The mean scores on the questionnaires were 5.5 ± 8.1 for the Pain Catastrophizing Scale, 9.0 ± 6.2 for the Beck-Depression Inventory II, 25.4 ± 10.9 for the State-Trait Anxiety State Scale, 27.6 ± 9.4 the State-Trait Anxiety Trait Scale and 21.3 ± 6.0 and 7.1

± 5.1 for the Positive and Negative Affect scales respectively. These mean scores do not indicate

clinical levels of pain catastrophizing, depressive or anxiety symptoms [7; 45; 70].

The FDI RMT was $60.5 \pm 8.6\%$ for the active session and $60.0 \pm 7.9\%$ for the sham session,

with moderate evidence for no difference between sessions ($BF_{10} = 0.23$). Similar to TA RMT

values from previous studies using the double cone coil [28; 67], the TA RMT in the present study

was $42.7 \pm 6.0\%$ for the active session and $43.0 \pm 6.6\%$ for the sham session, with moderate

evidence for no difference between sessions ($BF_{10} = 0.24$).

Capsaicin Pain NRS Ratings

Prior to rTMS (0-45 min post-capsaicin), there was extreme Bayesian evidence for an increase in

pain NRS ratings following capsaicin administration (main effect of timepoint: $BF_{10} = 6.4 \times 10^{32}$,

Fig. 2A), anecdotal evidence that pain NRS ratings did not differ between the active and sham

rTMS sessions (main effect of session: $BF_{10} = 0.57$), and moderate evidence that this difference did

not change across timepoints (session x timepoint interaction: $BF_{10} = 0.15$). After rTMS (50-90 min

post-capsaicin), there was strong evidence for lower pain NRS ratings in the active versus sham

rTMS session (main effect of session: $BF_{10} = 25.4$), and extreme evidence of this difference

becoming larger over time (sessions x timepoint interaction: $BF_{10} = 2.1 \times 10^8$). Follow-up t-tests

showed that, when comparing pain NRS ratings at 90 with 50 minutes, there was strong evidence

that pain NRS reduced in the active rTMS session ($BF_{10} = 14.30$, $d = -.76$), and anecdotal evidence

that pain NRS increased in the sham rTMS session ($BF_{10} = 1.70$, $d = 0.50$).

Thermal Pain Sensitivity Distant to Capsaicin Application Site

Table 1 shows CPT, HPT and evoked pain scores at each timepoint before being normalized to

baseline. There was anecdotal evidence for no difference in HPT ($BF_{10} = 0.81$) or CPT ($BF_{10} =$

0.45) between active and sham rTMS sessions at baseline. There was extreme Bayesian evidence

that active rTMS modulated ΔHPT relative to sham (session x timepoint interaction: $BF_{10} =$

6419.72, Fig. 2B). Follow-up Bayesian paired t-tests showed that for the active rTMS session, there

was extreme evidence that ΔHPT increased from the pain pre rTMS to pain post-rTMS timepoints

($BF_{10} = 145080.16$, $d = 1.8$). For the sham rTMS session, there was anecdotal evidence that ΔHPT

decreased from pain pre-rTMS to pain post-rTMS timepoints ($BF_{10} = 1.60$, $d = -.40$). Follow-up

Bayesian one sample t-tests showed that, at the pain pre-rTMS timepoint, there was moderate

evidence that ΔHPT = 0 in the active session ($BF_{10} = 0.25$, $d = .09$) and moderate evidence that

ΔHPT > 0 in the sham session ($BF_{10} = 3.15$, $d = .58$). At the pain post-rTMS timepoint, there was

strong evidence that ΔHPT > 0 for the active session ($BF_{10} = 40.28$, $d = 0.87$), and moderate

evidence that ΔHPT = 0 for the sham session ($BF_{10} = 0.26$, $d = -.11$). There was anecdotal evidence

for no alteration in ΔCPT following active relative to sham rTMS (session x timepoint interaction:

$BF_{10} = 0.96$, Fig. 2C).

Evoked Pain NRS Scores

Prior to capsaicin administration, the HPT at the target site was 44.1 ± 2.8 °C for the active session

and 45.1 ± 3.3 °C for the sham session, with moderate evidence of no difference between sessions

($BF_{10} = 0.24$). There was moderate evidence of no difference in heat-evoked pain NRS scores

between active and sham rTMS sessions at the baseline timepoint ($BF_{10} = 0.23$). There was

anecdotal evidence that active rTMS did not modulate Δ evoked pain relative to sham (session x

timepoint interaction: $BF_{10} = 0.93$, Fig. 2D).

Transcranial Magnetic Stimulation Evoked Electroencephalography

Figures 3 and 4 show the grand-average TEPs and scalp topographies at each timepoint for the

active and sham sessions respectively. Figure 5 shows the grand-average TEPs for the frontocentral

ROI at each timepoint for the active and sham sessions. When comparing active and sham sessions

at baseline, there was moderate evidence for no difference in the N15 ($BF_{10} = 0.24$), P30 ($BF_{10} =$

0.27), N45 ($BF_{10} = 0.24$), and P60 ($BF_{10} = 0.31$) peaks, anecdotal evidence for no difference in the

P180 peak ($BF_{10} = 0.45$), and anecdotal evidence for a difference in the N100 peak ($BF_{10} = 1.3$).

Figure 6 shows the ΔTEP peak amplitudes (relative to baseline) at each timepoint for the

active and sham sessions. There was strong evidence that active rTMS modulated ΔN45 relative to

sham (session x timepoint interaction: $BF_{10} = 18.65$). Follow-up Bayesian t-tests showed moderate

evidence that ΔN45 decreased from pain pre rTMS to pain post-rTMS timepoints following active

rTMS ($BF_{10} = 3.80$, *d* = -0.60) and anecdotal evidence of no change in ΔN45 from the pain pre-

rTMS to pain post-rTMS timepoints following sham rTMS ($BF_{10} = 0.85$, *d* = 0.39). A one sample t-

test showed that at the pain pre-rTMS timepoint, there was moderate evidence ΔN45 < 0 in the

active session ($BF_{10} = 3.03$, *d* = -0.57), and moderate evidence that the ΔN45 = 0 in the sham

session ($BF_{10} = 0.26$, *d* = -0. 11). At the pain post rTMS timepoint, there was moderate evidence

that ΔN45 = 0 for the active session ($BF_{10} = 0.24$, *d* = -0.07), and moderate evidence that ΔN45 < 0

for the sham session ($BF_{10} = 3.02$, *d* = 0.57). For all other peaks, there was no moderate evidence

for any modulation by active rTMS relative to sham (BF_{10}'s for all session x timepoint interactions

< 3).

Relationship between N45 peak changes and pain parameters following capsaicin administration

To further explore the link between ΔN45 and pain, we plotted the capsaicin pain NRS score at 45

minutes and ΔHPT at the pain pre-rTMS timepoint against ΔN45 at the pain pre-rTMS timepoint,

pooled across both sessions (Figure 7A and 7B). There was moderate evidence for no correlation

between ΔN45 and both capsaicin pain NRS score at 45 minutes ($r_{38} = 0.03$, $BF_{10} = 0.20$) and

ΔHPT from baseline at the pain pre-rTMS timepoint ($r_{38} = 0.11$, $BF_{10} = 0.24$).

Relationship between N45 peak changes and pain parameters following rTMS

To explore the link between ΔN45 and reductions or increases in pain following rTMS, we plotted

changes in pain NRS ratings (90 - 50 min) and changes in HPT (pain post-rTMS – pain pre-rTMS)

against the change in the N45 peak (pain post rTMS – pain pre rTMS), pooled across sessions

(Figure 7C and 7D). Across both sessions, we found extreme evidence that a larger reduction in

capsaicin pain NRS scores were associated with a larger decrease in the N45 peak ($r_{38} = 0.54$, BF_{10}

$= 110.61$), and moderate evidence that a larger increase in HPTs was associated with a larger

decrease in the N45 peak ($r_{38} = .38$, $BF_{10} = 3.4$).

Mediation Analysis

We determined whether the reductions in pain/increases in HPT following active vs. sham

PSI-rTMS were mediated by decreases in the N45 peak. Two models were investigated (Fig. 8),

where the outcome variable was Δpain-NRS (90 - 50 mins) or ΔHPT (pain post-rTMS – pain pre-

rTMS), the predictor variable was PSI-rTMS session (active vs sham) and the mediating variable

was ΔN45 (pain post-rTMS – pain pre-rTMS). When determining the total effect of PSI-rTMS on

Δpain-NRS scores, the mean and credibility interval was 2.65 [1.59, 3.76] (i.e. a 2.65 stronger

decrease in Δpain-NRS scores for the active vs. sham condition). The mean direct effect of rTMS

on Δpain-NRS was 2.0 [0.9, 3.1] and the mean effect of rTMS on ΔN45 was -1.0 [0.9, 3.1] (i.e. a 1

μV stronger decrease in the ΔN45 in the active vs. sham condition). The mean indirect effect via

ΔN45 was 0.65 [0.01, 1.61], suggesting 95% of plausible values of the indirect effect was above 0.

This provides evidence that the reductions in pain by active vs. sham PSI-rTMS were partially

mediated by decreases in the N45 peak. There was no evidence of mediation when analysing

changes in HPT following rTMS.

DISCUSSION

The present study aimed to determine whether tonic experimental pain and cortical inhibition

assessed by TMS-EEG are modulated by PSI-rTMS. Active PSI-rTMS led to a decrease in

capsaicin-induced pain intensity and increase in heat pain thresholds in body areas away from the

site of experimental pain compared to sham. For the active rTMS session, pain prior to rTMS led to

an increase in the frontal negative peak ~45 ms (N45) TEP relative to baseline. Following active

rTMS, there was a decrease in the N45 peak back to baseline levels. In contrast, the N45 was

increased relative to baseline following sham PSI-rTMS. Lastly, decreases in pain intensity

following active vs. sham rTMS were partially mediated by reductions in the N45 peak. Taken

together, our findings suggest that 10Hz PSI-rTMS for ~7.5 minutes has analgesic effects on

experimental tonic pain. Furthermore, active PSI-rTMS not only leads to a decrease in cortical

inhibition as indexed by the TEP N45 response, but these decreases partially mediate the effects of

rTMS on pain intensity. Overall, these findings provide further insight into the role of cortical

inhibitory processes during both pain and analgesia.

Analgesic Effects of PSI-rTMS

The insular cortex is a key region involved in pain perception. The PSI receives sensory input from

the spinal cord and thalamus [31], is activated during acute and chronic pain [6; 36], triggers painful

sensations in response to direct electrical stimulation, and reduces painful sensations when lesioned

[29]. The PSI is also believed to project descending inputs to GABAergic neurons within the brain

stem. When PSI activity is increased, this triggers a loss of descending inhibition to the spinal cord

leading to increased nociception [37]. rTMS is hypothesised to have a "blocking" effect on the PSI,

which produces an antinociceptive/analgesic effect due to disinhibition of brain stem GABAergic

neurons [37]. To determine whether PSI-rTMS does indeed have an antinociceptive/analgesic

effect, we and others have determined the effects of PSI-rTMS using experimental pain models in

healthy individuals and patients with neuropathic pain or epilepsy [17; 21; 27; 30; 37; 49]. The

findings of the present study are largely consistent with previous studies, with a decrease in tonic

pain intensity following active rTMS relative to sham and an increase in heat pain thresholds

following active rTMS, although contrasted with no effects on cold pain thresholds or evoked heat

pain NRS scores. In addition, this is the first study to demonstrate the effects of PSI-rTMS using a

prolonged tonic experimental pain model (capsaicin), with previous experimental research using

transient painful stimuli [49]. Furthermore, we demonstrate these analgesic effects for the first time

using the fast PSI method [17], which precludes MRI-guided neuronavigation. New rTMS targets

such as the PSI are actively being investigated as an alternative to M1 stimulation to increase the

number of people with chronic pain responding to rTMS. However, targeting the PSI has

traditionally required MRI guided neuronavigation, which can be time consuming and cost

inefficient. The fast PSI method [18] was developed recently to reduce target identification time,

and was shown to produce similar estimates of the PSI target compared to methods requiring

neuronavigation, with high intra and inter-rater reliability. The finding that PSI-rTMS produced

analgesia using the fast PSI method is promising for clinical application of the fast PSI method as

this would greatly reduce time and costs required for targeted brain stimulation.

Cortical Plasticity during PSI-rTMS analgesia

For the first time, we used TMS-EEG to investigate the potential neuroplastic changes during PSI-

rTMS analgesia. In the active rTMS session, the TEP N45 peak was increased relative to baseline

following capsaicin administration, which is consistent with previous work showing that tonic heat

pain results in an increase in the N45 peak [13]. Following PSI-rTMS, the N45 peak was then

decreased to baseline levels. This pattern of change in the N45 peak is consistent with previous

work showing that larger increases in the N45 peak during tonic heat pain were associated with

higher pain intensity [13]. In the sham session, we did not replicate an increase in the N45 peak at

the pain pre rTMS point. The inconsistency might relate to the overall lower pain intensities at the

pre-rTMS timepoint, with Figure 7A showing the majority of ratings were 3/10 or below which can

be considered mild pain [72]. Indeed, in the sham rTMS condition, where pain gradually increased

from the pre to post rTMS timepoint, the N45 peak was increased relative to baseline. Taken

together, we argue that the natural response of the N45 peak is to increase in response to ongoing

pain. When active rTMS is delivered, this tendency is reverted, with the N45 peak being brought

back to baseline levels. These findings suggest that increased pain is associated with increases in the

N45 peak, and analgesia is associated with decreases in the N45 peak.

To further unpack the association between the N45 peak and pain intensity, we determined

whether individual changes in the N45 peak following active and sham rTMS were associated with

increases or decreases in pain perception. As anticipated, across both sessions, increases in HPT and

decreases in pain following rTMS were both associated with decreases in the N45 peak, suggesting

that regardless of stimulation, the trajectory of pain correlates with the expected trajectory of the

N45 peak. Further supporting the link between these measures, we found evidence that the

reductions in pain NRS following active vs. sham rTMS were partially mediated by decreases in the

N45 peak. This, for the first time, shows evidence for a potential causal role of the TEP N45 peak in

the analgesic effects of rTMS. Currently, the mechanisms that mediate the analgesic effects of

rTMS remain poorly understood [50]. While evidence suggests that PSI-rTMS leads to increased

connectivity between cortical and subcortical structures directly involved in descending pain

modulation [44; 52; 58], whether or not these alterations at the cortical or subcortical level, in turn,

mediate the reductions in pain intensity is seldom investigated [50]. As such, our study provides

crucial knowledge regarding these mediating mechanisms. This can inform targeted pain

interventions, such that treatments that specifically reduce the TEP N45 peak may bring about

larger pain reduction effect sizes.

Exactly how the analgesic effect of PSI-rTMS is mediated by the TEP N45 peak remains

unclear. Source reconstruction showed that the TEP N45 peak might reflect activity within the

sensorimotor cortex, despite its frontocentral topography in electrode space [13; 32]. Furthermore,

pharmacological studies show that the TEP N45 peak reflects $GABA_A$ receptor activity [60]. This

suggests that the TEP N45 peak might reflect GABAergic activity within the sensorimotor cortex.

Increased GABAergic activity in the sensorimotor cortex have been reported in response to painful

thermal stimuli consistent with the present study [41]. Studies have shown that the primary and

secondary somatosensory and motor cortices are functionally connected with the PSI, and that this

connection is critical for the sensory discrimination aspect of pain processing [34; 35; 59]. Given

PSI-rTMS is believed to block PSI function, this might result in downregulation of sensorimotor

cortical GABA receptor activity resulting in antinociceptive effects. This hypothesis is speculative,

as further multimodal work is required to elucidate the mediating role of TEP N45 peak on the

analgesic effects of rTMS and the causal role of the TEP N45 peak in pain perception broadly.

Further caution is also advised in interpreting the mediation analysis, given its exploratory nature

and the relatively low sample size, and given a mediation of heat pain thresholds was not

demonstrated.

The present study did not reveal alterations in other TEP peaks, such as the N100. This is

inconsistent with research showing a decrease in the frontocentral N100 following 10 Hz rTMS to a

different target, namely the dlPFC, with these decreases associated with increases in cold pain

thresholds [77]. Beyond clear differences related to the targeting area, the effects on TEP peaks

other than N45 may have occurred later, rather than immediately after TMS, given the analgesic

effects of PSI-rTMS became larger overtime. Indeed, it has been suggested that the effects of M1

rTMS on pain seem to build up after at least the 1st hour following stimulation [22; 57]. In this

sense, it may have been suitable to record TEPs at multiple timepoints after rTMS to determine the

onset and duration of effects.

Strengths and limitations

This present study used a thorough experimental approach including a randomized sequence of

sham and active sessions, separated by ~2-3 weeks to minimize carry over effects, and TEP

measurement based on real-time monitoring to improve data quality. Furthermore, successful

blinding between active and sham sessions was achieved, and we reported no missing data.

However, some limitations require attention. First, active rTMS over the PSI area can induce strong

muscle activity in the temporalis and frontalis muscles. This may have, in turn, contributed to the

analgesic effects as opposed to stimulation of the PSI per se. Future studies are encouraged to use

control conditions that involve stimulation of the facial muscles. Another limitation is that visual

inspection was used to estimate the motor threshold of the TA muscle instead of EMG. While this

may have led to different estimation of the TA motor threshold, visual inspection has been shown to

be a reliable method of RMT determination and some studies have shown no difference in RMT

estimation between EMG and visual inspection [4; 62]. Moreover, the performance of visual

methods is further improved when using ML-PEST [54], as was done in this study

Conclusion

This study showed that PSI-rTMS reduces tonic experimental pain intensity and increases heat pain

thresholds. These effects are accompanied by a decrease in cortical inhibition assessed by the TEP

N45 response, with this decrease partially mediating the analgesic effects of rTMS. This study

expands our understanding of the effects of PSI stimulation in humans showing that not only pain

thresholds, but also experimental tonic pain is impacted by stimulating this target, and points to the

N45 as a potential marker and mediator of analgesic effects of rTMS.

Acknowledgements: Center for Neuroplasticity and Pain (CNAP) is supported by the Danish

National Research Foundation (DNRF121). DCA is supported by a Novo Nordisk Grant

(NNF21OC0072828). The present study was not pre-registered with an analysis plan

Conflict of Interest: The authors have no conflicts of interests to declare.

REFERENCES

[1] Alhajri N, Boudreau SA, Graven-Nielsen T. Decreased default mode network connectivity following 24 hours of capsaicin-induced pain persists during immediate pain relief and facilitation. The Journal of Pain 2023;24(5):796-811.

[2] Argaman Y, Granovsky Y, Sprecher E, Sinai A, Yarnitsky D, Weissman-Fogel I. Clinical effects of repetitive transcranial magnetic stimulation of the motor cortex are associated with changes in resting-state functional connectivity in patients with fibromyalgia syndrome. The Journal of Pain 2022;23(4):595-615.

[3] Awiszus F. Fast estimation of transcranial magnetic stimulation motor threshold: is it safe? Brain Stimulation: Basic, Translational, and Clinical Research in Neuromodulation 2011;4(1):58-59.

[4] Badran BW, Ly M, DeVries WH, Glusman CE, Willis A, Pridmore S, George MS. Are EMG and visual observation comparable in determining resting motor threshold? A reexamination after twenty years. Brain Stimul 2019;12(2):364-366.

[5] Baptista AF, Fernandes AMB, Sá KN, Okano AH, Brunoni AR, Lara-Solares A, Iskandar AJ, Guerrero C, Amescua-García C, Kraychete DC. Latin American and Caribbean consensus on noninvasive central nervous system neuromodulation for chronic pain management (LAC2-NIN-CP). Pain reports 2019;4(1).

[6] Barthas F, Sellmeijer J, Hugel S, Waltisperger E, Barrot M, Yalcin I. The anterior cingulate cortex is a critical hub for pain-induced depression. Biological psychiatry 2015;77(3):236-245.

[7] Beck AT, Steer RA, Brown GK. Beck depression inventory: Harcourt Brace Jovanovich New York:, 1987.

[8] Beck AT, Ward CH, Mendelson M, Mock J, Erbaugh J. An inventory for measuring depression. Archives of general psychiatry 1961;4(6):561-571.

[9] Belardinelli P, König F, Liang C, Premoli I, Desideri D, Müller-Dahlhaus F, Gordon PC, Zipser C, Zrenner C, Ziemann U. TMS-EEG signatures of glutamatergic neurotransmission in human cortex. Scientific reports 2021;11(1):1-14.

[10] Casarotto S, Fecchio M, Rosanova M, Varone G, D'Ambrosio S, Sarasso S, Pigorini A, Russo S, Comanducci A, Ilmoniemi RJ. The rt-TEP tool: real-time visualization of TMS-Evoked Potentials to maximize cortical activation and minimize artifacts. Journal of Neuroscience Methods 2022;370:109486.

[11] Cavaleri R, Chipchase LS, Summers SJ, Chalmers J, Schabrun SM. The Relationship Between Corticomotor Reorganization and Acute Pain Severity: A Randomized, Controlled Study Using Rapid Transcranial Magnetic Stimulation Mapping. Pain Medicine 2021;22(6):1312-1323.

[12] Chowdhury NS, Chang W-J, Millard SK, Skippen P, Bilska K, Seminowicz DS, Schabrn SM. The Effect of Acute and Sustained Pain on Corticomotor Excitability: A Systematic Review and Meta-analysis of group-and individual-level data. The Journal of Pain 2022;23(10):1680-1696.

[13] Chowdhury NS, Chiang AK, Millard SK, Skippen P, Chang W-J, Seminowicz DA, Schabrun SM. Combined transcranial magnetic stimulation and electroencephalography reveals alterations in cortical excitability during pain. Elife 2023;12:RP88567.

[14] Chowdhury NS, Rogasch NC, Chiang AK, Millard SK, Skippen P, Chang W-J, Bilska K, Si E, Seminowicz DA, Schabrun SM. The influence of sensory potentials on transcranial magnetic stimulation–Electroencephalography recordings. Clinical Neurophysiology 2022;140:98-109.

[15] Chowdhury NS, Skippen P, Si E, Chiang AK, Millard SK, Furman AJ, Chen S, Schabrun SM, Seminowicz DA. The reliability of two prospective cortical biomarkers for pain: EEG peak alpha frequency and TMS corticomotor excitability. Journal of neuroscience methods 2023;385:109766.

[16] Cruccu G, Garcia□Larrea L, Hansson P, Keindl M, Lefaucheur JP, Paulus W, Taylor R, Tronnier V, Truini A, Attal N. EAN guidelines on central neurostimulation therapy in chronic pain conditions. European Journal of Neurology 2016;23(10):1489-1499.

[17] da Cunha PHM, Dongyang L, Fernandes AM, Thibes RB, Sato J, Tanaka H, Dale C, da Silva Lapa JD, de Morais ADS, Soares FHC. Non-invasive insular stimulation for peripheral neuropathic pain: Influence of target or symptom? Neurophysiologie Clinique 2022;52(2):109-116.

[18] da Cunha PHM, Tanaka H, da Silva Lapa JD, Dongyang L, Sorte AAB, Pereira TMR, Soares FHC, Fernandes AM, Da Silva VA, Graven-Nielsen T. The fast-posterior superior insula (Fast-PSI): A neuronavigation-free targeting method for non-invasive neuromodulation. Brain Stimulation: Basic, Translational, and Clinical Research in Neuromodulation 2022;15(5):1178-1180.

[19] Darmani G, Ziemann U. Pharmacophysiology of TMS-evoked EEG potentials: A mini-review. Brain
Stimul 2019;12(3):829-831.
[20] Darmani G, Zipser CM, Böhmer GM, Deschet K, Müller-Dahlhaus F, Belardinelli P, Schwab M,
Ziemann U. Effects of the Selective α5-GABAAR Antagonist S44819 on Excitability in the Human
Brain: A TMS-EMG and TMS-EEG Phase I Study. The Journal of neuroscience : the official journal
of the Society for Neuroscience 2016;36(49):12312-12320.
[21] de Andrade DC, Galhardoni R, Pinto L, Lancelotti R, Rosi Jr J, Marcolin M, Teixeira M. Into the island:
a new technique of non-invasive cortical stimulation of the insula. Neurophysiologie
Clinique/Clinical Neurophysiology 2012;42(6):363-368.
[22] de Andrade DC, Mhalla A, Adam F, Texeira MJ, Bouhassira D. Neuropharmacological basis of rTMS-
induced analgesia: the role of endogenous opioids. Pain 2011;152(2):320-326.
[23] De Martino E, Casali A, Casarotto S, Hassan G, Rosanova M, Graven-Nielsen T, Ciampi de Andrade D.
Acute pain drives different effects on local and global cortical excitability in motor and prefrontal
areas: insights into interregional and interpersonal differences in pain processing. Cerebral Cortex
2023;33(18):9986-9996.
[24] De Martino E, Gregoret L, Zandalasini M, Graven-Nielsen T. Slowing in peak-alpha frequency
recorded after experimentally-induced muscle pain is not significantly different between high and
low pain-sensitive subjects. The Journal of Pain 2021;22(12):1722-1732.
[25] Delorme A, Makeig S. EEGLAB: an open source toolbox for analysis of single-trial EEG dynamics
including independent component analysis. Journal of neuroscience methods 2004;134(1):9-21.
[26] Delorme A, Sejnowski T, Makeig S. Enhanced detection of artifacts in EEG data using higher-order
statistics and independent component analysis. Neuroimage 2007;34(4):1443-1449.
[27] Denis D, Marouf R, Rainville P, Bouthillier A, Nguyen D. Effects of insular stimulation on thermal
nociception. European Journal of Pain 2016;20(5):800-810.
[28] Dharmadasa T, Matamala JM, Howells J, Simon NG, Vucic S, Kiernan MC. The effect of coil type and
limb dominance in the assessment of lower-limb motor cortex excitability using TMS. Neuroscience
Letters 2019;699:84-90.
[29] Dimov LF, Toniolo EF, Alonso-Matielo H, de Andrade DC, Garcia-Larrea L, Ballester G, Teixeira MJ,
Dale CS. Electrical stimulation of the insular cortex as a novel target for the relief of refractory pain:
an experimental approach in rodents. Behavioural brain research 2018;346:86-95.
[30] Dongyang L, Fernandes AM, da Cunha PHM, Tibes R, Sato J, Listik C, Dale C, Kubota GT, Galhardoni
R, Teixeira MJ. Posterior-superior insular deep transcranial magnetic stimulation alleviates
peripheral neuropathic pain—A pilot double-blind, randomized cross-over study. Neurophysiologie
Clinique 2021;51(4):291-302.
[31] Dum RP, Levinthal DJ, Strick PL. The spinothalamic system targets motor and sensory areas in the
cerebral cortex of monkeys. Journal of Neuroscience 2009;29(45):14223-14235.
[32] Farzan F, Bortoletto M. Identification and verification of a'true'TMS evoked potential in TMS-EEG.
Journal of neuroscience methods 2022;378:109651.
[33] Farzan F, Vernet M, Shafi MM, Rotenberg A, Daskalakis ZJ, Pascual-Leone A. Characterizing and
modulating brain circuitry through transcranial magnetic stimulation combined with
electroencephalography. Frontiers in neural circuits 2016;10:73.
[34] Frot M, Magnin M, Mauguière F, Garcia☐Larrea L. Cortical representation of pain in primary
sensory☐motor areas (S1/M1)—a study using intracortical recordings in humans. Human brain
mapping 2013;34(10):2655-2668.
[35] Frot M, Mauguiere F. Dual representation of pain in the operculo☐insular cortex in humans. Brain
2003;126(2):438-450.
[36] Fuchs PN, Peng YB, Boyette-Davis JA, Uhelski ML. The anterior cingulate cortex and pain processing.
Frontiers in integrative neuroscience 2014;8:35.
[37] Galhardoni R, Aparecida da Silva V, García-Larrea L, Dale C, Baptista AF, Barbosa LM, Menezes
LMB, de Siqueira SR, Valério F, Rosi Jr J. Insular and anterior cingulate cortex deep stimulation for
central neuropathic pain: Disassembling the percept of pain. Neurology 2019;92(18):e2165-e2175.
[38] Giannoni-Luza S, Pacheco-Barrios K, Cardenas-Rojas A, Mejia-Pando PF, Luna-Cuadros MA, Barouh
JL, Gnoatto-Medeiros M, Candido-Santos L, Barra A, Caumo W. Noninvasive motor cortex
stimulation effects on quantitative sensory testing in healthy and chronic pain subjects: a systematic
review and meta-analysis. Pain 2020;161(9):1955-1975.
[39] Gogolla N. The insular cortex. Current Biology 2017;27(12):R580-R586.

[40] Gordon EM, Chauvin RJ, Van AN, Rajesh A, Nielsen A, Newbold DJ, Lynch CJ, Seider NA, Krimmel SR, Scheidter KM, Monk J, Miller RL, Metoki A, Montez DF, Zheng A, Elbau I, Madison T, Nishino T, Myers MJ, Kaplan S, Badke D'Andrea C, Demeter DV, Feigelis M, Ramirez JSB, Xu T, Barch DM, Smyser CD, Rogers CE, Zimmermann J, Botteron KN, Pruett JR, Willie JT, Brunner P, Shimony JS, Kay BP, Marek S, Norris SA, Gratton C, Sylvester CM, Power JD, Liston C, Greene DJ, Roland JL, Petersen SE, Raichle ME, Laumann TO, Fair DA, Dosenbach NUF. A somato-cognitive action network alternates with effector regions in motor cortex. Nature 2023;617(7960):351-359.

[41] Gross J, Schnitzler A, Timmermann L, Ploner M. Gamma oscillations in human primary somatosensory cortex reflect pain perception. PLoS biology 2007;5(5):e133.

[42] Han X, Zhu Z, Luan J, Lv P, Xin X, Zhang X, Shmuel A, Yao Z, Ma G, Zhang B. Effects of repetitive transcranial magnetic stimulation and their underlying neural mechanisms evaluated with magnetic resonance imaging-based brain connectivity network analyses. European Journal of Radiology Open 2023;10:100495.

[43] Hernandez-Pavon JC, Veniero D, Bergmann TO, Belardinelli P, Bortoletto M, Casarotto S, Casula EP, Farzan F, Fecchio M, Julkunen P. TMS combined with EEG: Recommendations and open issues for data collection and analysis. Brain Stimulation 2023.

[44] Hosomi K, Seymour B, Saitoh Y. Modulating the pain network—neurostimulation for central poststroke pain. Nature Reviews Neurology 2015;11(5):290-299.

[45] Kayikcioglu O, Bilgin S, Seymenoglu G, Deveci A. State and Trait Anxiety Scores of Patients Receiving Intravitreal Injections. Biomedicine hub 2017;2(2):1-5.

[46] Lefaucheur J-P, Aleman A, Baeken C, Benninger DH, Brunelin J, Di Lazzaro V, Filipović SR, Grefkes C, Hasan A, Hummel FC. Evidence-based guidelines on the therapeutic use of repetitive transcranial magnetic stimulation (rTMS): An update (2014–2018). Clinical neurophysiology 2020;131(2):474-528.

[47] Lefaucheur J-P, André-Obadia N, Antal A, Ayache SS, Baeken C, Benninger DH, Cantello RM, Cincotta M, de Carvalho M, De Ridder D. Evidence-based guidelines on the therapeutic use of repetitive transcranial magnetic stimulation (rTMS). Clinical Neurophysiology 2014;125(11):2150-2206.

[48] Lefaucheur J-P, Nguyen J-P. A practical algorithm for using rTMS to treat patients with chronic pain. Neurophysiologie Clinique 2019;49(4):301-307.

[49] Lenoir C, Algoet M, Mouraux A. Deep continuous theta burst stimulation of the operculo□insular cortex selectively affects Aδ□fibre heat pain. The Journal of physiology 2018;596(19):4767-4787.

[50] Liu Y, Sun J, Wu C, Ren J, He Y, Sun N, Huang H, Chen Q, Liu D, Huang Y. Characterizing the opioidergic mechanisms of repetitive transcranial magnetic stimulation–induced analgesia: a randomized controlled trial. Pain 2022:10.1097.

[51] Liu Y, Yu L, Che X, Yan M. Prolonged continuous theta burst stimulation to demonstrate a larger analgesia as well as cortical excitability changes dependent on the context of a pain episode. Frontiers in Aging Neuroscience 2022;13:804362.

[52] Maarrawi J, Peyron R, Mertens P, Costes N, Magnin M, Sindou M, Laurent B, Garcia-Larrea L. Brain opioid receptor density predicts motor cortex stimulation efficacy for chronic pain. PAIN® 2013;154(11):2563-2568.

[53] Mazzola L, Isnard J, Peyron R, Guénot M, Mauguiere F. Somatotopic organization of pain responses to direct electrical stimulation of the human insular cortex. Pain 2009;146(1-2):99-104.

[54] Mishory A, Molnar C, Koola J, Li X, Kozel FA, Myrick H, Stroud Z, Nahas Z, George MS. The maximum-likelihood strategy for determining transcranial magnetic stimulation motor threshold, using parameter estimation by sequential testing is faster than conventional methods with similar precision. The journal of ECT 2004;20(3):160-165.

[55] Mutanen TP, Biabani M, Sarvas J, Ilmoniemi RJ, Rogasch NC. Source-based artifact-rejection techniques available in TESA, an open-source TMS–EEG toolbox. Brain Stimulation: Basic, Translational, and Clinical Research in Neuromodulation 2020;13(5):1349-1351.

[56] Mutanen TP, Metsomaa J, Liljander S, Ilmoniemi RJ. Automatic and robust noise suppression in EEG and MEG: The SOUND algorithm. Neuroimage 2018;166:135-151.

[57] Nahmias F, Debes C, de Andrade DC, Mhalla A, Bouhassira D. Diffuse analgesic effects of unilateral repetitive transcranial magnetic stimulation (rTMS) in healthy volunteers. PAIN® 2009;147(1-3):224-232.

[58] Pagano RL, Fonoff ET, Dale CS, Ballester G, Teixeira MJ, Britto LR. Motor cortex stimulation inhibits
thalamic sensory neurons and enhances activity of PAG neurons: possible pathways for
antinociception. PAIN® 2012;153(12):2359-2369.
[59] Peltz E, Seifert F, DeCol R, Dörfler A, Schwab S, Maihöfner C. Functional connectivity of the human
insular cortex during noxious and innocuous thermal stimulation. Neuroimage 2011;54(2):1324-
1335.
[60] Premoli I, Castellanos N, Rivolta D, Belardinelli P, Bajo R, Zipser C, Espenhahn S, Heidegger T,
Müller-Dahlhaus F, Ziemann U. TMS-EEG signatures of GABAergic neurotransmission in the
human cortex. Journal of Neuroscience 2014;34(16):5603-5612.
[61] Premoli I, Király J, Müller-Dahlhaus F, Zipser CM, Rossini P, Zrenner C, Ziemann U, Belardinelli P.
Short-interval and long-interval intracortical inhibition of TMS-evoked EEG potentials. Brain Stimul
2018;11(4):818-827.
[62] Pridmore S, Fernandes Filho JA, Nahas Z, Liberatos C, George MS. Motor threshold in transcranial
magnetic stimulation: a comparison of a neurophysiological method and a visualization of
movement method, Vol. 14: LWW, 1998. pp. 25-27.
[63] Rogasch NC, Sullivan C, Thomson RH, Rose NS, Bailey NW, Fitzgerald PB, Farzan F, Hernandez-
Pavon JC. Analysing concurrent transcranial magnetic stimulation and electroencephalographic data:
A review and introduction to the open-source TESA software. Neuroimage 2017;147:934-951.
[64] Rossi S, Hallett M, Rossini PM, Pascual-Leone A. Safety, ethical considerations, and application
guidelines for the use of transcranial magnetic stimulation in clinical practice and research. Clin
Neurophysiol 2009;120(12):2008-2039.
[65] Rossini PM, Barker A, Berardelli A, Caramia M, Caruso G, Cracco R, Dimitrijević M, Hallett M,
Katayama Y, Lücking C. Non-invasive electrical and magnetic stimulation of the brain, spinal cord
and roots: basic principles and procedures for routine clinical application. Report of an IFCN
committee. Electroencephalography and clinical neurophysiology 1994;91(2):79-92.
[66] Russo S, Sarasso S, Puglisi GE, Dal Palù D, Pigorini A, Casarotto S, D'Ambrosio S, Astolfi A,
Massimini M, Rosanova M. TAAC-TMS Adaptable Auditory Control: A universal tool to mask
TMS clicks. Journal of neuroscience methods 2022;370:109491.
[67] Schecklmann M, Schmaußer M, Klinger F, Kreuzer PM, Krenkel L, Langguth B. Resting motor
threshold and magnetic field output of the figure-of-8 and the double-cone coil. Sci Rep
2020;10(1):1644.
[68] Short EB, Borckardt JJ, Anderson BS, Frohman H, Beam W, Reeves ST, George MS. Ten sessions of
adjunctive left prefrontal rTMS significantly reduces fibromyalgia pain: a randomized, controlled
pilot study. Pain 2011;152(11):2477-2484.
[69] Silbert B, Patterson H, Pevcic D, Windnagel K, Thickbroom G. A comparison of relative-frequency and
threshold-hunting methods to determine stimulus intensity in transcranial magnetic stimulation.
Clinical Neurophysiology 2013;124(4):708-712.
[70] Sullivan MJ, Bishop SR, Pivik J. The pain catastrophizing scale: development and validation.
Psychological assessment 1995;7(4):524.
[71] Taylor JJ, Borckardt JJ, George MS. Endogenous opioids mediate left dorsolateral prefrontal cortex
rTMS-induced analgesia. Pain 2012;153(6):1219-1225.
[72] Utne I, Miaskowski C, Bjordal K, Paul SM, Jakobsen G, Rustøen T. Differences in the use of pain
coping strategies between oncology inpatients with mild vs. moderate to severe pain. Journal of pain
and symptom management 2009;38(5):717-726.
[73] van Doorn J, van den Bergh D, Böhm U, Dablander F, Derks K, Draws T, Etz A, Evans NJ, Gronau QF,
Haaf JM. The JASP guidelines for conducting and reporting a Bayesian analysis. Psychonomic
Bulletin & Review 2021;28:813-826.
[74] Voineskos D, Blumberger DM, Rogasch NC, Zomorrodi R, Farzan F, Foussias G, Rajji TK, Daskalakis
ZJ. Neurophysiological effects of repetitive transcranial magnetic stimulation (rTMS) in treatment
resistant depression. Clinical Neurophysiology 2021;132(9):2306-2316.
[75] Vuorre M, Bolger N. Within-subject mediation analysis for experimental data in cognitive psychology
and neuroscience. Behavior Research Methods 2018;50:2125-2143.
[76] Watson D, Clark LA, Tellegen A. Development and validation of brief measures of positive and
negative affect: the PANAS scales. Journal of personality and social psychology 1988;54(6):1063.
[77] Ye Y, Wang J, Che X. Concurrent TMS-EEG to reveal the neuroplastic changes in the prefrontal and
insular cortices in the analgesic effects of DLPFC-rTMS. Cerebral Cortex 2022;32(20):4436-4446.

FIGURE LEGENDS

Figure 1. Diagram of the experimental protocol.

Figure 2. Mean (n = 20) and Standard Errors for capsaicin pain numerical rating scale (NRS) scores (**A**), Δ cold pain thresholds (**B**), Δ heat pain thresholds (**C**) and Δ evoked pain NRS scores (**D**) at each timepoint. *, **, ***, **** indicates moderate, strong, very strong and extreme Bayesian evidence of a difference between conditions or difference from baseline.

Figure 3. A: The Grand-average TEPs (n = 20) for all 63 channels during the baseline, pain pre-rTMS, and pain post-rTMS timepoints of the active repetitive transcranial magnetic stimulation session. The red line represents the mean TEP for the frontocentral region of interest. **B**: Scalp topographies and estimated source activity at timepoints where TEP peaks are commonly observed, including the N15, P30, N45, P60, N100, and P180.

Figure 4. A: The Grand-average TEPs (n = 20) for all 63 channels during the baseline, pain pre-rTMS and pain post-rTMS timepoints of the sham repetitive transcranial magnetic stimulation session. The red line represents the mean TEP for the frontocentral region of interest. **B**: Scalp topographies and estimated source activity at timepoints where TEP peaks are commonly observed, including the N15, P30, N45, P60, N100, and P180.

Figure 5. The Grand-average TEPs (n=20) for the frontocentral ROI during the baseline, pain pre-rTMS and pain post rTMS timepoints of the active **(A)** and sham **(B)** repetitive transcranial magnetic stimulation sessions.

Figure 6. Mean (n = 20) and Standard Errors for ΔN15 (**A**), ΔP30 (**B**), ΔN45 (**C**), ΔP60 (**D**), ΔN100 (**E**) and ΔP180 (**F**) peaks normalized to baseline. * indicates moderate Bayesian evidence of a difference between conditions or a difference from baseline.

Figure 7. (A) Relationship between ΔN45 (pain pre-rTMS – baseline) and pain NRS at 45 minutes pooled across sessions (B). Relationship between ΔN45 (pain pre-rTMS – baseline) and Δ heat pain thresholds (pain pre-rTMS – baseline) pooled across sessions **(C)** Relationship between ΔN45 (pain post-rTMS – pain pre-rTMS) and Δ pain NRS (90– 50 mins). **(D)** Relationship between ΔN45 (pain post rTMS – pain pre rTMS) and Δ heat pain thresholds (pain post rTMS – pain pre rTMS). *, **, ***, **** indicates moderate, strong, very strong and extreme Bayesian evidence of a correlation.

Figure 8. (A) Mediation model with session (active vs. sham) as the predicting variable, Δ pain NRS (90– 50 mins) as the outcome variable, and ΔN45 (pain post-rTMS - pain pre-rTMS) as the mediating variable. **(B)** Mediation model with session (active vs. sham) as the predicting variable, Δ heat pain thresholds (pain post-rTMS-pain pre-rTMS) as the outcome variable, and ΔN45 (pain post-rTMS -pain pre-rTMS) as the mediating variable. Credibility intervals for the mean effects and standard deviations are shown in brackets. In both models, the total effect represents the effect of session on the outcome in the absence of the mediator. The c' path represents the direct effect of session on the outcome in the presence of the mediator, the a path represents the effect of session on ΔN45, and the b path represents the effect of ΔN45 on the outcome. The indirect effect determines the extent to which the effect of session on the outcome is accounted for by ΔN45.

FIGURES

A Capsaicin Pain Ratings

B Heat Pain Thresholds

C Cold Pain Thresholds

D Change in Evoked Pain

946 **TABLES**

947 *Table 1. Thermal sensitivity outcomes at each timepoint for active and sham rTMS sessions before*

948 *and after capsaicin-induced pain. CDT = Cold Detection Threshold, WDT = Warm Detection*

949 *Threshold, CPT = Cold Pain Threshold, HPT = Heat Pain Threshold.*

	Active rTMS			Sham rTMS		
	Baseline	Pain pre-rTMS	Pain post-rTMS	Baseline	Pain pre-rTMS	Pain post-rTMS
CPT (°C)	9.4±1.92	9.5±1.78	8.5 ±1.85	8.3±2.05	6.9±1.83	7.3±1.83
HPT (°C)	45.1±0.74	45.3±0.74	46.9 ±0.71	46.5±0.80	47.1 ±0.73	46.3 ±0.82
Evoked Pain (NRS/10)	4.8±0.50	6.1±0.46	6.5±0.39	4.8±0.46	6.4±0.56	7.3±0.41

Pain ratings collected every 5 minutes

A Capsaicin Pain Ratings

— Active — Sham

Pain Score (NRS/10)

Time after Capsaicin Administration (mins)

**

B Heat Pain Thresholds

Change in Threshold (°C)

****

*

**

Pain, pre-rTMS   Pain, post-rTMS

Timepoint

C Cold Pain Thresholds

Change in Threshold (°C)

Pain, pre-rTMS   Pain, post-rTMS

Timepoint

D Change in Evoked Pain

■ Active ■ Sham

Change in Pain Rating (NRS)

Pain, pre-rTMS   Pain, post-rTMS

Timepoint

A.

Sham Baseline

Sham Pain, pre-rTMS

Sham Pain, Post-rTMS

B.

N15   P30   N45   P60   N100   P180

Amplitude (μV)

-1

-2

Pre-rTMS vs. Baseline Correlations

A. Pain NRS Ratings — r = .04

B. Heat Pain Thresholds — r = .11

Post-rTMS vs. Pre-rTMS correlations

C. r = .54****

D. r = .38*

Sham
Active

Pain NRS Scores

Heat Pain Thresholds

Total effect = 2.65[1.59, 3.76]
Mediation effect = 0.65[0.01, 1.61]

ΔN45

a = -1
[-1.6, -0.37]
SD = 0.46
[0.029, 1.1]

b = -0.67
[-1.3, -0.056]
SD = 0.41
[0.019, 1.2]

Active
vs. Sham

c' = 2
[0.9, 3.1]
SD = 0.45
[0.017, 1.2]

Δpain
NRS

Total effect = -2.35[-1.44, -3.27]
Mediation effect = -0.09[-0.84, 0.62]

ΔN45

a = -1
[-1.6, -0.37]
SD = 0.51
[0.049, 1.1]

b = -0.004
[-0.63, 0.6]
SD = 0.62
[0.043, 1.3]

Active
vs. Sham

c' = -2.3
[-3.3, -1.3]
SD = 0.5
[0.028, 1.3]

ΔHPT