The *autophagy-related genes AtATG5* and *AtATG7* influence reserve mobilisation and response to ABA during seed germination

Estefanía Contreras[1], Inmaculada Sánchez-Vicente[3], Elena Pastor-Mora[1], Mar Aylón-Rodríguez[1], Mar G. Ceballos[1], Miguel Ángel Delgado-Gutiérrez[1], Óscar Lorenzo[3], JesúsVicente-Carbajosa[1,2]*, Raquel Iglesias-Fernández[1,2]*

E.C.: estefania.contreras@upm.es; https://orcid.org/0000-0002-1822-6839
I.S.V.: elfik@usal.es; https://orcid.org/0000-0001-6608-5578
E.P.M.: elena.pastor@inia.csic.es
M.A.R.: mar.aylon@upm.es
M.G.C.: mar.gceballos@upm.es
M.D.G.: md642046@gmail.com
O.L.: oslo@usal.es; https://orcid.org/0000-0001-9523-0789
J.V.C: jesus.vicente@upm.es; https://orcid.org/0000-0002-6332-1712
R.I.F.: raquel.iglesias@upm.es; https://orcid.org/0000-0003-4773-7176

*For correspondence: (email: raquel.iglesias@upm.es and jesus.vicente@upm.es)

[1]Centro de Biotecnología y Genómica de Plantas-Severo Ochoa (CBGP, UPM-INIA), Universidad Politécnica de Madrid (UPM) - Instituto Nacional de Investigación y Tecnología Agraria y Alimentaria (CSIC/INIA), 28223 Pozuelo de Alarcón, Spain
[2]Departamento de Biotecnología-Biología Vegetal, Escuela Técnica Superior de Ingeniería Agronómica, Alimentaria y de Biosistemas, UPM, 28040 Madrid, Spain
[3]Departamento de Botánica y Fisiología Vegetal, Instituto de Investigación en Agrobiotecnología (CIALE), Facultad de Biología, Universidad de Salamanca, C/ Río Duero 12, 37185 Salamanca, Spain.

Number of Figures: 10 (in colour Figs.: 1, 2, 4-10)

Word count: 7869 (including Materials and Methods)

Supplementary data: 10 Figs, 1 table and 5 data worksheets.

Summary

Autophagy is an intracellular recycling mechanism that generally degrades cytoplasmic components non-selectively, but it can also target specific substrates under certain conditions. Here, we investigate the impact of autophagy on Arabidopsis seed biology through autophagy-related (*ATG*) genes *AtATG5* and *AtATG7*, and their role in ABA responses. Seeds of *atg5* and *atg7* mutants germinate significantly slower than Col-0, especially under ABA, and show transcriptomic differences and histochemical alterations in the organization of lipid droplets and protein storage vacuoles. Notably, immunolocalization of ATG8 is observed in PSV of Col-0, but not in *atg* mutants. Differentially expressed genes in *atg7* compared to Col-0 in response to ABA include reported targets of the transcription factor (TF) ABI5, a master regulator of ABA signalling in the seed. Interestingly, the decrease in ABI5 normally observed in Col-0 seeds after imbibition is delayed in *atg* mutants, which also show altered accumulation in developing seeds of the bZIP67, an ABI5 homolog that regulates reserve biosynthesis. Yeast two-hybrid and co-immunoprecipitation assays confirmed a direct interaction with the autophagy machinery *in vitro* and *in vivo*, mediated through ATG8. Our data highlight the relevance of autophagy in seed reserve mobilization, and seed germination through ABA responses by a TF decay mechanism.

Keywords

Arabidopsis thaliana, Autophagy, ABA signalling, bZIPs, ABI5, Lipid metabolism, Protein storage vacuoles (PSV), Reserve mobilization, Seed germination

Introduction

Autophagy is a conserved recycling mechanism in eukaryotes that degrades cytoplasmic components, removing unnecessary or damaged elements. While traditionally seen as a non-selective process, it also targets specific proteins, lipids, and organelles in response to biotic and abiotic stresses (Su *et al.*, 2020; Qi *et al.*, 2021). Plant autophagy plays a crucial role in cellular homeostasis and facilitates adaptation to changing environments, making it essential for plant survival under various stresses and developmental programs (Liu *et al.*, 2009; Di Berardino *et al.*, 2018). In plant cells, autophagy occurs in the central vacuole via vesicular trafficking, mainly through macroautophagy and microautophagy. The most studied, macroautophagy, transports cargo in double-membrane autophagosomes from the endoplasmic reticulum (ER) to the vacuole (Michaeli *et al.*, 2016). In microautophagy, cytoplasmic elements such as lipid droplets (LD; also known as oil bodies and oleosomes), specialized organelles accumulating triacylglycerols (TAG) are directly engulfed by the central vacuole through tonoplast invagination (Siénko *et al.*, 2020).

In Arabidopsis, autophagy involves around 40 conserved autophagy-related proteins (ATGs) and other evolutionary conserved factors from over 17 protein families, including kinases, ubiquitin-like proteins, vacuolar sorting factors, and cysteine proteases (Ding *et al.*, 2018). *In A. thaliana*, ATGs are involved in three main complexes for autophagosome formation: (i) the ATG1 kinase complex, which induces autophagy via interaction with the Target Of Rapamycin (TOR) under nutrient starvation (Dobrenel *et al.*, 2016); (ii) the phosphatidylinositol 3-kinase (PI3K) complex, which initiates phagophore nucleation (Yang *et al.*, 2010); and (iii) the ATG9–ATG18–ATG2 complex, which recruits membranes for phagophore expansion (Sawa-Makarska *et al.*, 2020). The ATG8–ATG12 conjugation system attaches phosphatidylethanolamine (PE) to ATG8, enabling selective cargo recognition via ATG8-interacting motifs (AIMs) that bind to selective receptors like Neighbour of BRCA1 gene 1 (NBR1) and translocator protein (TSPO) (Kellner *et al.*, 2017; Ding *et al.*, 2018; Luo *et al.*, 2021). The autophagy-related factors AtATG5 and AtATG7 are crucial for the conjugation of PE to ATG8 in both animals and plants (Xie and Klionsky, 2007; Minina *et al.*, 2018; Fraiberg and Elazar, 2020). Genetic studies in *A. thaliana* have shown that *ATG5* and *ATG7* are essential for storage protein and lipid accumulation during embryogenesis and seed maturation. Their overexpression enhances seed production and nitrogen content, while *atg5* and *atg7* mutants display altered seed composition (Guiboileau *et al.*, 2013; Masclaux-Daubresse *et al.*, 2014; Di Berardino *et al.*, 2018; Minina *et al.*, 2018; Fan *et al.*, 2019). Similarly, in *Oryza sativa*, autophagy affects starch quality, influencing both seed size and appearance, as *Osatg7-1* produces fewer and small seeds with a chalky appearance and a low starch content in endosperm (Sera *et al.*, 2019; Iglesias-Fernández & Vicente-Carbajosa, 2022).

Seeds are essential for plant evolution, enabling dispersal, survival under harsh conditions, and regrowth in favourable environments (Finch-Savage & Bassel, 2016). Seed maturation, germination, and post-germination involve the accumulation of reserves like starch, seed storage proteins (SSP), and lipids (triacylglycerols; TAG) (Holdsworth *et al.*, 2008; Bewley *et al.*, 2013; Sajeev *et al.*, 2024). SSP synthesis and storage in maturing seeds occur in the ER, which transports them to seed-specific vacuoles, known as Protein Storage Vacuoles (PSV), through an autophagy-like process (Hills, 2004; Vitale *et al.*, 2022). Seed germination starts with water uptake and ends with the breakage of embryo-covering layers (i.e. endosperm rupture) that allows radicle emergence (germination *sensu stricto*). The post-germination stages involve the hydrolysis of seed storage reserves by *de novo* synthesised proteases, providing essential support for seedling growth until the photosynthetic capacity is acquired (Nakabayashi *et al.*, 2005). In imbibed Arabidopsis seeds, the degradation of SSP begins at the radicle, coinciding with cell expansion and differentiation. After radicle emergence (i.e. post-germination period), SSP catabolism occurs mainly in the cotyledons,

especially when the SSP reserves of the embryonic axis are nearly depleted (Müntz, 2007; Iglesias-
Fernández *et al.*, 2014). In *Vigna mungo* (Fabaceae), the ERvt (ER to vacuole trafficking) pathway
has been reported to transport cysteine proteases from the ER to PSV, encapsulated within KDEL
Vesicles (KV) (Okamoto *et al.*, 2003). Interestingly, the autophagy receptor TSPO plays a major
role in the regulation of fatty acids (FA) and LD in Arabidopsis germinating seeds and in seedlings.
Recently, studies in the knock-out mutant of LD structural protein -CALEOSIN 1 (CLO1) have
described its influence in the degradation of LD by autophagy (lipophagy) in germinating seeds of
Arabidopsis (Jurkiewicz *et al.*, 2018; Miklaszewska *et al.*, 2023; Qin *et al.*, 2023; Zhuang *et al.*,
2024).

In natural soil seedbeds, seeds encounter a complex environment with various stresses, requiring
mechanisms for survival and adaptation. Seed vigor, defined by the ability to germinate under
different conditions, is crucial for this adaptation (Finch-Savage and Bassel, 2016). The ER plays a
central role in detecting and managing stress responses. When stress occurs, the Unfolded Protein
Response (UPR) pathway is activated to prevent protein accumulation in the ER and restore
homeostasis (Ellgaard *et al.*, 2003; Yang *et al.*, 2016). Persistent stress leads to the selective
removal of ER domains via autophagy, helping manage protein overload (Stephani *et al.*, 2020).
While autophagy is essential for degradation in response to various stimuli, its role in seeds remains
underexplored. In *Medicago truncatula*, ER stress and autophagy, mediated by MtATGs, are key
for seed development and drought response (Yang *et al.*, 2021). Abscisic acid (ABA) regulates
plant adaptation to abiotic stresses and controls seed maturation and dormancy. Environmental
stress during embryo development can halt germination through an ABA-dependent mechanism. In
Arabidopsis thaliana, the autophagy receptor NBR1 interacts with ABA signaling transcription
factors (TFs: ABI3, ABI4, ABI5), and its overexpression reduces seed sensitivity to ABA
(Carbonero *et al.*, 2017; Tarnowski *et al.*, 2020; Luo *et al.*, 2021; Chen *et al.*, 2022; Matilla, 2022).
130
This study investigates the role of autophagy in seed biology by analyzing *Arabidopsis thaliana*
AtATG5 and *AtATG7* loss-of-function mutants. Seeds from *atg5* and *atg7* knockout mutants
germinate slower than wild type (WT, Col-0), both under normal conditions and with ABA.
Histochemical analysis showed alterations in storage compounds (proteins and lipids) in these
mutants compared to WT seeds at both dry and imbibed stages. Immunofluorescence revealed that
ATG8a did not localize to PSV in autophagy-deficient mutants, supporting the role of autophagy in
reserve mobilization. Transcriptomic analysis during germination identified significant
transcriptional differences between *atg7* and WT seeds, particularly in ABA-imbibed seeds, with
22.5% of differentially expressed genes linked to ABI5 TF response. Western blot analysis showed
altered levels of ABI5 and bZIP67 in autophagy mutants, indicating autophagy's regulatory role in

seed development and imbibition. Protein-protein interactions between ATG8 and ABI3/ABI5 were
confirmed through yeast two-hybrid assays (Y2H). Co-IP assays have *in vivo* confirmed the ABI5-
ATG8 interaction in Arabidopsis 4 day-old seedlings. Overall, the study highlights autophagy's
involvement in seed development, germination, and the seed's response to environmental signals,
such as ABA.

Materials and Methods

Plant material and growth conditions

Arabidopsis thaliana Columbia (Col-0) ecotype was used as WT. Knockout T-DNA insertion lines
were used for *ATG5* (*atg5-1;* SAIL_129B079) and *ATG7* (*atg7-2;* GK-655B06; Hofius *et al.*, 2009;
Yoshimoto *et al.*, 2009), kindly donated by C. Masclaux-Daubresse (IJPB, France), *ABI5* (*abi5-7*;
Nambara *et al.*, 2002; Albertos *et al.*, 2015), *35S:cMyc-ABI5* (Albertos et al, 2015) and *ProUBQ10-*
GFP-ATG8A (kindly provided by Dr. Cecilia Gotor and Dr. Luis C. Romero, Instituto de
Bioquímica Vegetal y Fotosíntesis-CSIC-Sevilla; Calero-Muñoz *et al.*, 2019; Nakamura *et al.*,
2021).

Seeds were surface sterilized in 70% ethanol for 15 minutes, followed by a 100% ethanol wash and
air-dried. They were then plated on solid half-strength Murashige and Skoog medium with 1%
sucrose (MS/2, Duchefa Biochemie, Haarlem, Netherlands) and stratified at 4ºC for 3 days. Seeds
were incubated for 2 weeks at 21ºC in a germination chamber under a long-day photoperiod (16h
light/8h dark) with a light intensity of 155 μmol photons/m²/s. Seedlings were transferred to
soil:vermiculite (3:1) pots and grown in a greenhouse under similar conditions. Siliques were
collected at different seed embryogenesis and maturation stages for RNA isolation. Mature brown
seeds were harvested, stored at 21ºC and 30% relative humidity, and considered after-ripened after
one month.

Germination assays

Three replicates of 50 after-ripened, non-stratified mature seeds were sterilized and then imbibed on
moist grid-patterned nitrocellulose filter paper (Whatman nº1; Thermo Fisher Scientific; Waltham,
MA, USA) placed on 55 mm Petri dishes containing solid MS/2, with or without 1 μM 2-cis, 4-
trans-(±) Abscisic acid (ABA; Merck, Rahway, NY, USA). Plates were kept in a germination
chamber under the specified conditions. Seed Coat Rupture (SCR) and Endosperm Rupture were
evaluated at 0, 24, 30, 36, 48, 54, 72, 76, and 80 hours of imbibition (hoi) using an Olympus stereo
microscope SZ2-ILST (Olympus; Tokyo, Japan). Statistical analysis and the time to reach 50%

germination (t50) were calculated using the GERMINATOR Microsoft Office Excel 16.9 package

(Joosen et al., 2010). Col-0 and *atg7* imbided seeds were also collected at selected time points (0,

24, 48, 96 hoi) in the absence and presence of 1 μM ABA and frozen in liquid nitrogen for RNA

isolation and subsequent RNA-seq analysis.

178 Generation of *PAtATG5::uidA* and *PAtATG7::uidA* transgenic lines and

179 histochemical GUS assays

AtATG5 (*At5g17290*) and *AtATG7* (*At5g45900*) native promoter sequences were amplified from *A.*

thaliana genomic DNA (from -404 bp and -340 bp before the initial ATG, respectively) using

oligonucleotides with *Bam*HI and *Sal*I restriction sites. PCR products were cloned into the pGEM

T-easy vector (Thermo Fisher Scientific) and transferred to the pBI101 vector (between *Bam*HI and

*Sal*I sites), which contains the *Escherichia coli uidA* reporter gene. Recombinant plasmids were

transformed into C51C1 *Agrobacterium tumefaciens* cells via electroporation (ECM 630

Electroporator; BTX; Harvard Bioscience, Holliston, MA, USA) and introduced into *A. thaliana*

Col-0 plants using Agrobacterium floral dip transformation (Clough and Bent, 1998). The presence

of the pBI101 vector in T1 plants was confirmed by hygromycin resistance (40 mg/L). Plants from

subsequent generations were self-fertilized until homozygous lines were identified in T3, with

100% germination on selection media. *AtATG5* and *AtATG7* promoter expression was evaluated in

a representative transgenic line by qualitative GUS staining (Jefferson et al., 1987) and visualized

under a Leica DMi8 light microscope (Leica; Wetzlar, Germany).

194 RNA isolation, cDNA synthesis, and quantitative PCR assays (qPCR)

Total RNA was purified from Col-0, *atg5*, and *atg7* at different germination time points (0, 12, 24,

30, 36, 42 hours of imbibition) and seed stages (early green, late green, and brown seeds) as

described by Oñate-Sánchez and Vicente-Carbajosa (2008). RNA quality and concentration were

assessed using a Nanodrop® ND-1000 spectrophotometer (Thermo Fisher Scientific). cDNA was

synthesized from 1 μg of RNA using the RevertAid First Strand cDNA Synthesis Kit (Thermo

Fisher Scientific) following the manufacturer's instructions, and samples were stored at -20ºC.

qPCR was performed on an Eco® Real-Time PCR System (Illumina, San Diego, CA, USA) using

primers for *AtATG5*, *AtATG7*, and *AtACT8* (Graeber et al., 2011; Iglesias-Fernández et al., 2014) as

a reference gene. The 10 μl reaction contained cDNA, FastStart SYBR Green Master (ROX;

Hoffmann-La Roche, Basel, Switzerland), primers, and water. Thermal cycling conditions were

95ºC for 10 min, followed by 40 cycles of 95ºC for 10 sec and 60ºC for 30 sec. A melting curve

was used to calculate melting temperatures. Primer efficiencies (E) were estimated from cDNA

serial dilutions, using the formula $E = (10^{-1}/\text{regression line slope} - 1) \times 100$. Expression levels

were determined by the cycle threshold (CT) in the exponential phase of PCR (Pfaffl, 2001).

RNA library construction and RNA-seq analysis

Total RNA was isolated from Col-0 and *atg7* seeds, with RNA integrity confirmed using a

Bioanalyzer 2100 (Agilent). Library construction and RNA sequencing (PE150) were performed on

Illumina NovaSeq™ 6000 platforms (Illumina, San Diego, CA, USA) by Novogene Genomics

Service (Novogene, Beijing, China) with three biological replicates per genotype and treatment.

mRNA was purified from total RNA using poly-T oligo-attached magnetic beads, fragmented, and

synthesized into cDNA. The cDNA was subjected to end repair, A-tailing, adapter ligation,

amplification, and purification. Library quantification was done using Qubit and real-time PCR,

with size distribution checked using a bioanalyzer. Clean reads were obtained by removing low-

quality reads and adapters and aligned to *the A. thaliana* reference genome (TAIR10) using Hisat2

v2.0.5 (Kim *et al*., 2019; Anders *et al*., 2015). Differential expression analysis was conducted with

DESeq2 (1.42.0; Love *et al*., 2014), with adjusted p-values (padj) ≤ 0.05 indicating differentially

expressed genes. Principal component analysis (PCA) was performed using R (version 4.3.1), and

visualization was done using the FactoExtra package. Gene set enrichment analysis (GSEA) was

carried out with the gseGO function in the clusterProfiler package, with visualization via ridge plots

using ggridges. Heatmaps were constructed using R studio and InstantClue v.0.10.10.dev-snap

(Nolte *et al*., 2018) and Gene Ontology (GO) term enrichment analysis was done using ShinyGO

0.80 (Ge *et al*., 2020).

Fixation, embedding, and sectioning of material for histochemistry

Arabidopsis Col-0, dry and imbibed (24, 48 hoi) seeds were infiltrated with formaldehyde: acetic

acid: ethanol: water (3.5:5:50:41.5 by vol.) for 45 min under vacuum (41 mbar) and then incubated

at 4ºC for 3 days with gentle shaking as described by Ferrándiz *et al.* (2000) with modifications.

Increasing concentrations of ethanol solutions were used to dehydrate sections and subsequently

ethanol was progressively replaced with HistoClear (National Diagnostics, Hessle Hull, England)

and then embedded in paraffin. Thin sections of 8 µm were performed with a Leica HistoCore

NANOCUT R microtome (Leica) and collected on glass slides. Before histological staining and

fluorescent immunolocalization, samples were de-waxed, by repeating the washes in reverse order.

Fluorescent immunolocalization

Seed histological sections were washed with Phosphate Buffered Saline (PBS) and digested with 1 mg/ml proteinase-K (Hoffmann-La Roche). Slides were then incubated for 30 minutes at room temperature with a blocking solution (Blocking Reagent Roche), followed by overnight incubation with primary Anti-APG8A/ATG8A antibody (ab77003; Abcam, Cambridge, UK) at a 1:1000 dilution in 3% BSA and 0.5% Triton X-100 (Merck). After washing with Tris-Buffered Saline (TBS) containing 5 mM Na-Azide, sections were incubated for 3 hours with secondary Horse Anti-Rabbit IgG Antibody, DyLight™ 488 (DI-1088-1.5; Thermo Fisher Scientific) at a 1:100 dilution in the same buffer. Finally, samples were washed in PBS and water, mounted in CitifluorTM (Sciences Services GmbH, München, Germany), and examined using a Zeiss LSM 880 confocal microscope (excitation 493 nm, emission 512 nm).

Polysaccharide and protein histological staining

Seed sections were stained with PAS-NBB: 0.5% (w/v) Periodic Acid (Merck) plus Schiff's reagent (Merck) to detect polysaccharides (pink staining), and with 1% (w/v) Naphthol Blue Black (Merck) to visualize proteins (blue staining). Microscopy analyses were performed on a Zeiss LSM 880 microscope in bright field mode, and the images were captured and processed with the Zen Blue Edition software (Zeiss).

Embryo lipid staining

The coat of Col-0, *atg5*, and *atg7* imbided seeds (24, 48 hoi) was dissected using hypodermic needles, and embryos were collected in PBS. Then, they were fixed in 4% paraformaldehyde in PBS for 1h under vacuum (41 mbar), washed twice in PBS, and incubated o/n in ClearSee solution (10% xylitol, 15% sodium deoxycholate, 25% urea in distilled water) with gentle agitation. Embryos were stained o/n with 0.05% Nile Red in ClearSee, washed for 30 min and then 1h in ClearSee solution, and mounted in the same solution for imaging on a Zeiss LSM 880 microscope (excitation 561, emission 620 nm) and subsequently processed with Airyscan. Lipid quantification in embryo cells was analyzed using Fiji software (Schindelin *et al.*, 2022) and has been expressed as a percentage of lipid area/cell area, in a minimum of 10 distinct embryos per line and germination time.

Western blot

Total protein was extracted from 20 mg of Col-0, *atg5*, and *atg7* seeds at different germination time points (0, 36, 48, 72 hoi). Seeds were frozen in liquid nitrogen, ground using a Mikro-Dismembrator S (Sartorius AG, Göttingen, Germany), and mixed with 200 µL of extraction buffer (100 mM Tris-HCl pH 6.8, 4% SDS, 20% glycerol, 200 mM DTT). The samples were vortexed, heated at 95ºC for 5 minutes, and centrifuged. Supernatants were recovered, and protein concentration was measured with the Bio-Rad Protein Assay (Bio-Rad, Hercules, CA, USA). Twenty-five µg of protein were analyzed using SDS-PAGE with a Mini-PROTEAN Tetra system (Bio-Rad), followed by transfer to a nitrocellulose membrane (Bio-Rad). Membranes were blocked in TBST with 5% BSA and probed with anti-ABI5 antibody (1:4,000; Agrisera, Vännäs, Sweden) overnight at 4ºC. After washing, membranes were incubated with goat anti-rabbit ALP-conjugated antibody (1:10,000; Agrisera) for 1 hour. Detection was done using NBT/BCIP solution (Merck), and band intensity was quantified using ImageJ software.

For western blot analysis of siliques, total protein was extracted from green siliques of entire stems as described in Sánchez-Vicente *et al.*, 2024. The tissue was homogenized in extraction buffer (100mM Tris-HCl, 150mM NaCl, 0.25% NP-40) with 1mM PMSF and 1X protease inhibitor. After centrifugation, protein concentration was quantified, and 70 µg were loaded on an SDS-PAGE. Proteins were transferred to a PVDF membrane (Millipore) and probed with antibodies, including anti-bZIP67 (Biomedal), anti-Actin (Sigma-Merck), and ECL-Peroxidase-labelled anti-rabbit and anti-mouse antibodies (Amersham). Detection was done using the ECL Advance kit (Amersham) and chemiluminescence was detected with the ChemiDoc Imaging System (Bio-Rad).

Yeast two-hybrid (Y2H) assays

For Y2H assays, a mating-based yeast method and the *HIS3* reporter gene were used. A *Saccharomyces cerevisiae* J694α (α mating type) library containing a plasmid (pDEST™22, GAL4 activation domain, AD; Thermo Fisher Scientific) with 1,200 Open Reading Frames of TFs in *A. thaliana* was employed (Castrillo et al., 2011). Yeasts containing AtABI3 and AtABI5 were selected as prey. The coding sequence of *AtATG8* was amplified from *A. thaliana* cDNA, subcloned into pENTR3C, and then cloned into the pDEST™32 vector (GAL4 binding domain, BD) by Gateway strategy. This construct was used to transform *S. cerevisiae* YM4271a (a mating type) by

300 PEG method and used as bait. Diploid cells were grown on solid DOB media (Bioworld, Dublin,

301 OH, USA) lacking leucine and tryptophan. If AtATG8 interacts with AtABI3 and AtABI5 fused to

302 BD, GAL4 TF is reconstituted, and Histidine (H) auxotrophy occurs. Increasing concentrations of

303 3-amino-1,2,4-triazole (3-AT; Merck) were added to avoid leakage of the *HIS*3 gene. Positive

304 colonies appeared after 2-5 days of incubation at 28ºC.

Semi *in vivo* co-immunoprecipitation (CoIP) and pull-down assays

For CoIP assays, proteins were extracted from transgenic lines *35S:cMyc-ABI5* and *ProUBQ10-*

GFP-ATG8A. Seedlings were grown in MS and then treated or not with 3-MA 5mM for 24 hours.

Proteins were extracted from 4-day-old Arabidopsis seedlings with a buffer containing 50mM Tris–

HCl, pH 7.5, 250mM NaCl, NP40 1%, 1mM PMSF and 1X cOmplete® EDTA-free proteases in-

hibitors (Sigma). Extracts were cleared by centrifugation and protein concentration was determined

by Bradford assay. 1,5 mg of total soluble proteins were maintained in darkness at room tempera-

ture for 30 min. After treatment, extracts were incubated for 30 min in ice and then GFP-tag was

magnetically labeled and immunoprecipitated using µMACS GFP Isolation Kit (MiltenyiBiotec)

following manufacturer's protocol. The proteins were visualized using anti-ABI5 (Biomedal,

1:10,000), anti-GFP-HRP (MiltenyiBiotec, 1:1,000) and ECL-Peroxidase-labelled anti-rabbit

(Amersham, 1:10,000) antibodies.

318
319

Results

The *AtATG5* and *AtATG7* genes are expressed in embryos of *Arabidopsis thaliana* during seed maturation and germination

To study the impact of autophagy on seed biology, a selection of 68 *A. thaliana* autophagy genes

was analyzed based on GO classification (see Materials and Methods). Their expression patterns

were obtained from the Bio-Analytic Resource for Plant Biology (BAR) public database, and a

heatmap representation was created (Fig. S1; Data S1). Among these genes, *ATG5* and *ATG7* were

chosen for further study, as they are single-copy genes in Arabidopsis (*AtATG5*: *At5g17290*,

AtATG7: *At5g45900*), and are essential components of the autophagy machinery in both plants and

animals (Xie and Klionsky, 2007; Minina *et al*., 2018; Fraiberg and Elazar, 2020). Both genes

showed high expression during late silique maturation and dry seed stages (DS) (Fig. 1a).

We further investigated the spatial expression patterns of *AtATG5* and *AtATG7* using stable transgenic Arabidopsis (Col-0) lines with *AtATG5* and *AtATG7* promoters driving GUS expression. Histochemical staining confirmed their ubiquitous expression in both adult plants (75 days) and seedlings (15 days), with GUS activity observed in the vascular elements of roots and leaves (Fig. 1b and 1c). Additionally, expression was also found in pollen grains and in seeds at late silique maturation stages. In seedlings, both genes were expressed in the stem apical meristem (SAM) and weakly in cotyledons (Fig. 1b and 1c).

Quantitative PCR (qPCR) was used to analyze *AtATG5* and *AtATG7* expression during embryogenesis, seed maturation, and germination, with *AtACT7* as a reference gene which showed stable expression throughout the period of study (Graeber *et al*., 2011; Fig. S2). Expression levels of both genes peaked during early green (EG) to late green (LG) and brown (B) stages of silique development, with maximal transcript accumulation in the B stage (Fig. 2b). During seed germination, *AtATG5* and *AtATG7* mRNAs were detected at high levels in the DS stage (Fig. 2c), followed by a sharp decline in the early stages of imbibition (12 hoi), and a slight increase between 24-30 hoi, when seeds reached t50 (time to 50% germination). GUS histochemical staining in *PAtATG5::uidA* and *PAtATG7::uidA* transgenic lines showed GUS activity at 24 and 48 hoi in the embryo but not in the endosperm. At 24 hoi, GUS activity was localized in the radicle transition zone, and by 48 hoi, the signal expanded to the hypocotyl and cotyledons, with increasing intensity as they developed (Fig. 2c). These findings suggest that *AtATG5* and *AtATG7* play important roles during seed maturation and germination.

Seed germination is impaired and significantly delayed in *atg5* and *atg7* knockout mutants in the presence of ABA during imbibition

To investigate the relevance of *AtATG5* and *AtATG7* genes in seed germination, we analyzed homozygous T-DNA insertion mutant lines *atg5-1* and *atg7-2* (Hofius *et al*., 2009; Yoshimoto *et al*., 2009; Avin-Wittenberg *et al*., 2015; Barros *et al*., 2017; Erlichman *et al*., 2023) under both control conditions and in the presence of 1 μM ABA. The ABA-insensitive *abi5-7* mutant (Nambara *et al*., 2002; Albertos *et al*., 2015) was used as a negative control. The absence of *AtATG5* and *AtATG7* transcripts in the *atg5* and *atg7* mutants was confirmed by qPCR, while *abi5-7* exhibited basal expression (Fig. S3).

Seed Coat Rupture (SCR) and Endosperm Rupture were scored in Col-0, *atg5*, and *atg7* KO mutant lines (Fig. 3). Under control conditions, *atg5*, *atg7*, and *abi5-7* mutants showed a significant delay

in germination, evidenced by an increased t50 for both SCR and ER compared to Col-0 (Fig. 3). While 1 µM ABA had minimal effect on Col-0 and *abi5-7* SCR, a notable delay in Endosperm Rupture was observed for *atg5* and *atg7* mutants, which did not reach t_{50} within the observed period. In contrast, Col-0 seeds reached t_{50} at 69.3 ± 4.2 h, while *abi5-7* seeds showed reduced ABA sensitivity, reaching t_{50} at 55.6 ± 1.2 h.

Analysis of *PAtATG5::uidA* and *PAtATG7::uidA* imbibed seeds revealed that the spatial expression and intensity of GUS activity were not significantly altered in the presence of 1 µM ABA compared to control conditions (Fig. S4a). Additionally, RNA-seq analysis (FPKM; Fragments Per Kilobase Million, see next section) of *AtATG5* and *AtATG7* expression in the presence of ABA during imbibition showed results consistent with those observed under control conditions (Fig. S4b).

Comprehensive transcriptomic profiling of *Arabidopsis thaliana atg7* mutant during seed imbibition reveals important changes in gene expression compared to the WT

To analyze transcriptional changes during seed germination in Col-0 and *atg7* mutants, under conditions with and without 1 µM ABA, RNA-seq analysis was performed on dry and imbibed seeds collected at 0, 24, 48, and 96 hours of imbibition (hoi) (Fig. S5a). These time-points correspond to t_0, t_{25}, t_{75}, and t_{100}, representing the time to reach 100% germination (Fig. S5a). Total RNA was extracted at each time point, and after library preparation, RNA sequencing was performed using the DESeq2 package in RStudio (see Materials and Methods). The raw DESeq2 output, including normalized counts, $\log_2$FoldChange, p-value, and adjusted p-value (padj; FDR corrected), is provided in Supplementary Data S2 and S4.

The hierarchical clustering analysis (Fig. S5b) revealed that the samples cluster primarily by their state (dry or imbibed) and secondarily by time points for imbibed samples. A clear distinction between mutant and WT samples was observed at each time point, especially for imbibed seeds. The principal component analysis (PCA) results (Figs. S5c-d) showed that the first Principal Component (PC1) explained most of the variance (78.2%), associated with differences between dry and imbibed samples. The second Principal Component (PC2) accounted for 15.9% of the variance and mainly reflected the time progression from 24h to 96h in imbibed seeds. Differences between the 48 hoi and 96 hoi time points may also be influenced by ABA treatment. Col-0 and *atg7* mutants were most clearly separated at 24 hours without ABA (Fig. S5c-d), and these conditions were chosen for differential gene expression analysis (DEG; Data S2).

The volcano plot of DEG (Fig. 4a) revealed higher expression of genes related to storage compound

synthesis and mobilization in *atg7* mutant compared to WT. Gene set enrichment analysis (GSEA)

(adjusted p-value < 0.05) identified ten pathways significantly induced and five significantly

repressed in the mutant (Fig. 4b; Supplementary Data S3a). GO categories like lipid storage and

seed maturation were overrepresented among induced categories, with lipid storage showing the

highest Normalized Enrichment Score (NES). In contrast, response to ER stress (ERS), including

autophagy, was repressed and showed the most negative NES. The repression of genes related to

ERS and the induction of lipid storage genes suggests a clear trend in the mutant, with genes in

these categories clustering at opposite ends of the list based on their Log_2FC.

Fig. 4c presents a lollipop plot illustrating the Log_2FC of significant genes within selected

categories (lipid storage, seed maturation, and ERS response). Genes involved in seed maturation,

like *AtCRC* (*CRU3*), *AtCRA1* (*12S SSP*, cruciferins), *AtCLO1*, and *SEIPIN1*, were identified.

AtCLO1 encodes a caleosin protein involved in lipid body formation, while *SEIPIN1* is crucial for

lipid droplet biogenesis in embryos. Additionally, lipid storage-related genes like *OLE1*, *OLE2*, and

OLE3 (oleosins) and *Late Embryogenesis Abundant* (*LEA*) protein-encoding genes (*SSLEA*, *LEA29*,

LEA7) were also identified. Interestingly, two aquaporin-encoding genes, including *TIP3;1*

(*ALPHA-TIP*), were differentially expressed. *TIP3;1* is notably the most expressed autophagy-

related gene in seeds (Fig. S1).

As expected, genes related to ERS, including *BIP3* (HSP70 family) and *IRE1C* (involved in UPR in

the ER), were repressed in the mutant (Fig. 4c). These findings suggest autophagy's involvement in

lipid and protein metabolism during early seed imbibition. Furthermore, 18 out of 38 genes in the

seed maturation category were identified as potential ABI5 target genes through DAP-seq analysis

(AGRIS Plant TF database linked to ShinyGO platform, Yilmaz *et al.*, 2011; O'Malley *et al.*, 2016;

Ge *et al.,* 2020) (Data S3c).

Although other pathways like the response to oxygen levels and light were also identified as over-

expressed in the mutant, lipid storage, ERS, and seed maturation categories were prioritized due to

their significant differences and relevance in seed biology.

Histochemical analysis of protein and lipids in *atg* mutants during seed

imbibition reveals autophagy-related changes in the accumulation and

dynamics of storage compounds

The transcriptomic analysis described above revealed a significant upregulation of genes associated
with storage compounds in imbibed seeds of the *atg7* mutant compared to the WT (Fig. 4). To
further investigate this finding, the detection of seed lipids and proteins was carried out in both *atg5*
and *atg7* mutants, as well as in the WT, at 24 and 48 hoi.

Microscopic examination of embryos using Nile red for lipid staining (Fig. 5; Fig. S6: negative
control) demonstrated similar levels in *atg5* and *atg7* mutants and Col-0 at 24 hoi (Fig. 5a).
However, a quantitative analysis of the percentage of lipid /cell area in the radicle region for each
line at 48 hoi (Fig. 5b) showed that the percentage of lipid/cell area is significantly lower in mutants
(*ca.* 30%) than in WT (ca. 45%). The *atg7* mutant exhibits significantly reduced expression of 15
genes related to fatty acid biosynthesis, in addition to 2 genes specifically involved in fatty acid
desaturation and 5 genes associated with fatty acid elongation, compared to Col-0 at 24 hours
(Figure S7a, Data S3d). These genes encode enzymes that catalyze 7 reactions involved in fatty acid
synthesis and 1 reaction in the extension of palmitoyl-CoA and stearoyl-CoA into very-long-chain
acyl-CoAs (KEGG; Figure S7b, c). Interestingly, three genes encoding lipases related to lipid
catabolism (*MPL1*, *MAGL4*, and *MAGL12*) as well as *ACX1*, which encodes the first enzyme in the
β-oxidation pathway, are significantly induced in the *atg7* mutant compared to the WT during
imbibition (Figure S7).

Likewise, our analyses revealed differences when *atg7* was compared with WT in the ontology term
of genes related to seed maturation proteins. Examples include *AtCRC* (*CRU3*) and *AtCRA1*, which
encode *12S-SSP* (cruciferins), highly abundant SSP in seeds of *A. thaliana.* These proteins
accumulate in PSV during seed maturation. To investigate SSP in Col-0, *atg5*, and *atg7* seeds, in
DS and imbibed stages, a histochemical analysis was performed. Longitudinal sections of *A.*
thaliana seeds (DS and water-imbibed at 24 and 48 hoi) were stained using Periodic acid–Schiff
(PAS) and Naphthol Blue Black to detect insoluble polysaccharides (mainly cellulose and starch)
and proteins, respectively. As shown in Fig. 6a–f, both DS and imbibed seeds exhibited abundant
proteins (blue stained). In Col-0 DS, proteins accumulated within well-defined PSV are
characterized by rounded structures (Fig. 6a-b). On the contrary, both *atg5* and *atg7* mutants,
although rich in proteins as well, exhibited altered organization in their cytoplasm, as the rounded
structures seen in WT embryo cells were not clearly distinguished, especially in the radicle (Fig. 6a-
r). These differences in cytoplasmic organization, relative to protein accumulation in radicle cells of
atg mutants, persist at 24 hoi (Fig. 6g-l). At 48 hoi, proteins were completely degraded in both Col-
0 and mutants after radicle emergence (Fig. 6m–r).
462

Immunolocalization of the ATG8 protein in PSVs of WT but not in *atg* mutants supports an autophagy-like mechanism involved in reserve mobilization during seed germination

The insights presented above from genetic, transcriptomic, and histochemical analyses during seed imbibition and reserve mobilization prompted us to explore the molecular link between storage compound mobilization and autophagy. To this end, we performed immunofluorescence labeling using a primary antibody specific to ATG8a protein (Miklaszewska *et al.*, 2023) participating in autophagosome capture of cargoes (Fig. 7a-l, Fig. S8). The presence of the ATG8a protein was detected in longitudinal sections of Col-0 seeds imbibed at 24 hoi. As illustrated in Figs. 7a-c, ATG8a is predominantly localized within PSVs in the radicle of Col-0 seeds. Interestingly, this localization was practically absent in *atg* mutants, specially in*atg7* embryos (Fig. 7e-g, i-k). Secondary/primary antibodies and empty negative controls did not show fluorescence signals as expected (Figs. S8a-i). Examining coarse sections (8 μm) of paraffin samples through differential interference contrast microscopic analysis revealed that the cellular organization of seed tissues had been well preserved (Fig. 7d, h, l; Fig. S8c, f, i). Although these data do not rule out the presence of other ATG8 proteins in PSVs in the radicle of *atg5* and *atg7* embryos, our histochemical assays support the participation of autophagy not only in lipid and protein metabolism, as suggested by our transcriptomic data, but also in its cytoplasmic organization of embryo cells throughout seed germination.

Transcriptomic profiling of *atg7* imbibed seeds reveals a connection of autophagy to ABA responses and genes regulated by ABI5

ABA is a hormone that regulates seed development and germination. The autophagy mutants exhibited delayed germination compared to WT, particularly with 1 μM ABA during imbibition (Fig. 3). To explore the molecular basis of ABA responses in the *atg7* mutant, we compared transcriptional differences during seed imbibition in the absence and presence of ABA between Col-0 and the *atg7* mutant (Fig. S5). The ABA treatment at 24 hoi had no significant effect (Fig. S5). Therefore, we compared the transcriptomes of germinating *atg7* and Col-0 seeds at 96 hoi with ABA and 48 hoi without ABA (Figs. 4, S5; Data S4 and S5). For differential expression analysis, we used the model: ~ Treatment + Mutant + Treatment:Mutant, and performed a likelihood ratio test against the reduced model ~ Treatment + Mutant to assess the interaction term's significance. Genes were con-

sidered significantly differentially expressed when the adjusted p-value was < 0.05 and the $\log_2$fold

change ($\log_2$FC) was $\geq \pm 1$.

In the volcano plot (Fig. 8a; Supplementary Data S4), DEGs show whether the effect of ABA dif-

fers between *atg7* and Col-0. A positive Log_2FC represents genes where ABA effect is more pro-

nounced in the mutant ("enhanced ABA effect genes"; n = 308), and a negative Log_2FC represents

genes with a reduced ABA effect in the mutant ("reduced ABA effect genes"; n = 885). Several

ABA-responsive genes, such as *EM1* and *AFP1* (Bensmihen *et al.*, 2002; López-Molina *et al.*,

2003; Du *et al.*, 2024), are highlighted.

GO term analysis (ShinyGO platform) revealed significant enrichment of genes related to the "re-

sponse to chemical" GO term in the *atg7*-reduced gene group (Fig. 8b, top right panel). In contrast,

the *atg7*-enhanced gene group showed enrichment for "response to abiotic stimulus," particularly

ABA response (Fig. 8b, top left panel).

268 of the 1,193 (22.5%) DEGs identified were previously identified as potential *ABI5* target genes

(Fig. 8b, bottom panels; Supplementary Data S5c and S5d). Of these, 36 genes were classified in

the ABA response category (Fig. 8c; Supplementary Data S5e, S5f). Fig. 8C presents a lollipop plot

illustrating the Log_2FC of these 36 genes (adjusted p-value < 0.05).

The autophagy protein ATG8 directly interacts with the TF ABI5 both *in*

vivo and *in vitro*, supporting its modulation through autophagy

The transcriptomic analysis previously described supported the role of ABA in seed development

and suggested the connection of the ABI5/ABI3 signalling to autophagy during seed germination.

To deepen our findings linking these TFs with the autophagy machinery, we focused on ABI5, a

bZIP TF primarily expressed in seeds and associated with suppression of germination and post-

germination growth under adverse conditions (López-Molina *et al.*, 2001; Albertos *et al.*, 2015).

ABI5 accumulation was systematically monitored in the *atg5* and *atg7* mutants, both in the DS and

during imbibition, compared to the WT seeds. Western blot assays were performed at 0, 36, 48, and

72 hoi using a primary antibody specific for the ABI5 protein (Fig. 9a). The signal intensities of the

ABI5 multiple forms in the Western blot assays were quantified using ImageJ software and plotted

in Fig. 9b. This Fig. shows that multiple forms of the ABI5 are clearly detected in WT seeds in DS

(0 hoi) but shows increased levels in autophagy mutants *atg5* (nearly 2-fold) and *atg7* (about 2.5-

fold). This difference in ABI5 accumulation was further monitored at different time points during imbibition (Figs. 9a and 9b) where it shows a temporal decline until untraceable levels around 72 hoi. These findings indicate that autophagy can be important in modulating ABI5 accumulation during seed imbibition. Furthermore, we verified that another bZIP factor expressed during seed development, bZIP67. This TF is the closest homolog of ABI5, and was previously described as a regulator of omega-3 fatty acid content in Arabidopsis seeds (Mendes *et al*., 2013; Sánchez-Vicente *et al*., 2024).bZIP67 also accumulates to a greater extent in autophagy mutants *atg5* and *atg7* and shows decreased levels in *ATG5* and *ATG7* overexpression lines (Fig. S9a). Moreover, bZIP67 contains three possible ATG8-interacting motifs (AIMs) within its sequence (Fig. S9b). Interestingly, genes such as *cruciferin 1* (*CRU1*), *CRU2*, *CRU3*, *seed storage albumin 1* (*AT2S1*) and *fatty acid elongation 1* (*FAE1*), previously recognized as bZIP67 target genes with roles in protein and lipid metabolism in seeds (Mendes *et al*., 2013), were enhanced by ABA in *atg7*-imbibed seeds compared to Col-0, as revealed by our transcriptomic analysis (Fig. S9c).

ABI5 serves as a master regulator in ABA signaling cascades within seeds as part of a complex with ABI3 (a B3 TF). These proteins play critical roles as seed-specific downstream ABA targets and are recognized as key repressors of germination (Sajeev *et al*., 2024). Sequence analysis revealed that both ABI3 and ABI5 contain several AIMs, with 6 and 5 potential AIMs, respectively (Fig. S10). This raised the question of whether ABI3 and ABI5 could physically interact with components of the autophagy machinery, particularly ATG8, during seed germination. To investigate this possibility, we examined their interaction with ATG8 using yeast two-hybrid (Y2H) assays (Fig. 9c). We focused on the ATG8H protein, the major ATG8 gene expressed during seed germination (https://bar.utoronto.ca/) and fused its coding sequence to the GAL4 DNA binding domain as bait. Yeast cells expressing the BD-ATG8H fusion along with either AD-ABI3 or AD-ABI5 were able to grow in selective medium containing more than 60 mM 3-aminotriazole (3-AT). This result, in contrast to the lack of growth in control cells or those expressing only AD-ATG8H, indicates that both ABI3 and ABI5 can indeed interact with ATG8 *in vitro* (Fig. 9c). With this information, we checked again the interaction by *in vivo* CoIP assays using the *35S:cMyc-ABI5* and *ProUBQ10-GFP-ATG8A* lines (Fig. 9d). Protein extracts from 4-day old seedlings, treated or not with the autophagy inhibitor (3-MA, 5 mM),were mixed and analyzed. The *GFP-ATG8A* protein was selected as bait and immunoprecipitated with anti-GFP beads, and the presence of *cMyc-ABI5* was detected by the anti-ABI5 antibody. Interestingly, the ABI5-ATG8A interaction was detected mainly in the presence of 3-MA, indicating that autophagy may control the ABI5 abundance during early stages of development.

Discussion

In recent years, autophagy has been increasingly linked to a diverse array of plant biological processes, including development, nutrient recycling, and responses to both abiotic and biotic stresses. Although several studies have acknowledged its role in seed development (e.g., Michaeli *et al.* 2016; Minina *et al.*, 2018; Tarnowski *et al.*, 2020), the molecular mechanisms and targets involved remain partially understood. In this work, our aim was to explore its impact on seed germination in *A. thaliana* through the study of autophagy mutants *atg5* and *atg7* with a special focus on their connections to ABA signaling.

Autophagy mutants *atg5* and *atg7* show a delayed seed germination phenotype and hypersensitivity to ABA

AtATG5 and *AtATG7* are single-copy genes in the Arabidopsis genome, and their gain- and loss-of-function mutants have been used to study autophagy-related phenotypes (Hofius *et al*., 2009; Yoshimoto *et al*., 2009; Minina *et al*., 2018; Ohlsson *et al*., 2023). To investigate their role in germination, we examined their expression patterns during seed maturation and germination. Transcriptomic data from the BAR database showed increased expression of both genes in dry seeds, early imbibed seeds, late silique maturation, and mature pollen. These results were confirmed by qPCR and GUS histochemical staining (Fig. 1 and 2). The *ATG5* and *ATG7* promoters drove expression in vascular elements of roots, leaves, mature pollen, and seeds in late silique maturation, consistent with previous reports in Arabidopsis and other species (Thompson *et al*., 2005; Li *et al*., 2015; Yu and Hua, 2022). Expression was also observed in the stem apical meristem and cotyledons. Autophagy, associated with stress-enhanced microspore embryogenesis, plays a role in pollen germination, tube growth, and sperm cell development (Yan *et al*., 2023; Testillano, 2019; Berenguer *et al*., 2021), and is involved in forming xylem and phloem elements (Michalak *et al*., 2024).

Our results support earlier studies showing increased expression of *ATG* genes during seed development, especially during maturation (Di Berardino *et al*., 2018; Bedu *et al*., 2020). *atg5* and *atg7* mutants exhibit reduced nitrogen content, possibly due to defective nitrogen remobilization. Evolutionary conservation of ATG function was also observed in species like *M. truncatula* and *Zea mays* (Guiboileau *et al*., 2012; Li *et al*., 2015; Barros *et al*., 2017; Minina *et al*., 2018; Sera *et al*., 2019; Liu *et al*., 2023). *AtATG5* and *AtATG7* expression peaks during late seed maturation (Fig. 2a), coinciding with ABA increase, storage compound synthesis, and primary dormancy induction (Ali

et al., 2021). These findings led us to investigate the germination capacity of *atg5* and *atg7* seeds.

Germination assays showed significant delays in mutants compared to WT, particularly under ABA

treatment. Slower germination in *atg* mutants was reflected by increased t_{50} values (Fig. 3).

In transgenic seeds expressing *uidA* under the control of the *AtATG5* and *AtATG7* promoters, GUS

activity was observed in the radicle transition zone at 24 hoi, especially for *AtATG7*, and extended

to the hypocotyl and cotyledons at 48 hoi. This pattern aligns with reserve mobilization involving

SSP hydrolysis first in the embryonic axis and then in cotyledons (Müntz, 2007; Iglesias-Fernández

et al., 2014). These results suggest that *AtATG5* and *AtATG7* influence germination by controlling

storage mobilization in the radicle. Interestingly, F1 seeds from maternal *atg* mutants germinated

earlier, likely due to increased water permeability and normal autophagy in the embryo. Maternal

atg mutant seeds showed reduced mucilage content and structural changes in seed coat analysis

(Erlichman *et al.*, 2023).

In the presence of ABA, *atg5* and *atg7* mutants exhibited slower germination compared to WT.

However, the spatiotemporal expression of *AtATG5* and *AtATG7* was unchanged by ABA in the

seed imbibition medium (Fig. S4), suggesting ABA does not affect their expression but alters other

factors impacting germination. Our data suggest a role for autophagy in seed germination progres-

sion under both non-stress and ABA-induced stress conditions.

Transcriptomic analysis and reserve mobilization in *atg7* mutant seeds dur-

ing imbibition

We compared the transcriptomes of WT and *atg7* seeds at 24 hoi to identify autophagy-related

changes. DEGs analysis revealed significant transcriptional differences, with GO enrichment in

processes linked to storage compounds, especially lipids and proteins. We thus performed histo-

chemical analyses of protein bodies and lipid droplets in dry and imbibed seeds.

In *A. thaliana*, lipids are stored as TAG in lipid droplets (LD) stabilized by proteins like oleosins

and caleosins (Miklaszewska *et al.*, 2023). *AtOLE1* and *AtOLE2* are key seed oleosins for LD for-

mation and degradation (Simada *et al.*, 2008; Chen *et al.*, 2019; Shao *et al.*, 2019). At 24 hoi, *atg7*

seeds show increased expression of *AtOLE1*, *AtOLE2*, and *AtOLE3*, suggesting a compensatory

mechanism. Overexpression or suppression of oleosins delays germination, confirming their role

(*Siloto et al.*, 2006; Yuan *et al.*, 2021). DEGs also include *AtSEIPIN*, involved in LD biogenesis

(Zhao *et al.*, 2023), and *AtCLO1*, required for LD degradation and LD-vacuole interaction (Naested

et al., 2000; Kim *et al*., 2011; Poxleitner *et al*., 2006). Interactions between *AtATG8* and both

AtCLO1 and *AtOLE1* via AIMs support autophagy's role in LD dynamics (Marshall *et al*., 2019;

Miklaszewska *et al*., 2023). The upregulation of LD-structural protein genes in *atg7* seeds supports

autophagy's involvement in lipid mobilization.

Autophagy mutants (*atg5, atg7*) exhibit reduced TAG and FA levels (Minina *et al*., 2018). Fan et al.

(2019) showed that this reduction stems from impaired lipid recycling, not synthesis. Havé et al.

(2019) reported altered phospholipid/sphingolipid accumulation in *atg5*, affecting ER and plasma

membranes. Consistent with these, our Fig. 5 shows that while LD numbers in *atg7* seeds at 24 hoi

are similar to WT, lipid degradation is faster at 48 hoi, highlighting a role for autophagy in early

lipid mobilization.

The *atg7* mutant also shows reduced expression of FA biosynthesis, desaturation, and elongation

genes, confirming autophagy's role in lipid homeostasis (Bouchnak *et al*., 2023). Conversely, genes

related to FA catabolism, such as *MPL1*, *MAGL4*, *MAGL12*, and *ACX1* (first β-oxidation enzyme),

are upregulated. Genes like *CER8*, *LACS2*, and *LACS9*, traditionally linked to lipid synthesis, also

participate in catabolism (Bai *et al*., 2021). Our RNA-seq at 24 hoi reflects early transcriptional

changes corresponding to lipid shifts observed at 48 hoi (Fig. 5a), in line with Barros *et al*. (2021).

Together, these suggest a shift toward lipid breakdown in *atg7*.

Arabidopsis seeds store 12S globulins (cruciferins) and 2S albumins (napins) in PSVs (Baud *et al*.,

2008). *AtCRU1* and *AtCRU3* transcripts are elevated in *atg7* at 24 hoi. Histochemical analysis at DS

and 24 hoi shows PSV disorganization in the radicle of *atg5* and *atg7* seeds, which resolves by 48

hoi. This suggests two protein trafficking routes to PSVs: Golgi-dependent and autophagy-like

ERvt (Hills, 2004; Vitale *et al*., 2022). ATG8 immunolocalization in Col-0 at 24 hoi confirms its

presence in PSVs, while it is absent in *atg* mutants, indicating impaired ERvt trafficking and protein

degradation in the embryonic axis.

A connection of the delayed germination phenotype and ABA responses to

the stability of ABI5 in the *atg7* mutant

ABA is a key phytohormone regulating seed maturation, dormancy, and stress responses (Ali *et al*.,

2021). We found that *atg5* and *atg7* mutants exhibit delayed germination (Fig. 3), especially under

ABA treatment. Transcriptomic analysis comparing *atg7* and Col-0 with/without ABA revealed

significant DEGs, with ABA response genes notably enriched among upregulated genes in *atg7*.

About 22.4% of these DEGs are predicted ABI5 targets, suggesting altered ABA signaling.

ABI5, together with ABI3, regulates ABA-responsive genes suppressing germination and control-

ling preharvest sprouting (Koornneef *et al.*, 1984; Finkelstein, 1994; Abraham *et al.*, 2016; Matilla,

2022; Sajeev *et al.*, 2024). ABA modulates germination based on environmental cues (López-

Molina *et al.*, 2001). A connection between ABA signaling and autophagy involves NBR1, a selec-

tive autophagy receptor that interacts with ABI3, ABI4, and ABI5 (Tarnowski *et al.*, 2020). Expres-

sion pattern of *ATG7* (Figure S4) and *ABI5* (Albertos *et al.*, 2015) co-localizes in the radicle zone,

showing additional evidence of this relationship. These TFs have also been detected in ATG8 im-

munoprecipitation assays (Dagdas Y, personal communication). Moreover, in our co-IP assays, the

ABI5–ATG8 interaction is observed *in vivo*, and it becomes more pronounced when the autophagy

inhibitor (3-MA) is added to the medium. This indicates a direct link between ABI5 and autopha-

gic processes in the early stages of development. We confirmed direct interactions between ATG8

and ABI3/ABI5 using Y2H assays (Fig. 9), supporting the possibility that these TFs are autophagy

targets. Additionally, ABI5 degradation during germination follows a profile consistent with ATG8-

mediated autophagy (Fig. 9). In *atg5* and *atg7*, this degradation is slower, aligning with their ABA

hypersensitivity and delayed germination. Autophagy also affects bZIP67, a bZIP TF that corre-

sponds to ABI5 homolog. In *atg7*, both bZIP67 levels and expression of its targets (*CRU1, CRU2,*

CRU3, AT2S1, FAE1) are increased (Mendes *et al.*, 2013; Fig. S10), suggesting autophagic regula-

tion of bZIP67 during seed maturation. Altogether, our data support a model where autophagy

modulates ABA responses during seed germination by controlling the abundance of key TFs, ena-

bling rapid adaptation to environmental signals.

Conclusions

The data presented here highlight the role of autophagy not only in the accumulation of reserves

during seed maturation but also in the mobilization of storage compounds during early seed

imbibition in *A. thaliana*. Our findings support the role of autophagy in cell vesicle trafficking in

the radicle, contributing to the cellular dynamics of LD (lipid turnover) and PSV (protein storage

and mobilisation). Our findings also indicate a role for autophagy in mediating the ABA response

during early imbibition. This response involves the regulation of protein levels of selected TFs, such

as ABI5 and bZIP67, which are pivotal in specific environmental responses. Consequently,

autophagy emerges as a broad mechanism that controls how germinating seeds respond rapidly to

environmental cues (Fig. 10). Therefore, autophagy likely plays a crucial role in the seed

development, and germination programs and seed environmental perception (seed vigour) in *A.*
thaliana. However, the question of whether these processes in *A. thaliana* can be extrapolated to
other dicotyledonous species or even monocotyledonous species, with different seed anatomy and
storage compounds, needs further investigation. Our work sheds light on seed autophagy and its
fundamental importance in enhancing plant yield. Targeting autophagy genes associated with high
seed yield and vigour will be crucial to achieving this goal.

Acknowledgments

This work was supported by projects PID2023-148279OB-I00 and BIO2017-82873-R (PIs: R.I-F.
and J.V.C and), Quinoa4Med (PRIMA-AEI, ref: PCI2022-132988; P.I.: R.I.F.), and PID2023-
149447OB-I00, funded by MCIU/AEI/10.13039/ 501100011033, and project SA142P23 from the
Regional Government of Castile and Leon and "Escalera de Excelencia" CLU-2018-04 co-funded
by the FEDER Operative Program of Castile and Leon 2014–2020 Spain (to O.L.). Research grant
funded by Fundación Memoria de D. Samuel Solórzano Barruso-Universidad de Salamanca Ref
FS/8-2024 (IP: I. S-V.). We also thank the Spanish Research Network *EvoDevoSigNet* (RED2022-
134917-T, MICIU).

The authors kindly thank Prof. A. Matilla and Rosina Saviaar (Mondego Science) for the critical
reading of the manuscript.

Competing interests

711   Authors declare no conflict of interest

Author contributions

R.I.F. and J.V.C. conceived the project and reviewed the manuscript. R.I.F., J.V.C. and E.C. wrote
the manuscript and prepared the figures with contributions of all authors. R.I.F. performed the
bioinformatic analysis. E.C., M.A., and E.P. conducted all the immunohistochemical assays under
the supervision of R.I.F. E.P. performed the expression analyses using qPCR. M.G.C. carried out
the cloning of the *pATG5/7:uidA* lines and those used for the yeast two-hybrid assays under
supervision of J.V.C. E.C. performed the Western blot assays for ABI5 and ABI3. O.L. and I.S.V.
conducted the assays for bZIP67 and co-IPs. M.A.D. performed the germination assays.

Data availability

The data obtained from the RNA sequencing (RNA-Seq) analysis have been uploaded to the Sequence Read Archive (SRA) of the National Center for Biotechnology Information (NCBI) under the reference PRJNA1190055. The codes used to perform the analysis with RStudio are deposited in a repository on GitHub (https://github.com/RaquelIglesias/Autophaghy-in-Arabidopsis-seed-imbibition).

References

Abraham Z, Iglesias-Fernández R, Martínez M, Rubio-Somoza I, Díaz I, Carbonero P, Vicente-Carbajosa J. 2016. A developmental switch of gene expression in the barley seed mediated by HvVP1 Viviparous-1 and HvGAMYB Interactions. *Plant Physiology* **170**: 2146–2158. doi: https://doi.org/10.1104/pp.16.00092

Albertos P, Romero-Puertas MC, Tatematsu K, Mateos I, Sánchez-Vicente I, Nambara E, Lorenzo O. 2015. S-nitrosylation triggers ABI5 degradation to promote seed germination and seedling growth. *Nature Communications* **6**: 8669. doi: https://doi.org/10.1038/ncomms9669

Ali F, Qanmber G, Li F, Wang Z. 2021. Updated role of ABA in seed maturation, dormancy, and germination. *Journal of Advanced Research* **35**: 199–214. doi: https://doi.org/10.1016/j.jare.2021.03.011

Anders S, Pyl PT, Huber W. 2015. HTSeq—a Python framework to work with high-throughput sequencing data. *Bioinformatics* **31**: 166–169. doi: https://doi.org/10.1093/bioinformatics/btu638

Avin-Wittenberg T, Bajdzienko K, Wittenberg G, Alseekh S, Tohge T, Bock R, Giavalisco P, Fernie AR. 2015. Global analysis of the role of autophagy in cellular metabolism and energy homeostasis in Arabidopsis seedlings under carbon starvation. *The Plant Cell* **27**: 306–22. doi: https://doi.org/10.1105/tpc.114.134205

Bai F, Yu L, Shi J, Li-Beisson Y, Liu J. 2022. Long-chain acyl-CoA synthetases activate fatty acids for lipid synthesis, remodeling and energy production in Chlamydomonas. *New Phytologist* **233**: 823-837. doi: https://doi.org/10.1111/nph.17813.

Barros JAS, Cavalcanti JHF, Medeiros DB, Nunes-Nesi A, Avin-Wittenberg T, Fernie AR, Araújo WL. 2017. Autophagy deficiency compromises alternative pathways of respiration following energy deprivation in *Arabidopsis thaliana. Plant Physiology* **175**: 62–76. doi: https://doi.org/10.1104/pp.16.01576

Barros JAS, Magen S, Lapidot-Cohen T, Rosental L, Brotman Y, Araújo WL, Avin-
Wittenberg T. 2021. Autophagy is required for lipid homeostasis during dark-induced senescence.
Plant Physiology **185**: 1542-1558. doi: https://doi.org/10.1093/plphys/kiaa120

Baud S, Dubreucq B, Miquel M, Rochat C, Lepiniec L. 2008. Storage reserve accumulation in
Arabidopsis: metabolic and developmental control of seed filling. In *Arabidopsis Book* **6**:
e0113.doi: https://doi.org/10.1199/tab.0113

Bedu M, Marmagne A, Masclaux-Daubresse C, Chardon F. 2020. Transcriptional plasticity of
autophagy-related genes correlates with the genetic response to nitrate starvation in *Arabidopsis*
thaliana. Cells **9**: 1021. doi: https://doi.org/10.3390/cells9041021.

Benjamini Y, Hochberg Y. 1995. Controlling the False Discovery Rate: a practical and powerful
approach to multiple testing. *Journal of the Royal Statistical Society, Series B (Methodology)* **57**:
289–300. doi: https://doi.org/10.3390/cells9041021

Bensmihen S, Rippa S, Lambert G, Jublot D, Pautot V, Granier F, Giraudat J, Parcy F. 2002.
The homologous ABI5 and EEL transcription factors function antagonistically to fine-tune gene
expression during late embryogenesis. *The Plant Cell* **146**: 1391–403. doi: https://doi.org/
10.1105/tpc.000869

Bentsink L, Jowett J, Hanhart CJ, Koornneef M. 2006. Cloning of DOG1, a quantitative trait
locus controlling seed dormancy in Arabidopsis. *Proceedings of the National Academy of Sciences*
of the United States of America **103**: 17042–17047. doi: https://doi.org/10.1073/pnas.0607877103

Bewley JD, Bradford KJ, Hilhorst HWM, Nonogaki H. 2013. Structure and composition In
Seeds, Bewley, JD, Bradford, KJ, Eds; Springer: New York, NY, USA, pp 85–131. doi:
http://dx.doi.org/10.1007/978-1-4614-4693-4

Bouchnak I, Coulon D, Salis V, D'Andréa S, Bréhélin C. 2023. Lipid droplets are versatile
organelles involved in plant development and plant response to environmental changes. *Frontiers in*
Plant Science **14**: 1193905. doi: http://dx.doi.org/10.3389/fpls.2023.1193905

Calero-Muñoz N, Exposito-Rodriguez M, Collado-Arenal AM, Rodríguez-Serrano M,
Laureano-Marín AM, Santamaría ME, Gotor C, Díaz I, Mullineaux PM, Romero-Puertas
MC, Olmedilla A, Sandalio LM. 2019. Cadmium induces reactive oxygen species-dependent
pexophagy in Arabidopsis leaves. *Plant Cell and Environment* 42, 2696-2714. doi:
10.1111/pce.13597

Carbonero P, Iglesias-Fernández R, Vicente-Carbajosa J. 2017. The AFL subfamily of B3
transcription factors: evolution and function in angiosperm seeds. *Journal of Experimental Botany*
68: 871–880. doi: https://doi.org/10.1093/jxb/erw458

Carrillo-Barral N, Rodríguez-Gacio MDC, Matilla AJ. 2020. Delay of Germination-1 DOG1: a
key to understanding seed dormancy. *Plants (Basel)* **9**: 480. doi: https://doi.org/
10.3390/plants9040480

Carlson M. 2019. org.At.tair.db: Genome wide annotation for Arabidopsis. R package version
3.8.2.

Castrillo G, Turck F, Leveugle M, Lecharny A, Carbonero P, Coupland G, Paz-Ares J,
Oñate-Sánchez L. 2011. Speeding *cis-trans* regulation discovery by phylogenomic analyses
coupled with screenings of an arrayed library of Arabidopsis transcription factors. *PloS ONE* **6**:
e21524. doi: https://doi.org/ 10.1371/journal.pone.0021524

Chen K, Yin Y, Liu S, Guo Z, Zhang K, Liang Y, Zhang L, Zhao W, Chao H, Li M. 2019.
Genome-wide identification and functional analysis of oleosin genes in *Brassica napus L. BMC*
Plant Biology **19**: 294. doi: https://doi.org/ 10.1186/s12870-019-1891-y

Chen L, Lv Q, Yang W, Yang H, Chen Q, Wang B, Lei Y, Xie Y. 2022. TaNBR1, a novel wheat
NBR1-like domain gene negatively regulates drought stress tolerance in transgenic Arabidopsis.
International Journal of Molecular Sciences **23**: 4519. doi: https://doi.org/10.3390/ijms23094519

Clough SJ, Bent AF. 1998. Floral dip: a simplified method for *Agrobacterium*-mediated
transformation of *Arabidopsis thaliana. The Plant Journal* **16**: 735–743. doi: https://doi.org/
0.1046/j.1365-313x.1998.00343.x

Di Berardino J, Marmagne A, Berger A, Yoshimoto K, Cueff G, Chardon F, Masclaux-
Daubresse C, Reisdorf-Cren M. 2018. Autophagy controls resource allocation and protein storage
accumulation in Arabidopsis seeds. *Journal of Experimental Botany* **69**: 1403–1414. doi:
https://doi.org/ 10.1093/jxb/ery012

Ding X, Zhang X, Otegui MS. 2018. Plant autophagy: new flavors on the menu. *Current Opinion*
in Plant Biology **46**: 113–121. doi: https://doi.org/ 10.1016/j.pbi.2018.09.004

Dobrenel T, Caldana C, Hanson J, Robaglia C, Vincentz M, Veit B, Meyer C. 2016. TOR
signaling and nutrient sensing. *Annual Review of Plant Biology* **67**: 261–285. doi: https://doi.org/
10.1146/annurev-arplant-043014-114648

Du C, Liu M, Yan Y, Guo X, Cao X, Jiao Y, Zheng J, Ma Y, Xie Y, Li H, Yang C, Gao C,
Zhao Q, Zhang Z. 2024. The U-box E3 ubiquitin ligase PUB35 negatively regulates ABA

signaling through AFP1-mediated degradation of ABI5. *The Plant Cell* **369**: 3277–3297. doi:
https://doi.org/ 10.1093/plcell/koae194

Ellgaard L, Helenius A. 2003. Quality control in the endoplasmic reticulum. *Nature Reviews*
Molecular Cell Biology **4**: 181–191. doi: https://doi.org/10.1038/nrm1052

Erlichman OA, Weiss S, Arkia MA, Ankary-Khaner M, Soroka Y, Jasinska W, Rosental L,
Brotman Y, Avin-Wittenberg T. 2023. Autophagy in maternal tissues contributes to Arabidopsis
seed development. *Plant Physiology* **193**: 611–626. doi: https://doi.org/10.1093/plphys/kiad350

Fan J, Yu L, Xu C. 2019. Dual role for autophagy in lipid metabolism in Arabidopsis. *The Plant*
Cell **31**: 1598–1613. doi: https://doi.org/10.1105/tpc.19.00170

Ferrándiz C, Liljegren SJ, Yanofsky MF. 2000. Negative regulation of the SHATTERPROOF
genes by FRUITFULL during Arabidopsis fruit development. *Science* **289**: 436–438. doi:
https://doi.org/10.1126/science.289.5478.436

Finch-Savage WE, Bassel GW. 2016. Seed vigour and crop establishment: extending performance
beyond   adaptation.   *Journal   of   Experimental   Botany*   **67**:   567–591.   doi:
https://doi.org/10.1093/jxb/erv490

Finkelstein RR. 1994. Mutations at two new Arabidopsis ABA response loci are similar to the *abi3*
mutations. *The Plant Journal* **5**: 765–771. doi: https://doi.org/10.1046/j.1365-313X.1994.5060765.x

Finkelstein RR, Gampala SS, Rock CD. 2002. Abscisic acid signaling in seeds and seedlings. *The*
Plant Cell **14**: S15–S45. doi: https://doi.org/ 10.1105/tpc.010441

Finkelstein RR. 2004. The role of hormones during seed development and germination. In: *Plant*
Hormones: Biosynthesis, Signal Transduction, Action! PJ Davies, ed (Dordrecht, The Netherlands:
Kluwer Academic Publishers), pp 549–573. doi: https://doi.org/ 0.1007/978-1-4020-2686-7_24

Fraiberg M, Elazar Z. 2020. Genetic defects of autophagy linked to disease. *Progress in*
Molecular   Biology   and   Translational   Science   **172**:   293–323.   doi: https://doi.org/
10.1016/bs.pmbts.2020.04.001

Ge SX, Jung D, Yao R. 2020. ShinyGO: a graphical gene-set enrichment tool for animals and
plants. *Bioinformatics* **36**: 2628–2629. doi: https://doi.org/10.1093/bioinformatics/btz931

Graeber K, Linkies A, Wood ATA, Leubner-Metzger GA. 2011. Guideline to family-wide
comparative state-of-the-art quantitative RT-PCR analysis exemplified with a Brassicaceae cross-
species seed germination case study. *The Plant Cell* **23**: 2045–2063. doi: https://doi.org/
10.1105/tpc.111.084103

Guiboileau A, Avila-Ospina L, Yoshimoto K, Soulay F, Azzopardi M, Marmagne A, Lothier

J, Masclaux-Daubresse C. 2013. Physiological and metabolic consequences of autophagy

deficiency for the management of nitrogen and protein resources in Arabidopsis leaves depending

on nitrate availability. *New Phytologist* **199**: 683–694. doi: https://doi.org/10.1111/nph.12307

Guiboileau A, Yoshimoto K, Soulay F, Bataillé M, Avice JC, Masclaux-Daubresse C. 2012.

Autophagy machinery controls nitrogen remobilization at the whole-plant level under both limiting

and ample nitrate conditions in Arabidopsis. *New Phytologist* **194**: 732–740. doi: https://doi.org/

10.1111/j.1469-8137.2012.04084.x

Havé M, Luo J, Tellier F, Balliau T, Cueff G, Chardon F, Zivy M, Rajjou L, Cacas JL,

Masclaux-Daubresse C. 2019. Proteomic and lipidomic analyses of the Arabidopsis atg5

autophagy mutant reveal major changes in endoplasmic reticulum and peroxisome metabolisms and

in lipid composition. *New Phytologist* **223**: 1461–1477. doi: https://doi.org/ 10.1111/nph.15913

Hills MJ. 2004. Control of storage-product synthesis in seeds. *Current Opinion in Plant Biology* **7**:

302–308. doi: https://doi.org/ 10.1016/j.pbi.2004.03.003

Hofius D, Schultz-Larsen T, Joensen J, Tsitsigiannis DI, Petersen NHT, Mattsson O,

Jørgensen LB, Jones JDG, Mundy J, Petersen M. 2009. Autophagic components contribute to

hypersensitive cell death in Arabidopsis. *Cell* **137**: 773–783. doi: https://doi.org/

10.1016/j.cell.2009.02.036

Holdsworth MJ, Bentsink L, Soppe WJJ. 2008. Molecular networks regulating Arabidopsis seed

maturation, after-ripening, dormancy and germination. *New Phytologist* **179**: 33–54. doi:

https://doi.org/10.1111/j.1469-8137.2008.02437.x

Iglesias-Fernández R, Vicente-Carbajosa J. 2022. A view into seed autophagy: from

development to environmental responses. *Plants* **11**: 3247. doi:

https://doi.org/10.3390/plants11233247

Iglesias-Fernández R, Wozny D, Iriondo-de Hond M, Oñate-Sánchez L, Carbonero P,

Barrero-Sicilia C. 2014. The AtCathB3 gene, encoding a cathepsin B-like protease, is expressed

during germination of *Arabidopsis thaliana* and transcriptionally repressed by the basic leucine

zipper protein GBF1. *Journal of Experimental Botany* **65**: 2009–2021. doi: https://doi.org/

10.1093/jxb/eru055

Jefferson RA, Kavanagh TA, Bevan MW. 1987. GUS fusions: beta-glucuronidase as a sensitive

and versatile gene fusion marker in higher plants. *The EMBO Journal* **6**: 3901–3907. doi:

https://doi.org/10.1002/j.1460-2075.1987.tb02730.x

Joosen RVL, Kodde J, Willems LAJ, Ligterink W, Van Der Plas LHV, Hilhorst HWM. 2010.
Germinator: a software package for high-throughput scoring and curve fitting of Arabidopsis seed
germination. *The Plant Journal* **62**: 148–159. doi: https://doi.org/ 10.1111/j.1365-
313X.2009.04116.x

**Jurkiewicz P, Melser S, Maucourt M, Ayeb H, Veljanovski V, Maneta-Peyret L, Hooks M,
Rolin D, Moreau P, Batoko H.** 2018. The multistress-induced Translocator protein TSPO
differentially modulates storage lipids metabolism in seeds and seedlings. *The Plant Journal* **96**:
274–286. doi: https://doi.org/10.1111/tpj.14028

Kellner R, De la Concepcion JC, Maqbool A, Kamoun S, Dagdas YF. 2017. ATG8 expansion:
A driver of selective autophagy diversification? *Trends in Plant Science* **22**: 204–214. doi:
https://doi.org/10.1016/j.tplants.2016.11.015

Kim, D, Paggi, JM, Park, C, Bennett, C, Salzberg, SL. 2019. Graph-based genome alignment
and genotyping with HISAT2 and HISAT-genotype. *Nature Biotechnology* **37**: 907–915. doi:
https://doi.org/10.1038/s41587-019-0201-4

Kim YY, Jung KW, Yoo KS, Jeung JU, Shin JS. 2011. A stress-responsive caleosin-like protein,
AtCLO4, acts as a negative regulator of ABA responses in Arabidopsis. *Plant and Cell Physiology*
52: 874–884. doi: https://doi.org/10.1093/pcp/pcr039

Koornneef M, Reuling G, Karssen CM. 1984. The isolation and characterization of abscisic acid-
insensitive mutants of *Arabidopsis thaliana*. *Physiologia Plantarum* **61**: 377–383. doi:
https://doi.org/ 10.1093/pcp/pcr039

**Li F, Chung T, Pennington JG, Federico ML, Kaeppler HF, Kaeppler SM, Otegui MS,
Vierstra RD.** 2015. Autophagic recycling plays a central role in maize nitrogen remobilization. *The
Plant Cell* **27**: 1389–1408. doi: https://doi.org/10.1105/tpc.15.00158

Liu Y, Xiong Y, Bassham DC. 2009. Autophagy is required for tolerance of drought and salt stress
in plants. *Autophagy* **5**: 954–963. doi: https://doi.org/10.4161/auto.5.7.9290

Liu Z, Yang Q, Wu P, Li Y, Lin Y, Liu W, Guo S, Liu Y, Huang Y, Xu P, Qian Y, Xie Q.
2023. Dynamic monitoring of TGW6 by selective autophagy during grain development in rice. *New
Phytologist* **240**: 2419–2435. doi: https://doi.org/10.1111/nph.19271

López-Molina L, Mongrand S, Chua NH. 2001. A postgermination developmental arrest
checkpoint is mediated by abscisic acid and requires the ABI5 transcription factor in Arabidopsis.
Proceedings of the National Academy of Sciences of the United States of America **98**: 4782–4787.
doi: https://doi.org/10.1073/pnas.081594298

Lopez-Molina L, Mongrand S, Kinoshita N, Chua NH. 2003. AFP is a novel negative regulator
of ABA signaling that promotes ABI5 protein degradation. *Genes and Development* **173**: 410–418.
doi: https://doi.org/10.1101/gad.1055803

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biology* **15**: 550. doi: https://doi.org/10.1186/s13059-014-
0550-8

Luo S, Li X, Zhang Y, Fu Y, Fan B, Zhu C, Chen Z. 2021. Cargo recognition and function of
selective autophagy receptors in plants. *International Journal of Molecular Sciences* **22**: 1013. doi:
https://doi.org/10.3390/ijms22031013

Marshall RS, Hua Z, Mali S, McLoughlin F, Vierstra RD. 2019. ATG8-binding UIM proteins
define a new class of autophagy adaptors and receptors. *Cell* **177** : 766–781e24. doi: https://doi.org/
10.1016/j.cell.2019.02.009. Erratum in: *Cell* (2022) **185**: 1101–1102

Masclaux-Daubresse C, Clément G, Anne P, Routaboul JM, Guiboileau A, Soulay F, Shirasu
K, Yoshimoto K. 2014. Stitching together the multiple dimensions of autophagy using
metabolomics and transcriptomics reveals impacts on metabolism, development, and plant
responses to the environment in Arabidopsis. *The Plant Cell* **26**: 1857–1877. doi: https://doi.org/
10.1105/tpc.114.124677

Matilla AJ. 2022. Exploring breakthroughs in three traits belonging to seed life. *Plants* **11**: 490.
doi: https://doi.org/10.3390/plants11040490

Maurel C, Boursiac Y, Luu DT, Santoni V, Shahzad Z, Verdoucq L. 2015. Aquaporins in
Plants. *Physiological Reviews* **95**: 1321–1358. doi: https://doi.org/10.1152/physrev.00008.2015

Mendes AT, Kelly AA, Erp HV, Shaw E, Powers SJ, Kurup S, Eastmondb PJ. 2013. bZIP 67
regulates the omega-3 fatty acid content of Arabidopsis seed oil by activating FATTY ACID
DESATURASE 3. *The Plant Cell* **25**: 3104–3116. doi: https://doi.org/10.1105/tpc.113.116343

Michaeli S, Galili G, Genschik P, Fernie AR, Avin-Wittenberg T. 2016. Autophagy in plants-
what's new on the menu? *Trends in Plant Science* **21**: 134–144. doi: https://doi.org/
10.1016/j.tplants.2015.10.008

Michalak KM, Wojciechowska N, Marzec-Schmidt K, Bagniewska-Zadworna A. 2024.
Conserved autophagy and diverse cell wall composition: Unifying features of vascular tissues in
evolutionarily distinct plants. *Annals of Botany* **133**: 559–572. doi: https://doi.org/
10.1093/aob/mcae015

Miklaszewska M, Zienkiewicz K, Klugier-Borowska E, Rygielski M, Feussner I, Zienkiewicz
A. 2023. CALEOSIN 1 interaction with AUTOPHAGY-RELATED PROTEIN 8 facilitates lipid
droplet microautophagy in seedlings. *Plant Physiology* **193**: 2361–2380. doi: https://doi.org/
10.1093/plphys/kiad471

Minina EA, Moschou PN, Vetukuri RR, Sanchez-Vera V, Cardoso C, Liu Q, Elander PH,
Dalman K, Beganovic M, Yilmaz JL, Marmon S, Shabala L, Suarez MF, Ljung K, Novák O,
Shabala S, Stymne S, Hofius D, Bozhkov PV. 2018. Transcriptional stimulation of rate-limiting
components of the autophagic pathway improves plant fitness. *Journal of Experimental Botany* **69**:
1415–1432. doi: https://doi.org/ 10.1093/jxb/ery010

Müntz K. 2007. Protein dynamics and proteolysis in plant vacuoles. *Journal of Experimental*
Botany **58**: 2391–2407. doi: https://doi.org/10.1093/jxb/erm089

Naested H, Frandsen GI, Jauh GY, Hernandez-Pinzon I, Nielsen HB, Murphy DJ, Rogers JC,
Mundy J. 2000. Caleosins: Ca2+-binding proteins associated with lipid bodies. *Plant Molecular*
Biology **44**: 463–476. doi: https://doi.org/ 10.1023/a:1026564411918

Nakabayashi K, Okamoto M, Koshiba T, Kamiya Y, Nambara E. 2005. Genome-wide profiling
of stored mRNA in *Arabidopsis thaliana* seed germination: Epigenetic and genetic regulation of
transcription in seed. *The Plant Journal* **41**: 697–709. doi: https://doi.org/ 10.1111/j.1365-
313X.2005.02337.x

Nakamura S, Hagihara S, Otomo K, Ishida H, Hidema J, Nemoto T, Izumi M. 2021.
Autophagy contributes to the quality control of leaf mitochondria. *Plant Cell and Physiology* 62:
229-247. doi: 10.1093/pcp/pcaa162

Nambara E, Suzuki M, Abrams S, McCarty DR, Kamiya Y, McCourt P. 2002. A screen for
genes that function in abscisic acid signaling in *Arabidopsis thaliana. Genetics* **1613**: 1247–1255.
doi: https://doi.org/10.1093/genetics/161.3.1247

Nolte H, MacVicar TD, Tellkamp F, Krüger M. 2018. Instant Clue: a software suite for
interactive data visualization and analysis. *Scientific Reports* **8**: 12648. doi:
https://doi.org/10.1038/s41598-018-31154-6

Ohlsson JA, Leong JX, Elander PH, Ballhaus F, Holla S, Dauphinee AN, Johansson J,
Lommel M, Hofmann G, Betnér S, Sandgren M, Schumacher K, Bozhkov PV, Minina EA.
2024. SPIRO - the automated Petri plate imaging platform designed by biologists, for biologists.
The Plant Journal **118**: 584–600. doi: https://doi.org/10.1111/tpj.16587

Okamoto T, Shimada T, Hara-Nishimura I, Nishimura M, Minamikawa T. 2003. C-terminal
KDEL sequence of a KDEL-tailed cysteine proteinase sulfhydryl-endopeptidase is involved in
formation of KDEL vesicle and in efficient vacuolar transport of sulfhydryl-endopeptidase. *Plant*
Physiology **132**: 1892–1900. doi: https://doi.org/10.1104/pp.103.021147

O'Malley RC, Huang SC, Song L, Lewsey MG, Bartlett A, Nery JR, Galli M, Gallavotti A,
Ecker JR. 2016. Cistrome and Epicistrome Features Shape the Regulatory DNA Landscape. *Cell*
1655, 1280–1292. Erratum in: *Cell* **1666**, 1598. doi: https://doi.org/10.1016/j.cell.2016.04.038

Oñate-Sánchez L, Vicente-Carbajosa J. 2008. DNA-free RNA isolation protocols for
Arabidopsis thaliana, including seeds and siliques. *BMC Research Notes* **1**: 93. doi: https://doi.org/
10.1186/1756-0500-1-93

Pfaffl MW. 2001. A new mathematical model for relative quantification in quantitative RT-PCR.
Nucleic Acids Research **29**: e45. doi: https://doi.org/10.1111/j.1365-313X.2006.02845.x

Poxleitner M, Rogers SW, Lacey Samuels A, Browse J, Rogers JC. 2006. A role for caleosin in
degradation of oil-body storage lipid during seed germination. *The Plant Journal* **47**: 917–933. doi:
https://doi.org/10.1111/j.1365-313X.2006.02845.x

Qi H, Xia FN, Xiao S. 2021. Autophagy in plants: Physiological roles and post-translational
regulation. *Journal of Integrative Plant Biology* **63**: 161–179. doi:
https://doi.org/10.1111/jipb.12941

Qin Z, Wang T, Zhao Y, Ma C, Shao, Q. 2023. Molecular machinery of lipid droplet degradation
and turnover in plants. *International Journal of Molecular Sciences* **24**: 16039. doi: https://doi.org/
10.3390/ijms242216039

Reeves WM, Lynch TJ, Mobin R, Finkelstein RR. 2011. Direct targets of the transcription
factors ABA-InsensitiveABI4 and ABI5 reveal synergistic action by ABI4 and several bZIP ABA
response factors. *Plant Molecular Biology* **75**: 347–363. doi: https://doi.org/10.1007/s11103-011-
9733-9

Sajeev N, Koornneef M, Bentsink L. 2024. A commitment for life: Decades of unraveling the
molecular mechanisms behind seed dormancy and germination. *The Plant Cell* **12**: koad328. doi:
https://doi.org/ 10.1093/plcell/koad328

Sánchez-Vicente I, Albertos P, Sanz C, Wybouw B, De Rybel B, Begara-Morales JC, Chaki
M, Mata-Pérez C, Barroso JB, Lorenzo O. 2024. Reversible S-nitrosylation of bZIP67 by
peroxiredoxin IIE activity and nitro-fatty acids determines the plant lipid profile. *Cell Reports* **43**:
114091. doi: https://doi.org/10.1016/j.celrep.2024.114091

Sawa-Makarska J, Baumann V, Coudevylle N, von Bülow S, Nogellova V, Abert C,
Schuschnig M, Graef M, Hummer G, Martens S. 2020. Reconstitution of autophagosome
nucleation defines Atg9 vesicles as seeds for membrane formation. *Science* **369**: eaaz7714. doi:
https://doi.org/ 10.1126/science.aaz7714

Sera Y, Hanamata S, Sakamoto S, Ono S, Kaneko K, Mitsui Y, Koyano T, Fujita N, Sasou A,
Masumura T, Saji H, Nonomura KI, Mitsuda M, Mitsui T, Kurusu T, Kuchitsu K. 2019.
Essential roles of autophagy in metabolic regulation in endosperm development during rice seed
maturation. *Scientific Reports* **9**: 18544. doi: https://doi.org/10.1038/s41598-019-54361-1

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S,
Rueden C, Saalfeld S, Schmid B, Tinevez JY, White DJ, Hartenstein V, Eliceiri K, Tomancak
P, Cardona A. 2012. Fiji: an open-source platform for biological-image analysis. *Nature Methods*
9: 676-82. doi: 10.1038/nmeth.2019

Shao Q, Liu X, Su T, Ma C, Wang P. 2019. New insights into the role of seed oil body proteins in
metabolism and plant development. *Frontiers in Plant Science* **10**: 1568. doi: https://doi.org/
10.3389/fpls.2019.01568

Siénko K, Poormassalehgoo A, Yamada K, Goto-Yamada S. 2020. Microautophagy in plants:
consideration of its molecular mechanism. *Cells* **9**: 887. doi: https://doi.org/10.3390/cells9040887

Siloto RM, Findlay K, Lopez-Villalobos A, Yeung EC, Nykiforuk CL, Moloney MM. 2006.
The accumulation of oleosins determines the size of seed oilbodies in Arabidopsis. *The Plant Cell*
18: 1961–1974. doi: https://doi.org/ 10.1105/tpc.106.041269

Stephani M, Picchianti L, Gajic A, Beveridge R, Skarwan E, Sanchez de Medina Hernandez
V, Mohseni A, Clavel M, Zeng Y, Naumann C, Matuszkiewicz M, Turco E, Loefke C, Li B,
Dürnberger G, Schutzbier M, Chen HT, Abdrakhmanov A, Savova A, Chia KS, Djamei A,
Schaffner I, Abel S, Jiang L, Mechtler K, Ikeda F, Martens S, Clausen T, Dagdas Y. 2020. A
cross-kingdom conserved ER-phagy receptor maintains endoplasmic reticulum homeostasis during
stress. *eLife* **9**, e58396. doi: https://doi.org/10.7554/eLife.58396

Su T, Li X, Yang M, Shao Q, Zhao Y, Ma C, Wang P. 2020. Autophagy: An intracellular
degradation pathway regulating plant survival and stress response. *Frontiers in Plant Science* **11**:
164. doi: https://doi.org/10.3389/fpls.2020.00164

Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, Paulovich A,
Pomeroy SL, Golub TR, Lander ES, Mesirov JP. 2005. Gene set enrichment analysis: a
knowledge-based approach for interpreting genome-wide expression profiles. *Proceedings of the*

National Academy of Sciences of the United States of America **102**: 15545–15550. doi: https://doi.org/ 10.1073/pnas.0506580102

Tarnowski L, Rodriguez MC, Brzywczy J, Piecho-Kabacik M, Krcˇkova Z, Martinec J, Wawrzynska A, Sirko A. 2020. A selective autophagy cargo receptor NBR1 modulates abscisic acid signalling in *Arabidopsis thaliana*. *Scientific Reports* **10**: 7778. doi: https://doi.org/ 10.1038/s41598-020-64765-z

Thompson AR, Doelling JH, Suttangkakul A, Vierstra RD. 2005. Autophagic nutrient recycling in Arabidopsis directed by the ATG8 and ATG12 conjugation pathways. *Plant Physiology* **138**: 2097–2110. doi: https://doi.org/10.1104/pp.105.060673

Tian R, Wang F, Zheng Q, Niza VMAGE, Downie AB, Perry SE. 2020. Direct and indirect targets of the arabidopsis seed transcription factor ABSCISIC ACID INSENSITIVE3. *The Plant Journal* **103**: 1679–1694. doi: https://doi.org/10.1111/tpj.14854

Vitale A, Pedrazzini E. 2022. StresSeed: The unfolded protein response during seed development. *Frontiers in Plant Science* **13**: 869008. doi: https://doi.org/10.3389/fpls.2022.869008

Wesch N, Kirkin V, Rogov VV. 2020. Atg8-family proteins-structural features and molecular interactions in autophagy and beyond. *Cells* **9**: 2008. doi: https://doi.org/10.3390/cells9092008

Xie Z, Klionsky D J. 2007. Autophagosome formation: core machinery and adaptations. *Nature Cell Biology* **9**: 1102–1109. doi: https://doi.org/10.1038/ncb1007-1102

Yan H, Zhuang M, Xu X, Li S, Yang M, Li N, Du X, Hu K, Peng X, Huang W, Wu H, Tse YC, Zhao L, Wang H. 2023. Autophagy and its mediated mitochondrial quality control maintain pollen tube growth and male fertility in Arabidopsis. *Autophagy* **19**: 768–783. doi: https://doi.org/ 10.1080/15548627.2022.2095838

Yang M, Wang L, Chen C, Guo X, Lin C, Huang W, Chen L. 2021. Genome-wide analysis of autophagy-related genes in *Medicago truncatula* highlights their roles in seed development and response to drought stress. *Scientific Reports* **11**: 22933. doi: https://doi.org/10.1038/s41598-021-02239-6

Yang X, Srivastava R, Howell SH, Bassham DC. 2016. Activation of autophagy by unfolded proteins during endoplasmic reticulum stress. *The Plant Journal* **85**: 83–95. doi: https://doi.org/ 10.1111/tpj.13091

Yang Z, Klionsky DJ. 2010. Mammalian autophagy: core molecular machinery and signaling regulation. *Current Opinion in Plant Biology* **22**: 124–131. doi: https://doi.org/ 10.1016/j.ceb.2009.11.014

Yilmaz A, Mejia-Guerra MK, Kurz K, Liang X, Welch L, Grotewold E. 2011. AGRIS:
Arabidopsis Gene Regulatory Information Server, an update. *Nucleic Acids Research* **39**: D1118–
D1122. doi: https://doi.org/10.1093/nar/gkq1120

Yoshimoto K, Jikumaru Y, Kamiya Y, Kusano M, Consonni C, Panstruga R, Ohsumi Y,
Shirasu K. 2009. Autophagy negatively regulates cell death by controlling NPR1-dependent
salicylic acid signaling during senescence and the innate immune response in Arabidopsis. *The*
Plant Cell **21**: 2914–2927. doi: https://doi.org/10.1105/tpc.109.068635

Yu P, Hua Z. 2022. The ubiquitin-26S proteasome system and autophagy relay proteome
homeostasis regulation during silique development. *The Plant Journal* **111**: 1324–1339. doi:
https://doi.org/10.1111/tpj.15891

Yuan Y, Cao X, Zhang H, Liu C, Zhang Y, Song XL, Gai S. 2021. Genome-wide identification
and analysis of Oleosin gene family in four cotton species and its involvement in oil accumulation
and germination. *BMC Plant Biology* **21**: 569. doi: https://doi.org/10.1186/s12870-021-03358-y

Zhao Y, Dong Q, Geng Y, Ma C, Shao Q. 2023. Dynamic regulation of lipid droplet biogenesis in
plant cells and proteins involved in the process. *International Journal of Molecular Sciences* **24**:
7476. doi: https://doi.org/10.3390/ijms24087476

Zhuang X, Li B, Jiang L. 2024. Autophagosome biogenesis and organelle homeostasis in plant
cells. *The Plant Cell* **36**: 3009–3024. doi: https://doi.org/10.1093/plcell/koae099

1083

Legends to figures

Fig. 1. Temporal and spatial expression of *AtATG5* and *AtATG7* genes in adult plants. (a) Expression data of *AtATG5* and *AtATG7* were obtained from the Gene Expression Tool at the Bio-Analytic Resource for Plant Biology (BAR) public database and represented as a heatmap (scale 0-500 GCOS units). GUS histochemical staining of different organs and tissues from *PAtATG5::uidA* (b) and *PAtATG7: uidA* (c) transgenic adult plants. C: Cotyledon, CL: Cauline Leaf, DS: Dry Seed, F: Flower, R: Root, MP: Mature Pollen.

Fig. 2. Expression patterns of *AtATG5* and *AtATG7* genes during embryogenesis, seed maturation and germination. (a) Images display the developmental stages of siliques used in the qPCR assays, along with corresponding patterns of silique weight, cell expansion, cell division, and ABA accumulation during silique development. Transcript analysis of *AtATG5* and *AtATG7* during seed embryogenesis and maturation is shown in (b), while (c) illustrates their expression during imbibition. (d) GUS histochemical staining of different *PAtATG5::uidA* and *PAtATG7::uidA* embryos at different time points (hours) of imbibition (hoi). Early Green stages 1 and 2 (EG1 and EG2), Late Green stages 1 and 2 (LG1 and LG2), and Brown (B) seeds. Data is shown as means ± standard error (SE) of three independent experiments, normalized to the expression of the *AtACT-7* gene. ABA peaks are shown in Fig. 2a, as described by Finkelstein *et al.* (2002, 2004).

Fig. 3. Germination assays in *atg5, atg7, abi5-7* mutant lines, and Col-0 (control) in the presence and absence of 1 μM ABA. Means ± SE of three independent replicates are represented. SCR: Seed Coat Rupture; ER: Endosperm Rupture, *sensu stricto* germination. In the inset, the time required to achieve 50% seed coat rupture (t_{50}) and endosperm rupture (t_{50}) is indicated.

Fig. 4. Differentially expressed genes (DEG) analysis in *atg7* in comparison to Col-0 at 24 hoi. (a) The volcano plot displays DEG as red dots, meeting the criteria of *p*-adj. ≤ 10e-16 and ± ≤ |2.| Some of the most significant DEGs such as *AtCLO1, CRC, ATPDI10, AtCRT1b, SEIPIN1, and ALPHA-TIP* are highlighted in boxes. (b) GSEA categorizes all DEG into different GOs represented as curves ranked according to their Log_2FC. The categories highlighted in red are further analyzed. (c) The lollipop plot visualizes the expression of DEGs that meet the criteria in (a) and belong to the selected GO categories from (b). Additional DEGs that do not fall into the selected categories are included in the plot as well. NS, non-significant; Log_2FC, Log_2 fold change; *p*-adj, *p*-adjusted. *Indicate genes present in two categories.

1118

Fig. 5. Differences in lipid content in *A. thaliana* Col-0, *atg5*, and *atg7* embryos at 24 and 48 hoi. (a) Differential Interference Contrast (DIC) images at x10 magnification show the entire embryos, with the radicle area used for lipid visualization and lipid content analysis marked with a rectangle. Confocal images of Nile Red lipid staining of individual embryo cells are visualized at x60 magnification. (b) Lipid content analysis of individual cells represented as % lipid area/cell area in the different *A. thaliana* lines. Data shown are means ± SE of at least 10 independent cells. Letters indicate statistically significant differences among the three Arabidopsis lines ($p \leq 0.05$, One-way ANOVA followed by Turkey's comparisons test). (c) Dynamics of lipid accumulation in seed embryos. The depicted scheme shows the balance between oil body biogenesis and degradation, indicating participating enzymes with altered gene expression levels in the *atg7* mutant at 24 hoi (red: repressed; green induced), see Fig. S7 for more details.

1130

Fig. 6. Differences in protein organization in *A. thaliana* Col-0, *atg5*, and *atg7* dry (0 hoi) and 24 and 48 hoi seed sections. Periodic acid (PAS)- Naphthol Blue Black (NBB) staining of protein and polysaccharides is shown for Col-0 (a-f), *atg5* (g-l), and *atg7* (m-r) seed transversal sections. Images of DS (a, g, m), at 24 hoi (c, i, o) and at 48 hoi (e, k, q) taken at 25x magnification, showing with a white rectangle indicating the close-up image taken at 60x magnification for DS (b, h, n), at 24 hoi (d, j, p) and 48 hoi (f, l, r). C, cotyledon; PSV, Protein Storage Vacuoles; R, radicle; SG, starch grain.

1138

Fig. 7. Differences in PSV organization in *A. thaliana* Col-0, *atg5*, and *atg7* germinating seed sections. Confocal images showing AtATG8 immunolocalization in Col-0 (a-c), *atg5* (e-g), and *atg7* (i-k) seed transversal sections at 40x magnification (a, e, i), 63x magnification (b, f, j) and a close-up image from x63 magnification, indicated with a white rectangle (c, g, k). DIC images at x40 magnification are also shown (d, h, l). C, cotyledon; PSV, Protein Storage Vacuoles; R, radicle.

1144

Fig. 8. Transcriptomic analysis of the specific ABA response in *atg7* and Col-0 imbibed seeds. (a) The volcano plot displays DEG as red dots, meeting the criteria of *p*-adj < 0.05 and $Log_2FC \geq |2.0|$. Some of the most significant DEGs such as *AFP1, AtCAX1, AtCLO4, AtFER2, EM1, and OLE1,* are highlighted in boxes. (b) Gene ontology (GO) analysis using ShinyGO platform of DEG in (a), key gene categories are marked in red, focusing on ABA and abiotic stress-related categories (top plots). Transcription factor (TF) analysis using the AGRIS Plant TF database linked to ShinyGO platform for potential regulators of the DEG in (a; bottom plots). (c) The lollipop plot visualizes the expression of DEGs identified as possible ABI5 targets and belong to ABA response

1153 GO term in (b). hoi, hours of imbibition; NS, non-significant; Log$_2$FC, Log$_2$ fold change; *p*-adj, *p*-
1154 adjusted. ABI5 target genes extracted from AGRIS Plant TF database are reported in Besmihen *et*
1155 *al.* (2002) and O'Malley *et al.* (2016).

1156

Fig. 9. AtATG8H specifically binds to AtABI3 and AtABI5 proteins in a Y2H experiment and
AtABI5 degradation is delayed in *atg5* and *atg7* mutants. (a) Western blot analysis of AtABI5 in
A. thaliana Col-0, *atg5* and *atg7* dry (0 hoi) and germinating seeds (36, 48, 72 hoi). (b) The ABI5
band intensity from western blot analysis in (b) is calculated by densitometry (means of the band
found in each sample). (c) Growth of yeast diploid cells expressing the GAL4 binding domain fused
to ATG8H (BD-ATG8H) and the GAL4 activation domain fused to ABI3 or ABI5 (AD-ABI3 or
AD-ABI5) in selective medium (–L, –W, –H) with increasing concentrations of 3-AT. The lack of
growth of the negative controls BD-□ and AD-□ is also displayed. (d) Interaction of ATG8-GFP
and cMyc-ABI5. CoIP assays between *35S:cMyc-ABI5* and *ProUBQ10-GFP-ATG8A* Arabidopsis
4-day-old seedling extracts. Plants were grown in MS and the treated or not with 3-MA for 24
hours. Inputs levels were determined with anti-GFP and anti-ABI5 antibodies. Pull-down was
revealed with anti-GFP antibody and the CoIP was revealed with anti-ABI5.

1169

Fig. 10. Proposed model of the integrative role of autophagy in early seed imbibition.
According to our data, under non-stress conditions (non-ABA conditions), autophagy is crucial in
controlling lipid turnover and protein storage mobilisation during germination. On the other hand,
the ABA signal transduction pathway leads to modifications of downstream targets like ABI5,
modulating their activity and participation in responses to environmental signals. The degradation
of this kind of TF is critical for seed germination, and the contribution of a selective autophagy
mechanism could contribute to modulate ABA signaling in response to adaptation to the
environment.

1178

1179 ## Legends to supporting figures, tables and data

1180

Fig. S1. Expression profile of the genes included in Autophagy GO term (GO:0006914;
Biological Process) at different seed developmental and germination stages, represented as a
heatmap. Gene expression is retrieved from the Gene Expression Tool at the Bio-Analytic
Resource for Plant Biology (BAR) public database (https://bar.utoronto.ca/) and is expressed as the
average log$_2$FC of three replicates, in comparison to their controls by using Heatmapper tool. Gene
set includes not only ATG, but also those genes encoding VPS, autophagy receptors (e.g., *NBR1*),
TFs (e.g., *WRKY33*), and factors involved in ER stress perception (e.g., *IRE1*). Expression remains

relatively constant across diverse samples, including various silique stages (3-10), DS, and seeds
imbibed for 1, 3, 6, 12 and 24 hoi. The *TIP3;1* gene (*At1g73190*; alpha-tonoplast intrinsic protein)
stands out for its high expression across all silique stages and dry seed. Gene copy number data
taken from the Plaza tool (https://bioinformatics.psb.ugent.be/plaza) are also included. DS, dry
seed; hoi, hours of imbibition.
1193
1194 **Fig. S2. Expression profile of *AtACT-7* reference gene.** Samples are from Col-0, *atg5*, *atg7*, and
1195 *abi5-7* germinating seeds, from Col-0 siliques at different seed maturation stages, and Col-0 seeds
1196 at different time points (hours) of imbibition (hoi). Early Green seeds stages 1 and 2 (EG1 and
1197 EG2), Late Green seeds stages 1 and 2 (LG1 and LG2), and Brown seeds (). Data are presented as
1198 means ± SE of two technical replicates of three biological samples.
1199
1200 **Fig. S3. Expression of *AtATG5, AtATG7* and *AtABI5* genes in Col-0, *atg5*, *atg7* and *abi5-7* DS.**
1201 Relative expression is referred to *AtACT-7* gene expression.
1202
Fig. S4. Expression of *AtATG5* and *AtATG7* genes during embryogenesis in the absence
(control) and the presence of ABA. (a) GUS histochemical staining of *PAtATG5::uidA* and
PAtATG7::uidA embryos at 24 and 48 hoi in the absence or presence of 1 µM ABA. (b) FPKMs
extracted from quantification analysis of RNA-seq data are represented. Data are presented as
means ± SE from three independent samples. Asterisks indicate significant difference (P-value<
0.05).
1209
Fig. S5. RNAseq experiment design and quality analysis. (a) Design of the RNAseq experiment.
Col-0 and *atg7* DS, as well as imbibed seeds in the absence (24, 48 hoi) and in the presence of ABA
(96 hoi), were collected for RNAseq analysis in triplicate. The germination status of the seeds when
collected (non-germinated or a mixture of germinated + non-germinated) are drawn. (b) Heatmap
showing the expression data of the 500 most variable genes. The colored bar on the top indicates
Col-0 samples in blue and *atg7* samples in coral. Abbreviations of the samples' name: D, dry seed;
A, in presence of ABA; 24, 24 hoi; 48, 48 hoi; 96, 96 hoi; _1, replicate 1; _2, replicate 2; _3,
replicate 3. (c) Two-dimensional PCA of the RNAseq samples performed with of the 500 most
variable genes. (d) Scree plot of RNAseq samples, showing the percentage of explained variances in
each dimension.
1220

Fig. S6. Negative controls of Nile Red lipid staining were performed on *A. thaliana* Col-0, *atg5*, and *atg7* unstained embryos at 24 and 48 hoi. DIC images of the entire embryos were captured at x10 magnification and confocal images were taken at 60x magnification under the same conditions as in Fig. 6.

Fig. S7. Lipid metabolism-related genes are differentially expressed in *atg7* compared to Col-0 at 24 hoi. (a) Lollipop plot showing the expression of differentially expressed genes (DEGs) involved in lipid metabolism in the *atg7* mutant vs. Col-0 at 24 hoi (Data S3d). (b) Schematic representation of fatty acid biosynthesis, elongation, and one cycle of the β-oxidation pathways, adapted from KEGG (http://bioinformatics.sdstate.edu/go/). Enzymatic activities corresponding to the DEGs in (a) are highlighted in red. (c) Correspondence between enzymatic activities and gene names. Abbreviations: ACP, acyl carrier protein; B-ox, β-oxidation; CPT2, carnitine palmitoyltransferase 2; FAB, fatty acid biosynthesis; FAD, fatty acid desaturation; FAE, fatty acid elongation; LIP, lipases.

Fig. S8. Negative controls of AtATG8 immunolocalization performed on germinating seed sections of *A. thaliana* Col-0. Confocal images taken at 40x magnification (a, d, g), at 60x magnification (b, e, h), and DIC images at 40x magnification (c, f, i) of the control samples incubated with primary antibody but not secondary (a-c), secondary antibody but not primary (d-f) and without any antibody (g, h, i). Images were taken in the same conditions as in Fig. 7. R, radicle; C, cotyledon.

Fig. S9. AtbZIP67 and its target genes accumulation in AtATG mutant lines. (a) Western blot analysis of AtbZIP67 in *A. thaliana* siliques from Col-0, *atg5*, *atg7* mutants and the overexpression lines *oeATG5* and *oeATG7* at different temperatures (15ºC and 21ºC). (b) AIMs found in the *AtbZIP67* gene sequence. (c) Expression of AtbZIP67 target genes in the RNAseq analysis comparing the effect of the presence of ABA in *atg7* and Col-0 lines (96 hoi in the presence of ABA vs 48 hoi in the absence).

Fig. S10. ATG8 Interacting Motifs (AIMs) found in AtABI3 and AtABI5. The consensus sequence of AIMs is considered as Θ-X-X-Γ, where Θ is an aromatic (W/F/Y) and Γ is a hydrophobic (L/I/V) residue (Wesch *et al.*, 2020).

Table S1. Sequences of primers used in this work.

1256

1257   **Data S1.** List of genes found in the AmiGO server using autophagy Gene Ontology term and their

1258   expression in siliques and seeds retrieved from the Gene Expression Tool at the Bio-Analytic

1259   Resource for Plant Biology (BAR) public database (https://bar.utoronto.ca/), expressed in GeneChip

1260   Operating System (GCOS) units (scale 0-10,000).

1261

1262   **Data S2.** RNA-seq differential expression analysis results (DESeq2) from the comparison of *atg7*

1263   vs Col-0 imbibed seeds at 24 hoi. Key columns include: Gene ID, Normalized counts (average

1264   normalized counts: BaseMean), $\log_2$FoldChange (expression change between conditions), p-value

1265   (statistical significance), adjusted p-value (padj; FDR corrected).

1266

Data S3. Gene Set Enrichment Analysis (GSEA) from the comparison of *atg7* vs Col-0

imbibed seeds at 24 hoi. (a-sheet): Key columns include: Description Gene Set, NES (Normalized

Enrichment Score), p-value (statistical significance), FDR q-value (false discovery rate),

Core_enrichment (subset of genes driving the enrichment). **(b-sheet)** contains selected enriched

gene sets and their corresponding results from the differential expression analysis: Lipid storage,

seed maturation, and response to ER stress categories. Genes that belong to selected categories are

listed and those meeting the cut-offs (*p*-adj < 0.05 and $\text{Log}_2\text{FC} \geq |2.0|$) are highlighted in green.

Data represented in Fig. 4c as Lollipop plot. (**c-sheet**) shows the analysis of genes belonging to the

Seed   Maturation   category   in   B-sheet   using   the   ShinyGO   platform

(http://bioinformatics.sdstate.edu/go/) with the AGRIS Plant TF DB database, displaying the

potential TFs that could regulate the expression of these genes. A column has been added in b-sheet

marking the possible target genes of ABI5. (**d-sheet**) shows the expression analysis of genes

belonging to the Lipid metabolism.

1280

1281

Data S4. RNA-seq differential expression analysis results (DESeq2) from the comparison of

atg7 vs Col-0 imbibed seeds at 96 hoi (1 μM ABA) and 48 hoi (DEGs responding to ABA

specifically in the *atg7* mutant compared to Col-0, at comparable physiological states, see Fig.

S5). Key columns include: Gene ID, Normalized counts (average normalized counts: BaseMean),

log2FoldChange (expression change between conditions), p-value (statistical significance), adjusted

p-value (padj; FDR corrected).

1288

Data S5. Gene Ontology and Transcription Factor Analysis in *atg7* vs. Col-0 in response to

ABA upon seed germination. Treatment. a-sheet: Gene ontology (GO) analysis using ShinyGO

for genes significantly downregulated in Data S4 (p-adj $\leq$ 0.05 and Log$_2$FC< - 1.0). The most

significant gene categories are highlighted in red, particularly those related to ABA response and

abiotic stress. **b-sheet**: Gene ontology (GO) analysis using ShinyGO for genes significantly

upregulated in Data S4 (p-adj $\leq$ 0.05 and Log$_2$FC> 1.0). As in sheet A, key gene categories are

marked in red, focusing on ABA and abiotic stress-related categories. **c-sheet**: Transcription factor

(TF) analysis using the AGRIS Plant TF database linked to ShyniGO platform for potential

regulators of the downregulated genes of Sheet-A. Potential targets of ABI5 are highlighted in red.

d-sheet: Transcription factor (TF) analysis for the upregulated genes in Data S4, using the same

AGRIS database than in C-Sheet. Potential ABI5 target genes are marked in red. **e-sheet:** Gene

ontology (GO) analysis of the potential ABI5 target genes obtained in C-Sheet, specifically

focusing on genes categorized under "ABA response" (highlighted in red). F-sheet: Differentially

Gene expression analysis results for the genes highlighted in red from sheets D and E (Data taken

from Data S4). These results correspond to the Lollipop chart displayed in Fig. 8c, illustrating the

expression levels of the selected genes.

1305

1306

Figure 1

(a)

| | CL | C | R | F | MP | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | DS | 24 hoi |

AtATG5
AtATG7

silique stage

Color Key

100   300   500

GCOS units

(b) *PAtATG5::uidA* (-404 bp from the ATG; intergenic region)

Adult Plant (75 days old)

LEAF · FLOWER · STAMEN AND SILIQUE · SILIQUE · SEED AT SILIQUE

Seedlings (15 days old)

APICAL MERISTEM · COTYLEDON · LEAF · ROOT TIP · ROOT

(c) *PAtATG7::uidA* (-340 bp from the ATG; intergenic region)

Adult Plant (75 days old)

LEAF · FLOWER · STAMEN AND SILIQUE · SILIQUE · SEED AT SILIQUE

Seedlings (15 days old)

APICAL MERISTEM · COTYLEDON · LEAF · ROOT TIP · ROOT

Figure 2

Figure 3

Figure 4

atg7 vs Col-0 (24 hoi)

Figure 5

Figure 6

Figure 7

Figure 8

Figure 9

(a)

(b)

(c)

(d)

Figure 10