In Search of the Perfect Model: How Cancer Cell Lines Relate to Native Cancers

Rahel Paloots[1,2], Ziying Yang[1,2], and Michael Baudis [1,2]

[1]Department of Molecular Life Sciences, University of Zurich, Winterthurerstrasse 190, Zurich, 8057, Switzerland
[2]Swiss Institute of Bioinformatics (SIB), Winterthurerstrasse 190, Zurich, 8057, Switzerland

Cancer cell lines are frequently used in biological and translational research to study cellular mechanisms and explore treatment options. However, cancer cell lines may display mutational profiles divergent from native cancers or may be misidentified or contaminated. We explored how similar cancer cell lines are to native cancers to find the most suitable representations for the corresponding diseases by utilising large collections of copy number variation (CNV) profiles and applied machine learning (ML) algorithms to predict cell line classifications.

Our results confirm that cancer cell lines indeed accumulate more mutations compared to native cancers but retain similar CNV profiles. We demonstrate that many relevant oncogenes and tumor suppressor genes are altered by CNV events in both cancers and their corresponding cell lines. Based on the similarities between the two groups and the predictions of the ML model, we provide some recommendations about cell lines with good potential to represent selected cancer types in *in vitro* studies.

copy number variants | cancer | cancer cell lines
Correspondence: *info@progenetix.org*

Introduction

Derived from cellular samples of primary or recurring tumors, hematopoietic neoplasias, or cancer metastases, cancer cell lines serve as tractable *in vitro* representations of human malignancies, enabling researchers to dissect cancer biology and evaluate potential therapeutic interventions. They are popular models due to established handling procedures and relatively low cost. To establish a cancer cell line, cells are isolated from a native cancer - *e.g.* a solid tumor or bone marrow in case of some lymphoid neoplasms. The ability of many cancer cells to divide perpetually is exploited for propagating them indefinitely for recurrent use in studies, potentially over the course of decades. For example, the first cancer cell line ever established, HeLa (1951) (1), is still one of the most commonly used *in vitro* model systems.

Since the establishment of HeLa and besides individual cell lines established in research projects and maintained at individual institutions, thousands of cell lines have been made available through commercial services, promising precise matching of cell lines to disease types. Many fixed sets of cell lines have been created to model different cancer types in a consistent manner under experimental conditions, e.g. for comparing drug activity or effects of targeted genetic modifications. A widely used collection of cancer cell lines is the NCI-60 panel that represents 9 distinct cancer types (2).

Cancer tissues are complex and genetically heterogeneous, frequently comprising various clonal populations (3, 4). Genomic aberrations in cancers enable disease progression as well as metastatic growth. A major class of these genomic alterations are copy number variations (CNVs) where large regions in the genomes have been modified by amplification or deletion of a section. These CNV profiles are also cancer type specific, *e.g.* duplication of chromosome 13 in colorectal carcinomas (5–7). Moreover, genomic alterations in cancers depend on the grade as well as the stage of the disease (3), suggesting different subtypes can be detected within disease classification. For example, medulloblastomas have four well known and characterized subtypes. Two of these well-known subsets are based on the expression of Wnt and Shh genes, the other two groups are "group 3" and "group 4" since the biology behind these types is not clear (8).

Due to their restricted origin, cancer cell lines inherently capture only a limited subpopulation of the original tumor's genetic diversity. This inherent limitation is further compounded by the selective pressure exerted by in vitro culture conditions. *In vitro* systems also lack the crucial interactions with other cell types that help shape neoplastic growths. Therefore, under these disparate conditions, cancer cell lines obtain novel features, such as bear a larger amount of copy number alterations compared to primary cancers (9, 10).

In this study, we evaluated cancer cell lines of different types and compared their profiles to their respective native cancer types. We applied statistical and machine learning models to identify CN patterns within 31 distinct diagnostic terms. We give an overview of the genomic differences between cancers and their cell lines and unravel the key differences in the genomic features of the two groups. Based on these results, we suggest the best available cell lines per diagnostic group.

Methods

Input Data. Cancer cell line CNV data used in this study originates from cancercelllines.org - our knowledge resource for cancer cell line variants. This database currently includes about 5,600 individual CNV samples (11). Progenetix, our source for native cancer samples includes around 112,000 cancer related CNV profiles (12). Following an initial analysis of cell line samples, a subset of 31 cancer types was chosen for further investigation. This selection was guided by the distribution of representative samples within the cancer cell lines set.

To reduce the impact of samples with limited quality of their CNV profiling data, all cell line samples were assessed visu-

ally and hlproblematic samples were flagged. Native cancer samples without any detected segments were excluded from the dataset automatically. Both native cancer and cell line samples labeled as Unspecified Tissue (NCIT:C132256) were also excluded from the dataset. A set of NCIT cancer types with a sufficient number of cancer cell line samples was used in the similarity assessment and machine learning models 2.

Binning of CNV Call Values. To bring cell line and tumor data to a uniform format, CNV data was transformed into a binned matrix. All bins are of equal specified length and represent an area in the genome. Inside each bin, CNV coverages are calculated. Duplication and deletion coverages of the bins are calculated separately. All CNV gains or losses in a bin are counted and the fraction of CNVs covering the bin are calculated. All bins with CNV gain fractions are then saved into an array, followed by the bins with deletion coverage fractions. An open-source Python package (bycon) was used for bin calculations. Genomic intervals were then calculated for different bin sizes from 1 to 10Mb. A bin size of 5 Mb was selected for further evaluations to avoid biases due to small and very large segments in the genome. This resulted in 635 bins for duplications and 635 bins for deletions, total 1270 bins.
Based on genomic bins the bycon package also supports the creation of CNV frequency maps, representing the occurrence of a CNV (gain or loss) for each segment as a percent value for all selected samples (%). For example, if all lung carcinoma samples have a duplication in 8q, the frequency in this region would be 100%. CNV frequencies are calculated based on a 1 Mb bin size to capture locations of focal events. Calculated frequencies were then used for similarity assessments and visualizations.

CNV Coverage Calculation. CNV coverage of cell line and tumor samples were retrieved from progenetix and cancercelllines databases. CNV coverage fraction is the amount of the genome that is affected by structural variants (gains and losses). To assess the levels of structural variants between cell lines and tumors of the same cancer types, the average CNV coverage fractions and subsequently fold changes between the two groups were calculated. Pre-calculated CNV coverages for both groups are included in cancercelllines and progenetix databases.

Cosine Similarity. Cosine similarity is a suitable measure for bins with values between 0 and 1 due to its scale invariance, which ensures consistent comparison regardless of vector magnitude. Its angle-based approach captures directional information, making it effective for assessing relative proportions or relationships between values, particularly in sparse datasets. Additionally, its normalized output provides easy interpretation and comparison across different datasets or dimensions. Pairwise cosine similarities between samples were calculated to detect outliers and assess the similarity between instances of the same cell line. Additionally, similarities between different cell lines as well as cell lines and native cancers were determined. Cosine similarity was calculated

by using open source python package scikit-learn (Version: 1.2.2). To assess the overall similarity of cell line and native cancer profiles, cosine similarities of CNV frequencies were calculated.

$$\text{Cosine Similarity}(\mathbf{A}, \mathbf{B}) = \frac{\mathbf{A} \cdot \mathbf{B}}{\|\mathbf{A}\| \cdot \|\mathbf{B}\|}$$

Supervised Machine Learning Algorithms. We trained support vector machine (SVM) and random forest (RF) algorithms on native cancer samples to be able to predict cancer cell line diagnostic classifications based on the profiles of neoplasms. Python package scikit-learn (Version: 1.2.2) SVM classifier with rbf kernel and RF classifiers (n_estimators=200) were used for training and prediction. Data matrices with bin sizes from 1-10 Mb were tested and 5 Mb bin size selected for further use. To find the accuracy of differentiating between cancer cell line and native cancer, equal numbers of neoplasia instances to cancer cell lines were picked at random, as the number of neoplasia samples is greater. To predict cell line diagnostic labels based on tumor profiles, machine learning models were trained on selected cancer types. Cancer types were chosen based on the number of cell line samples available per NCIT classification term and child terms of the types were included (Table 2). Testing size for all models was 0.25 with random_state=10. Table **??** outlines the individual stages of various models.

Feature Importances. We utilized scikit-learn's PermutationImportance to determine the contributions of each feature to predictions. The script fits a PermutationImportance object to the dataset and computes feature importances using permutation tests. Class feature importances ("Cell Line" and "Tumor") were then calculated for both SVM and RF models. Feature importance results included training with "Cell Line"- "Tumor" of equal numbers to identify general differences between all neoplasias and cell lines. We identified features important for distinct cancer types by training the algorithms on all cell line and neoplasia samples of the same diagnosis. The feature importances of RF model were used for this analysis because of the inherent feature importance measures of the model and the ability to capture complex relationships. Identified features were then mapped to the COSMIC set of signature genes `https://cancer.sanger.ac.uk/signatures/downloads/` (last accessed 2023-08-28) to identify relevant genes in these bins.

Matching Cell Lines with Cancer Subtypes. We partitioned cancer data by using the K-means clustering algorithm. Each cancer type (primary cancer samples only) was partitioned into clusters using 2-20 number of clusters. Median sample of each cluster in the partitioned data was calculated and the median sample of each cell line as well to represent each group. Then, cosine similarity between the medians was calculated. Only similarities equal to and above 0.7 were analyzed further.

Plotting and Visualization. All frequency maps and CNV profile plots were created with progenetix/cancercelllines on-

line software tools and bycon package software. Other graphs were created with python packages plotly, seaborn and matplotlib.

Results

Cancer cell lines display limited heterogeneity. A cancer cell line derived from a single source would exhibit minimal genetic variation within its population and within the samples of the same cell line. Even though cancer cell lines are not necessarily monoclonal, they still only represent a small subset of a tumor cell population. To determine the uniformity of the samples of the same cell line, we performed pairwise similarity calculations for all cell lines with at least 3 samples. Then, we compared computed indices to CNV sample plots to ascertain the efficiency of the measure by visual assessment. Figure 1 depicts 2 analyzed cell lines: prostate small cell carcinoma cell line NCI-H660 (CVCL_1576) and lung adenosquamous carcinoma cell line NCI-H596 (CVCL_1571). NCI-H660 is an example of a largely homogenous cell line with all similarity scores above 0.7 and there is a visual congruence among the samples on the CNV plot as well. NCI-H596 on the other hand portrays 2 distinct subsets that also brought about lower scores. Both subsets exhibit high similarity within the group but diverge from each other significantly. These results suggest that our measure can accurately determine the similarity of individual samples.

To disclose the homogeneity of the cell lines of a distinct cancer type, we calculated pairwise similarities of all samples within the same cell line (at least 3 samples) and cancer type (Supplementary Fig. A1). We demonstrate a quantifiable level of homogeneity in cell line samples for the majority of cancer types. The average level of homogeneity across these samples surpasses 0.6. Lower similarity was detected for cervical carcinoma and fibrosarcoma cell lines. These results suggest that samples of the same cell line are overall relatively homogeneous. To detect any outliers, we set the similarity threshold within the same cell line to 0.6. This threshold was set based on visual assessment of single sample plots in combination with calculated similarity indices. Therefore cell lines with at least one paired score below 0.6, would have an outlier. 53% of the cell lines had at least one outlier and only 2% diverged greatly with no similarities above threshold. Overall, the data indicated limited variability between the instances of the same cell line.

Cancer cell lines have higher CNV coverage compared to tumors of the same cancer type. Studies in breast and ovarian carcinoma cell lines have established that compared to tumors, cell lines accumulate more mutations (9, 10). For a systematic assessment of how cancer cell line genomes relate to neoplasias, we compared the CNV coverage fractions of both genomes. As expected, the CNV coverage in most cancer types was higher in cancer cell lines than in neoplasias (Fig.2) with the average fold change of 2.6. The exception was fibrosarcoma where fold change was below 1 and CNV coverage in tumors was greater. In chronic myeloge-

nous leukemia (CML) with BCR-ABL a 15-fold CNV fraction change was detected. This could be due to most cell lines being in the "blast phase" of CML when they acquire a higher mutation load due to increased genomic instability (13). The link between increased mutation load and blast phase becomes evident when considering the source of CML cell lines used in research, *e.g.* a study by Drexler et al. (2000) where most analyzed CML cell lines originated from patients in blast crisis (14). Interestingly, all three highest fold change values are in different types of leukemia, highlighting the importance of genetic fusion events in many hematopoietic leukemias. Another important fusion in leukemias is PML-RARA in acute myeloid leukemia (AML) but a variety of genes have been reported to be involved in gene fusions in leukemias (15).

Native cancers are highly heterogeneous but exhibit similar CNV profiles to cancer cell lines. It has been reported that despite harboring a higher burden of CNVs compared to their tissues of origin cancer cell lines remarkably retain a similar CNV signature to native cancers (9, 10). To systematically assess the concordance between neoplasia and cell line samples, we calculated pairwise similarity scores based on CNV coverage. Our analysis of the comparisons between cell lines and neoplasias, along with the internal consistency of the neoplasia data set, revealed scores that were lower than what we had anticipated.(Fig.3). Heterogeneity was higher within neoplasia samples than in instances of different cell lines of the same cancer type (Fig. 3, Supplementary Fig. A3). Standard deviations of similarities were higher for neoplasias in all cancer types except acute myeloid leukemia.

Pairwise comparisons of samples illustrate the heterogeneity but fail to reveal recurring patterns of similarity. Therefore, we compared the CNV frequency maps of neoplasias and corresponding cell lines. Frequency maps represent the occurrence of CNVs in the dataset in percentages - indicating the presence of CNVs in the proportion of samples. We calculated a similarity index based on CNV frequency data of cell lines and neoplasias and found high similarity indices for most cancer types (Supplementary Table A2). Clearly detectable patterns emerge between cell lines and its native cancer upon visual inspection of frequency plots as well (Supplementary Fig. A2). Our results confirm that both breast carcinomas and ovarian carcinomas exhibit similar CNV patterns to their cell lines (9, 10). Even though melanoma has bee reported as a highly heterogeneous disease (16, 17), it yielded a high similarity score (Supplementary Table A2, Supplementary Fig. A2) indicating an overall good representation of genomic patterns in aggregated cell line data.

Emerging cancer CNV patterns can be used to determine the origin of some cancer cell lines. Given the high occurrence of characteristic CNV patterns in a cancer type, our goal was to employ a method to forecast the diagnostic classification of a cell line. For that we trained a support vector machine (SVM) model on the 31 cancer types (Supplementary Table A1). In the evaluation of the model's

Fig. 1. CNV sample plots and similarity heatmaps for cell lines NCI-H660 (top) and NCI-H598 (bottom), indicating the regional coverage by copy number gains (yellow) and losses (blue) for chromosomes 1-22. For each of the 2 cell lines 5 individual instances are shown to visualize similarities and differences in CNV events. Inter-sample CNV cosine similarities are displayed on the right.

Fig. 2. Comparison of CNV coverage fractions between cell lines and patient derived samples from the same diagnostic groups. Fold change calculated per cancer type: average CNV coverage of cell line samples/average CNV coverage of native cancer samples. Error bars represent standard errors.

performance we found that for some diagnostic groups the CNV based diagnostic prediction was overall successful with the best results observed for breast adenocarcinoma, glioblastoma and colorectal carcinoma (Supplementary Table A3).

Fig. 3. Similarity heatmap of comparisons between cell lines and neoplasias. Each column shows average pairwise similarity for the cancer type: A - samples of different cell lines, B - cell line samples vs neoplasia samples, C - neoplasia samples.

We then applied the trained model to predict the diagnostic classifications of our collection of cell line CNV profiles. The prediction accuracies of the cell lines were largely similar to testing - highest percentage of correctly predicted samples belonging to breast adenocarcinoma, glioblastoma and colorectal carcinoma. Interestingly, the percentage of correctly predicted cell lines is higher for breast adenocarcinoma cell lines than tumor-tumor predictions, indicating that these cell lines are good representations of the majority of the disease. On the contrary, while the testing accuracy was the highest for acute myeloid leukemia (AML) (90.51%), the prediction accuracy of AML cell lines was below 20%. In fact, the results for all leukemia types in our data sets revealed poor performance. The fold change of CNV coverage was also the

highest among leukemias (Fig. 2), suggesting poor representation of the majority of native leukemia samples.

Co-occurrence of relevant cancer genes in features important for class determination. To interpret the performance of our machine learning model, we employed a feature importance analysis upon its implementation. This analysis identified the most significant regions within the genome that contribute to the model's classification capabilities.

Our initial objective was to examine whether any genomic features could be identified as influential in classifying "neoplasia" or "cell line". Therefore, we collected all cell line samples in our dataset and randomly picked the same number of neoplasia samples. This step ensured the same num-

Fig. 4. Frequency maps of all neoplasias and cancer cell lines. Blue - deletions, yellow - copy number gains, as percent of occurrence in analyzed samples. Locations of the top 30 genes with differences between cell lines and primary tumors in the ML model are indicated.

ber of samples to avoid introducing bias to our model. Distinguishing "neoplasia" from "cell line" with 89% accuracy (testing), this model also highlighted the most relevant features in the process (Fig. 4) which curiously include relevant cancer genes with functional involvement in many cancer types, such as TP53.

Next, we analyzed the important features in our selected 31 cancer types, by training our models with both cell line and neoplasia samples and finding the features relevant for each cancer type. Then, we matched identified features (genomic bins) with COSMIC cancer gene set, to identify any underlying oncogenes or tumor suppressor genes in the region. Of all the identified features, 56% are duplications and 44% are deletions. We then sorted the features by including only features with highest average values and that exist in at least 4 cancer types to have an overview of shared relevant features in the cancer types (Fig. 5). 42 out of 50 top shared features are duplications, indicating that duplications carry more weight in class identification. Many of these duplicated features include important genes such as BRCA1, EGFR, MAPK1. Wildtype-BRCA1 is a tumor suppressor gene and mutated BRCA1 increases risk for breast and ovarian carcinoma (18). BRCA1 is detected as an important feature for breast adenocarcinoma but not ovarian carcinoma (Fig. 5). EGFR- epidermal growth factor receptor, is an oncogene that promotes tumor progression. Notably, EGFR over-expression has been detected in lung adenocarcinomas but not in small cell lung carcinomas (SCLC) (19). Duplications in EGFR in lung adenocarcinomas and SCLCs are not marked as important features in our dataset. However, EGFR duplications are important features for determining breast adenocarcinoma sample types and breast cancers are also known to express EGFR (Fig. 5) (19). Another potent oncogene in cancers, particularly bladder carcinomas is FGFR3 (20). Interestingly, FGFR3 deletions are highlighted as important features for class detection in 5 cancer types,

including bladder carcinomas (Fig. 5).

Overall, the frequency of these features is lower in native cancers than in cell lines. This aligns with the known higher mutation rate observed in native cancers (Fig. 10, Fig. 11). While the frequencies of other features varied, the occurrence of mutations in certain genes remained consistent, e.g. COL1A1 in neuroblastomas and FHIT in lung small cell carcinomas. However, these genes were not deemed relevant for the classification algorithm.Different copy numbers in native cancers and cancer cell lines can be attributed to cancer cell lines only representing a small sub-population of the disease. The frequent use of cell lines derived from the same individual can lead to genetic drift, exacerbating these differences. Additionally, changes in copy numbers are observed in *in vitro* environments.

Our results suggest that several relevant cancer genes are an integral part for the determination of the class in our model. These genes and features are present in both neoplasias and cancer cell lines but at significantly higher levels in cell lines. These high level CN changes in cancer cell lines make them excellent models for studying the effects of these genes and testing for possible pharmaceuticals.

Modeling Tumor Heterogeneity: The Utility of Cell Lines. Cancers are heterogeneous diseases that consist of multiple subsets. For example, medulloblastoma is a form of brain cancer that includes several distinct subtypes with characteristic mutational profiles (8). Another well-known heterogeneous cancer type is melanoma (17) where a large population of subclones have been detected (21). Based on known intratumoral heterogeneity, we hypothesized that certain cell lines would more accurately represent the unique molecular characteristics of specific cancer subsets. Therefore, we partitioned our neoplasia data using K-means clustering and picking the median sample of each cluster. We then matched these cluster medians to median samples of cell

Fig. 5. Known oncogenes and tumor suppressor genes important for class determination. Deleted genes are marked with blue and amplified genes with yellow, grey indicating no change in copy numbers. Genes exhibiting a higher incidence in native cancers relative to cell lines are denoted by 'x' on the heatmap.

lines. Selecting the median sample allowed us to establish an "average" representation for each group. Indeed, we were able to match some subsets to some cell lines with high similarity, including melanomas and lung small cell carcinomas. For instance, Figure 6 displays a subset of lung small cell carcinoma tumor and cell line samples. We were able to identify three distinct cell lines that would be the best representations of this tumor subpopulation. Additionally, the subsets of melanoma, colorectal carcinoma, glioblastoma and kidney carcinoma matched with high similarity to several cell lines (Supplementary material).

Selection of representative cell lines. Our analysis, utilizing a SVM model trained on 31 distinct cancer types and complemented by visual assessment of CNV profiles, has yielded a shortlist of cell lines that serve as strong candidates for faithfully representing instances of 15 types of primary neoplasia (Table 1). Notably, these cell lines were consistently predicted across all 31 cancer types within the SVM model, suggesting a broader applicability rather than over-training towards a specific subtype match. Out of all 31 cancer types analysed, we were able to determine candid models for 15 types. The highest number of accurate models were identified for breast adenocarcinomas where prediction accuracy was also the highest. We were also able to pinpoint 3 models for AML despite overall low prediction accuracy for this hematologic neoplasia.

To further verify the quality of our selection we calculated pairwise cosine similarities between the native cancer samples and instances of each individual cell lines (Supplementary Materials). These data indicate that for some cancer types, such as prostate carcinoma, suggested cell lines indeed corresponded to predominant sample types. In general, although similarity scores between the suggested cell lines varied widely the trained model showed an ability to successfully identify promising candidates even with low overall CNV pattern similarity.

Discussion

We analyzed the molecular variability of over 500 cell lines (around 3,400 samples) using copy number variation profiles data. Overall, different instances of the same cell line displayed mostly limited variability. Due to the shared origin of these samples, this consistency in prediction is unsurprising. Higher similarity of different cancer cell lines of the same type to each other may be due to cell lines originating from a smaller clonal population and therefore being more stable. This trade-off presents a key challenge: While simplified models may offer stability and ease of analysis, they inherently struggle to capture the full spectrum of disease heterogeneity.

Studies with cancer cell lines have demonstrated that genetic drift occurs in cancer cell lines and significantly affects their pharmacogenetic properties (22, 23). Our results suggest that the heterogeneity of cancer cell lines is limited, especially when compared to native cancers. However, further analyses are required to determine the full scope of the genetic drift in the cell lines. Additionally, as apparent genetic drift in cancer cell lines can be the result of positive selection (22), some level of heterogeneity is inevitable.

The results of this study confirm that cancer cell lines indeed generally harbor a higher amount of CNVs compared to

Table 1. Cell lines with good correspondence to native cancer types

Cancer Types	Cell Lines
Acute Myeloid Leukemia	KO52 (CVCL_1321), KG-1 (CVCL_0374), MV4-11 (CVCL_0064)
Breast Adenocarcinoma	EFM-19 (CVCL_0253), ZR-75-1 (CVCL_0588), MDA-MB-134-VI (CVCL_0617), MDA-MB-361 (CVCL_0620), CAL-148 (CVCL_1106), CAL-85-1 (CVCL_1114), CAMA-1 (CVCL_1115), COLO 824 (CVCL_1136), DU4475 (CVCL_1183), Evsa-T (CVCL_1207), HCC1008 (CVCL_1244), HCC1143 (CVCL_1245), HCC1428 (CVCL_1252), HCC1500 (CVCL_1254), HCC1599 (CVCL_1256), HCC2218 (CVCL_1263), (CVCL_1267), HCC38 (CVCL_1270), MDA-MB-175-VII (CVCL_1400), MFM-223 (CVCL_1408), ZR-75-30 (CVCL_1661), UACC-812 (CVCL_1781), UACC-893 (CVCL_1782), EFM-192A (CVCL_1812), HDQ-P1 (CVCL_2067), JIMT-1 (CVCL_2077), KPL-1 (CVCL_2094), BT-483 (CVCL_2319), VP229 (CVCL_2754), VP267 (CVCL_2755), OCUB-F (CVCL_3352), HCC712 (CVCL_3378), SUM190PT (CVCL_3423), SUM44PE (CVCL_3424), MDA-MB-468GFP (CVCL_DH83)
Colorectal Carcinoma	COLO 320DM (CVCL_0219), SW403 (CVCL_0545), SW948 (CVCL_0632), LS1034 (CVCL_1382), NCI-H508 (CVCL_1564), SW1463 (CVCL_1718), SW626 (CVCL_1725), COLO 201 (CVCL_1987), COLO 206F (CVCL_1988), WiDr (CVCL_2760)
Diffuse Large B-Cell Lymphoma	HT (CVCL_1290), OCI-Ly10 (CVCL_8795), OCI-Ly19 (CVCL_1878), Ri-1 (CVCL_1885), SU-DHL-4 (CVCL_0539)), SU-DHL-5 (CVCL_1735)
Esophageal Carcinoma	KYSE-140 (CVCL_1347), KYSE-150 (CVCL_1348), KYSE-270 (CVCL_1350), KYSE-30 (CVCL_1351), KYSE-510 (CVCL_1354), OE21 (CVCL_2661), OE33 (CVCL_0471)
Glioblastoma	DK-MG (CVCL_1173), GaMG (CVCL_1226), SF295 (CVCL_1690), SNB-75 (CVCL_1706), YKG-1 (CVCL_1796)
Head and Neck Squamous Cell Carcinoma	SCC-25 (CVCL_1682), BICR 22 (CVCL_2310), CAL-27 (CVCL_1107)
Kidney Carcinoma	769-P (CVCL_1050), A-498 (CVCL_1056), A-704 (CVCL_1065), ACHN (CVCL_1067), CAL-54 (CVCL_1111), VMRC-RCW (CVCL_1790), VMRC-RCZ (CVCL_1791), UM-RC-6 (CVCL_2741), KMRC-1 (CVCL_2983), KMRC-2 (CVCL_2984), SLR26 (CVCL_V612)
Lung Adenocarcinoma	HCC1171 (CVCL_5126), HCC2450 (CVCL_5133), NCI-H1437 (CVCL_1472), NCI-H1693 (CVCL_1488), NCI-H1993 (CVCL_1512), NCI-H2030 (CVCL_1517), NCI-H2087 (CVCL_1524), NCI-H2291 (CVCL_1546), NCI-H650 (CVCL_1575), NCI-H820 (CVCL_1592), NCI-H920 (CVCL_1599), RERF-LC-KJ (CVCL_1654)
Lung Small Cell Carcinoma	COR-L279 (CVCL_1140)
Lung Squamous Cell Carcinoma	VMRC-LCP (CVCL_1788), LC-1/sq (CVCL_3008)
Melanoma	C32 (CVCL_1097), C32TG (CVCL_2324), COLO 741 (CVCL_1133), COLO 829 (CVCL_1137), G-361 (CVCL_1220), Hs 936.T (CVCL_1033), HT-144 (CVCL_0318), IGR-37 (CVCL_2075), Ma-Mel-36 (CVCL_A171), Malme-3M (CVCL_1438), SK-MEL-1 (CVCL_0068), SK-MEL-103 (CVCL_6069), SK-MEL-19 (CVCL_6025), SK-MEL-199 (CVCL_6104), SK-MEL-29 (CVCL_6031), SK-MEL-30 (CVCL_0039), UACC-257 (CVCL_1779), WM1193C (CVCL_C265), WM164 (CVCL_7928), WM1799 (CVCL_A341), WM266-4 (CVCL_2765), WM3060 (CVCL_6796), WM3066 (CVCL_C270), WM3208V (CVCL_L028), WM3248 (CVCL_6798), WM51 (CVCL_6995), WM852 (CVCL_6804), WM902B (CVCL_6807)
Ovarian Carcinoma	59M (CVCL_2291), COV318 (CVCL_2419), FU-OV-1 (CVCL_2047), OVCAR-4 (CVCL_1627), PEO1 (CVCL_2686), PEO4 (CVCL_2690), PEO6 (CVCL_2691)
Pancreatobiliary Carcinoma	SU.86.86 (CVCL_3881), Capan-1 (CVCL_0237)
Prostate Carcinoma	NCI-H660 (CVCL_1576), VCaP (CVCL_2235)

Fig. 6. Clustered sample plots of similar cell line and neoplasia subset samples. Here, a subset of small cell lung carcinoma (SCLC) samples is interspersed by instances of SCLC cell in the unsuper- vised clustering analysis, indicating the high degree of CNV pattern similarity between some SCLC samples and those in vitro models. Cell line samples are indicated in pink and neoplasia samples in black.

native cancers but also generally display CNVs shared with native cancers of the same diagnosis (Table 3, Fig. 8 and (9, 10)). The highest differences in CNV coverage were detected for leukemias, in particular for CML which can be attributed to the ease of using blast phase CML for generation of immortal cell lines (13) with inherent differences in genomic stability. Furthermore, this demonstrates that not all stages of cancers are well represented in *in vitro* models.

We examined CNV samples of 31 different cancer types and showed that neoplasias exhibit high molecular variability (Fig. 3, Fig. 9). A useful strategy in taking advantage of cell lines would be to partition primary cancer into subsets and match these subsets to cell lines by similarity, a strategy we successfully employed to match cell lines to a subset of lung small cell carcinoma samples (Fig. 6). Overall, the current model may not fully account for the significant variations associated with within different cancer subtypes and potentially could be improved by a general approach including partitioning of cancer types and identification of cell lines that closely resemble those rather than unselected, potentially heterogeneous diagnostic classes.

We demonstrated that the prediction of cell line's disease classification based on CNV patterns of native cancers was highly accurate for breast adenocarcinomas, colorectal carcinomas and glioblastomas. Using ML algorithms to classify cancers and cell lines also informs us about genomic features important for classifications. We showed that several well-known oncogenes and tumor suppressor genes might have influenced the decision-making process (Fig. 5). We also demonstrate that duplications are more important for class determination than deletions.

One effective approach to improving the performance of clas-

sification models on high-dimensional features like data from genomic screening experiments is to augment the sample size. For our particular analysis, addition of samples could lead to additional cell line matches while for underrepresented cancer types as well as a general approach consistent partitioning of cancer samples for subtype specific matching. Importantly our analyses hint at biases in cancer type representation through cell lines, at least such with accessible CNV profiling data.

In summary, despite the undeniable value of cancer cell lines in elucidating tumor biology and propelling advancements in precision medicine, the inherent genomic heterogeneity observed in cancer samples and across individuals needs to be accounted for. We provide a careful selection of the models corresponding best to the target disease to best capture the genomic intricacies of cancers. Data to create neoplasia subsets and match them to appropriate cell lines are available through the *Progenetix* and *cancercelllines.org* resources. Future steps in utilizing these resources could involve the creation of software tools to enable dynamic comparisons of cancer cell lines to native cancers.

Bibliography

1. WF Scherer, JT Syverton, and GO Gey. Studies on the propagation in vitro of poliomyelitis viruses. iv. viral multiplication in a stable strain of human malignant epithelial cells (strain hela) derived from an epidermoid carcinoma of the cervix. *J Exp Med*, 97(5):695–710, 1953.
2. Robert H Shoemaker. The nci60 human tumour cell line anticancer drug screen. *Nature Reviews Cancer*, 6(10):813–823, 2006.
3. Timothy N Clinton, Ziyu Chen, Hannah Wise, Andrew T Lenis, Shweta Chavan, Mark TA Donoghue, Nima Almassi, Carissa E Chu, Shawn Dason, Pavitra Rao, et al. Genomic heterogeneity as a barrier to precision oncology in urothelial cancer. *Cell reports*, 41(12), 2022.
4. Roland F Schwarz, Charlotte KY Ng, Susanna L Cooke, Scott Newman, Jillian Temple, Anna M Piskorz, Davina Gale, Karen Sayal, Muhammed Murtaza, Peter J Baldwin, et al.

Spatial and temporal heterogeneity in high-grade serous ovarian cancer: a phylogenetic analysis. *PLoS medicine*, 12(2):e1001789, 2015.

5. Kaja CG Berg, Peter W Eide, Ina A Eilertsen, Bjarne Johannessen, Jarle Bruun, Stine A Danielsen, Merete Bjørnslett, Leonardo A Meza-Zepeda, Mette Eknæs, Guro E Lind, et al. Multi-omics of 34 colorectal cancer cell lines-a resource for biomedical studies. *Molecular cancer*, 16:1–16, 2017.

6. Jordi Camps, Marian Grade, Quang Tri Nguyen, Patrick Hörmann, Sandra Becker, Amanda B Hummon, Virginia Rodriguez, Settara Chandrasekharappa, Yidong Chen, Michael J Difilippantonio, et al. Chromosomal breakpoints in primary colon cancer cluster at sites of structural variants in the genome. *Cancer research*, 68(5):1284–1295, 2008.

7. Eleanor J Douglas, Heike Fiegler, Andrew Rowan, Sarah Halford, David C Bicknell, Walter Bodmer, Ian PM Tomlinson, and Nigel P Carter. Array comparative genomic hybridization analysis of colorectal cancer cell lines and primary carcinomas. *Cancer research*, 64(14): 4817–4825, 2004.

8. Michael D Taylor, Paul A Northcott, Andrey Korshunov, Marc Remke, Yoon-Jae Cho, Steven C Clifford, Charles G Eberhart, D Williams Parsons, Stefan Rutkowski, Amar Gajjar, et al. Molecular subgroups of medulloblastoma: the current consensus. *Acta neuropathologica*, 123:465–472, 2012.

9. Richard M Neve, Koei Chin, Jane Fridlyand, Jennifer Yeh, Frederick L Baehner, Tea Fevr, Laura Clark, Nora Bayani, Jean-Philippe Coppe, Frances Tong, et al. A collection of breast cancer cell lines for the study of functionally distinct cancer subtypes. *Cancer cell*, 10(6): 515–527, 2006.

10. Silvia Domcke, Rileen Sinha, Douglas A Levine, Chris Sander, and Nikolaus Schultz. Evaluating cell lines as tumour models by comparison of genomic profiles. *Nature communications*, 4(1):2126, 2013.

11. Rahel Paloots and Michael Baudis. cancercelllines. org—a novel resource for genomic variants in cancer cell lines. *Database*, 2024:baae030, 2024.

12. Qingyao Huang, Paula Carrío-Cordo, Bo Gao, Rahel Paloots, and Michael Baudis. The progenetix oncogenomic resource in 2021. *Database*, 2021, 2021.

13. Jayastu Senapati, Elias Jabbour, Hagop Kantarjian, and Nicholas J Short. Pathogenesis and management of accelerated and blast phases of chronic myeloid leukemia. *Leukemia*, 37(1):5–17, 2023.

14. Hans G Drexler, Yoshinobu Matsuo, and Roderick AF MacLeod. Continuous hematopoietic cell lines as model systems for leukemia–lymphoma research. *Leukemia research*, 24(11): 881–911, 2000.

15. Abdul Mannan, Ibrahim N Muhsen, Eva Barragán, Miguel A Sanz, Mohamad Mohty, Shahrukh K Hashmi, and Mahmoud Aljurf. Genotypic and phenotypic characteristics of acute promyelocytic leukemia translocation variants. *Hematology/Oncology and Stem Cell Therapy*, 13(4):189–201, 2020.

16. Noemi Andor, Trevor A Graham, Marnix Jansen, Li C Xia, C Athena Aktipis, Claudia Petritsch, Hanlee P Ji, and Carlo C Maley. Pan-cancer analysis of the extent and consequences of intratumor heterogeneity. *Nature medicine*, 22(1):105–113, 2016.

17. Tomasz M Grzywa, Wiktor Paskal, and Paweł K Włodarski. Intratumor and intertumor heterogeneity in melanoma. *Translational oncology*, 10(6):956–975, 2017.

18. Amy X Zhong, Yumay Chen, and Phang-Lang Chen. Brca1 the versatile defender: Molecular to environmental perspectives. *International Journal of Molecular Sciences*, 24(18): 14276, 2023.

19. Elena Levantini, Giorgia Maroni, Marzia Del Re, and Daniel G Tenen. Egfr signaling pathway as therapeutic target in human cancers. In *Seminars in Cancer Biology*, volume 85, pages 253–275. Elsevier, 2022.

20. Claudia Maria Ascione, Fabiana Napolitano, Daniela Esposito, Alberto Servetto, Stefania Belli, Antonio Santaniello, Sarah Scagliarini, Felice Crocetto, Roberto Bianco, and Luigi Formisano. Role of fgfr3 in bladder cancer: Treatment landscape and future challenges. *Cancer Treatment Reviews*, page 102530, 2023.

21. Li Ding, Minjung Kim, Krishna L Kanchi, Nathan D Dees, Charles Lu, Malachi Griffith, David Fenstermacher, Hyeran Sung, Christopher A Miller, Brian Goetz, et al. Clonal architectures and driver mutations in metastatic melanomas. *PloS one*, 9(11):e111153, 2014.

22. Uri Ben-David, Benjamin Siranosian, Gavin Ha, Helen Tang, Yaara Oren, Kunihiko Hinohara, Craig A Strathdee, Joshua Dempster, Nicholas J Lyons, Robert Burns, et al. Genetic and transcriptional evolution alters cancer cell line drug response. *Nature*, 560(7718):325–330, 2018.

23. Rene Quevedo, Petr Smirnov, Denis Tkachuk, Chantal Ho, Nehme El-Hachem, Zhaleh Safikhani, Trevor J Pugh, and Benjamin Haibe-Kains. Assessment of genetic drift in large pharmacogenomic studies. *Cell systems*, 11(4):393–401, 2020.

Supplementary Note 1: Supplementary materials

Table 2. Cancer types used in our analysis

Code	Label	CL	T
NCIT:C5214	Breast Adenocarcinoma	829	10756
NCIT:C3224	Melanoma	689	1999
NCIT:C3512	Lung Adenocarcinoma	472	4172
NCIT:C2955	Colorectal Carcinoma	293	4787
NCIT:C4908	Ovarian Carcinoma	280	2851
NCIT:C4917	Lung Small Cell Carcinoma	212	318
NCIT:C8851	Diffuse Large B-Cell Lymphoma	141	2797
NCIT:C4911	Gastric Carcinoma	141	2278
NCIT:C9384	Kidney Carcinoma	133	2569
NCIT:C3058	Glioblastoma	115	3923
NCIT:C34447	Head and Neck Squamous Cell Carcinoma	112	1848
NCIT:C166418	Pancreatobiliary Carcinoma	107	1786
NCIT:C3270	Neuroblastoma	107	1761
NCIT:C3171	Acute Myeloid Leukemia	93	3805
NCIT:C3513	Esophageal Carcinoma	87	1738
NCIT:C3493	Lung Squamous Cell Carcinoma	80	1802
NCIT:C4912	Bladder Carcinoma	71	1689
NCIT:C4863	Prostate Carcinoma	67	3890
NCIT:C4450	Lung Large Cell Carcinoma	65	92
NCIT:C7376	Pleural Malignant Mesothelioma	56	211
NCIT:C9039	Cervical Carcinoma	54	905
NCIT:C2912	Burkitt Lymphoma	51	225
NCIT:C3043	Fibrosarcoma	50	138
NCIT:C4665	Plasma Cell Neoplasm	50	1617
NCIT:C7558	Endometrial Carcinoma	48	819
NCIT:C9145	Osteosarcoma	47	309
NCIT:C4817	Ewing Sarcoma	43	182
NCIT:C3183	T Acute Lymphoblastic Leukemia	40	238
NCIT:C3174	Chronic Myelogenous Leukemia, BCR-ABL1+	36	253
NCIT:C8644	B Acute Lymphoblastic Leukemia	32	275
NCIT:C9306	Soft Tissue Sarcoma	28	1196

Table 3. Cosine similarities of CNV frequencies between Tumor and Cell Line

Cancer Type	Cosine Similarity
Lung Small Cell Carcinoma	0.95
Esophageal Carcinoma	0.94
Gastric Carcinoma	0.94
Lung Adenocarcinoma	0.94
Melanoma	0.93
Head and Neck Squamous Cell Carcinoma	0.93
Ovarian Carcinoma	0.92
Pancreatobiliary Carcinoma	0.92
Breast Adenocarcinoma	0.92
Colorectal Carcinoma	0.90
Kidney Carcinoma	0.89
Lung Squamous Cell Carcinoma	0.89
Pleural Malignant Mesothelioma	0.88
Glioblastoma	0.87
Cervical Carcinoma	0.86
Lung Large Cell Carcinoma	0.86
Bladder Carcinoma	0.85
Prostate Carcinoma	0.84
Endometrial Carcinoma	0.83
Diffuse Large B-Cell Lymphoma	0.83
Osteosarcoma	0.82
Plasma Cell Neoplasm	0.81
Burkitt Lymphoma	0.80
Ewing Sarcoma	0.80
Neuroblastoma	0.80
Acute Myeloid Leukemia	0.78
Soft Tissue Sarcoma	0.73
T Acute Lymphoblastic Leukemia	0.70
Chronic Myelogenous Leukemia, BCR-ABL1+	0.58
Fibrosarcoma	0.55
B Acute Lymphoblastic Leukemia	0.39

Table 4. Accuracies for the SVM model. Correct test % shows the percentage of correct predictions in the testing phase. Correct % shows the percentage of correct predictions of cell line diagnoses.

Actual	Label	Correct Test %	Correct %
NCIT:C5214	Breast Adenocarcinoma	78.89	84.56
NCIT:C3058	Glioblastoma	78.32	63.48
NCIT:C2955	Colorectal Carcinoma	71.15	62.80
NCIT:C3513	Esophageal Carcinoma	32.83	56.32
NCIT:C9384	Kidney Carcinoma	72.22	54.89
NCIT:C3224	Melanoma	60.11	53.70
NCIT:C3270	Neuroblastoma	77.18	46.73
NCIT:C8851	Diffuse Large B-Cell Lymphoma	55.28	46.10
NCIT:C4665	Plasma Cell Neoplasm	68.00	44.00
NCIT:C4863	Prostate Carcinoma	64.36	41.79
NCIT:C3512	Lung Adenocarcinoma	62.00	37.08
NCIT:C34447	Head and Neck Squamous Cell Carcinoma	37.75	36.61
NCIT:C4908	Ovarian Carcinoma	56.56	35.36
NCIT:C166418	Pancreatobiliary Carcinoma	34.38	28.04
NCIT:C3171	Acute Myeloid Leukemia	90.51	19.35
NCIT:C9039	Cervical Carcinoma	33.02	16.67
NCIT:C4917	Lung Small Cell Carcinoma	40.51	15.09
NCIT:C3493	Lung Squamous Cell Carcinoma	56.37	10.00
NCIT:C4911	Gastric Carcinoma	30.90	7.09
NCIT:C4912	Bladder Carcinoma	42.30	7.04
NCIT:C7376	Pleural Malignant Mesothelioma	9.62	5.36
NCIT:C9145	Osteosarcoma	17.65	4.26
NCIT:C7558	Endometrial Carcinoma	18.72	4.17
NCIT:C8644	B Acute Lymphoblastic Leukemia	40.58	-
NCIT:C9306	Soft Tissue Sarcoma	36.33	-
NCIT:C4817	Ewing Sarcoma	13.46	-
NCIT:C2912	Burkitt Lymphoma	4.17	-
NCIT:C3043	Fibrosarcoma	2.86	-

Table 5. Accuracies for RF model. Correct test % shows the percentage of correct predictions in the testing phase. Correct % shows the percentage of correct predictions of cell line diagnoses.

Actual	Label	Correct Test %	Correct %
NCIT:C5214	Breast Adenocarcinoma	87.26	88.78
NCIT:C2955	Colorectal Carcinoma	72.83	70.31
NCIT:C3058	Glioblastoma	79.34	59.13
NCIT:C8851	Diffuse Large B-Cell Lymphoma	60.29	48.23
NCIT:C34447	Head and Neck Squamous Cell Carcinoma	37.40	41.07
NCIT:C9384	Kidney Carcinoma	69.22	40.60
NCIT:C3270	Neuroblastoma	79.92	39.25
NCIT:C3224	Melanoma	41.73	33.38
NCIT:C4908	Ovarian Carcinoma	48.65	32.86
NCIT:C4863	Prostate Carcinoma	67.26	28.36
NCIT:C3512	Lung Adenocarcinoma	62.91	27.33
NCIT:C4665	Plasma Cell Neoplasm	71.46	22.00
NCIT:C3513	Esophageal Carcinoma	27.08	21.84
NCIT:C3183	T Acute Lymphoblastic Leukemia	42.25	20.00
NCIT:C3493	Lung Squamous Cell Carcinoma	51.33	15.00
NCIT:C9145	Osteosarcoma	6.06	14.89
NCIT:C3171	Acute Myeloid Leukemia	84.51	13.98
NCIT:C166418	Pancreatobiliary Carcinoma	34.21	13.08
NCIT:C3174	Chronic Myelogenous Leukemia, BCR-ABL1+	36.99	11.11
NCIT:C4912	Bladder Carcinoma	28.16	8.45
NCIT:C4911	Gastric Carcinoma	29.09	7.09
NCIT:C9039	Cervical Carcinoma	24.70	5.56
NCIT:C4917	Lung Small Cell Carcinoma	29.90	5.19
NCIT:C7558	Endometrial Carcinoma	25.67	2.08
NCIT:C8644	B Acute Lymphoblastic Leukemia	58.75	-
NCIT:C9306	Soft Tissue Sarcoma	28.12	-
NCIT:C2912	Burkitt Lymphoma	22.39	-
NCIT:C4817	Ewing Sarcoma	13.33	-
NCIT:C7376	Pleural Malignant Mesothelioma	9.23	-
NCIT:C4450	Lung Large Cell Carcinoma	8.00	-

Fig. 7. Pairwise similaritites of the same cell line instances per cancer types (≥ 3 samples). Cell line averages are marked with a black dash. Abbreviations: AML - acute myeloid leukemia, B ALL - B acute lymphoblastic leukemia, CML - chronic myelogenous leukemia, DLBCL - diffuse large B-cell lymphoma, Plasma - plasma cell neoplasm, Pleural MM - pleural malignant mesothelioma, T ALL - T acute lymphoblastic leukemia

Fig. 8. Frequency map of primary melanoma samples and melanoma cell lines annotated with genes in identified important features

Fig. 9. Standard deviations of cosine similarities between cancer cell lines and within neoplasia samples.

Fig. 10. Percentage distribution of genes across native cancer samples of different types.

Fig. 11. Percentage distribution of genes across cell line samples of different types.

Supplementary Note 2: Cosine similarities

Violin plots of cosine similarities between native cancer and cell line samples are included in this appendix. Average cosine similarities are marked by black lines on the violins. Cell lines suggested in the article are shown in red font.
[pages=-]appendixB.pdf