Decoding glycosylation potential from protein structure across human glycoproteins with a multi-view recurrent neural network

Benjamin P. Kellman,[1,2,3,4,5,*] Julien Mariethoz,[7] Yujie Zhang,[1] Sigal Shaul,[1,2] Mia Alteri,[1,2] Daniel Sandoval,[6] Mia Jeffris,[1,2] Erick Armingol,[1,2,3] Bokan Bao,[1,3] Frederique Lisacek,[7,8] Daniel Bojar,[9,10,*] Nathan E. Lewis[1,2,5,*]

* Corresponding authors

[1] Department of Pediatrics, University of California, San Diego, La Jolla, CA 92093, USA
[2] Department of Bioengineering, University of California, San Diego, La Jolla, CA 92093, USA
[3] Bioinformatics and Systems Biology Graduate Program, University of California, San Diego, La Jolla, CA 92093, USA
[4] Augment Biologics, La Jolla, CA 92092
[5] Ragon Institute of MGH, MIT, and Harvard, Cambridge, MA, USA
[6] Department of Cellular and Molecular Medicine, University of California, San Diego, La Jolla, CA 92093, USA
[7] Proteome Informatics Group, Swiss Institute of Bioinformatics, CH-1227 Geneva, Switzerland
[8] Computer Science Department & Section of Biology, University of Geneva, route de Drize 7, CH-1227, Geneva, Switzerland
[9] Wallenberg Centre for Molecular and Translational Medicine, University of Gothenburg, Gothenburg 41390, Sweden
[10] Department of Chemistry and Molecular Biology, University of Gothenburg, Gothenburg 41390, Sweden
Correspondence to:
Benjamin P. Kellman,
Daniel Bojar,
Nathan E. Lewis, 9500 Gilman Dr. MC 0760, La Jolla, CA 92093, email: nlewisres@ucsd.edu

Abstract

Glycosylation is described as a non-templated biosynthesis. Yet, the template-free premise is antithetical to the observation that different N-glycans are consistently placed at specific sites. It has been proposed that glycosite-proximal protein structures could constrain glycosylation and explain the observed microheterogeneity. Using site-specific glycosylation data, we trained a hybrid neural network to parse glycosites (recurrent neural network) and match them to feasible N-glycosylation events (graph neural network). From glycosite-flanking sequences, the algorithm predicts most human N-glycosylation events documented in the GlyConnect database and proposed structures corresponding to observed monosaccharide composition of the glycans at these sites. The algorithm also recapitulated glycosylation in Enhanced Aromatic Sequons, SARS-CoV-2 spike, and IgG3 variants, thus demonstrating the ability of the algorithm to predict both glycan structure and abundance. Thus, protein structure constrains glycosylation, and the neural network enables predictive *in silico* glycosylation of uncharacterized or novel protein sequences and genetic variants.

Introduction

Glycosylation is difficult to study as the one supposedly non-templated biopolymer.[1] Unlike RNA, DNA, and proteins, glycan sequences are understood to be determined by local metabolic and enzymatic conditions, including the availability of charged nucleotide sugars, enzyme availability, Golgi localization, and substrate competition.[2] These well-supported claims do not explain how different glycosylation sites within one protein are consistently differentially glycosylated; a phenomenon called "microheterogeneity."[3]

Indications of protein structure bounded biosynthesis for glycans has existed for decades. After the N-glycosylation sequon (NX[S/T]) was defined, proximal-amino acid variation was found to impact glycosylation complexity,[4-6] occupancy,[7] efficiency,[8] and glycan class.[9] Conversely, amino acid sequence alignments of similarly glycosylated glycosites suggest the presence of glycosite-flanking sequence conservation.[10] In influenza and HIV, variation in glycosylation and

genetic variation proximal to glycosites can facilitate immune evasion.[11,12] Examples of how the

protein context can constrain glycosylation include observations of higher-order structures such

as β-sheets and α-helices,[13] accessibility,[14–16] and glycosylation kinetics,[17–20] all of which impact

glycan structure. We quantified associations between glycan substructures and local protein

structure, showing that protein structural constraints can predict glycosylation. Together, these

protein-glycan relations form a more comprehensive framework we call bounded biosynthesis,

wherein glycosylation is bounded by both metabolic conditions and genome-encoded protein

structural constraints.[21] That study describes protein structure as a major determinant of

glycosylation, but there is a need to functionalize the proteomic bounds on glycosylation such

that it can be leveraged with ease to predict glycosylation from protein structure.

Machine learning can be applied to the complex structures of glycans for the analysis of glycan

structure, function, and classification. For example, natural language processing can encode

glycans longitudinally from the reducing end.[22,23] The SweetTalk glycan embedding

recapitulated both antigenic glycans and microbial pathogenicity and phylogeny. Another study

leveraged the branched nonlinear glycan structure to scaffold graph convolutional neural

networks.[24] SweetNet identified glycan targets of viral lectins. Beyond glycan embedding,

biosynthetic constraints and outcomes have been modeled using neural networks.[25] Previous

attempts have been made to relate glycan branching with glycosite-proximal protein structure.[26]

In the absence of meaningful embeddings and biosynthetic-substructure decomposition like

SweetNet and GlyCompare,[27] previous observations were limited to the association between

surface accessibility and glycan complexity. With these new embeddings and the knowledge

that glycan biosynthesis is a protein structure guided process, we can now functionalize protein-

based glycan predictions.

Here we present the Interloping Saccharide Neural Network Extrapolation (InSaNNE) model,

which predicts N-glycosylation from glycosite-proximal protein features. Using long short-term

memory (LSTM) units,[28] a type of recurrent neural network, we analyze glycosite-proximal

amino acids and leverage the functional and biosynthetic glycan encodings of SweetTalk,

SweetNet, and GlyCompare to generate an accurate mapping of glycan structure to protein

sequence and structure. We train and validate our glycosite-glycan pairing model on empirically

observed site-specific glycosylation. The model is trained using data from UniCarbKB[29] and

validated using more extensively curated data from GlyConnect[30]. We further validate our

predictions on important glycosylation events on the coronavirus spike protein, immunoglobulin,

and the enhanced aromatic sequon. All N-glycan predictions are integrated in GlyConnect for

easy access. With InSaNNE, we leverage the new bounded biosynthesis paradigm to open

glycobiology to everyone by predicting expected and differential glycosylation onto their proteins

of interest.

78 Results

79 Graph convolutional neural networks accurately predict glycan-
80 glycosite pairs

We developed a model to predict the presence of specific glycans given the flanking amino acid

sequence at N-linked glycosylation sites. Specifically, glycan structures can be ranked to

indicate the most feasible glycosylation events at a glycosite of interest. To train, validate, and

test the model, we collected and annotated 1,721 unique glycosylation events across 75 human

glycoproteins from UniCarbKB[29] wherein glycan structure was previously fully determined (see

Methods). The model incorporates modules that analyzed both glycan structures (**Figure 1**a)

and the protein sequences (**Figure 1**b). To analyze the protein sequences, we used long short-

term memory (LSTM) units,[28] a recurrent neural network module effective at modeling protein

structure by asserting language-like processivity[31] (**Figure 1**b). Both sequence-proximal

(glycosite-flanking) and spatially proximal (within n-Angstroms) protein features are important for

predicting feasible glycosylation. We examined two separate LSTM-based modules into our

model for analyzing the sequence-proximal and spatially proximal amino acids, separately. For

the analysis of the glycan component, we tested three glycan embeddings: (1) a fully connected

neural network using GlyCompare glycan substructure features[27] as input, (2) a glycan-based

language model in the style of SweetTalk,[23] and (3) a graph convolutional neural network based

on SweetNet.[24]

On average, the model based on GlyCompare glycan substructure features achieved a 76.3%

accuracy in predicting which glycans have been observed at specific glycosites (**Table** 1). The

recurrent neural network (SweetTalk; 79.9%) or graph convolutional neural network (SweetNet;

83.1%) models further improved the performance, demonstrating that optimizing the glycan

analysis modules increases prediction performance. Choosing the SweetNet-based model as

our best-in-class performer, we used stochastic weight averaging (SWA; Izmailov et al., 2019)

to further optimize performance. SWA improved SweetNet-based model accuracy to 87.5%

(**Table** 1) and was therefore selected as our final model and used for all downstream analyses.

Table 1 – A model for glycan-glycosite matching was developed to predict permissible glycans on a glycosylation site.
Modules analyzing the glycosite-flanking protein sequence and additional spatially proximal amino acids consisted of
recurrent neural networks, while the module analyzing glycans was either a fully connected neural network using
GlyCompare substructure features as input (GlyCompare), a glycan-based language model (SweetTalk), or a graph
convolutional neural network (SweetNet). We further tested the effect of stochastic weight averaging (SWA) on model
performance. Removing the information about spatially proximal amino acids from the model input is denoted by "-
Spatial" while the addition of the whole protein sequence as an additional input for the model is indicated by
"+Whole". Results represent the mean values for accuracy and area under the curve (AUC) for the receiver-operator
curve (ROC) on our test set after five independent training runs.

Metric	GlyCompare	SweetTalk	SweetNet	SweetNet SWA	SweetNet SWA -Spatial	SweetNet SWA +Whole
Accuracy	0.763	0.799	0.831	0.875	0.861	0.879
ROC AUC	0.823	0.871	0.894	0.929	0.920	0.930

After optimizing the glycan analysis module, we analyzed the role of protein sequences on

prediction performance. We trained a model that only had access to the glycan structure and

the glycosite-flanking sequence, without additional spatially (3D) proximal amino acids.

Compared to the full InSaNNE model (87.5% accuracy), the model without spatially proximal

amino acids achieved a slightly worse performance (86.1%, **Table** 1). The marginal

performance loss suggests that, while spatially proximal information helps, the glycosite-flanking

residues are most important.

We next trained a model with access to the whole sequence of each protein, in addition to

glycosite-proximal amino acids, and glycan structures. The additional information from the

whole protein slowed training and inference, while providing a limited performance improvement

(87.9% accuracy, **Table** 1). We concluded that distant amino acids carry limited relevant

information for predicting permissible glycan structure that is not already captured in the nearby

sequence and spatially proximal amino acids.

Different glycosites prefer specific glycan features

True negatives, infeasible glycans, are hard to obtain experimentally, so we focused on recall
(True Positive Rate). InSaNNE achieved a recall of 84.8% for *N*-linked glycosylation events in
our dataset. The notable performance in these glycan-type-specific models, suggests that
InSaNNE performs with exceptional recall – recovering most permissible glycans at a given
glycosite.

Next, we examined which N-glycan motifs were more difficult for InSaNNE to predict. For this,
we calculated the average prediction accuracy for each glycan feature in the validation set.
Several rare glycan motifs (<10 observations) were more difficult to predict (**Figure 2**a).
However, InSaNNE exhibited a predictive accuracy of >80% for most motifs (**Figure 2**b). Since
glycan features represent a hierarchical feature set, rare motifs with low prediction accuracy are
not independent from each other and formed clusters based on glycan structure similarity
(**Figure 2**c). For example, glycan features with lower predictive performance were enriched for
oligomannose. Analogous to the glycan features, most glycosites exhibited an aggregate
predictive accuracy >90% (**Figure 2**d) and we found prediction performance correlated with the
number of observed glycans for similar glycosites (close in the embedding manifold;
Supplementary Figure 1). Predictions were robust to the removal of single amino acids or
short motifs, suggesting redundancy within glycosite-flanking sequences and soft boundaries on
the flanking window size (**Supplementary Figure** 2). Furthermore, the flanking residues, rather
than the central sequon-proximal residues, informed model predictions the most; ablation of
upstream residues was most impactful on performance (**Supplementary Figure** 2). In general,
given the consensus sequence of *N*-linked glycosylation, flanking residues are more variable,
and may carry more information for deep learning models, than more conserved sequon-
adjacent residues.

To illustrate the capabilities of InSaNNE, we used the model to predict the feasibility of all
glycans in our dataset at the glycosite GTVLTR<u>N</u>ETHATYS (P07911:N396) from human
uromodulin – the most abundant protein in human urine and relevant for chronic kidney
disease.[32] Notably, 58 of 61 experimentally observed glycans were placed in the top 80
predicted glycans (**Figure 2**e). Additionally, top glycans that were not previously reported at this
glycosite shared features with the observed glycans, such as a strong negative charge via
sialylation and/or sulfation. These results further demonstrate protein-sequence-based glycan
prediction and emphasize the value and relevance of our model.

Single amino acid changes modulate specific glycan features

While the ablation of individual glycosite-flanking amino acids does not substantially diminish model performance (**Supplementary Figure** 2), glycosylation efficiency and range can be impacted by glycosite-flanking mutations.[5,6,9,11] Therefore, we tested if InSaNNE can predict how changes to the glycosite-flanking sequence will impact glycosylation. This could facilitate glycoengineering and elucidate structural interactions between protein and glycan structures at the glycosylation site. We performed a deep mutational scan *in silico* (replacing each of the 14 glycosite-flanking amino acids with all amino acids) on every N-glycosite in our dataset. Using the modified glycosite sequences as inputs for InSaNNE, we analyzed the changes in predicted glycans compared to the wild-type sequence. To focus interpretation, we grouped glycans into "sialylated" and "fucosylated." This allowed us to track the changes in predicted probability for each of these features following specific glycosite-flanking mutation (**Figure 3**, **Supplementary Figure** 3). However, while these reflect general trends of individual glycosites across all proteins, amino acid substitutions may have effects that deviate from these general trends.

For multiple amino acid substitutions, we observed distinct changes in the predicted glycosylation of modified glycosites, with clear differences between changes to upstream and downstream regions. The introduction of some amino acids (e.g., tyrosine; **Figure 3**a) had the same qualitative effect regardless of where they were introduced. Meanwhile, other amino acids (e.g., cysteine; **Figure 3**b) have diverging effects, with a decrease in predicted complex glycans when introduced upstream and an increase when it is present downstream. We also observed that predicted changes in glycosylation were impacted more strongly by mutations in the distal parts of the glycosite-flanking sequence (e.g., glutamate; **Figure 3**c). These general trends of amino acid-glycan associations could be useful for glycosite-specific glycoengineering.

Uncharacterized glycoproteins and glycan compositions can be annotated with candidate glycan structures

Computational prediction to annotate protein features and functions is done routinely for newly discovered proteins, yet limited *in silico* characterizations exist for glycosylation. However, the relative speed of predicting glycosylation would make it invaluable for new, existing, or poorly characterized proteins; typical glycoprofiling approaches can otherwise take several months.

Even many well-characterized glycoproteins have only compositional measurements
(unstructured monosaccharide counts) since glycan structure measurement and
characterization are resource and expertise-intensive processes. Thus, InSaNNE could be
invaluable for annotating glycosylation sites.

Predicting glycosite location is one of the few high-confidence bioinformatic predictions involving
glycosylation.[33–37] To extend this capability, we predict the feasible glycan structures of 2,763
human N-linked glycosites in the GlyConnect database.[30] For this, we used InSaNNE to analyze
the annotated glycosylation sites together with the six upstream and seven downstream amino
acids. For each glycosite, we predicted the likelihood of 199 N-linked glycans (**Supplementary**
Dataset 1). Using our independent test set, we ascertain a threshold with an acceptable false-
positive rate (AUC 0.92, **Figure 4**a). A threshold of 0.6 (predicted presence) corresponded to a
false-positive rate <10% while maintaining a true positive rate >85%. This allowed us to assess
the recall or sensitivity of our predictions within GlyConnect by quantifying known glycan
structures that were successfully predicted (**Figure 4**b). Thus, InSaNNE could inform future
experiments and comparative analyses of structure-based constraints in glycosylation and
functional impacts.

InSaNNE predicts complex glycans in the enhanced aromatic sequon and the SARS-CoV-2 Spike

N-glycans are commonly grouped into categories, such as highly processed complex glycans,
hybrid glycans, and immature oligomannose glycans.[38] Previous work showed that an aromatic
residue located two-positions N-terminal from a glycosylation site results in less complex N-
glycosylation at the site, termed the enhanced aromatic sequon.[6] In this case, an L to F
substitution two residues upstream of the CD2 glycosylation site transformed the site from
predominantly complex (sialylated) and hybrid structures to low complexity (oligomannose)
structures. When InSaNNE evaluates the same sequences, the F allele sequence shows
significantly higher predicted presence for higher-mannose structures. We predict an
enrichment for 7-mannose structures (One-sided Mann-Whitney-Wilcoxon, p=0.017) and predict
an overall increase in oligomannose structure for the F allele (Linear model; Wald, p<0.001; F-
statistic, p=7.44×10^{-5}; **Figure 5a**). We see a corresponding decrease in sialylated structures in
the F allele (One-sided Mann-Whitney-Wilcoxon, p<1e-4; **Figure 5b**).

InSaNNE also recapitulates glycan types of SARS-CoV-2. These sites have been extensively
characterized throughout the pandemic.[15,39–41] N234, N717, and N801 are highly reproducible
oligomannose sites.[15] Oligomannose at N234 is consistently high (80-100%)[15] and appears
necessary to support the open ACE2-binding spike conformation.[42] Our predictions show strong
preference for Man5 and Man9 structures and a strong anticorrelation with sialylation (**Figure**
5c-d). Sites N717 and N801[15]) are predicted here to have almost no sialylation (**Figure** 5**c-d**).
Predictions for all glycosylation sites were mostly consistent with empirical observations
(**Supplementary Figure** 4).

We wondered if the spike protein of new strains shows predictable changes in glycosylation. We
examined InSaNNE predictions at site N616 in a simulated D614G variant (**Supplementary**
Figure 6) and N717 in a T716I variant (**Figure** 5**e-f**). We found distinct changes in predicted
glycosylation. T716I, between the furin cleavage site and the fusion peptide, is within the more
conserved S2 sequence and retains moderate antibody accessibility regardless of RBD
conformation.[43] To focus on relevant changes, we examined those with non-negligible ancestral
predicted-presence (>0.1) and substantial fold change (|logFC|>1) relative to the ancestral
spike. At site N717 in the T716I variant, many asialylated sugars with one to three galactose
residues decrease relative to ancestral (**Figure 5f**, blue points). Additionally, a small number of
sugars with zero to two sialic acids and one to four galactose residues increase. Though
InSaNNE predicts that site N717 becomes variably permissible to mono-, di-, tri- and tetra-
antennary sialylated and asialylated structures, empirically, it is an oligomannose site,
suggesting these terminal galactoses may not be visible without additional mutations to the site.
Distinctly, InSaNNE reveals few confident changes at site N616 in the D614G variant
(**Supplementary Figure** 6). If glycan structure can be predicted from primary sequence, site
occupancy may also be bound by these constraints.

InSaNNE predictions recapitulate biantennary abundance on human IgG3

Mutations can perturb glycosylation in IgG3.[9] Eight complex biantennary structures in human
IgG3 were measured for wildtype (*wt*) and glycosite (N297; P01860:N227) proximal mutants.
While the *wt* IgG3 showed a preference for core-fucose and a1-6-branch galactose, R301A

increased all terminal galactose, and Y296A accepted no galactosylation (**Figure** 6**a**). Thus,
primary protein structure can profoundly influence glycosylation.

We compared InSaNNE predictions for the R301A and Y296A mutants and found that
predicted-presence and change in predicted-presence were correlated with empirical
occupancy. Abundance-prediction correlation was high for the R301A mutant (R^2=0.876; **Figure**
6**b**) and moderate for *wt* abundance (R^2=0.25; **Figure** 6**b**). Predicted presence was consistent
with measured abundance in the Y296A mutant (R^2=0.33; **Figure** 6**b**). Interestingly, prediction
performance increased when we compared changes relative to *wt*. The predicted presence log
fold-change in R301A relative to *wt* was highly correlated with measured abundance log fold-
change (R^2=0.87; **Figure** 6**c**). Yet, the consistency in predicted vs observed change for Y296A
decreased dramatically (R<0, R^2=0.27; **Figure** 6**c**). To further probe the prediction failure in
Y296A, we removed glycans with small predicted changes (|logFC|<1). Without the low-
confidence changes, abundance prediction performance for *wt* (R^2=0.52), R301A (R^2=0.99),
and log fold-change (R301A vs. *wt*: R^2=0.95) improved (**Figure** 6**d-e**), while nearly all
predictions for Y296A dropped out. These results suggest that InSaNNE can predict occupancy
and occupancy change for non-small (|log fold-change|>1) changes.

Accessing InSaNNE predictions and continuous comparison through GlyConnect

We evaluated the agreement between InSaNNE predictions and GlyConnect data at the
compositional level. **Figure** 7**a** shows the protein-page d3 heatmap illustration comparing
GlyConnect-annotated glycosylation events for human coagulation factor XI (UniProt:P03951;
GlyConnect:818) with InSaNNE predictions; GlyConnect:818 is supported by four published
references. **Table 2** summarizes the comparison between GlyConnect annotation and InSaNNE
predictions for human coagulation factor XI.

	reported structures	*reported compositions*	*predicted structures*	*overlap*
Asn-90	7	0	2	2
Asn-126	5	4	4	2
Asn-163	2	1	9	4
Asn-450	4	4	4	2
Asn-491	10	5	6	6

Total glycans	42	27	16
Number of compositions	14	7	7

The first composition, H5N4 (five hexoses and four hexosamines), matches three structures with similar linkages recorded in GlyConnect. At site P03951:N491, in composition block H5N4, we see InSaNNE correctly predicts the presence of GlyConnect glycan 3471; the dashed-line compositional matches to glycans 2363 and 3233 are expected as all three glycans are members of the same composition block. Additionally, glycan 2363 is highly predicted at N491 suggesting a partial linkage resolution for the incompletely determined structure stored in GlyConnect. Likewise, structures matching the H5N4S2 (five hexoses, four hexosamines, and two sialic acids) compositions contain glycan 3353 predicted and observed at all sites. Within composition block H5N4S2, InSaNNE predicts a higher likelihood (>0.9) for glycan 1641 at N163 (biantennary α2,3-Neu5Ac). Glycan 1641 offers a complete resolution of structural ambiguity for H5N4S2 at N163. Prediction and annotation both involve flexible linkage definitions, particularly for non-core residues. In contrast, the prediction at site N163 is more extensive than reported data. Interestingly, N163 is a rare NXC sequon, which may explain the smaller number of reported structures and provides novel insights into the distinct preferences of this rare sequon.

For human coagulation factor XI, GlyConnect contains site-specific observations of 42 structures and compositions, and 14 additional distinct but structurally related glycans (Table 2). Compositional similarity was displayed using Compozitor (**Figure** 7**b**). The Compozitor graph shows 14 compositional nodes connected through the addition of a single monosaccharide. Two virtual nodes (green: H6N4S2 and H5N5S2) are needed to fully connect the graph.[44] All site-specific InSaNNe-predicted structures correspond to previously annotated site-specific compositions in GlyConnect (magenta). InSaNNE fails to predict structures corresponding to three previously reported compositions the H6N5S2, H6N5F1S2, and H6N5F1S23.

311 Interestingly, the glycan property distribution (**Figure** 7**c**) is similar between reported and
312 predicted compositions, suggesting a lack of systematic bias that would diminish expected
313 performance for specific glycotypes. Other compositions were found in large scale
314 glycoproteomics experiments without any precise structural features and may be less reliable
315 annotations.

Discussion

Here we present InSaNNE, the Interloping Saccharide Neural Network Extrapolation, for
predicting glycans on membrane-bound and secreted proteins. This approach employs a
recurrent neural network and a graph convolutional neural network with stochastic weight
averaging to predict feasible glycan structures based on the underlying protein sequence.
InSaNNE successfully predicts known glycan structures on a wide range of proteins and
assesses the impact of single amino acid substitutions on resulting glycan structures. Beyond
initial cross-validation and test-set validation, we successfully predicted glycans on uromodulin,
SARS-CoV2, IgG3, and across the GlyConnect database. We have added the glycan
predictions to the glycome database GlyConnect, making them accessible for further study of
this discovery. Importantly, InSaNNE further questions the premise of template-free glycan
biosynthesis. Glycosylation through the bounded biosynthesis paradigm, and its accessibility
through the InSaNNE framework, will facilitate more accurate and accessible study of diverse
glycoproteins and glycoproteomic behaviors.

InSaNNE enables the draft annotation of glycosylation on novel proteins, glycoprotein
composition analyses, glycoinformatics, and whole proteomes. By increasing the predictability
of glycans, we have reduced the challenge of measuring glycans. Mass spectrometry is the gold
standard in glycan measurement today, but these measurements may produce partially
ambiguous structures and topologies. Consequently, the field is rich with datasets and
databases of partially or minimally assembled glycoprofiles.[45–48] Combining measured glycan
compositions with site-specific predictions of feasible glycosylation should facilitate automated
glycoprofile assembly. These annotations can be completed for novel and existing glycoprofile
assemblies; because of the automated nature, structural glycoprofiles can be assembled for
single experiments or entire databases with comparable ease. The sequence-only nature of the
prediction is especially important, as many proteins lack experimental structural observations;
an algorithm that can operate on the primary sequence is considerably more portable than one

requiring structural information. A sequence-only prediction can even be used to quickly

compare different isoforms or predict glycans on newly discovered protein sequences.

We demonstrated our ability to glycosylate an entire proteome by predicting decoration

throughout GlyConnect. Newly glycosylated proteins can be used to identify lectin-binding,

glycan co-ligands, alternative charge, or steric conformations on proteins of interest, and

changes in protein dynamics. These predictions can be disseminated to enrich databases

detailing glycosylation[30,49,50] and other post-translational modifications,[51–53] protein structure,[54,55]

domains,[56,57] and interactions.[58–61] Future work will extend this approach to *O*-linked glycans, an

even more challenging endeavor due to less available data for training and a seeming absence

of a clear consensus sequence on the protein side.[62]

Predicted glycosylation can be used to inform large genetic and genome-wide studies. Genetic

variation can change protein function and resulting phenotype, but here we demonstrate that it

can impact glycosylation. InSaNNE can predict such changes and thus provide further

hypotheses for elucidating disease mechanisms. For example, adding predicted differential

glycosylation to a study of a high-heterogeneity critical immune gene like Human Leukocyte

Antigen (HLA) will be invaluable. This is because HLA has a functional binding-groove adjacent

glycosite[63,64] that could contribute to the behavior, accessibility, and peptide presentation. Some

HLA molecules have already been observed to carry allotype-specific glycans.[65] Beyond HLA,

understanding differential glycosylation on reference and variant molecules can help distinguish

benign from pathogenic mutations: characterized (e.g., ClinVar) or uncharacterized (e.g.,

precision medicine). Additionally, certain glycoforms can modulate secretion.[66,67] Because each

glycan may confer a change in behavior, phenotypes of highly diverse glycoproteins such as

secretion, protein-ligand interactions, cell-cell interactions, and extracellular protein complexes

can be enriched by knowledge of glycosylation. These are only a few of the studies that may

benefit from protein-predicted glycosylation potential.

Bounded biosynthesis provides a more complete picture of immune evasion by evolving

pathogens. Glycan-coated viruses have been responsible for many pandemics, while nearly

every decade has seen epidemic strains of viruses, such as influenza. Recent work has

highlighted the alignment of these fluctuations with changes in glycans decorating these

viruses.[12] Without specific glycoforms, it is not possible to determine which of these viruses

successfully disguised critical immune epitopes and which viruses created or maintained new

lectin-targeted epitopes. With specific glycan prediction, we may predict the most concerning
mutations, those that may reinforce a glycan shield,[11,68–70] stabilize virulence factors,[42] or
occlude immunogenic antigens.[71] Glycoform predictions can provide these missing data along
with previously inaccessible insight into the history and future of viral evolution.
381
382 In summary, bounded glycan biosynthesis, as functionalized by InSaNNE and made accessible
383 through GlyConnect, will enable investigators to easily consider glycosylation across many
384 areas of biological study. InSaNNE will thereby sharpen our understanding of the extracellular
385 space and innumerable intercellular phenotypes.
386

Acknowledgements

388 This work was supported by NIGMS (R35 GM119850, NEL), the Novo Nordisk Foundation
389 (NNF20SA0066621, NEL), and a Branco Weiss Fellowship – Society in Science awarded to
390 D.B.
391

Conflicts

393 This work is associated with a provisional patent filed by the authors, and Augment Biologics,
394 founded by BK and NEL.
395

Methods

Site-specific glycosylation training set construction

Empirical site-specific glycosylation data from humans was obtained from UnicarbKB[29] and
Glyconnect[72] with supplemental information from GlyGen.[73] The protein structure annotation
was done using the Structural Systems Biology (ssbio) package in python.[74] Protein structure
analysis was performed in Python v2.7.15 using ssbio v0.9.9.8 to retrieve and calculate: existing
empirical and homology models from PDB and SWISSMOD (PDBe SIFTS),[75] *de novo*
homology models (I-TASSER v5.1), sequence properties (EMBOS v6.6.0.0 pepstats), sequence
alignment (EMBOS v6.6.0.0 needle), secondary structure (DSSP v3.0.0, SCRATCHv1.1::sspro
and SCRATCHv1.1::sspro8), solvent accessibility (DSSPv3.0.0 and FreeSASAv2.0.2), and
residue depth (MSMSv2.2.6.1). Additional amino acid aggregate features were calculated using

R::seqinr. Glycan structures were annotated using a combination of glypy[76] and GlyCompare[27]

for structure parsing and comparison, respectively. All glycan substructures, a connected subset

of monosaccharides with and without linkage information, were extracted from each glycan,

merged to make a superset of substructures, then mapped to each glycan. This resulted in a

mapping from every glycan in the input database to shared substructures.

For the dataset used to train InSaNNE, we extracted 1,721 unique glycosylation events from

UniCarbKB.[29] This included the glycan structure that was observed and the glycosite-flanking

sequence (14 amino acids, with the glycosylated amino acid in the center) and structural

information in the form of additional amino acids within 6Å if structural simulations converged.

As negative examples, we generated the same number of combinations of glycosites and

glycans that have not been observed.

Model construction

All glycan-glycosite matching models comprised (1) a recurrent neural network that analyzed

the amino acid sequence of the glycosite, (2) another recurrent neural network analyzing the

amino acids of the three-dimensional glycosite surroundings, (3) a model analyzing the glycan

structure, described below, and (4) a part consisting of fully connected layers to use the

concatenated features generated by the previous modules to predict whether a glycan is

permissible at a glycosite. The recurrent neural networks consisted of a 128-dimensional

embedding layer followed by two bidirectional long short-term memory (LSTM) layers. The fully

connected model part consisted of a linear layer, a leaky ReLU (rectified linear unit) activation

function, a batch normalization layer, and a multi-sample dropout scheme[77] followed by a

sigmoid function.

We compared three different model architectures for the glycan analysis module. For assessing

GlyCompare,[27] the glycan analysis module comprised a fully connected neural network using

the 12,259 GlyCompare features as inputs for two linear layers interspersed with dropout, leaky

ReLU, and batch normalization layers. For the model containing a SweetTalk-based language

model for glycan analysis,[22] we converted glycans to glycowords and used a bidirectional

recurrent neural network for protein sequences. For the SweetNet-based model,[24] we converted

glycans to graphs by constructing a list of nodes (representing monosaccharides or linkages)

and edges to denote graph connectivity. All glycan processing for SweetTalk and SweetNet was

done using glycowork version 0.5.[78] The corresponding model contained an embedding layer
and three graph convolutional layers, interspersed by leaky ReLUs, Top-K pooling layers, and
both global mean and global maximum pooling operations. Model architectures and
hyperparameters were optimized using cross-validation.

Model training and prediction

All models were trained with an NVIDIA® Tesla® K80 GPU using PyTorch version 1.11.0.[79] We
split the data on a protein level into 80% for training and 20% for testing. For the RNNs, all
glycosite-flanking protein sequence and glycan structure were brought to the same length by
padding. Linear layers and RNNs were initialized using Xavier initialization[80] while SweetNet-
type models were initialized using a sparse initialization scheme with a sparsity of 10%.

We used a batch size of 64 for all models. As an optimizer, we used ADAM (adaptive moment
estimation) with a weight decay value of 0.00001 and a starting learning rate of 0.00001, which
was decayed according to a cosine function over 170 epochs. We trained models for a
maximum of 250 epochs, with an early stopping criterion of 25 epochs without a decrease in
validation loss. As a loss function, we used binary cross-entropy. Beginning from epoch 150, we
additionally employed stochastic weight averaging[81] with a learning rate of 0.0001.

The presence or absence of each glycan can be predicted from the trained InSaNNE model by
inputting a glycosite and glycans to predict whether these glycans could occur on this glycosite.
To heuristically boost signal for glycans with limited representation in the training set, we
generated a naturalistic background of predicted presence for each glycan. Predictions were
generated from all training-set glycosites to capture the biases and variation of the dataset as a
background predicted-presence distribution for each glycan. The background-adjusted
predicted-presence is the product of predicted presence and the predicted-presence cumulative
probability (statsmodels::ECDF v0.12.2) relative to the naturalistic background for that glycan.
462

Integration and display of predictions in GlyConnect

Using InSaNNe, we calculate the predicted presence of 512 N-linked glycans for each N-linked
glycosite in the GlyConnect dataset. Prediction data were processed to fit the requirements of

the GlyConnect database format, mainly storing association between glycans, glycoproteins and
glycosites.[30]

IUPAC-represented glycans,[82,83] output by InSaNNe, were transformed to GlycoCT[84] using the
GlyConnect   API   function,   convertIupacToGlycoct   (https://bitbucket.org/sib-pig/sugar-
converter/downloads/). Transformed prediction data was integrated in the database to enable
dynamic mapping through predefined queries for glycan structures and glycoprotein sites. Once
transformed, any update of the InSaNNe prediction will easily be reflected in the database.

JSON files resulting from querying GlyConnect REST API are used for data export and display.
A d3.js heatmap (https://d3-graph-gallery.com/heatmap) was selected as an appropriate data
visualizer. The dimensions are defined as glycan structures/compositions and glycoprotein sites
(designated by UniProt accession numbers and glycosylated amino acid sequence position).
Heatmaps are created in three types of pages: (1) protein page featuring all glycan structures
and compositions found attached to that protein, (2) structure page, featuring one structure and
the many proteins on which they are found attached, and (3) composition page, featuring all
matching glycan structures and the many proteins on which they are found attached. This data
can be exported as csv files. Prediction data can also be visualized and compared using
GlyConnect Compozitor.[44]

485
486
487

Figure Captions

Figure 1 InSaNNE model architecture. A) *Three model architectures were used to embed glycan structures in meaningful manifolds,[23,24,27] given a glycan, these models output glycan-specific coordinates within the embeddings. To analyze the GlyCompare features of glycans, we used a fully connected neural network, while a SweetTalk-based language model used linear glycan sequences and a SweetNet-based graph convolutional neural network relied on glycan connectivity (see Methods for details).* **B)** *Full model architecture of InSaNNE. The results of one of the glycan embedding modules (A) is concatenated with protein-structure and protein-sequence embeddings output by the two protein-language models. These outputs were analyzed by a fully connected neural network and yielded the predicted probability of a glycan-glycosite match. Specifically, InSaNNE takes in a 14 amino acid glycosite-flanking sequence, optional spatially proximal amino acids, and a comprehensive library of 700 representative glycans on which InSaNNE was trained. Glycan libraries containing non-represented glycans can be used following additional training.*

Figure 2 – Characterizing the glycan-glycosite-matching model InSaNNE. A) *Dependence of glycan feature prediction performance on occurrence. Using our trained InSaNNE model, we plotted the averaged prediction performance of glycan features against their counts in our dataset.* **B)** *Glycan feature accuracy distribution. A histogram of the prediction performance for all observed glycan features is shown.* **C)** *Clusters of difficult-to-predict glycan features. We used t-SNE to visualize the glycan representation learned by InSaNNE for all glycan features. Each feature was colored by its averaged prediction performance to identify structurally related clusters of glycan features that are more difficult to predict for InSaNNE (shown in brighter colors).* **D)** *Prediction performance depending on the glycosite was visualized using a t-SNE of the glycosite representations learned by InSaNNE. For all glycosites in our dataset, we averaged prediction performance over all glycans and colored glycosites by prediction performance to identify difficult to predict glycosite clusters.* **E)** *Experimentally observed and predicted glycans at a glycosylation site of human uromodulin were compared. GTVLTRNETHATYS (P07911:N396) was used to predict permissible glycans using the trained InSaNNE model, and the top 80 predicted glycans were analyzed and compared to previously observed glycans at that site [32].*

Figure 3 - Effects of amino acid substitutions on predicted glycosylation ranges. *A-C) For all N-linked glycosites in our dataset, we substituted each amino acid with tyrosine (A), cysteine (B), or glutamate (C) and input the modified glycosite-flanking sequences into our InSaNNE model and predicted feasible glycosylation. We then calculated the average change (predicted presence difference) compared to the predicted wild-type glycosylation glycosites; shown here with a 95% confidence interval. Lines for changes to fucosylated (red) and sialylated (purple) glycans are shown. See* **Supplementary Figure** *3 for analogous plots for other amino acid substitutions.*

Figure 4 - Enriching GlyConnect with InSaNNE predictions. A) *For classification thresholds between 0 and 1, we assessed true and false positive rates of InSaNNE predictions on the independent test set and compared it to a random classifier baseline.* **B)** *We validated InSaNNE predictions with existing structures on GlyConnect by investigating the influence of classification threshold on the hit rate (i.e., recall/sensitivity) of InSaNNE accurately predicting known glycan structures in GlyConnect. The grey dotted line marks the 0.6 threshold used.*

Figure 5 InSaNNE predicts complex glycans around the enhanced aromatic sequon and the SARS-CoV-2 spike protein. *(A-B) Boxplot distributions of predicted-presence for the L and F variants at N-2 stratified by number of* **(A)** *mannoses per glycan and* **(B)** *sialic acids per glycan.* **(C-D)** *Boxplots describing predicted glycosylation by (C) mannose per glycan and (D) sialic acid per glycan for three oligomannose sites in the SARS-CoV-2 spike glycoprotein. See* **Supplementary Figure** *4 for all SARS-CoV-2 spike glycosylation sites.* **(E-F)** *Fold changes of predicted glycans at site N717, labeled by number of (E) galactose and (F) sialic acid units, between the wild-type and B.1.1.7 spike protein. Predicted-presence fold-change (y-axis) is stratified by the basal predicted-presence for each glycan in the wild-type (x-axis). Predicted-presence fold-change from wild-type by galactose, mannose, GlcNAc, and sialic acid is provided for N717 and N616 in B.1.1.7 (***Supplementary Figure*** 5) and D615G (***Supplementary Figure*** 6) variants respectively. ns: p>0.05, *: p<0.05, **: p<0.01, **: p<0.001, ***: p<1e-3, ****:p<1e-4*

Figure *6 - **InSaNNE predictions of relative abundance on IgG3. A)** Heatmap showing the log-scale abundance of various glycan species observed in wt and mutant Fc on human IgG3.[9] **B)** The background-adjusted InSaNNE predicted-presence is compared with the empirical abundance in wild type (black), R301A mutant (blue), and the Y296A mutant (teal). **C)** Log fold change between glycan abundance for mutants relative to wildtype were compared between empirical and predicted abundance for all glycans. **D-E)** The bottom panels mirror panels **B-C** except glycans with a predicted absolute log fold-change less than 1 were removed.*

Figure *7 Predicted glycosylation pattern of human coagulation factor XI (P03951). H: hexose, N: hexosamine, F: fucose, S: sialic acid. (**A**) The heatmap displays the predicted presence for glycan structures at each known N-glycosite and indicates agreement with glycans previously observed at those sites retrieved from GlyConnect. The structures in each row are ordered by glycan composition; columns represent the five annotated N-glycosites of P03951. Site-specific glycan structure predictions are many-to-many relationships in the GlyConnect database since the same structure may be associated with several sites and conversely a single site may be predicted to present several similar yet non-mutually exclusive glycan structures. Composition blocks contain all structures matching a specific composition. Color indicates the strength of the predicted presence from 0.8 (lower-bound cutoff) to 1 (predicted presence upper-bound). A solid-line borders indicate exact structural matches (identical precise monosaccharides and identical linkages) while dashed lines indicate composition matches (monosaccharide category, e.g., hexose) with at least one non-identical linkage; composition-equivalent blocks (e.g., H5N4) are labelled. (**B**) A Compozitor graph representing compositional similarity between predicted and observed glycans. Fourteen glycan compositions are reported in GlyConnect for human coagulation factor XI. Nodes are connected via single monosaccharide additions represented as the edge label. Seven compositions are predicted and all included in the fourteen previously observed compositions (magenta). Two virtual nodes (green) were added to connect the graph. Numbers within the blue nodes express a correspondence in GlyConnect data between a composition and structures. When the number is absent it means we only have compositional data. The size of the non-blue nodes represents a comparison with the total content of GlyConnect to indicate the likelihood of the composition. For large nodes, the composition occurs often, irrespective of the protein where it is seen. (**C**) The bar chart represents glycan properties mapped in all subsets (database, predicted and virtual). It highlights the similarity across properties of predicted and stored structures.*

References

1. Pothukuchi, P., Agliarulo, I., Russo, D., Rizzo, R., Russo, F., and Parashuraman, S. (2019). Translation of genome to glycome: role of the Golgi apparatus. FEBS Lett. *593*, 2390–2411.

2. Kellman, B.P., and Lewis, N.E. (2021). Big-Data Glycomics: Tools to Connect Glycan Biosynthesis to Extracellular Communication. Trends Biochem. Sci. *46*, 284–300.

3. Johnson, R.L., and Deutsch, H.F. (1970). Preparation and studies of myeloma Fab subfractions. Immunochemistry *7*, 207–215.

4. Petrescu, A.-J., Milac, A.-L., Petrescu, S.M., Dwek, R.A., and Wormald, M.R. (2004). Statistical analysis of the protein environment of N-glycosylation sites: implications for occupancy, structure, and folding. Glycobiology *14*, 103–114.

5. Huang, Y.-W., Yang, H.-I., Wu, Y.-T., Hsu, T.-L., Lin, T.-W., Kelly, J.W., and Wong, C.-H. (2017). Residues Comprising the Enhanced Aromatic Sequon Influence Protein N-Glycosylation Efficiency. J. Am. Chem. Soc. *139*, 12947–12955.

6. Murray, A.N., Chen, W., Antonopoulos, A., Hanson, S.R., Wiseman, R.L., Dell, A., Haslam, S.M., Powers, D.L., Powers, E.T., and Kelly, J.W. (2015). Enhanced Aromatic Sequons Increase Oligosaccharyltransferase Glycosylation Efficiency and Glycan Homogeneity. Chem. Biol. *22*, 1052–1062.

7. Shakin-Eshleman, S.H., Spitalnik, S.L., and Kasturi, L. (1996). The Amino Acid at the X Position of an Asn-X-Ser Sequon Is an Important Determinant of N-Linked Core-glycosylation Efficiency (∗). J. Biol. Chem. *271*, 6363–6366.

8. Kasturi, L., Eshleman, J.R., Wunner, W.H., and Shakin-Eshleman, S.H. (1995). The Hydroxy Amino Acid in an Asn-X-Ser/Thr Sequon Can Influence N-Linked Core Glycosylation Efficiency and the Level of Expression of a Cell Surface Glycoprotein∗. J. Biol. Chem. *270*, 14756–14761.

9. Lund, J., Takahashi, N., Pound, J.D., Goodall, M., and Jefferis, R. (1996). Multiple interactions of IgG with its core oligosaccharide can modulate recognition by complement and human Fc gamma receptor I and influence the synthesis of its oligosaccharide chains. J. Immunol. *157*, 4963–4969.

10. Gastaldello, A., Alocci, D., Baeriswyl, J.-L., Mariethoz, J., and Lisacek, F. (2016). GlycoSiteAlign: Glycosite alignment based on glycan structure. J. Proteome Res. *15*, 3916–3928.

11. Yu, W.-H., Zhao, P., Draghi, M., Arevalo, C., Karsten, C.B., Suscovich, T.J., Gunn, B., Streeck, H., Brass, A.L., Tiemeyer, M., et al. (2018). Exploiting glycan topography for computational design of Env glycoprotein antigenicity. PLoS Comput. Biol. *14*, e1006093.

12. Altman, M.O., Angel, M., Košík, I., Trovão, N.S., Zost, S.J., Gibbs, J.S., Casalino, L., Amaro, R.E., Hensley, S.E., Nelson, M.I., et al. (2019). Human Influenza A Virus Hemagglutinin Glycan Evolution Follows a Temporal Pattern to a Glycan Limit. mBio *10*. 10.1128/mbio.00204-19.

13. Silverman, J.M., and Imperiali, B. (2016). Bacterial N-Glycosylation Efficiency Is Dependent on the Structural Context of Target Sequons. J. Biol. Chem. *291*, 22001–22010.

14. Thaysen-Andersen, M., and Packer, N.H. (2012). Site-specific glycoproteomics confirms that protein structure dictates formation of N-glycan type, core fucosylation and branching. Glycobiology *22*, 1440–1452. 10.1093/glycob/cws110.

15. Allen, J.D., Chawla, H., Samsudin, F., Zuzic, L., Shivgan, A.T., Watanabe, Y., He, W.-T., Callaghan, S., Song, G., Yong, P., et al. (2021). Site-specific steric control of SARS-CoV-2 spike glycosylation. Cold Spring Harbor Laboratory, 2021.03.08.433764. 10.1101/2021.03.08.433764.

16. García-García, A., Serna, S., Yang, Z., Delso, I., Taleb, V., Hicks, T., Artschwager, R., Vakhrushev, S.Y., Clausen, H., Angulo, J., et al. (2021). FUT8-directed core fucosylation of N-glycans is regulated by the glycan structure and protein environment. ACS Catal. *11*, 9052–9065.

17. Losfeld, M.-E., Scibona, E., Lin, C.-W., Villiger, T.K., Gauss, R., Morbidelli, M., and Aebi, M.
(2017). Influence of protein/glycan interaction on site-specific glycan heterogeneity. FASEB
J. *31*, 4623–4635.

18. Losfeld, M.-E., Scibona, E., Lin, C.-W., and Aebi, M. (2022). Glycosylation network
mapping and site-specific glycan maturation in vivo. iScience, 105417.

19. Mathew, C., Weiß, R.G., Giese, C., Lin, C.-W., Losfeld, M.-E., Glockshuber, R., Riniker, S.,
and Aebi, M. (2021). Glycan-protein interactions determine kinetics of N-glycan remodeling.
RSC Chem Biol *2*, 917–931.

20. Adams, T.M., Zhao, P., Chapla, D., Moremen, K.W., and Wells, L. (2022). Sequential in
vitro enzymatic N-glycoprotein modification reveals site-specific rates of glycoenzyme
processing. J. Biol. Chem., 102474.

21. Kellman Protein structure, a genetic encoding for glycosylation. Unpublished co-
submission.

22. Bojar, D., Camacho, D.M., and Collins, J.J. (2020). Using Natural Language Processing to
Learn the Grammar of Glycans. bioRxiv, 2020.01.10.902114. 10.1101/2020.01.10.902114.

23. Bojar, D., Powers, R.K., Camacho, D.M., and Collins, J.J. (2021). Deep-learning resources
for studying glycan-mediated host-microbe interactions. Cell Host Microbe *29*, 132-144.e3.

24. Burkholz, R., Quackenbush, J., and Bojar, D. (2021). Using graph convolutional neural
networks to learn a representation for glycans. Cell Rep. *35*, 109251.

25. Kotidis, P., and Kontoravdi, C. (2020). Harnessing the potential of artificial neural networks
for predicting protein glycosylation. Metabolic Engineering Communications, e00131.

26. Senger, R.S., and Karim, M.N. (2008). Prediction of N-linked glycan branching patterns
using artificial neural networks. Math. Biosci. *211*, 89–104.

27. Bao, B., Kellman, B.P., Chiang, A.W.T., Zhang, Y., Sorrentino, J.T., York, A.K.,
Mohammad, M.A., Haymond, M.W., Bode, L., and Lewis, N.E. (2021). Correcting for
sparsity and interdependence in glycomics by accounting for glycan biosynthesis. Nat.
Commun. *12*, 4988.

28. Sherstinsky, A. (2020). Fundamentals of recurrent neural network (RNN) and long short-
term memory (LSTM) network. Physica D *404*, 132306.

29. Campbell, M.P., Peterson, R., Mariethoz, J., Gasteiger, E., Akune, Y., Aoki-Kinoshita, K.F.,
Lisacek, F., and Packer, N.H. (2014). UniCarbKB: building a knowledge platform for
glycoproteomics. Nucleic Acids Res. *42*, D215-21.

30. Alocci, D., Mariethoz, J., Gastaldello, A., Gasteiger, E., Karlsson, N.G., Kolarich, D.,
Packer, N.H., and Lisacek, F. (2019). GlyConnect: Glycoproteomics Goes Visual,
Interactive, and Analytical. J. Proteome Res. *18*, 664–677.

31. Bepler, T., and Berger, B. (2021). Learning the protein language: Evolution, structure, and
function. Cell Syst. *12*, 654-669.e3.

32. Devuyst, O., Olinger, E., and Rampoldi, L. (2017). Uromodulin: from physiology to rare and complex kidney disorders. Nat. Rev. Nephrol. *13*, 525–544.

33. Gupta, R., and Brunak, S. (2002). Prediction of glycosylation across the human proteome and the correlation to protein function. Pac. Symp. Biocomput., 310–322.

34. Steentoft, C., Vakhrushev, S.Y., Joshi, H.J., Kong, Y., Vester-Christensen, M.B., Schjoldager, K.T.-B.G., Lavrsen, K., Dabelsteen, S., Pedersen, N.B., Marcos-Silva, L., et al. (2013). Precision mapping of the human O-GalNAc glycoproteome through SimpleCell technology. EMBO J. *32*, 1478–1488.

35. Pitti, T., Chen, C.-T., Lin, H.-N., Choong, W.-K., Hsu, W.-L., and Sung, T.-Y. (2019). N-GlyDE: a two-stage N-linked glycosylation site prediction incorporating gapped dipeptides and pattern-based encoding. Sci. Rep. *9*, 15975.

36. Taherzadeh, G., Dehzangi, A., Golchin, M., Zhou, Y., and Campbell, M.P. (2019). SPRINT-Gly: predicting N- and O-linked glycosylation sites of human and mouse proteins by using sequence and predicted structural properties. Bioinformatics *35*, 4140–4146.

37. Pakhrin, S.C., Aoki-Kinoshita, K.F., Caragea, D., and Kc, D.B. (2021). DeepNGlyPred: A deep neural network-based approach for human N-linked glycosylation site prediction. Molecules *26*, 7314.

38. Varki, A., Cummings, R.D., Esko, J.D., Stanley, P., Hart, G.W., Aebi, M., Mohnen, D., Kinoshita, T., and Packer, N.H. eds. (2022). Essentials of glycobiology, fourth edition 4th ed. (Cold Spring Harbor Laboratory Press).

39. Watanabe, Y., Allen, J.D., Wrapp, D., McLellan, J.S., and Crispin, M. (2020). Site-specific glycan analysis of the SARS-CoV-2 spike. Science *369*, 330–333.

40. Zhao, P., Praissman, J.L., Grant, O.C., Cai, Y., Xiao, T., Rosenbalm, K.E., Aoki, K., Kellman, B.P., Bridger, R., Barouch, D.H., et al. (2020). Virus-Receptor Interactions of Glycosylated SARS-CoV-2 Spike and Human ACE2 Receptor. Cell Host Microbe *28*, 586-601.e6.

41. Shajahan, A., Supekar, N.T., Gleinich, A.S., and Azadi, P. (2020). Deducing the N- and O-glycosylation profile of the spike protein of novel coronavirus SARS-CoV-2. Glycobiology *30*, 981–988. 10.1093/glycob/cwaa042.

42. Casalino, L., Gaieb, Z., Goldsmith, J.A., Hjorth, C.K., Dommer, A.C., Harbison, A.M., Fogarty, C.A., Barros, E.P., Taylor, B.C., McLellan, J.S., et al. (2020). Beyond Shielding: The Roles of Glycans in the SARS-CoV-2 Spike Protein. ACS Central Science *6*, 1722–1734. 10.1021/acscentsci.0c01056.

43. Harvey, W.T., Carabelli, A.M., Jackson, B., Gupta, R.K., Thomson, E.C., Harrison, E.M., Ludden, C., Reeve, R., Rambaut, A., COVID-19 Genomics UK (COG-UK) Consortium, et al. (2021). SARS-CoV-2 variants, spike mutations and immune escape. Nat. Rev. Microbiol. *19*, 409–424.

44. Robin, T., Mariethoz, J., and Lisacek, F. (2020). Examining and fine-tuning the selection of glycan compositions with GlyConnect Compozitor. Mol. Cell. Proteomics. 10.1074/mcp.RA120.002041.

45. Liu, M.-Q., Zeng, W.-F., Fang, P., Cao, W.-Q., Liu, C., Yan, G.-Q., Zhang, Y., Peng, C.,
Wu, J.-Q., Zhang, X.-J., et al. (2017). pGlyco 2.0 enables precision N-glycoproteomics with
comprehensive quality control and one-step mass spectrometry for intact glycopeptide
identification. Nat. Commun. *8*, 438.

46. Rojas-Macias, M.A., Mariethoz, J., Andersson, P., Jin, C., Venkatakrishnan, V., Aoki, N.P.,
Shinmachi, D., Ashwood, C., Madunic, K., Zhang, T., et al. (2019). Towards a standardized
bioinformatics infrastructure for N- and O-glycomics. Nat. Commun. *10*, 3275.

47. Riley, N.M., Hebert, A.S., Westphall, M.S., and Coon, J.J. (2019). Capturing site-specific
heterogeneity with large-scale N-glycoproteome analysis. Nat. Commun. *10*, 1311.

48. Yang, Y., Yan, G., Kong, S., Wu, M., Yang, P., Cao, W., and Qiao, L. (2021). GproDIA
enables data-independent acquisition glycoproteomics with comprehensive statistical
control. Nat. Commun. *12*, 6073.

49. Kahsay, R., Vora, J., Navelkar, R., Mousavi, R., Fochtman, B.C., Holmes, X., Pattabiraman,
N., Ranzinger, R., Mahadik, R., Williamson, T., et al. (2020). GlyGen data model and
processing workflow. Bioinformatics *36*, 3941–3943.

50. Yamada, I., Shiota, M., Shinmachi, D., Ono, T., Tsuchiya, S., Hosoda, M., Fujita, A., Aoki,
N.P., Watanabe, Y., Fujita, N., et al. (2020). The GlyCosmos Portal: a unified and
comprehensive web resource for the glycosciences. Nat. Methods. 10.1038/s41592-020-
0879-8.

51. Minguez, P., Letunic, I., Parca, L., Garcia-Alonso, L., Dopazo, J., Huerta-Cepas, J., and
Bork, P. (2015). PTMcode v2: a resource for functional associations of post-translational
modifications within and between proteins. Nucleic Acids Res. *43*, D494-502.

52. Craveur, P., Rebehmed, J., and de Brevern, A.G. (2014). PTM-SD: a database of
structurally resolved and annotated posttranslational modifications in proteins. Database
(Oxford) *2014*, bau041–bau041.

53. Li, Z., Li, S., Luo, M., Jhong, J.-H., Li, W., Yao, L., Pang, Y., Wang, Z., Wang, R., Ma, R., et
al. (2022). dbPTM in 2022: an updated database for exploring regulatory networks and
functional associations of protein post-translational modifications. Nucleic Acids Res. *50*,
D471–D479.

54. Goodsell, D.S. (2021). Fifty years of open access to PDB structures. RCSB Protein Data
Bank. 10.2210/rcsb_pdb/mom_2021_10.

55. UniProt Consortium (2021). UniProt: the universal protein knowledgebase in 2021. Nucleic
Acids Res. *49*, D480–D489.

56. Mistry, J., Chuguransky, S., Williams, L., Qureshi, M., Salazar, G.A., Sonnhammer, E.L.L.,
Tosatto, S.C.E., Paladin, L., Raj, S., Richardson, L.J., et al. (2021). Pfam: The protein
families database in 2021. Nucleic Acids Res. *49*, D412–D419.

57. Sigrist, C.J.A., Cerutti, L., de Castro, E., Langendijk-Genevaux, P.S., Bulliard, V., Bairoch,
A., and Hulo, N. (2010). PROSITE, a protein domain database for functional
characterization and annotation. Nucleic Acids Res. *38*, D161–D166.

58. Clerc, O., Deniaud, M., Vallet, S.D., Naba, A., Rivet, A., Perez, S., Thierry-Mieg, N., and
Ricard-Blum, S. (2019). MatrixDB: integration of new data with a focus on
glycosaminoglycan interactions. Nucleic Acids Res. *47*, D376–D381.

59. Chatr-Aryamontri, A., Breitkreutz, B.-J., Oughtred, R., Boucher, L., Heinicke, S., Chen, D.,
Stark, C., Breitkreutz, A., Kolas, N., O'Donnell, L., et al. (2015). The BioGRID interaction
database: 2015 update. Nucleic Acids Res. *43*, D470-8.

60. Szklarczyk, D., Gable, A.L., Lyon, D., Junge, A., Wyder, S., Huerta-Cepas, J., Simonovic,
M., Doncheva, N.T., Morris, J.H., Bork, P., et al. (2019). STRING v11: protein-protein
association networks with increased coverage, supporting functional discovery in genome-
wide experimental datasets. Nucleic Acids Res. *47*, D607–D613.

61. Franz, M., Rodriguez, H., Lopes, C., Zuberi, K., Montojo, J., Bader, G.D., and Morris, Q.
(2018). GeneMANIA update 2018. Nucleic Acids Res. *46*, W60–W64.

62. Malaker, S.A., Riley, N.M., Shon, D.J., Pedram, K., Krishnan, V., Dorigo, O., and Bertozzi,
C.R. (2022). Revealing the human mucinome. Nat. Commun. *13*, 3542.

63. Ryan, S.O., and Cobb, B.A. (2012). Roles for major histocompatibility complex
glycosylation in immune function. Semin. Immunopathol. *34*, 425–441.

64. Ilca, F.T., and Boyle, L.H. (2021). The glycosylation status of MHC class I molecules
impacts their interactions with TAPBPR. Mol. Immunol. *139*, 168–176.

65. Hoek, M., Demmers, L.C., Wu, W., and Heck, A.J.R. (2021). Allotype-specific glycosylation
and cellular localization of human leukocyte antigen class I proteins. J. Proteome Res. *20*,
4518–4528.

66. Sagt, C.M., Kleizen, B., Verwaal, R., de Jong, M.D., Müller, W.H., Smits, A., Visser, C.,
Boonstra, J., Verkleij, A.J., and Verrips, C.T. (2000). Introduction of an N-glycosylation site
increases secretion of heterologous proteins in yeasts. Appl. Environ. Microbiol. *66*, 4940–
4944.

67. Olczak, M., and Szulc, B. (2021). Modified secreted alkaline phosphatase as an improved
reporter protein for N-glycosylation analysis. PLoS One *16*, e0251805.

68. Harbison, A.M., Fogarty, C.A., Phung, T.K., Satheesan, A., Schulz, B.L., and Fadda, E.
(2022). Fine-tuning the spike: role of the nature and topology of the glycan shield in the
structure and dynamics of the SARS-CoV-2 S. Chem. Sci. *13*, 386–395.

69. Wei, C.-J., Boyington, J.C., Dai, K., Houser, K.V., Pearce, M.B., Kong, W.-P., Yang, Z.-Y.,
Tumpey, T.M., and Nabel, G.J. (2010). Cross-neutralization of 1918 and 2009 influenza
viruses: role of glycans in viral evolution and vaccine design. Sci. Transl. Med. *2*, 24ra21.

70. Go, E.P., Ding, H., Zhang, S., Ringe, R.P., Nicely, N., Hua, D., Steinbock, R.T., Golabek,
M., Alin, J., Alam, S.M., et al. (2017). Glycosylation Benchmark Profile for HIV-1 Envelope
Glycoprotein Production Based on Eleven Env Trimers. J. Virol. *91*. 10.1128/JVI.02428-16.

71. Grant, O.C., Montgomery, D., Ito, K., and Woods, R.J. (2020). Analysis of the SARS-CoV-2
spike protein glycan shield reveals implications for immune recognition. Sci. Rep. *10*,
14991.

72. Mariethoz, J., Alocci, D., Gastaldello, A., Horlacher, O., Gasteiger, E., Rojas-Macias, M.,
Karlsson, N.G., Packer, N.H., and Lisacek, F. (2018). Glycomics@ExPASy: Bridging the
Gap. Molecular & Cellular Proteomics *17*, 2164–2176. 10.1074/mcp.ra118.000799.

73. York, W.S., Mazumder, R., Ranzinger, R., Edwards, N., Kahsay, R., Aoki-Kinoshita, K.F.,
Campbell, M.P., Cummings, R.D., Feizi, T., Martin, M., et al. (2020). GlyGen:
Computational and Informatics Resources for Glycoscience. Glycobiology *30*, 72–73.

74. Mih, N., Brunk, E., Chen, K., Catoiu, E., Sastry, A., Kavvas, E., Monk, J.M., Zhang, Z., and
Palsson, B.O. (2018). ssbio: a Python framework for structural systems biology.
Bioinformatics *34*, 2155–2157.

75. Velankar, S., Dana, J.M., Jacobsen, J., van Ginkel, G., Gane, P.J., Luo, J., Oldfield, T.J.,
O'Donovan, C., Martin, M.-J., and Kleywegt, G.J. (2013). SIFTS: Structure Integration with
Function, Taxonomy and Sequences resource. Nucleic Acids Res. *41*, D483-9.

76. Klein, J., and Zaia, J. (2019). glypy: An Open Source Glycoinformatics Library. J. Proteome
Res. *18*, 3532–3537.

77. Inoue, H. (2019). Multi-sample dropout for accelerated training and better generalization.
arXiv [cs.NE].

78. Thomès, L., Burkholz, R., and Bojar, D. (2021). Glycowork: A Python package for glycan
data science and machine learning. Glycobiology *31*, 1240–1244.

79. Paszke, A., Gross, S., Massa, F., Lerer, A., Bradbury, J., Chanan, G., Killeen, T., Lin, Z.,
Gimelshein, N., Antiga, L., et al. (2019). PyTorch: An Imperative Style, High-Performance
Deep Learning Library. In Advances in Neural Information Processing Systems, H. Wallach,
H. Larochelle, A. Beygelzimer, F. d\textquotesingle Alché-Buc, E. Fox, and R. Garnett, eds.
(Curran Associates, Inc.).

80. Glorot, X., and Bengio, Y. (2010). Understanding the difficulty of training deep feedforward
neural networks. In Proceedings of the Thirteenth International Conference on Artificial
Intelligence and Statistics Proceedings of Machine Learning Research., Y. W. Teh and M.
Titterington, eds. (PMLR), pp. 249–256.

81. Izmailov, P., Podoprikhin, D., Garipov, T., Vetrov, D., and Wilson, A.G. (2018). Averaging
weights leads to wider optima and better generalization. arXiv [cs.LG].

82. Sharon, N. (1986). IUPAC-IUB Joint Commission on Biochemical Nomenclature (JCBN).
Nomenclature of glycoproteins, glycopeptides and peptidoglycans. Glycoconj. J. *3*, 123–
133.

83. McNaught, A.D. (1997). Nomenclature of carbohydrates. Carbohydr. Res. *297*, 1–92.

84. Herget, S., Ranzinger, R., Maass, K., and Lieth, C.-W.V.D. (2008). GlycoCT-a unifying
sequence format for carbohydrates. Carbohydr. Res. *343*, 2162–2171.

A

Glycompare
Output
...
...
...
Glycompare Features

Sweet Talk
Output
LSTM
Glycan

Sweet Net
Output
...
...
Glycan

B

Glycan Embedding

Glycan Library

Sequon Embedding
Output
LSTM
Glycosite Sequence

Structure Embedding
Output
LSTM
Glycosite Structure

N

Sequence Proximal

Spatially Proximal

Glycosylation Site

Leaky ReLU | Batch Normalization | Multi-Sample Dropout

0.1
0.5
0.9
0.8
0.9
0.1
0.2
0.3

Glycan-Glycosite Match Probability "Predicted Presence"

Input | InSaNNE | Output

A

Examples per substructure (count)

Accuracy

B

Number of Features

Accuracy

C

Average Accuracy
- 0.0
- 0.2
- 0.4
- 0.6
- 0.8

t-SNE Dim2

t-SNE Dim1

D

t-SNE Dim2

t-SNE Dim1

E

Sorted Glycans

Predicted Presence

58/61 observed glycans rank in the top 80 predicted glycans

Experimentally Observed

Similar to Experimentally Observed

Prediction Rank 12

Prediction Rank 121

Prediction Rank 15

Prediction Rank 106

Prediction Rank 32

Prediction Rank 84

Predicted Complex and Oligomannose Structures: L to F Mutant

A Mannose

B Neu5Ac

Predicted Glycosylation for SARS-CoV-2 Spike: wild-type/ancestral

C Mannose

D Neu5Ac

Predicted Differential Glycosylation for SARS-CoV-2 Spike, site N717: B.1.1.7 vs ancestral (T716I)

E Galactose

F Neu5Ac