Dissociable encoding of evolving beliefs and momentary belief updates in distinct neural decision signals

Authors: Elisabet Parés-Pujolràs[1], Simon P. Kelly[1], Peter R. Murphy[2]

 Affiliations:

[1]University College Dublin, Dublin, Ireland

[2]Maynooth University, Maynooth, Ireland

Abstract

Making accurate decisions in noisy or otherwise uncertain environments requires integrating evidence over time. Using simple tasks requiring rapid evaluation of a stationary sensory feature, two human neurophysiological signals have been found to evolve with similar integration dynamics, with one (centroparietal positivity; CPP) appearing to compute the running integral and continuously feed it to the other (motor beta lateralisation; MBL). However, it remains unknown whether and how these signals serve more distinct functional roles in more complex scenarios where information arrives discontinuously and requires more computational steps to form appropriate belief updates. We employed a volatile expanded judgement task that permits dissociation of the encoding of raw sensory information in each of a series of discrete stimuli ('objective evidence'), the appropriately transformed belief updates from that information ('effective evidence'), and the evolving belief itself (accumulated evidence). Whereas MBL traced the evolving belief across stimuli, the CPP was found only to locally encode the effective evidence associated with each individual stimulus. Furthermore, fluctuations in CPP amplitude after each single stimulus could account for variability in the encoding of accumulated evidence in MBL. These results suggest a flexible computational hierarchy in which effective evidence can be computed sample-by-sample at an intermediate processing level to drive downstream belief updates for protracted decisions about discrete stimuli.

1. Introduction

Perceptual decisions are widely thought to involve a core process of evidence accumulation (Gold & Shadlen, 2007; O'Connell & Kelly, 2021; Shadlen & Kiani, 2013). Extensive research across species has described multiple signals in distributed brain areas that are related to decision variables (DV), reflecting the evolving evidence tally during such accumulation (e.g. de Lafuente et al., 2015; Gherman et al., 2024; Hanks et al., 2015, see Hanks & Summerfield (2017) for a review). This multiplicity of neural DV correlates has arisen through research predominantly using simple tasks that require the minimum computational ingredients of encoding and accumulating a single sensory evidence feature, usually in a single, contiguous evaluation of a stationary stimulus. Researchers are naturally turning to the question of what distinct functional roles these distributed signals might serve, and there is increasing recognition that answers may be found in employing tasks with more complex structure, context-dependencies and/or integration strategies (Okazawa & Kiani, 2023). Here, we employ a volatile expanded judgement task in which individual sensory samples must be non-linearly transformed into appropriate belief updates, to investigate the potentially distinct computational roles of two human neurophysiological signatures of decision formation that normally exhibit similar dynamics during standard stationary-stimulus tasks: motor beta lateralisation (MBL) and the centroparietal positivity (CPP).

Beta-band activity (13-30Hz) over the motor cortex is a well-known marker of motor preparation. Beta power gradually decreases during preparation of a movement of a contralateral limb (Pfurtscheller & Lopes da Silva, 1999; Stancák & Pfurtscheller, 1995), and typically reaches a stereotyped level just before action consistent with a threshold-crossing event (O'Connell et al., 2012; Steinemann et al., 2018). In perceptual decision tasks, MBL emerges at a rate proportional to the strength of evidence, and when choices are freely reported with a manual response, peaks at the time of that response (de Lange et al., 2013; Donner et al., 2009; O'Connell et al., 2012). In delayed-response tasks, provided the stimulus-response mapping is known, MBL traces decision formation during evidence viewing and sustains even after commitment is reached, until the time of a subsequent response cue (Twomey et al., 2016). Further, in a decision-making context with volatility in the evidence source, MBL has been shown to approximate a DV computed through a non-linear accumulation process that is normative for this context (Murphy et al., 2021).

The CPP, on the other hand, is a positive-going event-related potential (ERP) that evolves with the same evidence-dependent, build-to-peak accumulation dynamics as MBL during decision-making tasks with continuous sensory evidence streams, for any sensory modality (O'Connell et al., 2012), and slightly in advance of motor preparation (Kelly & O'Connell, 2013). Previous research has already established some notable properties in the CPP that are distinct from motor preparation signals: it traces decision formation even when no motor response is required (O'Connell et al., 2012), and in delayed-response tasks, peaks and falls at the time of choice commitment rather than sustaining until response, regardless of whether the response mapping is known during presentation of the sensory evidence (Twomey et al., 2016). Furthermore, where strategic adjustments and biases are enacted through changes in motor preparation starting-levels, the CPP undergoes a corresponding shift not in its starting-level but rather in the amplitude it reaches at response(Geuzebroek et al., 2023; Steinemann et al., 2018).

The characteristics of these two signals are consistent with the view that the CPP reflects an accumulation process that is directly and continuously fed to the motor system, which executes no further computations on the sensory evidence but is subject to additional task-dependent biases and triggers a response once it reaches a stereotyped threshold level. However, since most previous studies have used continuous stimuli with stationary statistics and short integration times (Geuzebroek et al., 2023; O'Connell et al., 2012; Steinemann et al., 2018; Twomey et al., 2016), it remains unclear whether and how the roles of the CPP and MBL might dissociate in more complex tasks that require additional computational steps and/or integration over longer time scales.

Here, we used a token-based sequential sampling task first developed in previous work (Murphy et al., 2021) where the generative state of the task can change unpredictably during the formation of single decisions. This task has three key advantages for our purposes. First, the temporal separation between tokens (400 ms stimulus onset asynchrony) allows us to precisely compute how the theoretical DV changes after each token, a process we refer to here as 'belief updating', and to directly map related quantities onto neural responses (Balsdon et al., 2021; Murphy et al., 2021; Wyart et al., 2012, 2015). This contrasts with other frequently used decision-making paradigms (e.g. random dot kinematogram, RDK), where stimulus fluctuations are often too noisy and too fast to enable a precise link with computational variables and evoked potentials (Hyafil et al., 2023). Secondly, our token task unfolds over

longer timescales than traditional paradigms (i.e. trials lasting several seconds), allowing us to dissociate transient activities evoked by individual stimuli (i.e. responses to a particular sample) from sustained patterns of information encoding (i.e. an evolving DV that integrates over samples). Finally, the CPP has typically been studied in stationary environments where the optimal strategy is to weigh each piece of evidence equally and to integrate perfectly over time. In such scenarios, the information provided by each sample (objective evidence), and the change in the DV evoked by it (effective evidence) are identical. In volatile environments, instead, these two quantities can be dissociated. According to the normative model for evidence accumulation on our task (Glaze et al., 2015), for example, a sample providing strong evidence against an observer's current belief results in a bigger belief update than a sample providing equally strong evidence for the observer's current belief (Murphy et al., 2021). Our task thus disentangles objective from effective evidence, and allows us to test which of them may be captured by the CPP.

To preface our results, we find that centroparietal signals in this task context transiently encode the magnitude of belief updates evoked by each sample (effective evidence), but do not exhibit a longer-lasting, sustained component that tracks the DV over the course of the trial. By contrast, MBL reflects the evolving DV throughout the trial. We further show that single-trial fluctuations in both signals capture variability in behavioural choice over and above that accounted for by the sensory information on each trial, and that variability in the CPP's encoding of belief updates are directly linked to residual fluctuations in MBL's encoding of a decision variable. Our study provides evidence for a key functional dissociation between these well-studied decision signals in the task setting investigated here, and points toward a potentially flexible neural architecture for decision formation that may be reconfigured to meet the demands of different decision-making contexts.

2.   Results

2.1. Task & behaviour

In our task (N = 20, *Fig. 1A*), participants monitored a set of checkerboard patches ('samples') appearing anywhere along a semicircular arc in the lower visual hemifield and centred on a central fixation point. Sample locations (specifically, polar angles relative to the fixation point) were drawn from one of two overlapping Gaussian distributions (generative states) with equal standard deviations and means placed symmetrically to the left and right of the vertical meridian. Samples appearing close to the vertical meridian provided weak evidence for one or the other generative states, as they could be drawn from either distribution with similar probabilities. In contrast, samples appearing closer to the horizontal meridian provided strong evidence, since they were a lot more likely to be drawn from one of the two distributions than the other. Sample onset asynchrony was 400 ms. Participants viewed a maximum of 10 samples per trial, and the distribution from which sample locations were drawn could switch during the trial with a fixed probability (hazard rate, H) of 0.1. Thus, some trials contained one, some more than one, and the rest contained no changes in the generative distribution (*Fig. 1A, right*). The participants' task was to report what they believed to be the generative state (i.e. the 'active' distribution) at the end of each trial. They completed 18 blocks of 80 trials across 2 sessions.

The normative model for this task prescribes that participants transform their posterior beliefs after accumulating each sample to form the prior for the next sample, according to a non-linear function that depends on their estimate of the environmental volatility (*Fig. 1B*; see *Eq 1-2* in Methods). In effect, this non-linearity limits the degree of belief that can be attained for either alternative and facilitates fast revision of that belief following state changes (*Fig. 1C, upper panel*), particularly when the volatility is estimated to be high. Our participants performed the task better (83.8% ± 0.66%; mean ± s.e.m.) than expected if they followed simple strategies such as of choosing based on the last sample only (79.7% ± 0.14%; p < 0.001) or the sum of all available samples (i.e. perfect accumulation, 75.5% ± 0.22%; $p < 0.001$), but worse than the ideal observer employing the normative updating rule with perfect knowledge of the generative statistics and no noise or bias (87.5% ± 0.20%; $p < 0.001$; *Fig. 1D*).

The non-linear transformation of the evolving belief that is the key feature of the normative approach to solving this task has a number of behavioural consequences. First, its slope is

always <1, which produces leaky accumulation and a recency effect in evidence weighting. Further, it produces choice behaviour that is particularly sensitive to samples indicating that a potential change in the generative state has occurred (i.e. samples that clearly conflict with a strongly held current belief), as well as to samples encountered when the belief is generally weak (i.e. when uncertainty about the current generative state is high; Murphy et al., 2021; *Fig. 1C*). To test whether participants exhibited these behavioural hallmarks of normative belief updating, we fit logistic regression models estimating the average weight on choice of each sample in the trial sequence (*Eq. 7* in Methods). The evidence provided by each sample was quantified in terms of its log-likelihood ratio (LLR). Further, we computed the *surprise* associated with a sample (defined here as the analytically computed change-point probability, which quantifies the likelihood, given the existing belief, that a sample directly followed a change in generative state; see *Eqs. 4-6*), and the *uncertainty* in the belief at the time of encountering that sample (see *Methods*). As predicted by the normative model, we found that participants weighted evidence occurring later in the trial more heavily ($p < 0.001$, two-tailed permutation test on weights for the first 5 versus the last 5 samples), and also upweighted samples associated with high change-point probability ($p < 0.001$, two-tailed cluster permutation test against zero*; Fig. 1E*). By contrast, we did not find a reliable positive modulation by uncertainty ($p = 0.286$ positive effect at sample 7). This null effect is in contrast to previous data from similar versions of our task (Murphy et al. 2021), though a weaker effect in the present case is consistent with normative model predictions due to differences in task generative statistics (here, a higher H and higher signal-to-noise in generative distributions; see Methods). Note also that "surprise" and "uncertainty" are not quantities directly used in the normative computations, but rather sensitivities to these variables are diagnostic of the normative belief updating strategy that we can test for in our data. The normative model was able to account for these behavioural effects (*Fig. 1E*) with estimated participant-specific subjective hazard rates *(H)* that did not differ from the true generative H = 0.1 at the group level (M = 0.092, s.e.m. = 0.015; $t_{(19)} = 0.54$, $p = 0.59$). It also fit the data better than several alternative models (see *Fig. S1* for full parameter estimate and model comparison details).

Thus, consistent with previous work using the same task (Murphy et al. 2021), our results show that participants' behaviour approximated normative belief updating. This in turn allowed us to use the fits of the normative model to derive decision-relevant quantities with which to investigate our neural signals of interest: MBL and the CPP.

Figure 1. Experimental task design and behavioural results. **A.** In our task, participants were required to monitor a sequence of samples of fixed duration (max. 10 samples, 400 ms stimulus onset asymmetry (SOA)) and determine which generative distribution (left/right) was active at the end of the trial. Upon presentation of a response cue, they reported their decisions pressing a keyboard key with their left/right hand respectively. There was a 0.1 probability that the generative distribution might change after any given sample (Hazard rate, $H = 0.1$), so in most trials there were either no change points (CP) or only 1 (see inset). **B.** Normative belief update schematic. **C.** Time course of prior (*top*), surprise (*middle*) and uncertainty (*bottom*) over samples for an illustrative example case of trials with a single CP at sample 5, showing the average of 100 simulated trials. Generative sources for samples in these trials switched from left to right (*dashed line*) or right to left (*solid line*). An ideal observer's normative decision variable, tracked by the normative prior (Ψ), plateaus shortly after trial start, and rapidly changes sign after the change in the generative distribution. In contrast, a perfect accumulation strategy would produce a decision variable that grows linearly over time and requires more evidence to flip sign after a CP (*magenta*). Surprise values (i.e. change point probabilities) peak at the time of a CP, and uncertainty values (-$|\Psi|$) decrease as priors grow. Note that at sample

6, when the discrepant sample at the change point has fully countered the existing belief, beliefs are weakest (i.e. Ψ close to 0) and uncertainty is maximal. The bottom panel illustrates how normative beliefs (Ψ) and step-wise belief updates ($\Delta\Psi$) evolve in this subset of trials. While the mean LLR values (*objective evidence*) remain stable, as expected, throughout the duration of any given state (i.e. before and after sample 5 in this example), belief updates become smaller (*effective evidence*, $\Delta\Psi$) as beliefs grow stronger. **D.** Proportion of observed and model-predicted correct responses. White bars and overlaid coloured points represent individual data and fits, respectively. Filled bars display grand-averaged data $\pm$ s.e.m.. **E.** Observed and model-predicted (mean $\pm$ s.e.m.) weight on choice of each sample's LLR, and its interaction with change point probability (*surprise*) and uncertainty. *Asterisks along the bottom indicate samples where weighting significantly differed from zero ($p < 0.05$, two-tailed cluster permutation test).

2.2. Motor preparation signals track the ongoing state of a decision variable

We first investigated motor preparation signals. Beta power over the motor cortex progressively lateralized during stimulus viewing, indicating increasing motor preparation and commitment to a specific action plan as the trial unfolded (*Fig. 2A*). In trials where changes in the generative distribution occurred, the time at which the response-specific lateralisation became apparent varied with the time of the last change point (*Fig. 2B*).

To test the sensitivity of MBL signals to evidence- and decision-related quantities from the normative model fits, we segmented the motor beta lateralisation traces [Left-Right hemispheres] around each evidence sample s (see *Methods*) and conducted time-resolved regressions against a set of model-derived variables. In our first analysis (*Eq. 9*), we included the prior belief (Ψ_s), the evidence provided by the new sample (LLR), and its modulation by surprise (LLR*surprise) - terms that linearly sum to approximate the updated belief (Ψ_{s+1}) subject to the non-linear normative transformation as observed in our data (Murphy et al., 2021). Uncertainty modulations were not included in EEG analysis due to their lack of a reliable behavioural effect in our dataset. We constructed the regression models such that positive coefficients would indicate lateralisation for the response favoured by the normative model given the evidence. Based on previous work (with MEG rather than EEG) suggesting that MBL encodes the evolving belief during trials of our task (Murphy et al. 2021), we expected a specific pattern of effects across the three regressors: sustained prior (Ψ_s) coefficients across the entire sample epoch, indicative of stable encoding of the belief before presentation of the new sample; and a sample-evoked increase in the coefficients for each of the individual sample-related terms (LLR and its modulation by surprise) which, as described above, would linearly sum with prior encoding to reflect the updated, normatively transformed

belief after accumulating the new sample (Murphy et al., 2021). By contrast, a signal encoding the *objective* sensory evidence carried by each sample would display a reliable effect only for the LLR term in this regression model; whereas a signal encoding only the *effective* evidence for the current sample (that is, the magnitude of the belief update but not the overall belief) would show reliable effects for LLR and its modulations, but not the prior term.

Our analysis showed that MBL followed the pattern expected of a signal encoding the evolving belief. It showed reliable encoding of the model-estimated prior belief *(Ψ)* that was sustained across the entirety of the analysed sample epoch (*Fig 2C,D*; $p < 0.001$). Further, MBL showed an evoked lateralisation following each sample, the magnitude of which scaled with each sample's LLR ($p < 0.001$) and was further modulated by how surprising it was ($p = 0.004$).

To maintain consistency with our analyses of the CPP signal below (see *Methods*), we also ran a simplified regression including only prior belief ($Ψ_s$) and effective evidence ($|\Delta Ψ| = Ψ_{s+1} - Ψ_s$) as regressors (*Eq. 10*) – with the latter term capturing the full extent of sample-by-sample normative belief update as compared to its approximation by LLR with surprise modulation described above. This model similarly yielded both the sustained prior ($Ψ_s$) and post-sample belief update ($|\Delta Ψ|$) encoding of an approximately normative DV (all $p < 0.001$; *Fig. 2E*), and provided a comparable fit to the data (paired t-test of adjusted R squared values $t_{(19)} = 0.83$; $p = 0.41$). To confirm that MBL tracked the normative DV across samples better than it tracked momentary stimulus information, we compared the adjusted R^2 values of regressions that each included a single regressor: a pure decision variable (DV, $Ψ_{s+1}$), the effective evidence (EE, $\Delta Ψ$) or objective evidence (OE, LLR). MBL was indeed best described by a regression encoding the state of an evolving belief (*Fig 2F*).

In sum, our results showed that motor beta lateralisation approximated a normative DV, with a sustained pattern of activity reflecting integrated information across samples as well as sample-evoked responses that are indicative of a normative belief updating process. These evoked lateralisation activities resulted from a decrease in power over the hemisphere contralateral to the response supported by the evidence, and an increase in power over the ipsilateral one (*Fig S2*).

Figure 2. Beta power lateralisation over the motor cortex encodes the evolving decision variable. **A.** Grand-averaged (±s.e.m.) beta lateralisation over the motor cortex [left-right hemispheres] aligned to trial start, sorted by response. Topography illustrates [Left-Right] hand responses at the end of the trial [4.2-4.4 s]. **B.** Grand-averaged (±s.e.m.) relative motor preparatory activity measured as the difference in MBL [Left-Right hemispheres] between left- and right-hand responses for trials containing a single change-point (CP) at different latencies. Positive values indicate increasing motor preparation towards the final response. Overlaid orange traces indicate the estimated normative prior (Ψ, scaled for visualisation) for the same set of trials, and illustrate that MBL closely mirrored the evolution of a normative DV over trial time. For illustration purposes, only correct trials are included**. C.** Motor beta lateralisation aligned to sample onset and averaged over samples 2-10 (*left*), and split in quartiles according to the belief (Ψ) at sample onset (*right*). **D.** Standardised regression coefficients of the motor

lateralisation [Left - Right] activity resulting from the regression including prior, LLR, and the modulation of LLR encoding by surprise (*Eq. 9*), averaged over samples 2-10 and aligned to sample onset. *Asterisks indicate significant cluster periods (two-tailed cluster-based permutation test, p < 0.05). **E.** Standardised regression coefficients of the motor lateralisation activity resulting from the regression with only prior and effective evidence factors (*Eq. 10*), averaged over samples 2-10 and aligned to sample onset. *Asterisks indicate significant cluster periods (two-tailed cluster-based permutation test, p < 0.05). **F.** Single-regressor model comparison. Traces indicate the difference in adjusted R^2 values for regressions including only the updated normative decision variable (DV, Ψ_{s+1}) or only effective evidence (EE, $\Delta\Psi$) as regressors, minus a regression including only absolute objective evidence (OE, LLR)(see *Methods*). The adjusted R^2 of the OE regression is used as a reference, and appears as a flat line with value 0 in the plot. The normative DV regression explained the greatest proportion of MBL variance. *Asterisks along the bottom indicate significant cluster periods for the normative DV model compared to the objective evidence (OE) one (two-tailed cluster-based permutation test, p < 0.05). Bar graphs indicate Adj. R^2 values for each regression (mean ± s.e.m.), across the whole sample epoch.

2.3. Centroparietal potentials reflect normative belief updates

We next similarly tested for distinct slow-building (across samples) and transient (sample-evoked) components of evidence processing present in the CPP. We observed that the centroparietal ERP was predominantly characterised by transient positive-going responses to each individual evidence sample rather than protracted build-up over the duration of the trial (*Fig. 3A*). Although some channels did show a positive drift over the whole trial, these positive deflections were not consistent across neighbouring electrodes in comparison with the transient sample-evoked responses. In order to examine transient and slow-building signal components systematically, we conducted time-resolved regressions of ERP amplitude against relevant decision-related quantities. Following the same logic as for motor beta lateralization, in our first analysis (*Eq. 11*) we included the model-estimated prior belief ($|\Psi|$), the evidence provided by each sample ($|LLR|$), and its modulation by surprise ($|LLR|*$surprise). In this case, we took the absolute (i.e. unsigned) values of the prior and LLR to account for the fact that the CPP is positive-going in response to sensory evidence regardless of the decision alternative that it favours. To allow for slow-building effects over trial time, we baseline-corrected the data for this analysis prior to trial onset and employed a high-pass filter with a low cutoff frequency (0.1 Hz). If a slow-building positivity encodes the overall DV across samples, we would expect to find a sustained, positively weighted sensitivity to the absolute prior belief $|\Psi|$, similar to the encoding of the signed prior observed in motor beta lateralisation (see *Fig. 2*).

We found that centroparietal responses to each sample scaled with its |LLR| (p = 0.007, two-tailed cluster permutation), and were further modulated by how surprising the sample was (p = 0.017, two-tailed cluster permutation, *Fig. 3B; Eq. 11*). Conversely, we found no evidence for encoding of the unsigned prior (no positive clusters identified). Given that some channels showed a positive drift over trial time, we repeated the analysis on all 128 scalp electrodes without averaging over channels, but no significant clusters encoding the unsigned prior were found anywhere on the scalp (p > 0.05; two-tailed cluster permutation test; *Fig. 3B*).

The modulation by surprise indicates that for a given level of |LLR|, the sample-evoked CPP is enhanced for more surprising samples and relatively reduced for unsurprising ones. This suggests encoding not of the *objective* evidence carried by each sample per se (i.e., its |LLR|), but rather the *effective* evidence that determines the belief update evoked by each sample after, in the normative model, it is accumulated and the belief passed through the normative non-linearity (see *Eq. 12:* $|\Delta\Psi| = |\Psi_{s+1} - \Psi_s|$). Supporting this interpretation, a simpler regression including only the magnitude of the belief update $|\Delta\Psi|$ revealed significant encoding of this variable over the same group of centroparietal electrodes (p = 0.015; *Fig. 3C*).

Finally, to confirm that a model where centroparietal responses encode effective evidence provides a better fit relative to other candidates, we compared the adjusted R-squared values of this last regression (*effective* evidence, $|\Delta\Psi|$) to those of a regression with |LLR| as the only regressor (*objective* evidence) and those of a regression with the updated decision variable ($|\Delta\Psi|_{s+1}$). The effective evidence regression yielded significantly better fits for a centroparietal cluster of electrodes consistent with the topographic peak of the CPP (*Fig. 3D*). Comparing subsets of trials with equally high values of |LLR| but that differed in effective evidence, illustrates that the response to high $|\Delta\Psi|$ was greater than to low $|\Delta\Psi|$ (*Fig, 3E*), showing how the CPP's encoding of effective evidence is truly distinct from objective evidence (|LLR|). Note that even when the required belief update is close to 0 (lowest $|\Delta\Psi|$ quartile) it is expected that the CPP still shows an appreciable positive deflection: whereas noise in the underlying belief update computation may take either a positive or negative sign and would thus average out to 0 across trials, the CPP reflects the absolute value of this belief update quantity, and so the average still results in a positive-going potential. Supplementary analyses showed that the sensitivity of the CPP to effective evidence observed here cannot be accounted for by collinearity between this quantity and various forms of surprise (*Fig. S4*).

Figure 3. Centroparietal signals reflect normative belief updates. **A.** Grand-averaged ERP traces aligned to trial onset, with baseline-correction by the period preceding the trial onset, for the highlighted electrodes (mean ± s.e.m.). **B,C.** Standardised regression coefficients for a regression including $|\Psi|$, $|LLR|$ and the $|LLR|$ modulation by surprise (*Eq. 11*, **B**) or a simplified version including only effective evidence ($|\Delta\Psi|$) as a regressor (*Eq. 12*, **C**), averaged over samples 2-10 and aligned to sample onset (mean ± s.e.m.). Evoked responses in a pre-selected group of centroparietal electrodes tracked normative belief updates, but not prior beliefs. *Asterisks indicate significant cluster periods (two-tailed permutation cluster test, $p < 0.05$). Topographies show whole-scalp regression coefficients [0.25-0.3s] post-sample, averaged over samples 2-10. **D.** Centroparietal activity is best explained by an effective evidence model. Traces indicate the difference in adjusted R^2 values for regressions including only the updated normative decision variable (DV, $|\Psi|_{s+1}$) or only effective evidence (EE, $|\Delta\Psi|$) as regressors, minus a regression including only absolute objective evidence (OE, $|LLR|$)(see *Methods*). As in Fig. 2F, the adjusted R^2 of the OE regression is used as a reference, and appears as a flat line with value 0 in the plot. The effective evidence regression explained a greater proportion of EEG variance in a cluster of posterior electrodes. *Asterisks along the bottom indicate significant cluster periods for the effective evidence (EE) model compared to the objective evidence (OE) one (two-tailed cluster-based permutation test, $p < 0.05$). Topographies show average difference in the regressions' adjusted R^2 [Adj. $R^2(|\Delta\Psi|)$ - Adj. $R^2(|LLR|)$] within the time window indicated by the shaded area, averaged over samples 2-10. Bar graphs indicate Adj. R^2 values for each regression (mean ± s.e.m.). **E.** Standard (not regression-derived) grand-averaged ERP traces aligned to trial onset restricted to samples with $|LLR| > 4.75$ (coloured areas in the top right inset), and split into quartiles according to the magnitude of the belief

update (effective evidence) they evoked. Data are high-pass filtered (>1 Hz) and baselined at sample onset for illustration purposes. Samples appearing in the same absolute angular location (i.e. same objective evidence or |LLR|) evoked higher potentials when they resulted in a bigger normative belief update. Bar graphs (*right*) show mean (± s.e.m.) $|\Delta\Psi|$ (*top*) and $|LLR|$ (*bottom*) of each quartile.

2.4. Linking centroparietal potentials, motor beta lateralisation and behaviour

We next sought to directly link fluctuations in our neural signals of interest to one another and to variability in choice behaviour. To do so, we exploited the trial-by-trial variability in the neural signals. First, we extracted the residual fluctuations in EEG activity following a given sample that could not be explained by computational variables derived from the normative model, and investigated how these related to choice (schematic in *Fig. 4A*). In a separate analysis, we aimed to link the two neural signals at a single-trial level by testing whether trials where, for example, a particularly high centroparietal response was observed for a given combination of model variables, were accompanied by a particularly large belief update at the motor preparation level (*Fig. 4E*). In order to increase our signal to noise ratio for these single-trial analyses, we applied an additional high-pass filter (>1Hz) to the EEG data prior to re-running the regressions in *Eq. 12* (following previous work, see e.g. Wyart et al. 2015), and extracted the residuals from a subset of anterior & posterior electrodes where effects were found to be maximal (see *Methods & Fig. S3*).

Trial-by-trial variations in centroparietal responses modulated the weight that each sample had on choice ($p = 0.004$; two-tailed cluster-based permutation test; *Fig. 4B, left*). That is, samples that evoked larger-than-expected centroparietal responses for a given level of effective evidence (i.e. trials with positive residuals from *Eq. 12*) also had stronger weight on choices. No differences were observed between results restricted to the anterior or posterior clusters ($p > 0.05$; two-tailed cluster-based permutation test; *Fig 4B, right*), respectively, nor between early and late samples in the trial ($p > 0.05$; two-tailed cluster permutation early vs. late). In turn, residual fluctuations in motor preparatory activity occurring late in the trial influenced the overall choice probabilities (*Fig. 4D*, right; two-tailed cluster-based permutation test for early [2-7] vs. late samples [8-10], $p=0.001$; the effect was not significant ($p = 0.08$) when averaging across all samples; *Fig. 4D*, left).

Finally, we aimed to directly link variability in centroparietal encoding of belief updates to the evolving representation of the belief in motor beta lateralisation. We found that residual EEG

fluctuations in centroparietal channels significantly modulated the sample-evoked change in the motor preparation signals. Namely, trials where the evoked centroparietal responses were larger than expected given the computational variables from the normative model fits (i.e. trials with positive residuals from *Eq. 12*) were also accompanied by greater-than-expected changes in lateralisation at the motor preparation level ($p = 0.038$; *Fig. 4F, left*). This effect was not significantly different between early and late samples (*Fig 4F, right;* two-tailed cluster-based permutation test for early [1-7] vs. late samples [8-10]), nor did it differ between the two groups of electrodes (*Fig. S5*).

Note that the time-course of the modulatory effect of centroparietal residuals on evidence weighting (*Fig. 4B*), as well as its effect on motor lateralisation updates (*Fig. 4F*) matched that of the neural encoding of belief updates over centroparietal channels (see *Fig. 2*). The temporal specificity of these effects indicates that only residual fluctuations specifically in the encoding of belief updates in these signals have an impact on choice and motor preparation.

Figure 4. Single-trial fluctuations link centroparietal signals, motor lateralisation and choice. **A,B.** Single trial fluctuations in centroparietal evoked responses modulate sample weight on choice. **A.** Schematic of the residuals analysis linking centroparietal signals to weight on choice. Black lines represent hypothetical EEG traces & evidence weights on choice predicted by the normative model, and coloured lines represent positive (*green*) or negative (*red*) residuals in Eq. 12 (*left*) and their predicted effect on evidence weight modulation (*right*). **B.** *Left:* regression coefficients (mean ± s.e.m.) of Eq. 13, showing that trials where CPP potentials were higher than predicted by the normative model (i.e. positive residuals from Eq. 12) were related to increased evidence weights. CPP residuals were extracted from the highlighted electrodes. *Right:* the same analysis using only the anterior and only the posterior electrodes clusters. **C,D.** Single-trial fluctuations in motor lateralisation influence choice probabilities at the end of trial. **C.** Schematic of the residuals analysis linking motor lateralisation signals to choice probabilities. Black lines represent the beta lateralisation & choice probabilities predicted by the normative model at any point in time (here taken as a baseline), and coloured lines represent positive (*green*) or negative (*red*) residuals in Eq. 9 (*left*). These indicate lateralisation favouring right or left responses over and above any lateralisation predicted by the normative model, and their predicted effect on evidence weight modulation (*right*). **D.** Normalised regression coefficients (mean ± s.e.m.) of Eq. 14, showing that motor lateralisation residuals influence choice probabilities. The effect observed across all fitted samples [2-10] (*left*) was driven by residuals at the end of the trial (*right*). Trials where motor lateralisation

favoured right-hand preparation more than expected by the normative model were related to higher than expected p(right) choice probabilities. **E-F.** Centroparietal signals influence updates to the decision variable encoded in motor beta lateralisation. **E.** Schematic of the residuals analysis linking centroparietal signals to changes in beta lateralisation following each sample, and to final choice probabilities. The black lines represent hypothetical EEG traces/motor beta lateralisation, and coloured lines indicate positive (*green*) or negative (*red*) residuals in Eq. 12/9, *left/right* respectively). **F.** Standardised regression coefficients (mean ± s.e.m.) of Eq. 17, showing that variability in centroparietal responses influences motor beta lateralisation updates following each sample across all samples (*left*), and for early and late sets of samples (*right*). Motor beta lateralisation updates (ΔMBL) were computed as the difference in beta lateralisation during the initial 100ms in consecutive samples. The impact of centroparietal fluctuations on motor beta lateralisation did not systematically vary over trial time. *Asterisks in all panels indicate significant cluster periods in all panels. Where present, shaded areas indicate cluster test period; in all other panels the full 1s epoch was included.

3. Discussion

Making perceptual decisions about external stimuli often requires the integration of multiple pieces of information. While several brain signals have been linked to decision variables (DV) exhibiting integration dynamics during a contiguous evaluation of a stationary sensory feature, it is important to know whether and how functional roles dissociate in more complex tasks. In this study, our goal was to directly test which specific aspects of a belief updating process are encoded by two prominent human EEG signatures of evidence accumulation - the CPP and MBL - in a volatile expanded judgement task. Specifically, we investigated whether these signals tracked an evolving belief in a sustained manner across successive discrete evidence samples, or whether they reflected momentary sample information or the belief updates derived therefrom. We used a task that required participants to integrate evidence in a non-linear manner, and to transform stimulus information into an intermediate decision-relevant quantity. Further, the discrete, temporally separate evidence samples we used allowed us to directly map decisional quantities to evoked neural responses. We applied a recent formulation of a normative model that can account for how agents flexibly tune their accumulation strategies in environments with various degrees of volatility (Glaze et al., 2015; Murphy et al., 2021), and which allowed us to precisely quantify how a DV evolved over trial time, including the precise belief updates (i.e. changes in the DV) required after each sample in our task. This normative model provided a close fit to participants' behaviour, as expected from previous work (Murphy et al. 2021), and thus we could use model-derived quantities to test which aspects of a decision might be encoded by the CPP and MBL. We found that 1) MBL consistently reflects the state of an evolving normative DV, 2) the CPP encodes normative belief updates but not the overall state of the DV, and finally 3) single-trial fluctuations in the CPP can account for otherwise unexplained deviations from the normative DV in MBL signals, providing a direct link between the processes indexed by these two signals.

Our first finding replicates previous MEG findings (Murphy et al. 2021) that MBL approximates the normative DV (i.e. the strength of beliefs) on our volatile decision-making task, with a sustained pattern of activity that reflects across-sample integration as well as discrete sample-evoked responses reflecting a normative belief update. The magnitude and sign of MBL at any given time during the trial reflected the DV, and updates were implemented by modulating power in both hemispheres, according to the sign of the belief update (i.e. simultaneous ipsilateral increase and contralateral decrease of power, as a function of the sign of the DV; see *Fig. S2*). Next, we addressed the main question of how the CPP participates in

this task; specifically, whether slow, sustained components of the CPP encode a DV, and how fast centroparietal responses evoked by each sample relate to the information contained in that sample. In contrast to MBL, we found no sustained encoding of the DV across samples, but rather a transient encoding of the normative belief update derived from each sample.

Previous work comparing the dynamics and functional properties of the CPP with motor preparation activity had employed simple tasks requiring a rapid, contiguous evaluation of a stationary, elemental stimulus feature (e.g. motion coherence or stimulus contrast), where momentary evidence that is readily encoded in sensory cortex directly translates into appropriate increments in an accumulating decision process. In those contexts, it was shown that the CPP exhibits many key characteristics of the accumulation process tracing the overall DV: it builds positively from a stimulus-aligned onset, at a rate proportional to evidence strength, up to a peak at the time of decision commitment, and its amplitude at that time varies with choice accuracy (O'Connell et al., 2012; O'Connell & Kelly, 2021; Steinemann et al., 2018). Apart from additional task-dependent biases and urgency components found to be added at the motor level in some contexts, the evidence-dependent dynamics exhibited in the CPP were mirrored in downstream motor preparation, at first blush suggesting a degree of computational redundancy. Our current study demonstrates that the signals' computational roles can diversify in tasks with discontinuous evidence samples requiring more computational steps to transform stimulus information into decision-relevant quantities. This gives new insight into the flexible cognitive capabilities afforded by the two-level architecture for decision formation reflected in the CPP and motor preparation signatures. Specifically, our results suggest that in task contexts where evidence comes in discrete samples, an intermediate process computes appropriate belief updates, which are then accumulated downstream at the motor level. In the advancing area of decision research in the past 3 decades, a commonly noted hallmark of neural decision variable correlates that has distinguished them from sensory encoding of the evidence itself is their "freedom from immediacy," whereby they keep a running total of cumulative evidence for as long as it takes to arrive at an ultimate commitment. In this sense, the CPP in the context investigated here is revealed to occupy an interesting middle ground, not concerned with basic sensory processing but handling intermediate cognitive computations applied to sensory information in order to feed a more slowly evolving tally downstream.

Previous work had suggested that sample-evoked centroparietal signals can be better explained by momentary rather than cumulative evidence in expanded judgement tasks (Wyart et al., 2012, 2015). However, the possibility that trial-long dynamics could encode an evolving DV, which is predicted from the CPP account emerging from classical studies in stationary environments, had not been tested. More specifically, signal processing choices in previous work (i.e. the application of a 1Hz high-pass filter or decorrelation of the signal between consecutive samples in Wyart et al. 2012, 2015) would have removed any sustained, across-sample activity. Thus, our use of a low high-pass filter cutoff (0.1Hz) for the main analysis (*Fig. 3*) was critical for this aim in order to allow potentially sustained activity to evolve over the long trials in our task. Using a regression analysis analogous to the one we used for MBL, we found that the amplitude of centroparietal signals throughout trial time did not reflect evolving beliefs. The lack of significant $|\Psi|$ encoding indicates that although some channels exhibited a growing positivity over time (see *Fig 3A*), this increase did not correlate with changes in a normative DV. That is, it was not the case that a stronger positivity was linked to stronger beliefs.

Our second research question related to how fast, sample-evoked centroparietal potentials encode momentary evidence. Centroparietal signals have previously been shown to scale with momentary evidence in expanded judgement, token-based tasks (Wyart et al., 2012, 2015) in static environments where the optimal strategy is to perfectly accumulate the objective evidence in each token. The volatility in our task allowed us to go a step further and test whether centroparietal signals evoked by samples reflect *objective* evidence (here quantified as each sample's $|LLR|$), or rather *effective* evidence – that is, the belief update each piece of evidence imposes on a decision variable ($|\Delta\Psi| = |\Psi_{s+1} - \Psi_s|$). In static contexts, the normative accumulation strategy prescribes that *objective* and *effective* evidence be identical: if evidence is accumulated perfectly, the step-change in a decision variable should be identical to the strength of the incoming objective evidence. In contrast, the non-linear transformation required by the normative model (Glaze et al., 2015) in volatile contexts entails that changes in a decision variable depend not only on the objective evidence provided by each sample, but also on its (in)consistency with the existing belief and the uncertainty under which it is encountered (Murphy et al., 2021). Our results show that centroparietal signals are driven by a sample's effective evidence after taking these factors into account according to a normative integration strategy (see *Fig. 3E*). This suggests centroparietal signals do not merely reflect objective sensory information, but rather an appropriate update quantity given the current belief.

It is worth noting that our results show strong encoding of belief updates in a broad group of electrodes, encompassing conventional CPP topographies but also extending to more anterior regions. However, the difference topographies comparing the two simple regressions assuming sole encoding of objective vs. effective evidence (*Fig. 3D, Fig S4*) show that activity in posterior regions typically associated with the CPP is best captured by an effective evidence model, whereas anterior regions seem to be almost equally well-explained by objective evidence (*Fig S4A*). Objective evidence in our task is inversely correlated with overall stimulus probability (i.e. samples with high |LLR| values are generally less likely to occur over the task), which has long been associated with the frontal ERP component known as the P3a, most often linked with the detection of novel, but not necessarily decision-relevant, information (Comerchero & Polich, 1998; Polich, 2007). Taken together, these results suggest that it is the posterior signal component that primarily handles the encoding of appropriate update quantities to be integrated in the decision process, but a frontal process may encode more general stimulus information content in parallel, encompassing the objective evidence related to stimulus location in the current task.

Finally, we sought to directly link centroparietal signals to motor beta lateralisation and behaviour, exploiting variability in single-trial fluctuations (Wyart et al., 2012, 2015). We show that samples which evoked higher-than-expected ERP responses also had a greater weight on choice, and that in turn motor fluctuations towards the end of the trial bias choice probabilities, in agreement with previous work (Wyart et al., 2012, 2015). Moreover, the fact that single-trial fluctuations in centroparietal responses also modulated motor lateralisation updates additionally suggests that 1) weight modulations are implemented by a direct impact of single-trial fluctuations in centroparietal signals on motor beta, and 2) that single-trial variability in motor lateralisation can be at least partially accounted for by single-trial variability in centroparietal signals.

Our study leaves some open questions. First, our findings suggest that the CPP reflects an intermediary step in expanded judgement tasks, in which stimulus information is transformed into appropriate belief update quantities and is then passed on downstream to the motor system. The CPP in such contexts thus reflects momentary belief updates evoked by single samples rather than a total DV integrated across samples. This integrated DV is instead encoded in the motor system. However, recent work has suggested that multiple brain areas may engage differently in decision formation depending on contextual factors such as knowledge about

stimulus-response mapping, decision policy or integration strategy (Okazawa & Kiani, 2023). In our task, the stimulus-response mapping was known and constant throughout the task, so participants could directly forward appropriately transformed stimulus information into response preparation areas. However, in a task where stimulus-response mapping is unknown, or a different integration strategy is prescribed, the involvement of the CPP generators in decision formation & maintenance may be different. Further research is required to better understand if and when the CPP and MBL interactions observed here generalise to other conditions.

Further, our finding that the CPP reflects the normatively required belief update means that the activity in its source regions must receive input beyond just signals carrying the newly arrived sensory evidence. However, there are several ways in which the non-linear transformation from objective into effective evidence could be implemented in the brain. Because belief updates in the normative model are determined by an interaction of incoming evidence with the state of the existing DV, one option is that the additional input to CPP generators (on top of objective sensory evidence) could take the form of feedback directly from areas representing the DV - in the case of the current task, putatively motor preparatory areas. In this case, local computations would need to take place to translate this combination of new evidence and current DV inputs into a signal coding for the effective belief updates. An alternative possibility, inspired by our decomposition of belief updating into modulations of sensory evidence weighting by surprise and (at least in some contexts) uncertainty (Murphy et al., 2021), is that the CPP generators receive inputs that relay internal estimates of these quantities, and these inputs in turn serve to modulate the representation of incoming evidence. Work in predictive inference tasks with strong similarities to our own points to candidate cortical networks for representing surprise and uncertainty quantities (McGuire et al., 2014; Nassar et al., 2019), while our own previous work with the current task also implicates pupil-linked arousal systems (Murphy et al., 2021). A final plausible option is that evolving beliefs may bias early sensory responses, directly affecting the primary input of the CPP generators without the need for any additional mechanism. Previous work on simple decision tasks has shown that beliefs and expectations can bias early sensory representations of evidence (Kok et al., 2013), or influence the accumulation process itself (Kelly et al., 2021). In our task, a sample's location on screen indicated how much evidence it provided. However, we do not have a way of measuring the strength of sensory representations in a way that is independent from the evidentiary quantity they are associated with. Thus, testing which of these or other

plausible candidate explanations for the implementation of effective evidence encoding will be a matter to be addressed in future work.

In sum, this study sheds new light on the neural correlates of normative belief updating. We show that centroparietal signals in this task context encode effective evidence provided by each sample – that is, the impact of new information on beliefs - rather than tracking the evolving DV across samples. Our results expand our understanding of which precise aspect of decision formation is tracked by the process reflected in the CPP, and further extends its validity as a marker of decision processing to volatile contexts, which require non-linear integration and to the best of our knowledge had not been examined before. This broadens our understanding of the nature of centroparietal decision signals, and paves the way for future work studying the link between the different neural signals involved in decision making.

4. Methods

4.1. Participants:

Twenty subjects aged 19-35 (M=23.1, SD=3.55; 15 male) each took part in two EEG sessions while performing a decision-making task (*Fig. 1A*). All procedures were approved by the human research ethics committee of University College Dublin and participants were compensated for their time.

4.2.Behavioural task

Stimuli were coded using Psychtoolbox-3 (Brainard, 1997) for Matlab (Mathworks). Participants monitored a set of checkerboard patches ('samples') appearing anywhere along a semicircular arc in the lower visual hemifield and centred on fixation. Sample locations were drawn from one of two overlapping Gaussian distributions, with equal s.d. and means left vs right of the vertical meridian (mean polar angle relative to vertical meridian: +/- 26°, s.d.: 29°). Samples were presented for 300ms, and a blank 100 ms interval separated consecutive stimuli. Participants viewed a maximum of 10 samples per trial, and trials had a 0.3 probability of terminating early, in which case the number of samples was drawn from a uniform distribution between 1-9. The distribution from which sample locations were drawn could change during the trial with fixed probability (hazard rate, H) of 0.1. Participants' task was to report what they believed to be the 'active' distribution at the end of each trial (left-or right-centered).

Stimuli were presented against a grey background. Three placeholders were present throughout each trial: a light-grey vertical line extending downward from fixation; a light grey half-ring in the lower visual hemifield (polar angle, from −90 to +90°); and a fixation mark which changed colours to indicate trial time (*blue:* intertrial period and initial fixation, *red*: view samples, *green*: response cue). The position of samples along the grey half-ring indicated their LLR: samples appearing close to the vertical meridian (0°) provided weak evidence (i.e. small |LLR|), as they could be drawn from either the left- or right- centred distribution with similar probabilities. Instead, samples appearing close to the horizontal midline (-90/+90°) provided strong evidence (i.e. large |LLR|), as they were more likely to be drawn from one of the two distributions than the other. All trials started with a checkerboard patch presented at a polar angle of 0° (i.e. on the vertical midline) for 300ms, serving as a warning cue, followed

by the first sample and a change in the colour of the fixation point (*red*) after a 100 ms blank interval. Evidence samples consisted of black and white flickering checkerboards (temporal frequency, 10Hz; spatial frequency, 2°) within a circular aperture and varied in polar angle. At the end of the trial, the fixation point changed to *green* ('Response cue') 0.7s to 1.3s (uniform distribution) after the final sample offset. Participants reported their response using their left- or right-hand thumb to press the left or right arrow keys, respectively, on a standard keyboard. There was no response deadline and no time pressure to respond. Auditory feedback 0.1s after the response informed participants of their accuracy. An ascending tone (350Hz to 950Hz) indicated correct responses, and a descending one (950Hz to 350Hz) errors. The next trial started after a 2s intertrial interval.

4.3. Behavioural analyses

Normative belief updating requires the following computation:

$$L_s = L_{s-1} + LLR_s \tag{1}$$

$$\Psi_s = L_{s-1} + log\left(\frac{1-H}{H} + exp\left(-L_{s-1}\right)\right) - log\left(\frac{1-H}{H} + exp\left(L_{s-1}\right)\right) \tag{2}$$

Here, L_s is the observer belief after encountering the evidence in sample *s*, expressed in log-posterior odds of the alternative task states; H is the hazard rate; and Ψ_s was the prior expectation of the observer before encountering a sample at location X_s, and LLR_s is the log-likelihood ratio:

$$LLR_s = log\left(\frac{p(X_s|Right)}{p(X_s|Left)}\right) \tag{3}$$

where $p(X_s|Right)$ and $p(X_s|Left)$ refer to the conditional probability of the token appearing at the observed location X for the current sample, given the generative distribution at that time is centred on the right and left, respectively.

We used this model to compute a metric of *surprise* and a metric of *uncertainty* (Murphy et al., 2021). Surprise was quantified as the analytically computed change-point probability after each sample. Note that we adopt the term "surprise" here in place of change-point probability (Murphy et al., 2021) to avoid ambiguity in acronym use; CPP, in this paper, refers to centroparietal positivity. The change-point probability indicates the probability that a

change of state has just occurred based on the most recent sample's likelihood under both generative states, the belief prior to that sample, and the generative *H:*

$$Surprise_s = \left(\frac{H * \left(\big(p(X_s|Left) * p(Left_{s-1}) \big) + \big(p(X_{s-1}|Right) * p(Right_{s-1}) \big) \right)}{\begin{array}{c} H * \left(\big(p(X_s|Left) * p(Left_{s-1}) \big) + \big(p(X_s|Right) * p(Right_{s-1}) \big) \right) \\ + (1-H) * \left(\big(p(X_s|Left) * p(Left_{s-1}) \big) + \big(p(X_s|Right) * p(Right_{s-1}) \big) \right) \end{array}} \right) \tag{4}$$

where p(Left) and p(Right) represent the probability that the generative state is left or right based on the prior belief and are computed as follows:

$$p(Left_{s-1}) = \frac{1}{e^{L_{s-1}} + 1} \tag{5}$$

$$p(Right_{s-1}) = \frac{e^{L_{s-1}}}{e^{L_{s-1}} + 1} \tag{6}$$

Surprise values were logit transformed in order to reduce skewness prior to introducing them in the behavioural and EEG regressions.

Uncertainty was defined as the negative absolute prior $(-|\Psi|)$. Higher values of uncertainty thus correspond to weak prior beliefs. Surprise and uncertainty are not explicitly used in the normative computations, but provide an intuitive understanding of the impact the non-linear transform has on belief updates and have been used in previous research (Murphy et al. 2021). It is worth noting that as environmental volatility (i.e. H) increases, the normative nonlinearity becomes steeper, producing faster transitions between strong beliefs and reducing the amount of time spent in high uncertainty states. A similar effect will also be achieved by increasing the signal-to-noise ratio (SNR; difference in means divided by s.d.) of the generative distributions. The generative volatility (H = 0.1) and SNR (1.79) were both higher than those in previous work (e.g. H = 0.08 and SNR = 1.17 in Murphy et al. 2021). Thus, given the lower proportion of samples expected to be received in high uncertainty states with the current generative settings, the modulation by this factor is expected to be weaker in our task.

Weight on choice:

We quantified the impact of evidence at each sample position on observed or model-derived choices, and the modulation of this impact by surprise, via logistic regression:

$$Logit(P(choice_{trl} = right)) \sim \beta_0 + \sum_{i=1}^{10} \beta_{1,i} \, LLR_{i,trl}$$

$$+ \sum_{j=2}^{10} \begin{pmatrix} \beta_{2,j} LLR_{j,trl} * Surprise_{j,trl} & + \\ \beta_{3,j} LLR_{j,trl} * Uncertainty_{j,trl} \end{pmatrix} \qquad (7)$$

where i and j index sample position within sequences of 10 samples, and LLR was the true LLR. We restricted this analysis to full trials (i.e. trials where the maximum possible number of samples (10) were presented).

Computational modelling:

We fit the normative model to subjects' choices, along with several alternative models that capture some, but not all, of its features: a perfect accumulation model, a perfect accumulation model with non-absorbing bounds, and a leaky accumulator. All models included noise and evidence gain (i.e. linear multiplicative scaling of LLR) parameters. The noise term was used to corrupt the log-posterior odds at the end of each trial, and choice probabilities in all models were computed as 0.5 + 0.5*erf(L/noise).

Additionally, subjective hazard rate, bound and leak parameters were estimated for the normative, non-absorbing bounds and leaky accumulator models, respectively (see Murphy et al. 2021 for full details). We fitted the model parameters by minimising the cross-entropy between participant and model choices. Error was minimised via particle swarm optimization (PSO), setting wide bounds on all parameters and running 300 pseudorandomly initialised particles for a maximum of 1500 search iterations.

The goodness of fit of the investigated models was assessed by computing Bayes Information Criterion (BIC):

$$BIC = 2e + klog(n) \qquad (8)$$

where k is the number of free parameters, n the number of trials and e the model's error as measured by the cross-entropy value.

4.4. EEG data acquisition and analysis:

Preprocessing: 128-channel scalp EEG data were recorded with a sampling rate of 512 Hz (Biosemi). Data were processed using Matlab R2021a (MathWorks) and the EEGlab toolbox (Delorme & Makeig, 2004). EEG data were high-pass filtered above 0.1 Hz (3rd order, two-pass Butterworth), and notch filtered at 50Hz and 100Hz to remove line noise. Data were epoched from [-0.4 s to 5.4 s] around trial start, and were baselined using the 100 ms prior to the trial onset cue. Eye movements were identified and removed using EEGlab's *runica* function, which applies the extended Infomax algorithm for Independent Component Analysis (ICA) (Bell & Sejnowski, 1995). Additionally, channels showing $\pm150\mu V$ deviations from baseline were interpolated on a trial-by-trial basis using the TBT toolbox (Ben-Shachar, 2018). If any given epoch had more than 20 bad channels, it was marked for rejection. Further, if any given channel was marked as artifactual on >30% of epochs, it was interpolated across the whole experiment. Data were downsampled to 200 Hz and finally, we applied a Current Source Density (CSD) filter using the CSD toolbox (Kayser & Tenke, 2006) to the preprocessed data to reduce the spread of potential components on the scalp, through computing a second spatial derivative.

Spectral analysis. We used a sliding-window Fourier transform to compute time–frequency decompositions of the single-trial activity. We used a Hanning taper (window length, 0.4 s; time steps, 0.05 s; frequency steps, 1Hz; frequency smoothing, $\pm2.5Hz$) for the range 1–35Hz using the FieldTrip toolbox (Oostenveld et al., 2011). To compute motor lateralisation, we averaged a cluster of 6 left- and 6 right-hemisphere electrodes based on the response-aligned topography differences between left- and right-response trials, and computed an index of motor lateralisation as the difference in beta power (13-30Hz) between the two hemispheres (left minus right, electrodes highlighted in figures). Time-frequency data were decibel (dB) transformed prior to analysis (10 x log10 (power/baseline)),consistent with Murphy et al. 2021, where baseline was the trial-averaged power in the 100 ms preceding trial onset).

EEG modelling analysis:

Following Murphy et al. 2021, we ran a time-resolved regression to investigate whether motor beta lateralisation encoded computational variables relevant for normative belief updating :

$$MBL_{s,t} \sim \beta_0 + \beta_1 \Psi_s + \beta_2 LLR_s + \beta_3 LLR_s * Surprise_s + \beta_4 session \qquad (9)$$

The modulation by surprise is predicted by the non-linear prior scaling in the normative model and was observed in the behavioural data, but they are not explicitly used in the computations. Thus, to directly test whether changes in motor lateralisation following each sample reflected the normatively required belief update, we ran a second simplified regression using only the prior belief (Ψ_s) and the belief update ($\Psi_{s+1} - \Psi_s$) as regressors:

$$MBL_{s,t} \sim \beta_0 + \beta_1 \Psi_s + \beta_2 (\Psi_{s+1} - \Psi_s) + \beta_3 session \tag{10}$$

Next, we ran an equivalent analysis for the ERP to test whether centroparietal activity reflected evidence-related variables relevant to normative belief updating. (*Eq. 11,12*). To reduce noise in the estimated coefficients, EEG signals were low-pass filtered at 6Hz (two-pass Butterworth filter) prior to the analysis:

$$EEG_{e,s,trl} \sim \beta_0 + \beta_1 |\Psi|_s + \beta_2 |LLR|_s + \beta_3 |LLR|_s * Surprise_s + \beta_4 session \tag{11}$$

$$EEG_{e,s,trl} \sim \beta_0 + \beta_1 |\Psi_{s+1} - \Psi_s| + \beta_2 session \tag{12}$$

In all regressions (*Eq. 9-12*), the analysis was restricted to full trials (i.e. trials where the maximum possible number of samples (10) were presented), and a session covariate taking values 0 or 1 for sessions 1 and 2 respectively was included to control for spurious differences between the two recording sessions. Analyses were run for each sample separately, for a period spanning 1s after sample onset, and finally coefficients were averaged across samples. Note that for broadband EEG regressions (*Eq.* 11-12) absolute values for the and LLR regressors are used. This is because previous research has shown that centroparietal potentials evoked by decision-relevant evidence are always positive, irrespective of the response alternative supported by the evidence (O'Connell et al., 2012), consistent with a scheme where scalp amplitude reflects the absolute value of what is internally encoded as a signed decision variable (Afacan-Seref et al., 2018). Further, the initial sample was excluded from analysis in Eq. 9 and Eq. 11 because prior (Ψ) values are always 0 at the beginning of the trial.

EEG model comparison:

To test the extent to which a specific computational quantity explained the EEG variability better than an alternative one, we compared the adjusted R squared values of simple regressions including only the computational variable of interest to a reference model. In the main text, we

compared models reflecting three possible aspects of a belief updating process that could be captured by the CPP and MBL: a normative decision variable (Ψ), momentary objective evidence (OE, quantified as the LLR), or effective evidence (EE, quantified as $\Delta\Psi$ and indicating the extent to which a belief is updated following any given sample. Absolute and signed quantities were used for the CPP and MBL analysis, respectively. In an additional supplementary analysis for the CPP (*Fig. S4*), we further compared these models to an intercept-only model and to a model containing only surprise to investigate whether sensitivity to surprise alone could account for the observed results. Only trials where the maximum possible number of samples (10) were presented were included in all analyses. Further, in order to increase our signal to noise ratio, and since the key comparisons of interest pertained to the sample-wise evoked potentials, all these simple regressions were run after additionally high-pass filtering the preprocessed data above 1Hz (3rd order, two-pass Butterworth; see next section).

Residuals analysis:

Finally, we harnessed the single-trial variability of EEG signals to investigate whether intrinsic fluctuations in neural activity over and above the effects captured by the normative model influenced behaviour. We used the residuals from Eq. 9 and Eq. 12 as measures of trial-by-trial fluctuations not accounted for by the normative model.

For the analysis of residual variability in centroparietal signals, we tested whether samples evoking particularly big positivities (i.e. positive residuals) were given more weight on choices by including the residuals of Eq. 12 as an interaction term with LLR on a regression predicting choice (Eq. 13):

$$
Logit(P(choice_{trl} = right)) \sim \beta_0 + \sum_{i=1}^{10} \left(\begin{matrix} \beta_{1,i} LLR_{i,trl} + \\ \beta_{2,i} LLR_{i,trl} * Residual\ CPP_{i,trl} \end{matrix} \right) +
$$

$$
+ \sum_{j=2}^{10} \left(\begin{matrix} \beta_{3,j} LLR_{j,trl} * Surprise_{j,trl} + \\ \beta_{4,j} LLR_{j,trl} * Uncertainty_{j,trl} \end{matrix} \right) \tag{13}
$$

where *i* and *j* index sample position within sequences of 10 samples. Only trials where the maximum possible number of samples (10) were presented were included in this analysis. Note that, in order to increase our signal to noise ratio for this single-trial analysis of centroparietal signal residuals, we repeated the analysis in Eq. 12 after additionally high-pass filtering the

preprocessed data above 1Hz (3rd order, two-pass Butterworth). This method has been used for similar residuals analysis in the past (Wyart et al., 2015), and it overcomes two problems. First, it avoids the need to baseline-correct data prior to each individual sample within a trial, which might be problematic given the timings of our stimuli and overlapping potentials. Second, since our analysis involves averaging over samples occurring at various times after trial onset (ranging between 0 and 4.4s), removing slow drifts in the data, which are not of interest in this analysis, makes activity following early and late samples more comparable and increases signal to noise ratio. Importantly, the effects of interest did not change qualitatively or quantitatively when repeating the regression analysis on this additionally filtered dataset (see *Fig. S3*). Residuals were extracted from the analysis reported in *Fig S3B*, from a subset of anterior and posterior channels centred around the midline where effects were found to be maximal.

Regarding the impact of fluctuations in beta lateralisation on choices, we investigated whether variations in lateralisation following any given sample modulated choice probabilities by including the residuals from Eq. 9 as a main effect (Eq. 14).

$$
\begin{aligned}
Logit(P(choice_{trl} = right)) \sim \beta_0 \; + \; & \sum_{i=1}^{10} \beta_{1,i} \, LLR_{i,trl} \\
+ \sum_{j=2}^{10} & \begin{pmatrix} \beta_{2,j} LLR_{j,trl} * Surprise_{j,trl} \;\; + \\ \beta_{3,j} LLR_{j,trl} * Uncertainty_{j,trl} + \\ \beta_{4,j} Residual\ lateralisation_{j,trl} \end{pmatrix}
\end{aligned}
\tag{14}
$$

Finally, we aimed to directly link fluctuations in centroparietal activity to changes in motor beta lateralisation following each sample. Motor beta lateralisation [left-right] was averaged over the first 100ms following each sample, and we computed a measure of motor updates (ΔMBL) indicating how much beta lateralisation changed between consecutive samples (*Eq. 15*), as well as a measure of the normatively required update, defined as the change in normative priors between consecutive samples (*Eq. 16*).

$$
\Delta MBL_s \; = MBL_{s+1} - MBL_s
\tag{15}
$$

$$
\Delta \Psi_s \; = \Psi_{s+1} - \Psi_s
\tag{16}
$$

Then, we tested whether variability in motor updates could be partially explained by fluctuations in centroparietal signals. In other words, whether centroparietal fluctuations

modulated the extent to which motor updates deviated from the normatively required change. To test this, we fitted the following regression (*Eq. 17*):

$$\Delta MBL_s = \beta_0 + \beta_1 \Delta \Psi_s + \beta_2 \Delta \Psi_s * Residual\ CPP_{s,trl} \tag{17}$$

Cluster statistics:

EEG regression results were statistically evaluated using cluster-based permutations using the FieldTrip toolbox (Maris & Oostenveld, 2007; Oostenveld et al., 2011). All tests were two-tailed, cluster-corrected ($p<0.05$), 10000 permutations were run.. For analysis in *Eq. 11* and *Eq. 12*, regression coefficients were averaged over a cluster of preselected centroparietal electrodes prior to testing. The preselected electrodes encompassed areas where the centroparietal positivity is typically observed in evidence accumulation tasks.

References

Afacan-Seref, K., Steinemann, N. A., Blangero, A., & Kelly, S. P. (2018). Dynamic Interplay of Value and Sensory Information in High-Speed Decision Making. *Current Biology*, *28*(5), 795-802.e6. https://doi.org/10.1016/J.CUB.2018.01.071

Balsdon, T., Mamassian, P., & Wyart, V. (2021). Separable neural signatures of confidence during perceptual decisions. *ELife*, *10*. https://doi.org/10.7554/eLife.68491

Ben-Shachar, M. S. (2018). TBT: Reject and Interpolate channels on a trial-by-trial basis. Zenodo. https://doi.org/10.5281/zenodo.1241518

Brainard, D. H. (1997). The Psychophysics Toolbox. *Spatial Vision*, *10*(4), 433–436. https://doi.org/10.1163/156856897X00357

Comerchero, M. D., & Polich, J. (1998). P3a, perceptual distinctiveness, and stimulus modality. *Cognitive Brain Research*, *7*(1), 41–48. https://doi.org/10.1016/S0926-6410(98)00009-3

de Lafuente, V., Jazayeri, M., & Shadlen, M. N. (2015). Representation of accumulating evidence for a decision in two parietal areas. *Journal of Neuroscience*, *35*(10), 4306–4318. https://doi.org/10.1523/JNEUROSCI.2451-14.2015

de Lange, F. P., Rahnev, D. A., Donner, T. H., & Lau, H. (2013). Prestimulus oscillatory activity over motor cortex reflects perceptual expectations. *Journal of Neuroscience*, *33*(4), 1400–1410. https://doi.org/10.1523/JNEUROSCI.1094-12.2013

Delorme, A., & Makeig, S. (2004). EEGLAB: An open source toolbox for analysis of single-trial EEG dynamics including independent component analysis. *Journal of Neuroscience Methods*, *134*(1), 9–21. https://doi.org/10.1016/j.jneumeth.2003.10.009

Donner, T. H., Siegel, M., Fries, P., & Engel, A. K. (2009). Buildup of Choice-Predictive Activity in Human Motor Cortex during Perceptual Decision Making. *Current Biology*. https://doi.org/10.1016/j.cub.2009.07.066

Geuzebroek, A. C., Craddock, H., O'Connell, R. G., & Kelly, S. P. (2023). Balancing true and false detection of intermittent sensory targets by adjusting the inputs to the evidence accumulation process. *ELife*, *12*. https://doi.org/10.7554/ELIFE.83025

Gherman, S., Markowitz, N., Tostaeva, G., Espinal, E., Mehta, A. D., O'Connell, R. G., Kelly, S. P., & Bickel, S. (2024). Intracranial electroencephalography reveals effector-independent evidence accumulation dynamics in multiple human brain regions. *Nature Human Behaviour 2024 8:4*, *8*(4), 758–770. https://doi.org/10.1038/s41562-024-01824-9

Glaze, C. M., Kable, J. W., & Gold, J. I. (2015). Normative evidence accumulation in unpredictable environments. *ELife*, *4*(08825). https://doi.org/10.7554/eLife.08825.001

Gold, J. I., & Shadlen, M. N. (2007). The neural basis of decision making. *Annual Review of Neuroscience*, *30*, 535–574. https://doi.org/10.1146/annurev.neuro.29.051605.113038

Hanks, T. D., Kopec, C. D., Brunton, B. W., Duan, C. A., Erlich, J. C., & Brody, C. D. (2015). Distinct relationships of parietal and prefrontal cortices to evidence accumulation. *Nature 2015 520:7546*, *520*(7546), 220–223. https://doi.org/10.1038/nature14066

Hanks, T. D., & Summerfield, C. (2017). Perceptual Decision Making in Rodents, Monkeys, and Humans. *Neuron*, *93*(1), 15–31. https://doi.org/10.1016/J.NEURON.2016.12.003

Hyafil, A., De La Rocha, J., Pericas, C., Katz, L. N., Huk, A. C., Pillow, J. W., & Salinas, E. (2023). Temporal integration is a robust feature of perceptual decisions. *ELife*. https://doi.org/10.7554/eLife.84045

Kayser, J., & Tenke, C. E. (2006). Principal components analysis of Laplacian waveforms as a generic method for identifying ERP generator patterns: I. Evaluation with auditory oddball tasks. *Clinical Neurophysiology*, *117*(2), 348–368. https://doi.org/10.1016/j.clinph.2005.08.034

Kelly, S. P., Corbett, E. A., & O'Connell, R. G. (2021). Neurocomputational mechanisms of prior-informed perceptual decision-making in humans. *Nature Human Behaviour*, *5*(4), 467–481. https://doi.org/10.1038/s41562-020-00967-9

Kelly, S. P., & O'Connell, R. G. (2013). Internal and External Influences on the Rate of Sensory Evidence Accumulation in the Human Brain. *Journal of Neuroscience*, *33*(50), 19434–19441. https://doi.org/10.1523/jneurosci.3355-13.2013

Kok, P., Brouwer, G. J., van Gerven, M. A. J., & de Lange, F. P. (2013). Prior Expectations Bias Sensory Representations in Visual Cortex. *Journal of Neuroscience*, *33*(41), 16275–16284. https://doi.org/10.1523/JNEUROSCI.0742-13.2013

Maris, E., & Oostenveld, R. (2007). Nonparametric statistical testing of EEG- and MEG-data. *Journal of Neuroscience Methods*, *164*(1), 177–190. https://doi.org/10.1016/j.jneumeth.2007.03.024

Mars, R. B., Debener, S., Gladwin, T. E., Harrison, L. M., Haggard, P., Rothwell, J. C., & Bestmann, S. (2008). Trial-by-Trial Fluctuations in the Event-Related Electroencephalogram Reflect Dynamic Changes in the Degree of Surprise. *Journal of Neuroscience*, *28*(47), 12539–12545. https://doi.org/10.1523/JNEUROSCI.2925-08.2008

McGuire, J. T., Nassar, M. R., Gold, J. I., & Kable, J. W. (2014). Functionally Dissociable Influences on Learning Rate in a Dynamic Environment. *Neuron*, *84*(4), 870–881. https://doi.org/10.1016/J.NEURON.2014.10.013

Murphy, P. R., Wilming, N., Hernandez-Bocanegra, D. C., Prat-Ortega, G., & Donner, T. H. (2021). Adaptive circuit dynamics across human cortex during evidence accumulation in changing environments. *Nature Neuroscience*, *24*(7), 987–997. https://doi.org/10.1038/s41593-021-00839-z

Nassar, M. R., Bruckner, R., & Frank, M. J. (2019). Statistical context dictates the relationship between feedback-related EEG signals and learning. *ELife*, *8*. https://doi.org/10.7554/ELIFE.46975

O'Connell, R. G., Dockree, P. M., & Kelly, S. P. (2012). A supramodal accumulation-to-bound signal that determines perceptual decisions in humans. *Nature Neuroscience*, *15*(12), 1729–1735. https://doi.org/10.1038/nn.3248

O'Connell, R. G., & Kelly, S. P. (2021). *Neurophysiology of Human Perceptual Decision-Making*. https://doi.org/10.1146/annurev-neuro-092019

Okazawa, G., & Kiani, R. (2023). *Annual Review of Physiology Neural Mechanisms That Make Perceptual Decisions Flexible*. https://doi.org/10.1146/annurev-physiol-031722

Oostenveld, R., Fries, P., Maris, E., & Schoffelen, J. M. (2011). FieldTrip: Open source software for advanced analysis of MEG, EEG, and invasive electrophysiological data. *Computational Intelligence and Neuroscience*. https://doi.org/10.1155/2011/156869

Pfurtscheller, G., & Lopes da Silva, F. H. (1999). Event-related EEG/MEG synchronization and desynchronization: basic principles. *Clinical Neurophysiology*, *110*(11), 1842–1857. https://doi.org/10.1016/S1388-2457(99)00141-8

Polich, J. (2007). Updating P300: An integrative theory of P3a and P3b. *Clinical Neurophysiology*, *118*(10), 2128–2148. https://doi.org/10.1016/J.CLINPH.2007.04.019

Shadlen, M. N., & Kiani, R. (2013). Decision making as a window on cognition. In *Neuron* (Vol. 80, Issue 3, pp. 791–806). https://doi.org/10.1016/j.neuron.2013.10.047

Stancák, A., & Pfurtscheller, G. (1995). Desynchronization and recovery of β rhythms during brisk and slow self-paced finger movements in man. *Neuroscience Letters*, *196*(1–2), 21–24. https://doi.org/10.1016/0304-3940(95)11827-J

Steinemann, N. A., O'Connell, R. G., & Kelly, S. P. (2018). Decisions are expedited through multiple neural adjustments spanning the sensorimotor hierarchy. *Nature Communications*, *9*(1), 3627. https://doi.org/10.1038/s41467-018-06117-0

Twomey, D. M., Kelly, S. P., & O'Connell, R. G. (2016). Abstract and effector-selective decision signals exhibit qualitatively distinct dynamics before delayed perceptual reports. *Journal of Neuroscience*, *36*(28), 7346–7352. https://doi.org/10.1523/JNEUROSCI.4162-15.2016

Wyart, V., de Gardelle, V., Scholl, J., & Summerfield, C. (2012). Rhythmic Fluctuations in Evidence Accumulation during Decision Making in the Human Brain. *Neuron*, *76*(4), 847–858. https://doi.org/10.1016/j.neuron.2012.09.015

Wyart, V., Myers, N. E., & Summerfield, C. (2015). Neural mechanisms of human perceptual choice under focused and divided attention. *Journal of Neuroscience*, *35*(8), 3485–3498. https://doi.org/10.1523/JNEUROSCI.3276-14.2015

Supplementary material

Figure S1. **The normative model provides the best fit to the data. A.** Proportion of observed and model-predicted correct responses. White bars and overlaid coloured points represent individual data and fits, respectively. Filled bars display grand-averaged data ± s.e.m.. **B.** Goodness of fit (BIC) of the three best fitting models, relative to the normative one. Overlaid dots indicate individual data. The normative model produced the best quantitative fit to the data (all p < 0.05, Bonferroni corrected, BIC paired t-tests). **B.** Observed and model-predicted (mean ± s.e.m.) weight on choice of each sample's LLR, and its interaction with change point probability (*surprise*) and uncertainty. *Asterisks along the bottom indicate samples where weighting significantly differed from zero (p < 0.05, two-tailed cluster permutation test). Several models can account for some of the normative model's behavioural implications. A leaky accumulator captured the increasing weight on choice of later samples but not how each sample's weight is modulated by surprise or belief uncertainty. A perfect accumulator with non-absorbing bounds, which effectively operates as a piecewise-linear approximation to the normative transfer function, was able to recapitulate all of the same key qualitative features of the data as the normative model. **D.** Estimated normative model parameters (hazard rate (*H*, *left*), evidence gain (*middle*) and noise (*right*)) for each individual participant (coloured *dots*), ordered as a function of H estimate, and at the group level(*bar graph;* mean ± s.e.m.). The average estimate of the hazard rate (H) did not significantly differ from the generative statistics (H = 0.1; *p* = 0.59; one-sample t-test).

Figure S2. Standardized regression coefficients resulting from the analyses in *Eq. 9*, aligned to sample onset. Analyses were run on all electrodes separately, and regression coefficients were then averaged over the left- and right-cluster of electrodes over the motor cortex. Prior encoding and belief updates were implemented via modulation of both hemispheres. Regressions fitted separately for each sample, and then averaged over samples 2-10.

Figure S3. Reanalysis of data reported in Figure 3 after applying 1Hz high-pass filter for residuals analysis. **A.** Grand-averaged ERP traces aligned to trial start in, in the highlighted electrodes (mean ± s.e.m.). Note the absence of positive drift in anterior channels (c.f. *Fig. 3A*). **B.** Standardised regression coefficients resulting from the analyses in *Eq. 12* averaged over samples 2-10 and aligned to sample onset (mean ± s.e.m.). Evoked responses in a pre-selected cluster of centroparietal electrodes tracked normative belief updates. *Asterisks indicate significant cluster periods. Topographies show whole-scalp cluster statistics [0.25-0.3s] post-sample, averaged over samples 2-10. Key effects of interest remain significant and consistent with the analysis reported in the main manuscript. The solid line shows results for the same selection of centroparietal electrodes used throughout the manuscript. To increase signal to noise ratio for the single-trial analysis, we extracted the residuals from a subset of electrodes where effective evidence encoding was maximal (*dashed line*).

Figure S4. The effective evidence model ($|\Delta\Psi|$) provides a better fit to EEG data than other models. Other work has suggested that CPP-like signals specifically encode forms of surprise in learning and predictive inference contexts (Mars et al., 2008; Nassar et al., 2019). However, we observed that the effective evidence model provided a better fit to the EEG evoked potentials than alternative models with two different forms of surprise as a the sole regressors: |LLR| (***B***), which corresponds to low-level form of surprise on our task given the fact that high |LLR| stimuli are generally the least probable to occur; and change-point probability (referred to as 'surprise' above), which captures a high-level form of surprise conditional on the decision-maker's existing belief about the generative state (***C***). ***A.*** Adjusted R square values of single-regressor models, compared to an intercept-only model, for posterior (*left*) and anterior (*left*) electrode selections where effects were maximal. The effective evidence model provided a better fit in both anterior & posterior electrodes. ***B.*** Adjusted R squared values of the effective evidence model compared to the objective evidence one. The effective evidence model provided better fits in a posterior cluster of electrodes. ***C.*** Adjusted R squared values of the effective evidence model compared to a surprise-only model. Topographies in each inset show the whole-scalp distribution of the difference between the effective evidence model and the

reference model in each graph (***A,*** intercept; ***B,*** objective evidence; ***C,*** surprise), at the time indicated by the shaded area. Note: all models included a binary "session" regressor to control for any spurious differences between the two recording sessions.*Asterisks represent significant cluster periods (two-tailed cluster permutation test against 0, [0-1s]).Topographies illustrate the difference between the effective evidence model and the reference one in each panel, during the times indicated by the shaded areas.

Figure S5. Normalised regression coefficients (mean ± s.e.m.) of *Eq. 15*, showing that variability in centroparietal responses influences motor beta lateralisation updates following each sample across all samples (***A***), in two electrode clusters (***B***), and sorted by sample times (***C***).