Unexpected events modulate context signaling in VIP and excitatory cells of the visual cortex

Farzaneh Najafi[1,4 #], Simone Russo[3,5,6], Jérôme Lecoq[1,2 #]

[1] Allen Institute, Mindscope Program, [2] Neural Dynamics Program, [3] Brain and Consciousness Program, Seattle,

WA, USA

[4] Georgia Institute of Technology, Biological Sciences, [5] Wallace H Coulter Department of Biomedical

Engineering, Atlanta, GA, USA

[6] Department of Biomedical and Clinical Sciences, Università degli Studi di Milano, Milan 20157, Italy

Corresponding authors:
Jerome Lecoq: jeromel@alleninstitute.org
Farzaneh Najafi: fnajafi3@gatech.edu

ABSTRACT

The visual cortex predicts incoming sensory stimuli through internal models of the world. Unexpected stimuli

that violate these predictions update internal models and drive adaptation. Cortical inhibitory neurons,

particularly VIP (vasoactive intestinal peptide) interneurons, are suggested to play a key role in representing

unexpected stimuli, given their robust firing following unexpected omissions of familiar images. Importantly,

this response is stimulus non-specific, raising an important question about what information it conveys. Given

their unique connectivity with other cell types and brain areas, we hypothesized that during unexpected events,

VIP neurons encode contextual information, defined as neuronal activity that is not driven by the stimulus

itself. To test this hypothesis, we analyzed the Allen Institute Visual Behavior dataset, in which mice viewed

repeated familiar images and unexpected omissions of these images while two-photon calcium imaging data

from different cell types and visual areas were recorded. Using dimensionality reduction techniques, we found

that omissions trigger contextual signaling in VIP neurons in the primary visual cortex (V1) and the lateral

medial (LM) visual area, particularly in the superficial layers. Similarly, contextual coding was enhanced in

excitatory neurons following omissions. This contrasted sharply with the excitatory response to expected

images, during which contextual information was substantially suppressed. Our results suggest that unexpected

events activate VIP neurons, which subsequently propagate contextual information across the cortical network.

This potentially facilitates the integration of context within the cortical network, and leads to updated

predictions about our dynamic environment.

INTRODUCTION

The visual cortex is crucial in shaping our internal visual representations of the sensory environment. In the

mouse visual cortex, inhibitory neurons are diverse, with major classes including parvalbumin (PV), vasoactive

intestinal polypeptide (VIP), and somatostatin (SST) neurons, each with distinct subclasses as evidenced by

transcriptomic characterization[1]. VIP cells, presumed to exert a dis-inhibitory effect on pyramidal cells, may

play a key role in regulating information flow within the cortical network due to their connectivity with

associative cortical areas and neuromodulatory systems[2–5]. VIP neurons exist in superficial (layer 2/3

primarily), and deeper (layer 4) layers of the cortex, suggesting their involvement in both top-down and

bottom-up information processing, respectively[6].

Circuit models suggest that unexpected stimuli trigger computations across both excitatory and inhibitory

cortical cells to refine sensory representations[7]. Empirical data indicate that unexpected stimuli elicit responses

across multiple layers of the cortex, involving inhibitory neurons[8]. A recent study demonstrated that the activity

of VIP cells in the visual cortex peaks during unexpected omissions of visual stimuli[9], but the underlying nature

of these responses remains unclear. VIP cells may signal the presence of a prediction error signal; alternatively,

they may encode contextual changes that co-occur during omissions, including behavioral, external, or internal

states.

In the present study, we addressed this question using multi-area two-photon calcium imaging during a visually

guided behavior. This task contained both expected events (regular flashes of images) and unexpected events

(image omissions). We leveraged population activity decoding, dimensionality reduction and trial shuffling to

examine whether VIP activity during omissions may contain stimulus-specific prediction-error signals, or if it

was primarily influenced by context information. Importantly, "context" in our study broadly refers to shared

activity across brain areas that are independent of task-relevant features. This may involve behavioral features

such as the animal's locomotion, external inputs such as the sensory signals in the animal's environment, and

internal representations related to the animal's current state.

We found that most VIP neurons, except those in layer 4 of the primary visual area, encoded context

information. This suggests that VIP neurons represent a general 'surprise' signal enriched with contextual

details during unexpected events. In contrast, excitatory neurons displayed weaker modulation by context

during expected image presentations but were significantly influenced by context during unexpected

omissions. These results highlight the role of superficial VIP neurons in updating cortical activity with

contextual information in response to unexpected stimuli. This mechanism potentially facilitates the

association of contextual signals with unexpected events, thereby driving adaptive responses.

RESULTS

Distinct cortical pathways are thought to carry sensory and prediction signals[10,11], suggesting that cortical areas

and cell types might show unique responses to expected vs. unexpected events. To test this hypothesis, we

leveraged two-photon imaging data from the Allen Brain Observatory platform[9,12]. In this dataset, mice were

trained to perform a go/no-go, image-change detection task[9]. In brief, water-restricted animals were presented

with a constant stream of natural images (250 ms) interleaved with gray screens of matched luminance

(500 ms). When a change in image identity occurred, mice would receive a water reward if they licked

immediately after the image change. Importantly, during imaging sessions in well-trained mice, a pseudo-

random 5% of non-change images were replaced with a gray screen (image omissions; Fig. 1A). Since mice

had extensive experience with the highly regular timing of image presentations (every ~750 ms), the rare

stimulus omissions were unexpected events, as opposed to the expected, frequent, and repeated images. During

this task, the activity of excitatory neurons as well as inhibitory SST and VIP neurons were recorded separately,

using 3 distinct mouse lines (see Methods). Neuronal cells within each line were simultaneously imaged at 4

cortical depths in 2 visual areas (V1 and higher visual area: LM; Fig. 1B,C; Supplementary Fig. 1).

We first studied how excitatory and inhibitory subtypes (VIP, SST) respond to image presentations and

omissions, and how their responses depend on cortical area and depth. The population average of neural

activities demonstrated a clear difference across cell classes (Fig. 1D,E). Excitatory neurons and, more

robustly, SST neurons in all areas and layers were activated by expected images, i.e. the expected events.

Conversely, excitatory neurons, overall, did not respond to omissions, except for those in the middle depth of

LM. SST neuronal activity was slightly decreased after omissions in all recorded locations (Fig. 1D,E top,

middle). In sharp contrast to excitatory and SST neurons, VIP neurons in all layers and areas were robustly

activated after omissions (Fig. 1D,E bottom). Specifically, VIP cells demonstrated small ramping activity

approximately 250 ms prior to each image presentation that further increased when the image was not

presented (i.e. omission), and was immediately inhibited by image presentation, confirming previous results

using the same behavioral task[9].

This multi-area dataset allowed comparison of neuronal responses across cortical depths of V1 and LM in each

experiment (Fig. 1D,E). We found that excitatory neuronal responses to images were stronger in deeper layers

of both visual areas, particularly V1. SST responses to images were strongest at middle depths of V1,

corresponding to layer 2/3 and superficial layer 4, but did not differ among cortical layers of LM. VIP responses

to neither images nor omissions were significantly different across cortical layers or areas, as reported

previously[6], although the responses were slightly stronger in deeper layers. Therefore, in both V1 and LM,

images are more robustly represented by excitatory and SST neurons, particularly in deeper layers; while

unexpected omissions are represented by VIP neurons (Fig. 1D,E; Stats: Supplementary Table 1).

It is suggested that feedback pathways, from higher order cortical areas to superficial layers of V1, convey

prediction signals[10,11,13]. In light of these studies, we used simultaneous multi-plane recording to investigate if

cortical interactions might be different when expectations are violated, i.e. during omissions. As a proxy for

cortical interactions, we measured pairwise noise correlations across repeated image presentations, as well as

omissions, for each cell type. Correlations were computed within each area (V1-V1, LM-LM; Supplementary

Fig. 2-4), as well as across areas (V1-LM; Fig. 2). We also performed a shuffling analysis to control for firing

rate impacts on correlations (Methods; Supplementary Fig. 2-4)

In the excitatory network, V1-V1 and V1-LM coactivations increased after image presentations, but only in

deep layers (V1-LM: Fig. 2; V1-V1: Supplementary Fig. 2A,D). Interestingly, excitatory network coactivation

in LM-LM did not change after images, across any pair of cortical layers (LM-LM: Supplementary Fig. 2B,E).

Following omissions, there was no significant change in the coactivation of excitatory neurons across visual

layers and areas (Fig. 2; Supplementary Fig. 2A,D).

In the SST network, images increased coactivation more broadly across visual layers and areas, particularly in

our "middle" imaging depth, i.e., deep layer 2/3 and superficial layer 4 (Fig. 2; Supplementary Fig. 3).

Omissions, on the other hand, slightly decreased coactivation of neurons relative to the baseline across most

layers of the SST network (Fig. 2; Supplementary Fig. 3).

In the VIP network, coactivation patterns were strikingly different compared to other cell types: neurons in

superficial layers of V1 and LM were strongly coactivated following omissions. There was also a weak

coactivation prior to images in most layers and areas (Fig. 2; Supplementary Fig. 4). Notably, correlations were

overall much weaker among excitatory neurons compared to inhibitory neurons (Fig. 2), confirming previous

results and indicating stronger local connectivity between inhibitory neurons[14,15].

Our results suggest that the VIP network in the visual cortex is strongly recruited during unexpected events:

cells from all layers of V1 and LM are active during omissions and neural correlations are especially strong in

the superficial layers. One possible explanation is that prediction errors (i.e. mismatch between the animal's

expectation of a given image and its omission) generate a shared input signal that co-activates VIP cells in

superficial layers across areas, thereby synchronizing the local cortical network. An alternative hypothesis is

that VIP activity during omissions may be due to co-occurring changes triggered by unexpected events. Such

changes could include variations in animal's locomotion and arousal, as well as alterations in the representation

of environmental cues, or internal states. This hypothesis is corroborated by recent reports[16,17], including our

own[18], that have demonstrated a broad range of behavioral features modulate brain-wide network dynamics,

raising the possibility that our correlation findings could be contributed by global brain fluctuations driven by

widespread behavioral events. Importantly, in our previous work [9], we had found that the VIP population

carries no information about the identity of the previous image. These results demonstrated that VIP neurons

do *not* encode stimulus-specific prediction error signals; instead suggesting that VIP omission activity may

represent a broad non-specific surprise signal.

To address whether VIP correlated activity during omissions may be affected by global contextual changes,

we performed Tensor Component Analysis (TCA)[19] on real and trial-shuffled data, aligned on images and

omissions (number of sessions = 22, 15, 21 for excitatory, SST, and VIP cells, respectively). TCA allowed

decomposing neural population activity into distinct components, and trial shuffling allowed studying

contextual representation (see below). Each TCA component was characterized by weights indicating the

unique contributions from individual neurons, trials, and time points (Fig. 3A,B). Importantly, we defined TCA

components as image-specific or omission-specific if their trial weights were significantly higher on images

or omissions, respectively (ranksum test across image and omission weights; $p < 0.05$; Fig. 3E). Components

in which trial weights were not significantly different between images and omissions were excluded from

further analyses (number of non-specific components, pooled across sessions; mean +/- SD across 100 TCA

iterations; Excitatory: 38 +/- 4; SST: 20 +/- 3 ; VIP: 55 +/- 3).

To investigate the neural coding of images and omissions across visual areas and layers, we compared the

absolute magnitude of TCA neuronal weights among V1, LM, superficial and deep layers (imaging depth

smaller or bigger than 250μm, respectively; Fig 3C,D, red; for the details of TCA stats see Table S2). In

omission-specific components, VIP neurons had stronger weights in *superficial* layers of both V1 and LM.

This contrasted with excitatory and SST neurons which had stronger weights in deeper layers of V1 and LM

(Fig. 3C, red). In image-specific components, excitatory neurons in deeper layers of V1 and LM had stronger

weights; while SST neurons showed stronger weights in V1 compared to LM, independent of cortical layer

(Fig. 3D red). VIP neurons had few image-specific components, hence excluded from our analysis of neuronal

weights.

TCA neural weights for image- and omission-specific components reflect the coding strength of individual

neurons on image and omission trials, respectively. This neural coding could arise from the direct presentation

of images and omissions, or could be influenced by co-occurring behavioral and sensory features in adjacent

or remote areas, which we collectively term contextual features. To distinguish the neuronal coding of images

and omissions from these contextual influences, we performed TCA on trial shuffled data, for which trial

identities were shuffled *within* the image and omission conditions for each neuron (see Methods). This

shuffling procedure allowed us to preserve image- and omission-specific responses while removing any

contextual information that co-occurred across neurons during images and omissions. In contrast, in the non-

shuffle data, TCA components could encode the presence of images and omissions as well as any context-

dependent information present in the neuronal population.

Shuffling led to a general decrease of neuronal weight magnitudes for omission-specific components in all cell

types (Fig. 3C blue: shuffled; red: real data). This reduction was most prominent in VIP neurons in the

superficial layers, and absent from the VIP neurons in deep V1 (Fig. 3C, right). These findings suggest that

omissions are co-represented with contextual information in the visual cortex in excitatory, SST, and

particularly superficial VIP neurons. The major exception is that VIP neurons in the deeper layers of V1 show

a smaller contribution from contextual signals (Wicoxon ranksum test; p=0.009).

Interestingly, shuffling had a smaller effect on the neuronal weights of image-specific components in excitatory

and SST cells (Fig. 3D blue vs. red; Note the absence of VIP neurons from this analysis due to their few

numbers of image components). This finding suggests that during images, excitatory neurons primarily encode

image information, overriding contextual signals. This is aligned with previous studies that demonstrated

reduced neuronal variability during sensory inputs[20].

Indeed, all cell types, and particularly VIP neurons, had more TCA components that were omission-specific

vs. image-specific (Fig. 3E, black: images, gray: omissions). This finding suggests that omission responses in

our neuronal population exhibit more variability than image responses and is aligned with our TCA shuffled

analysis which demonstrated response variability during omissions was associated with the co-representation

of contextual features.

DISCUSSION

Recent work had demonstrated that the VIP population is functionally quite diverse, and strongly represents

behavioral features and unexpected image omissions[9]. Extending these findings, our current study

demonstrates that during unexpected omissions, VIP neurons in the visual cortex not only encode a non-

specific surprise signal but also a substantial amount of contextual information. This dual representation of

omissions and context is predominantly observed in superficial VIP neurons, with notably reduced context

coding in deeper layers.

Multiple lines of work have suggested the role of VIP neurons in encoding external and internal features. VIP

neurons can represent behavioral state, leading to gain modulation in sensory cortices[4,21]. VIP activity increases

with pupil dilation[22], locomotion[4,23], and whisking [3,24], leading to enhanced sensory modulation during high

attention and arousal states that follow surprising and reinforcement signals[4,21–23,25,26]. VIP neurons are driven

by neuromodulators, particularly cholinergic inputs[4], as well as long-range cortical and thalamic inputs[2,3]. VIP

neurons have also been shown to play a key role to surround suppression[27], when neuronal responses to

centered visual stimuli can be decreased by the presence of adjacent stimuli.

Our study suggests that the VIP network in primary and higher visual cortices comprises two subnetworks with

distinct roles, associated with superficial and deeper layers. In V1, the subnetwork in the deeper layers encodes

information about omissions, independently of context, while the subnetwork in the superficial layers

concurrently encodes both context and omission information. Importantly, the omission coding by VIP cells is

non-specific, lacking any information about which stimulus was omitted, as suggested by our recent work

(Garrett 2023). This supports the hypothesis that VIP neurons broadly encode surprise signals rather than

specific mismatches between expected and actual stimuli. Our findings suggest a potential specialization within

the VIP network that enhances the reliable representation of sensory features in deeper V1, while integrating

sensory and contextual features in the superficial layers of V1, hence facilitating adaptive responses to dynamic

environmental stimuli.

The ability of VIP cells to disinhibit excitatory neurons through its connectivity pattern, is ideally suited to

facilitate the propagation of selective activity patterns through the cortex, leading to attention towards

contextual sensory inputs[26] when needed, or the emergence of motor behaviors[28,29]. We propose a framework

in which VIP cells serve as gatekeepers of contextual information, which are dynamically activated when

transitioning from expected to unexpected stimuli (Fig. 4). This framework is in agreement with the numerous

studies on the activity of the inhibitory network. Future experiments should aim to simultaneously record from

all those cell types in the context of dynamic but controlled sensory uncertainty.

Figure 1. Excitatory and inhibitory cell types distinctly encode expected images and unexpected omissions across visual areas and layers. A) Allen Institute Visual Behavior dataset: calcium signals were simultaneously imaged across two cortical columns, including 4 depths (approximately 100 μm, 200 μm, 300 μm, and 375 μm), in visual areas V1 and LM, and in 3 distinct mouse lines tagged for excitatory, and 2 inhibitory cell types: VIP and SST. **B)** Mice, walking on a disk, view a familiar set of images, separated by a gray screen. Images are unexpectedly omitted on 5% of the presentation. **C)** Example population activity of excitatory neurons, recorded simultaneously from ~1180 neurons in V1 and LM. **D)** Population-averaged calcium responses to images (shaded rectangle) and omissions (dashed rectangle), across 2 visual areas (V1, LM), and 4 cortical depths (blue, cyan, green, red), in 3 different mouse lines (A-C: excitatory, SST, VIP). ΔF/F traces normalized to baseline standard deviation. Mean +/- SEM; n = 24, 22, 24 sessions for excitatory, SST, and VIP, respectively. **E)** Quantification of evoked responses averaged over 350ms after images, and 500ms after omissions (time-windows used for quantification of image and omission-evoked responses are indicated by orange and gray horizontal lines above top left panel).

Figure 2. Distinct cortical interactions occur across primary and higher-order visual areas following expected and unexpected events. A) Spearman correlation coefficients between V1 and LM neurons at different cortical depths (colors indicate LM depths; each subplot corresponds to a given V1 depth). B) Change in correlation coefficients during images (left) and omissions (right) relative to baseline correlation coefficient. Correlation coefficients quantified over 500ms after images, and 750ms after omissions. Traces and heatmaps: mean +/- SEM; n = 8, 6, 9 mice for excitatory, SST, and VIP cell types, respectively.

Figure 3. Unexpected events lead to enhanced context representation in excitatory and particularly VIP neurons in superficial layers. A) Schematic of the analysis pipeline. Imaging data were decomposed using Tensor Component Analysis (TCA; red: original; blue: shuffled). Shuffled data were obtained by permuting trials identities within each category (image and omission). Each TCA component was identified as image-

specific, omission-specific, or non-specific depending on the weight magnitude of the TCA trial component for images vs. omissions (Methods). **B)** Left to right: neuronal, temporal, and trials weights of a representative omission-specific TCA component (red: original; blue: shuffled). In this example, trials from 1 to 159 are images, while the remaining trials are omissions. TCA was performed on spikes extracted from DF/F traces aligned on images and omissions, including 300ms before to 700ms after images and omissions. **C)** Neuronal weight of the omission-specific components for excitatory, SST, and VIP neurons (left to right) extracted from original and shuffled data across areas (Wilcoxon ranksum test; all $p<0.001$ except for VIP V1 deep: $p=0.009$). **D)** Same as panel C for image-specific components for excitatory and SST neurons (left to right; Wilcoxon ranksum test; all $p<0.001$ except for excitatory V1 superficial: $p=0.8$ and excitatory LM superficial: $p=0.5$). **E)** Number of components pooled across sessions, for excitatory, SST, and VIP neurons (left to right) from original data compared between images and omissions (black and gray boxes, respectively; Wilcoxon ranksum test; all $p<0.001$).

Figure 4. Proposed mechanism: Triggered by unexpected events, VIP neurons gate contextual signal flow across cortical cell types and areas. Left) Expected stimuli, such as familiar images in our study, are encoded by excitatory and SST neurons. **Right)** When unexpected stimuli occur, a non-specific surprise signal activates VIP cells to promote contextual integration across all cell types. This contextual information may originate from nearby and distant brain areas, encoding a variety of signals, including internal (cognitive and motivational states), or external (sensory and behavioral) information. The presence of this contextual information facilitates the updating of the unexpected stimulus representation in the local network.

256 **ONLINE METHODS**

257 **Processing of calcium imaging movies**

258 The preprocessing of all calcium imaging data was done within the imaging processing pipeline described in

259 detail elsewhere[34] and crosstalk removal was applied after pre-processing. Briefly, all data was first corrected

260 for brain motion by performing rigid registration in two dimensions. Then, cell-segmentation was performed

261 to identify spatial masks of active neurons. Further, fluorescence from spatially-overlapping neuronal masks

262 was unmixed and corrected for neuropil contamination. Finally, mask-matching and crosstalk removal was

263 performed using FastICA[37] and ghost cells were filtered, and lastly, $\Delta F/F$ was computed on corrected masks.

264 **Description of datasets**

265 We used the publicly available Allen Institute Visual behavior dataset: https://portal.brain-map.org/circuits-

266 behavior/visual-behavior-2p. Details are explained in our previous work (Garrett 2023). In brief, this dataset

267 included 3 mouse lines: excitatory mouse line: Slc17a7-IRES2-Cre ; Camk2a-tTA ; Ai93(TITL-GCaMP6f),

268 inhibitory subpopulation VIP: Vip-IRES-Cre ; Ai148 (TIT2L-GC6f-ICL-tTA2), and inhibitory subpopulation

269 SST: Sst-IRES-Cre ; Ai148 (TIT2L-GC6f-ICL-tTA2). For the excitatory cell line, 24 sessions from 8 mice;

270 for the SST cell line, 22 sessions from 6 mice; for the VIP cell line, 24 sessions from 9 mice were recorded.

271 **Population averages of neuronal responses**

272 Traces were aligned to the onset of image omission; since images were presented regularly, omission alignment

273 also aligned the traces on images. The median of neural responses was computed across trials. The average

274 response was then computed across neurons, for each session. A grand average was then computed across

275 sessions (see Fig. 1D, traces). To quantify the image-evoked responses, the calcium trace of each neuron was

276 averaged over 350ms after image onset for excitatory and SST neurons, and over [-250, 100]ms relative to

277 image onset for VIP neurons to account for their anticipatory response. Response quantification was done on

278 the mean (across trials) trace of each neuron. The same quantification was performed for omission-evoked

279 responses, except a 500ms window was used for quantification. The responses were averaged across neurons,

280 for each session, and then a grand average was computed across sessions (see Fig. 1E; error-bars).

Correlation of neural responses across cortical planes

On omission-aligned traces, Spearman correlation coefficient was computed between pairs of neurons across

trials. Here, by trial we mean omissions. Correlations were computed for every individual frame, over [-1,

2] sec relative to the omission. This procedure was done for all pairs of neurons; then an average value was

computed across all pairs (Fig. 2). Neuron pairs were present within the same plane, or in 2 different planes.

This analysis allowed studying how the response of neurons (within the same plane or in different areas/layers)

covaried across trials, and how this coactivation changed at different moments (e.g. after images vs. omissions).

Spearman correlation coefficients were also computed on shuffled traces (Supplementary Fig. 2-4 d-f), which

were driven by independently shuffling trial orders for each neuron. For each neuron pair, shuffling was

repeated 50 times, resulting in a distribution of correlation coefficients for shuffled data. This shuffling analysis

controlled for firing rate impacts on our correlation results.

To compute "noise" correlations, we measured pairwise correlations of "signal"-removed traces. "Signal" was

computed by taking the average neural response to each image type (there were 8 images in each session), and

subtracting the average from the response of individual trials of that image type.

To quantify coactivation of neurons (Fig. 2, heatmaps; Supplementary Fig. 2-4d-f), correlations coefficients

("cc") were first averaged over 500 ms after images for excitatory and SST neurons, and over [-250, 250] ms

relative to image onset for VIP neurons, accounting for their anticipatory response. Then, correlation

coefficients were averaged across baseline frames. We call this quantity "image cc". To quantify omission-

evoked coactivation, we averaged correlation coefficients over 750 ms after omissions. We call this quantity

"omission cc". Next, we quantified baseline coactivation by averaging correlation coefficients across baselines

frames. Baseline was defined as the frame immediately preceding each image presentation, for excitatory and

SST neurons, and 250 ms earlier than each image presentation for VIP neurons. We call baseline quantification

of correlation "baseline cc". Finally, we measured the change in coactivation during images or omissions by

subtracting out "baseline cc" from "image cc", or "omission cc". This quantity is plotted in Fig. 2, heatmaps,

and Supplementary Fig. 2-4d-f, error-bars.

The Python package scipy (scipy.stats.spearmanr) was used for computing correlations, and p-values were

computed in 2 ways: 1) using the p-value output of the spearmanr package; 2) manually computing the p-value

by comparing the correlation coefficient of real (non-shuffled) data with the shuffled distribution using 2-sided,

1-sample t-test (using the Python package scipy.stats.ttest_1samp).

Tensor Component Analysis

Tensor Component Analysis (TCA) was performed on inferred spikes from the DF/F trace using the tensortool

package (https://github.com/neurostatslab/tensortools). TCA was set to extract 5 components from the trials

split from -300 to +600 msec relative to the image or omission onset. The resulting components were first

filtered according to their image- or omission-specificity. Specifically, we compared the weights of image and

omission trials for each component (Wilcoxon ranksum test, alpha <0.05) and identified as condition-specific

the components showing a significant difference between the two. On this subset of components, those with

larger absolute average weights during image trials were classified as image-specific, while those with larger

absolute average weights during omission trials were classified as omission-specific. To assess the variability

across trials, on each subset of components we computed the standard deviation across trial weights. This entire

procedure was performed 100 times to account for the variability between TCA decompositions. To obtain a

null distribution that discarded context information while preserving image- and omission-specific

information, we shuffled trials for each neuron keeping separated image and omission trials (e.g. the activity

of an omission trial was reassigned to another omission trial independently for each neuron). Similarly to the

original data, the procedure was performed 100 times to account for variability in shuffling and TCA

decomposition. While TCA explained a relatively small amount of the overall signal, especially in the case of

excitatory neurons, the components were highly reproducible across iterations (for TCA error and similarity

across iterations see Table S3). As an additional control, we performed the same analysis with a larger number

of reconstructed components by TCA and found similar results (Supplementary Fig. 5).

330 Statistical tests

331   We used two-way ANOVA, followed by Tukey HSD to compare population averages across cortical layers.

332   Two-sided t-test was used to compare correlations between real and shuffled data, for each cortical plane. A

333   p-value of 0.05 was used as the significance threshold. For comparison of the correlation coefficients, we used

334   two-tailed, two-sample Kolmogorov–Smirnov tests.

335 Data availability

336   Data are publicly available as Allen Institute Visual Behavior Dataset: https://portal.brain-map.org/circuits-

337   behavior/visual-behavior-2p

338 Code availability

339   Crosstalk removal was performed using custom routines employing FastICA which is available as part of

340   scikit-learn python package (https://scikit-learn.org/). Code to perform population average and correlation

341   analyses   are   available   at: https://github.com/farznaj/multiscope_fn.

342   Code   to   perform   TCA   and   the   related   statistical   tests   are   available   at:

343   https://github.com/SimoneRusso/Multiscope_TCA

345 Acknowledgements

We wish to thank the founder of the Allen Institute for Brain Science, Paul G. Allen, for his vision,

encouragement and support.

Competing interests

The dual-beam add-on module (J.L.) intellectual property has been licensed to Thorlabs. Inc., by the Allen

Institute. S.R. is the Chief Medical Officer of Manava Plus.

Correspondence and requests for materials should be addressed to J.L, and F.N..

References

References

1. Gouwens, N. W. *et al.* Integrated Morphoelectric and Transcriptomic Classification of Cortical

GABAergic Cells. *Cell* **183**, 935–953.e19 (2020).

2. Wall, N. R. *et al.* Brain-Wide Maps of Synaptic Input to Cortical Interneurons. *J. Neurosci.* **36**, 4000–

4009 (2016).

3. Lee, S., Kruglikov, I., Huang, Z. J., Fishell, G. & Rudy, B. A disinhibitory circuit mediates motor

integration in the somatosensory cortex. *Nat. Neurosci.* **16**, 1662–1670 (2013).

4. Fu, Y. *et al.* A cortical circuit for gain control by behavioral state. *Cell* **156**, 1139–1152 (2014).

5. Bastos, G. *et al.* Top-down input modulates visual context processing through an interneuron-specific

circuit. *Cell Rep.* **42**, 113133 (2023).

6. Garrett, M. *et al.* Experience shapes activity dynamics and stimulus coding of VIP inhibitory cells. *Elife*

9, (2020).

7. Hertäg, L. & Sprekeler, H. Learning prediction error neurons in a canonical interneuron circuit. *Elife* **9**,

(2020).

8. Hamm, J. P. & Yuste, R. Somatostatin Interneurons Control a Key Component of Mismatch Negativity

in Mouse Visual Cortex. *Cell Rep.* **16**, 597–604 (2016).

9. Garrett, M. *et al.* Stimulus novelty uncovers coding diversity in visual cortical circuits. *bioRxiv*

2023.02.14.528085 (2023) doi:10.1101/2023.02.14.528085.

10. Keller, G. B. & Mrsic-Flogel, T. D. Predictive Processing: A Canonical Cortical Computation. *Neuron*

100, 424–435 (2018).

11. Bastos, A. M. *et al.* Canonical microcircuits for predictive coding. *Neuron* **76**, 695–711 (2012).

12. de Vries, S. E. J. *et al.* A large-scale standardized physiological survey reveals functional organization

of the mouse visual cortex. *Nat. Neurosci.* **23**, 138–151 (2020).

13. Rao, R. P. & Ballard, D. H. Predictive coding in the visual cortex: a functional interpretation of some

extra-classical receptive-field effects. *Nat. Neurosci.* **2**, 79–87 (1999).

14. Najafi, F. *et al.* Excitatory and Inhibitory Subnetworks Are Equally Selective during Decision-Making

and Emerge Simultaneously during Learning. *Neuron* **105**, 165–179.e8 (2020).

15. Packer, A. M. & Yuste, R. Dense, unspecific connectivity of neocortical parvalbumin-positive

interneurons: a canonical microcircuit for inhibition? *J. Neurosci.* **31**, 13260–13271 (2011).

16. Steinmetz, N. A., Zatka-Haas, P., Carandini, M. & Harris, K. D. Distributed coding of choice, action and

engagement across the mouse brain. *Nature* **576**, 266–273 (2019).

17. Musall, S., Kaufman, M. T., Juavinett, A. L., Gluf, S. & Churchland, A. K. Single-trial neural dynamics

are dominated by richly varied movements. *Nat. Neurosci.* **22**, 1677–1686 (2019).

18. Ramadan, M. *et al.* A Standardized Nonvisual Behavioral Event Is Broadcasted Homogeneously across

Cortical Visual Areas without Modulating Visual Responses. *eNeuro* **9**, (2022).

19. Williams, A. H. *et al.* Unsupervised Discovery of Demixed, Low-Dimensional Neural Dynamics across

Multiple Timescales through Tensor Component Analysis. *Neuron* **98**, 1099–1115.e8 (2018).

20. Churchland, M. M. *et al.* Stimulus onset quenches neural variability: a widespread cortical phenomenon.

Nat. Neurosci. **13**, 369–378 (2010).

21. Pi, H.-J. *et al.* Cortical interneurons that specialize in disinhibitory control. *Nature* **503**, 521–524 (2013).

22. Reimer, J. *et al.* Pupil fluctuations track fast switching of cortical states during quiet wakefulness.

Neuron **84**, 355–362 (2014).

23. Pakan, J. M. *et al.* Behavioral-state modulation of inhibition is context-dependent and cell type specific

in mouse visual cortex. *Elife* **5**, (2016).

24. Yu, J., Hu, H., Agmon, A. & Svoboda, K. Recruitment of GABAergic Interneurons in the Barrel Cortex

during Active Tactile Behavior. *Neuron* **104**, 412–427.e4 (2019).

25. Letzkus, J. J. *et al.* A disinhibitory microcircuit for associative fear learning in the auditory cortex.

Nature **480**, 331–335 (2011).

26. Karnani, M. M. *et al.* Opening Holes in the Blanket of Inhibition: Localized Lateral Disinhibition by

VIP Interneurons. *J. Neurosci.* **36**, 3471–3480 (2016).

27. Keller, A. J. *et al.* A Disinhibitory Circuit for Contextual Modulation in Primary Visual Cortex. *Neuron*

108, 1181–1193.e8 (2020).

28. Arroyo, S., Barati, S., Kim, K., Aparicio, F. & Ganguly, K. Emergence of preparatory dynamics in VIP

interneurons during motor learning. *Cell Rep.* **42**, 112834 (2023).

29. Ren, C. *et al.* Global and subtype-specific modulation of cortical inhibitory neurons regulated by

acetylcholine during motor learning. *Neuron* **110**, 2334–2350.e8 (2022).

Supplementary material

Excitatory, V1 image responses	one-way ANOVA: F= 16.3, p<0.001; TUKEY HSD: p<0.01 for all pairwise comparisons except layer 1-2/3 and layer 1-4.
Excitatory, LM image responses	TUKEY HSD: p>0.05 for all pairwise comparisons, except for layer 2/3-5: p=0.001.
SST, V1 image responses	one-way ANOVA: F=12.5, p<0.001; TUKEY HSD: p<0.05 for layer1-2/3, layer 1-4, and layer 4-5 pairwise comparisons.
SST, LM image responses	one-way ANOVA: F= 1.9, p=0.14.
VIP, V1 image responses	one-way ANOVA: F=3.0, p=0.03; yet TUKEY HSD: p>0.05 for all pairwise comparisons.
VIP, LM image responses	one-way ANOVA: F=2.4, p=0.07. VIP, V1 omission responses: one-way ANOVA: F=3.0, p=0.03; yet TUKEY HSD: p>0.05 for all pairwise comparisons. VIP, LM omission responses: one-way ANOVA: F=1.6, p=0.19.

Supplementary Table 1. Statistical tests corresponding to Fig. 1.

SLC - images

ANOVA

statistic=974.9242425360012, pvalue=2.053417877366087e-182

Posthoc

Group1	Group2	p
0	1	<0.001
0	2	0.0045
0	3	<0.001
1	2	<0.001
1	3	0.17
2	3	<0.001

Ranksum

Group	p	effect size
1	0.5	-0.000559821
2	<0.001	0.005208667
3	0.8	-0.000677232
4	<0.001	-0.006407717

SLC - omissions

ANOVA

statistic=924.6397775278477, pvalue=2.0365188317079515e-178

Posthoc

Group1	Group2	p
0	1	<0.001
0	2	0.9
0	3	<0.001
1	2	<0.001
1	3	0.9
2	3	<0.001

Ranksum

Group	p	effect size
1	<0.001	-0.007043246
2	<0.001	-0.010607699
3	<0.001	-0.005858144
4	<0.001	-0.011777935

SST - images

ANOVA

statistic=562.5678946813928, pvalue=2.34542498531712e-142

Posthoc

Group1		p
0	1	0.26
0	2	<0.001
0	3	<0.001
1	2	<0.001
1	3	<0.001
2	3	<0.001

Ranksum

Group	p	effect size
1	<0.001	-0.020148597
2	<0.001	0.008925503
3	<0.001	-0.018708875
4	<0.001	-0.015717922

SST - omissions

ANOVA

statistic=388.69757476891175, pvalue=1.3433542133282506e-117

Posthoc

Group1		p
0	1	<0.001
0	2	<0.001
0	3	<0.001
1	2	<0.001
1	3	<0.001
2	3	<0.001

Ranksum

Group	p	effect size
1	<0.001	-0.021273879
2	<0.001	-0.032506687
3	<0.001	-0.010707941
4	<0.001	-0.061329306

VIP - omissions

ANOVA

statistic=152.09786442559385, pvalue=1.4251507357988139e-65

Posthoc

Group1		p
0	1	<0.001
0	2	<0.001
0	3	<0.001
1	2	<0.001
1	3	0.0058
2	3	<0.001

Ranksum

Group	p	effect size
1	<0.001	-0.054864618
2	<0.001	-0.017996513
3	<0.001	-0.068608776
4	0.009	-0.005409973

LEGEND

group 1	LM superficial
group 2	LM deep
group 3	V1 superficial
group 4	V1 deep

Supplementary Table 2. Statistical tables for TCA analysis.

Number of components	Celltype	Data	Error	Similarity
5	Excitatory	Original	0.9401	0.8879
		Shuffled	0.9681	0.8673
	SST	Original	0.7695	0.9227
		Shuffled	0.8365	0.9067
	VIP	Original	0.774	0.918
		Shuffled	0.8838	0.8886
10	Excitatory	Original	0.9084	0.8265
		Shuffled	0.9366	0.8038
	SST	Original	0.67	0.8695
		Shuffled	0.7481	0.858
	VIP	Original	0.6991	0.8515
		Shuffled	0.8238	0.8195

Supplementary Table 3. TCA error and similarity values across with across number of components, celltypes, and original vs shuffled data.

Supplementary Figure 1. Number of neurons per depth and area, for each cell type. Distribution of number of neurons recorded simultaneously across depths of V1 (top) and LM (bottom), in all recording sessions, for excitatory **(a)**, SST **(b)**, and VIP **(c)** mice. Note that our goal in this study was to chronically record the activity of distinct excitatory and inhibitory cell types, from the same field of view and across multiple sessions. Like in other studies involving the original Mesoscope[1,2], our system can record from a much larger number of neurons, however, our biological experiments were not designed for that purpose.

Supplementary Figure 2. Correlation of excitatory neurons within V1, within LM, and across V1-LM, during images and omissions. a-c Spearman correlation coefficients computed, at different moments in the trial, between activities of excitatory neurons located in different depths of V1-V1 (a), LM-LM (b), and V1-LM (c). **d-f** Change in correlation coefficients during images (top) and omissions (bottom) relative to the baseline correlation coefficient, computed for the real data (black), and trial shuffled data (gray). Correlation coefficients were quantified over 500 ms after images, and 750 ms after omissions. Red stars indicate statistical significance (two-sided t-test, real vs. shuffle data; p<0.05). Traces and error-bars: mean +/- SEM; n = 8 mice.

Supplementary Figure 3. Correlation **of SST neurons within V1, within LM, and across V1-LM, during images and omissions. a-c** Spearman correlation coefficients computed, at different moments in the trial, between activities of excitatory neurons located in different depths of V1-V1 (a), LM-LM (b), and V1-LM (c). **d-f** Change in correlation coefficients during images (top) and omissions (bottom) relative to the baseline correlation coefficient, computed for the real data (black), and trial shuffled data (gray). Correlation coefficients were quantified over 500 ms after images, and 750 ms after omissions. Red stars indicate statistical significance (two-sided t-test, real vs. shuffle data; p<0.05). Traces and error-bars: mean +/- SEM; n = 6 mice.

Supplementary Figure 4. Correlation of VIP neurons within V1, within LM, and across V1-LM, during images and omissions. a-c Spearman correlation coefficients computed, at different moments in the trial, between activities of excitatory neurons located in different depths of V1-V1 (a), LM-LM (b), and V1-LM (c). **d-f** Change in correlation coefficients during images (top) and omissions (bottom) relative to the baseline correlation coefficient, computed for the real data (black), and trial shuffled data (gray). Correlation coefficients were quantified over 500 ms after images, and 750 ms after omissions. Red stars indicate statistical significance (two-sided t-test, real vs. shuffle data; p<0.05). Traces and error-bars: mean +/- SEM; n = 9 mice.

Supplementary Figure 5. TCA with 10 components. a, b, and c depict the results for the TCA analysis for 10 components corresponding to Figure 3D,E,F, respectively.