Unexpected events modulate context signaling in VIP and excitatory cells of the visual cortex

Farzaneh Najafi[1,4 #], Simone Russo[3,5,6], Jérôme Lecoq[1,2 #]

[1] Allen Institute, Mindscope Program, [2] Neural Dynamics Program, [3] Brain and Consciousness Program, Seattle, WA, USA

[4] Georgia Institute of Technology, Biological Sciences, [5] Wallace H Coulter Department of Biomedical Engineering, Atlanta, GA, USA

[6] Department of Biomedical and Clinical Sciences, Università degli Studi di Milano, Milan 20157, Italy

\# Corresponding authors:

Jerome Lecoq: jeromel@alleninstitute.org

Farzaneh Najafi: fnajafi3@gatech.edu

Abstract

The visual cortex plays a significant role in constructing internal models that predict incoming sensory experiences. Unexpected stimuli that violate those predictions play an important role in how neural circuits learn these internal models. The cortical inhibitory network, particularly VIP cells, is hypothesized to act as a key player in representing unexpected stimuli due to its unique anatomical connectivity motifs; however, the exact information carried by VIP neurons during unexpected events remains unclear: to what extent does VIP activity reflect stimulus-specific prediction violations vs. broad context information such as shared network dynamics that may arise from the animal's internal representations or external cues signals? To address this question, we analyzed a dataset of cell-type specific, multi-area, two-photon calcium imaging during a behavioral task collected by the Allen Institute. Mice viewed repeated familiar images, as well as unexpected omissions of such images. Using dimensionality reduction, we extracted the contribution of context to the activity of individual neurons, defined as the shared neuronal activity across brain areas that is *not* driven by images or omissions. We demonstrate that during omissions the activity of VIP neurons in V1 and higher visual area LM is highly influenced by context, particularly in superficial layers. Excitatory neurons across all recorded layers and areas switched from weakly modulation by context during expected images to being largely modulated by context during unexpected omission. Our results suggest that during expected images context information is suppressed to allow for external stimuli

26   to drive neuronal activity. Conversely, during unexpected omissions, VIP neurons function as non-specific error encoders,

27   driving cortical networks to attend to context.

INTRODUCTION

The visual cortex is crucial in shaping our internal visual representations of the sensory environment. In the mouse visual

cortex, inhibitory neurons are diverse, with major classes including parvalbumin (PV), vasoactive intestinal polypeptide

(VIP), and somatostatin (SST) neurons, each with distinct subclasses as evidenced by transcriptomic characterization[1]. .

VIP cells, presumed to exert a dis-inhibitory effect on pyramidal cells, may play a key role in regulating information flow

within the cortical network due to their connectivity with associative cortical areas and neuromodulatory systems[2–4]. VIP

neurons exist in superficial (layer 2/3 primarily), and deeper (layer 4) layers of the cortex, suggesting their involvement

in both top-down and bottom-up information processing, respectively[5].

Circuit models suggest that unexpected stimuli trigger computations across both excitatory and inhibitory cortical cells to

refine sensory representations[6]. Empirical data indicate that unexpected stimuli elicit responses across multiple layers of

the cortex, involving inhibitory neurons[7]. A recent study demonstrated that the activity of VIP cells in the visual cortex

peaks during unexpected omissions of visual stimuli[8], but the underlying nature of these responses remains unclear. VIP

cells may signal the presence of a prediction error signal; alternatively, they may encode contextual changes that co-occur

during omissions, including behavioral, external, or internal states.

In the present study, we addressed this question using multi-area two-photon calcium imaging during a visually guided

behavior. This task contained both expected events (regular flashes of images) and unexpected events (image omissions).

We leveraged population activity decoding, dimensionality reduction and trial shuffling to examine whether VIP activity

during omissions may contain stimulus-specific prediction-error signals, or if it was primarily influenced by context

information. Importantly, "context" in our study broadly refers to shared activity across brain areas that are independent

of task-relevant features. This may involve behavioral features such as the animal's locomotion, external inputs such as

the sensory signals in the animal's environment, and internal representations related to the animal's current state.

We found that most VIP neurons, except those in layer 4 of the primary visual area, encoded context information. This

suggests that VIP neurons represent a general 'surprise' signal enriched with contextual details during unexpected events.

In contrast, excitatory neurons displayed weaker modulation by context during expected image presentations but were

significantly influenced by context during unexpected omissions. These results highlight the role of superficial VIP

neurons in updating cortical activity with contextual information in response to unexpected stimuli. This mechanism

potentially facilitates the association of contextual signals with unexpected events, thereby driving adaptive responses.

RESULTS

Distinct cortical pathways are thought to carry sensory and prediction signals[9,10], suggesting that cortical areas and cell

types might show unique responses to expected vs. unexpected events. To test this hypothesis, we leveraged two-photon

imaging data from the Allen Brain Observatory platform[8,11]. In this dataset, mice were trained to perform a go/no-go,

image-change detection task[8]. In brief, water-restricted animals were presented with a constant stream of natural images

(250 ms) interleaved with gray screens of matched luminance (500 ms). When a change in image identity occurred, mice

would receive a water reward if they licked immediately after the image change. Importantly, during imaging sessions in

well-trained mice, a pseudo-random 5% of non-change images were replaced with a gray screen (image omissions; Fig.

1A). Since mice had extensive experience with the highly regular timing of image presentations (every ~750 ms), the rare

stimulus omissions were unexpected events, as opposed to the expected, frequent, and repeated images. During this task,

the activity of excitatory neurons as well as inhibitory SST and VIP neurons were recorded separately, using 3 distinct

mouse lines (see Methods). Neuronal cells within each line were simultaneously imaged at 4 cortical depths in 2 visual

areas (V1 and higher visual area: LM; Fig. 1B,C; Supplementary Fig. 1).

We first studied how excitatory and inhibitory subtypes (VIP, SST) respond to image presentations and omissions, and

how their responses depend on cortical area and depth. The population average of neural activities demonstrated a clear

difference across cell classes. Excitatory neurons and, more robustly, SST neurons in all areas and layers were activated

by expected images, i.e. the expected events (Fig. 2A,B). Conversely, excitatory neurons, overall, did not respond to

omissions, except for those in the middle depth of LM (Fig. 2A). SST neuronal activity was slightly decreased after

omissions in all recorded locations (Fig. 2B). In sharp contrast to excitatory and SST neurons, VIP neurons in all layers

and areas were robustly activated after omissions (Fig. 2C). Specifically, VIP cells demonstrated small ramping activity

approximately 250 ms prior to each image presentation that further increased when the image was not presented (i.e.

omission), and was immediately inhibited by image presentation (Fig. 2C), confirming previous results using the same

behavioral task[8].

This multi-area dataset allowed comparison of neuronal responses across cortical depths of V1 and LM in each experiment. We found that excitatory neuronal responses to images were stronger in deeper layers of both visual areas, particularly V1 (Fig. 2A). SST responses to images were strongest at middle depths of V1, corresponding to layer 2/3 and superficial layer 4, but did not differ among cortical layers of LM (Fig. 2B). VIP responses to neither images nor omissions were significantly different across cortical layers or areas (Fig. 2C), as reported previously[5], although the responses were slightly stronger in deeper layers. Therefore, in both V1 and LM, images are more robustly represented by excitatory and SST neurons, particularly in deeper layers; while unexpected omissions are represented by VIP neurons (Stats: Supplementary Table 1).

It is suggested that feedback pathways, from higher order cortical areas to superficial layers of V1, convey prediction signals[9,10,12]. In light of these studies, we used simultaneous multi-plane recording to investigate if cortical interactions might be different when expectations are violated, i.e. during omissions. As a proxy for cortical interactions, we measured pairwise noise correlations across repeated image presentations, as well as omissions, for each cell type. Correlations were computed within each area (V1-V1, LM-LM; Supplementary Fig. 2-4), as well as across areas (V1-LM; Fig. 3). We also performed a shuffling analysis to control for firing rate impacts on correlations (Methods; Supplementary Fig. 2-4)

In the excitatory network, V1-V1 and V1-LM coactivations increased after image presentations, but only in deep layers (V1-LM: Fig. 3A; V1-V1: Supplementary Fig. 2A,D). Interestingly, excitatory network coactivation in LM-LM did not change after images, across any pair of cortical layers (LM-LM: Supplementary Fig. 2B,E). Following omissions, there was no significant change in the coactivation of excitatory neurons across visual layers and areas (Fig. 3A; Supplementary Fig. 2A,D).

In the SST network, images increased coactivation more broadly across visual layers and areas, particularly in our "middle" imaging depth, i.e., deep layer 2/3 and superficial layer 4 (Fig. 3B; Supplementary Fig. 3). Omissions, on the other hand, slightly decreased coactivation of neurons relative to the baseline across most layers of the SST network (Fig. 3B; Supplementary Fig. 3).

In the VIP network, coactivation patterns were strikingly different compared to other cell types: neurons in *superficial* layers of V1 and LM were strongly coactivated following omissions. There was also a weak coactivation prior to images in most layers and areas (Fig. 3C; Supplementary Fig. 4). Notably, correlations were overall much weaker among

excitatory neurons compared to inhibitory neurons (Fig. 3), confirming previous results and indicating stronger local

connectivity between inhibitory neurons[13,14].

Our results suggest that the VIP network in the visual cortex is strongly recruited during unexpected events: cells from

all layers of V1 and LM are active during omissions and neural correlations are especially strong in the superficial layers.

One possible explanation is that prediction errors (i.e. mismatch between the animal's expectation of a given image and

its omission) generate a shared input signal that co-activates VIP cells in superficial layers across areas, thereby

synchronizing the local cortical network. An alternative hypothesis is that VIP activity during omissions may be due to

co-occurring changes triggered by unexpected events. Such changes could include variations in animal's locomotion and

arousal, as well as alterations in the representation of environmental cues, or internal states. This hypothesis is

corroborated by recent reports[15,16], including our own[17], that have demonstrated a broad range of behavioral features

modulate brain-wide network dynamics, raising the possibility that our correlation findings could be contributed by global

brain fluctuations driven by widespread behavioral events. Importantly, in our previous work (Garrett et al. 2023), we had

found that the VIP population carries no information about the identity of the previous image. These results demonstrated

that VIP neurons do *not* encode stimulus-specific prediction error signals; instead suggesting that VIP omission activity

may represent a broad non-specific surprise signal.

To address whether VIP correlated activity during omissions may be affected by global contextual changes, we performed

Tensor Component Analysis (TCA)[18] on real and trial-shuffled data, aligned on images and omissions (number of sessions

= 22, 15, 21 for excitatory, SST, and VIP cells, respectively). TCA allowed decomposing neural population activity into

distinct components, and trial shuffling allowed studying contextual representation (see below). Each TCA component

was characterized by weights indicating the unique contributions from individual neurons, trials, and time points (Fig.

4A,B). Importantly, we defined TCA components as image-specific or omission-specific if their trial weights were

significantly higher on images or omissions, respectively (ranksum test across image and omission weights; $p < 0.05$; Fig.

4E). Components in which trial weights were not significantly different between images and omissions were excluded

from further analyses (number of non-specific components, pooled across sessions; mean +/- SD across 100 TCA

iterations; Excitatory: 38 +/- 4; SST: 20 +/- 3 ; VIP: 55 +/- 3).

To investigate the neural coding of images and omissions across visual areas and layers, we compared the absolute

magnitude of TCA neuronal weights among V1, LM, superficial and deep layers (imaging depth smaller or bigger than

250μm, respectively; Fig 4C,D red). In omission-specific components, VIP neurons had stronger weights in *superficial*

layers of both V1 and LM. This contrasted with excitatory and SST neurons which had stronger weights in deeper layers

of V1 and LM (Fig. 4C red). In image-specific components, excitatory neurons in deeper layers of V1 and LM had

stronger weights; while SST neurons showed stronger weights in V1 compared to LM, independent of cortical layer (Fig.

4D red). VIP neurons had few image-specific components, hence excluded from our analysis of neuronal weights.

TCA neural weights for image- and omission-specific components reflect the coding strength of individual neurons on

image and omission trials, respectively. This neural coding could arise from the direct presentation of images and

omissions, or could be influenced by co-occurring behavioral and sensory features in adjacent or remote areas, which we

collectively term contextual features. To distinguish the neuronal coding of images and omissions from these contextual

influences, we performed TCA on trial shuffled data, for which trial identities were shuffled *within* the image and omission

conditions for each neuron (see Methods). This shuffling procedure allowed us to preserve image- and omission-specific

responses while removing any contextual information that co-occurred across neurons during images and omissions. In

contrast, in the non-shuffle data, TCA components could encode the presence of images and omissions as well as any

context-dependent information present in the neuronal population.

Shuffling led to a general decrease of neuronal weight magnitudes for omission-specific components in all cell types (Fig.

4C blue: shuffled; red: real data). This reduction was most prominent in VIP neurons in the superficial layers, and absent

from the VIP neurons in deep V1 (Fig. 4C, right). These findings suggest that omissions are co-represented with contextual

information in the visual cortex in excitatory, SST, and particularly superficial VIP neurons. The major exception is that

VIP neurons in the deeper layers of V1 show a smaller contribution from contextual signals (Wicoxon ranksum test;

p=0.009).

Interestingly, shuffling had a smaller effect on the neuronal weights of image-specific components in excitatory and SST

cells (Fig. 4D blue vs. red; Note the absence of VIP neurons from this analysis due to their few numbers of image

components). This finding suggests that during images, excitatory neurons primarily encode image information,

overriding contextual signals. This is aligned with previous studies that demonstrated reduced neuronal variability during

sensory inputs[19].

Indeed, all cell types, and particularly VIP neurons, had more TCA components that were omission-specific vs. image-

specific (Fig. 4E, black: images, gray: omissions). This finding suggests that omission responses in our neuronal

population exhibit more variability than image responses, and is aligned with our TCA shuffled analysis which

demonstrated response variability during omissions was associated with the co-representation of contextual features. As

a control, we performed the same analysis with a larger number of reconstructed components by TCA and found similar

results (Supplementary Fig. 5).

DISCUSSION

Recent work had demonstrated that the VIP population is functionally quite diverse, and strongly represents behavioral

features and unexpected image omissions[8]. Extending these findings, our current study demonstrates that during

unexpected omissions, VIP neurons in the visual cortex not only encode a non-specific surprise signal but also a

substantial amount of contextual information. This dual representation of omissions and context is predominantly

observed in superficial VIP neurons, with notably reduced context coding in deeper layers.

Multiple lines of work have suggested the role of VIP neurons in encoding external and internal features. VIP neurons

can represent behavioral state, leading to gain modulation in sensory cortices[4,20]. VIP activity increases with pupil

dilation[21], locomotion[4,22], and whisking [3,23], leading to enhanced sensory modulation during high attention and arousal

states that follow surprising and reinforcement signals[4,20–22,24,25]. VIP neurons are driven by neuromodulators, particularly

cholinergic inputs[4], as well as long-range cortical and thalamic inputs[2,3]. VIP neurons have also been shown to play a key

role to surround suppression[26], when neuronal responses to centered visual stimuli can be decreased by the presence of

adjacent stimuli.

Our study suggests that the VIP network in primary and higher visual cortices comprises two subnetworks with distinct

roles, associated with superficial and deeper layers. In V1, the subnetwork in the deeper layers encodes information about

omissions, independently of context, while the subnetwork in the superficial layers concurrently encodes both context

and omission information. Importantly, the omission coding by VIP cells is non-specific, lacking any information about

which stimulus was omitted, as suggested by our recent work (Garrett 2023). This supports the hypothesis that VIP

neurons broadly encode surprise signals rather than specific mismatches between expected and actual stimuli. Our

findings suggest a potential specialization within the VIP network that enhances the reliable representation of sensory

features in deeper V1, while integrating sensory and contextual features in the superficial layers of V1, hence facilitating

adaptive responses to dynamic environmental stimuli.

The ability of VIP cells to disinhibit excitatory neurons through its connectivity pattern, is ideally suited to facilitate the

propagation of selective activity patterns through the cortex, leading to attention towards contextual sensory inputs[25] when

needed, or the emergence of motor behaviors[27,28]. We propose a framework in which VIP cells serve as gatekeepers of

contextual information, which are dynamically activated when transitioning from expected to unexpected stimuli (**Fig.**

5). This framework is in agreement with the numerous studies on the activity of the inhibitory network. Future experiments

should aim to simultaneously record from all those cell types in the context of dynamic but controlled sensory uncertainty.

Figure 1: Allen Institute Visual Behavior dataset: Multi-area, two-photon calcium imaging from visual areas V1 and LM during a visual task. A). Calcium signals were simultaneously imaged across two cortical columns, including 4 depths (approximately 100 µm, 200 µm, 300 µm, and 375 µm), in visual areas V1 and LM, and in 3 distinct mouse lines tagged for excitatory, and 2 inhibitory cell types: VIP and SST. **B).** Mice, walking on a disk, view a familiar set of images, separated by a gray screen. Images are unexpectedly omitted on 5% of the presentation. **C).** Example population activity of excitatory neurons, recorded simultaneously from ~1180 neurons in V1 and LM.

Figure 2. Excitatory and inhibitory cell types distinctly encode expected images and unexpected omissions across visual areas and layers. Left: Population-averaged calcium responses to images (shaded rectangle) and omissions (dashed rectangle), across 2 visual areas (V1, LM), and 4 cortical depths (blue, cyan, green, red), in 3 different mouse lines (A-C: excitatory, SST, VIP). ΔF/F traces normalized to baseline standard deviation. Mean +/- SEM; n = 24, 22, 24 sessions for excitatory, SST, and VIP, respectively. **Right:** quantification of evoked responses averaged over 350 ms after images, and 500 ms after omissions (time-windows used for quantification of image and omission-evoked responses are indicated by orange and gray horizontal lines above top left panel).

Figure 3. Distinct cortical interactions occur across primary and higher-order visual areas following expected and

unexpected events., Left: Spearman correlation coefficients between V1 and LM neurons at different cortical depths

(colors indicate LM depths; each subplot corresponds to a given V1 depth). **Right:** change in correlation coefficients

during images (left) and omissions (right) relative to baseline correlation coefficient. Correlation coefficients quantified

over 500 ms after images, and 750 ms after omissions. Traces and heatmaps: mean +/- SEM; n = 8, 6, 9 mice for

excitatory, SST, and VIP cell types, respectively.

Figure 4. Unexpected events lead to enhanced context representation in excitatory and particularly VIP neurons

in superficial layers. A) schematic of the analysis pipeline. The data were decomposed through Tensor Component

Analysis (TCA; red: original; blue: shuffled). The shuffled data were obtained by permuting trials identities within each

category (image and omission). Each TCA component was identified as image-specific, omission-specific, or non-specific

depending on the weight magnitude of the TCA trial component for images vs. omissions (Methods).**B)** left to right:

neuronal, temporal, and trials weights of a representative omission-specific TCA component (red: original; blue:

shuffled). In this example, trials from 1 to 159 are images, while the remaining trials are omissions. TCA was performed

on spikes extracted from DF/F traces aligned on images and omissions, including 300ms before to 700ms after images

and omissions. **C)** Neuronal weight of the omission-specific components in excitatory, SST, and VIP neurons (left to

right) extracted from original and shuffled data across areas (Wilcoxon ranksum test; all $p<0.001$ except for VIP V1 deep:

$p=0.009$). **D)** same as panel C for image-specific components in excitatory and SST neurons (left to right; Wilcoxon

ranksum test; all $p<0.001$ except for excitatory V1 superficial: $p=0.8$ and excitatory LM superficial: $p=0.5$). **E)** number

of components pooled across sessions, for excitatory, SST, and VIP neurons (left to right) from original data compared

between images and omissions (black and gray boxes, respectively; Wilcoxon ranksum test; all $p<0.001$).

Figure 5. <u>Proposed mechanism:</u> Triggered by unexpected events, VIP neurons gate contextual signal flow across cortical cell types and areas. Left) Expected stimuli, such as familiar images in our study, are encoded by excitatory and SST neurons. **Right)** When unexpected stimuli occur, a non-specific surprise signal activates VIP cells to promote contextual integration across all cell types. This contextual information may originate from nearby and distant brain areas, encoding a variety of signals, both internal, such as cognitive and motivational states, and external, such as sensory and behavioral information. The presence of this contextual information facilitates the updating of the unexpected stimulus representation in the local network.

ONLINE METHODS

Processing of calcium imaging movies

The preprocessing of all calcium imaging data was done within the imaging processing pipeline described in detail elsewhere[34] and crosstalk removal was applied after pre-processing. Briefly, all data was first corrected for brain motion by performing rigid registration in two dimensions. Then, cell-segmentation was performed to identify spatial masks of active neurons. Further, fluorescence from spatially-overlapping neuronal masks was unmixed and corrected for neuropil contamination. Finally, mask-matching and crosstalk removal was performed using FastICA[37] and ghost cells were filtered, and lastly, ΔF/F was computed on corrected masks.

Description of datasets

We used the publicly available Allen Institute Visual behavior dataset: https://portal.brain-map.org/circuits-behavior/visual-behavior-2p. Details are explained in our previous work (Garrett 2023). In brief, this dataset included 3 mouse lines: excitatory mouse line: Slc17a7-IRES2-Cre ; Camk2a-tTA ; Ai93(TITL-GCaMP6f), inhibitory subpopulation VIP: Vip-IRES-Cre ; Ai148 (TIT2L-GC6f-ICL-tTA2), and inhibitory subpopulation SST: Sst-IRES-Cre ; Ai148 (TIT2L-GC6f-ICL-tTA2). For the excitatory cell line, 24 sessions from 8 mice; for the SST cell line, 22 sessions from 6 mice; for the VIP cell line, 24 sessions from 9 mice were recorded.

Population averages of neuronal responses

Traces were aligned to the onset of image omission; since images were presented regularly, omission alignment also aligned the traces on images. The median of neural responses was computed across trials. The average response was then computed across neurons, for each session. A grand average was then computed across sessions (see Fig. 2, traces). To quantify the image-evoked responses, the calcium trace of each neuron was averaged over 350 ms after image onset for excitatory and SST neurons, and over [-250, 100] ms relative to image onset for VIP neurons to account for their anticipatory response. Response quantification was done on the mean (across trials) trace of each neuron. The same quantification was performed for omission-evoked responses, except a 500 ms window was used for quantification. The responses were averaged across neurons, for each session, and then a grand average was computed across sessions (see

278 Fig. 2,   Error-bars).

Correlation of neural responses across cortical planes

On omission-aligned traces, Spearman correlation coefficient was computed between pairs of neurons across trials. Here,

by trial we mean omissions. Correlations were computed for every individual frame, over [-1, 2] sec relative to the

omission. This procedure was done for all pairs of neurons; then an average value was computed across all pairs (Fig. 3).

Neuron pairs were present within the same plane, or in 2 different planes. This analysis allowed studying how the response

of neurons (within the same plane or in different areas/layers) covaried across trials, and how this coactivation changed

at different moments (e.g. after images vs. omissions). Spearman correlation coefficients were also computed on shuffled

traces (Supplementary Fig. 2-4 d-f), which were driven by independently shuffling trial orders for each neuron. For each

neuron pair, shuffling was repeated 50 times, resulting in a distribution of correlation coefficients for shuffled data. This

shuffling analysis controlled for firing rate impacts on our correlation results.

To compute "noise" correlations, we measured pairwise correlations of "signal"-removed traces. "Signal" was computed

by taking the average neural response to each image type (there were 8 images in each session), and subtracting the

average from the response of individual trials of that image type.

To quantify coactivation of neurons (Fig. 3, heatmaps; Supplementary Fig. 2-4d-f), correlations coefficients ("cc") were

first averaged over 500 ms after images for excitatory and SST neurons, and over [-250, 250] ms relative to image onset

for VIP neurons, accounting for their anticipatory response. Then, correlation coefficients were averaged across baseline

frames. We call this quantity "image cc". To quantify omission-evoked coactivation, we averaged correlation coefficients

over 750 ms after omissions. We call this quantity "omission cc". Next, we quantified baseline coactivation by averaging

correlation coefficients across baselines frames. Baseline was defined as the frame immediately preceding each image

presentation, for excitatory and SST neurons, and 250 ms earlier than each image presentation for VIP neurons. We call

baseline quantification of correlation "baseline cc". Finally, we measured the change in coactivation during images or

omissions by subtracting out "baseline cc" from "image cc", or "omission cc". This quantity is plotted in Fig. 3, heatmaps,

and Supplementary Fig. 2-4d-f, error-bars.

The Python package scipy (scipy.stats.spearmanr) was used for computing correlations, and p-values were computed in

2 ways: 1) using the p-value output of the spearmanr package; 2) manually computing the p-value by comparing the

correlation coefficient of real (non-shuffled) data with the shuffled distribution using 2-sided, 1-sample t-test (using the Python package scipy.stats.ttest_1samp).

Tensor Component Analysis

Tensor Component Analysis (TCA) was performed on inferred spikes from the DF/F trace using the tensortool package (https://github.com/neurostatslab/tensortools). TCA was set to extract 5 components from the trials split from -300 to +600 msec relative to the image or omission onset. The resulting components were first filtered according to their image- or omission-specificity. Specifically, we compared the weights of image and omission trials for each component (Wilcoxon ranksum test, alpha <0.05) and identified as condition-specific the components showing a significant difference between the two. On this subset of components, those with larger absolute average weights during image trials were classified as image-specific, while those with larger absolute average weights during omission trials were classified as omission-specific. To assess the variability across trials, on each subset of components we computed the standard deviation across trial weights. This entire procedure was performed 100 times to account for the variability between TCA decompositions. Finally, we wanted to obtain a null distribution that discarded context information while preserving image- and omission-specific information. To this aim, we shuffled trials for each neuron keeping separated image and omission trials (e.g. the activity of an omission trial was reassigned to another omission trial independently for each neuron). Similarly to the original data, the procedure was performed 100 times to account for variability in shuffling and TCA decomposition.

Statistical tests

We used two-way ANOVA, followed by Tukey HSD to compare population averages across cortical layers. Two-sided t-test was used to compare correlations between real and shuffled data, for each cortical plane. A p-value of 0.05 was used as the significance threshold. For comparison of the correlation coefficients, we used two-tailed, two-sample Kolmogorov–Smirnov tests.

Data availability

Data are publicly available as Allen Institute Visual Behavior Dataset: https://portal.brain-map.org/circuits-behavior/visual-behavior-2p

Code availability

Crosstalk removal was performed using custom routines employing FastICA which is available as part of scikit-learn python package (https://scikit-learn.org/). Code to perform population average and correlation analyses are available at: https://github.com/farznaj/multiscope_fn.

Acknowledgements

We wish to thank the founder of the Allen Institute for Brain Science, Paul G. Allen, for his vision, encouragement and support.

Competing interests

The dual-beam add-on module (J.L.) intellectual property has been licensed to Thorlabs. Inc., by the Allen Institute. S.R. is the Chief Medical Officer of Manava Plus.

Correspondence and requests for materials should be addressed to J.L, and F.N..

References

1.   Gouwens, N. W. *et al.* Integrated Morphoelectric and Transcriptomic Classification of Cortical GABAergic Cells.
Cell **183**, 935–953.e19 (2020).

2.   Wall, N. R. *et al.* Brain-Wide Maps of Synaptic Input to Cortical Interneurons. *J. Neurosci.* **36**, 4000–4009 (2016).

3.   Lee, S., Kruglikov, I., Huang, Z. J., Fishell, G. & Rudy, B. A disinhibitory circuit mediates motor integration in the
somatosensory cortex. *Nat. Neurosci.* **16**, 1662–1670 (2013).

4.   Fu, Y. *et al.* A cortical circuit for gain control by behavioral state. *Cell* **156**, 1139–1152 (2014).

5.   Garrett, M. *et al.* Experience shapes activity dynamics and stimulus coding of VIP inhibitory cells. *Elife* **9**, (2020).

6.   Hertäg, L. & Sprekeler, H. Learning prediction error neurons in a canonical interneuron circuit. *Elife* **9**, (2020).

7.   Hamm, J. P. & Yuste, R. Somatostatin Interneurons Control a Key Component of Mismatch Negativity in Mouse
Visual Cortex. *Cell Rep.* **16**, 597–604 (2016).

8.   Garrett, M. *et al.* Stimulus novelty uncovers coding diversity in visual cortical circuits. *bioRxiv* 2023.02.14.528085
(2023) doi:10.1101/2023.02.14.528085.

9.   Keller, G. B. & Mrsic-Flogel, T. D. Predictive Processing: A Canonical Cortical Computation. *Neuron* **100**, 424–
435 (2018).

10. Bastos, A. M. *et al.* Canonical microcircuits for predictive coding. *Neuron* **76**, 695–711 (2012).

11. de Vries, S. E. J. *et al.* A large-scale standardized physiological survey reveals functional organization of the
mouse visual cortex. *Nat. Neurosci.* **23**, 138–151 (2020).

12. Rao, R. P. & Ballard, D. H. Predictive coding in the visual cortex: a functional interpretation of some extra-
classical receptive-field effects. *Nat. Neurosci.* **2**, 79–87 (1999).

13. Najafi, F. *et al.* Excitatory and Inhibitory Subnetworks Are Equally Selective during Decision-Making and Emerge

Simultaneously during Learning. *Neuron* **105**, 165–179.e8 (2020).

14. Packer, A. M. & Yuste, R. Dense, unspecific connectivity of neocortical parvalbumin-positive interneurons: a

canonical microcircuit for inhibition? *J. Neurosci.* **31**, 13260–13271 (2011).

15. Steinmetz, N. A., Zatka-Haas, P., Carandini, M. & Harris, K. D. Distributed coding of choice, action and

engagement across the mouse brain. *Nature* **576**, 266–273 (2019).

16. Musall, S., Kaufman, M. T., Juavinett, A. L., Gluf, S. & Churchland, A. K. Single-trial neural dynamics are

dominated by richly varied movements. *Nat. Neurosci.* **22**, 1677–1686 (2019).

17. Ramadan, M. *et al.* A Standardized Nonvisual Behavioral Event Is Broadcasted Homogeneously across Cortical

Visual Areas without Modulating Visual Responses. *eNeuro* **9**, (2022).

18. Williams, A. H. *et al.* Unsupervised Discovery of Demixed, Low-Dimensional Neural Dynamics across Multiple

Timescales through Tensor Component Analysis. *Neuron* **98**, 1099–1115.e8 (2018).

19. Churchland, M. M. *et al.* Stimulus onset quenches neural variability: a widespread cortical phenomenon. *Nat.*

Neurosci. **13**, 369–378 (2010).

20. Pi, H.-J. *et al.* Cortical interneurons that specialize in disinhibitory control. *Nature* **503**, 521–524 (2013).

21. Reimer, J. *et al.* Pupil fluctuations track fast switching of cortical states during quiet wakefulness. *Neuron* **84**, 355–

362 (2014).

22. Pakan, J. M. *et al.* Behavioral-state modulation of inhibition is context-dependent and cell type specific in mouse

visual cortex. *Elife* **5**, (2016).

23. Yu, J., Hu, H., Agmon, A. & Svoboda, K. Recruitment of GABAergic Interneurons in the Barrel Cortex during

Active Tactile Behavior. *Neuron* **104**, 412–427.e4 (2019).

24. Letzkus, J. J. *et al.* A disinhibitory microcircuit for associative fear learning in the auditory cortex. *Nature* **480**,

331–335 (2011).

25.   Karnani, M. M. *et al.* Opening Holes in the Blanket of Inhibition: Localized Lateral Disinhibition by VIP

Interneurons. *J. Neurosci.* **36**, 3471–3480 (2016).

26.   Keller, A. J. *et al.* A Disinhibitory Circuit for Contextual Modulation in Primary Visual Cortex. *Neuron* **108**,

1181–1193.e8 (2020).

27.   Arroyo, S., Barati, S., Kim, K., Aparicio, F. & Ganguly, K. Emergence of preparatory dynamics in VIP

interneurons during motor learning. *Cell Rep.* **42**, 112834 (2023).

28.   Ren, C. *et al.* Global and subtype-specific modulation of cortical inhibitory neurons regulated by acetylcholine

during motor learning. *Neuron* **110**, 2334–2350.e8 (2022).