Synthetic diets containing a single polysaccharide disrupt gut microbial community

structure and microbial interaction networks in the American cockroach

Rachel Dockman[a], Elizabeth Ottesen[a#]

[a]Department of Microbiology, University of Georgia, Athens, Georgia, USA

Running Head: Synthetic Diets Modify Cockroach Microbiome

Address correspondence to Elizabeth Ottesen, ottesen@uga.edu

Abstract

Achieving and maintaining a healthy gut microbiome has numerous benefits for the host. Host

diet plays a key role in shaping the gut microbial community, and understanding how diet

composition influences gut microbiome structure and stability is key to developing effective

interventions to treat gut microbiome dysbiosis. We use the American cockroach (*Periplaneta*

americana) as a model system to dissect the response of gut microbes to host diet modification.

Here, we designed synthetic diets from lab-grade, purified ingredients to identify how the

cockroach gut community responds to different carbohydrate components (chitin,

methylcellulose, microcrystalline cellulose, pectin, starch, xylan) in otherwise balanced diets.

Using 16S rRNA gene sequencing, we show that synthetic diets produce replicable shifts in the

cockroach gut community diversity and phylogenetic composition, with xylan-fed insects

displaying the largest alterations. Comparison with cockroaches fed whole-food diets reveal that,

rather than introducing new microbes, synthetic diets alter microbiome composition by inducing

blooms among taxa present basally within the cockroach gut community. Synthetic diets are also

associated with less-robust, more fragmentary microbial co-occurrence networks compared to

cockroaches fed whole-food diets. Our results highlight the utility of lab-grade artificial diets in

microbiome research and shed light on how purified polysaccharides may exert more influence

over a stable gut community to generate noticeable change than whole food-derived fibers.

Importance

Cockroaches are unusual but effective model organisms for understanding how the gut

microbiome reacts to dietary changes. Previously, research in the cockroach gut microbiome

demonstrated surprisingly robust stability following extreme dietary shifts using whole foods,

but our work uncovers the potential of synthetic diets to induce change in these stable

communities. The observation that diets enriched in purified polysaccharides reduced gut

microbiome stability and undermined microbial interaction networks has important implications

for the use of prebiotics to induce targeted changes in gut microbiome composition and structure.

Introduction

The gut microbiome is a key player in host metabolism and homeostasis; it extracts energy from recalcitrant dietary components, provisions essential nutrients, and stimulates the host's immune system to protect against pathogens and toxins [1-4]. Microbiome research seeks to identify specific microbes and microbial community configurations that influence host wellbeing, as well as techniques to maintain a desirable community once achieved [5, 6]. Key drivers of gut microbiome community structure include host genetics [7, 8], environment, and most notably, diet [9, 10]. Diet has gained particular attention as the most easily manipulated of these factors, and a clear relationship exists between microbially derived metabolic products from the gut microbiome and overall host health [11, 12]. However, the complex nutritional needs of mammalian models limit the extent of dietary manipulation possible, while well-known insect models such as *Drosophila* have limited dietary range. To address this challenge, we are developing the omnivorous American cockroach (*Periplaneta americana*) as a model of microbiome dynamics that extends our understanding of human-relevant bacteria while offering the benefits of invertebrate research.

Omnivorous cockroaches such as *P. americana* are colonized by a complex hindgut microbiome that is taxonomically similar to the human colonic flora, consisting of many shared lineages of microbes, including diverse Bacteroidota, Firmicutes (now Bacillota), and Proteobacteria (now Pseudomonadota) [13-15]. The hindgut microbiota also plays an analogous role in host nutrition, with hindgut bacteria fermenting otherwise indigestible dietary components into volatile fatty acids that can be utilized as an energy supplement by their hosts or influence host development [16-19]. Further, the extreme dietary flexibility conferred by *Blattabacterium* endosymbionts and the unique ability for cockroaches to store nitrogen as uric acid for future use

facilitates extensive dietary manipulation [20-22]. Studies of cockroach gut microbiome

responses to diet have generated contrasting responses, with multiple large-scale studies finding

that the gut microbiome is highly stable following dietary shifts [14, 23, 24], while others have

demonstrated shifts in response to select diets [25-27]. Currently, there is no consensus on why

these studies produced differing results and comparison is difficult due to inconsistent use of

synthetic, natural, or whole food diets across studies. Whole food or natural diets may obscure

bacterial responses to nutritional contents given their complexity, while synthetic diets are more

amenable to precise dietary changes, thus controlling for more variables [28-31]. Artificial diets

have been successfully developed and used as replacement for natural diets in insects with far

more specialized dietary needs than cockroaches, implying that cockroaches are an ideal

candidate for use with lab-synthesized diets [32-35].

In order to facilitate precise manipulation of dietary composition in cockroaches, we have

developed a series of synthetic cockroach diets based on the work of early entomologists [36-

38]. These artificial diets serve as a nutritionally complete base to precisely evaluate the

influence of specific dietary components on the *P. americana* hindgut microbiome, which as

mentioned above had previously been shown to be strongly resistant to dietary manipulation

using whole food diets [14]. Using these synthetic diets as a base, we tested a spectrum of

polysaccharides as the primary carbon and energy source to identify if the hindgut microbiome

responds to specific fibers. We found that these diets resulted in much stronger impacts on gut

microbiome composition than even highly divergent whole food diets. In seeking to understand

these divergent responses, our results will facilitate future studies of gut microbiome responses to

diet in the cockroach and shed light on how gut microbiome stability fluctuates when challenged

with synthetic diets compared to complex whole food diets.

Results

Impacts of synthetic diets on gut microbiome diversity and community composition

We formulated a series of synthetic diets composed of a fixed base of 25% protein amended with dietary salts, vitamins, and cholesterol while differing only in complex carbohydrate type. Initial experiments utilized five alternative polysaccharide sources: chitin, methylcellulose, microcrystalline cellulose (MCC), pectin, or xylan. Following initial analysis of these results, we tested an additional starch-based diet. The prepared diets were readily consumed by the cockroaches in all cases.

To evaluate the impact of these diets on the gut community, each diet was fed to adult cockroaches for a period of 14 consecutive days, after which the insects were sacrificed, and their hindgut dissected out for 16S rRNA gene library sequencing. Following library preparation and sequencing, we used DADA2 to obtain 2,321,848 quality-controlled, assembled sequences assigned to 3308 amplicon sequence variants (ASVs) after removal of endosymbiont (*Blattabacterium* sp.) and mitochondrial sequences [39].

At the phylum level, at least 80% of each sample was dominated by Bacteroidota, Firmicutes, and Desulfobacterota, in agreement with previous studies on the cockroach gut microbiome (**Figure 1A**) [14, 40, 41]. The relative abundances of these three phyla were similar across all samples excluding xylan-fed cockroaches; these insects hosted notably more Firmicutes and less Desulfobacterota than cockroaches fed other diets (**Figure 1A**).

Alpha diversity, as measured by Shannon index, evenness, and community richness significantly differed across diet treatments (**Figure 1D-F;** Kruskal Wallis $p < 0.001$ for each). Pairwise analyses found that chitin-fed insects possessed higher Shannon index values ($p < 0.05$)

and community evenness ($p < 0.01$) than that of MCC- and starch-fed insects, while the xylan

diet resulted in lower alpha diversity measures than all other diets ($p < 0.05$ for each).

Beta diversity analyses using weighted and unweighted Bray Curtis dissimilarity revealed

significant impacts of our synthetic diets on gut microbiome composition. On average, between-

diet variation was greater than within-diet variation, with xylan-fed communities showing the

greatest dissimilarity (**Figure 1G and 1H**). Ordination analyses using non-metric

multidimensional scaling (NMDS) and PERMANOVA analysis showed that samples clustered

based on diet composition in both weighted (**Figure 1B;** PERMANOVA: R^2=0.42; $p<0.001$) and

unweighted (**Figure 1C;** PERMANOVA: R^2=0.329; $p<0.001$) Bray-Curtis metrics, with

especially clear separation of the xylan-based diet from other synthetic diets. Removing xylan-

fed samples from diversity calculations did not eliminate diet-based clustering for weighted

(**Figure S1A;** PERMANOVA: R^2=0.343; $p<0.001$) or unweighted (**Figure S1B;**

PERMANOVA: R^2=0.247; $p<0.001$) measures, suggesting that each carbohydrate source

enriched for a unique community composition. Interestingly, all samples produced significantly

larger weighted distances than unweighted from others within their diets, except for xylan-fed

samples (**Figure S2**; xylan vs xylan: $p = 0.1266$), suggesting that variation in microbial

abundance rather than microbial presence/absence may be driving much of the microbiome

response to diet.

We used DESeq2 to identify 76 microbes that exhibited significantly higher abundance in

a single synthetic diet across pairwise comparisons against all other treatments, which we termed

"diet-characteristic taxa" (**Figure S3**) [42]. Diet-characteristic ASVs were primarily assigned to

Firmicutes (n=48) and Bacteroidota (n=20); other phyla with diet-responsive taxa include

Fusobacteriota, Deferribacterota, Desulfobacterota, Fibrobacterota, and Spirochaetota. We found

that the chitin and methylcellulose diets were not associated with any diet-characteristic taxa by

this definition, while diets made with xylan, MCC, starch, and pectin enriched for 45, 10, 13, and

8 ASVs respectively.

Cohort effects on diet-driven differences in gut microbiome composition

To evaluate cohort effects on gut microbiome responses to diet composition, we

conducted a replicate experiment utilizing the MCC- and xylan-based diets. These diets were

selected for replication and follow-up experiments due to both their contrasting molecular

structure and the dissimilarity they generated in Bray-Curtis analyses (**Figure 1B-C**). Data from

the first and second experiments exhibited similar alpha diversity measurements (**Figure 2C-E**),

with replicates maintaining the significant shifts in alpha diversity ($p < 0.001$) observed in the

initial experiment between these two diets while showing no difference between replicates.

Beta diversity analysis showed that samples clustered by both replicate and diet (**Figure**

2A-B). Diet had large effects on both weighted (PERMANOVA: $R^2=0.34$; $p<0.001$) and

unweighted (PERMANOVA: $R^2=0.20$; $p<0.001$) Bray-Curtis dissimilarity, while replicate had a

much smaller effect significant for unweighted (PERMANOVA: $R^2=0.058$; $p<0.01$) but not

weighted (PERMANOVA: $R^2=0.037$; $p=0.1$) measures. Replicate cohorts differed primarily in

the representation of rare ASVs, with 27% of ASVs present uniquely in cohort 1 and 20.2% in

cohort 2 samples (**Figure 2F**). However, 66.7% and 61% of these taxa appeared only in one

sample (**Figure S4**), and overall represented 1.6-1.7% of total sequences obtained for each

cohort. Direct pairwise comparison of cohorts separated by diet treatment showed similar

patterns, with rare taxa contributing few reads despite comprising 19.7-33.7% of unique ASVs

(**Figure 2F**). Altogether, these results show that synthetic diets reproducibly alter the gut

microbiome composition in cockroaches.

Microbiome diversity comparison with whole food diets

The changes in community composition triggered by our synthetic diets were unexpected

given that previous experiments examining the impact of whole food diets with strongly

differing macronutrient profiles did not induce substantial change in gut microbiome

composition [14]. Therefore, we compared the samples from this current study ("synthetic" diet

type) to samples from the previous study ("whole food" diet type) of cockroaches fed butter,

tuna, honey, white flour, or whole wheat flour [14].

We found that gut microbiome samples from cockroaches fed synthetic diets exhibited

higher ASV richness ($p < 0.01$) but lower evenness ($p < 0.001$) and Shannon index ($p < 0.01$) than

those from insects fed whole foods (**Figure 3C-E**). Synthetic and whole food diets produced

distinct diet type clusters in NMDS ordinations (**Figure 3A-B**) for weighted (PERMANOVA:

$R^2 = 0.105$; $p < 0.001$) and unweighted (PERMANOVA: $R^2 = 0.152$; $p < 0.001$) Bray-Curtis

dissimilarities. When we analyzed the samples by diet, we found diet explained more variation in

NMDS ordination than diet type as interpreted from PERMANOVA R^2 values (weighted:

$R^2 = 0.393$; unweighted: $R^2 = 0.369$).

Beta dispersion analysis of variation within diet types showed that, together, the gut

microbiota of cockroaches fed synthetic diets was more variable than that observed among whole

food fed cockroaches (**Figure S5A-B**; Tukey's HSD: $p < 0.001$). However, when diets were

analyzed individually, they were equally dispersed in weighted Bray-Curtis dissimilarity (**Figure

S5A)** but not unweighted measures (**Figure S5B;** ANOVA: $p < 0.001$). Xylan-fed cockroaches

exhibited significantly greater within-group variability than all ten other diets (Tukey's HSD

range: $p = 0.037 - 3.17e^{-06}$), with no significant differences observed in pairwise comparisons of

all other diets. When synthetic diets were compared to whole food diets without including xylan-

fed samples, we observed no significant differences in unweighted beta dispersion (**Figure S5D)**

or Shannon index values (**Figure 3D, red boxes**). However, significant differences remained in

weighted Bray-Curtis dispersion (**Figure S5C**), richness and evenness (**Figure 3C, 3F**), and in

both weighted and unweighted Bray-Curtis ordination analysis **(Figure S6A-B)**, highlighting

that the altered microbiomes produced by the synthetic diet type were not solely due to biases

produced by xylan-fed samples.

Core taxa differences between synthetic and whole food diets

Given these strong differences in community structure, we utilized the R package

UpSetR to determine how the ASVs in different diet types overlap [43]. UpSet plots are akin to

Venn diagrams, considering only presence/absence of an ASV. Rarefied count tables were

aggregated by diet and ASVs were marked as either present or absent per diet. ASVs present in

the same subset of diets were grouped into "Sets", with the phylum-level composition per set

depicted as stacked bar charts labelled with the number of included ASVs (**Figure 4A**). We

supplemented the UpSet plot with pie charts illustrating the relative abundance (calculated as the

fraction of total reads recovered from the collapsed treatment group) of reads assigned to ASVs

within each set (**Figure 4A**). For simplicity's sake, **Figure 4** shows only the five largest

intersecting sets as well as all single-diet sets; additional sets are presented in **Figure S7**.

A total of 492 ASVs ("Set 1") were shared across all diet treatments (**Figure 4A**). These

ASVs made up over half of the sequences recovered for all diets except the MCC diet, for which

they represented 49% of sequences. Only 43 ASVs ("Set 4") were exclusive to the whole food

diets, contributing between 0.9% and 1.65% of reads in these diet sets. The 57 ASVs ("Set 3")

identified as exclusive to synthetic diets made up 1.6-3.4% of the reads recovered from starch-,

pectin-, chitin-, and methylcellulose-fed cockroaches, and 7% of xylan- and 13% of MCC- fed

cockroaches. Together, these results argue that the synthetic diets did not eliminate core taxa

present in the guts of cockroaches fed whole foods, nor did they result in hindgut colonization by

a large new set of microbial taxa.

Similarly, individual synthetic diets were not associated with hindgut colonization by

large groups of unique microbes. In general, taxa that were unique to individual synthetic or

whole food diets represented a very small proportion of sequences recovered (**Figure 4A**). ASVs

unique to the MCC diet formed the second largest set overall, with 160 diet-unique taxa, yet they

only represented 0.4% of total recovered sequences (**Figure 4A**). A xylan-based diet, which

repeatedly produced the largest community dissimilarities (**Figure 3A-B; Figure S5**), was

associated with 90 diet-specific taxa comprising only 0.63% of reads. Interestingly, our analysis

revealed that a xylan-based diet did result in the loss of 136 taxa that were present in all other

diets in abundances ranging from 4.78%-10.14% ("Set 2" in **Figure 4A**). However, other sets

that excluded individual diets were substantially smaller (**Figure S7A**) suggesting that this was

not a common mechanism underlying the diet-driven changes in gut microbiome composition.

Instead, synthetic diet-driven changes in gut microbiome composition were primarily associated

with changes in the relative abundance of individual taxa that were consistently present in the

cockroach gut regardless of dietary treatment.

To follow up on these observations, we used DESeq2 to assess enrichment of individual

ASVs between synthetic vs whole food fed cockroaches. Of the 492 ASVs present in at least one

sample belonging to each diet (Set 1), 95 ASVs were significantly enriched in cockroaches fed

synthetic diets, while 38 ASVs were enriched in cockroaches fed whole food diets (**Figure 4B**;

padj < 0.001). Set 1 ASVs were present in all diets, often at high abundances, and shifts across

diet type were modest (log fold change <5). Bacteroidota, Desulfobacterota, Proteobacteria and

other phyla showed similar enrichment distributions to each other, but more Firmicutes were

enriched by the synthetic diets.

We also examined enrichment of ASVs that fell outside of Set 1 but were both abundant

(baseMean > 1) and present in at least 5 samples (**Figure 4C**). These 1270 ASVs generally had

smaller abundances than members of Set 1, but greater log fold changes between diet types

(**Figure 4C**). The ASVs *Ruminococcus_NA* and *Fusobacterium ulcerans* were exceptions, being

both highly abundant (baseMean = 1020 and 651, respectively) and unique to the synthetic diets.

In contrast, the *Christensenellaceae* ASV (R7_NA.68) unique to whole food diets had high log-

fold change, but a baseMean of only 4.29. Apart from a few highly abundant ASVs in the

synthetic diets, most abundant, diet-enriched taxa were shared by all diets, supporting that

change is driven by common gut bacteria restructuring the community rather than interloping

bacteria disrupting the community.

Differential diet-based fluctuations of abundant Firmicutes and Bacteroidota ASVs

To explore the impact of diet on microbial taxa associated with fiber degradation, we

evaluated dietary responses of abundant taxa within the Firmicutes and Bacteroidota. For this

analysis, we selected the two most abundant representatives of each of these phyla from each diet

and examined their abundance across all diets (**Figure 5**). We identified 16 Firmicutes and 11

Bacteroidota as most or second-most abundant in at least one diet.

The most abundant taxa from both Firmicutes and Bacteroidota represented a small

fraction of reads across the whole foods diets, consistent with the higher observed Shannon

diversity and evenness of the gut microbiome for whole food-fed cockroaches (**Figure 3D, E**). In

contrast, the synthetic diets produced strong 'blooms' of individual ASVs, particularly among

the Firmicutes (**Figure 5A**). Several abundant Firmicutes were both present in Set 1 (**Figure 4B**)

and enriched in individual synthetic diets (**Figure S3**), namely *Lachnoclostridium_NA* (xylan),

Lachnoclostridium_NA.1 (xylan), *Enterococcus_NA* (xylan), *Enterococcaceae_NA* (xylan), and

ChristensenellaceaeR7_NA (pectin). In contrast, Firmicutes identified as enriched in the MCC

and Starch diets were not typically found across all diets: *Ruminococcus_NA* (MCC),

Lachnospiraceae_NA.5 (starch), and *Ruminococcaceae_NA* (starch). Among the Bacteroidota

(**Figure 5B**), we observed greater overlap in the most abundant taxa present in each diet group,

with all but one (*3M1PL1.52termite_NA*) of the abundant Bacteroidota ASVs belonging to Set 1,

and only two classified as diet-characteristic: *3M1PL1.52termite_NA* (MCC) and

Bacteroides_NA.12 (pectin).

Analysis of Microbial Co-Correlation Networks

As synthetic diets appeared to reduce gut microbiome compositional stability and induce

blooms of individual taxa, we constructed co-correlation networks with SparCC to examine the

community structure underlying synthetic and whole food microbiome data sets [44]. To filter

out noise and reduce spurious correlations, datasets were filtered to include ASVs present in at

least 25% of samples for each diet type, resulting in 976 ASVs for whole food diets and 700

ASVs for synthetic diets. Networks were further pruned to contain edge weights with absolute

values larger than 0.4, removing 75 and 168 ASVs from whole food and synthetic networks

respectively. Both positive and negative edges were retained for network layout formation

(**Figure S8**), but for the rest of analyses only positive edges were considered (**Figure 6, Figure**

S9). After negative edge removal, the whole food network contained 875 ASVs with 9515 edges

forming two connected components (**Figure 6A**), while the synthetic diet network contained 497

ASVs with 2536 edges that formed six connected components (**Figure 6B**).

Gut microbiota from whole food-fed cockroaches formed an extensive and dense

interaction network (**Figure 6A**), with higher edge counts (**Figure 6C**), node counts (**Figure**

6D), network strength (**Figure S9**), and betweenness scores (**Figure S9**) than synthetic diets at

most levels of filtering based off SparCC correlation values. High-degree nodes, or ASVs with

large numbers of neighbors, were present throughout the network structure, while in the

synthetic diets two primary clusters of ASVs appeared with fewer connecting ASVS. When a

range of inclusion cutoffs were considered, the synthetic diet network degraded more quickly

than the whole food network into separate connected components (**Figure 6E)** displaying greater

fragility and tendency to fragment. These overall network structures suggest that the synthetic

diets disrupt the stability of the cockroach microbiome.

Discussion

The core finding of this work is that synthetic diets dominated by a single carbohydrate type induced marked changes in hindgut microbial community composition in omnivorous cockroaches. Diets featuring the hemicellulose xylan induced the largest shifts in alpha diversity, inter-individual variability, and overall community composition (**Figure 1**). Xylan is a common structural polysaccharide containing a xylose backbone with branching arabinose, glucuronic acid, and/or acetyl groups that has been extensively investigated for prebiotic potential both in livestock and humans [45-48]. Research investigating xylan degradation ability in gut microbiota mainly focuses on *Bacteroides* [49-54] although select Firmicutes such as *Roseburia intestinalis* have been identified as key butyrate-producing xylan fermenters [55-57]. Interestingly, we found that in cockroaches, xylan-based diets decreased the relative abundance of Bacteroidota, while increasing Firmicutes (**Figure 1A**). Concurrent with this enrichment of specific taxa, xylan decreased the overall diversity of the gut community (**Figure 1**). This result contrasts with studies of gut microbiome composition changes in chickens [58] and mice [59] fed diets supplemented with xylooligosaccharides where gut diversity increased, although both studies reflected the increase in *Lachnospiraceae* observed in our experiments.

However, while xylan-based diets induced the largest change in hindgut community composition, samples clustered by diet treatment even when the xylan treatment group was excluded (**Supplement 1**). Most of these diets were associated with 'blooms' of specific Bacteroidetes and Firmicutes, excluding those made with methylcellulose and chitin, which were not associated with enrichment of any single taxa in pairwise comparisons with the other diets. This suggests that while the disproportionate impact on community structure due to xylan

warrants its own research, synthetic diets in general appear to reduce hindgut microbiome

stability compared to whole food-based diets.

These results stand in stark contrast to previous work from multiple investigators, who

found minimal to no change in hindgut microbial community composition in response to diet

alterations [14, 23, 24]. A commonality between these experiments is that the investigators

utilized complex, minimally processed or whole food diets that were highly biased in

macromolecular composition. On the other hand, a few investigators have previously observed

substantial influence of diet on the gut microbiome composition [25-27]. These experiments all

utilized synthetic diets that contained purified carbohydrate and protein sources. For example, in

experiments using *B. germanica*, Pérez-Cobas et al [26] prepared synthetic diets with a dextrin

and micronutrient base amended with either 50% cellulose or 50% casein while Zhu et al [27]

used diets composed of a cellulose and micronutrient base with supplemented with 40% by mass

purified starch, casein, or sesame oil. In *P. americana*, Bertino-Grimaldi et al [25] utilized

purified cellulose to compare with sugarcane bagasse, a complex dietary substrate.

A disproportionate influence of synthetic diets with highly purified components on gut

microbial composition has been observed beyond cockroach research as well. Termites, a close

relative to cockroaches, responded to single carbohydrate source diets with larger alterations in

gut community composition than termites fed mixed-carbohydrate diets [60]. Other insect model

systems produced similar results, such as in silkworms [61], ladybugs [62], waxworms [63], and

honeybees [64]. Among mammals, dogs provided a purified diet also exhibited reduced alpha

diversity compared to those fed a complex diet [65], wild-caught mice transitioned from natural

diets to laboratory diets lost large portions of native gut microbes [66], and humans given meal

replacement shakes showed loss of biodiversity in their microbiome compositions [67].

Comparison of our dataset with data recovered in previous whole food-based dietary

experiments suggest that synthetic diets altered the gut community by inducing overgrowth of

microbes already present in the cockroach gut microbiome (**Figure 4).** Taxa that were unique to

individual diets represented <1% of sequence reads in all diets but starch, of which they made up

2.84% (**Figure 4**). In contrast, 15 out of 20 highly abundant Firmicutes and Bacteroidota

associated with one or more diets were shared across all diet types, while the remaining 5 were

found sporadically in other diets (**Figure 5**).

We hypothesize that highly purified synthetic diets enabled microbial 'specialists' to

bloom beyond their former constraints in the whole food diets. While the whole food diets used

in Tinker and Ottesen [14] were highly biased in macromolecular composition, as natural foods

they may have had a more diverse nutritional profile according to the "eye" of a bacterium [51,

68, 69]. Both whole and white wheat flour, for example, are composed predominantly of

endosperm-derived starch, which physically impacts microbial adhesion due to variance in

particle size following milling [70, 71]. The bran, germ, and, to a lesser degree, endosperm

fractions of wheat contain structural polysaccharides and bioactive phytochemicals that are

targeted by gut bacteria and influence health parameters of the host [28, 72-74], while honey

contains complex mixtures of sugars (glucose, fructose, disaccharides) and

fructooligosaccharides in addition to organic acids, nitrogenous compounds, vitamins, and bee-

derived enzymes [75]. Tuna and butter are similarly high-complexity substrates that offer

resident gut microbes diverse metabolizable compounds that are lost in purified dietary

components such as those utilized in our study. The purified components used in our synthetic

diets are not entirely homogeneous but are far more processed than the 'natural' whole foods

used in our comparison sets and lack bioactive compounds present basally in these diets [68]. As

a result, the less complex dietary structures of our synthetic diets may not require as much

cooperative metabolism to digest while the overall loss in nutritional diversity edges out bacteria

that are unable to keep pace with specialist polysaccharide fermenters.

This conclusion is supported by network analysis of the diet types (**Figure 6**), where the

whole food network is highly interconnected with numerous paths from one ASV to the next,

while the synthetic diet network is easily fragmented into modules of microbes that are weakly

or negatively associated with the other network members (**Figure S9**). Hallmarks of a stable and

resilient gut microbiome often include metabolic redundancy, where many microbes are capable

of filling different niches and are rate limited by their partner activity but can adjust to shifts in

diet without extreme disruption [76, 77]. Synthetic diets may weaken the cooperative potential of

the gut microbiome, narrowing the niches that can flourish to fewer bacteria, and allow drastic

shifts such as those observed in the xylan diet that would otherwise not occur.

A key limitation of this study is the fact that the comparison group of "whole food" fed

cockroaches were from an earlier experiment and we lack 'contemporaneous' controls fed whole

food diets. However, an examination of cohort effects suggest that observed responses to

synthetic diets were highly conserved across cohorts in experiments conducted one year apart

(**Figure 2**). Further, despite the large time gap between the whole food and synthetic diet

experiments, a set of 492 ASVs (14% of total ASVs) shared by all treatment groups comprised

approximately 68% of the reads recovered (**Figure 4**), suggesting minimal drift in overall gut

microbiome composition due to time alone.

Conclusions

Overall, this study showed that synthetic diets that were highly enriched in a single carbohydrate type could induce large shifts in gut microbiome composition in the American cockroach, which has previously been shown to be highly resistant to diet-induced shifts in gut microbiome composition [14]. Changes in gut microbiome composition in response to synthetic diets were primarily associated with 'blooms' of diet responsive ASVs already present at lower levels in the gut rather than introduction new taxa. This alteration in gut microbiome composition is associated with fragmentation of gut microbiome interaction networks and increased inter-individual variability. Together, these results suggest that overconsumption of a single, purified class of polysaccharides can have destabilizing effects on the gut microbiota. This has major implications for the potential for prebiotics to induce changes in gut microbiome composition even in highly stable systems, as well as potential implications regarding the impact of highly processed foods on gut microbiome homeostasis. Future work will explore the functional and metabolic basis of observed shifts in microbial community composition.

Acknowledgements

This work was supported by the National Institute of General Medical Sciences of the National

Institutes of Health (NIH) under award number R35GM133789. We thank Darian Talamantes

for assistance with initial diet formulation. We also thank Dr. Kara Tinker for performing the

foundational work serving as the basis of this experiment.

Data Availability

Data associated with this study are available from the NCBI short-read database under

BioProjects PRJNA1096047 and PRJNA1105088. Data associated with the earlier study of the

impact of whole food diets on cockroaches is available under BioProject accession

PRJNA320546.

Materials and Methods

Insects and Experimental Conditions

Our *Periplaneta americana* colony has been maintained in captivity at the University of Georgia for over a decade. Mixed age and sex stock insects are maintained at room temperature in glass aquarium tanks with wood chip bedding and cardboard tubes for shelter in a 12:12 light:dark cycle. Water via cellulose sponge fit to a Tupperware reservoir and dog chow (Purina ONE chicken & rice formula, approximately 26% protein, 16% fat, and 3% fiber) are provided to stock colonies *ad libitum*.

Synthetic Diets

The synthetic diets created for dietary testing were designed to provide balanced nutrition while remaining malleable to component manipulation. Diets contained Vanderzant vitamin mix (MP Biomedicals, Irvine, CA, USA), Wesson salt mix (MP Biomedicals, Irvine, CA, USA), peptone (Bacto Peptone) and casein (Sigma-Aldrich, St. Louis, MO, USA). The peptone and casein utilized were high purity mixes with greater than 90% total protein content. Cholesterol (VWR International, Radnor, PA, USA) was added since insects are unable to synthesize sterols themselves. The dry ingredients were combined and suspended in minimal volumes of H_2O to create a batter or dough, formed into pellets, then dehydrated at 65°C until they were sufficiently dry to maintain shape. Food pellets were stored at -20°C until use.

In most experiments, the only component changed was the carbohydrate source. Polysaccharides used include microcrystalline cellulose (MCC; Sigma-Aldrich, St. Louis, MO, USA), methylcellulose (Sigma-Aldrich, St. Louis, MO, USA), xylan from corn core (TCI

Chemicals, Portland, OR, USA), pectin from apple (Sigma-Aldrich, St. Louis, MO, USA), starch

(Sigma-Aldrich, St. Louis, MO, USA), and chitin (Alfa Aesar, Ward Hill, MA, USA).

Experimental Design

Experimental conditions were prepared as described in [14]. Briefly, mixed-sex healthy

adult insects were transferred from the stock colony to plastic tanks containing pebbles and

bleached polyvinyl chloride tubes for footing and shelter, respectively. Food and water were

provided *ad libitum* in rigid plastic or glass dishes following two days of food restriction and

habituation. Dietary treatment lasted two weeks, during which debris, oothecae, and lethargic

insects were removed daily.

Upon completion of dietary treatments, each insect was isolated in a sterile culture plate

and placed on ice until torpid. Sternites were removed with sterile forceps to expose the intact

gut and fat body tissue was cleaned away. The cleaned gut was frozen on a sterile aluminum dish

on dry ice and divided into foregut, midgut, and hindgut sections for collection in 500-800μL

phosphate-buffered saline (1X PBS). Gut contents and tissue-attached bacteria were disrupted

with a sterile pestle, and the samples stored at –20°C until DNA extraction. For this study, only

the hindgut community was analyzed due to higher microbial density and activity than in other

gut regions.

DNA Extraction

DNA was extracted from 200μL aliquots of individual samples using the EZNA Bacterial

DNA Kit (Omega Biotek, Norcross, GA, USA) with some modifications. Sample aliquots were

centrifuged at 5000g for 10min, with the resulting pellet resuspended in 100μL TE buffer plus

10μL lysozyme (50mg/mL) and incubated for 30 min at 37°C. Following incubation, samples

were vortexed with glass beads (25mg, Omega Biotek) for 5 min at 3000rpm, then incubated at

55°C for one hour with 100μL TL buffer, 20μL proteinase K and continuous 600rpm shaking.

The kit protocol was followed for additional incubations with BL buffer and DNA isolation

using the provided column. DNA was eluted into 50μL of provided Elution Buffer and

quantified using either a Nanodrop Lite spectrophotometer (Thermo Scientific) or the Take3

plate for BioTek plate readers (Agilent).

16S rRNA Gene Library Preparation and Sequencing

The V4 region of the 16S rRNA gene was amplified via 2-step polymerase chain reaction

(PCR) from individual hindgut lumen samples as previously described in [14, 41, 78]. Both PCR

reactions used 0.02U/L Q5 Hot Start high-fidelity DNA polymerase (New England BioLabs,

Ipswich, MA, USA) with 200M dNTPs and 0.5M forward and reverse primers in 1M Q5

reaction buffer. The first 10μL reaction containing 3ng DNA and primers targeting the V4 region

(**515F**: GTGCCAGCMGCCGCGGTAA; **806R**: GGACTACHVGGGTWTCTAAT) was

performed under the following conditions: activation at 98°C for 30s; 15 cycles of 98°C for 10s,

52°C for 30s, and 72°C for 30s; final extension at 72°C for 2 min. Immediately following

amplification, 9μL of the first reaction was added to 21μL of Q5 reaction mix containing

barcoded primers with adaptor sequences for Illumina sequencing [79]. Cycling was performed

as follows: activation at 98°C for 30s; 4 cycles of 98°C for 10s, 52°C for 10s, and 72°C for 30s;

6 cycles of 98°C for 10s and 72°C for 1 min; final extension at 72°C for 2 min.

After product size verification via gel electrophoresis, samples were cleaned as instructed

in Omega Biotek's Cycle Pure kit, quantified, and pooled for equimolar representation of each

sample. Prepared libraries were sent to the Georgia Genomics and Bioinformatics Core at the

University of Georgia for 250 base pair paired-end Illumina MiSeq sequencing.

DNA Sequence Processing

Each dataset collected in this study was processed separately in R (version 4.2.1) by

sequencing run using R package DADA2 (version 1.24.0), with the cumulative Amplicon

Sequence Variants (ASVs) generated input as a priors table for each successive run [39, 80]. To

allow for comparison with this dataset, raw data from previous research in the Ottesen lab were

reprocessed to generate ASVs following the same procedures as in this current study [14]. All

sequence tables produced by these datasets were combined by ASV sequence prior to taxonomy

assignment to ensure continuity in naming. Taxonomy was assigned using DADA2 and the ARB

Silva v138 classifier to the species level, uniquely numbered, and filtered to remove sequences

matching eukaryotic (chloroplast, mitochondria) or endosymbiotic *Blattabacterium* DNA [39,

81].

Community analysis

Alpha and beta diversity analyses were performed via the R package vegan (version 2.6-

4) [82]. Samples were rarefied prior to diversity analysis to 7924 reads for comparisons between

synthetic and/or whole food diets, and 9685 reads for analysis of repeat xylan and MCC diet

experiments. Alpha diversity was measured via Shannon index, the count of ASVs observed in

rarefied samples, and Pielou's evenness (calculated as Shannon/log(Observed)). Weighted and

unweighted Bray-Curtis dissimilarities were calculated, assessed for dispersion, and plotted

using the vegan functions vegdist, betadisper, and metaMDS. Statistics for alpha diversity

indices were calculated with the Wilcoxon rank sum test (pairwise comparisons) and Kruskal-

Wallis test (multi-group comparisons). The significance of community composition differences

observed in beta diversity measures was assessed using PERMANOVA. Beta dispersion was

further examined through the Tukey's HSD test for pairwise comparisons and ANOVA for

multi-group comparisons.

Differential abundance analysis was conducted using DESeq2 (version 1.36) [42]. For

identification of 'diet-characteristic taxa", raw count data for the synthetic diet set (n=66) was

filtered to exclude ASVs present in less than 5 samples and run through the 'DESeq' command

with parameters 'fitType = "local"' and 'design = ~ Diet'. Pairwise result tables were obtained

for all diet comparisons and filtered for significant data, defined as having an adjusted p-value

smaller than 0.05 and a baseMean larger than 10. ASVs significantly upregulated for one diet vs

the other five diets were identified as diet-characteristic (n=76) and used to generate the heatmap

in **Supplement 3**. For comparison of "synthetic vs whole", raw count data for both diet sets

(n=125) were combined and filtered to exclude ASVs that appeared in fewer than 5 samples.

DESeq was run with parameters 'fitType = "local"' and 'design = ~ Diet_Type' to identify

differentially abundant ASVs between the diet types. The resulting baseMean and log2 fold

change were used to generate the MA plots in **Figure 4**.

UpSetR (version 1.4) was employed to visualize intersecting sets of taxa, providing

insights into the distribution of taxonomic features across samples [43]. For UpSet analysis,

samples were rarefied to 7924 reads then collapsed together to obtain total counts per diet. Both

a presence/absence table and a proportion table were generated from these data, with the

presence/absence table used for UpSet graph generation. Relative abundance of each set was

calculated using the proportion table with ASVs collapsed per set and visualized as pie charts

within the UpSet graph.

Co-correlation analysis was conducted to evaluate the impact of synthetic diets on

microbial interaction networks using the SparCC procedure [44]. Networks were constructed

separately for the synthetic diet group and the whole food diet group using sequence count tables

that were filtered to only include ASVs with at least 5 representatives present in 25% of samples

(synthetic: 17 samples; whole food: 15 samples), preventing spurious correlations from rare taxa.

SparCC was implemented in R with standard parameters, and the resultant networks were

characterized and analyzed with the igraph R package (version 1.5.1) [83, 84]. Networks were

pruned to contain only edges with a correlation absolute value of at least 0.4 and exported into

Cytoscape for visualization using the edge-weighted spring embedded layout method [85].

References

1.   Huttenhower, C., et al., *Structure, function and diversity of the healthy human microbiome.* Nature, 2012. **486**(7402): p. 207-214.

2.   Belkaid, Y. and O.J. Harrison, *Homeostatic Immunity and the Microbiota.* Immunity, 2017. **46**(4): p. 562-576.

3.   Thaiss, C.A., et al., *The microbiome and innate immunity.* Nature, 2016. **535**(7610): p. 65-74.

4.   Bäckhed, F., et al., *Host-bacterial mutualism in the human intestine.* Science, 2005. **307**(5717): p. 1915-20.

5.   Kolodziejczyk, A.A., D. Zheng, and E. Elinav, *Diet-microbiota interactions and personalized nutrition.* Nat Rev Microbiol, 2019. **17**(12): p. 742-753.

6.   Fassarella, M., et al., *Gut microbiome stability and resilience: elucidating the response to perturbations in order to modulate gut health.* Gut, 2021. **70**(3): p. 595-605.

7.   Goodrich, Julia K., et al., *Human Genetics Shape the Gut Microbiome.* Cell, 2014. **159**(4): p. 789-799.

8.   Kurilshikov, A., et al., *Host Genetics and Gut Microbiome: Challenges and Perspectives.* Trends Immunol, 2017. **38**(9): p. 633-647.

9.   Sonnenburg, E.D., et al., *Diet-induced extinctions in the gut microbiota compound over generations.* Nature, 2016. **529**(7585): p. 212-5.

10.   Kurilshikov, A., et al., *Large-scale association analyses identify host factors influencing human gut microbiome composition.* Nat Genet, 2021. **53**(2): p. 156-165.

11.   Wolter, M., et al., *Leveraging diet to engineer the gut microbiome.* Nat Rev Gastroenterol Hepatol, 2021. **18**(12): p. 885-902.

12.   Tanes, C., et al., *Role of dietary fiber in the recovery of the human gut microbiome and its metabolome.* Cell Host Microbe, 2021. **29**(3): p. 394-407.e5.

13.   Schauer, C., C.L. Thompson, and A. Brune, *The bacterial community in the gut of the Cockroach Shelfordella lateralis reflects the close evolutionary relatedness of cockroaches and termites.* Appl Environ Microbiol, 2012. **78**(8): p. 2758-67.

14.   Tinker, K.A. and E.A. Ottesen, *The Core Gut Microbiome of the American Cockroach, Periplaneta americana, Is Stable and Resilient to Dietary Shifts.* Appl Environ Microbiol, 2016. **82**(22): p. 6603-6610.

15.   Cruden, D.L. and A.J. Markovetz, *Microbial ecology of the cockroach gut.* Annu Rev Microbiol, 1987. **41**: p. 617-43.

16.   Jahnes, B.C. and Z.L. Sabree, *Nutritional symbiosis and ecology of host-gut microbe systems in the Blattodea.* Current Opinion in Insect Science, 2020. **39**: p. 35-41.

17.   Zurek, L. and B.A. Keddie, *Contribution of the colon and colonie bacterial flora to metabolism and development of the american cockroach Periplaneta americana L.* Journal of Insect Physiology, 1996. **42**(8): p. 743-748.

18.   Ayayee, P.A., et al., *The role of gut microbiota in the regulation of standard metabolic rate in female Periplaneta americana.* PeerJ, 2018. **6**: p. e4717.

19.   Vera-Ponce de León, A., et al., *Microbiota Perturbation or Elimination Can Inhibit Normal Development and Elicit a Starvation-Like Response in an Omnivorous Model Invertebrate.* mSystems, 2021. **6**(4): p. e0080221.

20.   Ayayee, P.A., T. Larsen, and Z. Sabree, *Symbiotic essential amino acids provisioning in*
the American cockroach, Periplaneta americana (Linnaeus) under various dietary
conditions. PeerJ, 2016. **4**: p. e2046.

21.   Cochran, D.G., D.E. Mullins, and K.J. Mullins, *Cytological Changes in the Fat Body of*
the American Cockroach, Periplaneta aniericana,1 in Relation to Dietary Nitrogen
Levels. Annals of the Entomological Society of America, 1979. **72**(2): p. 197-205.

22.   Sabree, Z.L., S. Kambhampati, and N.A. Moran, *Nitrogen recycling and nutritional*
provisioning by Blattabacterium, the cockroach endosymbiont. Proc Natl Acad Sci U S
A, 2009. **106**(46): p. 19521-6.

23.   Schauer, C., C. Thompson, and A. Brune, *Pyrotag sequencing of the gut microbiota of*
the cockroach Shelfordella lateralis reveals a highly dynamic core but only limited effects
of diet on community structure. PLoS One, 2014. **9**(1): p. e85861.

24.   Lampert, N., A. Mikaelyan, and A. Brune, *Diet is not the primary driver of bacterial*
community structure in the gut of litter-feeding cockroaches. BMC Microbiol, 2019.
19(1): p. 238.

25.   Bertino-Grimaldi, D., et al., *Bacterial community composition shifts in the gut of*
Periplaneta americana fed on different lignocellulosic materials. Springerplus, 2013. **2**:
p. 609.

26.   Pérez-Cobas, A.E., et al., *Diet shapes the gut microbiota of the omnivorous cockroach*
Blattella germanica. FEMS Microbiol Ecol, 2015. **91**(4).

27.   Zhu, J., et al., *Diet Influences the Gut Microbial Diversity and Olfactory Preference of*
the German Cockroach Blattella germanica. Curr Microbiol, 2022. **80**(1): p. 23.

28.   Okarter, N. and R.H. Liu, *Health benefits of whole grain phytochemicals.* Crit Rev Food
Sci Nutr, 2010. **50**(3): p. 193-208.

29.   Tuncil, Y.E., et al., *Delayed utilization of some fast-fermenting soluble dietary fibers by*
human gut microbiota when presented in a mixture. Journal of Functional Foods, 2017.
32: p. 347-357.

30.   Koropatkin, N.M., E.A. Cameron, and E.C. Martens, *How glycan metabolism shapes the*
human gut microbiota. Nat Rev Microbiol, 2012. **10**(5): p. 323-35.

31.   Adesogan, A.T., et al., *RUMINANT NUTRITION SYMPOSIUM: Improving cell wall*
digestion and animal performance with fibrolytic enzymes1. Journal of Animal Science,
2014. **92**(4): p. 1317-1330.

32.   Piper, M.D., et al., *A holidic medium for Drosophila melanogaster.* Nat Methods, 2014.
11(1): p. 100-5.

33.   Talyuli, O.A., et al., *The use of a chemically defined artificial diet as a tool to study*
Aedes aegypti physiology. J Insect Physiol, 2015. **83**: p. 1-7.

34.   Gonzales, K.K., et al., *The Effect of SkitoSnack, an Artificial Blood Meal Replacement,*
on Aedes aegypti Life History Traits and Gut Microbiota. Sci Rep, 2018. **8**(1): p. 11023.

35.   Majumder, R., et al., *Artificial Larval Diet Mediates the Microbiome of Queensland Fruit*
Fly. Front Microbiol, 2020. **11**: p. 576156.

36.   Noland, J.L., J.H. Lilly, and C.A. Baumann, *Vitamin Requirements of the Cockroach*
Blatella Germanica (L.)1. Annals of the Entomological Society of America, 1949. **42**(2):
p. 154-164.

37.   Haydak, M.H., *Influence of the Protein Level of the Diet on the Longevity of*
Cockroaches. Annals of The Entomological Society of America, 1953. **46**: p. 547-560.

38.　Hamilton, R.L. and C. Schal, *Effects of Dietary Protein Levels on Reproduction and*
Food Consumption in the German Cockroach (Dictyoptera: Blattellidae). Annals of the
Entomological Society of America, 1988. **81**(6): p. 969-976.
39.　Callahan, B.J., et al., *DADA2: High-resolution sample inference from Illumina amplicon*
data. Nat Methods, 2016. **13**(7): p. 581-3.
40.　Dukes, H.E., K.A. Tinker, and E.A. Ottesen, *Disentangling hindgut metabolism in the*
American cockroach through single-cell genomics and metatranscriptomics. Front
Microbiol, 2023. **14**: p. 1156809.
41.　Tinker, K.A. and E.A. Ottesen, *Phylosymbiosis across Deeply Diverging Lineages of*
Omnivorous Cockroaches (Order Blattodea). Appl Environ Microbiol, 2020. **86**(7).
42.　Love, M.I., W. Huber, and S. Anders, *Moderated estimation of fold change and*
dispersion for RNA-seq data with DESeq2. Genome Biol, 2014. **15**(12): p. 550.
43.　Conway, J.R., A. Lex, and N. Gehlenborg, *UpSetR: an R package for the visualization of*
intersecting sets and their properties. Bioinformatics, 2017. **33**(18): p. 2938-2940.
44.　Friedman, J. and E.J. Alm, *Inferring Correlation Networks from Genomic Survey Data.*
PLOS Computational Biology, 2012. **8**(9): p. e1002687.
45.　Jefferson, A. and K. Adolphus, *The Effects of Intact Cereal Grain Fibers, Including*
Wheat Bran on the Gut Microbiota Composition of Healthy Adults: A Systematic Review.
Frontiers in Nutrition, 2019. **6**.
46.　Broekaert, W.F., et al., *Prebiotic and other health-related effects of cereal-derived*
arabinoxylans, arabinoxylan-oligosaccharides, and xylooligosaccharides. Crit Rev Food
Sci Nutr, 2011. **51**(2): p. 178-94.
47.　Jana, U.K., N. Kango, and B. Pletschke, *Hemicellulose-Derived Oligosaccharides:*
Emerging Prebiotics in Disease Alleviation. Front Nutr, 2021. **8**: p. 670817.
48.　Smith, M.M. and J. Melrose, *Xylan Prebiotics and the Gut Microbiome Promote Health*
and Wellbeing: Potential Novel Roles for Pentosan Polysulfate. Pharmaceuticals (Basel),
2022. **15**(9).
49.　Zhang, M., et al., *Xylan utilization in human gut commensal bacteria is orchestrated by*
unique modular organization of polysaccharide-degrading enzymes. Proc Natl Acad Sci
U S A, 2014. **111**(35): p. E3708-17.
50.　Déjean, G., et al., *Adaptation of Syntenic Xyloglucan Utilization Loci of Human Gut*
Bacteroidetes to Polysaccharide Side Chain Diversity. Appl Environ Microbiol, 2019.
85(20).
51.　Centanni, M., et al., *Differential growth of bowel commensal Bacteroides species on*
plant xylans of differing structural complexity. Carbohydrate Polymers, 2017. **157**: p.
1374-1382.
52.　Dodd, D., R.I. Mackie, and I.K. Cann, *Xylan degradation, a metabolic property shared*
by rumen and human colonic Bacteroidetes. Mol Microbiol, 2011. **79**(2): p. 292-304.
53.　Kmezik, C., et al., *A polysaccharide utilization locus from the gut bacterium*
Dysgonomonas mossii encodes functionally distinct carbohydrate esterases. J Biol Chem,
2021. **296**: p. 100500.
54.　Pereira, G.V., et al., *Degradation of complex arabinoxylans by human colonic*
Bacteroidetes. Nature Communications, 2021. **12**(1): p. 459.
55.　Mohand-Oussaid, O., et al., *The extracellular xylan degradative system in Clostridium*
cellulolyticum cultivated on xylan: evidence for cell-free cellulosome production. J
Bacteriol, 1999. **181**(13): p. 4035-40.

56.   Hershko Rimon, A., et al., *Novel clostridial cell-surface hemicellulose-binding CBM3*
654   *proteins.* Acta Crystallogr F Struct Biol Commun, 2021. **77**(Pt 4): p. 95-104.

655   57.   Leth, M.L., et al., *Differential bacterial capture and transport preferences facilitate co-*
656   *growth on dietary xylan in the human gut.* Nature Microbiology, 2018. **3**(5): p. 570-580.

657   58.   Zhou, J., et al., *Dietary supplemental xylooligosaccharide modulates nutrient*
658   *digestibility, intestinal morphology, and gut microbiota in laying hens.* Anim Nutr, 2021.
659   **7**(1): p. 152-162.

660   59.   Berger, K., et al., *Xylooligosaccharides Increase Bifidobacteria and Lachnospiraceae in*
661   *Mice on a High-Fat Diet, with a Concomitant Increase in Short-Chain Fatty Acids,*
662   *Especially Butyric Acid.* J Agric Food Chem, 2021. **69**(12): p. 3617-3625.

663   60.   Miyata, R., et al., *Influence of feed components on symbiotic bacterial community*
664   *structure in the gut of the wood-feeding higher termite Nasutitermes takasagoensis.*
665   Biosci Biotechnol Biochem, 2007. **71**(5): p. 1244-51.

666   61.   Dong, H.L., et al., *Differences in gut microbiota between silkworms (Bombyx mori)*
667   *reared on fresh mulberry (Morus alba var. multicaulis) leaves or an artificial diet.* RSC
668   Adv, 2018. **8**(46): p. 26188-26200.

669   62.   Xie, B.H., et al., *Analysis of gut microbiota of ladybug beetle (Harmonia axyridis) after*
670   *feeding on different artificial diets.* BMC Microbiol, 2024. **24**(1): p. 5.

671   63.   Gohl, P., C.M.R. LeMoine, and B.J. Cassone, *Diet and ontogeny drastically alter the*
672   *larval microbiome of the invertebrate model Galleria mellonella.* Can J Microbiol, 2022.
673   **68**(9): p. 594-604.

674   64.   Powell, J.E., et al., *The microbiome and gene expression of honey bee workers are*
675   *affected by a diet containing pollen substitutes.* PLoS One, 2023. **18**(5): p. e0286070.

676   65.   Allaway, D., et al., *Rapid Reconstitution of the Fecal Microbiome after Extended Diet-*
677   *Induced Changes Indicates a Stable Gut Microbiome in Healthy Adult Dogs.* Appl
678   Environ Microbiol, 2020. **86**(13).

679   66.   Martínez-Mota, R., et al., *Natural diets promote retention of the native gut microbiota in*
680   *captive rodents.* Isme j, 2020. **14**(1): p. 67-78.

681   67.   Gurry, T., et al., *Predictability and persistence of prebiotic dietary supplementation in a*
682   *healthy human cohort.* Scientific Reports, 2018. **8**(1): p. 12699.

683   68.   Puhlmann, M.-L. and W.M. de Vos, *Intrinsic dietary fibers and the gut microbiome:*
684   *Rediscovering the benefits of the plant cell matrix for human health.* Frontiers in
685   Immunology, 2022. **13**.

686   69.   Hernot, D.C., et al., *In Vitro Digestion Characteristics of Unprocessed and Processed*
687   *Whole Grains and Their Components.* Journal of Agricultural and Food Chemistry, 2008.
688   **56**(22): p. 10721-10726.

689   70.   Lin, J., Y. Gu, and K. Bian, *Bulk and Surface Chemical Composition of Wheat Flour*
690   *Particles of Different Sizes.* Journal of Chemistry, 2019. **2019**: p. 5101684.

691   71.   Fernando, W.M., et al., *The influence of environmental factors on the adhesion of*
692   *combinations of probiotics to rice fibre fractions.* World J Microbiol Biotechnol, 2012.
693   **28**(6): p. 2293-302.

694   72.   Cardona, F., et al., *Benefits of polyphenols on gut microbiota and implications in human*
695   *health.* J Nutr Biochem, 2013. **24**(8): p. 1415-22.

696   73.   Parkar, S.G., T.M. Trower, and D.E. Stevenson, *Fecal microbial metabolism of*
697   *polyphenols and its effects on human gut microbiota.* Anaerobe, 2013. **23**: p. 12-9.

74.   Adom, K.K., M.E. Sorrells, and R.H. Liu, *Phytochemicals and antioxidant activity of*
milled fractions of different wheat varieties. J Agric Food Chem, 2005. **53**(6): p. 2297-
306.
75.   Sultana, S., et al., *A Review of the Phytochemistry and Bioactivity of Clover Honeys*
(Trifolium spp.). Foods, 2022. **11**(13): p. 1901.
76.   Pereira, F.C. and D. Berry, *Microbial nutrient niches in the gut.* Environ Microbiol, 2017.
19(4): p. 1366-1378.
77.   Ze, X., et al., *Some are more equal than others.* Gut Microbes, 2013. **4**(3): p. 236-240.
78.   Tinker, K.A. and E.A. Ottesen, *Differences in Gut Microbiome Composition Between*
Sympatric Wild and Allopatric Laboratory Populations of Omnivorous Cockroaches.
Front Microbiol, 2021. **12**: p. 703785.
79.   Caporaso, J.G., et al., *Global patterns of 16S rRNA diversity at a depth of millions of*
sequences per sample. Proc Natl Acad Sci U S A, 2011. **108 Suppl 1**(Suppl 1): p. 4516-
22.
80.   RStudioTeam, *RStudio: Integrated Development for R.* 2020.
81.   Quast, C., et al., *The SILVA ribosomal RNA gene database project: improved data*
processing and web-based tools. Nucleic Acids Res, 2013. **41**(Database issue): p. D590-
6.
82.   Jari Oksanen, F.G.B., Michael Friendly, Roeland Kindt, Pierre Legendre, Dan McGlinn,
Peter R. Minchin, R. B. O'Hara, Gavin L. Simpson, and M.H.H.S. Peter Solymos, Eduard
Szoecs, Helene Wagner *vegan: Community Ecology Package.* R package version 2.5-6,
2019(https://CRAN.R-project.org/package=vegan).
83.   Csardi, G. and T. Nepusz, *The igraph software package for complex network research.*
InterJournal, 2006. **Complex Systems**: p. 1695.
84.   Csárdi, G., et al., *igraph: Network Analysis and Visualization in R.* 2024.
85.   Shannon, P., et al., *Cytoscape: a software environment for integrated models of*
biomolecular interaction networks. Genome Res, 2003. **13**(11): p. 2498-504.
725

Figure 1

Figure 1: Composition of gut microbiomes from cockroaches fed synthetic diets. (**A**) Barplot

showing the relative abundance of phyla across samples for each of the synthetic carbohydrate

diets. Bars represent individual hindgut samples, clustered and labeled by diet polysaccharide

source. Phyla present at an abundance greater than 1% in at least one sample are plotted. Non-

metric multidimensional scaling (NMDS) was used to plot (**B**) weighted and (**C**) unweighted

Bray-Curtis dissimilarity. The alpha diversity measures (**D**) Shannon index, (**E**) Pielou's

evenness, and (**F**) number of observed taxa were plotted. Boxplots of (**G**) weighted and (**H**)

unweighted Bray-Curtis display each diet vs self (colored boxes) and the other five synthetic

diets (white boxes). Samples were rarefied to a constant depth of 7924 sequences for alpha and

beta diversity calculations. For alpha diversity measures, pairwise statistics were calculated with

Wilcoxon rank-sum tests and multivariate analysis was performed using Kruskal-Wallis tests.

PERMANOVA was used to generate statistics for ordination analyses. "all" indicates $p<0.05$ vs

all other diets; * = $p<0.05$; ** = $p<0.01$

Figure 2: Analysis of cohort effects on gut microbiome responses to MCC and Xylan diets.

NMDS ordinations of (**A**) weighted and (**B**) unweighted Bray-Curtis dissimilarity. R^2 and p-values calculated by PERMANOVA for diet ("MCC" and "Xylan") and cohort ("Cohort 1" and "Cohort 2") separately and for interactions. Boxplots show observed (**C**) ASVs, (**D**) Shannon index, and (**E**) Pielou's evenness with Kruskal-Wallis p-values. The last panel (**F**) contains Venn diagrams of shared and unique ASVs between cohorts for both diets together as well as separately, constructed using rarefied count tables. Grey pie slices represent the percent of ASVs observed that are cohort-unique, while black pie slices represent the percentage of sequence reads assigned to the indicated unique ASVs. Xylan and MCC-fed samples from replicate experiments were rarefied to 9685 ASVs for alpha and beta diversity assessment. ** p<0.01 ***p<0.001

Figure 3: Whole food and synthetic diets differ in alpha and beta diversity measures.

NMDS ordinations of (**A**) weighted and (**B**) unweighted Bray-Curtis distances between

cockroach hindgut samples from this study ("Synthetic") and data from Tinker and Ottesen

(2016) for cockroaches fed tuna, butter, honey, wheat flour, and white flour ("Whole Food

Diets"). R^2 and p-values for diet type comparisons were calculated using PERMANOVA.

Boxplots show (**C**) observed ASVs, (**D**) Shannon index**,** and (**E**) Pielou's evenness for each diet

type, with red boxes representing the synthetic diets minus xylan-fed samples. Wilcoxon rank-

sum test was used for pairwise statistical analysis between diet types. The final panels display

boxplots of (**F**) weighted and (**G**) unweighted Bray-Curtis dissimilarity, with red boxes

763   representing synthetic diets excluding xylan. Raw sequence data from Tinker and Ottesen (2016)

764   were reprocessed using the methods in this experiment to generate comparable ASVs. All

765   samples were rarefied to 7924 ASVs for alpha and beta diversity analysis. ** =p <0.01; ***

766   =p<0.001, ns =no significance

Figure 4

Figure 4: Whole-food diets share more ASVs than synthetic diets. The rarefied samples from

Figure 3 were used to generate an UpSet plot of **(A)** ASVs present in at least one sample per diet;

the five largest intersections (Sets 1-5) and diet-unique sets are displayed. Pie charts represent

the percent of reads within a diet that originate from each set (red slices), and the bar charts are

colored to display the phylum-level distribution of ASVs assigned to each set. Pie charts to the

right of the UpSet plot represent the reads from ASVs that are shared by all diets in a diet type,

regardless of presence in the other diet type. For the MA plots in panels **(B)** and **(C)**, raw

sequence count tables were filtered to include ASVs present in at least 5 samples (out of 125

total) and analyzed using DESeq2 with diet type as the design factor. Panel **(B)** includes only

ASVs from Set 1, while panel **(C)** displays the ASVs, excluding Set 1, with an adjusted p-value

smaller than 0.001. The ASVs in both plots are scaled to the same baseMean size and colored

based on phylum. Note that the plots differ in their x-axis scale ($\log_2$ fold change).

Figure 5

A. Top Firmicutes ASV relative abundance

B. Top Bacteroidota ASV relative abundance

Figure 5. Individual ASVs explain large differences between synthetic but not whole-food

diets. Variance-stabilized count data from DESeq2 were used to determine the top two ASVs for

every diet belonging to **(A)** Firmicutes and **(B)** Bacteroidota, and the relative abundances of

ASVs belonging to the combined 'top ASV' set were plotted for all samples. Grey bars include

all Firmicutes or Bacteroidota not named in the key. * indicates "Other Firmicutes" extends

beyond 60% relative abundance; please refer to **Figure 1A** for full values. Bolded names

indicate diet-characteristic ASVs from **Supplement 3**, and italics indicate Set 1 ASVs from

Figure 4.

Figure 6

Figure 6: Synthetic diet correlation networks are smaller and less interconnected than whole food diet correlations networks. Networks were calculated by SparCC from filtered count tables (ASVs present in > 25% of samples per diet set) for synthetic and whole food diets separately to create two distinct networks containing 976 (whole food) or 700 (synthetic) nodes. Networks were imported into Cytoscape and edges with absolute values < 0.4 were removed to generate panels **(A)** and **(B)**. Negative edges were included during initial layout generation with the edge-weighted spring embedded layout method and are displayed in **Supplement 8.** The

800   number of nodes, edges, and connected components that remain when the networks are filtered

801   by increasing correlation values are charted in **C**, **D,** and **E** respectively.