Title: Significance of extended N-terminal region and a first motif residue in a third

transmembrane helix of a novel pump-like cation channelrhodopsin HulaCCR

Authors:

Shunki Takaramoto[1], Shai Fainsod[2], Takashi Nagata[1], Andrey Rozenberg[2], Oded Béjà[2,3], Keiichi

Inoue[1].*

Affiliations:

1. The Institute for Solid State Physics, The University of Tokyo, Kashiwa, Chiba 277-8581, Japan

2. Faculty of Biology, Technion — Israel Institute of Technology, Haifa 3200003, Israel

3. The Nancy and Stephen Grand Technion Energy Program (GTEP), Technion—Israel Institute of

Technology, Haifa 3200003, Israel

Corresponding author:

Correspondence to Keiichi Inoue. E-mail: inoue@issp.u-tokyo.ac.jp; telephone: +81-4-7136-3230;

postal address: Room A401, The Institute for Solid State Physics, The University of Tokyo, 5-1-5,

Kashiwanoha, Kashiwa, Chiba 277-8581, Japan.

Abstract

Microbial rhodopsins are a family of photoreactive membrane proteins that have seven

transmembrane helices and a retinal chromophore. Among them, channelrhodopsins are light-

induced ion channels making them popular tools for modulating neuronal activity in optogenetics.

Cation channelrhodopsins (CCRs), first recognized as the main photoreceptors in the model

chlorophyte *Chlamydomonas reinhardtii*, have since been identified in diverse species of green

algae, as well in other unicellular eukaryotes. The CCRs from non-chlorophyte species are

commonly referred to as the bacteriorhodopsin-like channelrhodopsins, or BCCRs, as most of them

feature the three characteristic amino acid residues of the "DTD motif" in the third transmembrane

helix (TM3 or helix C) matching the canonical DTD motif of the well-studied archaeal light-driven

proton pump bacteriorhodopsin. Here, we report characterization of HulaCCR, a novel BCCR

identified through metatranscriptomic analysis of a unicellular eukaryotic community in Lake Hula,

Israel. Interestingly, HulaCCR has an ETD motif in TM3 in which the first residue of the canonical

motif is substituted for glutamate. Electrophysiological measurements of the wild-type and a

mutant with a DTD motif of HulaCCR suggest the critical role of the first glutamate in spectral

tuning and gating. Additionally, HulaCCR exhibits long extensions at the N- and C-termini.

Photocurrents recorded from a truncated variant without the signal peptide predicted at the N-

terminus were diminished, and membrane localization of the truncated variant significantly

decreased, which indicates that the signal peptide is important for membrane trafficking of

HulaCCR. These characteristics of HulaCCR would be related to a new biological significance in

the original unidentified species, distinct from those known for other BCCRs.

Introduction

Microbial rhodopsins are a family of photoreactive membrane proteins that have seven

transmembrane helices and a retinal chromophore covalently binding to a conserved lysine residue

in the 7th transmembrane helix (called TM7 or helix G).[1–3] Among them, channelrhodopsins

(ChRs) are light-induced ion channels and are extensively utilized in optogenetics.[4,5]

Channelrhodopsins were first discovered in green algae when the photoreceptors responsible for

phototactic behavior in *Chlamydomonas reinhardtii* were identified as microbial rhodopsin

channels transporting cations, or cation channelrhodopsins (CCRs).[6–8] In contrast to

Chlamydomonas, two different types of ChRs were subsequently identified in the cryptophyte

Guillardia theta: a distinct family of CCRs, as well as a family of anions-transporting ChRs, or

anion channelrhodopsins (ACRs), which was the first indication of the high diversity of ChRs

among unicellular eukaryotes.[9,10] Natural and modified CCRs and ACRs have since been used

as molecular tools in optogenetic research to stimulate or inhibit neuronal firing by light,

respectively.[11–14]

Many microbial rhodopsins, including the well-characterized light-driven proton pump

bacteriorhodopsin from *Halobacterium salinarum* (*Hs*BR), feature three characteristic amino acid

residues together constituting the canonical "DTD motif" in the third transmembrane helix (called

TM3 or helix C). This motif is composed of two aspartic acid residues and a threonine residue:

D85/D96 and T89 in *Hs*BR, respectively.[15] One remarkable feature of the cryptophyte CCRs

from *Guillardia theta* was the presence of the DTD motif identical to that of *Hs*BR.[16,17]

Following the discovery of the CCRs from *G. theta*, many CCRs possessing the DTD or similar

motifs were found in other species, collectively referred to as the bacteriorhodopsin-like

channelrhodopsins (BCCRs). Among them, ChRmine from the cryptophyte *Rhodomonas lens*

exhibiting high photocurrents and high light sensitivity has emerged as the model BCCR and has

found its application in deep transcranial optogenetics.[18–20] Recent years have also witnessed

the discovery of highly potassium selective BCCRs, the kalium channelrhodopsins (KCRs) from

the stramenopiles *Hyphochytrium catenoides* (*Hc*KCR1 and *Hc*KCR2)[21] and *Wobblia lunata*

(WiChR)[22], as well as other heterotrophic flagellates. Since KCRs efficiently induce

hyperpolarization by K^+ efflux, they are used for precise optogenetic inhibition.

Here, we characterize HulaCCR, a novel microbial rhodopsin identified through

metatranscriptomic analysis of the unicellular eukaryotic community of Lake Hula, Israel. This

rhodopsin has an ETD motif in TM3 and belongs to the family of stramenopile CCRs which

includes KCRs and other CCRs from cultured heterotrophic flagellates and environmental sources

[22, 23] (Figs. 1a and b). While uncommon among BCCRs, the ETD motif is not an idiosyncrasy

of HulaCCR and is found in the related proteins mgCCR1[23] and P1ChR1[22]. The structure of

HulaCCR predicted with AlphaFold2 Colab [24] features an unusually long N-terminal extension

with a signal peptide and a prominent partially structured long C-terminal extension (Fig. 1b).

79   In this study, we investigated the function of HulaCCR using electrophysiological techniques.

As a result, it was revealed that this rhodopsin exhibits non-specific cation channel activity. To

understand the role of the substitution of glutamate for the ancestral aspartate at the first position

of the TM3 motif in HulaCCR, we examined the effects of this residue on the channel activity. In

addition, we investigated the role of the long extensions at the N- and C-termini of HulaCCR.

Results

Identification of HulaCCR

HulaCCR was identified as part of an ongoing metatranscriptomic study of the unicellular

eukaryotic community of Lake Hula in Northern Israel. This protein was chosen as a representative

of a BCCR clade abundant in the Hula assembly and to which the partially characterized ChRs

mgCCR1–3[23] also belong (Figs. 1a and b). Ideal for the study of the role of the carboxylate

amino acid at the first position of the TM3 motif, this clade demonstrates both DTD and ETD

variants, with mgCCR1 and HulaCCR having the otherwise uncommon glutamate. HulaCCR and

mgCCR1 are the closest relatives, with a modest overall pairwise sequence identity of 48%, least

similar in the N-terminal extension. The identity of the organism possessing these ChRs remains

unknown.

Ion transport activity of HulaCCR

Upon establishing the expression of HulaCCR in COS-1 cells, we investigated the absorption spectra of the protein solubilized from the membrane fraction of the cells with a solution containing n-dodecyl-β-D-maltoside (DDM). The absorption maximum wavelength (λ_{max}) of HulaCCR was determined to be 545 nm by a hydroxyl amine (HA) bleaching method[25] (Fig. 1c). To investigate the molecular function of HulaCCR, electrophysiological measurements were conducted. Based on the strong EYFP fluorescence on the outlines of transfected cultured cells, we determined that HulaCCR was well expressed in the plasma membrane (Fig. 1d). Whole-cell patch clamp recordings were performed using these cells. With standard pipette ($[K^+]$ = 128 mM) and extracellular ($[Na^+]$ = 142 mM) solutions, a significant photocurrent appeared upon illumination with 542-nm light (Fig. 1e). The photocurrent depended on the holding potential and exhibited an inversion from positive to negative currents along with increasing the holding potential from −80 to 60 mV, indicating that HulaCCR exhibits light-dependent ion channel activity. Changing the cations in both the patch pipette and extracellular solutions to N-methyl-D-glucamine (NMG) resulted in a profound decrease in photocurrent intensity (Fig. 1e, right). Furthermore, in this measurement, despite the anion conditions being asymmetric between the external and internal side of the membrane, where the pipette and extracellular solutions contained 90 mM glutamate and 151 mM Cl^-, respectively, the reversal potential remained at around 0 mV, excluding the possibility of anion channel activity and indicating H^+ channel activity. We conclude that HulaCCR possesses cation channel transport activity.

Next, to investigate the ion selectivity of HulaCCR, photocurrents were measured in extracellular solutions containing different cations (Fig. 2a), and current–voltage (*I–V*) curves were plotted based on peak current intensities (Fig. 2b). The *I–V* curves were fitted by a quadratic function, then the reversal potentials (E_{rev}) of the photocurrents were estimated. Taking the results

with NMG-containing extracellular solution as a reference, the reversal potential shift (ΔE_{rev}) was

calculated. ΔE_{rev} values increased with monovalent cations, while only low values were observed

with divalent cations, Mg^{2+} and Ca^{2+}. Hence, Mg^{2+} and Ca^{2+} hardly permeate through the channel

pore in HulaCCR. Assuming that only monovalent cations in the extracellular/pipette solutions

permeate, the relative cation permeability (P_{X+}/P_{H+}) was calculated using Goldman–Hodgkin–Katz

(GHK) equation (eq. 1) from ΔE_{rev} values,

$$\Delta E_{rev} = \frac{RT}{F} \ln \left(\frac{P_X[X^+]_o + P_H[H^+]_o}{P_H[H^+]_o}\right), \tag{1}$$

where R is the gas constant, T is the temperature, and F is the Faraday constant. The obtained

values P_{X+}/P_{H+} are shown in Fig. 2c. P_{Mg2+}/P_{H+} and P_{Ca2+}/P_{H+} were not determined by GHK flux

equation[26] for divalent cations at equilibrium (eq. 2) because the calculated values of P_{Mg2+}/P_{H+}

and P_{Ca2+}/P_{H+} were considerably smaller than the experimental error.

$$4P_{Mg} \frac{[Mg^{2+}]_i \exp\left(\frac{2E_{rev}F}{RT}\right) - [Mg^{2+}]_o}{1 - \exp\left(\frac{2E_{rev}F}{RT}\right)} + 4P_{Ca} \frac{[Ca^{2+}]_i \exp\left(\frac{2E_{rev}F}{RT}\right) - [Ca^{2+}]_o}{1 - \exp\left(\frac{2E_{rev}F}{RT}\right)} + P_H \frac{[H^+]_i \exp\left(\frac{E_{rev}F}{RT}\right) - [H^+]_o}{1 - \exp\left(\frac{E_{rev}F}{RT}\right)} = 0$$

(2)

From these results, it can be concluded that HulaCCR exhibits a strong ion selectivity for

monovalent cations, and this property is similar to those of GtCCR4[27] and ChRmine[28].

Moreover, there was not much difference in the selectivity among monovalent alkali metal cations

(Fig. 2c).

Mutational analysis of spectral tuning and gating kinetics in HulaCCR

Many BCCRs share the DTD motif in TM3 identical to the residues D85, T89, and D96 of HsBR.

As mentioned above, in HulaCCR, a glutamate (E176) is present at the position of the first motif

residue, forming an ETD motif (Fig. 1a). To investigate the roles of the amino acid residues near

the retinal chromophore, including E176 (Fig. 3a), photocurrents of several single mutants were

measured (Figs. 3b–e and S2). While the E176D mutant, in which the DTD motif is restored,

exhibited photocurrents similar to those of the wild-type protein (WT), its action spectrum slightly

blue-shifted compared to the WT spectrum (Fig. 3b), indicating the red-shifting spectral-tuning

effect of E176 in HulaCCR. To determine whether this residue is crucial for channel gating,

photocurrents were measured with a higher time resolution using nanosecond-pulsed laser flashes

(Fig. 3c). The time course of photocurrents due to channel opening and closing was reproduced

with a quadruple-exponential function. The obtained time constants were shown in Fig. 3d. Notably,

the closing rate of outward current in E176D was significantly slower compared to that of the WT,

suggesting the importance of E176 in the closing kinetics.

In chlorophyte channelrhodopsins and BCCRs, a cysteine residue is conserved at the position

corresponding to T90 in *Hs*BR in TM3 (Fig. 1a). This cysteine together with D156 in TM4

constitutes the "DC gate" in *Cr*ChR2 [29], and HulaCCR mutants in which C181 were substituted

with alanine or serine (C181A and C181S) exhibited significantly slower channel closing after 200-

ms illumination (Fig. S2d). Both mutants showed a highly prolonged channel closing rate in the

nanosecond-pulse excitation experiments, as observed in cysteine mutants of other

channelrhodopsins (Fig. 3e).[30,31]

Next, we attempted to tune the spectrum of HulaCCR by mutating amino acid residues near

the retinal chromophore. First we noticed that HulaCCR has C303 adjacent to K304 that forms the

Schiff-base linkage with the retinal chromophore in TM7, a trait shared most notably with the

KCRs (Figs. 3a, S1),[21] while most of the other BCCRs have alanine at this position similarly to

*Hs*BR. Importantly, in the case of *Cr*ChR2, substituting alanine for serine at this position causes a

red shift in the absorption spectrum.[32] Nevertheless, the action spectrum of the photocurrents of

the HulaCCR C303A mutant showed no significant red shift compared to the WT (Fig. S2c). Taking

a different approach, we attempted to introduce negative charges on the β-ionone ring side of the

retinal which is known to stabilize the electronically excited state, thus leading to a red shift in the

absorption spectrum.[33] Hence, the action spectra of mutants of residues shown in Fig. S2b,

L208M, A232C, A232S, and P276T were measured. This strategy did result in significantly red-

shifted action spectra compared to that of the WT (Fig. S2c).

Although the expected spectral shift did not occur in C303A, we found a significant difference

in other photocurrent behaviors in this mutant. The decay rate of photocurrents after light-off in

C303A was slower compared to that of the WT (Fig. S2a). The slower channel closing might be

the result of a slower photocycle, as seen for example in DC-gate mutants of CrChR2.[30,34] The

dependence of photocurrent intensity on excitation light power was also investigated under

continuous light illumination. In this case, it is anticipated that for channelrhodopsins with slower

photoreaction cycles, the photocurrent intensity will saturate at a lower light intensity compared to

those with faster photocycles. Hence, photocurrent intensities for the WT and C303A were

measured during continuous light illumination for 100 ms and were plotted against power density

of the excitation light (Fig. 3f). As a result, the C303A mutant with the slower channel closing rates

exhibited a lower EC_{50} value than that of the WT. Additionally, when compared to another BCCR,

ChRmine, which has a slower closing rate than that of the HulaCCR WT (Fig. S2a), the trend that

longer closing rates lead to lower EC_{50} values was maintained (Fig. 3f).

Previous studies on BCCRs suggested that amino acid residues on the intracellular side are

also crucial for the gating mechanism. In GtCCR2, a mutant in which D98 (equivalent to D96 in

HsBR) in TM3 was mutated to asparagine showed no significant channel current.[35] The effect

of the identical mutation has been investigated in other BCCRs, including ChRmine[28] and

HcKCR1[36], where the motif residues DTD are changed to DTN. It resulted in a decrease in cation

channel activity. Moreover, it has been previously suggested for GtCCR2 that proton transfer from

D98 (corresponding to D96 in BR) to an unknown cytoplasmic residue and back govern the

opening and closing of the channel, respectively.[35] We notice that BCCRs have characteristic

well-conserved acidic amino acids on the C-terminal side of TM7 (Fig. S1). We thus hypothesized

that a potential proton exchange between the third motif residue (D) in TM3 and acidic amino acid

residues at the cytoplasmic side of TM7 in BCCRs plays a critical role in the channel gating. To

test this hypothesis, we generated mutants corresponding to these residues in HulaCCR: D187N in

TM3, and E319Q, D320N, and E322Q, in TM7, and analyzed their photocurrents (Figs. S3a–c).

The D187N and D320N mutants exhibited low expressions in the plasma membrane probably due

to a significant decrease in protein stability, resulting in no observable photocurrents (Fig. S3c).

Whereas E319Q exhibited photocurrent intensity and opening/closing rates similar to that of the

WT (Figs. S3b and c), a decrease in the closing rate after 200-ms illumination was indeed observed

for E322Q (17 ± 2 ms, 15 ± 1 ms, and 38 ± 2 ms for the WT, E319Q, and E322Q at -60 mV holding

potential, respectively, $n = 5$, Fig. S3b).

Next we compared photocurrent signals upon nanosecond laser-pulse illumination to precisely

determine the difference in the time constants of opening and closing (Figs. S3d and e).

Additionally, to clarify the relationship between the proton transfer reaction and channel dynamics,

the kinetic isotope effect (KIE) in heavy water (D_2O) was investigated by calculating a ratio

between the time constants with D_2O and H_2O solutions with respect to the channel opening and

closing kinetics. If a proton transfer is the rate-limiting elementary reaction step, the reaction rate

decreases in D_2O, thereby exhibiting a significant KIE value.[37–39] For the HulaCCR WT and

E322Q, the time courses of photocurrents at membrane potentials of -60 mV and 40 mV are shown

in Fig. S3d. The photocurrent kinetic traces of E322Q were fitted with a quadruple-exponential

function, similar to the WT, to determine the time constants of channel opening and closing (Fig.

S3e). By calculating the ratio of closing time constants in D_2O to those in H_2O, the KIE was

determined to be 1.61 ± 0.06 for the WT and 1.18 ± 0.03 for E322Q, respectively. Hence, E322Q

showed a smaller KIE for channel closing compared to that of the WT. Unfortunately, information

about D187, which is homologous to *Hs*BR D96 and thought to be involved in the proton transfer

with E322, could not be obtained due to the lack of significant photocurrents for D187N. However,

considering the proton transfer mechanism proposed in the previous study for *Gt*CCR2, where

protonation of the cytoplasmic D in TM3 controls channel closing, we hypothesized that a similar

mechanism might occur in HulaCCR and the proton transfer from E322 to D187 governs the

channel closing there as well.

Long N/C-terminal extension of HulaCCR

One of the prominent structural features of HulaCCR is the long N-terminal extension. A careful

comparison of HulaCCR to other BCCRs, revealed that the first 12 residues of the expressed

construct might not belong to the ORF, which still leaves 79 residues before the first helix of the

rhodopsin domain, which exceeds the length of this region in any other BCCRs. The N-terminus

harbors a distinctive amino acid sequence, which is predicted as signal peptides by signal peptide

prediction methods. The exact cleavage site varied depending on the software used: SignalP 6.0

[40] predicted it to be between residues 43 and 44, while TOPCONS[41] predicted the cleavage

site between 30 and 31. Similar signal peptides were found in other members of the mgCCR1–3

clade. To investigate the role of the N-terminal extension in HulaCCR, two truncated variants were

constructed: variant I without the predicted signal sequence region (residues 41–428) and variant

II lacking most of the N-terminal region (residues 72–428).

No stop codon was observed in the 1285-nt contig encoding HulaCCR, yet the 108 residues

of the C-terminal extension were available for analysis. Similarly to other BCCRs, no homology

could be detected to known functional domains in the C-terminal extension of HulaCCR, but the

structure modeled using AlphaFold2 exhibited two additional helices (Fig. 4a). Based on the

structural model, two C-terminal truncations: variants III and IV, were produced by truncating

immediately after the predicted TM7 helix (residues 1–341) and after the additional predicted

helices after TM7 (residues 1–372), respectively. Patch-clamp measurements were conducted for

these four truncated variants (I–IV).

Based on the fluorescence images (Fig. 4b), truncated variant I (residues 41–428) exhibited

significantly decreased plasma membrane trafficking in cells. When cells showing strong

fluorescence were used, no significant photocurrent was obtained. However, significant

photocurrents were observed with cells showing relatively lower fluorescence intensity, indicating

that mild protein expression improves membrane trafficking of the protein (Fig. 4c). The shape of

photocurrents of the truncated variant I did not differ from that of the full-length protein. Hence,

the N-terminal extension appears to actually function as a signal peptide enhancing the membrane

trafficking of HulaCCR. Intriguingly, truncated variant II, which has a further truncated N-terminus

(residues 72–428), membrane translocation was restored, showing an expression pattern and

photocurrents similar to those of the full-length protein. We conclude that the region 41–71 between

the predicted secretion signal and the rhodopsin domain appears to reduce the localization in the

plasma membrane. However, the decreased membrane localization is compensated by the secretion

signal peptide (residues 1–41) in the full-length protein. The region spanning the residues 41–71 is

predicted by AlphaFold2 to have a disordered structure and is not conserved even among mgCCR1-

like ChRs.

For the C-terminal truncated variants, truncated variant III, which is composed of the residues

1–341 and truncated immediately after TM7, showed an increase in fluorescence intensity inside

the cells (Fig. 4b), indicating a decreased membrane localization compared to that of the full-length

HulaCCR. Truncated variant IV, in which a predicted disordered region beyond an additional

cytoplasmic helix was removed, exhibited higher membrane localization than that of truncated

variant III. The photocurrents of truncated variants III and IV were similar to that of the full-length

HulaCCR, indicating that the C-terminal extended region is not as essential as the N-terminal one

(Fig. 4c).

HulaCCR is able to drive neuronal firing

Our findings suggest that HulaCCR exhibits cation conductance and ion selectivity similar to those

of *Gt*CCR4 and ChRmine, indicating its potential as an optogenetic tool to induce neuronal firing

using light as other CCRs. To assess the practical utility of HulaCCR as an optogenetic tool,

primary cultures of rat hippocampal neurons were transfected with HulaCCR, and its expression

was confirmed based on the fluorescence signal (Fig. 5a). HulaCCR was well expressed throughout

the neuronal cells, including the axon. Patch-clamp recordings were performed on neurons

expressing HulaCCR (Fig. 5b), and upon 200 ms light stimulation, neuronal firings were elicited.

Additionally, even with 5 ms light stimulation at 10 Hz, continuous neuronal firing was observed,

demonstrating HulaCCR's ability to control neuronal activity through light stimulation. This is

likely facilitated by the relatively high conductance of HulaCCR to sodium.

Discussion

In this study, HulaCCR was identified as a new CCR featuring the ETD motif. Considering cation

selectivity, P_{Na+}/P_{H+} was previously estimated for *Cr*ChR2 and several other CCRs. While

HulaCCR exhibits a P_{Na+}/P_{H+} value of $(1.5 \pm 0.1) \times 10^{-5}$, P_{Na+}/P_{H+} of *Cr*ChR2 is approximately 1

$\times 10^{-6}$.[8] *Gt*CCR4, with a value of about 5×10^{-5}, has been reported as a CCR with low proton

selectivity,[27] suggesting that HulaCCR also has relatively low proton selectivity like *Gt*CCR4

among CCRs. The P_{X+}/P_{H+} values obtained for HulaCCR do not differ significantly between the

different alkali-metal monovalent cations, indicating similar permeability levels for all of them.

The photocurrents derived from Cs^+ in HulaCCR do not significantly differ from that of Na^+,

whereas the photocurrents from Cs^+ are approximately half of that from Na^+ in GtCCR4.[27] Hence,

HulaCCR likely possesses a relatively large channel pore facilitating the transport of large

monovalent cations. This is further supported by the fact Li^+, which is strongly hydrated in

solution,[42] showed permeability comparable to other monovalent cations. Considering the

expected large pore size capable of efficient Cs^+ transport, it is likely that partially hydrated Li^+ is

stabilized within the pore rather than undergoing complete dehydration. mgCCR1[23] despite

sharing the ETD motif with HulaCCR and having a 48% amino acid sequence identity, exhibits

relatively high proton selectivity compared to HulaCCR. The significant difference in ion

selectivity despite the relatively high amino acid sequence identity is intriguing. A comparison of

predicted structures using AlphaFold2 and cavities potentially involved in channel gating did not

suggest significant differences in the size of channel pore (Fig. S4). Hence, although it is anticipated

that the structural changes in the open state of HulaCCR are larger than that in mgCCR1, leading

an opening of a larger channel that is capable of conducting large monovalent cations, further

investigation is needed to clarify the reason of different cation selectivity between HulaCCR and

mgCCR1.

Considering the role of the first motif residue E, it was found to influence the absorption

spectrum and gating (Figs. 3b–d). In the E176D mutant, the time constants of channel opening and

closing were altered compared to the WT, and regarding the fast channel closing component, the

closing time constant was 1.8 and 1.3 times slower than that of the wild type at a +40 mV and a

−60 mV membrane potential, respectively, indicating the difference in the closing rate between

HulaCCR E176D and the WT becomes larger at positive membrane potential. Hence, in the WT,

E187 appears to enhance inward rectification. Indeed, when observing the photocurrent upon 200

ms illumination, the positive photocurrent intensity in the WT is slightly suppressed compared to

E176D (Figure S2d). This suggests that the glutamate is one of the important residues defining the

rectification properties of HulaCCR.

Looking at the photocurrents of the C303A mutant involving the residue immediately before

the retinal-binding lysine residue in TM7 at 200 ms light illumination, we find that the decay rate

after light off is significantly slower compared to that of the WT (WT = 17 ± 2 ms, C303A = 130

± 20 ms, at -60 mV, $n = 5$) (Fig. S2d). GtACR1, a member of the cryptophyte ACR family, also

has a cysteine at this position, C237, and an analogous mutation, C237A, was reported to eliminate

one of the two closing components and to show approximately 1.8 times slower closing rate for the

remaining component.[43] However, the retardation effect of the corresponding mutation on the

closing rate is much higher in HulaCCR (7.6 times slower), indicating that C303 is likely to

influence the closing rate through a mechanism different from that of GtACR1. Since C303 is

located close to the retinal Schiff base, the mutation of this residue may affect the photochemical

reactions around the chromophore. Nevertheless, we observed no significant spectral shift between

HulaCCR C303A and the WT based on the action spectra of the photocurrent, contrasting with the

red shift in C237A of GtACR1[43] (Fig. S2c). Hence, the interaction between the retinal

chromophore and C303 in HulaCCR is different from that in GtACR1, and the effect on the closing

process differs even with the identical mutation.

The N-terminal extension, outside the seven TM helices constituting the rhodopsin domain,

contains a sequence predicted as a secretion signal sequence, and our results suggest that this

secretion signal sequence rescues the decrease in membrane translocation caused by the sequence

between residues 41 and 71. mgCCR1 has a similarly long N-terminal extension, although it is

shorter and demonstrates low homology to that of HulaCCR. Analogously, the N-terminal

extension of mgCCR1 is predicted to have signal peptide cleaved between positions 24 and 25

(according to SignalP 6.0). Hence, it is anticipated that a similar mechanism maintains protein

membrane translocation in mgCCR1 as well. Regarding the extension at the C-terminus, although

not as prominent as the N-terminus, a slight decrease in membrane translocation compared to the

full-length protein was observed in variant III (residues 1–341) (Fig. 4b). Hence, the C-terminal

region may contribute to maintaining the stability of proteins in the membrane by interacting with

the cytoplasmic surface of the rhodopsin part. The long C-terminal extensions of green algal CCRs,

prasinophyte ACRs and labyrinthulean ACRs have been found to contain response regulator-like

domains[44] as well as additional conserved domains of unknown function,[44,45] but we could

not identify any such functional domains in HulaCCR or other BCCRs. It has been previously

reported that multiple arginines within the C-terminal extension interact with the rhodopsin part

and regulate the closing rate.[46] HulaCCR also has several arginines between residues 352 and

358 that were truncated in variant III. Moreover, a slightly slower decay rate after light off was

observed in variant III compared to that of the WT upon 200 ms light illumination (WT = 17 ± 2

ms, variant III = 21 ± 1 ms, at -60 mV, $n = 5$), exhibiting behavior similar to KnChR. Hence, it is

conceivable that in structural stabilization in the membrane is facilitated by electrostatic

interactions between the C-terminal extension and the cytoplasmic side of the rhodopsin domain

mediated by the arginines, potentially influencing gating kinetics.

Figures

Fig. 1. Characterization of HulaCCR

a Amino acid alignment of TM3 of HulaCCR with *Hc*KCR1, ChRmine, *Gt*CCR4, *Cr*ChR2, and

*Hs*BR. **b** (left) The phylogenetic tree of known ion-pumping rhodopsins, channelrhodopsins, and

HulaCCR. The phylogenetic tree was inferred using the neighbor-joining method of MEGA X

software (version 11.0.13). (right) Schematic illustration of HulaCCR structure based on

AlphaFold2 prediction. **c** Difference UV–visible absorption spectra showing the λ_{max} of HulaCCR.

The difference spectra were measured upon the hydrolysis reaction of the retinal Schiff-base

linkage induced by hydroxylamine with visible light illumination (0, 1, or 2 min). **d** Fluorescent

image of ND7/23 cells expressing HulaCCR which is C-terminally labeled with EYFP (Scale bar

= 20 μm). **e** Photocurrents of HulaCCR at 142 mM $[Na^+]_{out}$ and 128 mM $[K^+]_{in}$ at pH = 7.4 (left)

 and at 146 mM [NMG$^+$]$_{out}$ and 130 mM [NMG$^+$]$_{in}$ at pH = 7.4 (right).

Fig. 2. Ion selectivity of HulaCCR

a Representative photocurrent traces of HulaCCR at 146 mM [X$^+$]$_{out}$ and 130 mM [NMG$^+$]$_{in}$ (top),

and at 70 mM [Mg^{2+}]$_{out}$ /72.5 mM [Ca^{2+}]$_{out}$ /146 mM [NMG$^+$]$_{out}$ and 130 mM [NMG$^+$]$_{in}$ (bottom).

 b The *I–V* plot of HulaCCR photocurrents with different extracellular cations (mean ± S.E., *n* = 6).

 d Difference in E_{rev} between different cations and H$^+$ (ΔE_{rev}) (left) and relative permeability of

 different cations against H$^+$ (P_{X+}/P_{H+}) (right) (mean ± S.E., *n* = 6).

	Open$_{fast}$ /ms (contribution)	Open$_{slow}$ /ms (contribution)	Close$_{fast}$ /ms (contribution)	Close$_{slow}$ /ms (contribution)
WT 40mV	0.40 ± 0.01 (93.6 $_{± 0.4}$%)	3.6 ± 0.2 (6.4 $_{± 0.4}$%)	14.6 ± 0.1 (96.0 $_{± 0.4}$%)	42 ± 2 (4.0 $_{± 0.4}$%)
E176D 40 mV	0.33 ± 0.01 (89.7 $_{± 0.5}$%)	7.3 ± 0.3 (10.3 $_{± 0.5}$%)	26.8 ± 0.1 (99.2 $_{± 0.1}$%)	300 ± 40 (0.8 $_{± 0.1}$%)
WT −60mV	0.34 ± 0.01 (87 $_{± 1}$%)	4.4 ± 0.4 (13 $_{± 1}$%)	15.8 ± 0.5 (87$_{±1}$%)	47 ± 2 (13$_{±1}$%)
E176D −60 mV	0.26 ± 0.01 (87.5 $_{± 0.5}$%)	8.4 ± 0.4 (12.5 $_{± 0.5}$%)	20.4 ± 0.1 (99.5 $_{± 0.1}$%)	200 ± 20 (0.5 $_{± 0.1}$%)

 Fig. 3. Effect of point mutation

 a AlphaFold2 structural model for HulaCCR showing the mutated residues. The retinal structure

 (orange) and the side chain of conjugated lysine are of *Hc*KCR1 (PDB ID: 8H86). **b** Normalized

 action spectra of the HulaCCR WT and E176D (mean ± S.E., *n* = 5). **c** Photocurrent traces of the

 WT and E176D upon nanosecond laser flash excitation at holding potentials of 40 mV (positive)

 and −60 mV (negative). Photocurrents were recorded at 133 mM [Na$^+$]$_{out}$ and 120 mM [Na$^+$]$_{in}$ at

pH = 7.4. Fitting curves are shown as broken lines. **d** Time constants of channel opening and closing of the WT HulaCCR and E176D (mean ± S.E., *n* = 9). **e** Photocurrent traces of the HulaCCR WT, C181A, and C181S upon nanosecond laser flash excitation recorded with standard pipette and extracellular solutions. **f** Light-power dependency of the photocurrent of HulaCCR WT, C303A, and ChRmine at −40 mV with standard pipette and extracellular solutions (mean ± S.E., *n* = 5).

Fig. 4. Long N/C-terminal extensions of HulaCCR

a Truncated sites shown in the AlphaFold2 structural model of HulaCCR. **b** Fluorescent images of truncated variants expressed in ND7/23 cells. Scale bars, 20 μm. **c** Representative photocurrent traces of truncated variants recorded with standard pipette and extracellular solutions.

Fig. 5. Light stimulation of hippocampal neurons expressing HulaCCR

a Fluorescent images of cell body (top) and axon (bottom) of a hippocampal neuron expressing HulaCCR. **b** Voltage traces showing depolarization and spikes of the neuron in response to 200 ms green (542 nm) light stimulation (left) and to 10 Hz stimulation with 5 ms pulses (right).

Materials and Methods

Sampling, RNA extraction, and Rhodopsin screening

Peat lake sampling was performed on February 23rd, 2021 in The Hula Nature Reserve, Israel (33°04'33.8"N 35°36'40.8"E). 20 L of water from the surface and 0.5 m depth was filtered through a 50 µm mesh net to remove large particles, filtered onto a GF/D filter (Whatman), and flash-frozen using liquid nitrogen on-site.

Total RNA was extracted from frozen filters, corresponding to ~5 L per sample (RNeasy PowerSoil Total RNA Kit; Qiagen). The RNA was tested with Bioanalyzer, and PolyA-selected libraries were constructed and sequenced at the Weizmann Institute of Science, on the NovaSeq platform (300 cycles, PE).

De-Novo assembly was done post trimming (Trimgalore; v. 0.6.6 [DOI 10.5281/zenodo.5127898], Cutadapt; v. 3.4[47] with MEGAHIT (v. 1.2.9)[48] and then clustered for redundancy (cd-hit-est; v. 4.8.1) [49]. Rhodopsin screening was done with HMMsearch (v. 3.3.2)[50] with the microbial rhodopsin profiles from https://github.com/BejaLab/RhodopsinProfiles.

DNA constructs

Coding sequence (Table S1) of full-length HulaCCR was cloned into pCMV3.0-enhanced fluorescent yellow protein (EYFP) vector between EcoRI and BamHI sites with a Kir2.1 membrane trafficking signal, EYFP, and an ER-export signal[51] fused at the C-terminus of HulaCCR. For point mutations, the QuikChange site-directed mutagenesis method (Agilent Technologies, CA) was employed according to a manufacturer standard protocol. The truncated mutants were PCR-based cloned using EcoRI-BamHI restriction sites into the pCMV3.0-EYFP vector. Primers used in DNA work are listed in Table S2.

UV–visible spectroscopy

Cell culture and transfection for UV–visible spectroscopy were conducted as described

elsewhere.[52] Briefly, COS-1 cells (cell number: JCRB9082; Japanese Collection of Research

Bioresources Cell Bank, Osaka, Japan) were transfected by the polyethyleneimine method.[53]

Cells were supplemented with 2.5 μM all-*trans*-retinal (ATR) on the following day and harvested

after 48 h of transfection. The cell membranes were solubilized with a phosphate buffer (pH 7.0)

containing (in mM) 66.5 phosphate, 133 NaCl with 3% *n*-dodecyl-β-D-maltoside. After

centrifugation ($21,600 \times g$, 20°C, 5 min), the supernatant was subjected for the measurement of

UV–visible absorption spectra using a spectrophotometer (V-730, JASCO, Japan). Rhodopsin was

bleached by hydroxylamine (50 mM) and illumination with visible light from a 1-kW Xe lamp

(MAX-303, Asahi Spectra, Japan) through a long-pass filter (Y52, AGC Techno Glass, Japan).

Difference spectra were calculated by subtracting post-illumination spectra from the pre-

illumination spectrum. To correct for spectral changes derived from non-rhodopsin components in

the sample, the corresponding spectra of untransfected cell membranes that underwent the same

treatment as the sample were subtracted from the sample spectra.

Cell culture and transfection for electrophysiology

ND7/23 cells were grown in Dulbecco's modified Eagle's medium (D-MEM, FUJIFILM Wako

Pure Chemical Co., Japan) supplemented with 5% fetal bovine serum (FBS) under a 5% CO_2

atmosphere at 37°C. ND7/23 cells were attached onto a collagen-coated 12-mm coverslips (IWAKI,

cat. 4912-010, Japan) placing in a 4-well cell culture plate (SPL Life Sciences, cat. 30004, Korea).

The expression plasmids were transiently transfected in ND7/23 cells using Lipofectamine 3000

transfection reagent (Thermo Fisher Scientific Inc., MA). Seven to eight hours after the transfection,

the medium was replaced with D-MEM containing 10% horse serum (New Zealand origin, Thermo

Fisher Scientific Inc., MA), 50 ng mL^{-1} nerve growth factor-7S (Sigma-Aldrich, MO), 1 mM N^6,2'-

O-dibutyryladenosine-3',5'-cyclic monophosphate sodium salt (Nacalai tesque, Japan), 1 μM

cytosine-1-β-D(+)-arabinofuranoside (FUJIFILM Wako Pure Chemical Co., Japan), and 2.5 μM

all-*trans*-retinal. Electrophysiological recordings were conducted at 1–3 days after the transfection.

The transfected cells were identified by observing the EYFP fluorescence under an up-right

microscope (BX50WI, Olympus, Japan).

Hippocampi were isolated from postnatal day 1 (P1) Wistar-ST rats (by Sankyo Labo

Service Co., Japan) and treated with papain dissociation system (Worthington Biochemical Co.,

NJ) according to the company protocol. Briefly, dissected hippocampi were treated with papain for

1 h at 37°C. The dissociated cells were washed with Earle's Balanced Salt Solution (EBSS)

supplemented with ovomucoid protease inhibitor with bovine serum albumin. The dissociated cells

were resuspended in culture medium (Neurobasal A containing 2% B-27 plus (Thermo Fisher

Scientific Inc., MA)). Approximately 150,000 cells were plated on 12-mm poly-L-lysine-coated

glass coverslips (IWAKI, cat. 4912-040, Japan) in the 4-well cell culture plate. Transfection was

done by adeno-associated virus (AAV), and the plasmid and virus packaging were synthesized by

Vectorbuilder Inc. The plasmid pAAV vector has a CaMKIIa promotor containing the sequence of

HulaCCR, C-terminally fused to the Kir2.1 membrane trafficking signal followed by EYFP and

the ER export signal. For transfection, AAV8 was added to each well (4–8 × 10^8 genome copies

well^{-1}) 7 days after plating. The electrophysiological recordings were performed 15–22 days after

transfection.

Electrophysiology

All experiments were carried out at room temperature (20–22°C). Currents were recorded using an

EPC-8 amplifier (HEKA Electronic, Germany) under a whole-cell patch clamp configuration. The

data were filtered at 1 kHz, sampled at 50 kHz (Digidata1440 A/D, Molecular Devices Co., CA)

and stored in a computer (pClamp11.1, Molecular Devices). The standard internal pipette solutions

for the whole-cell voltage clamp recordings from the ND7/23 cells contained (in mM) 119 KF, 10

KOH, 5 Na_2EGTA, 14 HEPES, 0.0025 ATR (pH 7.4 adjusted with HCl). The standard extracellular

solution contained (in mM): 138 NaCl, 3 KCl, 2.5 $CaCl_2$, 1 $MgCl_2$, 10 HEPES, 4 NaOH, and 11

glucose (pH 7.4 adjusted with HCl). In the ion selectivity measurement, extracellular solution

contained (in mM) 146 XCl ($X = Na^+$, K^+, Li^+, Rb^+, Cs^+), 6 N methyl-D-glucamine (NMG), 2.5

$CaCl_2$, 10 HEPES, 11 glucose (pH 7.4 adjusted with HCl). For the divalent cation selectivity,

extracellular solution contained (in mM) 70 XCl_2 ($X = Ca^{2+}$, Mg^{2+}), 6 NMG, 2.5 $CaCl_2$, 10 HEPES,

11 glucose (pH 7.4 adjusted with HCl). The NMG-extracellular solution contained (mM) 146 NMG,

146 HCl, 2.5 $CaCl_2$, 10 HEPES, and 11 glucose (pH 7.4 adjusted with NMG). The NMG-pipette

solution contained (in mM) 130 NMG, 90 glutamate, 6 $(NMG)_2EGTA$, 50 HEPES, 2.5 $MgSO_4$, 2.5

MgATP, 0.0025 ATR (pH 7.4 adjusted with H_2SO_4). The liquid junction potentials (LJPs) were

calculated by pClamp 11.1 software (Table. S3) and estimated reversal potentials were

compensated by calculated LJPs.

For whole-cell voltage clamp, illumination at 377 ± 25, 438 ± 12, 472 ± 15, 510 ± 5, 542

± 13, 575 ± 12 or 643 ± 10 nm was carried out using a SpectraX light engine (Lumencor Inc., OR)

controlled by pClamp 11.1 software. HulaCCR was illuminated through an objective lens

(LUMPlan FL 40x, NA 0.80W, Olympus, Japan). For the standard photocurrent detection, the

power of the 542-nm light was directly measured under a microscope using a visible light-sensing

thermopile (MIR178 101Q, SSC Co., Ltd., Japan) and was adjusted to 13.3 mW mm^{-2}. The action

spectrum was measured at a holding potential of -40 mV at wavelengths with equivalent power

density of 0.8 mW mm^{-2}. Each action spectrum was estimated by the maximal amplitude of the

photocurrent scaled by the light power density under the assumption of a linear relationship

between the photocurrent intensity and the light power density.

For Laser-flash patch clamp experiment, a laser flash (3–5 ns) at 532 nm (Nd:YAG laser, Minilite II, Continuum, CA) was used.[39] For measuring the KIE on the gating dynamics, the internal pipette solutions were made of H_2O or D_2O solutions containing (in mM) 60 Na_2SO_4, 25 NMG, 60 mannitol, 5 EGTA, 10 HEPES, 2.5 $MgSO_4$, 2.5 MgATP, and 0.0025 ATR adjusting pH to 7.4 with HCl or DCl, respectively. The extracellular solutions were made of H_2O or D_2O solutions containing (in mM) 133 NaCl, 3 KCl, 5 NMG, 10 HEPES, 5 $CaCO_3$ (solved by HCl or DCl solutions), 2.5 $MgSO_4$ and 11 glucose, adjusting pH to 7.4 with HCl or DCl, respectively. The pipette resistance was adjusted to 3–5 $M\Omega$ with a series resistance of 7–19 $M\Omega$ (WT; 12 ± 1, $n = 9$ for H_2O; 16 ± 1, $n = 9$ for D_2O, E176D; 9 ± 1, $n = 9$ for H_2O, E322Q; 10 ± 1, $n = 9$ for H_2O; 15 ± 1, $n = 9$ for D_2O) and a cell capacitance of 30–90 pF (WT; 63 ± 4, $n = 9$ for H_2O; 58 ± 3, $n = 9$ for D_2O, E176D; 50 ± 2, $n = 9$ for H_2O, E322Q; 43 ± 3, $n = 9$ for H_2O; 44 ± 2, $n = 9$ for D_2O). In every experiment, the series resistance was compensated for by 70%. Obtained signals had electrical noise from the laser system. Therefore, the noise signal recorded without light excitation was subtracted from raw photocurrent traces.

For whole-cell recordings in cultured hippocampal neurons, patch pipettes were filled with (in mM) 90 potassium gluconate, 30 KOH, 49 HEPES, 2.5 $MgCl_2$, 2.5 MgATP, 5 Na_2EGTA, and 0.0025 ATR, titrated to pH 7.4. The extracellular solution contained (in mM) 138 NaCl, 3 KCl, 2.5 $CaCl_2$, 1 $MgCl_2$, 11 glucose, and 10 HEPES, 2 kynurenic acid, titrated to pH 7.4.

Acknowledgements

We thank Ariel Chazan, Shirley Larom, David Iluz, Said Abu-Ghosh, and the park rangers David Zaguri and Ido Shaked for help in sampling, and the Israel Nature and Parks Authority for the permit to sample in The Hula Nature Reserve. This work was supported by JSPS KAKENHI Grants-in-Aid (grant JP 23K14151 to S.T., JP 21H01875, JP23K18090, and JP23H04404 to K.I.),

JST CERST (grant JPMJCR22N2 to K.I.), and MEXT Promotion of Development of a Joint Usage/

Research System Project: Coalition of Universities for Research Excellence Program (CURE)

(grant JPMXP1323015482 to K.I.), the European Commission, under Horizon Europe's research

and innovation programme (Bluetools project, Grant Agreement No 101081957 to O.B.), the Israel

Science Foundation (Research Center grant 3131/20 to O.B.), and the Nancy and Stephen Grand

Technion Energy Program (GTEP). O.B. holds the Louis and Lyra Richmond Chair in Life

Sciences.

References

[1] A. Rozenberg, K. Inoue, H. Kandori, O. Béjà, Microbial rhodopsins: The last two decades, Annu. Rev. Microbiol. 75 (2021) 427–447.

[2] O.P. Ernst, D.T. Lodowski, M. Elstner, P. Hegemann, L.S. Brown, H. Kandori, Microbial and animal rhodopsins: structures, functions, and molecular mechanisms, Chem. Rev. 114 (2014) 126–163.

[3] T. Nagata, K. Inoue, Rhodopsins at a glance, J. Cell Sci. 134 (2021) jcs258989.

[4] K. Deisseroth, P. Hegemann, The form and function of channelrhodopsin, Science 357 (2017) eaan5544.

[5] V. Emiliani, E. Entcheva, R. Hedrich, P. Hegemann, K.R. Konrad, C. Lüscher, M. Mahn, Z.-H. Pan, R.R. Sims, J. Vierock, O. Yizhar, Optogenetics for light control of biological systems, Nat. Rev. Methods Primers 2 (2022) 55.

[6] O.A. Sineshchekov, K.-H. Jung, J.L. Spudich, Two rhodopsins mediate phototaxis to low- and high-intensity light in *Chlamydomonas reinhardtii*, Proc. Natl. Acad. Sci. U. S. A. 99 (2002) 8689–8694.

[7] G. Nagel, D. Ollig, M. Fuhrmann, S. Kateriya, A.M. Musti, E. Bamberg, P. Hegemann, Channelrhodopsin-1: a light-gated proton channel in green algae, Science 296 (2002) 2395–2398.

[8] G. Nagel, T. Szellas, W. Huhn, S. Kateriya, N. Adeishvili, P. Berthold, D. Ollig, P. Hegemann, E. Bamberg, Channelrhodopsin-2, a directly light-gated cation-selective membrane channel, Proc. Natl. Acad. Sci. U. S. A. 100 (2003) 13940–13945.

[9] E.G. Govorunova, O.A. Sineshchekov, R. Janz, X. Liu, J.L. Spudich, Natural light-gated anion channels: A family of microbial rhodopsins for advanced optogenetics, Science 349 (2015) 647–650.

[10] E.G. Govorunova, O.A. Sineshchekov, J.L. Spudich, Structurally distinct cation channelrhodopsins from cryptophyte algae, Biophys. J. 110 (2016) 2302–2304.

[11] H. Hagio, W. Koyama, S. Hosaka, A.D. Song, J. Narantsatsral, K. Matsuda, T. Shimizu, S. Hososhima, S.P. Tsunoda, H. Kandori, M. Hibi, Optogenetic manipulation of neuronal and cardiomyocyte functions in zebrafish using microbial rhodopsins and adenylyl cyclases, eLife 12 (2023) e83975.

[12] M. Mahn, L. Gibor, P. Patil, K. Cohen-Kashi Malina, S. Oring, Y. Printz, R. Levy, I. Lampl, O. Yizhar, High-efficiency optogenetic silencing with soma-targeted anion-conducting channelrhodopsins, Nat. Commun. 9 (2018) 4125.

[13] S. Huang, M. Ding, M.R.G. Roelfsema, I. Dreyer, S. Scherzer, K.A.S. Al-Rasheid, S. Gao, G. Nagel, R. Hedrich, K.R. Konrad, Optogenetic control of the guard cell membrane potential and stomatal movement by the light-gated anion channel *Gt*ACR1, Sci. Adv. 7 (2021) eabg4619.

[14] Y. Zhou, M. Ding, S. Gao, J. Yu-Strzelczyk, M. Krischke, X. Duan, J. Leide, M. Riederer, M.J. Mueller, R. Hedrich, K.R. Konrad, G. Nagel, Optogenetic control of plant growth by a microbial rhodopsin, Nat. Plants 7 (2021) 144–151.

[15] H. Kandori, Ion-pumping microbial rhodopsins, Front. Mol. Biosci. 2 (2015) 52.

[16] E.G. Govorunova, O.A. Sineshchekov, J.L. Spudich, Structurally distinct cation channelrhodopsins from cryptophyte algae, Biophys. J. 110 (2016) 2302–2304.

[17] Y. Yamauchi, M. Konno, S. Ito, S.P. Tsunoda, K. Inoue, H. Kandori, Molecular properties of a DTD channelrhodopsin from *Guillardia theta*, Biophys Physicobiol. 14 (2017) 57–66.

[18] J.H. Marshel, Y.S. Kim, T.A. Machado, S. Quirin, B. Benson, J. Kadmon, C. Raja, A. Chibukhchyan, C. Ramakrishnan, M. Inoue, J.C. Shane, D.J. McKnight, S. Yoshizawa, H.E.

Kato, S. Ganguli, K. Deisseroth, Cortical layer-specific critical dynamics triggering perception, Science 365 (2019) eaaw5202.

[19] R. Chen, F. Gore, Q.-A. Nguyen, C. Ramakrishnan, S. Patel, S.H. Kim, M. Raffiee, Y.S. Kim, B. Hsueh, E. Krook-Magnusson, I. Soltesz, K. Deisseroth, Deep brain optogenetics without intracranial surgery, Nat. Biotechnol. 39 (2021) 161–164.

[20] H.E. Kato, Structure–Function Relationship of Channelrhodopsins, in: H. Yawo, H. Kandori, A. Koizumi, R. Kageyama (Eds.), Optogenetics: Light-sensing proteins and their applications in neuroscience and beyond, Springer Singapore, Singapore, 2021: pp. 35–53.

[21] E.G. Govorunova, Y. Gou, O.A. Sineshchekov, H. Li, X. Lu, Y. Wang, L.S. Brown, F. St-Pierre, M. Xue, J.L. Spudich, Kalium channelrhodopsins are natural light-gated potassium channels that mediate optogenetic inhibition, Nat. Neurosci. 25 (2022) 967–974.

[22] J. Vierock, E. Shiewer, C. Grimm, A. Rozenberg, I.-W. Chen, L. Tillert, A.G. Castro Scalise, M. Casini, S. Augustin, D. Tanese, B.C. Forget, R. Peyronnet, F. Schneider-Warme, V. Emiliani, O. Béjà, P. Hegemann, WiChR, a highly potassium-selective channelrhodopsin for low-light one- and two-photon inhibition of excitable cells, Sci. Adv. 8 (2022) eadd7729.

[23] E.G. Govorunova, O.A. Sineshchekov, L.S. Brown, A.-N. Bondar, J.L. Spudich, Structural foundations of potassium selectivity in channelrhodopsins, MBio 13 (2022) e0303922.

[24] M. Mirdita, K. Schütze, Y. Moriwaki, L. Heo, S. Ovchinnikov, M. Steinegger, ColabFold: making protein folding accessible to all, Nat. Methods 19 (2022) 679–682.

[25] K. Inoue, M. Karasuyama, R. Nakamura, M. Konno, D. Yamada, K. Mannen, T. Nagata, Y. Inatsu, H. Yawo, K. Yura, O. Béjà, H. Kandori, I. Takeuchi, Exploration of natural red-shifted rhodopsins using a machine learning-based Bayesian experimental design, Commun. Biol. 4 (2021) 362.

[26] C.A. Lewis, Ion-concentration dependence of the reversal potential and the single channel conductance of ion channels at the frog neuromuscular junction, J. Physiol. 286 (1979) 417–445.

[27] S. Shigemura, S. Hososhima, H. Kandori, S.P. Tsunoda, Ion channel properties of a cation channelrhodopsin, *Gt*_CCR4, Appl. Sci. 9 (2019) 3440.

[28] K.E. Kishi, Y.S. Kim, M. Fukuda, M. Inoue, T. Kusakizako, P.Y. Wang, C. Ramakrishnan, E.F.X. Byrne, E. Thadhani, J.M. Paggi, T.E. Matsui, K. Yamashita, T. Nagata, M. Konno, S. Quirin, M. Lo, T. Benster, T. Uemura, K. Liu, M. Shibata, N. Nomura, S. Iwata, O. Nureki, R.O. Dror, K. Inoue, K. Deisseroth, H.E. Kato, Structural basis for channel conduction in the pump-like channelrhodopsin ChRmine, Cell 185 (2022) 672-689.e23.

[29] M. Nack, I. Radu, M. Gossing, C. Bamann, E. Bamberg, G.F. von Mollard, J. Heberle, The DC gate in Channelrhodopsin-2: crucial hydrogen bonding interaction between C128 and D156, Photochem. Photobiol. Sci. 9 (2010) 194–198.

[30] A. Berndt, O. Yizhar, L.A. Gunaydin, P. Hegemann, K. Deisseroth, Bi-stable neural state switches, Nat. Neurosci. 12 (2009) 229–234.

[31] E.G. Govorunova, O.A. Sineshchekov, H. Li, Y. Wang, L.S. Brown, J.L. Spudich, RubyACRs, nonalgal anion channelrhodopsins with highly red-shifted absorption, Proc. Natl. Acad. Sci. U. S. A. 117 (2020) 22833–22840.

[32] S.A. Tennigkeit, R. Karapinar, T. Rudack, M.-A. Dreier, P. Althoff, D. Eickelbeck, T. Surdin, M. Grömmke, M.D. Mark, K. Spoida, M. Lübben, U. Höweler, S. Herlitze, K. Gerwert, Design of an ultrafast G protein switch based on a mouse melanopsin variant, Chembiochem 20 (2019) 1766–1771.

[33] A.A. Shtyrov, D.M. Nikolaev, V.N. Mironov, A.V. Vasin, M.S. Panov, Y.S. Tveryanovich, M.N. Ryazantsev, Simple models to study spectral properties of microbial and animal

rhodopsins: Evaluation of the electrostatic effect of charged and polar residues on the first absorption band maxima, Int. J. Mol. Sci. 22 (2021) 3029.

[34] K. Stehfest, E. Ritter, A. Berndt, F. Bartl, P. Hegemann, The branched photocycle of the slow-cycling channelrhodopsin-2 mutant C128T, J. Mol. Biol. 398 (2010) 690–702.

[35] O.A. Sineshchekov, E.G. Govorunova, H. Li, J.L. Spudich, Bacteriorhodopsin-like channelrhodopsins: Alternative mechanism for control of cation conductance, Proc. Natl. Acad. Sci. U. S. A. 114 (2017) E9512–E9519.

[36] S. Tajima, Y.S. Kim, M. Fukuda, Y. Jo, P.Y. Wang, J.M. Paggi, M. Inoue, E.F.X. Byrne, K.E. Kishi, S. Nakamura, C. Ramakrishnan, S. Takaramoto, T. Nagata, M. Konno, M. Sugiura, K. Katayama, T.E. Matsui, K. Yamashita, S. Kim, H. Ikeda, J. Kim, H. Kandori, R.O. Dror, K. Inoue, K. Deisseroth, H.E. Kato, Structural basis for ion selectivity in potassium-selective channelrhodopsins, Cell 186 (2023) 4325–4344.

[37] L.S. Brown, R. Needleman, J.K. Lanyi, Origins of deuterium kinetic isotope effects on the proton transfers of the bacteriorhodopsin photocycle, Biochemistry 39 (2000) 938–945.

[38] K. Inoue, S. Tahara, Y. Kato, S. Takeuchi, T. Tahara, H. Kandori, Spectroscopic study of proton-transfer mechanism of inward proton-pump rhodopsin, *Parvularcula oceani* Xenorhodopsin, J. Phys. Chem. B 122 (2018) 6453–6461.

[39] K. Shibata, K. Oda, T. Nishizawa, Y. Hazama, R. Ono, S. Takaramoto, R. Bagherzadeh, H. Yawo, O. Nureki, K. Inoue, H. Akiyama, Twisting and protonation of retinal chromophore regulate channel gating of channelrhodopsin C1C2, J. Am. Chem. Soc. 145 (2023) 10779–10789.

[40] F. Teufel, J.J. Almagro Armenteros, A.R. Johansen, M.H. Gíslason, S.I. Pihl, K.D. Tsirigos, O. Winther, S. Brunak, G. von Heijne, H. Nielsen, SignalP 6.0 predicts all five types of signal peptides using protein language models, Nat. Biotechnol. 40 (2022) 1023–1025.

[41] K.D. Tsirigos, C. Peters, N. Shu, L. Käll, A. Elofsson, The TOPCONS web server for consensus prediction of membrane protein topology and signal peptides, Nucleic Acids Res. 43 (2015) W401-7.

[42] J. Mähler, I. Persson, A study of the hydration of the alkali metal ions in aqueous solution, Inorg. Chem. 51 (2012) 425–438.

[43] Y.S. Kim, H.E. Kato, K. Yamashita, S. Ito, K. Inoue, C. Ramakrishnan, L.E. Fenno, K.E. Evans, J.M. Paggi, R.O. Dror, H. Kandori, B.K. Kobilka, K. Deisseroth, Crystal structure of the natural anion-conducting channelrhodopsin *Gt*ACR1, Nature 561 (2018) 343–348.

[44] A. Rozenberg, J. Oppermann, J. Wietek, R.G. Fernandez Lahore, R.-A. Sandaa, G. Bratbak, P. Hegemann, O. Béjà, Lateral gene transfer of anion-conducting channelrhodopsins between green algae and giant viruses, Curr. Biol. 30 (2020) 4910-4920.e5.

[45] A. Kianianmomeni, K. Stehfest, G. Nematollahi, P. Hegemann, A. Hallmann, Channelrhodopsins of *Volvox carteri* are photochromic proteins that are specifically expressed in somatic cells under control of light, temperature, and the sex inducer, Plant Physiol. 151 (2009) 347–366.

[46] R. Tashiro, K. Sushmita, S. Hososhima, S. Sharma, S. Kateriya, H. Kandori, S.P. Tsunoda, Specific residues in the cytoplasmic domain modulate photocurrent kinetics of channelrhodopsin from *Klebsormidium nitens*, Commun. Biol. 4 (2021) 235.

[47] M. Martin, Cutadapt removes adapter sequences from high-throughput sequencing reads, EMBnet.Journal 17 (2011) 10–12.

[48] D. Li, C.-M. Liu, R. Luo, K. Sadakane, T.-W. Lam, MEGAHIT: an ultra-fast single-node solution for large and complex metagenomics assembly via succinct de Bruijn graph, Bioinformatics 31 (2015) 1674–1676.

[49] W. Li, A. Godzik, Cd-hit: a fast program for clustering and comparing large sets of protein or
nucleotide sequences, Bioinformatics 22 (2006) 1658–1659.
[50] L.S. Johnson, S.R. Eddy, E. Portugaly, Hidden Markov model speed heuristic and iterative
HMM search procedure, BMC Bioinformatics 11 (2010) 431.
[51] V. Gradinaru, F. Zhang, C. Ramakrishnan, J. Mattis, R. Prakash, I. Diester, I. Goshen, K.R.
Thompson, K. Deisseroth, Molecular and cellular approaches for diversifying and extending
optogenetics, Cell 141 (2010) 154–165.
675
676 [52] N. Morimoto, T. Nagata, K. Inoue, Reversible photoreaction of a retinal photoisomerase,
677   retinal G-protein-coupled receptor RGR, Biochemistry 62 (2023) 1429–1432.
678 [53] J.F. Fay, D.L. Farrens, Chapter fifteen - Purification of functional CB1 and analysis by site-
679   directed fluorescence labeling methods, in: P.H. Reggio (Ed.), Methods in Enzymology,
680   Academic Press, 2017: pp. 343–370.
681