Comparative Analysis of the Techniques for the Determination of Binding Affinity between a Small Molecule Inhibitor and a Protein Target

Yalan Luo, Yijun Chen*

Laboratory of Chemical Biology, School of Life Science and Technology,

China Pharmaceutical University, Nanjing, Jiangsu 211198, China

*Corresponding author:

Yijun Chen

Tel: +86-25-86185919

Email: yjchen@cpu.edu.cn;

Abstract

The binding affinity constant (K_D) between a small molecule inhibitor and a protein target is pivotal parameter for target identification and early drug discovery. Despite the extensive applications of three major techniques, namely SPR, ITC and MST, the K_D values for a specifically defined binding measured by these techniques are drastically different, which poses a remarkable difficulty to deal with these ambiguous data. Here, we report the evaluation of the accuracy of K_D values from SPR, ITC and MST compared with enzyme kinetics. To enable an objective comparison, we theoretically proved that enzyme competitive inhibition constant (K_i) could directly reflect the binding affinity. Using purine nucleoside phosphorylase, its substrate inosine and competitive inhibitor immucillin-H, we determined respective K_D and K_i values to make a direct comparison. Moreover, we found that the K_D value measured by SPR is more relevant to its K_i value. This study highlights the urgent need on the development of new technologies for the determination of binding affinity between small molecule inhibitors and protein targets.

Keywords: binding affinity, enzyme kinetics, SPR, ITC, MST, competitive inhibition

1. Introduction

As commonly known, binding affinity represents the strength of ligand-protein interaction, which quantifies the extent of protein occupancy. Generally, affinity calculation is based on following three methods: a. $\Delta G_{binding} = RTln(K_D)$, which relies on thermodynamics [1]; b. $K_D = \frac{k_{off}}{k_{on}}$, which depends on kinetics [2]; and c. $K_D = \frac{[R][L]}{[RL]}$, which is based on equilibrium dynamics [3]. These formulas are calculation basis of different detection methods for affinity, but they all principally rely on the binding affinity between dissociated components [2, 4]. When the interaction of a small molecule inhibitor and its target protein is concerned in the case of early drug discovery, it is preferable to accurately measure inhibitor-target affinity because it is an indicator to associate with the efficacy of inhibitors and their off-target effects. Consequently, as an important parameter, binding affinity constant (K_D) has been a valuable *in vitro* measure for various preclinical evaluation, including the selection of lead compounds, the optimization of leads, and the predictions of *in vivo* potency, pharmacokinetic properties and safely profiles [5]. Therefore, it is of great important to obtain accurate K_D values of the inhibitors.

Despite the complexity of biochemical reactions in all living processes, every type of physical and chemical transformations results in a series of changes in dynamics and thermodynamics. As a result, the emerging technologies for the determination of binding affinity between different molecules are literally designed to measure these changes [6]. Currently, there are three major techniques for this purpose, namely Microscale Thermophoresis (MST), Isothermal Titration Calorimetry

(ITC) and Surface Plasmon Resonance (SPR). MST determines binding affinity through the variation of thermophoresis of a fluorescent labeled molecule, and such a variation is usually caused by the changes of molecular properties, including size, charge, hydration state, and/or structural/conformational changes of the proteins, under constant buffering conditions upon the binding with small molecule inhibitors [7]. ITC focuses on the thermodynamics of molecular interaction where the contacts between two molecules produce exothermic or endothermic reaction to generate the information on association equilibrium constant, enthalpy change and stoichiometry from the binding [8]. The principle of SPR utilizes optical phenomena for detecting the association and dissociation of analytes on a biosensor chip by real-time monitoring [9]

In recent years, the binding affinity values and their measurement techniques have greatly contributed to accelerating the process of target identification and early drug discovery. Unfortunately, as a conventional wisdom and common practice, the choice of using which technique largely depends upon the availability of instrumentation rather than data reliability, resulting in unprecedented uncertainty on the reliability and/or reproducibility of K_D values for a pair of ligand-protein across different techniques [10]. Especially, for the interaction between high weight proteins and small chemical compounds (typically molecular weight < 500), the significant difference on molecular weight of the analytes leads to inconspicuous signal from the interaction, causing an exceptional discrepancy. For example, ABT -737 is one of the inhibitors of BCL-2, and its binding affinity to BCL-2 was determined by SPR and

ITC to be 0. 6 nM and 20.5 nM, respectively [11, 12]. The affinity of MCL-1 inhibitor A-1210477 was reported as ~740 nM by MST [13], whereas another study showed 3.5 nM by SPR [14], an over 200-fold difference. More intriguingly, using the same technique for the same interaction produces even larger difference, such as the case between PDL-1 and BMS-202 measured by MST to be 1.1 μM and 37 nM in two different investigations [15, 16]. Therefore, these obviously discrepant data have generated a serious question on the reliability of reported K_D values and their associated techniques.

When enzyme kinetics is concerned, the inhibition constant (K_i), particularly in the case of competitive inhibition, is extremely similar to K_D value to represent the binding affinity. K_i is a thermodynamics parameter, an important parameter and gold standard for the efficiency of inhibitors, and the true affinity between an inhibitor and an enzyme [17]. Previous studies have also verified that binding affinity and enzyme inhibition are closely correlated in thermodynamics, where K_i values strongly correlate with K_D values. Meanwhile, competitive K_i is quite close to K_D when the structures of substrate and inhibitor are remarkably similar [18]. Thus, we speculate that competitive K_i can serve as the reference to estimate K_D under consistent and reproducible conditions. In this study, we demonstrated the close correlation between competitive K_i and K_D by a comparative analysis, which could facilitate the development of more reliable technologies for K_D determination.

2. Methods and Materials

2.1 Expression and purification of human purine nucleoside phosphorylase (hPNP)

BL 21 (DE3) cells were transformed with pET28a (+)-hPNP plasmid and grown in 20 ml Luria-Bertani medium with kanamycin (50 μg/ml) overnight, then 2 ml culture was inoculated to 200 ml fresh LB medium. The diluted cultures were grown at 37℃ to approximately $OD_{600} = 0.6$ for the induction with IPTG (250 μM final concentration). After 20 h of additional growth at 20℃, cells (~1 g) were harvested by configuration at 4 ☐.

Approximately 12 g cells were suspended in 50 mM potassium phosphate buffer (pH 8.0) containing 0.1 mM PMSF. After ultrasonication and centrifugation at 4☐, the supernatant was incubated with Ni-NTA affinity resin for 1 h at 4 ☐. hPNP was eluted with 14 mL 50 mM potassium phosphate buffer (pH 8.0) containing 300 mM imidazole. Approximately 80 mg protein purified via Ni-affinity chromatography was loading on Q-High performance anion exchange chromatography column for further purification, in which the column was equilibrated with 50 mM HEPES (pH 8.0), and then hPNP was eluted with a linear gradient of 0-0.25 M NaCl. The purity of hPNP was examined by 12 % SDS-PAGE with Coomassie blue staining.

2.2 Enzyme assay and determination of competitive inhibition

The increase of uric acid was determined by SpectraMax iD5 instrument in a microtiter 96-well plate. The reaction mixture contained 1.4 nM hPNP, 25 μM to 5mM inosine as substrate and 0 -5 nM immucillin-H (ImmH) in the wells at 25 ☐. Michaelis-Menten equation was employed to obtain K_m values, and the equation

$v_i = \frac{V_{max}[S]}{\left(1+\frac{[I]}{K_i}\right)K_m+[S]}$ was used to generate K_i values. The data were analyzed by Graph

Pad Prism9.

2.3 Binding affinity assays

SPR experiments were performed with a Biacore T200 instrument in 50 mM

potassium phosphate buffer (pH 7.4) at 25 □. Purified hPNP was diluted to 20 μg/ml

in 10mM sodium acetate solution (pH 5.5) and immobilized on the sensor chip to

reach a final immobilization level of 7000 RU. ImmH was diluted to 1.2 - 925 nM

with flow rate of 30 μl/min and contact time of 120 s in each injection cycle.

ITC experiments were performed with a PEAQ-ITC instrument in 50 mM

potassium phosphate buffer (pH 7.4) at 25 □. The starting concentrations of hPNP

and ImmH were 5 μM and 150 μM. One Sit of Sites model was used with a setting of

Reference Power, FeedBack, Stir Speed, Initial Delay by default. In addition,

injections and the volume of each injection and spacing time were 15, 2.5 μl and 150 s

during each experiment.

For MST determinations, 200 nM hPNP was incubated with 100 nM fluorescent

tags for 30 min in His-Tag Labeling Kit RED-tris-NTA 2nd generation to reach

enough raw fluorescent counts (> 200). The concentration of ImmH was varied in a

dilution series starting at 1 mM, and then mixed with fluorescent-labeled hPNP in 50

mM potassium phosphate buffer (pH 7.4). After loading the samples into standard

capillaries, scanning was performed under 40 % MST power and 40% LED power at

25□.

3. Results

3.1 The relationship between enzyme competitive inhibition and biding affinity

Different types of reversible inhibitions in enzyme kinetics exhibit various relationships between K_m^{obs} and K_i (Scheme 1a) [19]. Among the inhibitions, competitive inhibition provides the simplest and an intuitive linear relationship:

$$K_m^{obs} = K_m + \frac{K_m}{K_i}[I] \qquad (4)$$

In competitive inhibition, a substrate competes with an inhibitor to bind the active site of the enzyme. There are three equilibriums in competitive inhibition (Scheme 1b):

$$K_m = \frac{[E][S]}{[ES]} = \frac{k_2}{k_1} \text{ and } K_i = \frac{[E][I]}{[EI]} = \frac{k_4}{k_3} \qquad (5)$$

$$\frac{K_m}{K_i} = \frac{k_2 k_3}{k_1 k_4} \qquad (6)$$

Apparent Michaelis constant K_m^{obs} contains two equilibriums and can be calculated in accordance with Eqs. (1) and (3). Therefore, apparent inhibition constant K_i^{obs} could be derived from Eq. (2) and reverse equilibrium of Eq. (3). Subsequent rearrangement generates Eqs. (7) and (8):

$$K_m^{obs} = K_m + K_m \frac{[EI]}{[E]} \qquad (7)$$

$$K_i^{obs} = K_i + K_i \frac{[ES]}{[E]} \qquad (8)$$

Apparently, the K_i value from theoretical calculation represents the initial binding state of an enzyme and an inhibitor, which is the true affinity to differentiate from apparent inhibition constant K_i^{obs}. Meanwhile, K_D denotes similar equilibrium equation and biochemical meaning as K_i based on $P + L \rightleftharpoons PL$ and $K_D = \frac{[P][L]}{[PL]} =$

$\frac{k_d}{k_a}$, in which the dissociation constant reflects ligand concentration at the point that half of the protein is occupied under usual conditions. Consequently, K_i values in enzyme competitive inhibitions are convertible to the K_D values of proteins and small molecules if the inhibitors are structurally similar to ligands.

3.2 The correlation of competitive inhibition and binding affinity

Since hPNP is a well-studied enzyme with known kinetic properties and enzyme inhibition [20], hPNP and its substrate inosine and competitive inhibitor immucillin-H (ImmH) were chosen as a model of competitive enzyme inhibition. Additionally, the computational simulation on theoretical dynamic information for K_m and K_D could also be served as reference data [21-22]. Thus, it was a convenient and reliable model system for the determination of competitive enzyme inhibition.

Under strictly controlled conditions, the K_m value for inosine was determined to be 68.92 ± 4.08 μM, which is consistent to previous report [21]. Product formation by hPNP in the presence of ImmH exhibited a decrease of initial rates with the increase of inhibitor concentration (Fig. 1). The value of K_i was calculated to be $5.21 \pm$ 1.74 nM, which is also in the accordance with previous values [22].

For affinity measurements, hPNP and ImmH were used as a pair of ligand-protein in the absence of inosine (Table 1). After labeling hPNP and scanning fluorescent signal, the affinity values (K_D) by MST were determined to be $336.40 \pm$ 312.90 nM after fitting to MO Affinity Analysis software (Fig. 2). Next, after optimizing concentrations and titration conditions, the K_D values measured by ITC

were 69.10 ± 18.41 nM (Fig. 3). In the case of SPR, the K_D values were $7.07 \pm$

4.23 nM from the analysis by Biacore evaluation software. (Fig. 4)

4. Discussion

According to previous studies, the determination of binding affinity between macromolecules by currently available techniques displays reasonable accuracy and reproducibility. However, in the area of small molecule inhibitor discovery and target identification, the measurements of binding affinity between a small molecule and a protein have been unreliable to result in dramatic differences by different techniques, and even a technique from different determinations. Such ambitious K_D data could be easily misinterpreted or misled to a wasteful effort on target identification and early drug discovery. Currently, there are three major techniques for the determination of binding affinity, namely SPR, ITC and MST, and typically only one technique would be utilized to assess K_D values. Therefore, there is a natural question on which technique can produce the most reliable data on biding affinity. To address this issue, a direct comparison of these techniques with a valid reference is certainly necessary. In order to have a valid reference, we speculated that competitive enzyme inhibition might be able to resemble binding affinity, and subsequently competitive K_i value could be used to objectively compare the K_D values generated by these techniques.

Although enzyme kinetics represent kinetic behaviors during the catalysis, the binding between enzyme and substrate is a prerequisite. Typically, the structure of a competitive inhibitor is very similar to that of a substrate for competing to bind the same binding site. Thus, it is highly possible to use the model of competitive enzyme inhibition to resemble the binding between a small molecule inhibitor and a protein target. Indeed, after theocratical conversion, we demonstrated the correlation between

competitive K_i and binding affinity constant K_D, which principally are convertible based on thermodynamic characteristics. This provided a unique opportunity for us to make a direct comparison on the accuracy and reliability of different techniques used to determine binding affinity constants. Accordingly, when the measured K_D value is equivalent or lower than its K_i value, this biding affinity constant is likely more reliable, even closer to its true affinity value. In the present study, when hPNP, inosine and ImmH were used as a model system [20-22], the K_D value between hPNP and ImmH was closer to the K_D value determined by SPR compared to MST and ITC, suggesting that real-time monitoring through resonance signal might be more suitable for detecting slight changes from the interaction of a small molecule inhibitor and a protein. The K_D values obtained from ITC were slightly larger than K_i values, which could be resulted from the heat transfer effects. However, MST showed poor reproducibility on K_D values (approximately 64 times larger than K_i), which could be caused by various factors [23]. SPR monitors the entire interaction process, whereas MST just compares the variation after binding. In addition, the utilization of SPR in equilibrium analysis is highly compatible with the rapid reversible binding process. Therefore, based on the complexity of the assay mixtures and processes and the differences on individual principle, the present study only provided an estimation on the accuracy of these techniques by comparative analysis regardless of the significant differences.

Given the fact that most proteins for target identification and early drug discovery are not enzymes, it would be impossible to utilize enzyme kinetics to assess

binding affinity constants. Additionally, our present results indicated that currently available techniques are not able to accurately reflect the real affinity of small molecule inhibitors and proteins. Moreover, K_D values will continue to be an important parameter for the prediction of the draggability of small molecule inhibitors. Collectively, the present study alarms the need of developing new technique(s) for accurately, reliably and reproducibly determine the binding affinity constants for small molecule inhibitors.

Conflicts of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this communication.

Acknowledgements

This work was supported by grants from the National Key R&D Program of China (2018YFA0902000), Key Research and Development Project of Guangdong Province (2022B1111070004), the Double "First-Class" University Project (CPU2022QZ08), and the Project Program of the State Key Laboratory of Natural Medicines, China Pharmaceutical University (SKLNMZZ202201).

References

[1] S. Geschwindner, J. Ulander, P. Johansson, Ligand binding thermodynamics in drug discovery: still a hot tip? J. Med. Chem. 58 (2015) 6321-6335. https://doi.org/10.1021/jm501511f

[2] A.P. Ijzerman, D. Guo, Drug–target association kinetics in drug discovery, Trends Biochem. Sci. 44 (2019) 861-871. https://doi.org/10.1016/j.tibs.2019.04.004

[3] T.Y. Wang, H. Ji, D. Everton, A.T.H. Le, S.M. Krylova, R. Fournier, S.N. Krylov, Fundamental determinants of the accuracy of equilibrium constants for affinity complexes, Anal. Chem. 95 (2023) 15826-15832. https://doi.org/10.1021/acs.analchem.3c03557

[4] S.R.J. Hoare, B.A. Fleck, J.P. Williams, D.E. Grigoriadis, The importance of target binding kinetics for measuring target binding affinity in drug discovery: a case study from a CRF(1) receptor antagonist program, Drug Discov. Today 25 (2020) 7-14. https://doi.org/10.1016/j.drudis.2019.09.011

[5] Y. Yin, S.L. Zhao, D. Rane, Z. Lin, M. Wu, B.R. Peterson, Quantification of binding of small molecules to native proteins overexpressed in living cells, J. Am. Chem. Soc. (2023). https://doi.org/10.1021/jacs.3c07488

[6] W. Ma, L. Yang, L. He, Overview of the detection methods for equilibrium dissociation constant K(D) of drug-receptor interaction, J. Pharm. Anal. 8 (2018) 147-152. https://doi.org/10.1016/j.jpha.2018.05.001

[7] M. Jerabek-Willemsen, T. André, R. Wanner, H.M. Roth, S. Duhr, P. Baaske, D. Breitsprecher, MicroScale Thermophoresis: Interaction analysis and beyond, J. Mol. Struct. 1077 (2014) 101-113. https://doi.org/10.1016/j.molstruc.2014.03.009

[8] M. Bastos, O. Abian, C.M. Johnson, F. Ferreira-da-Silva, S. Vega, A. Jimenez-Alesanco, D. Ortega-Alarcon, A. Velazquez-Campoy, Isothermal titration calorimetry, Nat. Rev. Methods Primers 3 (2023). https://doi.org/10.1038/s43586-023-00199-x

[9] J. Homola, Present and future of surface plasmon resonance biosensors, Anal. Bioanal. Chem. 377 (2003) 528-539. https://doi.org/10.1007/s00216-003-2101-0

[10] V. Kairys, L. Baranauskiene, M. Kazlauskiene, D. Matulis, E. Kazlauskas, Binding affinity in drug design: experimental and computational techniques, Expert Opin. Drug Discov. 14 (2019) 755-768. https://doi.org/10.1080/17460441.2019.1623202

[11] B.B. Toure, K. Miller-Moslin, N. Yusuff, L. Perez, M. Dore, C. Joud, et al., The role of the acidity of N-heteroaryl sulfonamides as inhibitors of bcl-2 family protein-protein interactions, ACS Med. Chem. Lett. 4 (2013) 186-190. https://doi.org/10.1021/ml300321d

[12] T. Song, G. Chai, Y. Liu, X. Yu, Z. Wang, Z. Zhang, Bcl-2 phosphorylation confers resistance on chronic lymphocytic leukaemia cells to the BH3 mimetics ABT-737, ABT-263 and ABT-199 by impeding direct binding, Br. J. Pharmacol. 173 (2016) 471-483. https://doi.org/10.1111/bph.13370

[13] M. Milani, D.P. Byrne, G. Greaves, M. Butterworth, G.M. Cohen, P.A. Eyers, S. Varadarajan, DRP-1 is required for BH3 mimetic-mediated mitochondrial

fragmentation and apoptosis, Cell Death Dis. 8 (2017) e2552. https://doi.org/10.1038/cddis.2016.485

[14] A. Kotschy, Z. Szlavik, J. Murray, J. Davidson, A.L. Maragno, et al., The MCL1 inhibitor S63845 is tolerable and effective in diverse cancer models, Nature 538 (2016) 477-482. https://doi.org/10.1038/nature19830

[15] G. Donati, V.M. D'Amore, P. Russomanno, L. Cerofolini, J. Amato, S. Marzano, et al., Theoretical and experimental studies on the interaction of biphenyl ligands with human and murine PD-L1: Up-to-date clues for drug design, Comput.Struct. Biotechnol. J. 21 (2023) 3355-3368. https://doi.org/10.1016/j.csbj.2023.06.006

[16] R. Le Biannic, R. Magnez, F. Klupsch, N. Leleu-Chavain, B. Thiroux, et al., Pyrazolones as inhibitors of immune checkpoint blocking the PD-1/PD-L1 interaction, Eur. J. Med. Chem. 236 (2022) 114343. https://doi.org/10.1016/j.ejmech.2022.114343

[17] A. Meister, Advances in enzymology and related areas of molecular biology, John Wiley & Sons 2009.

[18] P.Y. Tong, S. Kornfeld, Ligand interactions of the cation-dependent mannose 6-phosphate receptor, J. Biol. Chem. 264 (1989) 7970-7975. https://doi.org/10.1016/S0021-9258(18)83137-4

[19] A.G. Marangoni, Enzyme kinetics: a modern approach, John Wiley & Sons 2003

[20] A. Bzowska, E. Kulikowska, D.J.P. Shugar, Purine nucleoside phosphorylases: properties, functions, and clinical aspects, Pharmacol. ther. 88 (2000) 349-425. https://doi.org/10.1016/S0163-7258(00)00097-8

[21] M. Ghanem, N. Zhadin, R. Callender, et al., Loop-tryptophan human purine nucleoside phosphorylase reveals submillisecond protein dynamics, Biochenmistry 48 (2009) 3658-3668. https://doi.org/10.1021/bi802339c

[22] J. Suarez, A.M. Haapalainen, S.M. Cahill, M.C. Ho, F. Yan, et al., Catalytic site conformations in human PNP by 19F-NMR and crystallography, Chem. Biol. 20 (2013) 212-222. https://doi.org/10.1016/j.chembiol.2013.01.009

[23] B. Lopez-Mendez, S. Uebel, L.P. Lundgren, A. Sedivy, Microscale Thermophoresis and additional effects measured in NanoTemper Monolith instruments, Eur. Biophys. J. 50 (2021) 653-660. https://doi.org/10.1007/s00249-021-01529-1

Scheme 1. Related equilibrium equations and formulas in enzyme kinetics

(a)

(b)

$$E + S \underset{k_2}{\overset{k_1}{\rightleftharpoons}} ES \xrightarrow{k_{cat}} E + P \tag{1}$$

$$E + I \underset{k_4}{\overset{k_3}{\rightleftharpoons}} EI \tag{2}$$

$$E + S \underset{k_1}{\overset{k_2}{\rightleftharpoons}} E \underset{k_4}{\overset{k_3}{\rightleftharpoons}} E + I \tag{3}$$

(a) linear relationship between of   and   in different types of inhibition; (b) three equilibriums in competitive inhibition.

Table 1. The binding affinity constants of ImmH with hPNP by different assays and comparison with competitive inhibition constant[a]

Method	K_d (nM)	K_i (nM)
		5.21 ± 1.74
MST	336.40 ± 312.90	
ITC	69.10 ± 18.41	
SPR	7.07 ± 4.23	

[a] Each value was determined by three independent experiments and expressed as average value with standard deviation.

Figure 1. Determination of the dissociation constants for competitive inhibition of hNPN by ImmH.   (5.21 ± 1.74 nM) was obtained from a replot of initial inhibition rate (   ) as a function of inhibitor concentration.

Figure 2. The binding affinity between hPNP and ImmH determined by MST. (A) The relative fluorescence time trace recorded by the MST instrument. (B) The binding curve of ImmH to hPNP was fitted by a standard *Kd* model.

(A)

(B)

Figure 3. Results of the ITC measurements with ImmH and hPNP. (A) The titration of raw data after sequential injection of ImmH. (B) The integrated heat after the correlation between enthalpy changes versus the molar ratios of ImmH (150 μM) to hPNP (10 μM)) was plotted.

Figuree 4. Binding affinity of ImmH to hPNP measured by SPR. (A) Plots of equilibrium response versus concentration of ImmH. (B) SPR sensorgrams by affinity analysis of the binding of ImmH to hPNP on sensor chip surface.