High-throughput screening of more than 30,000 compounds for anthelmintics against gastrointestinal nematode parasites

Mostafa A. Elfawal[1*], Emily Goetz[1#], You-Mie Kim[1#], Paulina Chen[1], Sergey N. Savinov[2], Leonard Barasa[3], Paul R. Thompson[3], Raffi V. Aroian[1*]

[1]Program in Molecular Medicine, University of Massachusetts Chan Medical School, Worcester, MA, 01605, USA

[2]Department of Science, Rivier University, Nashua, NH 03060, USA

[3]Department of Chemical Biology, University of Massachusetts Chan Medical School Worcester, MA, 01605, USA

* For Correspondence: raffi.aroian@umassmed.edu , mostafa.elfawal@umassmed.edu

# Equal contributing second authors

Abstract:

Gastrointestinal nematodes (GINs) are amongst the most common parasites of humans, livestock, and companion animals. GIN parasites infect 1-2 billion people worldwide, significantly impacting hundreds of millions of children, pregnant women, and adult workers, thereby perpetuating poverty. Two benzimidazoles with suboptimal efficacy are currently used to treat GINs in humans as part of mass drug administrations, with many instances of lower-than-expected or poor efficacy and possible resistance. Thus, new anthelmintics are urgently needed. However, screening methods for new anthelmintics using human GINs typically have low throughput. Here, using our novel screening pipeline that starts with human hookworms, we screened 30,238 unique small molecules from a wide range of compound libraries, including ones with generic diversity, repurposed drugs, natural derivatives, known mechanisms of action, as well as multiple target-focused libraries (e.g., targeting kinases, GPCRs, and neuronal proteins). We identified 55 compounds with broad-spectrum activity against adult stages of two evolutionary divergent GINs, hookworms (*Ancylostoma ceylanicum*) and whipworms (*Trichuris muris*). Based on known databases, the targets of these 55 compounds were predicted in nematode parasites. One novel scaffold from the diversity set library, F0317-0202, showed good activity (high motility inhibition) against both GINs. To better understand this novel scaffold's structure-activity relationships (SAR), we screened 28 analogs and created SAR models highlighting chemical and functional groups required for broad-spectrum activity. These studies validate our new and efficient screening pipeline at the level of tens of thousands of compounds and provide an important set of new GIN-active compounds for developing novel and broadly-active anthelmintics.

Keywords:

Gastrointestinal nematode parasites, soil-transmitted helminths, anthelmintic discovery, drug screening, hookworms, whipworms

Human gastrointestinal nematodes (GINs, also known as soil-transmitted helminths) include the roundworm *Ascaris lumbricoides*, the whipworm *Trichuris trichiura*, and the hookworms *Necator americanus*, *Ancylostoma duodenale*, and *Ancylostoma ceylanicum*; they are amongst the most prevalent infections of the so-called Neglected Tropical Diseases (NTDs)[1], affecting people in the world's poorest communities[2-4]. GINs are successful parasites of vertebrates, adapted to parasitize humans, livestock, and companion animals[5]. Morbidities associated with GIN infections in humans include anemia, malnutrition, growth stunting, cognitive impairment, pregnancy complications, and lowered immunity against other infectious diseases[2, 4, 6]. The negative impacts of GINs are hard to measure, but they are minimally responsible for > 2-3 million disability-adjusted life years (DALY)[3, 6, 7], with hookworms accounting for almost one-half of those DALYs[3, 6]. Children who received hookworm treatment had increased school attendance and adult income[4, 8]. GINs also have enormous impacts on the health and productivity of farm animals[9].

The World Health Organization (WHO) has recommended annual or semi-annual Mass Drug Administration (MDA) targeting preschool and school-age children to reduce worm burden and associated morbidity[10]. Four anthelmintics (levamisole, pyrantel pamoate, albendazole, and mebendazole), initially developed for livestock, are currently on the WHO list to control human GINs, with the two benzimidazoles, albendazole and mebendazole, most widely used because of their better efficacies and ease of dosing[11, 12]. In companion and livestock animals, GINs are controlled via frequent and routine treatment using a repertoire of anthelmintics, which has led to widespread and sometimes multi-drug anthelmintic resistance[13-16]. Benzimidazole resistance is well-established in veterinary parasites[17-20]. Single nucleotide polymorphisms (SNPs) in the beta-

tubulin gene linked to the benzimidazole resistance in veterinary parasites have been found in human GINs[21-24], indicating that with high drug pressure, there is likely to be an emergence for benzimidazole resistance in human GINs. In Haiti and Kenya, the frequency of benzimidazole-resistance codon 200 in *T. trichiura* was significantly increased following local MDA programs[22]. There are also multiple reports of lowered efficacy of benzimidazoles against human GINs and warnings against drug resistance[25-27]. Notwithstanding these challenges, the WHO has set an ambitious goal to eliminate human GINs by 2030[10]. Therefore, potent and broadly active anthelmintics with new modes of action are urgently needed for long-term control and elimination. One promising new drug is emodepside, which was discovered >30 years ago [28]. However, it has not yet been approved for human use, and, as with all single anthelmintics, resistance is a concern[29, 30]. Given the lead time for new drugs to enter the clinic, it is urgent that new lead anthelmintics be put into the pipeline.

Because understanding of these parasites' biology is still limited, whole organism phenotypic screening remains a preferred approach for anthelmintic drug discovery[31, 32], although more recently some more directed screens, *e.g.*, based on detailed knowledge of metabolic pathways, have shown promise[33-35]. Phenotypic screening using GIN parasites for anthelmintic has included screening against *ex vivo* adult parasites, egg hatch assay, larval stages, and infective third-stage larvae[32]. However, regarding GINs that infect humans, the throughput of these screens has been limited[32, 36-38]. The free-living nematode *Caenorhabditis elegans* has been used as a surrogate model for GINs that infect humans[32, 39-42]. However, we found limitations to using *C. elegans* as a model for anthelmintic screening targeting human GINs, most notably a significant false negative hit rate[31]. Thus, *C. elegans* is likely to miss some of the most promising

compounds. Using a pilot study with 1,280 compounds, we devised an alternative screening

model using egg-to-larva development of the free-living stages of *A. ceylanicum* human

hookworm parasites, which we found predicted better anthelmintic activity against GIN adult

parasites than *C. elegans* [31]. In this work we further extended this primary screen with secondary

screens using the adult stages of evolutionarily divergent hookworm and whipworm parasites

and proposed an entire screening pipeline based on these results (Figure 1)[31].

Here, we scale up this anthelmintic discovery pipeline and screen 38,293 conditions (76,586 in

duplicate) representing 30,238 unique compounds. We identified 55 small molecules with *in

vitro* efficacy against adult hookworms and whipworms. We focused on one novel scaffold and

performed Structure-Activity Relationship (SAR) studies. Based on analogies from other

systems, we propose essential GIN targets for further studies.

Figure 1: Novel anthelmintic discovery pipeline for human GINs. Our pipeline is centered around the zoonotic hookworm, *A. ceylanicum,* as a primary (larval) and secondary (adult) screening model, followed by tertiary screening against the murine whipworm *Trichuris muris* (adult) to identify broadly active compounds. Hits are subjected to deep data mining, and analogs of potential candidates are tested to generate the Structure-Activity Relationships. This report covers the results from the first five stages of the screening pipeline (in blue).

Results

In our pilot screen[31], we found that only ~10% (4/39) of the compounds that targeted adult

parasites were found by inhibiting *A. ceylanicum* egg hatch (the other 90% were found as *A.

ceylanicum* larval hits). Furthermore, of those four compounds, three were retested against larval

stages and were found to effectively target that stage (the fourth was not tested). On the other

hand, we found that the consistency of the screen improved starting with hatched and

synchronized *A. ceylanicum* first larval (L1) stages versus starting with eggs. We therefore modified the screen to start with *A. ceylanicum* L1s. We scaled up the same primary screen to include 30,000+ compounds in 13 chemical libraries (Figure 2). These libraries encompass many structural features, including diversity scaffolds, natural product derivatives, repurposing drugs, known modes of action, and target-based libraries (Table 1). In total, we screened 30,238 unique compounds in duplicate. Some compounds were screened at multiple doses due to plate format availability, making the total number of individual conditions screened 38,293 (76,586 when considered in duplicate). All libraries were initially screened in duplicate at 10 µM. As per our pipeline, primary screen hits (in both duplicate wells) were screened at 30 µM against adult *A. ceylanicum* hookworms. Adult hookworm hits were then screened at 30 µM against adult *T. muris* whipworms. A breakdown of hits "library by library" is given below and summarized in Table 1.

Figure 2: Primary screening using the free-living larval stages of *A. ceylanicum* hookworms. To work around variability in egg hatch rates, we found that starting with the first larval stage (L1) led to more consistent and robust results (see Results section). Isolated and purified eggs were allowed to hatch in S-media into synchronized L1. Each 96-well assay plate includes four wells of ivermectin (10µM) as positive control and four wells of 0.25% DMSO as negative controls. After seven days of incubation at 28°C, the plates were scored under a light microscope. A test compounds was considered active (hit) if at least 90% of all larvae in the well were killed or

failed to develop. Scale bar 1 mm.

Table 1: Screening summary of 30,238 compounds from 13 different libraries with information on screening doses, the number of hits per screening model, and the hit rates.

Library	Unique Compounds	Screened doses µM	Test wells screened in duplicate	*A. ceylanicum* L1 at 10µM (% hits)	*A. ceylanicum* adults 30µM (% hits)	*T. muris* Adults 30µM (% hits)
SMSF Life Chemicals Diversity Set	15360	10	15360	491 (3.2)	33 (0.21)	7 (0.05)
Broad Institute REPO 1, 2 and 3	6743	10	6743	230 (3.4)	96 (1.42)	36 (0.53)
ICCB Longwood MOA	1245	10, 2, 0.4, 0.08	4980	65 (5.3)	17 (1.36)	9 (0.72)
ICCB Selleck Neuronal signaling	1031	10	1031	29 (2.8)	12 (1.16)	2 (0.19)
ICCB SYNthesis Kinase Inhibitor 2	96	10, 2, 0.4	288	8 (8.3)	1 (1.04)	1 (1.04)
ICCB EMD Kinase Inhibitor 1	244	10	244	10 (4.1)	2 (0.81)	2 (0.81)
ICCB ChemBridge Focused Kinase	250	10	250	2 (0.8)	0	0
ICCB ChemBridge Focused GPCR	250	10	250	2 (0.8)	0	0
ICCB LINCS 1 Kinase inhibitors	88	10, 2, 0.4, 0.08	352	4 (4.5)	2 (2.27)	1 (1.13)
ICCB LINCS 2 Kinase inhibitors	88	10, 2, 0.4, 0.08	352	2 (2.25)	0	0
ICCB LINCS 4 Kinase inhibitors	169	10, 2, 0.4, 0.08	676	8 (4.7)	3 (1.77)	3 (1.7)
SMSF Kinase Inhibitors	492	10	492	13 (3)	5 (1.01)	3 (0.69)
SMSF TIMTEC Natural Compounds	4240	10	4240	14 (0.33)	0	0
Total	**30238**		**38293**	**878 (2.9)**	**171 (0.56)**	**64 (0.21)**

Hits from the Life Chemicals Diversity Set:

Of the 15,360 compounds in the Diversity Set tested against *A. ceylanicum* hookworm larvae, 491 were active, with a hit rate of 3.2%. These 491 larval hits were tested against the early adult stages of *A. ceylanicum* hookworm, yielding 33 hookworm adult-active compounds. An example of screening against the early adult stages of *A. ceylanicum* hookworms is shown in Figure 3. These 33 *A. ceylanicum* hookworm adult-active compounds were screened against *T. muris* adult whipworms, yielding seven hits active against the two evolutionarily divergent parasites. An example of screening against adult *T. muris* whipworms is shown in Figure 4.

Figure 3. Young adult *A. ceylanicum* hookworm screens. The top panel shows healthy and motile worms from the DMSO-treated or nonactive test compounds. The lower panel shows dead or intoxicated worms from ivermectin or active compounds. Healthy worms with smooth and fully expanded cuticles were capable of controlling their body muscles, were motile, were elongated, and formed more bends. Dead or intoxicated worms were shortened, had wrinkled and shrunken cuticles, were immobile or barely mobile, and appeared more internally darkened due to damaged intestines. Because young adult hookworms are much smaller than whipworms, we manually scored these versus using the Worminator. Scale bar 1 mm.

Figure 4: Adult parasite screening of *T. muris* whipworm**. (A), Adult whipworms, male and female treated with 30µM Levamisole or 1% DMSO. Scale bar 1 mm. (B): Motility readout screenshot from 48-well plate with single adult whipworm using Wormassay-6 software and the Worminator imaging setup. Red depicts pixel changes from frame to frame (movement) and blue depicts no change in pixel (dead or very low movement). (C). Example of calculated motility readout, showing some with adult-parasitic activity with normalized motility ~ 50% compared to the DMSO control.

Hits from the Repurposing and Mechanism of Action (MoA) Libraries

Of the 6,743 compounds tested from the repurposing library, we identified 230 *A. ceylanicum* hookworm larval-active compounds (hit rate of 3.4%), from which 96 showed activity against adult *A. ceylanicum* hookworms (hit rate 1.42%). Of these 96 compounds, 36 were active against adult *T. muris* whipworms with a final hit rate of 0.53%. Similarly, The ICCB-Longwood MoA library contained 1,245 high-quality compounds with known mechanisms of action. We identified 65 *A. ceylanicum* hookworm larval-active compounds with a 5.2% hit rate, the highest among all tested libraries. Seventeen of these compounds carried adulticidal activity against *A.*

ceylanicum hookworms, from which nine were active against *T. muris* whipworm adults, with a final hit rate of 0.72%.

Hits from the Neuronal and the GPCR libraries:

Of the 1031 compounds with known modulation of neuronal signaling, we identified 29 *A. ceylanicum* hookworm larval-active compounds (hit rate of 2.8%). Of these 29 compounds, 12 showed activity against adult *A. ceylanicum* hookworms (1.1% of the total), from which two were active against *T. muris* whipworm adults with a final hit rate of 0.19%. We also screened 250 known GPCR inhibitors and identified two compounds with larval activity against *A. ceylanicum;* however, these compounds did not show activity against the adult stages of parasites (suggesting these two compounds only target proteins essential in larval development). The general lack of success with this library could be explained by the selectivity of these GPCR inhibitors, which were tailored initially for binding to human GPCRs.

Hits from the Kinase Inhibitor Libraries:

Of seven kinase inhibitor libraries with 1427 compounds (Table 1), we identified 47 with *A. ceylanicum* larval activity, giving a hit rate of 3.2%. Of those 47 compounds, 13 showed activities against the adult stages of *A. ceylanicum* hookworms (0.9%), of which 10 compounds were also active against *T. muris* whipworm adults with a hit rate of 0.7%.

Hits from the Natural Products Library:

Of 4240 compounds from TimTec natural products libraries, we identified 14 compounds with *A. ceylanicum* larval activity, giving the lowest primary screen hit rate of 0.33%. None of these

14 larvicidal compounds showed activity against the adult stages of *A. ceylanicum* hookworms. We hypothesize that the low hit rate and lack of adult activity are due to the nature of these libraries (see Materials and Methods).

Summary of 30,238 compound screen

Collectively, we identified 878 compounds with *A. ceylanicum* larval activity, giving an overall primary screen hit rate of 2.9% (Table 1). In our secondary screen, we tested all larval-active compounds at 30 µM against the adult stages of *A. ceylanicum* hookworm parasites. Of those, 171 (19.5% of the larval hits) had anthelmintic activity toward adult hookworms, representing a hit rate of 0.5% based on the total number of compounds in the initial libraries. These 171 hookworm-adult actives were tested against adult *T. muris* whipworms at 30 µM. Sixty-four compounds (37% of the adult hookworm hits and 0.21% based on the total number of compounds in all the libraries) showed activity against whipworm adults (Table 1). Of the 64 compounds identified from all libraries, nine were redundant (included independently in multiple libraries), yielding 55 unique active compounds showing a motility index of ≤1 or intoxicated phenotype in hookworms and relative motility of ~ 50% in whipworms compared to the DMSO control, listed in Table 2.

Screened libraries contained known anthelmintic compounds, and we successfully identified albendazole, oxibendazole, flubendazole, methiazole, oxfendazole, thiabendazole, ivermectin, abamectin, dormamectin, emamectine, eprinomectin, selamectin, emodepside, closantel, monepantel, and moxidectin as part of the primary and secondary *A. ceylanicum* hookworm screens. However, many of these anthelmintics didn't show activity in our tertiary screening

against the *T. muris* whipworms at 30µM, highlighting the whipworms' resilience to anthelmintics compared to hookworms. For example, the published 50% inhibitory concentrations (IC$_{50}$) of albendazole and ivermectin against *T. muris* whipworms are both > 200µg/ml[43]. Only anthelmintic compounds that showed activity at 30µM against *A. ceylanicum* and *T. muris* are shown in Table 2.

Table 2: List of 55 potent and broadly active compounds including chemical structure, source library, *in vitro* activity against *A. ceylanicum* hookworms (HW) with motility index (M.I.) of ≤1 or intoxicated phenotype, where 1 is intoxicated worms moving only aftet being touched and 0 is completely immotile), *in vitro* activity against *T. muris* whipworms (WW) with percent Relative Motility (% R.M.) of ~ 50%, where100% is healthy; 0% is non-motile) to the DMSO control, known mode of action in eukaryotic organisms, and the orthologous targets in nematodes if any. (--) no known target.

Drug Names	Structure	Source Library	HW M.I. 30μM	WW % R.M. 30μM	Known MoA /Target	Target orthologue in Nematodes
Emodepside		REPO	0	8.6	Anthelminthic, bind to nematode Latrophilin-like receptor (LAT-1), Voltage-gated activated potassium channels (SLO-1) [44, 45]	Lat-1 and slo-1in *C. elegans*, TMP05851 and TMP00134 in *T. muris*
Metrifonate; Trichlorfon		REPO	0.75	1.3	Anthelminthic[46], Acetylcholinesterase (ACHE) inhibitor [47]	ace-1 in *C. elegans*, TMP09383 in *T. muris*
Tyrphostin-A9; Malonoben		REPO, SMSF kinase	0	9.1	Nematicidal [48], Protein tyrosine kinase receptor inhibitor/ Epithelial Growth Factor Receptor (EGFR), Platelet-Derived Growth Factor Receptor (PDGFR) [49]	Let-23, kin-30, in *C. elegans*, TMP05029 and TMP12675 in *T. muris*
Brilliant-green		REPO	0	5.1	--	--
Alexidine		REPO	1.75	1.6	Protein Tyrosine Phosphatase localized to the Mitochondrion 1(PTPMT1) [50]	F28C6.8 in *C. elegans*, TM12441 in *T. muris*
Octenidine		REPO	0.75	7.3	--	--
Walrycin-B		REPO	0	8.6	--	--

Drug Names	Structure	Source Library	HW M.I. 30μM	WW % R.M. 30μM	Known MoA /Target	Target orthologue in Nematodes
EVP4593; QNZ		REPO, ICCB MOA, ICCB EMD kinase, ICCB LINCS Kinase	0	35.8	Nematicidal[51], TNF-α and NF$_k$β pathway inhibitor [52]	--
AG-1024		REPO	2	7.5	Protein tyrosine kinase receptor inhibitor/ Insulin growth factor receptor inhibitor/ (IGF1R) [53]	Daf-2 in *C. elegans*, TMP10632 in *T. muris*
Tyrphostin AG879		REPO, SMSF kinase	0	20	Nematicidal [48], Protein tyrosine kinase receptor inhibitor/Nerve Growth Factor Receptor (NGFR) [49]	--
Diphenyleneiodonium		REPO	0	15.2	Nitric oxide synthase inhibitor/ NOS2, NOS3 [54]	Lacked in *C. elegans* NOS-Like in *A. suum*
ESI-09		REPO	0.5	16.2	Exchange protein activated by cAMP isoform 2 (EPAC2) inhibitor [55, 56]	epac-1 in *C. elegans*, TMP12712 in *T. muris*
Obatoclax		REPO, ICCB MOA	0.5	10.7	Induce apoptosis via the inhibition of the pro-survival protein/ BCL2[57]	ced-9 in *C. elegans*, TMP02363 in *T. muris*
Oligomycin-A		REPO	1.25	15.9	Mitochondrial ATP synthesis inhibitor/ ATP5A[58]	atp-1 in *C. elegans*, TMP09383 in *T. muris*
Pyrithione-zinc		REPO	0	3.6	KCNQ Kv7 potassium channels agonist [59]	kqt-1 in *C. elegans*, TMP05091 in *T. muris*

Drug Names	Structure	Source Library	HW M.I. 30μM	WW % R.M. 30μM	Known MoA /Target	Target orthologue in Nematodes
NH125		REPO, ICCB LINCS 1 kinase	1.25	8	Nematicidal[51], Eukaryotic translation elongation factor 2 kinase (EEF2K) inhibitor [60]	efk-1 in *C. elegans* and TMP13626 in *T. muris*
Tetramisole		REPO	2.5	9.4	Anthelminthic, acetylcholine receptor agonist [61]	acetylcholine receptor agonist
Levamisole		REPO	1	10.1	Anthelminthic, acetylcholine receptor agonist [61]	acetylcholine receptor agonist
Triclocarban		REPO	0	12.7	--	--
Semapimod		REPO	1.75	39.7	Inhibit Toll Like Receptors (TLR) signaling by targeting TLRs chaperone gp96 [62]	*enpl-1* in *C. elegans* and TMP11927 in *T. muris*
Amiselimod		REPO	1	3.3	Sphingosine 1-phosphate receptor (S1PR) modulator [63]	Spin-2 in *C. elegans* and TMP13635 in *T. muris*
Cetylpyridinium		REPO	0.25	15.1	Nematicidal [31], HSP-90 inhibitor [64]	Hsp-90 in *C. elegans*, TMP11927 in *T. muris*
Domiphen		REPO	0	11.5	Blockade of human ether-a-go-go related gene (HERG) potassium channels [65]	unc-103 in *C. elegans*, TMP12566 in *T. muris*
Sulconazole		REPO	1	3.1	Nematicidal [31], Sterol 14 alpha Demethylase cytochrome (CYP-51) inhibitor [66]	--
MLN8054		REPO	1.5	18	Aurora Kinase inhibitor/ AURKA [67]	Air-2 in *C. elegans*, TMP01390 in *T. muris*
Cetrimonium		REPO	0.25	2.7	Nematicidal [31] --	--

Drug Names	Structure	Source Library	HW M.I. 30μM	WW % R.M. 30μM	Known MoA /Target	Target orthologue in Nematodes
Econazole		REPO	1.2	7.6	Nematicidal [31], Sterol 14 alpha Demethylase cytochrome (CYP-51) inhibitor [66]	--
Crystal-violet		REPO	0.25	5.3	Nematicidal [31], HSP-90 inhibitor [64]	Hsp-90 in C. elegans TMP11927 in *T. muris*
Chlorpromazine		REPO	0.2	2.7	Nematicidal[68], Dopamine D2 receptor antagonist [69]	dop-2 and dop-3 in C. elegans and TMP13188 in *T. muris*
Dioscin		REPO	1.5	15.3	Antagonize chemokine receptor CXCR3 block interaction to IP-10 cytokine [70]	Npr-20 in *C. elegans*, TMP12387 in *T. muris*
Siramesine; Lu28-179		REPO	2	2.1	σ_2R receptor agonist [71]	--
Selamectin		REPO	1	17.1	Anthelminthic, Neuromuscular paralysis via activation of Glutamate-gated chloride channels [72-74]	avr-14 in *C. elegans*, TMP04454 in *T. muris*
Isoconazole		REPO	0.4	3.1	Sterol 14 alpha Demethylase cytochrome (CYP-51) inhibitor [66]	--

Drug Names	Structure	Source Library	HW M.I. 30µM	WW % R.M. 30µM	Known MoA /Target	Target orthologue in Nematodes
ML281		REPO	1.5	16.7	Serine/threonine kinase inhibitor/ STK33[75]	--
Abamectin		REPO	0	13.1	Anthelminthic, Neuromuscular paralysis via activation of Glutamate-gated chloride channels [72-74]	avr-14 in *C. elegans*, TMP04454 in *T. muris*
Squalamine		REPO	1	3.1	protein tyrosine phosphatase (ptpn1b) inhibitor [76]	Sdf-1 in *C. elegans*, TMP05082 in *T. muris*
Trodusquemine		REPO	0.25	2.1	protein tyrosine phosphatase (ptpn1b) inhibitor[76]	Sdf-1 in C. elegans, TMP05082 in *T. muris*
Bafilomycin A1		ICCB MOA	0.5	52.7	Disrupt Autophagic flux by inhibiting V-ATPase and Ca-P60A/SERCA[77]	Vha-19 and sca-1 in *C. elegans*, TMP10618 and TMP09741 in *T. muris*
NMS-873		ICCB MOA	0	34.5	Allosteric inhibitor of the ATPase valosin-containing protein (VCP/p97) [78]	cdc-48 in *C. elegans*, TMP02658 in *T. muris*
1-Phenyl-3-[4-(trifluoromethox) phenyl]urea		ICCB MOA	0	50.2	--	--
2,3-Diphenyl-5-[(2-methoxyphenyl)amino]-1,2,4-thiadiazole-2-ium		ICCB MOA	0	41.7	--	--

Drug Names	Structure	Source Library	HW M.I. 30μM	WW % R.M. 30μM	Known MoA /Target	Target orthologue in Nematodes
Furan-2-ylmethyl-(5-phenyl-thieno[2,3-d]pyrimidin-4-yl)-amine		ICCB MOA	0	41.9	--	--
4-Bromo-N-[2,2,2-trichloro-1-(4-chloroanilino)ethyl]benzamide; AMG7160		ICCB MOA	2	27.6	Ca^{2+} Cation channel, transient receptor potential ankyrin 1 (TRPA1) antagonist [79]	Trpa-1 in *C. elegans*, TMP11515 in *T. muris*
T0901317		ICCB Selleck Neuronal	2	37.6	Liver X receptor LXR (Nr1H2, Nr1H3) agonist [80-83]	nhr-247 and nhr-8 in *C. elegans*, TMP09010 and TMP06463 in *T. muris*
Vortioxetine; Lu AA21004		ICCB Selleck Neuronal	2	30.9	5-HT3-R antagonist [84, 85] and serotonin transporter (SERT) inhibitor [86]	Lgc-7, lgc-8 *and mod-5* in *C. elegans,* TMP06220, TMP00636 and TMP10599 in *T. muris*
Sorafenib		SMSF kinase, ICCB SYNthesis2, ICCB LINCS 1 kinase	0	20.6	Inhibit Raf-1, BRAF (serine/threonine Kinase), VEGFR, PDGFR, KIT, FLT3, FGFR1(tyrosine kinase) [87]	Lin-45, ver-1, ver-4, kin-30, old-2 and igcm-4 in *C. elegans* TMP03531, TMP12675, TMP13825, TMP00128, TMP02580, TMP02073 in *T. muris*

| Staurosporine | | ICCB EMD Kinase | 1.25 | 27.7 | Promiscuous Protein Kinase inhibitor [88] | |
Drug Names	Structure	Source Library	HW M.I. 30µM	WW % R.M. 30µM	Known MoA /Target	Target orthologue in Nematodes
NVP-BGT226; HMSL10349-106		ICCB LINCS 1 Kinase	1.5	0	phosphatidylinositol 3-kinase (PI3K) inhibitor and MTOR [89]	age-1 in *C. elegans*, TMP06498 in *T. muris*
F0317-0202		SMSF Diversity set	0	10.7	--	--
F1059-0041		SMSF Diversity set	1.75	19.5	--	--
F1620-0033		SMSF Diversity set	1	13	--	--
F9995-0076		SMSF Diversity set	0	3.7	--	--
F9995-0233; dithiazanine iodide		SMSF Diversity set	0	1.75	Anthelminthic activity via inhibition of oxygen consumption and interreference with glucose uptake [90-94].	--

F0900-4647		SMSF Diversity set	0	3	--	--
F1345-0623; HBX		SMSF Diversity set	0.5	9	USP7 inhibitor [95]	math-33 in *C. elegans*, TMP04955 in *T. muris*

Data Mining

The 55 broadly active compounds were subjected to a literature search to identify, if known, their possible mode of action and the eukaryotic drug targets in other organisms and the orthologous targets in phylum Nematoda, focusing on *C. elegans* in clade V and *T. muris* in clade I (Table 2). The compounds encompass a wide range of putative target classes including potassium channel, glutamate-gated chloride channel, acetylcholinesterase, acetylcholine receptor, tyrosine kinases, aurora kinase, phosphatidylinositol 3-kinase (PI3K), tyrosine phosphatase, insulin growth factor receptor (IGF1-R), nitric oxide synthase (NOS), Exchange Protein Activated by Cyclic AMP (EPAC), pro-survival protein (BCL2), ATP synthase, translation elongation factor, Enoyl-acyl carrier protein reductase (ENR), p38 mitogen-activated protein Kinase (p38 MAPK), sphingosine 1-phosphate receptor (S1PR), Heat Shock Protein-90 (HSP-90), Dopamine D2 receptor, liver X receptor (LXR), serotonin (5-HT)3 receptor, chemokine CXCR3 receptor, and ubiquitin specific peptidase 7 (USP-7) and Phosphatase localized to Mitochondrion 1(PTPMT1). Compounds identified here with well-known targets and mechanisms of action in other organisms may be used as probes for druggable and essential ortholog targets in nematode parasites (Table 2). For example, alexidine was shown to induce selective responses similar to the knockdown of its target PTPMT1[96]. Similarly, T0901317 is a well-known selective agonist of LXR used in many studies to investigate the physiological

effects of LXR on cholesterol and lipid homeostasis; notably, T0901317 induction of LXR was found to reduce cholesterol biosynthesis and to induce cholesterol breakdown into bile acids [80-82]. Nematodes can't synthesize cholesterol and rely on exogenous sources from the environment or their hosts[97, 98]. Thus, ligand activation of the LXRs ortholog in parasitic nematodes could result in cholesterol depletion and negatively impact parasite lipid homeostasis. As examples, we discuss putative nematode targets of four of these-- BCL2, NOS, PTPMT1, and LXRs, selected because these targets are known to be essential (See Supplementary Discussion).

Novel anthelmintic scaffold

We identified seven compounds with broad anthelmintic activity from the Life Chemicals Diversity Set, which is compiled of *de novo* and newly synthesized molecules developed for initial phenotypic screening. Because of the novelty of this library, we chose to study one of the hits further as an initial foray into follow up studies. Among the seven actives from this library of novel scaffolds, we focused on F0317-0202 as a lead for further studies because given its better drug-like structure over the other six (and see Discussion). To further investigate, 28 commercially available analogs of F0317-0202 were identified using structural similarity searches and then purchased from Life Chemicals. These could serve as training sets for ligand-based optimization and generate the SAR model. We screened this set of analogs against adult *T. muris* whipworms, followed by older and larger *A. ceylanicum* hookworm adults, which could be read more easily in the Worminator.

Of the 28 analogs, we identified 12 compounds (41%) that inhibit *T. muris* whipworms motility by $\geq$ 50% relative to the DMSO control (Figure S1). All 12 were tested and found to be active

against *A. ceylanicum* hookworm adults (Figure S1). To rank these 12 analogs, they were tested *in vitro* in dose-response against hookworms and whipworms. Of these 12 broadly active compounds, six analogs (F0317-0161, F0317-0013, F0328-0210, and F0317-0009, F0317-0019, and F0317-0160), along with the parent, showed IC_{50}s of $\leq 5\mu M$ against adult hookworms and $\leq 25\mu M$ against adult whipworms (Figure 5). Note for comparison, the IC_{50} values of albendazole and ivermectin, the first-line anthelmintics, against *T. muris* whipworms were > 753 and $228\mu M$, respectively [43].

Based on these data, an F0317-0202 initial SAR model for potent and broad anthelmintic activity was built and found to be quite strict— slight modifications could abolish the anthelmintic activity entirely. The functional core identified in this potent scaffold was the (z)-2-(2,3-dihydrobenzo[*d*]thiazol-2-yl)-3-hydroxybut-2-enenitrile (Figure 6A) linked to an aromatic ring, either thiophene or benzene (Figure 6. B). Interestingly, replacing thiophene with benzene resulted in a dramatic reduction in the activity against adult whipworms. Analogs with a benzene ring and one or more of the following groups: nitrile, ethylbenzene, chlorine, bromine, or anisole (Figure 6. C) showed potent *in vitro* activity against both parasites (Figure 5). F-0317-0161, with a benzonitrile, showed the most potent *in vitro* activity against hookworms with an IC_{50} of $0.77\mu M$. These findings suggest that further optimization and finetuning on the variable aromatic ring is feasible to design analogs with broad-spectrum anthelmintic activity.

F0317-0202 lead HW IC$_{50}$: 1.63µM WW IC$_{50}$: 1.46µM											
HW IC$_{50}$	WW IC$_{50}$	HW IC$_{50}$	WW IC$_{50}$	HW IC$_{50}$	WW IC$_{50}$	HW IC$_{50}$	WW IC$_{50}$	HW IC$_{50}$	WW IC$_{50}$	HW IC$_{50}$	WW IC$_{50}$
F0317-0161		F0317-0013		F0328-0210		F0317-0009		F0317-0019		F0317-0160	
0.77 µM	9.11 µM	2.1 µM	10.97 µM	4.0 µM	11.1 µM	4.7 µM	14.5 µM	4.6 µM	24.9 µM	5.0 µM	21.7 µM
F0463-0179		F0317-0175		F0336-0103		F0317-0012		F0328-0550		F0882-0289	
6.9 µM	15.5 µM	11.24 µM	-	12.35 µM	47.81 µM	16.06 µM	15.25 µM	23.55 µM	17.95 µM	54.19 µM	-

Figure 5: IC$_{50}$ values of 12 F0317-0202 broadly active analogs: Chemical structure of the top tested analogs and the motility IC$_{50}$ values, against *A. ceylanicum* hookworms (HW) and *T. muris* whipworms (WW). (-) indicate that we were not able to calculate an IC$_{50}$ values as the highest dose used did not reach 50% inhibition.

Figure 6: F0317-0202 Structure-activity relationship (SAR) model. (A) The core and essential structure (red) and the position of the variable rings of F0317-0202 .

(B) variable rings and the associate nematicidal activity. (C) Analogs with a benzene ring and variable groups ranked based on the IC_{50} against the adult stages of hookworms.

Discussion

Human GINs are among the most prevalent infections of the NTDs. Because understanding of these parasites' basic biology is still limited, whole organism screening (*i.e.*, phenotypic screening vs. target-based screening) remains the most effective approach in finding effective anthelmintics. We previously validated a novel screening pipeline using a set of 1280 FDA-approved compounds from a single library [31]. This screening pipeline utilizes two phylogenetically diverse nematode parasites, *A. ceylanicum* hookworms and *T. muris* whipworms, located at opposite ends of the phylum Nematoda phylogenetic tree and that are either a human parasite (*A. ceylanicum*) or closely related to a human parasite (*T. muris* vs *T. trichiura*). By selecting these diverse nematodes, we aim to identify compounds with broad anthelmintic activity. Here, using much larger and more diverse sets of compound libraries, we used our pipeline to screen more than 30,000 compounds, which is unprecedented for targeting the GINs of humans.

The life chemicals diversity set library comprises 15,360 compounds representing 1,717 chemical scaffolds from novel and newly synthesized small molecules. Using our screening pipeline, we uncovered seven molecules with broad activity against parasitic nematodes, giving a hit rate of 0.045%. Among these seven, four are polymethine cyanine dyes: F1059-0041, F9995-0076, F0900-4647, and dithiazanine iodide (F9995-0233). Dithiazanine iodide is known to be active against hookworms, threadworms, roundworms, pinworms, and whipworms [90-94],

providing additional validation for our pipeline. Of the other three non-polymethine cyanine dyes, F1345-0623 has been identified as a potential inhibitor (IC_{50} in the sub-micromolar range) of the deubiquitinase USP7 [95]; homologous proteins have been found to play essential roles in *C. elegans* (math-33)[99] and *Schistosoma mansoni* (SmUSP-7) [100].

Of the remaining two novel compounds, F0317-0202 showed better IC_{50}s than F01620-0033 against hookworms (1.6 compared to 14µM) and whipworms (1.4 compared to 5.8µM) and better drug-like structure and thus was selected for further studies. F0317-0202 includes two aromatic ring systems (thiophene and benzothiazole), listed among the most frequently used ring systems from small molecules in the FDA Orange Book [101]. F0317-0202 also has two functional groups (OH and CN), listed among the most frequent functional groups occurring in bioactive molecules [102]. F0317-0202's physicochemical property better agrees with Lipinski's rule of five for drug-likeness than F01620-0033[103], with a molecular weight of 318.8 compared to 449g/mol, one hydrogen bond donor compared to two, five hydrogen bond acceptor compared to seven. Finally, F0317-0202 was predicted to have better absorption with 113 □² of Topological Polar Surface Area (TPSA) than F01620-0033 (142 □²), as compounds with smaller TPSA are predicted to have better absorption [104]. Out of 28 similar analogs, 12 compounds (42%) showed potent activity against whipworm and hookworm parasites, with multiple compounds showing promising activity against the two parasites. We consider F0317-0202 and at least six of its analogs, F0317-0161, F0317-0013, F0328-0210, and F0317-0009, F0317-0019, and F0317-0160, as well-positioned for further characterization, mode of action studies, lead optimization, cytotoxicity studies in mammalian cells, and *in vivo* validation.

In contrast to screening diversity sets or random compound collections, we also screened repurposing libraries, which contain compounds already approved or associated with clinical trials and which can serve as a source for novel therapeutics for new indications [105]. The Broad Institute repurposing library, which we used in our screen, contains 6,743 compounds from molecules with high structural diversity and with 2006 known protein targets, including but not limited to the G protein-coupled receptors (GPCR), acetylcholine receptors, protein kinases, oxygenases, voltage-gated channels, and nuclear hormone receptors[106]. In addition, screening compounds with known targets can be highly beneficial, especially against NTDs where a deep understanding of their biology is still lacking. Compounds with known modes of action offer a significant advantage in drug development because they can act as probes for essential targets, assuming the targets are known. For this reason, we also screened the ICCB-Longwood mechanism of action library containing 1,245 high-quality compounds with known targets.

The total number of compounds tested from both libraries was 7,988, uncovering 45 compounds with putative broad activity against parasitic nematodes, giving a hit rate of 0.56%, 12.4 times that obtained from the generic diversity set library. These results are the first to thoroughly compare a generic dissimilarity-based library type with repurposing/known mode of action compounds against parasitic nematodes using the same screening conditions and pipeline. Our results indicate that repurposing/known mode of action libraries are significantly more (>10X; $P < 0.00001$ Fisher's exact test) likely to uncover active compounds.

The plethora of the chemical universe is expected to contain 10^{12} to 10^{180} drug-like compounds[107], which are impossible to screen against any single organism, not to mention GINs.

Therefore, techniques and strategies to design chemical libraries covering essential pharmacological targets are highly recommended. Among these approaches is the compilation of a target-focused library of compounds, which can be fitted by molecular docking to bind and inhibit the function of a specific target or protein family[104, 108]. This strategy has been previously validated using a kinase-targeted library of 1440 compounds tailored to fit in a total of 41 kinases from five different families, showing a 6.7-fold increase in the overall hits compared to the generic collection of molecules[109]. Target-focused screening improves efficiency by excluding compounds unlikely to modulate known target proteins of interest and is associated with increased potency and specificity[110].

We found that target-based libraries could serve as good sources for anthelmintic discovery. We screened nine target-focused libraries, including one for neuronal targets, one for GPCRs, and seven libraries targeting protein kinases. Neuronal signaling and the neuromuscular system are essential for nematodes' survival, including feeding, mating, egg laying, and locomotion; thus, many anthelmintics drugs target the nematode nervous system[111]. Neuropeptide GPCRs have been identified as potential and druggable targets and were found to play critical roles in the helminth neuromuscular function[112]. Kinases are the second most targeted protein class after GPCRs, with over 200 kinase inhibitors having reached the clinical phases and 52 FDA-approved drugs targeting protein kinases[39, 113].

Together, these nine libraries contain 2,708 compounds, and within these, we identified 12 compounds with putative broad activity, giving a hit rate of 0.44%, statistically the same as that obtained from the combined repurposing and mode of action libraries (P=0.54, Fisher's exact

test). It is worth noting that three of the nine target-based libraries resulted in no hits with activity against hookworms and whipworms: the ChemBridge Focused kinase, GPCR, and the LINCS 2 kinase inhibitors, which could be explained by the lack of specificity to the parasitic kinases and GPCRs, as these libraries were fitted to selectively binding specific human targets.

Historically, natural products have been the primary source of remedies for human ailments, and purified natural molecules or direct derivatives have contributed to at least 34% of all drugs approved by the FDA between 1981 and 2010[114]. Using our novel pipeline, we screened TimTec libraries of 4,240 purified compounds derived from or synthetically based upon natural products (see Materials and Methods). We identified 14 compounds with *A. ceylanicum* larval activity. However, none of these larvicidal compounds carried activity against adult parasites. This result may reflect biases of these particular libraries (see Materials and Methods).

In summary, we have developed a novel high throughput screening pipeline to identify anthelmintic compounds for GIN parasites and screened 13 different libraries with a total of 30,238 compounds that led to the discovery of 55 compounds with potent activity against two species of evolutionary distant nematode parasites. Repurposing, mode of action, and target-based libraries were more profitable than the diversity set for anthelmintic screening. For these 55 compounds, we reported the known modes of action and related targets in other organisms, and potential counterparts in the nematode, if available. As an example, we highlighted four nematode targets as potential candidates for target-based drug screening. We also identified thiazolyl oxopropanenitrile as a novel chemical scaffold with broader anthelmintic activity identified from the diversity set library and established the initial SAR models for potent activity

against hookworms and whipworms. Our findings open the doors for multiple follow-up approaches, including target identification, target-based screening, and ligand or structure-based pharmacophore optimization for developing safe and parasite-selective molecules.

Materials and Methods:

Ethics statement:

All experiments involving animals were conducted per the recommendation of the National Institute of Health (USA) Guide for Care and Use of Laboratory Animals and the Animal Welfare Act. The animal protocols used in this study (PROTO202000044 and PROTO202000071) were approved by the Institutional Animal Care and Use Committee (IACUC) of UMASS Chan Medical School.

Parasite maintenance in laboratory animals: Following standard protocols, *A. ceylanicum* hookworm parasites were maintained in golden Syrian hamsters[115], and *T. muris* whipworm parasites were maintained in STAT6-/- mice [116].

Primary screening using Larval Developmental Assay:

Feces from *A. ceylanicum*-infected hamsters were collected ~18 days post-infection and about 20 grams of feces were processed for egg isolation and purification following salt/sugar flotation protocol[117, 118]. Purified and surface sterilized eggs were incubated in S-medium[119] at 28 °C, without any food source for about 20 hours, allowing eggs to hatch into synchronized first-stage larvae [118]. Newly hatched L1 were washed three times in a S-medium. About 30-35 L1 larvae were dispensed into each well of 96-well assay plates containing 90µL of S-media supplemented

with 100U/ml of penicillin, 100µg/ml streptomycin, and 2.5µg/ml amphotericin, *Escherichia coli*

OP50 with final OD600 of 0.3 and 0.15µl Dimethylsulfoxide (DMSO). Library plates were

diluted in S-media to 100µM and 1%DMSO. Using a multichannel pipette, 10µL of the library

plates were transferred into each well of the assay plates, giving a final concentration of 10μM of

test drug and 0.25% DMSO. Plates were prepared in duplicates. Test plates were incubated at 28

°C for seven days and then scored for development under a dissecting microscope. Test drugs

were considered hit if > 90% of L1 larvae failed to develop in duplicate wells. Each assay plate

includes four wells of 10µM ivermectin as positive control and four wells of 0.25% DMSO as

negative control. For a group of 12 hamsters collecting feces twice weekly, the throughput for

this assay is ~2200 compounds/week in duplicate. Cherry-picked compounds for follow up study

(secondary screen), were obtained from the original library sources (Broad, Harvard, UMASS

Chan Med) independent of the initial library plates given to us for the primary screen.

Secondary screening using young adults of hookworm parasites:

We next screened hits from the primary screening (larval development) against the adult stages

of hookworm parasites. *A. ceylanicum* hookworm young adults were harvested 10-11 days post-

inoculation from the small intestines of infected hamsters. Harvested worms were washed and

incubated in prewarmed hookworm culture medium (RPMI1640 of pH 7.2 supplemented with

10% fetal bovine serum, 100U penicillin, 100µg/ml streptomycin, 2.5µg/ml amphotericin). Two

adults of *A. ceylanicum* hookworm were manually picked into each well of the 96-well assay

plates, containing 70µL of hookworm medium supplemented with 0.7µl DMSO. Using a

multichannel pipette, 30µL from the library plates diluted in hookworm medium to a final

concentration of 100µM and 1% DMSO were dispensed into screening wells, giving a final

concentration in assay plates of 30µM of test compounds and 1% DMSO with duplicate wells/drug. Assay plates were incubated for 48 hours at 37 °C and 5% CO_2. Drug activity was determined using the standard motility index [31, 116]. A motility index of 3 was given to vigorous healthy adults, 2 for motile but slow adults, 1 for immotile adults that moved after stimulation by touching, and 0 for non-motile adults even after touching. The drug was considered a hit if the motility index of at least three of the four *A. ceylanicum* hookworm adults were scored ≤1 and/or if all the parasites in the well had significant morphological defects (*e.g.*, shrunken cuticles, darkened intestines (Figure 3). Each assay plate included four positive control wells (30µM ivermectin) and four negative control wells (1% DMSO). Cherry-picked compounds for follow up study (tertiary screen), were obtained from the original library sources (Broad, Harvard, UMASS Chan Med) independent of the library plates given to us for the primary and secondary screens.

Tertiary screening using adult whipworm parasites:

We next screened *A. ceylanicum* hookworm adult hits against *T. muris* whipworm parasites. *T. muris* adult whipworms were harvested from the cecum and large intestines of infected STAT6 -/- mice. Single worms were manually picked into each well of a 48-well plate, six wells/ test drug, and a final volume of 291µL of whipworm culture medium with 2.1µl DMSO (like hookworm culture medium but supplemented with 5% fetal bovine serum). 9µL from library plates diluted in whipworm medium to a final concentration of 1mM and 10% DMSO were dispensed into each well of the screening plates, giving a final concentration of 30µM test drug and 1% DMSO. Assay plates were incubated at 37 °C, and 5% CO_2. *T. muris* whipworm motility was measured using an in-house assembled worminator and worm assay software[120] for

3 minutes after 24 and 48 hours. Drug activity was determined using the percent of relative motility compared to DMSO control. Drugs were considered active if the 24- and/or 48-hour average motility of six wells (1 worm/well) was reduced by at least 50% compared to negative control (Figure 4).

Chemical Libraries Screened

Libraries are provided by the Broad Institute (REPO), Harvard Medical School (ICCB), and UMASS Chan Small Molecules Screening Facility (SMSF), all of which have the purity and identity of their libraries assured by the source company or in-house using liquid chromatography-mass spectrometry.

Life Chemicals 15K diversity set:

Life Chemical pre-plated diversity set library was obtained from SMSF, UMASS Chan Medical School. The library contains 15,360 compounds representing 1,717 chemical scaffolds with an average Tanimoto similarity index of 0.409. Library compounds provided in 96-well plates with 10 mM stock concentration in 100% DMSO. Hits from the primary and secondary screening were cherrypicked into 96-well plates for *A. ceylanicum* hookworms or *T. muris* whipworms screening.

Broad Institute Repurposing Hub:

The drug repurposing hub includes 6,743 compounds which were from a curated and annotated collection of preclinical drugs in phase 1 (REPO 1) or phases 2-3 (REPO 2) and FDA-approved and launched drugs (REPO 3). REPO drugs cover a wide range of disease areas, including but

not limited to infectious diseases, allergies, cardiology, dermatology, endocrinology, metabolism, neurology, and gastroenterology. The library covers 1,133 different modes of action and 2,183 different drug targets[106]. Library compounds provided in 96-well plates with 10 mM stock concentration in 100% DMSO. Hits from the primary and secondary screening were cherrypicked from the source for *A. ceylanicum* hookworms or *T. muris* whipworms screening.

ICCB Screening libraries:

The Institute of Chemistry and Cell Biology-Longwood (ICCB-L) Screening Program at Harvard Medical School offers a wide range of well-annotated drug screening libraries. We have screened multiple ICCB-L libraries, including mechanism of action, neuronal targets, GPCR-based inhibitors, and six libraries of kinase inhibitors.

ICCB-L MOA: the mechanism of the action library is a collection of high-quality annotated compounds with biological activity against known targets with target-compound pairs based on high target specificity and potency. The library contains 1,245 unique compounds, each plated in 4 doses (10, 2, 0.4, 0.08 mM) in 100% DMSO. *ICCB-L Selleck Neuronal Targets:* the neuronal targets library was initially composed of Selleck chemicals from FDA-approved compound sets that were determined to have biological activity in neurologic research and neurological assays. Library compounds were provided in 96-well plates with 10mM stock concentration in 100% DMSO.*ICCB-L ChemBridge focused GPCR:* This focused library of 250 GPCR inhibitors was selected from 13,000 GPCR inhibitors assembled by ChemBridge. The library has 11 different scaffolds, each containing 19-24 compounds. Library compounds were provided in 96-well plates with 10mM stock concentration in 100% DMSO.

Kinase inhibitor libraries: in total, we screened focused sets of 1428 kinase inhibitors from 7 different kinase libraries, some of which were screened at multiple doses.

ICCB-L SYNthesis kinase inhibitor: this library was assembled by SYNthesis Med Chem of 96 compounds known to inhibit different kinases, each plated in 3 doses (10, 2, 0.4 mM) in 100%DMSO. *ICCB-L EMD Kinase inhibitors:* this library was built by EMD of 244 protein kinase inhibitors. Library compounds were provided in 96-well plates with 10mM stock concentration in 100% DMSO. *ICCB-L ChemBridge Focused Kinase inhibitors:* This library of 250 compounds was created and selected from the ChemBridge kinase-biased library of 6000 compounds. These compounds were selected based on the pharmacophore properties of known kinase inhibitors or from compounds predicted to interact with the ATP binding site of kinases. In total, these compounds represent 16 different scaffolds that cover 34 kinase targets. Library compounds were provided in 96-well plates with 10mM stock concentration in 100% DMSO. *ICCB-L LINCS 1,2 and 4 Kinase inhibitors:* these three libraries were assembled by LINCS together, including 345 compounds with known or predicted kinase inhibitors, each plated in 4 doses (10, 2, 0.4, and 0.08 mM) in 100%DMSO.

SMSF kinase inhibitor library: This library includes 429 kinase inhibitors with activity and safety confirmed by preclinical research and clinical trials. Library compounds were provided in 96-well plates with 10 mM stock concentration in 100% DMSO.

Natural Products: we screened the TimTec library of 4240 purified compounds from natural products or derivatives. Library compounds were provided in 96-well plates with 10 mM stock

concentration in 100% DMSO. The TimTec library was obtained from the SMSF and includes

two types of libraries with no overlap between them: (1) the Natural Products Library (NPL),

which contains compounds are primarily sourced from plants with the remaining samples from

bacteria, fungus, and animal source; and (2) the Natural Derivatives Library (NDL), which is a

hybrid between pure natural molecules and synthetic organic chemistry, derived from

publications and TimTec in-house material. The NDL elaborates on structural diversity of pure

natural compounds including natural derivatives, analogs, semi-natural compounds, and mimics.

Structure-Activity Relationship (SAR) of anthelmintic hits: Using vendor databases and

online tools, 28 analogs of F0317-0202 were identified and purchased from Life Chemicals.

Each analog has slight structural modification compared to the parent molecule. These analogs

were tested against the adult stages of *T. muris* whipworm and *A. ceylanicum* hookworm

parasites at 30μM and scored using the Worminator at 24 and 48 hours. Hence, *A. ceylanicum*

hookworm parasites used in testing F0317-0202 analogs for SAR model were mature adults

harvested 18 days post-infection.

Dose-response against hookworms and whipworms:

 To better evaluate the 12 potential analogs of F0317-0202 for broad-spectrum activity compared

to their parent molecules, these analogs were screened against hookworms and whipworms using

a dose-response of 0, 1, 3, 10, 30, and 90μM. Each dose was tested against 6 adult worms. Drug

activity was determined using a worminator motility readout. GraphPad Prism generated the IC_{50}

values utilizing the 48-hour average normalized motility of 6 individual worms.

Authors contributions:

M.A.E and R.V.A., conceived, designed, supervised, and interpreted the experiments. M.A.E and R.V.A. wrote the manuscript. M.A.E, Y.M.K., E.G., and P.C. performed the *in vitro* screening. M.A.E and S.S. performed data mining and analyzed SAR results. P.R.T. and L.B. contributed libraries, technical advice, and inputs on the manuscript. M.A.E. designed the graphic abstract. R.V.A designed the TOC graphic. R.V.A. secured funding.

Supporting information:

Supporting Information Available free of charge: Additional details for the potential nematode targets and additional experimental details for F0317-0202 tested analogs, chemical structures, and motility scores (Figure S1) (PDF).

Notes:

The authors declare the following competing financial interest(s): MAE and RVA are inventors on patents related to the compounds discussed in this manuscript, which have been filed by UMASS Chan Medical School. These patents cover the use of these compounds in the treatment of parasitic worm infections. No other conflicts of interest are reported.

Acknowledgments:

This work was financially supported by the National Institute of Health (USA) – National Institute of Allergy and Infectious Diseases grants R01 AI50866 and R01 AI056189 both to R.V.A.

References

1.	Hotez, P. J.; Kamath, A., Neglected tropical diseases in sub-saharan Africa: review of their prevalence, distribution, and disease burden. *PLoS Negl Trop Dis* **2009,** *3* (8), e412.

2.	Jourdan, P. M.; Lamberton, P. H. L.; Fenwick, A.; Addiss, D. G., Soil-transmitted helminth infections. *Lancet* **2018,** *391* (10117), 252-265.

3.	Loukas, A.; Maizels, R. M.; Hotez, P. J., The yin and yang of human soil-transmitted helminth infections. *Int J Parasitol* **2021,** *51* (13-14), 1243-1253.

4.	Hamory, J.; Miguel, E.; Walker, M.; Kremer, M.; Baird, S., Twenty-year economic impacts of deworming. *Proc Natl Acad Sci U S A* **2021,** *118* (14).

5.	Blaxter, M.; Koutsovoulos, G., The evolution of parasitism in Nematoda. *Parasitology* **2015,** *142 Suppl 1* (Suppl 1), S26-39.

6.	Weatherhead, J. E.; Hotez, P. J.; Mejia, R., The Global State of Helminth Control and Elimination in Children. *Pediatr Clin North Am* **2017,** *64* (4), 867-877.

7.	Pullan, R. L.; Smith, J. L.; Jasrasaria, R.; Brooker, S. J., Global numbers of infection and disease burden of soil transmitted helminth infections in 2010. *Parasit Vectors* **2014,** *7*, 37.

8.	Bleakley, H., Disease and Development: Evidence from Hookworm Eradication in the American South. *Q J Econ* **2007,** *122* (1), 73-117.

9.	Stromberg, B. E.; Gasbarre, L. C.; Ballweber, L. R.; Dargatz, D. A.; Rodriguez, J. M.; Kopral, C. A.; Zarlenga, D. S., Prevalence of internal parasites in beef cows in the United States: Results of the National Animal Health Monitoring System's (NAHMS) beef study, 2007-2008. *Can J Vet Res* **2015,** *79* (4), 290-5.

10.	World Health Organization, Ending the Neglect to Attain the Sustainable Development Goals: A Road Map for Neglected Tropical Diseases 2021-2030. *World Health Organization* **2020**.

11.	Vercruysse, J.; Albonico, M.; Behnke, J. M.; Kotze, A. C.; Prichard, R. K.; McCarthy, J. S.; Montresor, A.; Levecke, B., Is anthelmintic resistance a concern for the control of human soil-transmitted helminths? *Int J Parasitol Drugs Drug Resist* **2011,** *1* (1), 14-27.

12.	Moser, W.; Schindler, C.; Keiser, J., Efficacy of recommended drugs against soil transmitted helminths: systematic review and network meta-analysis. *BMJ* **2017,** *358*, j4307.

13.	Kotze, A. C.; Gilleard, J. S.; Doyle, S. R.; Prichard, R. K., Challenges and opportunities for the adoption of molecular diagnostics for anthelmintic resistance. *Int J Parasitol Drugs Drug Resist* **2020,** *14*, 264-273.

14.	Wolstenholme, A. J.; Fairweather, I.; Prichard, R.; von Samson-Himmelstjerna, G.; Sangster, N. C., Drug resistance in veterinary helminths. *Trends Parasitol* **2004,** *20* (10), 469-76.

15.	Kaplan, R. M., Drug resistance in nematodes of veterinary importance: a status report. *Trends Parasitol* **2004,** *20* (10), 477-81.

16.	Kaplan, R. M.; Vidyashankar, A. N., An inconvenient truth: global worming and anthelmintic resistance. *Vet Parasitol* **2012,** *186* (1-2), 70-8.

17.	Geurden, T.; Hoste, H.; Jacquiet, P.; Traversa, D.; Sotiraki, S.; Frangipane di Regalbono, A.; Tzanidakis, N.; Kostopoulou, D.; Gaillac, C.; Privat, S.; Giangaspero, A.; Zanardello, C.; Noe, L.; Vanimisetti, B.; Bartram, D., Anthelmintic resistance and multidrug

resistance in sheep gastro-intestinal nematodes in France, Greece and Italy. *Vet Parasitol* **2014,** *201* (1-2), 59-66.

18.   Torres-Acosta, J. F.; Mendoza-de-Gives, P.; Aguilar-Caballero, A. J.; Cuellar-Ordaz, J. A., Anthelmintic resistance in sheep farms: update of the situation in the American continent. *Vet Parasitol* **2012,** *189* (1), 89-96.

19.   Coles, G. C.; Jackson, F.; Pomroy, W. E.; Prichard, R. K.; von Samson-Himmelstjerna, G.; Silvestre, A.; Taylor, M. A.; Vercruysse, J., The detection of anthelmintic resistance in nematodes of veterinary importance. *Vet Parasitol* **2006,** *136* (3-4), 167-85.

20.   Rose, H.; Rinaldi, L.; Bosco, A.; Mavrot, F.; de Waal, T.; Skuce, P.; Charlier, J.; Torgerson, P. R.; Hertzberg, H.; Hendrickx, G.; Vercruysse, J.; Morgan, E. R., Widespread anthelmintic resistance in European farmed ruminants: a systematic review. *Vet Rec* **2015,** *176* (21), 546.

21.   Zuccherato, L. W.; Furtado, L. F.; Medeiros, C. D. S.; Pinheiro, C. D. S.; Rabelo, E. M., PCR-RFLP screening of polymorphisms associated with benzimidazole resistance in Necator americanus and Ascaris lumbricoides from different geographical regions in Brazil. *PLoS Negl Trop Dis* **2018,** *12* (9), e0006766.

22.   Diawara, A.; Halpenny, C. M.; Churcher, T. S.; Mwandawiro, C.; Kihara, J.; Kaplan, R. M.; Streit, T. G.; Idaghdour, Y.; Scott, M. E.; Basanez, M. G.; Prichard, R. K., Association between response to albendazole treatment and beta-tubulin genotype frequencies in soil-transmitted helminths. *PLoS Negl Trop Dis* **2013,** *7* (5), e2247.

23.   Rashwan, N.; Scott, M.; Prichard, R., Rapid Genotyping of beta-tubulin Polymorphisms in Trichuris trichiura and Ascaris lumbricoides. *PLoS Negl Trop Dis* **2017,** *11* (1), e0005205.

24.   Rashwan, N.; Bourguinat, C.; Keller, K.; Gunawardena, N. K.; de Silva, N.; Prichard, R., Isothermal Diagnostic Assays for Monitoring Single Nucleotide Polymorphisms in Necator americanus Associated with Benzimidazole Drug Resistance. *PLoS Negl Trop Dis* **2016,** *10* (12), e0005113.

25.   Krucken, J.; Fraundorfer, K.; Mugisha, J. C.; Ramunke, S.; Sifft, K. C.; Geus, D.; Habarugira, F.; Ndoli, J.; Sendegeya, A.; Mukampunga, C.; Bayingana, C.; Aebischer, T.; Demeler, J.; Gahutu, J. B.; Mockenhaupt, F. P.; von Samson-Himmelstjerna, G., Reduced efficacy of albendazole against Ascaris lumbricoides in Rwandan schoolchildren. *Int J Parasitol Drugs Drug Resist* **2017,** *7* (3), 262-271.

26.   Muller, I.; Beyleveld, L.; Gerber, M.; Puhse, U.; du Randt, R.; Utzinger, J.; Zondie, L.; Walter, C.; Steinmann, P., Low efficacy of albendazole against Trichuris trichiura infection in schoolchildren from Port Elizabeth, South Africa. *Trans R Soc Trop Med Hyg* **2016,** *110* (11), 676-678.

27.   Soukhathammavong, P. A.; Sayasone, S.; Phongluxa, K.; Xayaseng, V.; Utzinger, J.; Vounatsou, P.; Hatz, C.; Akkhavong, K.; Keiser, J.; Odermatt, P., Low efficacy of single-dose albendazole and mebendazole against hookworm and effect on concomitant helminth infection in Lao PDR. *PLoS Negl Trop Dis* **2012,** *6* (1), e1417.

28.   Harder, A.; Holden-Dye, L.; Walker, R.; Wunderlich, F., Mechanisms of action of emodepside. *Parasitol Res* **2005,** *97 Suppl 1*, S1-S10.

29.   Shaver, A. O.; Wit, J.; Dilks, C. M.; Crombie, T. A.; Li, H.; Aroian, R. V.; Andersen, E. C., Variation in anthelmintic responses are driven by genetic differences among diverse C. elegans wild strains. *PLoS Pathog* **2023,** *19* (4), e1011285.

30.   Wit, J.; Rodriguez, B. C.; Andersen, E. C., Natural variation in Caenorhabditis elegans responses to the anthelmintic emodepside. *Int J Parasitol Drugs Drug Resist* **2021,** *16*, 1-8.

31.   Elfawal, M. A.; Savinov, S. N.; Aroian, R. V., Drug Screening for Discovery of Broad-spectrum Agents for Soil-transmitted Nematodes. *Sci Rep* **2019,** *9* (1), 12347.

32.   Herath, H.; Taki, A. C.; Rostami, A.; Jabbar, A.; Keiser, J.; Geary, T. G.; Gasser, R. B., Whole-organism phenotypic screening methods used in early-phase anthelmintic drug discovery. *Biotechnol Adv* **2022,** *57*, 107937.

33.   Tyagi, R.; Rosa, B. A.; Lewis, W. G.; Mitreva, M., Pan-phylum Comparison of Nematode Metabolic Potential. *PLoS Negl Trop Dis* **2015,** *9* (5), e0003788.

34.   Tyagi, R.; Maddirala, A. R.; Elfawal, M.; Fischer, C.; Bulman, C. A.; Rosa, B. A.; Gao, X.; Chugani, R.; Zhou, M.; Helander, J.; Brindley, P. J.; Tseng, C. C.; Greig, I. R.; Sakanari, J.; Wildman, S. A.; Aroian, R.; Janetka, J. W.; Mitreva, M., Small Molecule Inhibitors of Metabolic Enzymes Repurposed as a New Class of Anthelmintics. *ACS Infect Dis* **2018,** *4* (7), 1130-1145.

35.   Tyagi, R.; Elfawal, M. A.; Wildman, S. A.; Helander, J.; Bulman, C. A.; Sakanari, J.; Rosa, B. A.; Brindley, P. J.; Janetka, J. W.; Aroian, R. V.; Mitreva, M., Identification of small molecule enzyme inhibitors as broad-spectrum anthelmintics. *Sci Rep* **2019,** *9* (1), 9085.

36.   Keiser, J.; Panic, G.; Adelfio, R.; Cowan, N.; Vargas, M.; Scandale, I., Evaluation of an FDA approved library against laboratory models of human intestinal nematode infections. *Parasit Vectors* **2016,** *9* (1), 376.

37.   Tritten, L.; Nwosu, U.; Vargas, M.; Keiser, J., In vitro and in vivo efficacy of tribendimidine and its metabolites alone and in combination against the hookworms Heligmosomoides bakeri and Ancylostoma ceylanicum. *Acta Trop* **2012,** *122* (1), 101-7.

38.   Coghlan, A.; Partridge, F. A.; Duque-Correa, M. A.; Rinaldi, G.; Clare, S.; Seymour, L.; Brandt, C.; Mkandawire, T. T.; McCarthy, C.; Holroyd, N.; Nick, M.; Brown, A. E.; Tonitiwong, S.; Sattelle, D. B.; Berriman, M., A drug repurposing screen for whipworms informed by comparative genomics. *PLoS Negl Trop Dis* **2023,** *17* (9), e0011205.

39.   Knox, J.; Joly, N.; Linossi, E. M.; Carmona-Negron, J. A.; Jura, N.; Pintard, L.; Zuercher, W.; Roy, P. J., A survey of the kinome pharmacopeia reveals multiple scaffolds and targets for the development of novel anthelmintics. *Sci Rep* **2021,** *11* (1), 9161.

40.   Partridge, F. A.; Brown, A. E.; Buckingham, S. D.; Willis, N. J.; Wynne, G. M.; Forman, R.; Else, K. J.; Morrison, A. A.; Matthews, J. B.; Russell, A. J.; Lomas, D. A.; Sattelle, D. B., An automated high-throughput system for phenotypic screening of chemical libraries on C. elegans and parasitic nematodes. *Int J Parasitol Drugs Drug Resist* **2018,** *8* (1), 8-21.

41.   O'Reilly, L. P.; Luke, C. J.; Perlmutter, D. H.; Silverman, G. A.; Pak, S. C., C. elegans in high-throughput drug discovery. *Adv Drug Deliv Rev* **2014,** *69-70*, 247-53.

42.   Burns, A. R.; Luciani, G. M.; Musso, G.; Bagg, R.; Yeo, M.; Zhang, Y.; Rajendran, L.; Glavin, J.; Hunter, R.; Redman, E.; Stasiuk, S.; Schertzberg, M.; Angus McQuibban, G.; Caffrey, C. R.; Cutler, S. R.; Tyers, M.; Giaever, G.; Nislow, C.; Fraser, A. G.; MacRae, C. A.; Gilleard, J.; Roy, P. J., Caenorhabditis elegans is a useful model for anthelmintic discovery. *Nat Commun* **2015,** *6*, 7485.

43.   Keiser, J.; Tritten, L.; Adelfio, R.; Vargas, M., Effect of combinations of marketed human anthelmintic drugs against Trichuris muris in vitro and in vivo. *Parasit Vectors* **2012,** *5*, 292.

44.   Buxton, S. K.; Neveu, C.; Charvet, C. L.; Robertson, A. P.; Martin, R. J., On the mode of action of emodepside: slow effects on membrane potential and voltage-activated currents in Ascaris suum. *Br J Pharmacol* **2011,** *164* (2b), 453-70.

45.   Holden-Dye, L.; Crisford, A.; Welz, C.; von Samson-Himmelstjerna, G.; Walker, R. J.; O'Connor, V., Worms take to the slo lane: a perspective on the mode of action of emodepside. *Invert Neurosci* **2012,** *12* (1), 29-36.

46.   Robson Duarte, E.; Duarte Matias, A.; Almeida Bastos, G.; Cordeiro Maia, R.; Soares Martins, V. J.; Maia Soares, A. C.; Dos Santos Magaco, F.; de Oliveira, N. J. F.; Dos Santos, T. A. X., Anthelmintic efficacy of trichlorfon and blood parameters of young lambs infected with Haemonchus contortus. *Vet Parasitol* **2019,** *272*, 40-43.

47.   Camacho, M.; Tarrab-Hazdai, R.; Espinoza, B.; Arnon, R.; Agnew, A., The amount of acetylcholinesterase on the parasite surface reflects the differential sensitivity of schistosome species to metrifonate. *Parasitology* **1994,** *108 (Pt 2)*, 153-60.

48.   Schwendeman, A. R.; Shaham, S., A High-Throughput Small Molecule Screen for C. elegans Linker Cell Death Inhibitors. *PLoS One* **2016,** *11* (10), e0164595.

49.   Levitzki, A.; Gazit, A., Tyrosine kinase inhibition: an approach to drug development. *Science* **1995,** *267* (5205), 1782-8.

50.   Nath, A. K.; Ryu, J. H.; Jin, Y. N.; Roberts, L. D.; Dejam, A.; Gerszten, R. E.; Peterson, R. T., PTPMT1 Inhibition Lowers Glucose through Succinate Dehydrogenase Phosphorylation. *Cell Rep* **2015,** *10* (5), 694-701.

51.   Liu, M.; Landuyt, B.; Klaassen, H.; Geldhof, P.; Luyten, W., Screening of a drug repurposing library with a nematode motility assay identifies promising anthelmintic hits against Cooperia oncophora and other ruminant parasites. *Vet Parasitol* **2019,** *265*, 15-18.

52.   Al-Gayyar, M. M. H.; Alattar, A.; Alshaman, R.; Hamdan, A. M., QNZ alleviated hepatocellular carcinoma by targeting inflammatory pathways in a rat model. *Cytokine* **2021,** *148*, 155710.

53.   Parrizas, M.; Gazit, A.; Levitzki, A.; Wertheimer, E.; LeRoith, D., Specific inhibition of insulin-like growth factor-1 and insulin receptor tyrosine kinase activity and biological function by tyrphostins. *Endocrinology* **1997,** *138* (4), 1427-33.

54.   Stuehr, D. J.; Fasehun, O. A.; Kwon, N. S.; Gross, S. S.; Gonzalez, J. A.; Levi, R.; Nathan, C. F., Inhibition of macrophage and endothelial cell nitric oxide synthase by diphenyleneiodonium and its analogs. *FASEB J* **1991,** *5* (1), 98-103.

55.   Almahariq, M.; Tsalkova, T.; Mei, F. C.; Chen, H.; Zhou, J.; Sastry, S. K.; Schwede, F.; Cheng, X., A novel EPAC-specific inhibitor suppresses pancreatic cancer cell migration and invasion. *Mol Pharmacol* **2013,** *83* (1), 122-8.

56.   Chen, H.; Tsalkova, T.; Chepurny, O. G.; Mei, F. C.; Holz, G. G.; Cheng, X.; Zhou, J., Identification and characterization of small molecules as potent and specific EPAC2 antagonists. *J Med Chem* **2013,** *56* (3), 952-62.

57.   Goard, C. A.; Schimmer, A. D., An evidence-based review of obatoclax mesylate in the treatment of hematological malignancies. *Core Evid* **2013,** *8*, 15-26.

58.   Symersky, J.; Osowski, D.; Walters, D. E.; Mueller, D. M., Oligomycin frames a common drug-binding site in the ATP synthase. *Proc Natl Acad Sci U S A* **2012,** *109* (35), 13961-5.

59.   Xiong, Q.; Sun, H.; Li, M., Zinc pyrithione-mediated activation of voltage-gated KCNQ potassium channels rescues epileptogenic mutants. *Nat Chem Biol* **2007,** *3* (5), 287-96.

60.   Arora, S.; Yang, J. M.; Kinzy, T. G.; Utsumi, R.; Okamoto, T.; Kitayama, T.; Ortiz, P. A.; Hait, W. N., Identification and characterization of an inhibitor of eukaryotic elongation factor 2 kinase against human cancer cell lines. *Cancer Res* **2003,** *63* (20), 6894-9.

61.   Martin, R. J.; Robertson, A. P., Mode of action of levamisole and pyrantel, anthelmintic resistance, E153 and Q57. *Parasitology* **2007,** *134* (Pt 8), 1093-104.
62.   Wang, J.; Grishin, A. V.; Ford, H. R., Experimental Anti-Inflammatory Drug Semapimod Inhibits TLR Signaling by Targeting the TLR Chaperone gp96. *J Immunol* **2016,** *196* (12), 5130-7.
63.   Sugahara, K.; Maeda, Y.; Shimano, K.; Mogami, A.; Kataoka, H.; Ogawa, K.; Hikida, K.; Kumagai, H.; Asayama, M.; Yamamoto, T.; Harada, T.; Ni, P.; Inoue, S.; Kawaguchi, A., Amiselimod, a novel sphingosine 1-phosphate receptor-1 modulator, has potent therapeutic efficacy for autoimmune diseases, with low bradycardia risk. *Br J Pharmacol* **2017,** *174* (1), 15-27.
64.   Shahinas, D.; Debnath, A.; Benedict, C.; McKerrow, J. H.; Pillai, D. R., Heat shock protein 90 inhibitors repurposed against Entamoeba histolytica. *Front Microbiol* **2015,** *6*, 368.
65.   Long, Y.; Chen, W.; Lin, Z.; Sun, H.; Xia, M.; Zheng, W.; Li, Z., Inhibition of HERG potassium channels by domiphen bromide and didecyl dimethylammonium bromide. *Eur J Pharmacol* **2014,** *737*, 202-9.
66.   Choi, J. Y.; Podust, L. M.; Roush, W. R., Drug strategies targeting CYP51 in neglected tropical diseases. *Chem Rev* **2014,** *114* (22), 11242-71.
67.   Sells, T. B.; Chau, R.; Ecsedy, J. A.; Gershman, R. E.; Hoar, K.; Huck, J.; Janowick, D. A.; Kadambi, V. J.; LeRoy, P. J.; Stirling, M.; Stroud, S. G.; Vos, T. J.; Weatherhead, G. S.; Wysong, D. R.; Zhang, M.; Balani, S. K.; Bolen, J. B.; Manfredi, M. G.; Claiborne, C. F., MLN8054 and Alisertib (MLN8237): Discovery of Selective Oral Aurora A Inhibitors. *ACS Med Chem Lett* **2015,** *6* (6), 630-4.
68.   Weeks, J. C.; Roberts, W. M.; Leasure, C.; Suzuki, B. M.; Robinson, K. J.; Currey, H.; Wangchuk, P.; Eichenberger, R. M.; Saxton, A. D.; Bird, T. D.; Kraemer, B. C.; Loukas, A.; Hawdon, J. M.; Caffrey, C. R.; Liachko, N. F., Sertraline, Paroxetine, and Chlorpromazine Are Rapidly Acting Anthelmintic Drugs Capable of Clinical Repurposing. *Sci Rep* **2018,** *8* (1), 975.
69.   Harrold, M. W.; Chang, Y. A.; Wallace, R. A.; Farooqui, T.; Wallace, L. J.; Uretsky, N.; Miller, D. D., Charged analogues of chlorpromazine as dopamine antagonists. *J Med Chem* **1987,** *30* (9), 1631-5.
70.   Ondeykal, J. G.; Herath, K. B.; Jayasuriya, H.; Polishook, J. D.; Bills, G. F.; Dombrowski, A. W.; Mojena, M.; Koch, G.; DiSalvo, J.; DeMartino, J.; Guan, Z.; Nanakorn, W.; Morenberg, C. M.; Balick, M. J.; Stevenson, D. W.; Slattery, M.; Borris, R. P.; Singh, S. B., Discovery of structurally diverse natural product antagonists of chemokine receptor CXCR3. *Mol Divers* **2005,** *9* (1-3), 123-9.
71.   Perregaard, J.; Moltzen, E. K.; Meier, E.; Sanchez, C., Sigma ligands with subnanomolar affinity and preference for the sigma 2 binding site. 1. 3-(omega-aminoalkyl)-1H-indoles. *J Med Chem* **1995,** *38* (11), 1998-2008.
72.   Mellin, T. N.; Busch, R. D.; Wang, C. C., Postsynaptic inhibition of invertebrate neuromuscular transmission by avermectin B1a. *Neuropharmacology* **1983,** *22* (1), 89-96.
73.   Martin, R. J., An electrophysiological preparation of Ascaris suum pharyngeal muscle reveals a glutamate-gated chloride channel sensitive to the avermectin analogue, milbemycin D. *Parasitology* **1996,** *112 (Pt 2)*, 247-52.
74.   Schaeffer, J. M.; Haines, H. W., Avermectin binding in Caenorhabditis elegans. A two-state model for the avermectin binding site. *Biochem Pharmacol* **1989,** *38* (14), 2329-38.
75.   Sun, E. L.; Liu, C. X.; Ma, Z. X.; Mou, X. Y.; Mu, X. A.; Ni, Y. H.; Li, X. L.; Zhang, D.; Ju, Y. R., Knockdown of human serine/threonine kinase 33 suppresses human small

cell lung carcinoma by blocking RPS6/BAD signaling transduction. *Neoplasma* **2017,** *64* (6), 869-879.

76.   Carmona, S.; Brunel, J. M.; Bonier, R.; Sbarra, V.; Robert, S.; Borentain, P.; Lombardo, D.; Mas, E.; Gerolami, R., A squalamine derivative, NV669, as a novel PTP1B inhibitor: in vitro and in vivo effects on pancreatic and hepatic tumor growth. *Oncotarget* **2019,** *10* (62), 6651-6667.

77.   Mauvezin, C.; Neufeld, T. P., Bafilomycin A1 disrupts autophagic flux by inhibiting both V-ATPase-dependent acidification and Ca-P60A/SERCA-dependent autophagosome-lysosome fusion. *Autophagy* **2015,** *11* (8), 1437-8.

78.   Magnaghi, P.; D'Alessio, R.; Valsasina, B.; Avanzi, N.; Rizzi, S.; Asa, D.; Gasparri, F.; Cozzi, L.; Cucchi, U.; Orrenius, C.; Polucci, P.; Ballinari, D.; Perrera, C.; Leone, A.; Cervi, G.; Casale, E.; Xiao, Y.; Wong, C.; Anderson, D. J.; Galvani, A.; Donati, D.; O'Brien, T.; Jackson, P. K.; Isacchi, A., Covalent and allosteric inhibitors of the ATPase VCP/p97 induce cancer cell death. *Nat Chem Biol* **2013,** *9* (9), 548-56.

79.   Klionsky, L.; Tamir, R.; Gao, B.; Wang, W.; Immke, D. C.; Nishimura, N.; Gavva, N. R., Species-specific pharmacology of Trichloro(sulfanyl)ethyl benzamides as transient receptor potential ankyrin 1 (TRPA1) antagonists. *Mol Pain* **2007,** *3*, 39.

80.   Houck, K. A.; Borchert, K. M.; Hepler, C. D.; Thomas, J. S.; Bramlett, K. S.; Michael, L. F.; Burris, T. P., T0901317 is a dual LXR/FXR agonist. *Mol Genet Metab* **2004,** *83* (1-2), 184-7.

81.   Lehmann, J. M.; Kliewer, S. A.; Moore, L. B.; Smith-Oliver, T. A.; Oliver, B. B.; Su, J. L.; Sundseth, S. S.; Winegar, D. A.; Blanchard, D. E.; Spencer, T. A.; Willson, T. M., Activation of the nuclear receptor LXR by oxysterols defines a new hormone response pathway. *J Biol Chem* **1997,** *272* (6), 3137-40.

82.   Schultz, J. R.; Tu, H.; Luk, A.; Repa, J. J.; Medina, J. C.; Li, L.; Schwendner, S.; Wang, S.; Thoolen, M.; Mangelsdorf, D. J.; Lustig, K. D.; Shan, B., Role of LXRs in control of lipogenesis. *Genes Dev* **2000,** *14* (22), 2831-8.

83.   Jakel, H.; Nowak, M.; Moitrot, E.; Dehondt, H.; Hum, D. W.; Pennacchio, L. A.; Fruchart-Najib, J.; Fruchart, J. C., The liver X receptor ligand T0901317 down-regulates APOA5 gene expression through activation of SREBP-1c. *J Biol Chem* **2004,** *279* (44), 45462-9.

84.   Ladefoged, L. K.; Munro, L.; Pedersen, A. J.; Lummis, S. C. R.; Bang-Andersen, B.; Balle, T.; Schiott, B.; Kristensen, A. S., Modeling and Mutational Analysis of the Binding Mode for the Multimodal Antidepressant Drug Vortioxetine to the Human 5-HT(3A) Receptor. *Mol Pharmacol* **2018,** *94* (6), 1421-1434.

85.   Dale, E.; Grunnet, M.; Pehrson, A. L.; Frederiksen, K.; Larsen, P. H.; Nielsen, J.; Stensbol, T. B.; Ebert, B.; Yin, H.; Lu, D.; Liu, H.; Jensen, T. N.; Yang, C. R.; Sanchez, C., The multimodal antidepressant vortioxetine may facilitate pyramidal cell firing by inhibition of 5-HT(3) receptor expressing interneurons: An in vitro study in rat hippocampus slices. *Brain Res* **2018,** *1689*, 1-11.

86.   Nackenoff, A. G.; Simmler, L. D.; Baganz, N. L.; Pehrson, A. L.; Sanchez, C.; Blakely, R. D., Serotonin Transporter-Independent Actions of the Antidepressant Vortioxetine As Revealed Using the SERT Met172 Mouse. *ACS Chem Neurosci* **2017,** *8* (5), 1092-1100.

87.   Gong, L.; Giacomini, M. M.; Giacomini, C.; Maitland, M. L.; Altman, R. B.; Klein, T. E., PharmGKB summary: sorafenib pathways. *Pharmacogenet Genomics* **2017,** *27* (6), 240-246.

88.   Gani, O. A.; Engh, R. A., Protein kinase inhibition of clinically important staurosporine analogues. *Nat Prod Rep* **2010,** *27* (4), 489-98.

89.   Baumann, P.; Schneider, L.; Mandl-Weber, S.; Oduncu, F.; Schmidmaier, R., Simultaneous targeting of PI3K and mTOR with NVP-BGT226 is highly effective in multiple myeloma. *Anticancer Drugs* **2012,** *23* (1), 131-8.

90.   Garaguso, P., [Dithiazanine and Ancylostoma or Necator]. *Sem Med* **1961,** *119*, 2016-21.

91.   Ho Thi, S.; Brumpt, L.; Lapierre, J., [Treatment of strongyloidiasis by dithiazanine by prolonged administration. The necessity of combining all efficient technics for the control of results]. *Bull Soc Pathol Exot Filiales* **1961,** *54*, 1059-75.

92.   Mendes, F.; Fonseca, E. E.; Salum, N.; Guedes, R.; Garrido, I., [Dithiazanine in the treatment of intestinal helminthiasis]. *Arq Bras Med* **1961,** *51*, 179-86.

93.   Vakil, B. J.; Mehta, B. J., Effect of dithiazanine in ascariasis, trichuriasis, enterobiasis and ankylostomiasis. *J Indian Med Assoc* **1961,** *37*, 219-22.

94.   Villadolid, L. S., Treatment of helminthiasis with dithiazanine iodide. *N Y State J Med* **1961,** *61*, 1539-41.

95.   Colland, F.; Formstecher, E.; Jacq, X.; Reverdy, C.; Planquette, C.; Conrath, S.; Trouplin, V.; Bianchi, J.; Aushev, V. N.; Camonis, J.; Calabrese, A.; Borg-Capra, C.; Sippl, W.; Collura, V.; Boissy, G.; Rain, J. C.; Guedat, P.; Delansorne, R.; Daviet, L., Small-molecule inhibitor of USP7/HAUSP ubiquitin protease stabilizes and activates p53 in cells. *Mol Cancer Ther* **2009,** *8* (8), 2286-95.

96.   Doughty-Shenton, D.; Joseph, J. D.; Zhang, J.; Pagliarini, D. J.; Kim, Y.; Lu, D.; Dixon, J. E.; Casey, P. J., Pharmacological targeting of the mitochondrial phosphatase PTPMT1. *J Pharmacol Exp Ther* **2010,** *333* (2), 584-92.

97.   Kurzchalia, T. V.; Ward, S., Why do worms need cholesterol? *Nat Cell Biol* **2003,** *5* (8), 684-8.

98.   Valdes, V. J.; Athie, A.; Salinas, L. S.; Navarro, R. E.; Vaca, L., CUP-1 is a novel protein involved in dietary cholesterol uptake in Caenorhabditis elegans. *PLoS One* **2012,** *7* (3), e33962.

99.   Heimbucher, T.; Liu, Z.; Bossard, C.; McCloskey, R.; Carrano, A. C.; Riedel, C. G.; Tanasa, B.; Klammt, C.; Fonslow, B. R.; Riera, C. E.; Lillemeier, B. F.; Kemphues, K.; Yates, J. R., 3rd; O'Shea, C.; Hunter, T.; Dillin, A., The Deubiquitylase MATH-33 Controls DAF-16 Stability and Function in Metabolism and Longevity. *Cell Metab* **2015,** *22* (1), 151-63.

100.   Pereira, R. V.; de, S. G. M.; Olmo, R. P.; Souza, D. M.; Cabral, F. J.; Jannotti-Passos, L. K.; Baba, E. H.; Andreolli, A. B.; Rodrigues, V.; Castro-Borges, W.; Guerra-Sa, R., Ubiquitin-specific proteases are differentially expressed throughout the Schistosoma mansoni life cycle. *Parasit Vectors* **2015,** *8*, 349.

101.   Shearer, J.; Castro, J. L.; Lawson, A. D. G.; MacCoss, M.; Taylor, R. D., Rings in Clinical Trials and Drugs: Present and Future. *J Med Chem* **2022,** *65* (13), 8699-8712.

102.   Ertl, P.; Altmann, E.; McKenna, J. M., The Most Common Functional Groups in Bioactive Molecules and How Their Popularity Has Evolved over Time. *J Med Chem* **2020,** *63* (15), 8408-8418.

103.   Lipinski, C. A.; Lombardo, F.; Dominy, B. W.; Feeney, P. J., Experimental and computational approaches to estimate solubility and permeability in drug discovery and development settings. *Adv Drug Deliv Rev* **2001,** *46* (1-3), 3-26.

104.   Orry, A. J.; Abagyan, R. A.; Cavasotto, C. N., Structure-based development of target-specific compound libraries. *Drug Discov Today* **2006,** *11* (5-6), 261-6.

105.   Pushpakom, S.; Iorio, F.; Eyers, P. A.; Escott, K. J.; Hopper, S.; Wells, A.; Doig, A.; Guilliams, T.; Latimer, J.; McNamee, C.; Norris, A.; Sanseau, P.; Cavalla, D.; Pirmohamed,

M., Drug repurposing: progress, challenges and recommendations. *Nat Rev Drug Discov* **2019,** *18* (1), 41-58.

106.   Corsello, S. M.; Bittker, J. A.; Liu, Z.; Gould, J.; McCarren, P.; Hirschman, J. E.; Johnston, S. E.; Vrcic, A.; Wong, B.; Khan, M.; Asiedu, J.; Narayan, R.; Mader, C. C.; Subramanian, A.; Golub, T. R., The Drug Repurposing Hub: a next-generation drug library and information resource. *Nat Med* **2017,** *23* (4), 405-408.

107.   Gorse, A. D., Diversity in medicinal chemistry space. *Curr Top Med Chem* **2006,** *6* (1), 3-18.

108.   Harris, C. J.; Hill, R. D.; Sheppard, D. W.; Slater, M. J.; Stouten, P. F., The design and application of target-focused compound libraries. *Comb Chem High Throughput Screen* **2011,** *14* (6), 521-31.

109.   Gozalbes, R.; Simon, L.; Froloff, N.; Sartori, E.; Monteils, C.; Baudelle, R., Development and experimental validation of a docking strategy for the generation of kinase-targeted libraries. *J Med Chem* **2008,** *51* (11), 3124-32.

110.   Maly, D. J.; Choong, I. C.; Ellman, J. A., Combinatorial target-guided ligand assembly: identification of potent subtype-selective c-Src inhibitors. *Proc Natl Acad Sci U S A* **2000,** *97* (6), 2419-24.

111.   Wolstenholme, A. J., Ion channels and receptor as targets for the control of parasitic nematodes. *Int J Parasitol Drugs Drug Resist* **2011,** *1* (1), 2-13.

112.   Atkinson, L. E.; McCoy, C. J.; Crooks, B. A.; McKay, F. M.; McVeigh, P.; McKenzie, D.; Irvine, A.; Harrington, J.; Rosa, B. A.; Mitreva, M.; Marks, N. J.; Maule, A. G.; Mousley, A., Phylum-Spanning Neuropeptide GPCR Identification and Prioritization: Shaping Drug Target Discovery Pipelines for Nematode Parasite Control. *Front Endocrinol (Lausanne)* **2021,** *12*, 718363.

113.   Roskoski, R., Jr., Properties of FDA-approved small molecule protein kinase inhibitors: A 2020 update. *Pharmacol Res* **2020,** *152*, 104609.

114.   Harvey, A. L.; Edrada-Ebel, R.; Quinn, R. J., The re-emergence of natural products for drug discovery in the genomics era. *Nat Rev Drug Discov* **2015,** *14* (2), 111-29.

115.   Hu, Y.; Zhan, B.; Keegan, B.; Yiu, Y. Y.; Miller, M. M.; Jones, K.; Aroian, R. V., Mechanistic and single-dose in vivo therapeutic studies of Cry5B anthelmintic action against hookworms. *PLoS Negl Trop Dis* **2012,** *6* (11), e1900.

116.   Hu, Y.; Ellis, B. L.; Yiu, Y. Y.; Miller, M. M.; Urban, J. F.; Shi, L. Z.; Aroian, R. V., An extensive comparison of the effect of anthelmintic classes on diverse nematodes. *PLoS One* **2013,** *8* (7), e70702.

117.   Mes, T. H.; Ploeger, H. W.; Terlou, M.; Kooyman, F. N.; Van der Ploeg, M. P.; Eysker, M., A novel method for the isolation of gastro-intestinal nematode eggs that allows automated analysis of digital images of egg preparations and high throughput screening. *Parasitology* **2001,** *123* (Pt 3), 309-14.

118.   Li, H.; Gazzola, D.; Hu, Y.; Aroian, R. V., An efficient method for viable cryopreservation of hookworms and other gastrointestinal nematodes in the laboratory. *bioRxiv* **2023**.

119.   Stiernagle, T., Maintenance of C. elegans. *WormBook* **2006**, 1-11.

120.   Marcellino, C.; Gut, J.; Lim, K. C.; Singh, R.; McKerrow, J.; Sakanari, J., WormAssay: a novel computer application for whole-plate motion-based screening of macroscopic parasites. *PLoS Negl Trop Dis* **2012,** *6* (1), e1494.