Combined physical and pharmacological anabolic osteoporosis therapies increase bone response and mechanoregulation in female mice

Authors: Friederike A. Schulte[1], Francisco C. Marques[1], Julia K. Griesbach[1], Claudia Weigt[1], Marcella von Salis-Soglio[1], Floor M. Lambers[1], Clemens Kreutz[2], Michaela Kneissel[2], Peter J. Richards[2,3], Gisela A. Kuhn[1], Ralph Müller[1]*

[1]Institute for Biomechanics, ETH Zurich; Zurich, Switzerland.

[2]Novartis Biomedical Research, Novartis Campus; Basel, Switzerland.

[3]Bone and Stem Cell Research Group, CABMM, University of Zurich; Zurich, Switzerland.

*Corresponding author. Email: ram@ethz.ch

Abstract: Bone's ability to adapt to mechanical demands is governed by mechanoregulation, the process by which cells sense and respond to mechanical stimuli to maintain skeletal integrity. In osteoporosis, increased bone resorption activity leads to structural deterioration and elevated fracture risk. While existing pharmacological therapies aim to restore bone mass to reduce fracture risk, it is unclear how they modulate mechanoregulation, especially when combined with physical interventions. Here, we investigate the joint effects of load-bearing physical and pharmacological treatment in a female mouse model of osteoporosis using longitudinal *in vivo* micro-computed tomography and computational mechanics. We demonstrate that mechanical loading additively and synergistically enhanced predicted strength, bone volume, and mechanoregulation parameters when combined with anabolic therapies (parathyroid hormone and sclerostin antibody) but not with anti-catabolic treatments (bisphosphonates). Increases in predicted strength are associated with reductions in bone resorption rates, shifts in the (re)modeling thresholds as anticipated by Frost in the mechanostat theory, and the modeling capacity of anabolic pharmacological treatments. These findings underscore the therapeutic potential of combining anabolic pharmacological therapies with load-bearing physical activity, particularly in early treatment phases, to optimize bone adaptation and fracture prevention in osteoporosis management.

Introduction

Postmenopausal osteoporosis in women is characterized by increased bone resorption, leading to structural deterioration of the bone micro-architecture and increased susceptibility to fractures[1]. Exercise is fundamental to bone health, but its role alongside pharmacological treatment remains an area of active inquiry[2]. Patients with diagnosed osteoporosis often question the compatibility of exercise with pharmacological treatment, some fearing interference with drug efficacy, others assuming that their potent medications obviate the need for physical activity. This highlights the need for evidence on whether mechanical loading improves outcomes for bone strength beyond pharmacological monotherapy.

Mechanical loads are transmitted from the organ to bone cells, where osteocytes act as mechanosensors, osteoblasts form, and osteoclasts resorb bone. This interplay drives both remodeling (formation followed by resorption) and modeling (formation without prior resorption), the latter potentially via lining cell redifferentiation in the absence of sclerostin[3]. Key signaling pathways between bone cells include Wnt/β-catenin[4] (osteocyte-osteoblast) and RANKL/OPG[5] (osteoblast-osteoclast). To study these dynamic processes *in vivo*, high-resolution imaging techniques such as micro-computed tomography (micro-CT) are commonly used. While micro-CT does not resolve individual cells or molecular signals, it captures the net microstructural changes resulting from formation and resorption, which are collectively referred to as bone (re)modeling[6].

On the cellular level, anti-resorptive bisphosphonates (BIS) prevent osteoclast attachment and activity by binding to hydroxyapatite crystals in the bone matrix, and are characterized by low osteoanabolic capacity[7]. We hypothesized that mechanical loading would stimulate bone formation, but bisphosphonates might blunt this response by inhibiting (re)modeling.

Intermittent parathyroid hormone (PTH) exerts anabolic effects by binding to PTH-receptors on osteoblasts and osteocytes, leading to suppression of sclerostin expression and subsequent activation of the Wnt/β-catenin signaling pathway. This promotes osteoblastic activity and indirectly enhances osteoclast recruitment via upregulation of RANKL expression by osteoblasts[8]. PTH and mechanical loading have shown synergistic effects in cortical bone[9, 10]. We hypothesized that in trabecular bone, their combination would suppress otherwise increased resorption, and concurrently stimulate bone formation[11].

Sclerostin antibodies (SclAB) inhibit a Wnt/β-catenin-signaling antagonist protein, promoting osteoblast activity and suppressing osteoclast activity[12]. Wnt/β-catenin-signaling is initiated when a Wnt ligand binds to a Frizzled family receptor and its co-receptor LRP5/6 on the surface of a cell, resulting in β-catenin stabilization and target gene transcription inside the cell[11]. We hypothesized mechanical loading in combination with SclAB would increase formation and decrease resorption further. Synergy may arise because mechanical loading downregulates sclerostin expression in osteocytes[13], while SclAB pharmacologically neutralizes it, amplifying the same pathway from two directions, especially in trabecular bone which particularly adapts its microarchitecture to mechanical demands. It should be noted that, similar to mechanical loading, sclerostin antibody effects saturate after a certain time, which limits their long-term use as a standalone therapy[14].

Frost's mechanostat theory[15, 16, 17] proposes that bone adapts to maintain mechanical strain within a defined setpoint range - much like a thermostat. The setpoints separate the four following states: the disuse state for bone resorption; the adapted state for quiescence, the overload state for formation, and the fracture state for microdamage and fractures. The mechanostat theory was introduced more than 30 years ago. In this theoretical construct, anti-catabolic bisphosphonates help inhibit increased bone resorption in underloaded areas by raising the mechanical setpoints towards higher strains. Anabolic treatments theoretically

promote bone formation by lowering the mechanical setpoints required to initiate formation, provided mechanical loading is also encouraged. Measuring the "setpoints" of osteoporosis medications in the context of Frost's mechanostat theory, however, is not straightforward, as the setpoints themselves are theoretical constructs that describe the mechanical thresholds at which bones adjust their structure, and measuring them experimentally would require complex assessments of both bone volume and mechanical strain under highly controlled conditions.

Here we provide experimental insights into this interaction using *in vivo* micro-computed tomography (micro-CT) and finite-element modeling in a C57BL/6 mouse model of osteoporosis. The aim of this study was to explore whether combining pharmacological treatments with mechanical loading leads to improved pharmacological treatment outcomes, i.e. stronger bones. We tested short-term effects of BIS, PTH, and SclAB treatment with or without mechanical loading of the 6th caudal vertebra, aiming to maximize bone accrual in early treatment phases. The choice of selected therapies was based on clinical relevance and distinct modes of action in the bone (re)modeling process.

Results

Micro-architectural and (re)modeling differences reveal treatment-specific effects on bone recovery after ovariectomy

As shown in Fig. 1a, C57BL/6 mice were ovariectomized at 15 weeks-of-age, and bone loss was confirmed in all animals (n=105) via *in vivo* micro-CT of the 6th caudal vertebra (-26.8 ± 0.95% change in bone volume fraction (BV/TV) from baseline). From week 20 to 24, animals received pharmacological and/or mechanical treatment through an established tail vertebra loading model[18].

In the VEH-group, BV/TV continued to decline from week 20, while all monotreatment groups showed significant increases after four weeks of treatment (p<0.001, Fig. 1b). Trabecular thickness (Tb.Th) did not increase significantly in BIS compared to VEH, whereas all anabolic treatments led to increases, but lower in PTH than in ML and SclAB (p<0.01, Fig. 1c). Trabecular number (Tb.N) was preserved in BIS but was significantly lower in all other groups compared to BIS (p<0.001, Fig. 1d). Connectivity density (Conn.D) was preserved in BIS and PTH and declined in VEH, ML, and SclAB (p<0.05, Fig. 1e). A full summary of these microstructural parameters is provided in Supplementary Table S.2. As previously described[18], mechanical loading increases BV/TV mainly through trabecular thickening despite a reduction in trabecular number, explainable by a full loss of thin trabeculae - a pattern also seen in the SclAB-group (Fig. 1c,d). In contrast, PTH increased BV/TV through moderate thickening and sustained preservation of trabecular number, resulting in a smaller loss of connectivity. BIS maintained BV/TV by preserving existing trabeculae, including thinner ones, with minimal thickening (Fig. 1, b-e).

The micro-architectural changes resulted from the time course of formation and resorption (Fig. 1f). In week interval w20-22 (see also Supplementary Table S.5 for results from the second interval, w22–24), bone formation rate per bone volume (BFR) was significantly increased in ML and PTH compared to VEH, with no significant increases in BIS or SclAB (Fig. 1g). Mineral apposition rate (MAR) did not differ between groups (Fig. 1h), while mineralizing surface per bone surface (MS) was elevated in all treatment groups compared to VEH (p<0.05, Fig. 1i). Bone resorption rate per bone volume (BRR) was significantly reduced in all treatment groups versus VEH (Fig. 1j), where BRR in PTH was the least reduced (significantly higher than VEH, ML, and BIS), consistent with the known anabolic effects of intermittent PTH treatment. Surprisingly, BRR was found to be significantly lower than VEH, while not as low as in ML and BIS. This decrease, even more pronounced in the second time interval (see

Supplementary material Table S.5), may result from micro-CT missing parts of the reversal phase between resorption and formation, thus potentially underrepresenting remodeling at the same sites. As expected, mineral resorption rate (MRR) was significantly lower in the BIS-group compared to VEH and unchanged in the anabolic groups (Fig. 1k), suggesting bisphosphonates preserve Tb.N (Fig. 1d) by preventing full trabecular resorption. Eroded surface per bone surface (ES), reflecting the proportion of bone surface previously affected by resorption, was significantly lower in ML and SclAB compared to VEH, BIS, and PTH ($p < 0.05$, Fig. 1l).

Together, these results confirm that each monotherapy exerted effects consistent with its known mode of action - bisphosphonates preserving trabecular structure by suppressing bone resorption, and anabolic agents increasing bone mass predominantly through formation - while also highlighting treatment-specific differences in trabecular architecture and dynamic turnover.

Mechanical loading modulates structural and functional responses to pharmacological interventions

We next assessed how additional mechanical loading influenced the response to pharmacological treatment. In the BIS + ML-group, BV/TV gains were significantly less than the expected additive effects (treatment:loading: $p < 0.05$, Fig. 2a), and did not differ significantly from BIS alone (Supplementary Table S.2). In contrast, anabolic treatment combined with mechanical loading (PTH + ML, SclAB + ML) resulted in additive increases, as supported by a non-significant treatment:loading interaction (Fig. 2a). ΔTb.Th was significantly higher in all combination groups compared to monotreatments after four weeks (Fig. 2b). ΔTb.N was preserved in BIS + ML, despite the known Tb.N loss with ML alone (Fig. 2c), suggesting BIS blunted this effect. The loss of Tb.N observed in SclAB was not rescued in

SclAB + ML (Fig. 2c). Conn.D as a measure of the structural integrity and interconnectedness

of the trabecular network, was preserved the most in BIS + ML, and PTH + ML (Fig. 2d). It

should be noted that structural parameters such as Tb.Th and Tb.N are interdependent; apparent

increases in Tb.Th may arise from the loss of complete thinner trabeculae. Synergy was found

in total area (Tt.Ar), leading to additive increases in cortical area fraction (Ct.Ar/Tt.Ar) and

cortical thickness (Ct.Th), found in Supplementary Fig. S.1 and Table S.4. Predicted strength

increased in all treatment groups, with loading-induced increases being additive in the PTH +

ML group and synergistic in the SclAB + ML group ($p < 0.05$, Fig. 2e). Strain energy density

(SED) decreased in all treatment groups, being most pronounced in the combination groups,

with less than the expected additive reduction in BIS + ML (treatment:loading: $p < 0.05$, Fig. 2f,

Table 1). Illustrating the effect of decreasing SED, the visualization of the 6^{th} caudal vertebra

in a SclAB + ML animal shows a highly loaded trabecula at week 20, followed by structural

reorganization and reduced SED at week 24 (see Fig. 2g). Linear regression analysis revealed

that the changes in predicted strength did not correlate with baseline BV/TV ($R = 0.010$) and

moderately with Tb.Th ($R = 0.424$, Fig. 2h), indicating that initial BV/TV or Tb.Th serve as

poor or only moderate predictors of future changes in strength.

Combination treatments affect bone formation and resorption dynamics differentially

Bone formation rate per bone volume (BFR) was lowest in VEH and highest in the anabolic

combination groups ($p < 0.05$, Fig. 3a). MAR showed synergy in SclAB + ML

(treatment:loading: $p < 0.05$, Fig. 3b). MS increased additively in PTH + ML and SclAB + ML,

and antagonistically in BIS + ML (treatment:loading: $p < 0.01$, Fig. 3c). Bone resorption rate per

bone volume (BRR) was reduced in all treatment groups ($p < 0.05$, Fig. 3d). In BIS + ML, the

BRR reduction was antagonistically affected by mechanical loading (treatment:loading,

$p < 0.05$). In contrast, BRR decreased synergistically in PTH+ML (treatment:loading: $p < 0.05$,

Fig. 3d) and additively in SclAB + ML (Fig. 3d). MRR remained suppressed in BIS + ML and was unaffected by mechanical loading (Fig. 3e). ES decreased synergistically in PTH + ML treatment:loading: $p<0.05$, Fig. 3f) and SclAB + ML (treatment:loading: $p<0.01$, Fig. 3f). This synergistic effect was, at least in SclAB + ML, still present in the second time interval (see Supplementary Fig. S.2 and Table S.5 for results from w22–24). Representative trabecular microstructures and insets of the combination groups at both time intervals are depicted in Fig. 3g. Changes in predicted strength correlated linearly with BFR (R = 0.830) and decayed according to a power-law relationship with BRR ($\rho = 0.760$), indicating that small reductions in bone resorption rates under treatment can result in large increases in predicted strength (Fig. 3h).

Bone formation and resorption occur in mechanically distinct strain windows

To address the question whether pharmacological treatments target bone (re)modeling to the places that mechanically need it, we correlated the underlying mechanical signals in the trabecular compartment with subsequent formation and resorption events. Visually, spatial agreement was found between high SED and regions of subsequent bone formation, and low SED and regions of subsequent bone resorption (Fig. 4a). Quantitative analysis (Fig. 4b) showed that mean SED was significantly higher at formation sites ($p<0.05$) and significantly lower at resorption sites ($p<0.05$) compared to quiescent surfaces, when presented as percentage change to the average quiescent SED across all animals per group. The addition of mechanical loading to the monotreatments resulted in marginal changes, indicating that bone formation and resorption continue to occur in mechanically distinct strain windows despite possible influences of pharmacological interventions. Levene's test revealed significantly smaller variances between BIS and BIS + ML and between VEH and ML ($p<0.05$), indicative of more consistent mechanosensitive responses in these groups in the presence of mechanical loading.

Conditional probabilities of formation, quiescence, and resorption events for a given SED-value were calculated and fitted with exponential functions. The probability for bone formation increased with increasing SED, and the probability for bone resorption decreased with increasing SED (Fig. 4c, Supplementary Fig. S.3 and Table S.6 for results in w22-24). Especially the PTH + ML and SclAB + ML conditional probability curves opened up compared to their monotreatments, indicating a higher responsiveness to mechanical signals (Fig. 4c). The asymptotic resorption probability value min_R represents the point at which further increases in local mechanical loading stimulus do not further suppress bone resorption. This value was significantly lower in ML, PTH + ML, and in SclAB + ML than in VEH ($p<0.05$, Fig. 4d, Table S.8). Further comparisons of the coefficients from fitted functions to the conditional probabilities are provided in Supplementary Tables S.7 and S.8. The presence of few significant differences between groups suggests mechanosensitivity of the system is largely preserved regardless of the presence or absence of mechanical loading. Combined anabolic treatments resulted in higher correct classification rate (CCR) values compared to VEH and the BIS groups (Fig. 4e), with CCR being a metric used to evaluate the accuracy of classifying local bone formation or resorption events based on predicted mechanical signals. However, CCR only correlated moderately with the increases in strength, indicating increased mechanosensitivity of the system may not fully explain the ability of bone to increase strength (Fig. 4f).

Frost's mechanostat setpoints are shifted towards lower values for mono- and combined treatments

To assess potential shifts in the setpoints due to different drug or combination treatments, (re)modeling velocity (RmV) curves and mechanostat parameters[19] were calculated for each treatment group as an approximation to the mechanostat theory. These curves show the net RmV for a given surface effective strain, where a negative value means net resorption and a

positive value means net formation (Fig. 5a). From these curves, formation saturation levels

(FSL), resorption saturation levels (RSL), and (re)modeling thresholds (RmTs) were derived.

FSL was increased in all treatments compared to VEH ($p<0.001$, Fig. 5b, Supplementary Table

S.9), and was significantly higher in anabolic combination groups compared to the respective

monotreatments ($p<0.001$). This suggests an increased mechanosensitivity of the cells involved

in bone formation. RSL was increased in the BIS and all combination groups compared to VEH

($p<0.05$, Fig. 5c). The increase in RSL under bisphosphonate treatment may be explained by

the suppression of bone turnover: at comparable strain levels, resorption would typically occur,

resulting in a lower RSL; however, due to inhibited osteoclast activity, this resorptive response

is blunted, making the system appear saturated at a higher RSL value. Regarding the role of

additional mechanical loading, only the PTH + ML group showed a significant increase

compared to PTH alone ($p<0.05$, Supplementary Table S.9). This suggests that, under strain

conditions that would typically lead to bone resorption, the balance has shifted towards bone

formation. RmT values, representing the setpoints in the mechanostat theory, were decreased

in all pharmacological treatment groups compared to VEH ($p<0.05$), with no significant

improvements observed in the combination groups receiving mechanical loading (Fig. 5d).

These results were even more pronounced in the second time interval (Supplementary Fig. 4d,

Table S.9). The reduction in RmT suggests an upregulation of osteocytic mechanosensitivity,

an expected outcome in the anabolic treatment groups, but an unexpected finding in the

bisphosphonate-treated groups. This may reflect a compensatory osteocytic response to the

suppressed bone resorption induced by bisphosphonate treatment. Furthermore, FSL correlated

strongly with predicted strength increases ($R = 0.888$), while RmT exhibited a moderate inverse

correlation ($\rho = 0.609$) when fitted with a power-law function. These findings suggest that a

reduction in RmT may be a prerequisite for strength gains; however, actual increases in strength

require anabolic potential (Fig. 5e,f).

The amount of modeling correlates highly with increases in predicted strength for all treatments

Next, we distinguished between the remodeling and the modeling ability of the different treatments. Visually, in several areas in VEH, high SED values decreased over time, but in other areas they increased and the resorption of a single trabecula led to a more heterogeneous SED distribution (Fig. 6a). While SED on average decreased over time for BIS + ML, this did not particularly affect areas with high SED, as there remained several highly strained areas in week 24. In contrast, both PTH + ML and SclAB + ML showed large decreases in SED, especially in areas with high SED values in week 20 (Fig. 6a).

The remodeling ability - as represented by the amount of resorption followed by formation - was in BIS, BIS + ML, PTH and SclAB comparable to VEH, and significantly lower in ML and the anabolic combination groups compared to VEH ($p<0.05$, Fig. 6b). In contrast, the modeling ability was significantly higher compared to VEH in all but the BIS group, with the highest amount in the anabolic combination treatments ($p<0.001$, Fig. 6c). Both remodeling and modeling capacity showed antagonistic effects in BIS + ML and additive effects in anabolic combination treatments (Fig. 6b,c). Remodeling showed a moderate inverse correlation with increases in predicted strength when fitted with a power-law function ($\rho = 0.476$, Fig. 6d), while modeling correlated strongly in a linear fit ($R = 0.928$, Fig. 6e). Fig. 6f illustrates the distribution of surface events relative to increasing effective strain. Remodeling events appeared strain-independent, and modeling increased with higher mechanical stimuli, particularly in the anabolic combination groups (Fig. 6f). There is also a non-linear increase in modeling with mechanical loading in comparison to the mono-treatment, with a stronger effect for low effective strain regions, especially for SclAB. Fig. 6g summarizes the trabecular bone response across treatments. Bisphosphonates, alone or with mechanical loading, help preserve trabecular

interconnectedness. Anabolic treatments combined with mechanical loading effectively

enhance predicted strength by promoting formation and suppressing resorption. These additive

effects across parameters likely underlie the observed synergistic gains in predicted strength, at

least in SclAB.

Discussion

The main goal of osteoporosis therapy is to restore bone mass to reduce fracture risk. To achieve

this efficiently, existing bone must be preserved and new bone formed at mechanically relevant

sites. In healthy bone, bone (re)modeling is proposed to be regulated via the mechanostat[6, 19, 20,

21]. Here, we investigated how three current pharmacological treatments influence this regulation

in the initial treatment phases in osteoporotic mice, particularly when combined with

mechanical loading. Our findings suggest that all three investigated treatments induce bone

formation and inhibit bone resorption at the mechanically relevant sites, but it was only in

anabolic therapies that mechanical loading additively (PTH) or synergistically (SclAB)

enhanced predicted bone strength.

When combined with sclerostin antibody treatment, mechanical loading led to a synergistic

increase in predicted bone strength and an additive increase in bone volume fraction. Mineral

apposition rates and reductions in eroded surfaces were synergistic, consistent with previous

findings[12, 22, 23]. This response may be explained by dual suppression of the activity of sclerostin,

a protein secreted by osteocytes that antagonizes Wnt signaling[11]. While mechanical loading

independently suppresses sclerostin expression in osteocytes[13], SclAB pharmacologically

neutralizes any residual sclerostin[11]. Together, these effects enhance Wnt/β-catenin activity,

and likely contribute to the pronounced anabolic response[24]. Additional pathways may also be

involved, as suggested by studies in SOST-KO mice, which still showed loading-induced bone

formation[25]. One candidate is Dickkopf-1 (Dkk1), another Wnt inhibitor[26, 27, 28]. Robling et al.[13]

reported that mechanical loading reduced both *Sost* mRNA (-73%) and *Dkk1* (-49%) transcripts.

Since both proteins compete at LRP5/6 receptors in osteoblasts and osteocytes, their

simultaneous inhibition may underlie the observed synergy between SclAB and mechanical

loading.

We found additive increases in predicted strength and increased sensitivity of

mechanoregulation mechanisms for PTH + ML, which is in line with previous results where

PTH increased the mechanoresponse of osteoblasts only when combined with fluid flow[29].

Hagino et al.[30] hypothesized that PTH lowers the (re)modeling threshold of the (re)modeling

velocity function, which our findings support (Fig. 5). In previous animal studies, mechanical

loading combined with intermittent PTH (1-34) treatment led to synergistic effects in the

trabecular compartment of the healthy tibia, but it was only prevalent in regions that were not

remodeled, but modeled[31].

Bisphosphonates were the only treatment that preserved Tb.N but showed no additive effects

with mechanical loading, consistent with clinical studies reporting limited benefit of combining

bisphosphonates and exercise[32, 33]. Since mineralizing surface was significantly increased in the

bisphosphonate group compared to the vehicle group, our results suggest that bisphosphonates

not only affect the activity of osteoclasts but also that of osteoblasts, possibly through disruption

of the coupling process via elevated osteoclast apoptosis[34], potentially impairing the spatial

guidance of osteoblasts by the mechanical signal. The observation that the BIS and BIS+ML

groups exhibit RmT values as low as other treatment groups despite showing the least overall

remodeling response may indicate a compensatory increase in osteocytic mechanosensitivity,

potentially allowing selective resorption of mechanically less relevant areas.

This study focuses on osteoblast and osteoclast surface activities using *in vivo* micro-CT, which

allows longitudinal tracking of bone (re)modeling. Histological analysis could offer further

insight into cellular and tissue-level mechanisms to support the cell-specific interpretations

derived from the micro-CT data. However, due to the method of tissue preservation and the

limited number of available samples, it was not considered feasible in the current study. It's

worth noting though that previous work demonstrated a significant correlation between micro-

CT-derived and histomorphometry-based MS and MAR (R ≈ 0.59 – 0.78)[35, 36]. Furthermore,

Birkhold et al.[37] have demonstrated the ability of longitudinal *in vivo* micro-CT to reliably

quantify bone (re)modeling, i.e. separation into modeling and remodeling, in both trabecular

and cortical bone. While *in vivo* micro-CT does not allow direct visualization of osteocytes, our

measurements reflect the functional output of the bone cells in which osteocytes play an

important role through orchestration of bone formation and resorption in response to local

mechanical stimuli[6].

Regions with high local strain are linked to higher occurrences of microscopic tissue damage[38]

of which even small amounts can lead to large reductions in strength[39] and increase fracture

risk[40]. Reductions in strain energy density (SED) over time do not necessarily imply a decrease

in bone formation capacity but rather reflect system saturation, consistent with mechanostat

theory, where bone formation plateaus once strain thresholds are reached.

As a limitation, our developed system does not yet incorporate the level of the transcriptome[41].

Other cell types, particularly immune cells and endothelial cells, also contribute to bone

(re)modeling via inflammatory signaling and vascularization, respectively. Recent work

originating from fracture healing highlights the role of immune and vascular interactions,

particularly in the context of aging and mechanical loading[42]. While these cell types were not

analyzed here, future multimodal approaches could complement imaging-based assessments.

Given the biological differences among the therapies selected, including administration routes

and duration limitations, direct long-term comparisons are not the aim of this study. Instead,

our focus on short-term effects aimed to investigate potential synergistic improvements in bone

strength and to gain mechanistic insight into how pharmacological and mechanical stimuli

modulate mechanoregulation. The findings support mechanostat theory: treatments lower

mechanical setpoints, enabling bone formation at lower strains. However, only anabolic

treatments effectively reinforce highly loaded trabeculae and reduce peak strains, underscoring

the importance of modeling capacity for strength gains. And while our results highlight the

promise of mechanical loading as an adjunct to anabolic therapies, clinical studies will

ultimately be required to determine optimal exercise modalities, just as pharmacological dosing

requires careful optimization.

In conclusion, the current study revealed that drug treatments target trabecular bone

(re)modeling at mechano-functionally relevant places, and osteoanabolic treatments in

combination with mechanical loading additively and synergistically increased the bone

response. The additive (PTH) and synergistic (SclAB) increases in predicted strength can be

explained by reductions in resorption rates, shifts in the (re)modeling thresholds as described

by Frost in the mechanostat theory more than 30 years ago, and by the modeling ability of

anabolic treatments. Being able to link organ-level changes induced by physical or

pharmacological therapies to the mechanosensitivity of the cells provides information on how

well skeletal structures can adapt to mechanical demands under pathological or treatment

conditions, a task for which the body has optimized itself over millions of years of evolution.

Load-bearing exercise may represent a promising adjunct option for physically active

osteoporosis patients on osteoanabolic therapy.

Methods

Study design

The study design is illustrated in Fig. 1a. In total, 120 female 15-week-old C57BL/6 mice

(n=60, C57BL/6JRccHsd, RCC/Harlan; Füllinsdorf, Switzerland/ Horst, The Netherlands;

n=60, C57BL/6JRj mice (Janvier Labs, Le Genest-Saint-Isle, France) were subjected to

bilateral ovariectomy. A basal micro-CT image of the 6[th] caudal vertebra was obtained (vivaCT

40, Scanco Medical, Brüttisellen, Switzerland). Recovery lasted five weeks after surgery, as in

this period, an osteopenic state was established in the trabecular compartment of the 6[th] caudal

vertebra in a previous study[43]. Pinning of the adjacent vertebrae as preparation for the loading

procedure took place in week 17. At week 20, the animals were divided into 8 groups and the

following treatments were administered for the following four weeks: BIS (n=9), BIS + ML

(n=9), PTH (n=10), PTH + ML (n=9), SclAB (n=9), SclAB + ML (n=8), ML (n=26), VEH

(n=25). 10 mice were excluded due to swollen tails or loose pins. In the SclAB study, 4 mice

showed a periosteal reaction, possibly due to injury of the periosteum during intravenous

injection at the tail vein. One mouse in the BIS + ML group did not lose bone volume after

ovariectomy. These mice were excluded from evaluation, resulting in 105 animals included.

Micro-CT was performed in weeks 15, 20, 22 and 24. Surgeries, imaging, and loading were

performed under isoflurane anesthesia (induction 4%, maintenance 1.5-2.5%, Attane, Piramal

Healthcare, India). Vehicle control was an isotonic buffer (pH 5.3). While the data was gained

in four experiments (at different times), study and loading parameters were not different except

for the administered pharmacological treatments, thus combined in this study. The bone

responses of the VEH and the ML groups of both mouse lines (JRccHsd and 6JRj) showed no

differences in reactions ($\Delta BV/TV_{w20-24;}$ VEH: p = 0.937; ML: p = 0.206; two-way ANOVA

followed by Tukey's post-hoc test), thus ML groups from all experiments and VEH groups

from all experiments were merged, explaining the larger sample size in these groups. The

merging was conducted because separate vehicle group presentations would not have yielded

additional insights but substantially would have increased data complexity (see Supplementary

Table S.3). All animal experiments were performed with approval of the local authority

(Kantonales Veterinäramt Zürich, Zurich, Switzerland, license number 171/2008, 108/2011,

078/2012) and reported according to ARRIVE 2.0 guidelines.

Animal care and monitoring

Animals were kept in groups of 3-5 mice in filter top cages in Scantainers (Scanbur, Karslunde, Denmark) with *ad libitum* access to tap water and chow (KLIBA NAFAG 3437, Granivit AG, Kaiseraugst Switzerland), wood chips bedding and a 12h:12h light:dark cycle. Room temperature was 22 ± 2 °C and humidity was not controlled. Cages were changed weekly by the experimenters. Microbiological status of the facility (Animal Imaging Center, ETH Zurich) was not monitored, because it housed only animals during experiments. Animals were checked and scored daily for three days after OVX surgery and pinning and at least three times per week during the further experiment. Meloxicam 2 mg kg^{-1} was injected s.c. for three days after OVX and pinning.

Mechanical loading

A validated *in vivo* mouse tail loading model was used to induce trabecular bone adaptation under controlled mechanical conditions[43, 44, 45]. Stainless steel pins were implanted into adjacent vertebrae under fluoroscopic guidance three weeks prior to loading. An 8N cyclic axial load (3000 cycles, 10 Hz) was applied to the 6th caudal vertebra (CV6) using a custom loading device (caudal vertebra axial loading device, CVAD[44]) three times a week. Control animals (0N groups) underwent identical procedures, including pin placement and anesthesia but no load was applied. A force of 8 N was used in order to maximize the effect within a short timeframe. It was the maximum force that could be applied based on what the pins within the bone could bear, and represents twice the force exerted on mouse vertebrae under physiological conditions[46]. Pistoia simulations estimate a maximum failure load of 24.5 N at CV6 in osteoporotic mice at week 20 (predicted trabecular strength, see Table S.2).

Pharmacological treatment

The BIS group received 100 µg kg^{-1} zoledronate (Zometa 4 mg/5 ml; Novartis Pharma Schweiz AG, Bern) subcutaneously once in week 20. Corresponding VEH (0.9% NaCl, B. Braun Medical AG, Switzerland) was injected at the same time point. Mice in the PTH study received 40 µg kg^{-1} PTH (hPTH 1-34, Bachem AG, Switzerland) or the according VEH (0.9% NaCl) subcutaneously daily over four weeks. SclAB (100 mg kg^{-1}, Novartis Pharma AG, Basel, Switzerland) or the corresponding VEH (isotonic buffer, pH 5.3) was injected i.v. weekly into the lateral tail vein over 4 weeks.

BIS and SclAB doses were selected based on established protocols in the literature[24, 47]. The PTH dose was determined in a dose-response study comparing 10 µg kg^{-1} d^{-1} and 40 µg kg^{-1} d^{-1}, following an initial 80 µg kg^{-1} d^{-1} dose proposed in the literature[48] that led to excessive increases in bone volume fraction (BV/TV) by the end of the experiment. The 40 µg kg^{-1} d^{-1} dose was chosen to obtain BV/TV increases comparable to those observed with BIS and SclAb treatments, enabling a more balanced comparison across treatment groups (see Fig. 1b).

Bone morphometry

Micro-CT images were obtained with a vivaCT 40 (Scanco Medical AG, Brüttisellen, Switzerland) operated at 55 kVp and 146 µA. 1000 projection images were obtained with an integration time of 200 ms. Grayscale images were reconstructed using a Feldkamp algorithm, with automated beam hardening correction, resulting in an isotropic voxel size of 10.5 µm. The image of week 20 was aligned in the main loading direction. Follow up scans were registered to the previous scan. A constrained 3D Gauss filter (sigma 1.2, support 1) and a global thresholding procedure of 22% (approximately 580 mg HA cm^{-3}) were applied. Automated masks were created to extract the trabecular bone volumes (trabecular mask truncated by 10% of length at both ends to exclude epiphyses and primary spongiosa; middle 26% removed

458 because this area is almost void of trabeculae)[18]. The cortical mask included 75% of the length

459 of the inner mask[18]. Static trabecular bone morphometric parameters were calculated according

460 to Bouxsein et al.[49]. Dynamic bone morphometry was performed by registering and

461 superimposing segmented bone volume images of consecutive time points. Bone formation and

462 resorption rates were calculated[36] over two two-week periods (w20-22 and w22-24).

Finite element analysis of mechanical properties

Finite-element simulations were used to analyze mechanical properties of the vertebrae. 3D

micro-FE models were generated by converting all voxels of the micro-CT image to 8 node

hexahedral elements, with each model consisting of approximately 1.8 million elements. A

Young's modulus of 14.8 GPa and a Poisson's ratio of 0.3[44] were assigned. Each model was

solved with ParOSol[50] running on the Euler cluster operated by Scientific IT Services at ETH

Zurich. Cylindric intervertebral disks (10% of absolute height) were applied and assigned the

same Young's modulus and Poisson's ratio as bone. The top of the model was displaced by 1%

of the total vertebral length, while the bottom was fixed. The resulting reaction force was

rescaled to the value of the force applied in the experimental setup; consequently, all the FE

outcomes were rescaled accordingly (4 N for non-loaded[46] and 8 N for loaded mice). Strain

energy density (SED), defined as the increase in energy associated with the tissue deformation

per unit volume, and effective strain were used as the mechanical signal. Trabecular bone

strength was estimated for each mouse at each time point by using the Pistoia criterion[51] which

assumes that bone failure is initiated if a certain percentage of the bone tissue (2%) is strained

beyond the tissue yield effective strain (7000 $\mu\varepsilon$). This criterion has been shown applicable to

the mouse vertebra as well[52]. For the predicted bone strength estimation, the full bone was

modeled but only the effective strains in the trabecular compartment were analyzed.

Local mechanoregulation analysis

The local mechanical environment was derived from micro-FE simulations of the baseline scan. Here, SED is used as a mathematical term to describe the (re)modeling stimulus phenomenologically. To quantify the relationship between the local SED and cellular activity on the bone surface, the bone formation or resorption sites determined by rigid registration were projected onto the surface of the baseline scan, resulting in three masks representing three different clusters of formed (F), quiescent (Q) or resorbed (R) bone surfaces[6]. In short, the mean SED value was calculated in each of these three masks and for each mouse, and normalized to the mean SED of the quiescent surface as absolute SED values differ per mouse and over time due to differences in BV/TV. To establish a quantitative description of the mechanoregulatory system, the relation between increasing mechanical stimuli and consequent (re)modeling events was assessed by calculating the relative percentage of voxels being formed, quiescent and resorbed for each value of SED binned at 1% step size of the maximum SED, and assuming that each (re)modeling event has the same occurrence probability (i.e., formation, resorption and quiescent regions were virtually rescaled to have the same amount of voxels) to remove dependencies on imbalances in the (re)modeling process due to differences between animals or treatments. The conditional probabilities were fitted per animal by exponential functions. Correct classification rate (CCR) values were calculated from the mean conditional probabilities from all animals, returning a scalar value between 0.33 and 1 assessing overall mechanoregulation, with 0.33 meaning (re)modeling is independent of the mechanical signal and 1 meaning it is entirely dependent on the mechanical signal. (Re)modeling velocities (RmVs) and their mechanostat parameters were calculated according to Marques et al. for each animal[19]. Minimum and maximum RmV, represented as formation saturation level (FSL) and resorption saturation level (RSL), as well as (re)modeling thresholds (RmTs) were extracted from the resulting curves.

Statistical methods

Statistical analysis was performed using R (R, Auckland, New Zealand[53]). Data are presented as group means ± standard error unless otherwise specified. P-values lower than 0.05 are considered significant, statistical significance was defined as $*p<0.05$, $**p<0.01$, $***p<0.001$. Normality was assessed using the Shapiro-Wilk test (in R, with $\alpha = 0.05$).

Static morphometric and mechanical parameters of trabecular bone in the 6th caudal vertebra were measured at weeks 15, 20, 22, and 24, with Δ representing the percentage change between weeks 20 and 24. Two-way ANOVA followed by Tukey's post-hoc test was used to compare Δw20-24. Synergy (S) or antagonism (I) between treatment and mechanical loading was evaluated using the interaction term ("treatment:loading") in a linear model.

A synergistic effect was defined as a combined treatment response that exceeded the sum of the individual effects of treatment and mechanical loading. This implies that the two interventions may act through the same or facilitating biological pathways. Statistically, such synergy is indicated by a significant and positive interaction term between the main effect (treatment) and mechanical loading. An antagonistic effect was defined as a combined treatment response that was less than the expected additive effect of treatment and mechanical loading. Statistically, such antagonism was reflected by a significant and negative interaction term between the main effect (treatment) and mechanical loading. The animal provider from the third (SclAB) and fourth (PTH) experiment was different, therefore we adjusted the model for this.

Dynamic morphometric parameters were analyzed during weeks 20-22 and 22-24. One-way ANOVA followed by Tukey's post-hoc test was applied to absolute values in each time interval. Synergy or antagonism was assessed using the interaction term from a linear model of $\log_2$-transformed values in each time interval.

Strain Energy Density (SED) in formation (F), quiescence (Q), and resorption (R) surfaces was

reported for each group in each time interval. Differences between F, Q, and R were compared

using two-sided Wilcoxon signed-rank tests with Bonferroni correction after testing for

normality.

For conditional probabilities, exponential curves were fitted to the formation and resorption

slopes. For formation, a saturating exponential model $f(x) = y_0 + a \cdot (1 - \exp(-bx))$ was

applied to the first 40% of normalized SED data for each animal individually, with parameters

y_0, a, and b representing level of formation in the absence of mechanical stimulus (i.e. at SED

= 0), maximal formation, and sensitivity to mechanical stimulus, respectively. For resorption,

a decaying exponential model $f(x) = y_0 + a \cdot \exp(-bx)$ was used to assess the rate of

resorption. Resorption asymptote min_R represents the saturation level of resorption at high

mechanical signal, i.e. the minimum resorption probability ($x \to \infty$), indicative of the amount of

non-targeted resorption. Coefficient a represents the magnitude of the response, i.e. the

difference between the initial and the minimum resorption probability. Coefficient b is the rate

of decay, representing how quickly the probability of resorption drops as SED increases.

Synergy or antagonism was assessed using the interaction term from a linear model applied to

$\log_2$-transformed coefficients, with significance determined by two-way ANOVA followed by

Tukey's post-hoc test.

Formation and resorption saturation levels (FSL, RSL) and remodeling thresholds (RmTs) were

derived from hyperbola remodeling velocity curves, fitted with the function $f(x) = y_0 + \frac{a}{(b-x)}$,

where x represents the mechanical signal, y_0 the FSL, and a the (re)modeling velocity modulus.

The remodeling threshold (RmT) is defined as: $RmT = b + \frac{a}{y_0}$. One-way ANOVA followed by

Tukey's post-hoc test was used for statistical comparisons, with synergy or antagonism assessed

using the interaction term from a linear model applied to absolute values in each time interval,

because FSL and RSL can obtain negative values.

Modeling and remodeling amounts were calculated by monitoring spatial agreement of voxels

over the time intervals w20-22 and w22-24, following the approach presented by Birkhold *et*

al.[37]. Modeling amount was defined as formation without prior resorption in either the first or

the second interval, divided by the amount of surface voxels. Remodeling amount was defined

as resorption followed by formation, divided by the amount of surface voxels.

Linear correlations between the %-change in predicted strength and parameter outcomes were

assessed using Pearson's correlation coefficient R. For nonlinear relationships, we employed a

power-law model of the form $y = a \cdot x^b$, which was linearized via logarithmic transformation

to $\ln(y) = \ln(a) + b \cdot \ln(x)$. Parameters a and b were estimated using linear regression on

the transformed data, and the Spearman's correlation coefficient ρ between the log-transformed

observed and predicted values was used to assess the goodness of fit. Since the fitted power-

law model adequately described the nonlinear relationships, we did not perform a formal model

selection procedure to compare alternative models, such as those based on the Akaike

information criterion (AIC).

Data availability

573 The primary data supporting the findings of this study are available within the paper. Source

574 data are provided with this paper. The raw data generated during the study are available in a

575 Zenodo repository (see link below).

Code availability

578 All data and code are available in a Zenodo repository: 10.5281/zenodo.17256047.

References

1. Kanis JA, McCloskey EV, Johansson H, Oden A, Melton LJ, Khaltaev N. A reference standard for the description of osteoporosis. *Bone* **42**, 467-475 (2008).

2. Schumm AK*, et al.* Does adding exercise or physical activity to pharmacological osteoporosis therapy in patients with increased fracture risk improve bone mineral density and lower fracture risk? A systematic review and meta-analysis. *Osteoporosis Int* **34**, 1867-1880 (2023).

3. Kim SW*, et al.* Sclerostin Antibody Administration Converts Bone Lining Cells Into Active Osteoblasts. *Journal of Bone and Mineral Research* **32**, 892-901 (2017).

4. Duan P, Bonewald LF. The role of the wnt/beta-catenin signaling pathway in formation and maintenance of bone and teeth. *Int J Biochem Cell Biol* **77**, 23-29 (2016).

5. Kim JM, Lin C, Stavre Z, Greenblatt MB, Shim JH. Osteoblast-Osteoclast Communication and Bone Homeostasis. *Cells* **9**, (2020).

6. Schulte FA*, et al.* Local mechanical stimuli regulate bone formation and resorption in mice at the tissue level. *PLoS One* **8**, e62172 (2013).

7. Dell RM*, et al.* Incidence of atypical nontraumatic diaphyseal fractures of the femur. *J Bone Miner Res* **27**, 2544-2550 (2012).

8. Kulkarni NH*, et al.* Effects of parathyroid hormone on Wnt signaling pathway in bone. *J Cell Biochem* **95**, 1178-1190 (2005).

9. Sugiyama T*, et al.* Mechanical loading enhances the anabolic effects of intermittent parathyroid hormone (1-34) on trabecular and cortical bone in mice. *Bone* **43**, 238-248 (2008).

10. Rooney AM, McNeill TJ, Ross FP, Bostrom MPG, van der Meulen MCH. PTH Treatment Increases Cortical Bone Mass More in Response to Compression than Tension in Mice. *Journal of Bone and Mineral Research* **38**, 59-69 (2023).

11. Baron R, Kneissel M. WNT signaling in bone homeostasis and disease: from human mutations to treatments. *Nat Med* **19**, 179-192 (2013).

12. Ominsky MS, Boyce RW, Li X, Ke HZ. Effects of sclerostin antibodies in animal models of osteoporosis. *Bone* **96**, 63-75 (2017).

13. Robling AG*, et al.* Mechanical stimulation of bone in vivo reduces osteocyte expression of Sost/sclerostin. *J Biol Chem* **283**, 5866-5875 (2008).

623   14.   Rauner M, Taipaleenmaki H, Tsourdi E, Winter EM. Osteoporosis Treatment with Anti-
624   Sclerostin Antibodies-Mechanisms of Action and Clinical Application. *J Clin Med* **10**,
625   (2021).

626
627   15.   Frost HM. The mechanostat: a proposed pathogenic mechanism of osteoporoses and the bone
628   mass effects of mechanical and nonmechanical agents. *Bone Miner* **2**, 73-85 (1987).

629
630   16.   Frost HM. Bone "mass" and the "mechanostat": a proposal. *Anat Rec* **219**, 1-9 (1987).

631
632   17.   Frost HM. Bone's mechanostat: a 2003 update. *Anat Rec A Discov Mol Cell Evol Biol* **275**,
633   1081-1101 (2003).

634
635   18.   Lambers FM, Schulte FA, Kuhn G, Webster DJ, Müller R. Mouse tail vertebrae adapt to
636   cyclic mechanical loading by increasing bone formation rate and decreasing bone resorption
637   rate as shown by time-lapsed in vivo imaging of dynamic bone morphometry. *Bone* **49**, 1340-
638   1350 (2011).

639
640   19.   Marques FC, Boaretti D, Walle M, Scheuren AC, Schulte FA, Müller R. Mechanostat
641   parameters estimated from time-lapsed micro-computed tomography data of mechanically
642   driven bone adaptation are logarithmically dependent on loading frequency. *Front Bioeng*
643   *Biotech* **11**, (2023).

644
645   20.   Christen P*, et al.* Bone remodelling in humans is load-driven but not lazy. *Nat Commun* **5**,
646   4855 (2014).

647
648   21.   Griesbach JK, Schulte FA, Schadli GN, Rubert M, Müller R. Mechanoregulation analysis of
649   bone formation in tissue engineered constructs requires a volumetric method using time-lapsed
650   micro-computed tomography. *Acta Biomater*, (2024).

651
652   22.   Taylor S*, et al.* Time-dependent cellular and transcriptional changes in the osteoblast lineage
653   associated with sclerostin antibody treatment in ovariectomized rats. *Bone* **84**, 148-159 (2016).

654
655   23.   Ominsky MS*, et al.* Romosozumab Improves Bone Mass and Strength While Maintaining
656   Bone Quality in Ovariectomized Cynomolgus Monkeys. *Journal of Bone and Mineral*
657   *Research* **32**, 788-801 (2017).

658
659   24.   Morse A, Schindeler A, McDonald MM, Kneissel M, Kramer I, Little DG. Sclerostin
660   Antibody Augments the Anabolic Bone Formation Response in a Mouse Model of Mechanical
661   Tibial Loading. *J Bone Miner Res* **33**, 486-498 (2018).

662
663   25.   Morse A*, et al.* Mechanical load increases in bone formation via a sclerostin-independent
664   pathway. *J Bone Miner Res* **29**, 2456-2467 (2014).

665
666   26.   Witcher PC*, et al.* Sclerostin neutralization unleashes the osteoanabolic effects of Dkk1
667   inhibition. *Jci Insight* **3**, (2018).

27.　Morse A, *et al.* KO Mice Treated with Sclerostin Antibody Have Additional Increases in Bone
Volume. *Calcified Tissue Int* **103**, 298-310 (2018).

28.　Florio M, *et al.* A bispecific antibody targeting sclerostin and DKK-1 promotes bone mass
accrual and fracture repair. *Nature Communications* **7**, (2016).

29.　Ryder KD, Duncan RL. Parathyroid hormone modulates the response of osteoblast-like cells
to mechanical stimulation. *Calcif Tissue Int* **67**, 241-246 (2000).

30.　Hagino H, Okano T, Akhter MP, Enokida M, Teshima R. Effect of parathyroid hormone on
cortical bone response to in vivo external loading of the rat tibia. *J Bone Miner Metab* **19**, 244-
250 (2001).

31.　Robinson ST, Shyu PT, Guo XE. Mechanical loading and parathyroid hormone effects and
synergism in bone vary by site and modeling/remodeling regime. *Bone* **153**, 116171 (2021).

32.　Uusi-Rasi K, *et al.* Effect of alendronate and exercise on bone and physical performance of
postmenopausal women: a randomized controlled trial. *Bone* **33**, 132-143 (2003).

33.　Kistler-Fischbacher M, Yong JS, Weeks BK, Beck BR. A Comparison of Bone-Targeted
Exercise With and Without Antiresorptive Bone Medication to Reduce Indices of Fracture
Risk in Postmenopausal Women With Low Bone Mass: The MEDEX-OP Randomized
Controlled Trial. *J Bone Miner Res* **36**, 1680-1693 (2021).

34.　Sims NA, Martin TJ. Osteoclasts Provide Coupling Signals to Osteoblast Lineage Cells
Through Multiple Mechanisms. *Annu Rev Physiol* **82**, 507-529 (2020).

35.　Birkhold AI, Razi H, Duda GN, Weinkamer R, Checa S, Willie BM. Mineralizing surface is
the main target of mechanical stimulation independent of age: 3D dynamic in vivo
morphometry. *Bone* **66**, 15-25 (2014).

36.　Schulte FA, Lambers FM, Kuhn G, Müller R. In vivo micro-computed tomography allows
direct three-dimensional quantification of both bone formation and bone resorption parameters
using time-lapsed imaging. *Bone* **48**, 433-442 (2011).

37.　Birkhold AI, Razi H, Weinkamer R, Duda GN, Checa S, Willie BM. Monitoring in vivo
(re)modeling: a computational approach using 4D microCT data to quantify bone surface
movements. *Bone* **75**, 210-221 (2015).

38.　Acevedo C, Stadelmann VA, Pioletti DP, Alliston T, Ritchie RO. Fatigue as the missing link
between bone fragility and fracture. *Nat Biomed Eng* **2**, 62-71 (2018).

39.　Hernandez CJ, Lambers FM, Widjaja J, Chapa C, Rimnac CM. Quantitative relationships
between microdamage and cancellous bone strength and stiffness. *Bone* **66**, 205-213 (2014).

40. Lambers FM, Bouman AR, Rimnac CM, Hernandez CJ. Microdamage caused by fatigue
loading in human cancellous bone: relationship to reductions in bone biomechanical
performance. *PLoS One* **8**, e83662 (2013).

41. Mathavan N*, et al.* Spatial transcriptomics in bone mechanomics: Exploring the
mechanoregulation of fracture healing in the era of spatial omics. *Sci Adv* **11**, eadp8496
(2025).

42. Duda GN, Geissler S, Checa S, Tsitsilonis S, Petersen A, Schmidt-Bleek K. The decisive early
phase of bone regeneration. *Nat Rev Rheumatol* **19**, 78-95 (2023).

43. Lambers FM, Kuhn G, Schulte FA, Koch K, Müller R. Longitudinal assessment of in vivo
bone dynamics in a mouse tail model of postmenopausal osteoporosis. *Calcif Tissue Int* **90**,
108-119 (2012).

44. Webster DJ, Morley PL, van Lenthe GH, Müller R. A novel in vivo mouse model for
mechanically stimulated bone adaptation--a combined experimental and computational
validation study. *Comput Methods Biomech Biomed Engin* **11**, 435-441 (2008).

45. Scheuren AC*, et al.* Mechano-Regulation of Trabecular Bone Adaptation Is Controlled by the
Local in vivo Environment and Logarithmically Dependent on Loading Frequency. *Front
Bioeng Biotechnol* **8**, 566346 (2020).

46. Christen P, van Rietbergen B, Lambers FM, Muller R, Ito K. Bone morphology allows
estimation of loading history in a murine model of bone adaptation. *Biomech Model
Mechanobiol* **11**, 483-492 (2012).

47. Stadelmann VA, Bonnet N, Pioletti DP. Combined effects of zoledronate and mechanical
stimulation on bone adaptation in an axially loaded mouse tibia. *Clin Biomech (Bristol, Avon)*
26, 101-105 (2011).

48. Alexander JM*, et al.* Human parathyroid hormone 1-34 reverses bone loss in ovariectomized
mice. *J Bone Miner Res* **16**, 1665-1673 (2001).

49. Bouxsein ML, Boyd SK, Christiansen BA, Guldberg RE, Jepsen KJ, Müller R. Guidelines for
Assessment of Bone Microstructure in Rodents Using Micro-Computed Tomography. *Journal
of Bone and Mineral Research* **25**, 1468-1486 (2010).

50. Flaig C, Arbenz P. A scalable memory efficient multigrid solver for micro-finite element
analyses based on CT images. *Parallel Comput* **37**, 846-854 (2011).

51. Pistoia W, Van Rietbergen B, Lochmüller EM, Lill CA, Eckstein F, Rüegsegger P. Estimation
of distal radius failure load with micro-finite element analysis models based on three-
dimensional peripheral quantitative computed tomography images. *Bone* **30**, 842-848 (2002).

52.   Nyman JS, *et al.* Predicting mouse vertebra strength with micro-computed tomography-
derived finite element analysis. *Bonekey Rep* **4**, 664 (2015).

761
762 53.   Team RDC. R: A language and environment for statistical computing. . In: *R Foundation for*
763   *Statistical Computing*) (2007).

Acknowledgements

We acknowledge Kathleen Koch for assistance and performance of animal experiments of the

PTH and BIS groups, and Dr. Ulrike Kettenberger for assistance and performance of animal

experiments of the PTH group. Nirujan Pasupathy and Júlia van den Nest Molina have

supported the analyses of modeling and mechanoregulation mechanisms. This work was funded

by the European Union grant VPHOP (FP7-ICT-2008-223865; FAS, FML, CW, GAK) and the

European Research Council ERC Advanced grant (MechAGE ERC-2016-ADG-741883; FCM,

GAK).

Author contributions

775 RM, GAK, PJR designed the experiment. CW, GAK, MS, FML performed the experiments.

776 FAS, FCM, JG, CW, MS, FML, GAK performed image processing and analysis. MK provided

777 pharmacological agents. RM, GAK, PJR, CK supervised the project. FAS wrote the original

778 draft of the manuscript. All authors contributed to data interpretation, were involved in

779 discussions throughout the study, and participated in manuscript revision and preparation.

Competing interests

782 MK, PJR, and CK are employees of Novartis. The other authors declare they have no competing

783 interests.

Figure legends

Figure 1: Osteoporosis monotreatments enhance trabecular bone micro-architecture by different mechanisms of action.

(a) Experimental setup. 15-week-old mice are ovariectomized, and treatment is administered between week 20 (w20) and 24

(w24). (b-e, g-l) Monotreatments (VEH: n=25, BIS: n=9, PTH: n=10, SclAB: n=9; each data point represents one animal

(biological replicate)), boxplots show percentage change. The box spans the interquartile range (IQR) from the 25th to the

75th percentile. The median is shown as a line inside the box. Whiskers extend to the most extreme data points within 1.5 ×

IQR, points beyond this range are plotted as outliers. # indicates that the delta of the group is different from zero, as tested

with t-test with null-hypothesis. (b) bone volume fraction ΔBV/TV, (c) trabecular thickness ΔTb.Th, (d) trabecular number

ΔTb.N (e) connectivity density ΔConn.D. (f) Representative micro-computed tomography overlays at intervals w20-22 and

w22-24, with formation and resorption volumes (g-l) (Re)modeling rates calculated from time interval w20-22 (g) bone

formation rate per bone volume BFR (h) mineral apposition rate MAR (i) mineralizing surface per bone surface MS (j) bone

resorption rate per bone volume BRR (k) mineral resorption rate MRR (l) eroded surface per bone surface ES. Significant

differences are depicted with a symbol in the color of the corresponding group (o: to monotreatment, ●: to combined

treatment, see Supplementary Table S.1 for color encoding, p<0.05). Source data are provided as a Source Data file.

Figure 2: Mechanical loading increases the morphological effects of anabolic and anti-catabolic treatments, contributing to increased predicted strength. (a-f) Changes from baseline. Data are presented as mean ± standard error. Each data point represents one animal (biological replicate; VEH: n=25, ML: n=26, BIS: n=9, BIS +ML: n=9, PTH: n=10, PTH +ML: n=9, SclAB: n=9, SclAB +ML: n=8) (a) bone volume fraction (ΔBV/TV) (b) trabecular thickness (ΔTb.Th) (c) trabecular number (ΔTb.N) (d) connectivity density (ΔConn.D) (e) predicted strength (ΔStrength), and (f) strain energy density (ΔSED). Bars show group means of sum of single vs. combined treatment indicating: antagonistic (I), additive (A) or synergistic (S) effects determined via treatment:loading interaction in a linear model on the %-change. (g) Example of SED distribution in the 6th caudal vertebra of a SclAB + ML animal at week 20, showing reduced SED after structural reorganization from week 20 to 24. (h) Linear regression of ΔStrength with BV/TV and Tb.Th at week 20. Source data are provided as a Source Data file.

Figure 3: Mechanical loading increases bone formation and decreases resorption rates of anabolic and anti-catabolic

pharmacological treatments which contribute to increased predicted strength. (a) bone formation rate per bone volume (BFR)

(b) mineral apposition rate (MAR) (c) mineralizing surface per bone surface (MS) (d) bone resorption rate per bone volume

(BRR) (e) mineral resorption rate (MRR) (f) eroded surface per bone surface (ES). (a-f) Each data point represents one animal

(biological replicate; VEH: n=25, ML: n=26, BIS: n=9, BIS +ML: n=9, PTH: n=10, PTH +ML: n=9, SclAB: n=9, SclAB

+ML: n=8). The box spans the interquartile range (IQR) from the 25th to the 75th percentile. The median is shown as a line

inside the box. Whiskers extend to the most extreme data points within 1.5 × IQR, points beyond this range are plotted as

outliers. Δ denotes difference from VEH as no baseline value for dynamic bone morphometry rates was available. Bars show

group means of sum of single vs. combined treatment indicating: antagonistic (I), additive (A) or synergistic (S) effects

determined via treatment:loading interaction in a linear model on the log$_2$-transformed values. Significances were calculated

using one-way ANOVA followed by Tukey's post-hoc test; significant differences are depicted with a symbol in the color of the

corresponding group (o: to monotreatment, ●: to combined treatment, see Supplementary Table S.1 for color encoding,

p<0.05). (g) Visual representations of combination groups BIS + ML, PTH + ML, and SclAB + ML in week intervals w20-22

*and w22-24 (**h**) Linear regression of ΔStrength with BFR, and exponential regression analysis with BRR in week interval w20-*

22. Source data are provided as a Source Data file.

***Figure 4: Formation aligns with high mechanical signal, resorption with low**: (**a**) Spatial comparison of local mechanical*

*signal (SED at week 20) with sites of trabecular bone formation and resorption (w20-22). (**b**) SED at formation, resorption,*

and quiescent sites[6], normalized to the group mean of quiescence to allow statistical testing with this group. Each data point

represents one animal (biological replicate; VEH: n=25, ML: n=26, BIS: n=9, BIS +ML: n=9, PTH: n=10, PTH +ML: n=9,

SclAB: n=9, SclAB +ML: n=8). The box spans the interquartile range (IQR) from the 25th to the 75th percentile. The median

is shown as a line inside the box. Whiskers extend to the most extreme data points within 1.5 × IQR, points beyond this range

are plotted as outliers. *p<0.05, ***<0.001 using two-sided paired Wilcoxon-signed rank test with Bonferroni correction;

*exact p-values are found in Supplementary Table S.6. (**c**) The plots show the exponential fitting functions (taking all animals*

per group into account) for conditional probability of formation and resorption as a function of SED/SEDmax. Dashed lines

represent monotreatment (0N, including VEH), and solid lines represent combined treatment with mechanical loading (8N,

including ML). VEH is indicated by black dotted line. Values are cropped above 40% due to low voxel count[6]. *(**d-e**) Each*

data point represents one animal (biological replicate; VEH: n=25, ML: n=26, BIS: n=9, BIS +ML: n=9, PTH: n=10, PTH

+ML: n=9, SclAB: n=9, SclAB +ML: n=8). The box spans the interquartile range (IQR) from the 25th to the 75th percentile.

The median is shown as a line inside the box. Whiskers extend to the most extreme data points within 1.5 × IQR, points

beyond this range are plotted as outliers. [-] denotes unitless, Δ denotes change from VEH; sum of single vs. combined

*treatment group means indicating antagonistic (I), additive (A) or synergistic (S) effects. (**d**) Asymptotic resorption*

*probability value (minR). (**e**) CCR as a measure of mechanoregulated (re)modeling, with the ΔCCR bar for the BIS + ML*

*group close to zero, making it barely visible. (**f**) Linear regression analysis in week interval w20-22 of ΔStrength with CCR.*

***Figure 5: (Re)modeling velocity curves as an approximation of the mechanostat theory. (a)** (re)modeling velocity (RmV)*

curve fits (taking all animals per group into account) for VEH / ML, BIS / BIS + ML, PTH / PTH + ML, SclAB / SclAB + ML;

***(b)** formation saturation level (FSL) **(c)** resorption saturation level (RSL), **(d)** (re)modeling threshold (RmT) **(b-d)** Each data*

point represents one animal (biological replicate; VEH: n=25, ML: n=26, BIS: n=9, BIS +ML: n=9, PTH: n=10, PTH +ML:

n=9, SclAB: n=9, SclAB +ML: n=8). The box spans the interquartile range (IQR) from the 25th to the 75th percentile. The

median is shown as a line inside the box. Whiskers extend to the most extreme data points within 1.5 × IQR, points beyond this

range are plotted as outliers. Δ denotes difference from VEH as no baseline value from week 20 was available. Bars show

group means of sum of single vs. combined treatment indicating: antagonistic (I), additive (A) or synergistic (S) effects

determined via treatment:loading interaction in a linear model on the absolute values. Significances were calculated using

one-way ANOVA followed by Tukey's post-hoc test; significant differences are depicted with a symbol in the color of the

corresponding group (o: to monotreatment, ●: to combined treatment, see Supplementary Table S.1 for color encoding,

*p<0.05). **(e)** Linear regression of ΔStrength with FSL, and **(f)** power-law regression with RmT from week interval w20-22.*

Source data are provided as a Source Data file.

Figure 6: Modeling capacity correlates highly with the increase in predicted strength. (a) Heterogeneity of the strain distribution decreases to different extents in different treatments. Qualitative comparison of changing SED-distribution for VEH, BIS + ML, PTH + ML and SclAB + ML over the course of 4 weeks of treatment. (b) Remodeling as calculated from formation following resorption at the same location over 3 measurement time points (w20, w22, w24). (c) Modeling as calculated from formation following quiescence at the same location over 3 measurement time points (w20, w22, w24) (b-c) Each data point represents one animal (biological replicate; VEH: n=25, ML: n=26, BIS: n=9, BIS +ML: n=9, PTH: n=10, PTH + ML: n=9, SclAB: n=9, SclAB + ML: n=8). The box spans the interquartile range (IQR) from the 25th to the 75th percentile. The median is shown as a line inside the box. Whiskers extend to the most extreme data points within 1.5 × IQR, points beyond this range are plotted as outliers. Δ denotes change from VEH; sum of single vs. combined treatment group means indicating graphically antagonistic (I), additive (A) or synergistic (S) effects. Significant differences are depicted with a symbol in the color of the corresponding group (o: to monotreatment, ●: to combined treatment, see Supplementary Table S.1 for color encoding, p<0.05). (d) Remodeling amount in % correlated to increases in predicted strength fitted using a power-law function. (e) Modeling amount in % correlated to increases in predicted strength fitted using linear regression. (f) Surface

876　*event proportion (modeling or remodeling) in relation to the mechanical signal. **(g)** Spider plots summarizing the mean of*

877　*outcome parameters of the analyzed treatments. Values were standardized to facilitate visual comparisons between groups.*

878　*Source data are provided as a Source Data file.*