1 *In vivo* sequential mutagenesis in germinal center B cells using a dual-
2 recombinase approach: FOXO1 re-expression upon FOXO1 knockout rescues
3 class switch recombination

Carlota Farré Díaz[1,2,3,4,#], Eleni Kabrani[1,5,#], Wiebke Winkler[1,2,3,4], Claudia Salomon[1], F.
Thomas Wunderlich[6,7,8], Martin Janz[2,3,4] and Klaus Rajewsky[1]

[1]Immune Regulation and Cancer, Max Delbrück Center for Molecular Medicine in the
Helmholtz Association, Berlin 13125, Germany.
[2]Biology of Malignant Lymphomas, Max Delbrück Center for Molecular Medicine in the
Helmholtz Association, Berlin 13125, Germany.
[3]Experimental and Clinical Research Center, a cooperation between the Max Delbrück
Center for Molecular Medicine in the Helmholtz Association and the Charité –
Universitätsmedizin Berlin, Berlin 13125, Germany.
[4]Hematology, Oncology and Cancer Immunology, Charité – Universitätsmedizin Berlin,
Berlin 13125, Germany.
[5]Genome Diversification & Integrity, Max Delbrück Center for Molecular Medicine in the
Helmholtz Association, Berlin 13125, Germany.
[6]Faculty of Medicine and University Hospital Cologne, Center for Molecular Medicine
Cologne (CMMC), University of Cologne, Cologne 50931, Germany
[7]Cologne Excellence Cluster for Stress Responses in Ageing-Associated Diseases
(CECAD), University of Cologne, Cologne 50931, Germany.
[8]Max-Planck-Institute for Metabolism Research, Cologne 50931, Germany
[#] These authors contributed equally

Corresponding author: Klaus Rajewsky, Max-Delbrück-Center, Robert-Rössle-
Straße 10, 13125 Berlin-Buch, klaus.rajewsky@mdc-berlin.de

Keywords: germinal center, sequential mutagenesis, dual-recombinase, Cre, Dre

Abbreviations: **4-OHT:** 4-hydroxytamoxifen, **BFP:** blue fluorescent protein, **Cγ1:**
immunoglobulin heavy constant gamma 1, **CSR:** class switch recombination, **DZ:** dark
zone, **GC:** germinal center, **GCBC**: GC B cell; **KO:** knockout, **LZ:** light zone, **MBC**:
memory B cell, **NP-CGG**: 4-hydroxy-3-nitrophenylacetyl-conjugated chicken gamma
globulin, **PC**: plasma cell, **R26:** Rosa 26, **SHM:** somatic hypermutation, **TAM:** tamoxifen.

Abstract

Modeling complex (patho)physiological processes by sequential mutagenesis in mice is limited by the lack of optimized genetic tools and complex breeding strategies. We present a new Cre/DreERT2 dual-recombinase germinal center B-cell (GCBC)-specific strain, with co-expression of the recombinases from a single allele. This enables highly efficient Cre-mediated FOXO1 knockout followed by time-controlled, efficient Dre-mediated FOXO1 re-expression and functional rescue in GCBCs, demonstrating suitability for precise targeted sequential mutagenesis *in vivo*.

Site-specific recombination systems, such as Cre/loxP, are key technologies to study
gene function and (patho)physiological processes *in vivo*. Dual-recombinase
approaches represent a further technical development that enables more precise
disease modelling, in particular with respect to multistep genetic manipulations and
lineage tracing[1,2]. However, one limitation of the currently available systems is that
recombinases are expressed from different loci, hereby requiring the generation of
complex compound mutant mice – containing separate transgenes encoding the
recombinases together with the respective target alleles – and thus complicating the
analysis both in terms of breeding time frame and fidelity of sequential mutagenesis at
the level of individual cells. Here, we describe a dual-recombinase mouse strain in the
context of germinal center (GC) B cells (GCBCs), where Cre and the tamoxifen (TAM)-
dependent DreERT2 recombinases are co-expressed from the immunoglobulin heavy
constant gamma 1 (*Ighg1/Cγ1*) locus and thus at the initiation of the GC response. By
sequentially targeting the endogenous *Foxo1* locus during the GCBC reaction as a proof-
of-concept experiment, we demonstrate highly efficient Cre-mediated FOXO1 knockout
(KO) with perturbation of GC physiology, followed by time-controlled Dre-mediated
FOXO1 re-expression and phenotypic rescue, demonstrating that this system can be
used *in vivo* as a genetic recombination switch that is precisely controlled with respect
to cellular compartment and time.

Upon antigen encounter, mature B cells undergo clonal expansion in distinct histological
structures called GCs. During the early phases of GC formation, activated B cells
diversify their immunoglobulin repertoire via class switch recombination (CSR), and later,
in the course of the GC reaction, via somatic hypermutation (SHM) of antibody V region
genes[3]. Positively selected GCBCs differentiate into memory B cells (MBCs) or antibody-
secreting plasma cells (PCs), thus shaping the humoral immune response[3]. Both CSR
and SHM require the introduction and the efficient repair of DNA strand breaks in the
immunoglobulin loci[4]. Errors during these processes can lead to oncogenic mutations
and/or translocations, rendering (post)-GCBCs the origin of most B cell malignancies[5].
In order to study the physiology and malignant transformation of GCBCs, a number of
Cre mouse lines have been generated[6,7,8,9,10]. However, none of the currently available
lines allows for the genetic manipulation of GCBC differentiation in a stepwise fashion.
This technical limitation prevents not only the *in vivo* investigation of the role and
cooperation of genes at different time points during the GC reaction, but also the study
of the sequential acquisition of oncogenic events in this microenvironment, underlying
malignant transformation.

Addressing this experimental challenge, we have generated a novel dual-recombinase
GCBC-specific mouse strain, in which Cre and the TAM-inducible DreERT2
recombinases are concomitantly expressed from the *Ighg1/Cγ1* locus, hereafter called
the *Cγ1-CDE* strain (Fig. 1A). *Cγ1* encodes the constant region of IgG1. Upon T cell-
dependent immunization, most activated B cells that enter the GC produce germline γ1
transcripts in response to IL4[6], allowing the expression of both recombinases – and thus
Cre activity – from the *Cγ1* locus early on. Cells that have completed switch
recombination to IgG1 continue to express the recombinases during the course of the
GC reaction. Since DreERT2 activation is dependent on TAM administration, the second
(Dre-mediated) recombination event can be induced in a time-controlled manner,
selectively in cells having undergone Cre induction. To functionally validate the newly
generated strain, *Cγ1-CDE* mice were crossed to the fluorescent reporter strains *Rosa*
26 (R26)-BFP[stopF][11] and *R26-ZsGreen*[stopRox][12] for detection of Cre- and Dre-mediated
recombination, respectively. *Cγ1-CDE*, *R26-BFP*[stopF]*, R26-ZsGreen*[stopRox] compound
mutant mice were immunized with NP-CGG and TAM was administered at days 2-5, 9-
12 or 15-18 after immunization with subsequent flow cytometric analysis at days 7, 14 or
21, respectively (Fig. 1B-G). Consistent with the previous characterization of the *Cγ1-*
Cre allele[6], up to 96% of splenic GCBCs were labelled through Cre-mediated
recombination (BFP[+] GCBCs, Fig. 1B-D, Supplemental Fig.1A-B). Further gating on
these BFP[+] GCBCs showed a robust and highly efficient Dre-mediated recombination at
all time points analyzed (Fig. 1B-D), with the highest dual labelling efficiency on day 7 of
analysis (up to 60% ZsGreen[+] cells within the BFP[+] GCBC population, Fig. 1B). GCs are
organized into two phenotypically and functionally distinct compartments, the dark zone
(DZ) and the light zone (LZ)[3]. GCBCs in both compartments were successfully labeled
(Fig. 1E-G). Importantly, in the absence of TAM, no Dre recombination was detected,
demonstrating that Dre-mediated recombination is tightly controlled *in vivo* (Fig. 1E-G).

Germline γ1 transcripts are detected in the activated B cell pool early on upon
immunization, and their presence is independent of the subsequent IgG1 CSR event *per*
se[13,6]. Accordingly, Cre-mediated followed by Dre-mediated sequential mutagenesis is
achieved not only in IgG1[+] (up to 70% ZsGreen[+] within BFP[+] cells) but also in IgM[+] (up
to 42% ZsGreen[+] within BFP[+] cells) GCBCs in our mouse model at all three time points
analyzed (Supplemental Fig. 1E-F).

In addition, analysis of the post-GC compartments revealed the feasibility of labeling
these cell types as well. MBCs exhibited solid Cre-mediated and Dre-mediated
sequential mutagenesis (up to 84% BFP[+] cells and 60% ZsGreen[+] within BFP[+] cells, Fig.
1B-D, Supplemental Fig. 1C). Analysis of the splenic plasmablast/PC compartment (Fig.

1B-D, Supplemental Fig. 1C) exhibited the highest labelling efficiency for this
compartment at day 7 of analysis (up to 75% BFP⁺ cells and 50% ZsGreen⁺ within BFP⁺
cells, Fig. 1B), indicating the feasibility to target and thus study extrafollicular PCs with
our *Cγ1-CDE* strain. However, the fraction of labelled PCs is reduced when analyzing
the immunized mice at later time points (day 14 and day 21, Fig. 1C-D), possibly due to
the short lifespan of extrafollicular PCs.

The FOXO1 transcription factor, although dispensable for GC
development/maintenance, positively regulates both GC compartmentalization (DZ
compartment) and CSR[14,15]. To this end, and as a proof-of-concept experiment, we
aimed at genetically manipulating the endogenous *Foxo1* locus in a sequential manner.
In addition to the already published Cre-inducible *Foxo1* null allele (*Foxo1^{fl}*)[16], we
generated a rox-controlled, Dre-inducible *Foxo1* allele (*Foxo1^{stopRox}*). The *Foxo1^{stopRox}*
allele carries a STOP cassette flanked by rox sites in front of the endogenous *Foxo1*
open reading frame, leading to the inactivation of the allele in the absence of Dre activity
(Supplemental Fig. 2A). Of note, while hemizygous *Foxo1* mice are viable and fertile,
homozygous germline deletion of FOXO1 results in embryonic lethality[17,18]. In order to
verify that the *Foxo1^{stopRox}* allele is functional upon deletion of the STOP cassette, splenic
B cells from *Cγ1-CDE*, *Foxo1$^{stopRox/wt}$* compound mutant mice were activated *in vitro* and
cultured in the presence or absence of 4-hydroxytamoxifen (4-OHT), followed by
sequencing analysis of *Foxo1* transcripts (Fig. 2A). Indeed, expression of the targeted
allele was observed solely in the presence of 4-OHT, demonstrating not only that the
Foxo1^{stopRox} allele is functional, but also the tightness of the system. *In vivo*, crossing of
Cγ1-CDE, *Foxo1^{stopRox}* mice with *Foxo1^{fl}* transgenic mice (*Cγ1-CDE*, *Foxo1$^{fl/stopRox}$*) would
generate a scenario in which upon NP-CGG immunization, and therefore Cre activation,
the *Foxo1^{fl}* allele would be deleted, resulting in a FOXO1-KO condition in GCBCs (Fig.
2B). GCBC-specific FOXO1 deletion is expected to lead to loss of the DZ compartment
and to defects in CSR, while maintaining the GCBC phenotype[11,12]. In a second step
upon TAM treatment – and hence Dre nuclear translocation and activation –, re-
expression of FOXO1 would be achieved from the *Foxo1^{stopRox}* allele, thus leading to a
rescue of the previously induced FOXO1 deficiency (Fig. 2B). To address whether the
sequential deletion and re-activation of FOXO1 from the endogenous locus could restore
GC compartmentalization and CSR, experimental *Cγ1-CDE*, *Foxo1$^{fl/stopRox}$, R26-*
ZsGreen^{stopRox} (in short *Foxo1$^{fl/stopRox}$*) and control *Cγ1-CDE*, *Foxo1$^{fl/wt}$, R26-*
ZsGreen^{stopRox} (in short *Foxo1$^{fl/wt}$*) mice were immunized with NP-CGG, subsequently
treated with TAM (day 10-12 after immunization) and analyzed on day 14. The *R26-*
ZsGreen^{stopRox} allele was included as an indirect reporter to identify Dre-recombined

cells, i.e., FOXO1 re-expression in *Foxo1*^{stopRox} mice (Fig. 2C). Strikingly, TAM

administration with subsequent FOXO1 re-activation in the *Foxo1*$^{fl/stopRox}$ group

completely restored the DZ compartment in ZsGreen$^+$ GCBCs, up to the same levels as

in *Foxo1*$^{fl/wt}$ control mice (Fig. 2D). This observation also suggests a high efficiency of

Dre-mediated recombination in this setting, since all rox sites available (here the rox

sequences in *Foxo1*^{stopRox} and *R26-ZsGreen*^{stopRox} alleles) are targeted in the same cell.

In contrast, ZsGreen$^-$ GCBCs in the experimental *Foxo1*$^{fl/stopRox}$ group, which had failed

to recombine the *Foxo1*^{stopRox} allele by Dre and had remained a full FOXO1-KO, lost the

physiological levels of the DZ/LZ ratio, at the expense of DZ cells (Fig. 2D) and in

agreement with previous studies[14,15]. Additionally, CSR to IgG1 was robustly restored in

ZsGreen$^+$ GCBCs in the *Foxo1*$^{fl/stopRox}$ group, albeit at a lower frequency compared to

controls (Fig. 2E). We attribute this lower level to the fact that reporter ZsGreen$^+$ cells

from the *Foxo1*$^{fl/stopRox}$ mice have resided in the GC only for a maximum of 4 days (after

TAM administration), in contrast to the control cells, which have participated in the GC

reaction from the beginning. This considerable rescue of class-switching in GCBCs at

the peak of the GC reaction suggests that CSR can robustly be induced during the GC

response, irrespective of its frequent occurrence at the pre-GCBC stage[13].

In summary, our newly established *Cγ1-CDE* allele allows for highly efficient Cre and

subsequent Dre recombination in the context of GCBCs. Thus, this tool can be used *in*

vivo not only to model the sequential acquisition of genetic alterations acquired during

malignant transformation but also to study the interplay of two or more genetic events or

the function of a specific gene at different time points during the GC reaction. A similar

strategy of sequential targeted mutagenesis could be applied in other cellular contexts.

1.   Li, H., Weng, W. & Zhou, B. Perfect duet: Dual recombinases improve genetic resolution.
Cell Prolif. 56, 1–12 (2023).

2.   Sket, T., Falcomatà, C. & Saur, D. Dual Recombinase–Based Mouse Models Help
Decipher Cancer Biology and Targets for Therapy. *Cancer Res.* 83, (2023).

3.   Victora, G. D. & Nussenzweig, M. C. Germinal Centers. *Annu. Rev. Immunol.* 40, (2022).

4.   Chi, X., Li, Y. & Qiu, X. V(D)J recombination, somatic hypermutation and class switch
recombination of immunoglobulins: mechanism and regulation. *Immunology* vol. 160,
(2020).

5.   Küppers, R., Klein, U., Hansmann, M. L. & Rajewsky, K. Cellular origin of human B-cell
lymphomas. *N. Engl. J. Med.* (1999).

6.   Casola, S. *et al.* Tracking germinal center B cells expressing germ-line immunoglobulin γ1
transcripts by conditional gene targeting. *Proc. Natl. Acad. Sci. U. S. A.* (2006).

7.   Weber, T. *et al.* A novel allele for inducible Cre expression in germinal center B cells. *Eur.*
J. Immunol. (2019).

8. Dogan, I. *et al.* Multiple layers of B cell memory with different effector functions. *Nat.*
Immunol. 10, (2009).

9. Shinnakasu, R. *et al.* Regulated selection of germinal-center cells into the memory B cell
compartment. *Nat. Immunol.* (2016).

10. Callahan, D. *et al.* Memory B cell subsets have divergent developmental origins that are
coupled to distinct imprinted epigenetic states. *Nat. Immunol.* (2024).

11. Sommermann, T. *et al.* Functional interplay of Epstein-Barr virus oncoproteins in a mouse
model of B cell lymphomagenesis. *Proc. Natl. Acad. Sci. U. S. A.* 117, (2020).

12. Biglari, N. *et al.* Functionally distinct POMC-expressing neuron subpopulations in
hypothalamus revealed by intersectional targeting. *Nat. Neurosci.* 24, (2021).

13. Roco, J. A. *et al.* Class-Switch Recombination Occurs Infrequently in Germinal Centers.
Immunity (2019).

14. Dominguez-Sola, D. *et al.* The FOXO1 Transcription Factor Instructs the Germinal Center
Dark Zone Program. *Immunity* 43, (2015).

15. Sander, S. *et al.* PI3 Kinase and FOXO1 Transcription Factor Activity Differentially Control
B Cells in the Germinal Center Light and Dark Zones. *Immunity* 43, (2015).

16. Paik, J. H. *et al.* FoxOs Are Lineage-Restricted Redundant Tumor Suppressors and
Regulate Endothelial Cell Homeostasis. *Cell* 128, (2007).

17. Hosaka, T. *et al.* Disruption of forkhead transcription factor (FOXO) family members in
mice reveals their functional diversification. *Proc. Natl. Acad. Sci. U. S. A.* 101, (2004).

18. Furuyama, T. *et al.* Abnormal angiogenesis in Foxo1 (Fkhr)-deficient mice. *J. Biol. Chem.*
279, (2004).

Conflict of interest: The authors declare no conflict of interest.

Acknowledgements: We thank all members of the K. Rajewsky lab for discussion, R.
Kühn (MDC transgenic facility), H.P. Rahn (MDC FACS facility), G. Natale (K. Rajewsky
lab, MDC) for excellent technical support and the MDC animal caretakers for their
outstanding technical help. Figure schematics were created using images from
BioRender. This work was supported by the European Research Council (ERC
Advanced Grant 268921) to K.R., the Deutsche Krebshilfe (grant #70112800) to K.R.
and M.J., and the Berlin School of Integrative Oncology (BSIO) to C.F.D. We apologize
to colleagues whose work could not be cited owing to space limitation.

Author contributions: C.F.D., E.K. and W.W. designed the experiments; C.F.D., E.K.,
W.W. and C.S. performed experiments; T.W. provided the *R26-ZsGreen*^{stopRox} strain;
C.F.D. and E.K. analyzed data and wrote the manuscript; M.J and K.R. supervised all
aspects of the study, and reviewed and edited the manuscript.

FIGURE 1

Figure 1. Characterization of the *Cγ1-CDE* strain. (A) Targeting strategy for the insertion of the IRES-Cre_T2A_DreERT2 cassette into the 3'-UTR of the mouse *Cγ1* locus. (B-G) Experimental scheme using the *Cγ1-CDE, R26-BFP^{stopF}, R26-ZsGreen^{stopRox}* compound mutant mice, immunizing with NP-CGG at day 0, administrating TAM at day 2-5 (B and E), 9-12 (C and F) or 15-18 (D and G) and analyzing at day 7, 14 or 21, respectively. (B-D) Percentage of Cre-mediated BFP$^+$ (left panel) and Dre-recombined ZsGreen$^+$ (within the BFP$^+$) cells (middle panel) among splenic Fas$^+$CD38$^-$ GCBCs, splenic Fas$^-$CD38$^+$IgD$^-$IgG1$^+$ MBCs and CD138$^+$TACI$^+$ PCs from spleen and BM (from one femur and tibia); as well as corresponding absolute numbers of the BFP$^+$ZsGreen$^+$ cells (right panel). (E-G) Percentage of ZsGreen$^+$ (within BFP$^+$) CXCR4^{hi}DC86^{low} DZ and CXCR4^{low}CD86^{hi} LZ GCBCs at day 7 (E), day 14 (F) and day 21 (G) after NP-CGG immunization. Data points for GCBCs without TAM administration are compiled from the three time points analyzed. Statistics: Mann-Whitney test *$P \leq 0.05$; **$P \leq 0.01$; ***$P \leq 0.001$; **** $P \leq 0.0001$; ns, not significant. Each dot represents one mouse. Data from at least two independent experiments. Horizontal lines indicate the median.

FIGURE 2

Figure 2. **Sequential mutagenesis at the endogenous *Foxo1* locus.** (A) Schematic representation of *in vitro* induction of the *Foxo1^{stopRox}* allele by *Cγ1-CDE* (left) and cDNA sequencing in activated *Cγ1-CDE, Foxo1^{stopRox}* splenic B cells (right). Expression of the targeted allele is observed only in the presence of 4-OHT, as demonstrated by the detection of the synonymous PAM mutation introduced during the targeting strategy of the transgenic strain. (B) Genetic scheme of recombination events after Cre and Dre activation at the *Foxo1* locus of *Foxo1$^{fl/stopRox}$* animals. (C) Experimental scheme (top) and representative gating strategy (bottom) employed for the phenotypic analysis of the GC compartment. (D-E) Top: Representative flow cytometry plots measuring the percentage of DZ and LZ (D) and IgG1$^+$ (E) splenic GCBCs of mice of the indicated genotypes at day 14 after immunization. Bottom: Quantification of the DZ/LZ ratio (D) and the percentage of IgG1$^+$ (E) ZsGreen$^-$ (grey) and ZsGreen$^+$ (green) splenic GCBCs. Statistics: Mann–Whitney test **$P \leq 0.01$; ***$P \leq 0.001$; ****$P \leq 0.0001$; ns, not significant. Each symbol represents one mouse. Data from three independent experiments. Horizontal lines indicate the median.