Article Title: Indirect genetic effects increase the heritable variation available to selection

and are largest for behaviours: a meta-analysis

Authors: Santostefano Francesca[1,2*], Maria Moiron[3,4*], Alfredo Sánchez-Tójar[4*], David N

Fisher[5*]

* All authors contributed equally

Affiliations :

[1] University of Exeter, Centre for Ecology and Conservation, University of Exeter, Penryn

Campus, Cornwall, U.K.

[2] Université du Québec à Montréal, Département des Sciences Biologiques, Montréal,

Canada

[3] Institute of Avian Research, Wilhelmshaven, 26386, Germany

[4] Department of Evolutionary Biology, Bielefeld University, 33615, Bielefeld, Germany

[5] School of Biological Sciences, University of Aberdeen, King's College, Aberdeen, United

Kingdom

Corresponding author contact information:

Francesca Santostefano, PhD

Centre for Ecology and Conservation, University of Exeter,

Penryn Campus, Penryn, Cornwall

TR10 9FE, UK

F.Santostefano2@Exeter.ac.uk

Title: Indirect genetic effects increase the heritable variation available to selection and are largest for behaviours: a meta-analysis

Abstract

The evolutionary potential of traits is governed by the amount of heritable variation available to selection. While this is typically quantified based on genetic variation in a focal individual for its own traits (direct genetic effects, DGEs), when social interactions occur, genetic variation in interacting partners can influence a focal individual's traits (indirect genetic effects, IGEs). Theory and studies on domesticated species have suggested IGEs can greatly impact evolutionary trajectories, but whether this is true more broadly remains unclear. Here we perform a systematic review and meta-analysis to quantify the amount of trait variance explained by IGEs and the contribution of IGEs to predictions of adaptive potential. We identified 180 effect sizes from 47 studies across 21 species and found that, on average, IGEs account for a small but statistically significant amount of phenotypic variation (0.03). As IGEs affect the trait values of each interacting group member and due to a typically positive – although statistically nonsignificant – correlation with DGEs ($r_{DGE-IGE}$ = 0.26), IGEs ultimately increase trait heritability substantially from 0.27 (narrow-sense heritability) to 0.45 (total heritable variance). This 66% average increase in heritability suggests IGEs can increase the amount of genetic variation available to selection. Furthermore, whilst showing considerable variation across studies, IGEs were most prominent for behaviours, and to a lesser extent for reproduction and survival, in contrast to morphological, metabolic, physiological, and development traits. Our meta-analysis therefore shows that IGEs tend to enhance the evolutionary potential of traits, especially for those tightly related to interactions with other individuals such as behaviour and reproduction.

Lay Summary

Predicting evolutionary change is important for breeding better livestock and crops, for understanding how biodiversity arises and how populations respond to environmental change. Normally, these predictions are based on how the genetic variants in an organism influence its own traits (characteristics). However, when organisms socially interact, for instance by fighting or cooperating, then the genes in one individual can influence the traits of others, therefore affecting the potential for evolutionary change. We compared 47 studies across 21 animal species and found that the effect of the genes of others is small but statistically significant for evolutionary potential. These effects are particularly important for the evolution of animals' behaviours and reproductive traits, but less so for other traits such as body size and physiology. We also found that, because an

individual can interact with many others and influence them all, social interactions can substantially
increase the potential for a population to evolve from generation to generation. Our results show
how social interactions can potentially alter the evolution of those traits known to respond to social
interactions in comparison to standard expectations.

Introduction

Traits can evolve when they are underpinned by heritable genetic variation and subject to selection.
The (narrow-sense) heritability of a trait is defined as the ratio of the additive genetic variance to the
total phenotypic variance within a population, with a higher ratio indicating greater potential for
evolutionary change in response to selection (Falconer & Mackay, 1996; Fisher, 1930; Lynch &
Walsch, 1998). Environmental variation generally makes up the majority of the remainder of the
total phenotypic variation and is thought not to contribute to evolved changes. However, the
environment of an individual can also include its conspecifics and their traits, such as the phenotype
of the breeding partner, a neighbouring individual, or a competitor. Most organisms live at least part
of their lives socially, interacting when they mate, fight, cooperate, and compete for access to
resources (Frank, 2007; Székely et al., 2010; Westneat & Fox, 2010). Through these social
interactions, individuals can influence the phenotypic expression of traits in conspecifics. If an
individual's effect on others has a heritable component, then genes in one individual influence the
phenotype of others, known as indirect genetic effects (hereafter IGEs; Fig. 1; Dickerson, 1947;
Griffing, 1967; Moore et al., 1997; Scott & Fuller, 1965; Wolf et al., 1998). These IGEs can contribute
to phenotypic evolution alongside the additive genetic variance stemming from an organism's own
genes (direct genetic effects; hereafter DGEs; Fig. 1). However, when measuring the evolutionary
potential of traits, researchers typically only quantify DGEs. Only considering DGEs neglects the total
genetic variance in trait values stemming from other sources, such as IGEs, and can lead to
inaccurate evolutionary predictions (Wolf et al., 1998).

Although the extra genetic variance generated by IGEs can speed up evolutionary change,
IGEs can also reduce, remove, or even reverse the response to selection if DGEs and IGEs are
negatively correlated (Griffing, 1967; Moore et al., 1997). A positive DGE-IGE correlation indicates
that individuals with genes for high trait values elicit higher trait values in their partners (e.g.
aggressive individuals instigating higher aggression in others, Alemu et al., 2014; Wilson et al., 2009).
In contrast, a negative correlation indicates individuals with high trait values are associated with low
trait values in partners (e.g. individuals good at acquiring limited resources cause others to acquire
fewer resources, Wilson, 2014). IGEs have long been appreciated in livestock breeding, where they
can be exploited to increase productivity and improve welfare (see Ellen et al., 2014 for a qualitative
review), as well as in agriculture to increase yield through intercropping (Wright 1985, Bourke et al.

2021). IGEs have also been studied in the context of maternal genetic effects (the heritable effect a mother has on the traits of her offspring beyond the direct effect of the genes they inherit from her; Dickerson, 1947; McAdam et al., 2014; Mousseau & Fox, 1998), as well as implicitly within social and sexual runaway dynamics (Bailey & Kölliker, 2019). Recently, however, we have begun to appreciate that IGEs are applicable to a much broader array of traits and contexts (McAdam et al., 2014). Examples include the influence of a male bird on the lay date of his mate (Brommer & Rattiste, 2008; Evans et al., 2020), the elicitation of aggression from an opponent during a contest (Santostefano et al., 2017; Wilson et al., 2009), and the growth rate and infection status of neighbours (Costa E Silva et al., 2013). IGEs therefore are *expected* to alter the evolutionary potential of a wide range of traits, but whether they do is still unclear. In the wild, trait evolution frequently does not match expectations based on the ratio of direct genetic variance to phenotypic variance and the strength of selection observed in wild populations (Merilä et al., 2001; Pujol et al., 2018). Under artificial selection, the observed response may also not match the expectations based on calculations including only DGEs and the strength of imposed selection (Ellen et al., 2014; Falconer, 1981; Muir, 2005). Studying the role of IGEs in trait evolution more broadly is therefore essential to increase the predictability of evolutionary change and explain these apparent patterns of "evolutionary stasis", both in the wild and under artificial conditions.

The recent increase in the number of studies providing estimates of IGEs across taxa presents us a timely opportunity to quantitatively assess the role that IGEs play in trait evolution and their importance relative to DGEs. Further, by comparing standardised estimates across contexts and among different trait types, we can assess when IGEs are particularly prevalent, and when trait evolution will diverge most from that predicted by DGEs alone. For example, behavioural traits are a key element of any social interaction and are typically considered to be highly plastic. Behavioural traits may therefore show large IGE variance (Bailey et al., 2018; see Hunt et al., 2019 for a review of IGEs on behaviours). However, behavioural traits may also vary more in response to the non-social environment, in which case the proportion of variance in phenotypes due to IGEs may not be so high. IGEs could also impact traits related to competition such as growth, resource-dependent life-history traits, or fitness components that are not social traits *per se* (Fisher & McAdam, 2019; Wilson, 2014), but still depend on the phenotypes of conspecifics. Many studies on IGEs have taken place on captive populations housed at relatively high densities where individuals cannot avoid social interactions (reviewed in Ellen et al., 2014). In contrast, organisms in the wild typically live at lower densities and may have the option of avoiding social interactions when beneficial. There is therefore the untested possibility that IGEs in the wild are weaker than in captive populations, and so may not contribute as much to the trait's evolutionary potential, and ultimately, change.

We performed a systematic review and meta-analysis to quantitatively assess the overall

magnitude and generality, and several patterns of variation, in IGEs across the published literature.

We did not however consider maternal effects, as they have recently been reviewed elsewhere (see

Moore et al., 2019). Further, we only analysed effect sizes stemming from the "variance-

partitioning" approach used to estimate IGEs (see Methods for further details) since Bailey and

Desjonquères (2022) recently reviewed IGEs estimates using the "trait-based" approach. A glossary

with definitions of terms used throughout is available in Table 1.

Our aims were as follows:

1.   Quantify the relative magnitude of IGEs across studies (*social* h^2 = variance in IGEs [V_{IGE}] /

total phenotypic variance [V_P]). We had no clear prediction about its magnitude, but

based on our understanding of the literature we expected social h^2 to be relatively low.

2.   Test whether IGEs differ among (a) types of traits, (b) ages, (c) sexes, and (d) wild and

captive populations. Our predictions for (a) were that, because traits closely linked to

fitness such as life-history traits typically have low heritabilities, but high evolvabilities

(see Hansen et al., 2011; Houle, 1992; Postma 2014, for detailed metric interpretations),

the pattern for IGEs variance would be similar. Instead, we expected that behavioural

traits would be highly plastic and therefore may show larger IGEs, but also high

environmental variances, giving low heritabilities, but high evolvabilities (Bailey et al.,

2018). We also predicted for (d) higher effects in captive populations compared to wild

populations given the difficulty of avoiding interactions in captivity. We had no specific

predictions for the remaining sub-aims.

3.   Quantify the magnitude of IGEs in comparison to DGEs. Specifically, we compared: (a)

direct additive genetic variance (V_A) vs. V_{IGE}, (b) heritability (h^2) vs. *social* h^2, and (c) I_{DGE}

vs. I_{IGE} (where $I_{DGE} = V_A/\bar{x}^2$ and $I_{IGE} = V_{IGE}/\bar{x}^2$; Bailey et al., 2018). The latter relates

variances to the trait mean to assess evolvability (i.e. a trait's evolutionary potential in

response to selection, Hansen et al., 2011; Houle, 1992). IGEs are mediated by social

interactions, giving a less direct relationship with trait expression than DGEs, and thus

we expected them to be smaller than DGEs.

4.   Test whether IGEs alter evolutionary trajectories, which we evaluated in two ways: (a)

by assessing the overall size and direction of the DGE-IGE correlation ($r_{DGE-IGE}$) to

understand whether IGEs are expected to increase (positive correlations) or decrease

(negative correlations) the rate of evolutionary change, and (b) by comparing narrow-

sense heritability (h^2) with total heritable variance (τ^2). Since h^2 only considers DGEs,

while τ^2 considers DGEs, IGEs, and their covariance, larger τ^2 compared to h^2 values

162    would indicate that IGEs increase the amount of genetic variation available to selection

163    and can potentially speed up evolution, while smaller τ^2 values would indicate IGEs can

164    potentially slow down evolution.

Materials and methods
165

Systematic review
166

167 We performed our systematic literature search on the 8th of August 2019 in both *Web of Science*

168 *Core Collection* (databases: SCI-EXPANDED 1900-present, SSCI 1956-present, A&HCI 1975-present,

169 ESCI 2015-present) and *Scopus*, and consisted of three different steps. First, we searched for studies

170 on IGEs published across all years using the string: (("indirect genetic effect*") OR ("social genetic

171 effect*") OR ("associative genetic effect*") OR ("interacting phenotype*")). Second, we performed a

172 backward search to identify references cited in two key recent reviews on the topic: Ellen et al.

173 (2014) and Bailey et al. (2018). Third, we performed a forward search to identify references citing at

174 least one of three seminal theoretical papers on the topic: Bijma & Wade (2008), Moore et al.

175 (1997), and Wolf et al. (1998). We downloaded all the references and found and removed duplicates

176 using the R package 'revtools' v.0.3.0 (Westgate, 2019). We screened the titles and abstracts of 1058

177 unique references using Rayyan (Ouzzani et al., 2016), of which 199 references were selected for

178 full-text screening. Full-text screening identified 111 studies, of which 63 applied an 'animal model'

179 approach to analyse their data and were then subjected to data extraction for our meta-analysis. All

180 authors equally performed both the title-and-abstract and the full-text screening using predefined

181 decision trees detailing the inclusion and exclusion criteria (Figures S1 and S2). To reduce any

182 potential observer biases, the title and abstract of 27% of the references (N = 284), and the full text

183 of 22% of the references (N = 44) were screened by more than one observer. Conflicting decisions

184 (title-and-abstract: 7% [N = 20]; full-text: 16% [N = 7]) were collectively discussed and resolved. For

185 more detailed information see our PRISMA diagram (Figure S3; Moher et al., 2009) and the provided

186 data (https://github.com/ASanchez-Tojar/meta-analysis_IGEs).

Data extraction
188

189 Data were extracted from the text and tables in the main text and supplementary materials of the

190 studies identified by our search. All authors equally performed data extraction and the data

191 extraction of all the papers was double-checked by another author. The original and second

192 observer discussed and resolved unclear decisions to ensure reliability. To ensure this process was

193 uniform, all authors extracted data from four selected papers as a pilot at the beginning of this step.

194   From each included study, we extracted information about the article (year of publication,

195 journal, author information), study subjects (species, population and years studied, population type

[captive, semi-captive {defined as a population kept in captivity for no more than five generations},
wild], sex [males, females, both] and age [juveniles, adults, both] of included subjects) and type of
trait(s) studied (behaviour, development, metabolism and physiology, morphology, reproduction,
survival). In addition, we extracted the information needed to interpret the statistical estimates of
each animal model, namely whether response variables were mean- or variance-standardized, the
error distribution and any transformations used, mean group size used to estimate IGEs, mean and
standard deviation (SD) of the trait(s) studied (i.e. the response variables), and whether fixed effects
pertaining to focal or partner individual traits were included in the models. Lastly, we extracted
sample sizes (e.g. number of individuals studied, number of observations), pedigree-related
information, and the estimates from the animal models (e.g. direct [V_A] and indirect genetic effect
variances [V_{IGE}], narrow-sense [h^2] and total heritability [τ^2], direct-indirect genetic correlation [$r_{DGE\text{-}}$
$_{IGE}$], and direct-indirect genetic covariance [$CoV_{A\text{-}IGE}$]).

After data extraction, we excluded 16 studies: (i) 8 studies (k = 13 effect sizes) that did not
contain usable data (i.e. because of being a conference abstract, reporting the same data as other
studies, not estimating IGEs using the animal model approach, or solely reporting estimates from
existing literature), (ii) 1 study (k = 1 effect size) on humans, (iii) 2 studies (k = 7 effect sizes) on
Eucalyptus globulus because we did not find data for any other plant species, and (iv) 5 studies (k =
10 effect sizes) that exclusively analysed non-Gaussian traits. Notably, several wild studies fell into
this latter exclusion category, reducing the sample size for that category (see below). Requests for
missing or partially reported data were sent to the corresponding authors of 38 studies using a
standardized email template (provided in Supplementary Materials), from which we obtained
additional data for 25 studies. Finally, note that our meta-analytic dataset included 13 out of the 21
studies included in the qualitative literature review by Ellen et al. (2014; see Table 4 therein), with
the difference being explained by the inclusion/exclusion criteria explained above. For a complete
list   of   studies   and   extracted   variables   see   Table   2   and   the   provided   data
(https://github.com/ASanchez-Tojar/meta-analysis_IGEs).

Effect size calculation
Our effect sizes of interest were: (i) relative magnitude of DGEs (narrow-sense heritability or h^2) and
IGEs (*social h^2*; aims 1, 3 and 4), (ii) unstandardized variances associated with DGEs (V_A) and IGEs
(V_{IGE}), which we log-transformed prior to the analyses to better match model assumptions (aim 3),
(iii) DGE and IGE evolvability (I_{DGE} and I_{IGE}; aim 3), (iv) DGE-IGE correlation ($r_{DGE\text{-}IGE}$; aim 4), which
when not reported, was calculated as $COV_{A\text{-}IGE}$ divided by the square root of the product of V_A and
V_{IGE}, and (v) total heritable variance, τ^2 (i.e. total proportion of variance explained by DGEs, IGEs, and

their covariance, aim 4), which was calculated as $[V_A + Cov_{A\text{-}IGE}*2*(n\text{-}1) + V_{IGE}*(n\text{-}1)^2] / V_P$ (Bijma,

2010; Bijma et al., 2007). As in previous meta-analyses (e.g. Dochtermann et al. 2019, Holtmann et

al. 2017), for most of our analyses we treated intra-class correlations as equivalent to Pearson's

correlation coefficients (r) and applied Fisher's Z transformation to obtain our main effect size of

interest (i.e. Zr and its associated sampling variance, VZr). Therefore, those meta-analytic estimates

correspond to Zr instead of r throughout; however, we present back-transformed values (i.e. to r) in

the figures to aid biological interpretation. The variance in Zr (i.e. sampling variance) was calculated

as: $VZr = 1 / (n\text{-}3)$, where n represents the number of unique individuals phenotyped. The different

levels of replication analyzed (e.g., number of groups, number of replicates per group) in the original

mixed-effects models from which we obtained the estimates may lead to an underestimation of

sampling variance. To test the robustness of our results to an increase in sampling variance, we

performed sensitivity analyses by rerunning several analyses after having artificially increased

sampling variance by a factor of 10 (i.e., $VZr_{new} = VZr_{original} * 10$). This increase did not substantially

change our conclusions (Supplementary Material). If social h^2 was not directly reported in the

corresponding study, we calculated it ourselves as V_{IGE} / V_P. If V_P was not reported, we calculated it

ourselves as V_A / h^2 when these variables were reported, and then calculated social h^2. These

calculations allowed us to add 13 additional effect sizes to our dataset. Additionally, there were nine

estimates of V_{IGE} reported as "0" in the primary literature. To avoid removing those values (as zeros

cannot be log-transformed), for the comparison of V_A and V_{IGE} only (aim 3) we converted the

reported V_{IGE} values to the minimum possible rounded value based on how the "0" was reported

(e.g. reported 0.00 transformed to 0.001). We note that small variances in our dataset might be

upwardly biased due to the likely prevalence of estimates from REML models and/or the likely

preference for reporting means and medians rather than modes in Bayesian models (see Pick et al.

2023).

Meta-analyses and meta-regressions

We first ran an intercept-only meta-analytic model to estimate the overall effect size (i.e. meta-

analytic mean) for social h^2. We then ran a meta-regression for each of our hypotheses, giving eight

meta-regressions, each containing a single moderator at a time. All meta-regressions were

complete-case meta-regressions (i.e. effect sizes with unreported moderator data were discarded).

In addition, we also ran an intercept-only meta-analysis to estimate the overall effect size for $r_{DGE\text{-}IGE}$.

All models included the following six random effects: (i) phylogeny, which consisted of a

phylogenetic relatedness correlation matrix (Cinar et al., 2022), (ii) species ID, encompassing effect

sizes obtained from the same species and accounts for among-species differences not explained by

phylogeny, (iii) population ID, encompassing effect sizes obtained from the same population, (iv)

study ID, encompassing effect sizes extracted from the same study, (v) group ID, encompassing

effect sizes derived from the same group of animals and allowing us to model different experimental

groups within the same study, and (vi) record ID (unit-level effect), which represents residual and/or

within-study variance in meta-analytic models. The phylogeny was built by searching for the species

in the Open Tree Taxonomy (Rees & Cranston, 2017) and retrieving their phylogenetic relationships

from the Open Tree of Life (Hinchliff et al., 2015) using the R package 'rotl' v.3.0.11 (Michonneau et

al., 2016). We estimated branch lengths following (Grafen, 1989) as implemented in the function

'compute.brlen()' of the R package 'ape' v.5.4-1 (Paradis & Schliep, 2019) and included the

phylogenetic variance–covariance matrix as a random effect in all models. For the meta-regressions

used to compare the (relative) magnitude of DGEs and IGEs as well as to compare narrow-sense

heritability (h^2) to total heritable variance (τ^2; hypotheses 3a, 3b, 3c and 4b), since each pair of

estimates (e.g. V_{IGE} and V_A) was obtained from the same model, we included an additional random

effect (vii; model ID) that encompassed effect sizes obtained from the same animal model to

account for that extra level of nonindependence.

279   To evaluate the importance of each potential source of variation, for all meta-analytic

280   (intercept-only) models, we calculated four heterogeneity metrics (σ^2, I^2, CV, M) to understand total

281   and random effect-specific heterogeneities following the pluralistic approach suggested by Yang et

282   al. (2023). For all meta-regressions, we estimated the percentage of heterogeneity explained by the

283   moderator as $R^2_{marginal}$ (Nakagawa & Schielzeth, 2013).

284   We used the programming language R v.4.0.4 (R Core Team, 2021) throughout. All meta-

analytic and meta-regression models were fitted in the R package 'metafor' v.2.5-82 (Viechtbauer,

2010). Results were graphically represented as orchard plots using the R package 'orchaRd' v.2.0

(Nakagawa et al., 2023).

Testing for sources of bias, including publication bias

We confirmed the robustness of our results by running several sensitivity analyses that tested: a) the

effect of including fixed effects pertaining to social partners in the original animal models, b)

whether conclusions remained when only using originally reported estimates (i.e. excluding our own

calculations of some effect sizes; see above), c) evidence for publication bias both small-study and

decline effects (following Nakagawa et al., 2022; Sánchez-Tójar et al., 2018) and d) the effect of

potentially underestimating sampling variance. None of these factors proved to be significant

sources of bias, see the Supplementary Materials for further details on our approach and the results.

Results

Our full dataset contained 180 effect sizes from 47 studies published between 1997 and 2019 and involving 21 species. The median and mean number of studies per species was 1 and 2.2, respectively, with *Sus scrofa* (N = 15 studies) and *Gallus gallus* (N = 8 studies) being the most commonly studied species. Note that since we performed complete-case analyses throughout, sample sizes differ slightly across models and research questions.

Aim 1: What is the relative magnitude of IGEs (social h^2)?

The intercept-only model revealed a small, but statistically significant estimate for social h^2 (Zr = 0.031, 95% confidence intervals, CI, [0.012, 0.051], 95% prediction intervals, PI, [-0.061, 0.124], p = 0.002; k = 146 effect sizes, N = 40 studies, 19 species; Figure 2), but heterogeneity was high (I^2_{total} = 93.2%; CV_{total} = 1.46; M_{total} = 0.70). I^2_{total} and CV_{total} indicated that heterogeneity in our dataset is, on average, around 14 times larger than statistical noise and around 1.5 times larger than the meta-analytic mean. Random effect-specific I^2 estimates showed that 62.9% of heterogeneity was attributed to study ID, 20.1% to species ID not linked to the phylogenetic tree, and 10.2% to record ID. Negligible or zero relative heterogeneity was attributable to the phylogenetic tree, group ID, and population ID.

Aim 2: Does the magnitude of IGEs vary with intrinsic or extrinsic factors?

2a: Trait category

Social h^2 differed among trait categories (Figure 3a) with the moderator explaining 32.1% of the heterogeneity ($R^2_{marginal}$ = 0.321). The only trait categories with a mean estimate statistically significantly different from zero were behaviour (Zr = 0.061, 95% CI [0.032, 0.091]) and reproduction (Zr = 0.053, 95% CI [0.014, 0.093]; Figure 3a). Behavioural traits had statistically significantly higher social h^2 values than all other trait categories except for reproduction and survival (see Table S1 for p-values from all pair-wise tests). The post-hoc Wald tests showed that most of the differences between the remaining trait categories were not statistically significant, with only reproduction either being significantly higher than both development and morphology, or showing a trend in that direction when compared to metabolism and physiology (Table S1).

2b: Age

Social h^2 differed among age categories and was statistically significantly larger for effect sizes combining both age classes (Zr = 0.091, 95% CI [0.022, 0.160]) compared to those from juveniles-

only (Zr = 0.013, 95% CI [-0.029, 0.055], p = 0.049) and adults-only (Zr = 0.028, 95% CI [-0.05, 0.062],

p = 0.026; Table S1, Figure 3b). Estimates for juveniles-only and adults-only were not statistically

significantly different from each other (p = 0.403; Table S1, Figure 3b). Overall, the moderator age

explained 6.8% of the heterogeneity ($R^2_{marginal}$ = 0.068). Note that there were only four effect sizes

for the category combining both age classes, and therefore caution should be applied when drawing

conclusions from those differences.

2c: Sex

Social h^2 differed among sex categories, with social h^2 for females-only effect sizes being larger (Zr =

0.063, 95% CI [0.041, 0.084]) than when including both sexes (Zr = 0.006, 95% CI [-0.011, 0.023], p <

0.001) or for only males (though not statistically significantly so; Zr = 0.038, 95% CI [-0.036, 0.112], p

= 0.524; Table S1, Figure 3c). Sex as a moderator explained a large amount of heterogeneity (31.1%;

$R^2_{marginal}$ = 0.311); however, there were only seven effect sizes for male-only data, and, out of the 40

female-only effect sizes, 28 were from chickens in a production setting, which limits our ability to

generalise these results.

2d: Population type

Social h^2 did not differ among population types (Table S1, Figure 3d) with the moderator explaining

1.5% of the heterogeneity ($R^2_{marginal}$ = 0.015). Social h^2 was statistically significantly different from

zero in captive populations (Zr = 0.030, 95% CI [0.006, 0.053]), however, the estimate did not

statistically differ from zero in semicaptive populations (Zr = 0.020, 95% CI [−0.037, 0.076]) and only

marginally so for wild populations (Zr = 0.048, 95% CI [−0.000, 0.097]: Table S1, Figure 3d). The pair-

wise comparisons revealed no statistically significant differences between all three levels (p > 0.448

in all cases; Table S1, Figure 3d). Note, however, that there were only six studies (k = 6 effect sizes)

from wild populations and four (k = 29 effect sizes) from semicaptive populations.

Aim 3: What is the relative magnitude of IGEs compared to DGEs?

We ran three meta-regressions to test for the difference in magnitude between V_{IGE} and V_A (aim 3a);

social h^2 and h^2 (aim 3b); and I_{IGE} and I_{DGE} (aim 3c). We found that regardless of whether IGEs were

estimated as raw variance ($\log(V_{IGE})$ = -3.775, 95% CI [-6.562, -0.988] vs. $\log(V_A)$ = -0.019, 95% CI [-

2.806, 2.767], p < 0.001), variance-standardized variance (*social* h^2: Zr = 0.042, 95% CI [-0.027, 0.111]

vs. h^2: Zr = 0.263, 95% CI [0.194, 0.331], p < 0.001), or mean-standardized variance (I_{IGE} = 0.045, 95%

CI [-0.134, 0.223] vs. I_{DGE} = 0.224, 95% CI [0.045, 0.402], p = 0.003), IGEs were smaller than DGEs

(Figure 4). The moderator explained a considerable amount of variance in each meta-regression
(13.5%, 26.8%, and 4.0%, respectively).

Aim 4: Do IGEs alter evolutionary trajectories?

The intercept-only model revealed an overall positive but statistically non-significant correlation
between DGEs and IGEs ($r_{DGE-IGE}$; $Zr = 0.270$, 95% CI [-0.081, 0.621]; Figure 5b), but heterogeneity was
high ($I^2_{total} = 99.98$; $CV_{total} = 3.25$; $M_{total} = 0.85$). This high heterogeneity explains the large 95%
prediction intervals observed (95% PI [-1.497, 2.036]), which shows that, although positive on
average, $r_{DGE-IGE}$ should be expected to range from very negative to very positive depending on the
particular context investigated.

Total heritable variance (τ^2) was stastistically significantly larger ($r = 0.449$, 95% CI [0.241,
0.656]) than narrow-sense heritability (h^2; $r = 0.270$, 95% CI [0.062, 0.478], p < 0.001) and the
moderator explained 6.7% of the heterogeneity ($R^2_{marginal} = 0.067$; Figure 5a).

To better understand whether the increase in τ^2 compared to h^2 was driven by the IGEs
variance and its relationship with group size (n) or due to the seemingly positive $r_{DGE-IGE}$ detected, as
a post-hoc investigation we calculated the two components of τ^2 that stem from V_{IGE}, following Bijma
et al., 2007 (*eq. 6*) and qualitatively compared their overall magnitude. These are the "covariance
component" ($Cov_{A-IGE}*2*[n-1]$) and the "IGE variance component" ($V_{IGE}*[n-1]^2$). This additional
analysis showed that the IGEs variance component was seemingly larger (mean = 693.14) than the
covariance component (mean = 241.75), suggesting that it is the IGEs variance component that
drives the increase in τ^2.

In addition, τ^2 represents how much genetic variation is available to *all* types of selection
(including group-level selection). To test how much difference IGEs make to potential responses to
direct, individual-level only selection, we calculated the difference between the predicted response
to direct selection on unrelated individuals in the presence and absence of IGEs, following eq. 15 in
Bijma and Wade (2008), assuming that individuals interact in groups and a direct selection gradient
of 1 (i.e. 1* [V_A + (n-1) Cov_{A-IGE}]). This post-hoc analysis showed that the response to direct,
individual-level only selection among unrelated individuals was not different in the presence of IGEs
($Zr = 0.183$, 95% CI [0.048, 0.319]) and in their absence ($Zr = 0.225$, 95% CI [0.089, 0.360], p = 0.470,
$R^2_{marginal} = 0.002$).

Discussion

We performed a meta-analysis to assess the overall strength and patterns of variation in indirect genetic effects across the published literature. The magnitude of IGEs is small but statistically significant, with social h^2 accounting for, on average, around 3% of the phenotypic variation. A recent meta-analysis concluded that genetic and non-genetic maternal effects together explained 11% of the phenotypic variance (Moore et al., 2019). Maternal *genetic* effects alone (a type of IGEs excluded from our own meta-analysis) must be smaller than 11% and, therefore, likely comparable to our IGEs estimate. The high levels of heterogeneity among effects that we found show that, as expected, the importance of IGEs varies across contexts, with the 95% prediction intervals suggesting that once estimate precision is accounted for, *social h^2* is expected to range from 0 to 0.12 in most cases. Indeed, we found that factors such as trait type as well as age and sex of individuals explain substantial heterogeneity in *social h^2* estimates across studies. Notably, IGEs were more prominent for behaviours (0.06) and traits related to reproduction (0.05), supporting the hypothesis that labile traits are most strongly affected by social interactions. Furthermeore, as expected, IGEs were smaller in magnitude compared to DGEs, with estimates of direct narrow-sense heritability (h^2 = 0.27) being, on average, 6 to 8 times (depending on the statistical model) larger than *social h^2*. However, despite this large difference, accounting for IGEs and their correlations with DGEs (i.e. total heritable variance, τ^2) leads to a 66% increase in the total amount of genetic variation available to selection (τ^2 = 0.45) when compared to narrow-sense h^2. This difference is mostly driven by the relationship between IGEs and interacting group size, but also due to the overall positive correlation bewtween DGEs and IGEs ($r_{DGE-IGE}$ = 0.26; although not statistically different from zero). We detected no evidence of publication bias, and our results were robust to data imputations, potential underestimation of the sampling variance, and several aspects of model structure in the original studies (see Supplementary Materials). Overall, our results show the general importance of IGEs for animal populations, the consequences of which we outline below.

IGEs can enhance the evolutionary potential of traits

The most impactful consequence of IGEs on traits evolution to emerge from our meta-analysis is that IGEs enhance evolutionary potential by increasing the amount of genetic variation available to selection. In their review, Ellen et al., (2014) suggested that total heritable variance (τ^2) seems to be larger than narrow-sense heritability (h^2) in most domesticated populations; we here confirm this quantitatively and statistically. Two separate but not mutually exclusive mechanisms appear to facilitate this increase in adaptive potential: (1) the influence of IGEs on many interaction partners, and (2) a positive DGEs-IGEs correlation ($r_{DGE-IGE}$).

With a post-hoc analysis, we showed that the increase in τ^2 compared to h^2 is mainly driven

by the IGEs variance component ($V_{IGE}[n-1]^2$) rather than covariance component ($Cov_{A-IGE}*2*[n-1]$).

This result suggests that there may be greater evolutionary potential when organisms interact in

larger groups, as then the impact of V_{IGE} is magnified; however, V_{IGE} may also decrease, if the

influence of any two random individuals on each other is diluted as n increases (see Arango et al.,

2005; Hadfield & Wilson, 2007; Wade et al., 2010). Bijma (2010) presents a general model for the

relationship of IGEs with group size, but we could not determine where studies included in our meta-

analysis used it (but see Duijvesteijn et al. 2012 and Canario et al. 2017 for examples). The suggested

importance of group size supports results from previous modelling approaches revealing that both

DGEs and IGEs can depend largely on both group size and interaction strength (Liu & Tang, 2016;

Trubenová & Hager, 2012). How interactions occur and whether group size will enhance or dilute

IGEs will be specific to the biology of the species and the setup of the study. In most captive studies,

experimenters enforce one group size across the experiment. Therefore, caution should be exercised

when trying to extrapolate estimates of IGEs to situations where group sizes are outside the range of

those used to estimate the IGEs in the first place (see also attempts to weight interaction strengths

with, for example, social- or spatial-based neighbour networks, and determine effective group size:

Bijma, 2010; Costa E Silva et al., 2013; Fisher et al., 2019; Santostefano et al., 2021). Furthermore,

differences among individuals in their effect on group mates (e.g. "keystone" individuals: Modlmeier

et al., 2014; or among individual variation in "social impact": Araya-Ajoy et al., 2020; de Groot et al.,

2023) could also become important and modulate the magnitude of the effect of IGEs in a group.

The overall positive (though uncertain) correlation between DGEs and IGEs ($r_{DGE-IGE}$ = 0.26)

also indicates that heritable social effects tend to facilitate trait evolution rather than impede it. This

suggests that IGEs may not be a general explanation for phenotypic stasis in the face of directional

selection (Kruuk et al., 2008; Merilä et al., 2001; Pujol et al., 2018; Wilson, 2014). Our result aligns

with a review of the interacting phenotypes coefficient Ψ (Bailey & Desjonquères, 2022), which

quantifies the effect of a trait on the expression of the same or a different trait in social partners.

The sign of Ψ was generally positive, indicating positive feedback between the trait expression in

interacting partners, and suggesting a predicted response to selection that is *greater* than in traits

where Ψ is zero or without IGEs (Wolf et al., 1998). We then estimated the potential response to

selection as an additional comparison for narrow-sense h^2 and τ^2, which revealed an important

caveat. The response to individual-level direct selection in the presence of IGEs (in the absence of

relatedness and multilevel selection) was not statistically different from that predicted in their

absence. This lack of statistical difference, in contrast to the results for narrow-sense h^2 and τ^2,

appears because it is non-zero relatedness and multilevel selection that magnify the influence of

V_{IGE} on the response to selection. Relatedness between the focal individual and the individuals
affecting its fitness is crucial for harnessing the heritable variation for adaptive evolution (see Bijma
2011 for further discussion). Therefore, if relatives interact, the contribution of V_{IGE} to traits
evolution will be even greater. That IGEs do not enhance the response to selection when only
individual-level selection occurs may partially explain the surprising lack of response to artificial
selection for improvement of socially-affected traits (Craig & Muir, 1996; Ellen et al., 2014; Wade,
1976, 1977), and suggests that multilevel selection may be necessary to "unlock" the evolutionary
potential of IGEs (Bijma & Wade, 2008; Muir et al., 2013). However, multilevel selection is not
typically quantified in studies of selection in wild populations, nor always applied in breeding
designs. We therefore do not yet know the general prevalence of multilevel selection and how that
impacts the evolutionary potential of populations, and call researchers in the field to consider
quantifying its influence in future studies.

IGEs are smaller than DGEs

As expected, DGEs were larger than IGEs (e.g. narrow-sense h^2 [0.27] was more than 6 times larger
than *social* h^2 [0.03]) across metrics. Our estimates of h^2 are in accordance with other estimates of
additive genetic variance derived from animal-model based studies (e.g. 30%: Postma, 2014; 10–
30%: Wood et al., 2016; 22%: Moore et al., 2019). Overall, this result confirms our prediction that
the effect of an individual's own genes on their own phenotype should be larger than the effect of
the genotypes of interacting individuals, even for traits that are often considered highly plastic in
response to conspecifics, such as behaviours (Bailey et al., 2018; West-Eberhard, 2003).

IGEs are largest for behaviours and reproductive traits

IGEs are larger for behaviours and reproductive traits (*social* h^2 = 0.06 and 0.05, respectively),
compared to other traits categories such as morphology, development, survival, metabolism and
physiology (*social* h^2 = 0 to 0.02). This higher susceptibility may occur as behavioural and
reproductive traits likely involve feedback dynamics between interacting partners, such as mutual
escalation of aggression in contests (McGlothlin et al., 2010; Trubenová et al., 2015). Similarly,
reproductive traits such as lay date or offspring production should heavily depend on the influence
of the mating partner as the reproductive outcome is in part a joint phenotype, contributed to by
both parents. Further, behavioural traits often have relatively low estimates of h^2 (Dochtermann et
al., 2015; Moore et al., 2019; van Oers & Sinn, 2014) and so our analyses shows that traits with
higher *social* h^2 are those that often have low direct h^2. Therefore, while behaviours may show
limited response to direct selection, due to the prominence of IGEs they may respond more to social,
group-level, and other types of selection. Altogether, these findings suggest that behaviours and
reproductive traits might follow a substantially different evolutionary pathway (including indirect

effects), and that evolution for these traits is expected to diverge most from that predicted by DGEs
alone.

Other sources of heterogeneity in IGEs

The additional moderators that we tested to try to explain the high heterogeneity observed
showed varying levels of importance. Although life stage explained a non-negligible amount of
heteregenity across effect sizes, the magnitude of IGEs did not clearly differ between early and late
life stages, which could be because our meta-analysis did not include maternal genetic effects, a
type of IGEs expected to be strong for juvenile traits (Moore et al., 2019). The sex of the studied
individuals explained a high amount of heterogeneity across effect sizes, with IGEs being most
important in female only estimates rather than in those including both sexes. However, we note that
we did not have enough male-only estimates for a meaningful comparison (7 effect sizes from only 2
studies) and that 30 out of the 40 female-only estimates corresponded to captive chickens, hence
more diversity in study systems is needed to directly compare sexes and confirm this trend. Indeed,
considering sex differences in the magnitude of IGEs in future studies will be useful as the sexes
often differ in their patterns of social interaction and competition, which is associated with sexually
dimorphic traits (Fairbairn et al., 2007; Shine, 1989).

The scarcity of wild studies hampered our assessment of whether IGEs estimates differ
compared to studies on captive animals. Fortunately, logistical challenges such as capturing
individuals and monitoring their social interactions, along with large sample sizes required by
quantitative genetics models (Charmantier et al., 2014), are gradually being met in recent years. The
development of large collaborative datasets (e.g. Culina et al. 2021), refined tools for spatial and
social network analyses (e.g. Fisher et al. 2019; Radersma 2021), tracking technology (e.g. Nathan et
al. 2022), as well as the ability to estimate genetic variances from genomic rather than pedigree data
(e.g. Gienapp et al. 2017, Johnston et al. 2022), are expected to lead to more studies estimating IGEs
in the wild. These findings mirror results of a meta-analysis estimating DGEs, where there was no
support for heritabilities differing between laboratory and field conditions, nor among wild,
domestic, or semi-domestic species (Dochtermann et al., 2019), although previous studies show that
behaviours measured in the field tended to exhibit higher heritabilities (Mousseau & Roff, 1987; van
Oers & Sinn, 2014; Stirling et al., 2002).

Conclusions
Our meta-analysis across 21 animal species highlights the small but statistically significant relative
magnitude of IGEs, yet also highlights their potentially large contribution to adaptive potential. IGEs

were most prominent for behavioural and reproduction traits, possibly suggesting a different

evolutionary route for these traits, despite the considerable variation across studies. Not accounting

for IGEs underestimates the total amount of genetic variance available to selection in most contexts.

IGEs can instead enhance the evolutionary potential of traits and speed up evolutionary rates by

facilitating a response to selection, possibly promoting the adaptation of populations to rapidly

changing environmental conditions. The increase in adaptive potential was mostly due to the

relationship between IGEs and group size, but also due to the correlation between IGEs and DGEs

tending to be, on average, positive across studies. However, it may take the action of multilevel

selection to realise a population's full potential to respond to selection in the presence of IGEs, and

therefore understanding the interplay between IGEs, multilevel selection, and relatedness will be

key for accurately predicting evolutionary change. Additionally, future studies should expand our

knowledge to underrepresented taxonomic groups, particularly focusing on wild populations, which

still remain severely understudied.

References

Adams, M. J., Robinson, M. R., Mannarelli, M., & Hatchwell, B. J. (2015). Social genetic and social environment effects on parental and helper care in a cooperatively breeding bird. *Proceedings of the Royal Society B: Biological Sciences*, *282*(1810), 20150689. https://doi.org/10.1098/rspb.2015.0689

Alemu, S. W., Berg, P., Janss, L., & Bijma, P. (2016). Estimation of indirect genetic effects in group-housed mink (Neovison vison) should account for systematic interactions either due to kin or sex. *Journal of Animal Breeding and Genetics*, *133*(1), 43–50. https://doi.org/10.1111/jbg.12163

Alemu, S. W., Bijma, P., Møller, S. H., Janss, L., & Berg, P. (2014). Indirect genetic effects contribute substantially to heritable variation in aggression-related traits in group-housed mink (Neovison vison). *Genetics Selection Evolution*, *46*, 30. https://doi.org/10.1186/1297-9686-46-30

Arango, J., Misztal, I., Tsuruta, S., Culbertson, M., & Herring, W. (2005a). Estimation of variance components including competitive effects of Large White growing gilts. *Journal of Animal Science*, *83*(6). https://doi.org/10.2527/2005.8361241x

Arango, J., Misztal, I., Tsuruta, S., Culbertson, M., & Herring, W. (2005b). Threshold-linear estimation of genetic parameters for farrowing mortality, litter size, and test performance of Large White sows. *Journal of Animal Science*, *83*(3). https://doi.org/10.2527/2005.833499x

Araya-Ajoy, Y. G., Westneat, D. F., & Wright, J. (2020). Pathways to social evolution and their evolutionary feedbacks. *Evolution*, *74*(9), 1894–1907. https://doi.org/10.1111/EVO.14054

Bailey, N. W., & Desjonquères, C. (2022). The Indirect Genetic Effect Interaction Coefficient ψ: Theoretically Essential and Empirically Neglected. *Journal of Heredity*, *113*(1). https://doi.org/10.1093/jhered/esab056

Bailey, N. W., & Kölliker, M. (2019). Social runaway: Fisherian elaboration (or reduction) of socially selected traits via indirect genetic effects. *Evolution*, *73*(8). https://doi.org/10.1111/evo.13791

Bailey, N. W., Marie-Orleach, L., & Moore, A. J. (2018). Indirect genetic effects in behavioral ecology: Does behavior play a special role in evolution? In *Behavioral Ecology*. https://doi.org/10.1093/beheco/arx127

Bergsma, R., Kanis, E., Knol, E. F., & Bijma, P. (2008). The contribution of social effects to heritable variation in finishing traits of domestic pigs (Sus scrofa). *Genetics*, *178*(3). https://doi.org/10.1534/genetics.107.084236

Bergsma, R., Mathur, P. K., Kanis, E., Verstegen, M. W. A., Knol, E. F., & Van Arendonk, J. A. M. (2013). Genetic correlations between lactation performance and growing-finishing traits in pigs. *Journal of Animal Science*, *91*(8). https://doi.org/10.2527/jas.2012-6200

Bijma, P. (2010). Estimating indirect genetic effects: Precision of estimates and optimum designs. *Genetics*, *186*(3). https://doi.org/10.1534/genetics.110.120493

Bijma, P., Muir, W. M., Ellen, E. D., Wolf, J. B., & Van Arendonk, J. A. M. (2007). Multilevel selection 2: Estimating the genetic parameters determining inheritance and response to selection. *Genetics*, *175*(1), 289–299. https://doi.org/10.1534/genetics.106.062729

Bijma, P., Muir, W. M., & Van Arendonk, J. A. M. (2007). Multilevel selection 1: Quantitative genetics of inheritance and response to selection. *Genetics*, *175*(1), 277–288. https://doi.org/10.1534/genetics.106.062711

Bijma, P., & Wade, M. J. (2008). The joint effects of kin, multilevel selection and indirect genetic
effects on response to genetic selection. *Journal of Evolutionary Biology*, *21*, 1175–1188.
https://doi.org/10.1111/j.1420-9101.2008.01550.x

Bourke, P.M., Evers, J.B., Bijma, P., van Apeldoorn, D.F., Smulders, M.J., Kuyper, T.W., Mommer, L.
and Bonnema, G., 2021. Breeding beyond monoculture: Putting the "intercrop" into
crops. *Frontiers in Plant Science*, *12*, p.734167. https://doi.org/10.3389/fpls.2021.734167

Bouwman, A. C., Bergsma, R., Duijvesteijn, N., & Bijma, P. (2010). Maternal and social genetic
effects on average daily gain of piglets from birth until weaning. *Journal of Animal Science*,
88(9). https://doi.org/10.2527/jas.2009-2494

Brinker, T., Bijma, P., Visscher, J., Rodenburg, T. B., & Ellen, E. D. (2014). Plumage condition in laying
hens: Genetic parameters for direct and indirect effects in two purebred layer lines. *Genetics
Selection Evolution*, *46*(1). https://doi.org/10.1186/1297-9686-46-33

Brinker, T., Ellen, E. D., Veerkamp, R. F., & Bijma, P. (2015). Predicting direct and indirect breeding
values for survival time in laying hens using repeated measures. *Genetics Selection Evolution*,
47(1). https://doi.org/10.1186/s12711-015-0152-2

Brommer, J. E., & Rattiste, K. (2008). "Hidden" reproductive conflict between mates in a wild bird
population. *Evolution*, *62*, 2326–2333. https://doi.org/10.1111/j.1558-5646.2008.00451.x

Canario, L., Lundeheim, N., & Bijma, P. (2017). The early-life environment of a pig shapes the
phenotypes of its social partners in adulthood. *Heredity*, *118*(6).
https://doi.org/10.1038/hdy.2017.3

Chakrabarty, A., van Kronenberg, P., Toliopoulos, N., & Schielzeth, H. (2019). Direct and indirect
genetic effects on reproductive investment in a grasshopper. *Journal of Evolutionary Biology*,
32(4). https://doi.org/10.1111/jeb.13417

Charmantier, A., Garant, D., & Kruuk, L. E. B. (2014). *Quantitative Genetics in the Wild*. Oxford
University Press.

Chen, C. Y., Kachman, S. D., Johnson, R. K., Newman, S., & Van Vleck, L. D. (2008). Estimation of
genetic parameters for average daily gain using models with competition effects. *Journal of
Animal Science*, *86*(10). https://doi.org/10.2527/jas.2007-0660

Cinar, O., Nakagawa, S., & Viechtbauer, W. (2022). Phylogenetic multilevel meta-analysis: A
simulation study on the importance of modelling the phylogeny. *Methods in Ecology and
Evolution*, *13*(2). https://doi.org/10.1111/2041-210X.13760

Costa E Silva, J., Potts, B. M., Bijma, P., Kerr, R. J., & Pilbeam, D. J. (2013). Genetic control of
interactions among individuals: Contrasting outcomes of indirect genetic effects arising from
neighbour disease infection and competition in a forest tree. *New Phytologist*.
https://doi.org/10.1111/nph.12035

Craig, J. V., & Muir, W. M. (1996). Group selection for adaptation to multiple-hen cages: Beak-
related mortality, feathering, and body weight responses. *Poultry Science*, *75*(3).
https://doi.org/10.3382/ps.0750294

David, I., Sánchez, J. P., & Piles, M. (2018). Longitudinal analysis of direct and indirect effects on
average daily gain in rabbits using a structured antedependence model. *Genetics Selection
Evolution*, *50*(1). https://doi.org/10.1186/s12711-018-0395-9

de Groot, C., Wijnhorst, R. E., Ratz, T., Murray, M., Araya-Ajoy, Y. G., Wright, J., & Dingemanse, N. J.
(2023). The importance of distinguishing individual differences in 'social impact' versus 'social

responsiveness' when quantifying indirect genetic effects on the evolution of social plasticity.
Neuroscience and Biobehavioral Reviews, 144.
https://doi.org/10.1016/J.NEUBIOREV.2022.104996

Dewan, I., Garland, T., Hiramatsu, L., & Careau, V. (2019). I Smell a Mouse: Indirect Genetic Effects
on Voluntary Wheel-Running Distance, Duration and Speed. *Behavior Genetics, 49*(1).
https://doi.org/10.1007/s10519-018-9930-2

Dickerson, G. (1947). Composition of hog carcasses as influenced by heritable differences in rate
and economy of gain. *Iowa Agric Exp Stn Res Bull, 354,* 492–524.

Dochtermann, N. A., Schwab, T., Anderson Berdal, M., Dalos, J., & Royauté, R. (2019). The
Heritability of Behavior: A Meta-analysis. *Journal of Heredity, 110*(4).
https://doi.org/10.1093/jhered/esz023

Dochtermann, N. A., Schwab, T., & Sih, A. (2015). The contribution of additive genetic variation to
personality variation: heritability of personality. *Proceedings of the Royal Society B: Biological*
Sciences, 282(1798), 20142201.

Duijvesteijn, N., Knol, E. F., & Bijma, P. (2012). Direct and associative effects for androstenone and
genetic correlations with backfat and growth in entire male pigs. *Journal of Animal Science,*
90(8). https://doi.org/10.2527/jas.2011-4625

Edward, D. A., Poissant, J., Wilson, A. J., & Chapman, T. (2014). Sexual conflict and interacting
phenotypes: A quantitative genetic analysis of fecundity and copula duration in drosophila
melanogaster. *Evolution, 68*(6). https://doi.org/10.1111/evo.12376

Ellen, E. D., Bas Rodenburg, T., Albers, G. A. A., Elizabeth Bolhuis, J., Camerlink, I., Duijvesteijn, N.,
Knol, E. F., Muir, W. M., Peeters, K., Reimert, I., Sell-Kubiak, E., van Arendonk, J. A. M.,
Visscher, J., & Bijma, P. (2014). The prospects of selection for social genetic effects to improve
welfare and productivity in livestock. In *Frontiers in Genetics, 5*(377).
https://doi.org/10.3389/fgene.2014.00377

Ellen, E. D., Peeters, K., Verhoeven, M., Gols, R., Harvey, J. A., Wade, M. J., Dicke, M., & Bijma, P.
(2016). Direct and indirect genetic effects in life-history traits of flour beetles (Tribolium
castaneum). *Evolution, 70*(1), 207–217. https://doi.org/10.1111/evo.12835

Ellen, E. D., Visscher, J., Van Arendonk, J. A. M., & Bijma, P. (2008). Survival of laying hens: Genetic
parameters for direct and associative effects in three purebred layer lines. *Poultry Science,*
87(2). https://doi.org/10.3382/ps.2007-00374

Evans, S. R., Postma, E., & Sheldon, B. C. (2020). It takes two: Heritable male effects on reproductive
timing but not clutch size in a wild bird population. *Evolution.*
https://doi.org/10.1111/evo.13980

Fairbairn, D. J., Blanckenhorn, W. U., & Székely, T. (2007). Sex, Size and Gender Roles: Evolutionary
Studies of Sexual Size Dimorphism. In *Sex, Size and Gender Roles: Evolutionary Studies of*
Sexual Size Dimorphism. https://doi.org/10.1093/acprof:oso/9780199208784.001.0001

Falconer, D. (1981). *Introduction to Quantitative Genetics* (2nd ed.). Longman.

Falconer, D. S., & Mackay, T. F. C. (1996). *Introduction to quantitative genetics* (Longman, Ed.).

Fisher, D. N., Haines, J. A., Boutin, S., Dantzer, B., Lane, J. E., Coltman, D. W., & McAdam, A. G.
(2019). Indirect effects on fitness between individuals that have never met via an extended
phenotype. In *Ecology Letters* (Vol. 22, Issue 4). https://doi.org/10.1111/ele.13230

Fisher, D. N., & McAdam, A. G. (2019). Indirect genetic effects clarify how traits can evolve even when fitness does not. *Evolution Letters*. https://doi.org/10.1002/evl3.98

Fisher, D. N., Wilson, A. J., Boutin, S., Dantzer, B., Lane, J. E., Coltman, D. W., Gorrell, J. C., & McAdam, A. G. (2019). Social effects of territorial neighbours on the timing of spring breeding in North American red squirrels. *Journal of Evolutionary Biology*, *32*(6), 559–571. https://doi.org/10.1111/jeb.13437

Fisher, R. (1930). *The genetical theory of natural selection*. Clarendon.

Frank, S. A. (2007). All of life is social. *Current Biology,* 17(16). https://doi.org/10.1016/j.cub.2007.06.005

Germain, R. R., Wolak, M. E., Arcese, P., Losdat, S., & Reid, J. M. (2016). Direct and indirect genetic and fine-scale location effects on breeding date in song sparrows. *Journal of Animal Ecology*, *85*(6). https://doi.org/10.1111/1365-2656.12575

Gienapp, P., Fior, S., Guillaume, F., Lasky, J. R., Sork, V. L., & Csilléry, K. (2017). Genomic quantitative genetics to study evolution in the wild. *Trends in Ecology & Evolution*, 32(12), 897-908.

Grafen, A. (1989). The phylogenetic regression. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, *326*(1233). https://doi.org/10.1098/rstb.1989.0106

Griffing, B. (1967). Selection in reference to biological groups. I. Individual and group selection applied to populations of unordered groups. *Australian Journal of Biological Sciences*, *20*, 127–139. https://doi.org/10.1071/BI9670127

Hadfield, J. D., & Wilson, A. J. (2007). Multilevel selection 3: Modeling the effects of interacting individuals as a function of group size. *Genetics*, *177*(1), 667–668. https://doi.org/10.1534/genetics.107.075622

Holtmann, B., Lagisz, M., & Nakagawa, S. (2017). Metabolic rates, and not hormone levels, are a likely mediator of between-individual differences in behaviour: a meta-analysis. Functional Ecology, 31(3), 685-696.

Han, C. S., Tuni, C., Ulcik, J., & Dingemanse, N. J. (2018). Increased developmental density decreases the magnitude of indirect genetic effects expressed during agonistic interactions in an insect. *Evolution*, *72*(11). https://doi.org/10.1111/evo.13600

Hansen, T. F., Pélabon, C., & Houle, D. (2011). Heritability is not Evolvability. *Evolutionary Biology*. https://doi.org/10.1007/s11692-011-9127-6

Hong, J. K., Kim, Y. M., Cho, K. H., Cho, E. S., Lee, D. H., & Choi, T. J. (2019). Genetic association between sow longevity and social genetic effects on growth in pigs. *Asian-Australasian Journal of Animal Sciences*, *32*(8). https://doi.org/10.5713/ajas.18.0789

Hong, J. K., Kim, Y. M., Cho, K. H., Park, J. C., & Lee, D. H. (2017). Reproductive performance of sows selected for divergent social genetic effects for growth. *Animal Reproduction*, *14*. https://doi.org/10.21451/1984-3143-AR946

Houle, D. (1992). Comparing evolvability and variability of quantitative traits. In *Genetics*.

Hsu, W. L., Johnson, R. K., & van Vleck, L. D. (2010). Effect of pen mates on growth, backfat depth, and longissimus muscle area of swine. *Journal of Animal Science*, *88*(3). https://doi.org/10.2527/jas.2009-1879

Hunt, J., Rapkin, J., House, C. M., & Wilson, A. J. (2019). Beyond genes and environments: Indirect
genetic effects and the evolution of behaviour. In *Genes and Behaviour: Beyond Nature-*
Nurture. https://doi.org/10.1002/9781119313663.ch4

Johnston, S. E., Chen, N., & Josephs, E. B. (2022). Taking quantitative genomics into the wild.
Proceedings of the Royal Society B, 289(1989), 20221930.

Khaw, H. L., Ponzoni, R. W., Yee, H. Y., Aziz, M. A. bin, & Bijma, P. (2016). Genetic and non-genetic
indirect effects for harvest weight in the GIFT strain of Nile tilapia (Oreochromis niloticus).
Aquaculture, *450*. https://doi.org/10.1016/j.aquaculture.2015.07.033

Kjaer, J. B., & Sørensen, P. (1997). Feather pecking behaviour in White Leghorns, a genetic study.
British Poultry Science, *38*(4). https://doi.org/10.1080/00071669708417999

Kruuk, L. E. B., Slate, J., & Wilson, A. J. (2008). New answers for old questions: The evolutionary
quantitative genetics of wild animal populations. In *Annual Review of Ecology, Evolution, and*
Systematics (Vol. 39). https://doi.org/10.1146/annurev.ecolsys.39.110707.173542

Liu, J., & Tang, G. (2016). Investigating the contribution of social genetic effect to longer selection
response in a ten generations breeding programme simulated. *Italian Journal of Animal*
Science, *15*(4). https://doi.org/10.1080/1828051X.2016.1248868

Luan, S., Luo, K., Chai, Z., Cao, B., Meng, X., Lu, X., Liu, N., Xu, S., & Kong, J. (2015). An analysis of
indirect genetic effects on adult body weight of the Pacific white shrimp Litopenaeus
vannamei at low rearing density. *Genetics Selection Evolution*, *47*(1).
https://doi.org/10.1186/s12711-015-0164-y

Lynch, M., & Walsch, B. (1998). *Genetics and analysis of quantitative traits*. Sinauer.

Mcadam, A. G., Garant, D., & Wilson, A. J. (2014). *Chapter 6. The effects of others' genes: maternal*
and other indirect genetic effects. 1–33.

McGlothlin, J. W., Moore, A. J., Wolf, J. B., & Brodie, E. D. (2010). Interacting phenotypes and the
evolutionary process. III. Social evolution. *Evolution*, *64*(9), 2558–2574.
https://doi.org/10.1111/j.1558-5646.2010.01012.x

Merilä, J., Sheldon, B. C., & Kruuk, L. E. B. (2001). Explaining stasis: Microevolutionary studies in
natural populations. *Genetica*. https://doi.org/10.1023/A:1013391806317

Michonneau, F., Brown, J. W., & Winter, D. J. (2016). rotl: an R package to interact with the Open
Tree of Life data. *Methods in Ecology and Evolution*, *7*(12). https://doi.org/10.1111/2041-
210X.12593

Modlmeier, A. P., Keiser, C. N., Watters, J. V., Sih, A., & Pruitt, J. N. (2014). The keystone individual
concept: An ecological and evolutionary overview. In *Animal Behaviour* (Vol. 89).
https://doi.org/10.1016/j.anbehav.2013.12.020

Moher, D., Liberati, A., Tetzlaff, J., Altman, D. G., Antes, G., Atkins, D., Barbour, V., Barrowman, N.,
Berlin, J. A., Clark, J., Clarke, M., Cook, D., D'Amico, R., Deeks, J. J., Devereaux, P. J., Dickersin,
K., Egger, M., Ernst, E., Gøtzsche, P. C., … Tugwell, P. (2009). Preferred reporting items for
systematic reviews and meta-analyses: The PRISMA statement. In *PLoS Medicine* (Vol. 6, Issue
7). https://doi.org/10.1371/journal.pmed.1000097

Moore, A. J., Brodie, E. D. I., & Wolf, J. B. (1997). Interacting Phenotypes and the Evolutionary
Process : I . Direct and Indirect Genetic Effects of Social Interactions. *Evolution*, *51*(5), 1352–
1362.

Moore, M. P., Whiteman, H. H., & Martin, R. A. (2019). A mother's legacy: the strength of maternal
effects   in   animal   populations.   In   *Ecology   Letters*   (Vol.   22,   Issue   10).
https://doi.org/10.1111/ele.13351

Mousseau, T. A., & Fox, C. W. (1998). The adaptive significance of maternal effects. In *Trends in*
Ecology and Evolution (Vol. 13, Issue 10). https://doi.org/10.1016/S0169-5347(98)01472-4

Mousseau, T. A., & Roff, D. A. (1987). Natural selection and the heritability of fitness components.
Heredity. https://doi.org/10.1038/hdy.1987.113

Muir, W. M. (2005). Incorporation of competitive effects in forest tree or animal breeding
programs. *Genetics*. https://doi.org/10.1534/genetics.104.035956

Muir, W. M., Bijma, P., & Schinckel, A. (2013). Multilevel selection with kin and non-kin groups,
experimental   results   with   japanese   quail   (coturnix   japonica).   *Evolution*,   *67*(6).
https://doi.org/10.1111/evo.12062

Nakagawa, S., Lagisz, M., Jennions, M. D., Koricheva, J., Noble, D. W. A., Parker, T. H., Sánchez-Tójar,
A., Yang, Y., & O'Dea, R. E. (2022). Methods for testing publication bias in ecological and
evolutionary meta-analyses. In *Methods in Ecology and Evolution* (Vol. 13, Issue 1).
https://doi.org/10.1111/2041-210X.13724

Nakagawa, S., Lagisz, M., O'Dea, R., Pottier, P., Rutkowska, J., Senior, A., Yang, Y., & Noble, D.
(2023). orchaRd 2.0: an R package for visualizing meta-analyses with orchard plots.
EcoEvoRxiv.

Nakagawa, S., & Schielzeth, H. (2013). A general and simple method for obtaining R2 from
generalized   linear   mixed-effects   models.   *Methods   in   Ecology   and   Evolution*,   *4*(2).
https://doi.org/10.1111/j.2041-210x.2012.00261.x

Nathan, R., Monk, C. T., Arlinghaus, R., Adam, T., Alós, J., Assaf, M., ... & Jarić, I. (2022). Big-data
approaches lead to an increased understanding of the ecology of animal movement. *Science*,
375(6582), eabg1780.

Nielsen, H. M., Ask, B., & Madsen, P. (2018). Social genetic effects for growth in pigs differ between
boars and gilts. *Genetics Selection Evolution*, *50*(1). https://doi.org/10.1186/s12711-018-0375-
0

Nielsen, H. M., Monsen, B. B., Ødegård, J., Bijma, P., Damsgård, B., Toften, H., & Olesen, I. (2014).
Direct and social genetic parameters for growth and fin damage traits in Atlantic cod (Gadus
morhua). *Genetics Selection Evolution*, *46*(1). https://doi.org/10.1186/1297-9686-46-5

Oers, K. Van, & Sinn, D. L. (2014). Quantitative and Molecular Genetics of Animal Personality. In
Animal Personalities. https://doi.org/10.7208/chicago/9780226922065.003.0007

Ouzzani, M., Hammady, H., Fedorowicz, Z., & Elmagarmid, A. (2016). Rayyan-a web and mobile app
for systematic reviews. *Systematic Reviews*, *5*(1). https://doi.org/10.1186/s13643-016-0384-4

Paradis, E., & Schliep, K. (2019). Ape 5.0: An environment for modern phylogenetics and
evolutionary   analyses   in   R.   *Bioinformatics*,   *35*(3).
https://doi.org/10.1093/bioinformatics/bty633

Peeters, K., Ellen, E. D., & Bijma, P. (2013). Using pooled data to estimate variance components and
breeding values for traits affected by social interactions. *Genetics Selection Evolution*, *45*(1).
https://doi.org/10.1186/1297-9686-45-27

Peeters, K., Eppink, T. T., Ellen, E. D., Visscher, J., & Bijma, P. (2012). Indirect genetic effects for
survival in domestic chickens (Gallus gallus) are Magnified in crossbred genotypes and show a
parent-of-origin effect. *Genetics*, *192*(2). https://doi.org/10.1534/genetics.112.142554

Petfield, D., Chenoweth, S. F., Rundle, H. D., & Blows, M. W. (2005). Genetic variance in female
condition predicts indirect genetic variance in male sexual display traits. *Proceedings of the*
National Academy of Sciences of the United States of America, *102*(17).
https://doi.org/10.1073/pnas.0409378102

Pick, J.L., Kasper, C., Allegue, H., Dingemanse, N.J., Dochtermann, N.A., Laskowski, K.L., Lima, M.R.,
Schielzeth, H., Westneat, D.F., Wright, J. and Araya-Ajoy, Y.G. (2023). Describing posterior
distributions of variance components: Problems and the use of null distributions to aid
interpretation. Methods in Ecology and Evolution, 14(10), 2557-2574.
https://doi.org/10.1111/2041-210X.14200

Piles, M., David, I., Ramon, J., Canario, L., Rafel, O., Pascual, M., Ragab, M., & Sánchez, J. P. (2017).
Interaction of direct and social genetic effects with feeding regime in growing rabbits. *Genetics*
Selection Evolution, *49*(1). https://doi.org/10.1186/s12711-017-0333-2

Postma, E. (2014). Four decades of estimating heritabilities in wild vertebrate populations:
improved methods, more data, better estimates? In *Quantitative Genetics in the Wild*.
https://doi.org/10.1093/acprof:oso/9780199674237.003.0002

Pujol, B., Blanchet, S., Charmantier, A., Danchin, E., Facon, B., Marrot, P., Roux, F., Scotti, I.,
Teplitsky, C., Thomson, C. E., & Winney, I. (2018). The Missing Response to Selection in the
Wild. In *Trends in Ecology and Evolution* (Vol. 33, Issue 5).
https://doi.org/10.1016/j.tree.2018.02.007

R Core Team. (2021). R core team (2021). In *R: A language and environment for statistical*
computing. R Foundation for Statistical Computing, Vienna, Austria. URL http://www. R-
project. org.

Radersma, R. (2021). Estimating heritability of social phenotypes from social networks. *Methods in*
Ecology and Evolution, 12(1), 42-53.

Ragab, M., Piles, M., Quintanilla, R., & Sánchez, J. P. (2019). Indirect genetic effect model using
feeding behaviour traits to define the degree of interaction between mates: An
implementation in pigs growth rate. *Animal*, *13*(2).
https://doi.org/10.1017/S1751731118001192

Rees, J. A., & Cranston, K. (2017). Automated assembly of a reference taxonomy for phylogenetic
data synthesis. *Biodiversity Data Journal*, *5*(1). https://doi.org/10.3897/BDJ.5.e12581

Rostellato, R., Sartori, C., Bonfatti, V., Chiarot, G., & Carnier, P. (2013). Comparison between direct
and competitive models to investigate variation of carcass and ham quality traits in heavy pigs.
Agriculturae Conspectus Scientificus, *78*(3).

Rostellato, R., Sartori, C., Bonfatti, V., Chiarot, G., & Carnier, P. (2015). Direct and social genetic
effects on body weight at 270 days and carcass and ham quality traits in heavy pigs. *Journal of*
Animal Science, *93*(1). https://doi.org/10.2527/jas.2014-8246

Sánchez-Tójar, A., Nakagawa, S., Sánchez-Fortún, M., Martin, D. A., Ramani, S., Girndt, A., Bókony,
V., Kempenaers, B., Liker, A., Westneat, D. F., Burke, T., & Schroeder, J. (2018). Meta-analysis
challenges a textbook example of status signalling and demonstrates publication bias. *ELife*, *7*.
https://doi.org/10.7554/eLife.37385

Santostefano, F., Allegue, H., Garant, D., Bergeron, P., & Réale, D. (2021). Indirect genetic and
environmental effects on behaviors, morphology, and life-history traits in a wild Eastern
chipmunk population. *Evolution*, *75*(6). https://doi.org/10.1111/evo.14232

Santostefano, F., Wilson, A. J., Niemelä, P. T., & Dingemanse, N. J. (2017). Indirect genetic effects: A
key component of the genetic architecture of behaviour. *Scientific Reports*, *7*(1).
https://doi.org/10.1038/s41598-017-08258-6

Sartori, C., & Mantovani, R. (2012). Effects of inbreeding on fighting ability measured in Aosta
chestnut and Aosta black pied cattle. *Journal of Animal Science*, *90*(9).
https://doi.org/10.2527/jas.2011-4010

Scott, J., & Fuller, J. (1965). *Genetics and the Social Behavior of the Dog*. University of Chicago Press.

Shine, R. (1989). Ecological causes for the evolution of sexual dimorphism: a review of the evidence.
Quarterly Review of Biology, *64*(4). https://doi.org/10.1086/416458

Stirling, D. G., Réale, D., & Roff, D. A. (2002). Selection, structure and the heritability of behaviour.
Journal of Evolutionary Biology, *15*(2), 277–289.

Sun, Y., Ellen, E. D., Van Der Poel, J. J., Parmentier, H. K., & Bijma, P. (2014). Modelling of feather
pecking behavior in beak-trimmed and non-beak-trimmed crossbred laying hens: Variance
component and trait-based approach. *Poultry Science*, *93*(4). https://doi.org/10.3382/ps.2013-
03367

Székely, T., Moore, A. J., & Komdeur, J. (2010). *Social behaviour: genes, ecology and evolution*.
Cambridge University Press.

Teplitsky, C., Mills, J. A., Yarrall, J. W., & Merilä, J. (2010). Indirect genetic effects in a sex-limited
trait: The case of breeding time in red-billed gulls. *Journal of Evolutionary Biology*.
https://doi.org/10.1111/j.1420-9101.2010.01959.x

Trubenová, B., & Hager, R. (2012). Phenotypic and Evolutionary Consequences of Social Behaviours:
Interactions among Individuals Affect Direct Genetic Effects. *PLoS ONE*, *7*(11).
https://doi.org/10.1371/journal.pone.0046273

Trubenová, B., Novak, S., & Hager, R. (2015). Indirect genetic effects and the dynamics of social
interactions. *PLoS ONE*, *10*(5). https://doi.org/10.1371/journal.pone.0126907

Van Vleck, L. D., Cundiff, L. V., & Kocht, R. M. (2007). Effect of competition on gain in feedlot bulls
from Hereford selection lines. *Journal of Animal Science*, *85*(7).
https://doi.org/10.2527/jas.2007-0067

Viechtbauer, W. (2010). Conducting Meta-analysis in R with the metafor package. *Journal of
Statistical Software*, *36*(3).

Wade, M. J. (1976). Group selection among laboratory populations of Tribolium. *Proceedings of the
National Academy of Sciences of the United States of America*, *73*(12).
https://doi.org/10.1073/pnas.73.12.4604

Wade, M. J. (1977). An Experimental Study of Group Selection. *Evolution*, *31*(1).
https://doi.org/10.2307/2407552

Wade, M. J., Bijma, P., Ellen, E. D., & Muir, W. (2010). Group selection and social evolution in
domesticated animals. *Evolutionary Applications*, *3*, 453–465. https://doi.org/10.1111/j.1752-
4571.2010.00147.x

West-Eberhard, M. J. (2003). *Developmental plasticity and evolution*. Oxford University Press.

Westgate, M. J. (2019). revtools: An R package to support article screening for evidence synthesis.
In *Research Synthesis Methods* (Vol. 10, Issue 4). https://doi.org/10.1002/jrsm.1374

Westneat, D. F., & Fox, C. (2010). *Evolutionary behavioral ecology*. Oxford University Press.

Wilson, A. J. (2014). Competition as a source of constraint on life history evolution in natural
populations. *Heredity*, *112*(1), 70–78. https://doi.org/10.1038/hdy.2013.7

Wilson, A. J., Gelin, U., Perron, M., & Réale, D. (2009). Indirect genetic effects and the evolution of
aggression in a vertebrate system Indirect genetic effects and the evolution of aggression in a
vertebrate system. *Proceedings of the Royal Society B: Biological Sciences*, *276*, 533–541.
https://doi.org/10.1098/rspb.2008.1193

Wolf, J. B., Brodie, Cheverud, J. M., Moore, A. J., & Wade, M. J. (1998). Evolutionary consequences
of indirect genetic effects. *Trends in Ecology and Evolution*, *13*(2), 64–69.

Wood, J. L. A., Yates, M. C., & Fraser, D. J. (2016). Are heritability and selection related to
population size in nature? Meta-analysis and conservation implications. In *Evolutionary*
Applications (Vol. 9, Issue 5). https://doi.org/10.1111/eva.12375

Wright, A. J. (1985). Selection for improved yield in inter-specific mixtures or intercrops. Theoretical
and applied genetics, 69, 399-407.

Wu, P., Wang, K., Yang, Q., Zhou, J., Chen, D., Liu, Y., Ma, J., Tang, Q., Jin, L., Xiao, W., Lou, P., Jiang,
A., Jiang, Y., Zhu, L., Li, M., Li, X., & Tang, G. (2019). Whole-genome re-sequencing association
study for direct genetic effects and social genetic effects of six growth traits in Large White
pigs. *Scientific Reports*, *9*(1). https://doi.org/10.1038/s41598-019-45919-0

Yang, Y., Noble, D. W., Spake, R., Senior, A. M., Lagisz, M., & Nakagawa, S. (2023). Measuring
biological generality in meta-analysis: a pluralistic approach to heterogeneity quantification
and stratification. *EcoEvoRxiv*. Preprint. https://doi.org/10.32942/X2RG7X

Author contributions

FS: Conceptualization; Methodology; Investigation; Data Curation; Writing – original draft preparation; Writing – review and editing; Project administration

MM: Conceptualization; Methodology; Validation; Formal analysis; Investigation; Writing – original draft preparation; Writing – review and editing; Visualization

AST: Conceptualization; Methodology; Formal analysis; Investigation; Data Curation; Writing – original draft preparation; Writing – review and editing; Visualization

DF: Conceptualization; Methodology; Validation; Formal analysis; Investigation; Writing – original draft preparation; Writing – review and editing; Visualization

Acknowledgements

We would like to thank Denis Reale, Clint Kelly, and Pierre-Olivier Montiglio for their feedback in the earlier stages of the project, and Daniel W.A. Noble for statistical advice. We thank the participants of the 2023 WAMBAM conference and Alastair Wilson for useful discussion. We also thank Allen Moore, Joel Pick, and one anonymous reviewer for their constructive feedback, which improved the manuscript. We are grateful to the following authors who provided additional data for the meta-analysis when we contacted them: Alastair Wilson, Anasuya Chakrabarty, Cristina Sartori, Esther Ellen, Chang Han, Jane Reid, João Costa e Silva, Jennifer Morinay, Mark Adams, Miriam Piles, Moha Ragab, Ingrid David, Paolo Carnier, Ryan Germain, Setegn Alemu, Simon Evans, and Céline Teplitsky.

Funding

FS was supported by the European Union's Horizon 2020 research and innovation programme under the Marie Sklodowska-Curie Individual Fellowship (INTERACTIVE, Grant Agreement Number 101023262). MM was funded by a Marie Skłodowska-Curie Individual Fellowship (PLASTIC TERN, Grant Agreement Number 793550) and an Alexander von Humboldt Research Fellowship for Postdoctoral Researchers. AST was partially funded by the German Research Foundation (DFG: Deutsche Forschungsgemeinschaft) as part of the SFB TRR 212 (NC3)—Project no. 316099922 and 396782608.

Data and code availability statement

All data and code are available at the following GitHub repository (https://github.com/ASanchez-Tojar/meta-analysis_IGEs). Upon acceptance, all data and code will be provided with a doi via Zenodo.

Conflict of interest statement

The authors declare no competing interests.

Tables

930 **Table 1.** Glossary of notation used in the manuscript with associated names and formulae.

Notation	Name	Formula
DGEs	Direct genetic effects	-
IGEs	Indirect genetic effects	-
V_A	Direct additive genetic variance	-
V_{IGE}	Indirect genetic effects variance	-
V_P	Total phenotypic variance	-
h^2	Heritability (narrow sense)	V_A / V_P
social h^2	Social heritability (narrow sense)	V_{IGE} / V_P
I_{DGE}	Evolvability	$V_A / \bar{x}^2$
I_{IGE}	Social evolvability	$V_{IGE} / \bar{x}^2$
$COV_{DGE\text{-}IGE}$	DGE-IGE covariance	-
$r_{DGE\text{-}IGE}$	DGE-IGE correlation	$COV_{DGE\text{-}IGE} / \sqrt{(V_A * V_{IGE})}$
τ^2	Total heritable variance	$[V_A + Cov_{A\text{-}IGE}*2*(n-1) + V_{IGE}*(n-1)^2] / V_P$
-	"Covariance component" of τ^2	$(Cov_{A\text{-}IGE}*2*[n-1])$
-	"IGE variance component" of τ^2	$(V_{IGE}*[n-1]^2)$
-	Response to individual selection among unrelated individuals	$1 * [V_A + (n-1) Cov_{A\text{-}IGE}]$

 Table 2. List of studies (n = 47) from which data was extracted and included in our meta-analysis on indirect genetic effects.

Study ID	Year	Journal	Number of effect sizes (k)	Species	Reference
IGE0057	2015	Genetics Selection Evolution	1	*Penaeus vannamei*	(Luan et al., 2015)
IGE0144	2013	Agriculturae Conspectus Scientificus	3	*Sus scrofa*	(Rostellato et al., 2013)
IGE0195	2012	Journal Of Animal Science	3	*Sus scrofa*	(Duijvesteijn et al., 2012)*
IGE0196	2016	Journal Of Animal Ecology	1	*Melospiza melodia*	(Germain et al., 2016)
IGE0197	2016	Evolution	1	*Tribolium castaneum*	(Ellen et al., 2016)
IGE0199	2019	Journal Of Evolutionary Biology	3	*Chorthippus biguttulus*	(Chakrabarty et al., 2019)
IGE0202	2015	Journal Of Animal Science	10	*Sus scrofa*	(Rostellato et al., 2015)
IGE0203	2014	Genetics Selection Evolution	19	*Gadus morhua*	(Nielsen et al., 2014)*
IGE0242	2007	Journal Of Animal Science	8	*Bos taurus*	(Van Vleck et al., 2007)*
IGE0243	2010	Journal Of Animal Science	3	*Sus scrofa*	(Hsu et al., 2010)*
IGE0253	2012	Journal Of Animal Science	1	*Bos taurus*	(Sartori & Mantovani, 2012)
IGE0291	2008	Journal Of Animal Science	1	*Sus scrofa*	(Chen et al., 2008)*
IGE0292	2016	Journal Of Animal Breeding And Genetics	3	*Neovison vison*	(Alemu et al., 2016)
IGE0294	2005	Journal Of Animal Science	1	*Sus scrofa*	(Arango et al., 2005)
IGE0347	1997	British Poultry Science	6	*Gallus gallus*	(Kjaer & Sørensen, 1997)
IGE0380	2016	Aquaculture	1	*Oreochromis niloticus*	(Khaw et al., 2016)
IGE0385	2019	Asian-Australasian Journal Of Animal Sciences	1	*Sus scrofa*	(Hong et al., 2019)
IGE0391	2013	Journal Of Animal Science	3	*Sus scrofa*	(Bergsma et al., 2013)*
IGE0406	2005	Proceedings Of The National Academy Of Sciences Of The United States Of America	16	*Drosophila serrata*	(Petfield et al., 2005)

IGE0468	2019	Behavior Genetics	24	*Mus musculus*	(Dewan et al., 2019)
IGE0483	2018	Evolution	6	*Gryllus bimaculatus*	(Han et al., 2018)
IGE0493	2019	Ecology Letters	1	*Tamiasciurus hudsonicus*	(Fisher, Haines, et al., 2019)
IGE0494	2019	Animal	1	*Sus scrofa*	(Ragab et al., 2019)
IGE0501	2009	Proceedings Of The Royal Society B-Biological Sciences	5	*Peromyscus maniculatus sonoriensis*	(Wilson et al., 2009)*
IGE0505	2014	Genetics Selection Evolution	3	*Neovison vison*	(Alemu et al., 2014)*
IGE0507	2012	Genetics	4	*Gallus gallus*	(Peeters et al., 2012)*
IGE0509	2010	Journal Of Evolutionary Biology	1	*Chroicocephalus scopulinus*	(Teplitsky et al., 2010)
IGE0515	2017	Scientific Reports	1	*Gryllus bimaculatus*	(Santostefano et al., 2017)
IGE0537	2017	Genetics Selection Evolution	2	*Oryctolagus cuniculus*	(Piles et al., 2017)
IGE0572	2018	Genetics Selection Evolution	5	*Oryctolagus cuniculus*	(David et al., 2018)
IGE0598	2010	Journal Of Animal Science	1	*Sus scrofa*	(Bouwman et al., 2010)*
IGE0640	2014	Poultry Science	2	*Gallus gallus*	(Sun et al., 2014)
IGE0654	2007	Genetics	1	*Gallus gallus*	(Bijma et al., 2007)
IGE0657*	2013	Evolution	1	*Coturnix japonica*	(Muir et al., 2013)*
IGE0746	2014	Genetics Selection Evolution	8	*Gallus gallus*	(Brinker et al., 2014)*
IGE0752	2015	Genetics Selection Evolution	4	*Gallus gallus*	(Brinker et al., 2015)
IGE0786	2017	Animal Reproduction	1	*Sus scrofa*	(Hong et al., 2017)
IGE0824	2014	Evolution	2	*Drosophila melanogaster*	(Edward et al., 2014)
IGE0852	2019	Journal Of Evolutionary Biology	1	*Tamiasciurus hudsonicus*	(Fisher, Wilson, et al., 2019)
IGE0857	2015	Proceedings Of The Royal Society B: Biological Sciences	1	*Aegithalos caudatus*	(Adams et al., 2015)

IGE0858	2018	Genetics Selection Evolution	2	*Sus scrofa*	(Nielsen et al., 2018)
IGE0889*	2008	Poultry Science	3	*Gallus gallus*	(Ellen et al., 2008)*
IGE0920	2008	Genetics	5	*Sus scrofa*	(Bergsma et al., 2008)
IGE0927	2017	Heredity	1	*Sus scrofa*	(Canario et al., 2017)
IGE1023	2013	Genetics Selection Evolution	2	*Gallus gallus*	(Peeters et al., 2013)
IGE1038	2019	Scientific Reports	6	*Sus scrofa*	(Wu et al., 2019)
IGE1050	2008	Evolution	1	*Larus canus*	(Brommer & Rattiste, 2008)

934   * References included in the qualitative review performed by Ellen et al. (2014: Table 4 in that review).

Figures

Figure 1. Indirect genetic effects occur when the genotype of one individual influences the expression of a phenotype in a conspecific, mediated by social interactions.

Figure 2. Social heritability (*social h²*) from a phylogenetic multilevel meta-analysis with 40 studies
and 146 effect sizes (Aim 1). Orchard plot showing the back-transformed meta-analytic mean as *r*,
95% confidence intervals (thick whisker), 95% prediction intervals (thin whisker) and individual effect
sizes (*r*) scaled by their precision (semitransparent circles). *k* corresponds to the number of effect
sizes, the number of studies is shown in brackets. Note that uncertainty measures (i.e., 95% CI and
95% PI) are generated assuming a normal distribution around the meta-analytic mean, explaining
why some intervals may overlap zero substantially despite the data analyzed being bounded
between 0 and 1.

Figure 3. Social heritability (*social h²*) meta-analytic estimates from phylogenetic multilevel uni-moderator meta-regressions testing the following

moderators (Aim 2): (a) Trait category, (b) Age, (c) Sex, (d) Population type. Orchard plot showing the back-transformed meta-analytic mean as *r*, 95%

confidence intervals (thick whisker), 95% prediction intervals (thin whisker) and individual effect sizes (*r*) scaled by their precision (semitransparent circles).

k corresponds to the number of effect sizes, the number of studies is shown in brackets. Note that uncertainty measures (i.e., 95% CI and 95% PI) are

generated assuming a normal distribution around the meta-analytic mean, explaining why some intervals may overlap zero substantially despite the data

analyzed being bounded between 0 and 1.

Social h²

Figure 4. IGEs and DGEs meta-analytic estimates from phylogenetic multilevel uni-moderator meta-regressions (Aim 3) comparing: (a) V_A vs. V_{IGE}, (b)

narrow-sense h^2 vs. *social* h^2, and (c) I_{DGE} vs. I_{IGE} (evolvability). Orchard plot showing the back-transformed meta-analytic mean as r, 95% confidence intervals

(thick whisker), 95% prediction intervals (thin whisker) and individual effect sizes (r) scaled by their precision (semitransparent circles). k corresponds to the

number of effect sizes, the number of studies is shown in brackets. Note that for (b) and (c) uncertainty measures (i.e., 95% CI and 95% PI) are generated

assuming a normal distribution around the meta-analytic mean, explaining why some intervals may overlap zero substantially despite the data analyzed

being bounded between 0 and 1.

Figure 5. Meta-analytic estimates for Aim 4: (a) phylogenetic multilevel meta-regression comparing narrow-sense heritability (h^2) to total heritable variance

(τ^2); (b) phylogenetic multilevel meta-analysis of $r_{DGE\text{-}IGE}$. Orchard plot showing the original (a) and back-transformed meta-analytic mean as r, 95%

confidence intervals (thick whisker), 95% prediction intervals (thin whisker) and individual effect sizes (r) scaled by their precision (semitransparent circles).

k corresponds to the number of effect sizes, the number of studies is shown in brackets. Note that for (a) uncertainty measures (i.e., 95% CI and 95% PI) are

generated assuming a normal distribution around the meta-analytic mean, explaining why some intervals may overlap zero substantially despite the data

analyzed being bounded between 0 and 1.