**Article Title:** Indirect genetic effects increase the heritable variation available to selection

and are largest for behaviours: a meta-analysis

**Authors**: Francesca Santostefano[1,2*], Maria Moiron[3,4*], Alfredo Sánchez-Tójar[4*], David N

Fisher[5*]

* All authors contributed equally

**Affiliations** :

[1] University of Exeter, Centre for Ecology and Conservation, University of Exeter, Penryn

Campus, Cornwall, U.K.

[2] Université du Québec à Montréal, Département des Sciences Biologiques, Montréal,

Canada

[3] Institute of Avian Research, Wilhelmshaven, 26386, Germany

[4] Department of Evolutionary Biology, Bielefeld University, 33615, Bielefeld, Germany

[5] School of Biological Sciences, University of Aberdeen, King's College, Aberdeen, United

Kingdom

**Corresponding author contact information:**

Francesca Santostefano, PhD

Centre for Ecology and Conservation, University of Exeter,

Penryn Campus, Penryn, Cornwall

TR10 9FE, UK

F.Santostefano2@Exeter.ac.uk

# Title: Indirect genetic effects increase the heritable variation available to selection and are largest for behaviours: a meta-analysis

## Abstract

The evolutionary potential of traits is governed by the amount of heritable variation available to selection. While this is typically quantified based on genetic variation in a focal individual for its own traits (direct genetic effects, DGEs), when social interactions occur, genetic variation in interacting partners can influence a focal individual's traits (indirect genetic effects, IGEs). Theory and studies on domesticated species have suggested IGEs can greatly impact evolutionary trajectories, but whether this is true more broadly remains unclear. Here we perform a systematic review and meta-analysis to quantify the amount of trait variance explained by IGEs and the contribution of IGEs to predictions of adaptive potential. We identified 180 effect sizes from 47 studies across 21 species and found that, on average, IGEs of a single social partner account for a small but statistically significant amount of phenotypic variation (0.03). As IGEs affect the trait values of each interacting group member and due to a typically positive – although statistically nonsignificant – correlation with DGEs ($r_{DGE-IGE}$ = 0.26), IGEs ultimately increase trait heritability substantially from 0.27 (narrow-sense heritability) to 0.45 (total heritable variance). This 66% average increase in heritability suggests IGEs can increase the amount of genetic variation available to selection. Furthermore, whilst showing considerable variation across studies, IGEs were most prominent for behaviours, and to a lesser extent for reproduction and survival, in contrast to morphological, metabolic, physiological, and development traits. Our meta-analysis therefore shows that IGEs tend to enhance the evolutionary potential of traits, especially for those tightly related to interactions with other individuals such as behaviour and reproduction.

## Lay Summary

Predicting evolutionary change is important for breeding better livestock and crops, for understanding how biodiversity arises and how populations respond to environmental change. Normally, these predictions are based on how the genetic variants in an organism influence its own traits (characteristics). However, when organisms socially interact, for instance by fighting or cooperating, then the genes in one individual can influence the traits of others, therefore affecting the potential for evolutionary change. We compared 47 studies across 21 animal species and found that the effect of the genes of a single social partner is small but statistically significant, while the total contribution of social genetic effects to evolutionary potential is large. These effects are particularly important for the evolution of animals' behaviours and reproductive traits, but less so

58  for other traits such as body size and physiology. We also found that, because an individual can

59  interact with many others and influence them all, social interactions can substantially increase the

60  potential for a population to evolve from generation to generation. Our results show how social

61  interactions can potentially alter the evolution of those traits known to respond to social

62  interactions in comparison to standard expectations.

## Introduction

Traits can evolve when they are underpinned by heritable genetic variation and subject to selection.

The (narrow-sense) heritability of a trait is defined as the ratio of the additive genetic variance to the

total phenotypic variance within a population, with a higher ratio indicating greater potential for

evolutionary change in response to selection (Falconer & Mackay, 1996; Fisher, 1930; Lynch &

Walsch, 1998). Environmental variation generally makes up the majority of the remainder of the

total phenotypic variation and is thought not to contribute to evolved changes. However, the

environment of an individual can also include its conspecifics and their traits, such as the phenotype

of the breeding partner, a neighbouring individual, or a competitor. Most organisms live at least part

of their lives socially, interacting when they mate, fight, cooperate, and compete for access to

resources (Frank, 2007; Székely et al., 2010; Westneat & Fox, 2010). Through these social

interactions, individuals can influence the phenotypic expression of traits in conspecifics. If an

individual's effect on others has a heritable component, then genes in one individual influence the

phenotype of others, known as indirect genetic effects (hereafter IGEs; Fig. 1; Dickerson, 1947;

Griffing, 1967; Moore et al., 1997; Scott & Fuller, 1965; Wolf et al., 1998). These IGEs can contribute

to phenotypic evolution alongside the additive genetic variance stemming from an organism's own

genes (direct genetic effects; hereafter DGEs; Fig. 1). However, when measuring the evolutionary

potential of traits, researchers typically only quantify DGEs. Only considering DGEs neglects the total

genetic variance in trait values stemming from other sources, such as IGEs, and can lead to

inaccurate evolutionary predictions (Wolf et al., 1998).

Although the extra genetic variance generated by IGEs can speed up evolutionary change,

IGEs can also reduce, remove, or even reverse the response to selection if DGEs and IGEs are

negatively correlated (Griffing, 1967; Moore et al., 1997). A positive DGE-IGE correlation indicates

that individuals with genes for high trait values elicit higher trait values in their partners (e.g.

aggressive individuals instigating higher aggression in others, Alemu et al., 2014; Wilson et al., 2009).

In contrast, a negative correlation indicates individuals with high trait values are associated with low

trait values in partners (e.g. individuals good at acquiring limited resources cause others to acquire

fewer resources, Wilson, 2014). IGEs have long been appreciated in livestock breeding, where they

can be exploited to increase productivity and improve welfare (see Ellen et al., 2014 for a qualitative

review), as well as in arable farming to increase yield through intercropping (Wright 1985, Bourke et

al. 2021), and recently in human medicine (Baud et al. 2017). IGEs have also been studied in the

context of maternal genetic effects (the heritable effect a mother has on the traits of her offspring

beyond the direct effect of the genes they inherit from her; Dickerson, 1947; McAdam et al., 2014;

Mousseau & Fox, 1998), as well as implicitly within social and sexual runaway dynamics (Bailey &

Kölliker, 2019). Recently, however, we have begun to appreciate that IGEs are applicable to a much

broader array of traits and contexts (McAdam et al., 2014). Examples include the influence of a male

bird on the lay date of his mate (Brommer & Rattiste, 2008; Evans et al., 2020), the elicitation of

aggression from an opponent during a contest (Santostefano et al., 2017; Wilson et al., 2009), and

the growth rate and infection status of neighbours (Costa E Silva et al., 2013). IGEs therefore are

*expected* to alter the evolutionary potential of a wide range of traits, but whether they do is still

unclear. In the wild, trait evolution frequently does not match expectations based on the ratio of

direct genetic variance to phenotypic variance and the strength of selection observed in wild

populations (Merilä et al., 2001; Pujol et al., 2018). Under artificial selection, the observed response

may also not match the expectations based on calculations including only DGEs and the strength of

imposed selection (Ellen et al., 2014; Falconer, 1981; Muir, 2005). Studying the role of IGEs in trait

evolution more broadly is therefore essential to increase the predictability of evolutionary change

and explain these apparent patterns of "evolutionary stasis", both in the wild and under artificial

conditions.

The recent increase in the number of studies providing estimates of IGEs across taxa

presents us a timely opportunity to quantitatively assess the role that IGEs play in trait evolution and

their importance relative to DGEs. Further, by comparing standardised estimates across contexts and

among different trait types, we can assess when IGEs are particularly prevalent, and when trait

evolution will diverge most from that predicted by DGEs alone. For example, behavioural traits are a

key element of any social interaction and are typically considered to be highly plastic. Behavioural

traits may therefore show large IGE variance (Bailey et al., 2018; see Hunt et al., 2019 for a review of

IGEs on behaviours). However, behavioural traits may also vary more in response to the non-social

environment, in which case the proportion of variance in phenotypes due to IGEs may not be so

high. IGEs could also impact traits related to competition such as growth, resource-dependent life-

history traits, or fitness components that are not social traits *per se* (Fisher & McAdam, 2019;

Wilson, 2014), but still depend on the phenotypes of conspecifics. Many studies on IGEs have taken

place on captive populations housed at relatively high densities where individuals cannot avoid

social interactions (reviewed in Ellen et al., 2014). In contrast, organisms in the wild typically live at

lower densities and may have the option of avoiding social interactions when beneficial. There is

therefore the untested possibility that IGEs in the wild are weaker than in captive populations, and

so may not contribute as much to the trait's evolutionary potential, and ultimately, change.

We performed a systematic review and meta-analysis to quantitatively assess the overall

magnitude and generality, and several patterns of variation, in IGEs across the published literature.

We did not however consider maternal effects, as they have recently been reviewed elsewhere (see

Moore et al., 2019). Further, we only analysed effect sizes stemming from the "variance-

partitioning" approach used to estimate IGEs (see Methods for further details) since Bailey and

Desjonquères (2022) recently reviewed IGEs estimates using the "trait-based" approach. A glossary

with definitions of terms used throughout is available in Table 1.

Our aims were as follows:

1.   Quantify the relative magnitude of IGEs across studies (social $h^2$ = variance in IGEs [$V_{IGE}$] /

total phenotypic variance [$V_P$]). We had no clear prediction about its magnitude, but

based on our understanding of the literature we expected social $h^2$ to be relatively low.

2.   Test whether IGEs differ among (a) types of traits, (b) ages, (c) sexes, and (d) wild and

captive populations. Our predictions for (a) were that, because traits closely linked to

fitness such as life-history traits typically have low heritabilities, but high evolvabilities

(see Hansen et al., 2011; Houle, 1992; Postma 2014, for detailed metric interpretations),

the pattern for IGEs variance would be similar. Instead, we expected that behavioural

traits would be highly plastic and therefore may show larger IGEs, but also high

environmental variances, giving low heritabilities, but high evolvabilities (Bailey et al.,

2018). We also predicted for (d) higher effects in captive populations compared to wild

populations given the difficulty of avoiding interactions in captivity. We had no specific

predictions for the remaining sub-aims.

3.   Quantify the magnitude of IGEs in comparison to DGEs. Specifically, we compared: (a)

direct additive genetic variance ($V_A$) vs. $V_{IGE}$, (b) heritability ($h^2$) vs. social $h^2$, and (c) $I_{DGE}$

vs. $I_{IGE}$ (where $I_{DGE} = V_A/\bar{x}^2$ and $I_{IGE} = V_{IGE}/\bar{x}^2$; Bailey et al., 2018). The latter relates

variances to the trait mean to assess evolvability (i.e. a trait's evolutionary potential in

response to selection, Hansen et al., 2011; Houle, 1992). IGEs are mediated by social

interactions, giving a less direct relationship with trait expression than DGEs, and thus

we expected them to be smaller than DGEs.

4.   Test whether IGEs alter evolutionary trajectories, which we evaluated in two ways: (a)

by assessing the overall size and direction of the DGE-IGE correlation ($r_{DGE-IGE}$) to

understand whether IGEs are expected to increase (positive correlations) or decrease

(negative correlations) the rate of evolutionary change in response to direct individual

161            level selection, and (b) by comparing narrow-sense heritability ($h^2$) with total heritable

162            variance ($\tau^2$). Since $h^2$ only considers DGEs, while $\tau^2$ considers DGEs, IGEs, and their

163            covariance, larger $\tau^2$ compared to $h^2$ values would indicate that IGEs increase the

164            amount of genetic variation available to selection and can potentially speed up

165            evolution, while smaller $\tau^2$ values would indicate IGEs can potentially slow down

166            evolution.

## Materials and methods

**Systematic review**

We performed our systematic literature search on the 8[th] of August 2019 in both *Web of Science*

*Core Collection* (databases: SCI-EXPANDED 1900-present, SSCI 1956-present, A&HCI 1975-present,

ESCI 2015-present) and *Scopus*, and consisted of three different steps. First, we searched for studies

on IGEs published across all years using the string: (("indirect genetic effect*") OR ("social genetic

effect*") OR ("associative genetic effect*") OR ("interacting phenotype*")). Second, we performed a

backward search to identify references cited in two key recent reviews on the topic: Ellen et al.

(2014) and Bailey et al. (2018). Third, we performed a forward search to identify references citing at

least one of three seminal theoretical papers on the topic: Bijma & Wade (2008), Moore et al.

(1997), and Wolf et al. (1998). We downloaded all the references and found and removed duplicates

using the R package 'revtools' v.0.3.0 (Westgate, 2019). We screened the titles and abstracts of 1058

unique references using Rayyan (Ouzzani et al., 2016), of which 199 references were selected for

full-text screening. Full-text screening identified 111 studies, of which 63 applied an 'animal model'

approach to analyse their data and were then subjected to data extraction for our meta-analysis. All

authors equally performed both the title-and-abstract and the full-text screening using predefined

decision trees detailing the inclusion and exclusion criteria (Figures S1 and S2). To reduce any

potential observer biases, the title and abstract of 27% of the references (N = 284), and the full text

of 22% of the references (N = 44) were screened by more than one observer. Conflicting decisions

(title-and-abstract: 7% [N = 20]; full-text: 16% [N = 7]) were collectively discussed and resolved. For

more detailed information see our PRISMA diagram (Figure S3; Moher et al., 2009) and the provided

data (https://github.com/ASanchez-Tojar/meta-analysis_IGEs).

**Data extraction**

Data were extracted from the text and tables in the main text and supplementary materials of the

studies identified by our search. All authors equally performed data extraction and the data

extraction of all the papers was double-checked by another author. The original and second

observer discussed and resolved unclear decisions to ensure reliability. To ensure this process was
uniform, all authors extracted data from four selected papers as a pilot at the beginning of this step.

From each included study, we extracted information about the article (year of publication,
journal, author information), study subjects (species, population and years studied, population type
[captive, semi-captive {defined as a population kept in captivity for no more than five generations},
wild], sex [males, females, both] and age [juveniles, adults, both] of included subjects) and type of
trait(s) studied (behaviour, development, metabolism and physiology, morphology, reproduction,
survival). In addition, we extracted the information needed to interpret the statistical estimates of
each animal model, namely whether response variables were mean- or variance-standardized, the
error distribution and any transformations used, mean group size used to estimate IGEs, mean and
standard deviation (SD) of the trait(s) studied (i.e. the response variables), and whether fixed effects
pertaining to focal or partner individual traits were included in the models. Lastly, we extracted
sample sizes (e.g. number of individuals studied, number of observations), pedigree-related
information, and the estimates from the animal models (e.g. direct [$V_A$] and indirect genetic effect
variances [$V_{IGE}$], narrow-sense [$h^2$] and total heritability [$\tau^2$], direct-indirect genetic correlation [$r_{DGE\text{-}IGE}$],
$_{IGE}$], and direct-indirect genetic covariance [$CoV_{A\text{-}IGE}$]).

After data extraction, we excluded 16 studies: (i) 8 studies (k = 13 effect sizes) that did not
contain usable data (i.e. because of being a conference abstract, reporting the same data as other
studies, not estimating IGEs using the animal model approach, or solely reporting estimates from
existing literature), (ii) 1 study (k = 1 effect size) on humans, (iii) 2 studies (k = 7 effect sizes) on
*Eucalyptus globulus* because we did not find data for any other plant species, and (iv) 5 studies (k =
10 effect sizes) that exclusively analysed non-Gaussian traits. Notably, several wild studies fell into
this latter exclusion category, reducing the sample size for that category (see below). Requests for
missing or partially reported data were sent to the corresponding authors of 38 studies using a
standardized email template (provided in Supplementary Materials), from which we obtained
additional data for 25 studies. Finally, note that our meta-analytic dataset included 13 out of the 21
studies included in the qualitative literature review by Ellen et al. (2014; see Table 4 therein), with
the difference being explained by the inclusion/exclusion criteria explained above. For a complete
list    of    studies    and    extracted    variables    see    Table    2    and    the    provided    data
(https://github.com/ASanchez-Tojar/meta-analysis_IGEs).

**Effect size calculation**
Our effect sizes of interest were: (i) relative magnitude of DGEs (narrow-sense heritability or $h^2$) and
IGEs (*social $h^2$*; aims 1, 3 and 4), (ii) unstandardized variances associated with DGEs ($V_A$) and IGEs

($V_{IGE}$), which we log-transformed prior to the analyses to better match model assumptions (aim 3),

(iii) DGE and IGE evolvability ($I_{DGE}$ and $I_{IGE}$; aim 3), (iv) DGE-IGE correlation ($r_{DGE-IGE}$; aim 4), which

when not reported, was calculated as $COV_{A-IGE}$ divided by the square root of the product of $V_A$ and

$V_{IGE}$, and (v) total heritable variance, $\tau^2$ (i.e. the ratio of total heritable variation and phenotypic

variance, aim 4), which was calculated as $[V_A + Cov_{A-IGE}*2*(n-1) + V_{IGE}*(n-1)^2] / V_P$, where n is group

size (Bijma, 2010; Bijma et al., 2007). As in previous meta-analyses (e.g. Dochtermann et al. 2019,

Holtmann et al. 2017), for most of our analyses we treated intra-class correlations as equivalent to

Pearson's correlation coefficients ($r$) and applied Fisher's $Z$ transformation to obtain our main effect

size of interest (i.e. $Zr$ and its associated sampling variance, $VZr$). Therefore, those meta-analytic

estimates correspond to $Zr$ instead of $r$ throughout; however, we present back-transformed values

(i.e. to $r$) in the figures to aid biological interpretation. The variance in $Zr$ (i.e. sampling variance) was

calculated as: $VZr = 1 / (n-3)$, where n represents the number of unique individuals phenotyped. The

different levels of replication analyzed (e.g., number of groups, number of replicates per group) in

the original mixed-effects models from which we obtained the estimates may lead to an

underestimation of sampling variance. To test the robustness of our results to an increase in

sampling variance, we performed sensitivity analyses by rerunning several analyses after having

artificially increased sampling variance by a factor of 10 (i.e., $VZr_{new} = VZr_{original} * 10$). This increase

did not substantially change our conclusions (Supplementary Material). If social $h^2$ was not directly

reported in the corresponding study, we calculated it ourselves as $V_{IGE} / V_P$. If $V_P$ was not reported,

we calculated it ourselves as $V_A / h^2$ when these variables were reported, and then calculated social

$h^2$. These calculations allowed us to add 13 additional effect sizes to our dataset. Additionally, there

were nine estimates of $V_{IGE}$ reported as "0" in the primary literature. To avoid removing those values

(as zeros cannot be log-transformed), for the comparison of $V_A$ and $V_{IGE}$ only (aim 3) we converted

the reported $V_{IGE}$ values to the minimum possible rounded value based on how the "0" was reported

(e.g. reported 0.00 transformed to 0.001). We note that small variances in our dataset might be

upwardly biased due to the likely prevalence of estimates from REML models and/or the likely

preference for reporting means and medians rather than modes in Bayesian models (see Pick et al.

2023).

**Meta-analyses and meta-regressions**

We first ran an intercept-only meta-analytic model to estimate the overall effect size (i.e. meta-

analytic mean) for social $h^2$. We then ran a meta-regression for each of our hypotheses, giving eight

meta-regressions, each containing a single moderator at a time. All meta-regressions were

complete-case meta-regressions (i.e. effect sizes with unreported moderator data were discarded).
In addition, we also ran an intercept-only meta-analysis to estimate the overall effect size for $r_{DGE\text{-}IGE}$.

All models included the following six random effects: (i) phylogeny, which consisted of a
phylogenetic relatedness correlation matrix (Cinar et al., 2022), (ii) species ID, encompassing effect
sizes obtained from the same species and accounts for among-species differences not explained by
phylogeny, (iii) population ID, encompassing effect sizes obtained from the same population, (iv)
study ID, encompassing effect sizes extracted from the same study, (v) group ID, encompassing
effect sizes derived from the same group of animals and allowing us to model different experimental
groups within the same study, and (vi) record ID (unit-level effect), which represents residual and/or
within-study variance in meta-analytic models. The phylogeny was built by searching for the species
in the Open Tree Taxonomy (Rees & Cranston, 2017) and retrieving their phylogenetic relationships
from the Open Tree of Life (Hinchliff et al., 2015) using the R package 'rotl' v.3.0.11 (Michonneau et
al., 2016). We estimated branch lengths following (Grafen, 1989) as implemented in the function
'compute.brlen()' of the R package 'ape' v.5.4-1 (Paradis & Schliep, 2019) and included the
phylogenetic variance–covariance matrix as a random effect in all models. For the meta-regressions
used to compare the (relative) magnitude of DGEs and IGEs as well as to compare narrow-sense
heritability ($h^2$) to total heritable variance ($\tau^2$; hypotheses 3a, 3b, 3c and 4b), since each pair of
estimates (e.g. $V_{IGE}$ and $V_A$) was obtained from the same model, we included an additional random
effect (vii; model ID) that encompassed effect sizes obtained from the same animal model to
account for that extra level of nonindependence.

To evaluate the importance of each potential source of variation, for all meta-analytic
(intercept-only) models, we calculated four heterogeneity metrics ($\sigma^2$, $I^2$, $CV$, $M$) to understand total
and random effect-specific heterogeneities following the pluralistic approach suggested by Yang et
al. (2023). For all meta-regressions, we estimated the percentage of heterogeneity explained by the
moderator as $R^2_{marginal}$ (Nakagawa & Schielzeth, 2013).

We used the programming language R v.4.0.4 (R Core Team, 2021) throughout. All meta-
analytic and meta-regression models were fitted in the R package 'metafor' v.2.5-82 (Viechtbauer,
2010). Results were graphically represented as orchard plots using the R package 'orchaRd' v.2.0
(Nakagawa et al., 2023).

**Testing for sources of bias, including publication bias**
We confirmed the robustness of our results by running several sensitivity analyses that tested: a) the
effect of including fixed effects pertaining to social partners in the original animal models, b)
whether conclusions remained when only using originally reported estimates (i.e. excluding our own

calculations of some effect sizes; see above), c) evidence for publication bias both small-study and

decline effects (following Nakagawa et al., 2022; Sánchez-Tójar et al., 2018) and d) the effect of

potentially underestimating sampling variance. None of these factors proved to be significant

sources of bias, see the Supplementary Materials for further details on our approach and the results.

## Results

301 Our full dataset contained 180 effect sizes from 47 studies published between 1997 and 2019 and

302 involving 21 species. The median and mean number of studies per species was 1 and 2.2,

303 respectively, with *Sus scrofa* (N = 15 studies) and *Gallus gallus* (N = 8 studies) being the most

304 commonly studied species. Note that since we performed complete-case analyses throughout,

305 sample sizes differ slightly across models and research questions.

**Aim 1: What is the relative magnitude of IGEs (social $h^2$)?**

The intercept-only model revealed a small, but statistically significant estimate for social $h^2$ ($Zr$ =

0.031, 95% confidence intervals, CI, [0.012, 0.051], 95% prediction intervals, PI, [-0.061, 0.124], $p$ =

0.002; k = 146 effect sizes, N = 40 studies, 19 species; Figure 2), but heterogeneity was high ($I^2_{total}$ =

93.2%; $CV_{total}$ = 1.46; $M_{total}$ = 0.70). $I^2_{total}$ and $CV_{total}$ indicated that heterogeneity in our dataset is, on

average, around 14 times larger than statistical noise and around 1.5 times larger than the meta-

analytic mean. Random effect-specific $I^2$ estimates showed that 62.9% of heterogeneity was

attributed to study ID, 20.1% to species ID not linked to the phylogenetic tree, and 10.2% to record

ID. Negligible or zero relative heterogeneity was attributable to the phylogenetic tree, group ID, and

population ID.

**Aim 2: Does the magnitude of IGEs vary with intrinsic or extrinsic factors?**

**2a: Trait category**

Social $h^2$ differed among trait categories (Figure 3a) with the moderator explaining 32.1% of the

heterogeneity ($R^2_{marginal}$ = 0.321). The only trait categories with a mean estimate statistically

significantly different from zero were behaviour ($Zr$ = 0.061, 95% CI [0.032, 0.091]) and reproduction

($Zr$ = 0.053, 95% CI [0.014, 0.093]; Figure 3a). Behavioural traits had statistically significantly higher

social $h^2$ values than all other trait categories except for reproduction and survival (see Table S1 for

p-values from all pair-wise tests). The post-hoc Wald tests showed that most of the differences

between the remaining trait categories were not statistically significant, with only reproduction

either being significantly higher than both development and morphology, or showing a trend in that
direction when compared to metabolism and physiology (Table S1).

**2b: Age**

Social $h^2$ differed among age categories and was statistically significantly larger for effect sizes
combining both age classes ($Zr$ = 0.091, 95% CI [0.022, 0.160]) compared to those from juveniles-
only ($Zr$ = 0.013, 95% CI [-0.029, 0.055], p = 0.049) and adults-only ($Zr$ = 0.028, 95% CI [-0.05, 0.062],
p = 0.026; Table S1, Figure 3b). Estimates for juveniles-only and adults-only were not statistically
significantly different from each other (p = 0.403; Table S1, Figure 3b). Overall, the moderator age
explained 6.8% of the heterogeneity ($R^2_{marginal}$ = 0.068). Note that there were only four effect sizes
for the category combining both age classes, and therefore caution should be applied when drawing
conclusions from those differences.

**2c: Sex**

Social $h^2$ differed among sex categories, with social $h^2$ for females-only effect sizes being larger ($Zr$ =
0.063, 95% CI [0.041, 0.084]) than when including both sexes ($Zr$ = 0.006, 95% CI [-0.011, 0.023], p <
0.001) or for only males (though not statistically significantly so; $Zr$ = 0.038, 95% CI [-0.036, 0.112], p
= 0.524; Table S1, Figure 3c). Sex as a moderator explained a large amount of heterogeneity (31.1%;
$R^2_{marginal}$ = 0.311); however, there were only seven effect sizes for male-only data, and, out of the 40
female-only effect sizes, 28 were from chickens in a production setting, which limits our ability to
generalise these results.

**2d: Population type**

Social $h^2$ did not differ among population types (Table S1, Figure 3d) with the moderator explaining
1.5% of the heterogeneity ($R^2_{marginal}$ = 0.015). Social $h^2$ was statistically significantly different from
zero in captive populations ($Zr$ = 0.030, 95% CI [0.006, 0.053]), however, the estimate did not
statistically differ from zero in semicaptive populations ($Zr$ = 0.020, 95% CI [−0.037, 0.076]) and only
marginally so for wild populations ($Zr$ = 0.048, 95% CI [−0.000, 0.097]: Table S1, Figure 3d). The pair-
wise comparisons revealed no statistically significant differences between all three levels ($p$ > 0.448
in all cases; Table S1, Figure 3d). Note, however, that there were only six studies (k = 6 effect sizes)
from wild populations and four (k = 29 effect sizes) from semicaptive populations.

**Aim 3: What is the relative magnitude of IGEs compared to DGEs?**

We ran three meta-regressions to test for the difference in magnitude between $V_{IGE}$ and $V_A$ (aim 3a);

social $h^2$ and $h^2$ (aim 3b); and $I_{IGE}$ and $I_{DGE}$ (aim 3c). We found that regardless of whether IGEs were

estimated as raw variance ($\log(V_{IGE})$ = -3.775, 95% CI [-6.562, -0.988] vs. $\log(V_A)$ = -0.019, 95% CI [-

2.806, 2.767], p < 0.001), variance-standardized variance (social $h^2$: $Zr$ = 0.042, 95% CI [-0.027, 0.111]

vs. $h^2$: $Zr$ = 0.263, 95% CI [0.194, 0.331], p < 0.001), or mean-standardized variance ($I_{IGE}$ = 0.045, 95%

CI [-0.134, 0.223] vs. $I_{DGE}$ = 0.224, 95% CI [0.045, 0.402], p = 0.003), IGEs were smaller than DGEs

(Figure 4). The moderator explained a considerable amount of variance in each meta-regression

(13.5%, 26.8%, and 4.0%, respectively).

**Aim 4: Do IGEs alter evolutionary trajectories?**

The intercept-only model revealed an overall positive but statistically non-significant correlation

between DGEs and IGEs ($r_{DGE-IGE}$; $Zr$ = 0.270, 95% CI [-0.081, 0.621]; Figure 5b), but heterogeneity was

high ($I^2_{total}$ = 99.98; $CV_{total}$ = 3.25; $M_{total}$ = 0.85). This high heterogeneity explains the large 95%

prediction intervals observed (95% PI [-1.497, 2.036]), which shows that, although positive on

average, $r_{DGE-IGE}$ should be expected to range from very negative to very positive depending on the

particular context investigated.

Total heritable variance ($\tau^2$) was stastistically significantly larger ($r$ = 0.449, 95% CI [0.241,

0.656]) than narrow-sense heritability ($h^2$; $r$ = 0.270, 95% CI [0.062, 0.478], p < 0.001) and the

moderator explained 6.7% of the heterogeneity ($R^2_{marginal}$ = 0.067; Figure 5a).

To better understand whether the increase in $\tau^2$ compared to $h^2$ was driven by the IGEs

variance and its relationship with group size ($n$) or due to the seemingly positive $r_{DGE-IGE}$ detected, as

a post-hoc investigation we calculated the two components of $\tau^2$ that stem from $V_{IGE}$, following Bijma

et al., 2007 (*eq. 6*) and qualitatively compared their overall magnitude. These are the "covariance

component" ($Cov_{A-IGE}*2*[n-1]$) and the "IGE variance component" ($V_{IGE}*[n-1]^2$). This additional

analysis showed that the IGEs variance component was seemingly larger (mean = 693.14) than the

covariance component (mean = 241.75), suggesting that it is the IGEs variance component that

drives the increase in $\tau^2$.

In addition, $\tau^2$ represents how much genetic variation is available to *all* types of selection

(including group-level selection). To test how much difference IGEs make to potential responses to

direct, individual-level only selection, we calculated the difference between the predicted response

to direct selection on unrelated individuals in the presence and absence of IGEs, following eq. 15 in

Bijma and Wade (2008), assuming that individuals interact in groups of size $n$ and a direct selection

gradient of 1 (i.e. $1*[V_A + (n-1) Cov_{A-IGE}]$). This post-hoc analysis showed that the response to direct,

individual-level only selection among unrelated individuals was not different in the presence of IGEs

($Zr$ = 0.183, 95% CI [0.048, 0.319]) and in their absence ($Zr$ = 0.225, 95% CI [0.089, 0.360], p = 0.470,

$R^2_{marginal}$ = 0.002).

## Discussion

We performed a meta-analysis to assess the overall strength and patterns of variation in indirect

genetic effects across the published literature. The magnitude of IGEs is small but statistically

significant, with social $h^2$ accounting for, on average, around 3% of the phenotypic variation. A

recent meta-analysis concluded that genetic and non-genetic maternal effects together explained

11% of the phenotypic variance (Moore et al., 2019). Maternal *genetic* effects alone (a type of IGEs

excluded from our own meta-analysis) must be smaller than 11% and, therefore, likely comparable

to our IGEs estimate. The high levels of heterogeneity among effects that we found show that, as

expected, the importance of IGEs varies across contexts, with the 95% prediction intervals

suggesting that once estimate precision is accounted for, *social $h^2$* is expected to range from 0 to

0.12 in most cases. Indeed, we found that factors such as trait type as well as age and sex of

individuals explain substantial heterogeneity in *social $h^2$* estimates across studies. Notably, IGEs were

more prominent for behaviours (0.06) and traits related to reproduction (0.05), supporting the

hypothesis that labile traits are most strongly affected by social interactions. Furthermeore, as

expected, IGEs were smaller in magnitude compared to DGEs, with estimates of direct narrow-sense

heritability ($h^2$ = 0.27) being, on average, 6 to 8 times (depending on the statistical model) larger

than *social $h^2$*. However, despite this large difference, accounting for IGEs and their correlations with

DGEs (i.e. total heritable variance, $\tau^2$) leads to a 66% increase in the total amount of genetic variation

available to selection ($\tau^2$ = 0.45) when compared to narrow-sense $h^2$. This difference is mostly driven

by the relationship between the contribution of IGEs to $\tau^2$ and group size, but also due to the overall

positive correlation bewtween DGEs and IGEs ($r_{DGE\text{-}IGE}$ = 0.26; although not statistically different from

zero). We detected no evidence of publication bias, and our results were robust to data imputations,

potential underestimation of the sampling variance, and several aspects of model structure in the

original studies (see Supplementary Materials). Overall, our results show the general importance of

IGEs for animal populations, the consequences of which we outline below.

*IGEs can enhance the evolutionary potential of traits*

The most impactful consequence of IGEs on traits evolution to emerge from our meta-analysis is that

IGEs enhance evolutionary potential by increasing the amount of genetic variation available to

selection. In their review, Ellen et al., (2014) suggested that total heritable variance ($\tau^2$) seems to be

larger than narrow-sense heritability ($h^2$) in most domesticated populations; we here confirm this

quantitatively and statistically. Two separate but not mutually exclusive mechanisms appear to

facilitate this increase in adaptive potential: (1) the influence of IGEs on many interaction partners,

and (2) a positive DGEs-IGEs correlation ($r_{DGE-IGE}$).

With a post-hoc analysis, we showed that the increase in $\tau^2$ compared to $h^2$ is mainly driven

by the IGEs variance component ($V_{IGE}[n-1]^2$) rather than covariance component ($Cov_{A-IGE}*2*[n-1]$).

This result suggests that there may be greater evolutionary potential when organisms interact in

larger groups, as then the impact of $V_{IGE}$ is magnified; however, $V_{IGE}$ may also decrease, if the

influence of any two random individuals on each other is diluted as n increases (see Arango et al.,

2005; Hadfield & Wilson, 2007; Wade et al., 2010). Bijma (2010) presents a general model for the

relationship of IGEs with group size, but we could not determine where studies included in our meta-

analysis used it (but see Duijvesteijn et al. 2012 and Canario et al. 2017 for examples). The suggested

importance of group size supports results from previous modelling approaches revealing that both

DGEs and IGEs can depend largely on both group size and interaction strength (Liu & Tang, 2016;

Trubenová & Hager, 2012). How interactions occur and whether group size will enhance or dilute

IGEs will be specific to the biology of the species and the setup of the study. In most captive studies,

experimenters enforce one group size across the experiment. Therefore, caution should be exercised

when trying to extrapolate estimates of IGEs to situations where group sizes are outside the range of

those used to estimate the IGEs in the first place (see also attempts to weight interaction strengths

with, for example, social- or spatial-based neighbour networks, and determine effective group size:

Bijma, 2010; Costa E Silva et al., 2013; Fisher et al., 2019; Santostefano et al., 2021). Furthermore,

differences among individuals in their effect on group mates (e.g. "keystone" individuals: Modlmeier

et al., 2014; or among individual variation in "social impact": Araya-Ajoy et al., 2020; de Groot et al.,

2023) could also become important and modulate the magnitude of the effect of IGEs in a group.

The overall positive (though uncertain) correlation between DGEs and IGEs ($r_{DGE-IGE}$ = 0.26)

also indicates that heritable social effects tend to facilitate trait evolution rather than impede it. This

suggests that IGEs may not be a general explanation for phenotypic stasis in the face of directional

selection (Kruuk et al., 2008; Merilä et al., 2001; Pujol et al., 2018; Wilson, 2014). Our result aligns

with a review of the interacting phenotypes coefficient Ψ (Bailey & Desjonquères, 2022), which

quantifies the effect of a trait on the expression of the same or a different trait in social partners.

The sign of Ψ was generally positive, indicating positive feedback between the trait expression in

interacting partners, and suggesting a predicted response to selection that is *greater* than in traits

where Ψ is zero or without IGEs (Wolf et al., 1998). We then estimated the potential response to

selection as an additional comparison for narrow-sense $h^2$ and $\tau^2$, which revealed an important

caveat. The response to individual-level direct selection in the presence of IGEs (in the absence of

relatedness and multilevel selection) was not statistically different from that predicted in their

absence. This lack of statistical difference, in contrast to the results for narrow-sense $h^2$ and $\tau^2$,

appears because it is non-zero relatedness and multilevel selection  that magnify the influence of

$V_{IGE}$ on the response to selection. Relatedness between the focal individual and the individuals

affecting its fitness is crucial for  harnessing the heritable variation for adaptive evolution (see Bijma

2011 for further discussion). Therefore, if relatives interact, the contribution of $V_{IGE}$ to traits

evolution will begreater. That IGEs do not enhance the response to selection when only individual-

level selection occurs may partially explain the surprising lack of response to artificial selection for

improvement of socially-affected traits (Craig & Muir, 1996; Ellen et al., 2014; Wade, 1976, 1977),

and suggests that multilevel selection may be necessary to "unlock" the evolutionary potential of

IGEs (Bijma & Wade, 2008; Muir et al., 2013). However, multilevel selection is not typically

quantified in studies of selection in wild populations, nor always applied in breeding designs. We

therefore do not yet know the general prevalence of multilevel selection and how that impacts the

evolutionary potential of populations, and call researchers in the field to consider quantifying its

influence in future studies.

*IGEs are smaller than DGEs*

As expected, DGEs were larger than IGEs (e.g. narrow-sense $h^2$ [0.27] was more than 6 times larger

than *social $h^2$* [0.03]) across metrics. Our estimates of $h^2$ are in accordance with other estimates of

additive genetic variance derived from animal-model based studies (e.g. 30%: Postma, 2014; 10–

30%: Wood et al., 2016; 22%: Moore et al., 2019). Overall, this result confirms our prediction that

the effect of an individual's own genes on their own phenotype should be larger than the effect of

the genotype of a single social partner, even for traits that are often considered highly plastic in

response to conspecifics, such as behaviours (Bailey et al., 2018; West-Eberhard, 2003).

*IGEs are largest for behaviours and reproductive traits*

IGEs are larger for behaviours and reproductive traits (*social $h^2$* = 0.06 and 0.05, respectively),

compared to other traits categories such as morphology, development, survival, metabolism and

physiology (*social $h^2$* = 0 to 0.02). This higher susceptibility may occur as behavioural and

reproductive traits likely involve feedback dynamics between interacting partners, such as mutual

escalation of aggression in contests (McGlothlin et al., 2010; Trubenová et al., 2015). Similarly,

reproductive traits such as lay date or offspring production should heavily depend on the influence

of the mating partner as the reproductive outcome is in part a joint phenotype, contributed to by

both parents. Further, behavioural traits often have relatively low estimates of $h^2$ (Dochtermann et

al., 2015; Moore et al., 2019; van Oers & Sinn, 2014) and so our analyses shows that traits with

higher *social $h^2$* are those that often have low direct $h^2$. Therefore, while behaviours may show
limited response to direct selection, due to the prominence of IGEs they may respond more to social,
group-level, and other types of selection. Altogether, these findings suggest that behaviours and
reproductive traits might follow a substantially different evolutionary pathway (including indirect
effects), and that evolution for these traits is expected to diverge most from that predicted by DGEs
alone.

*Other sources of heterogeneity in IGEs*

The additional moderators that we tested to try to explain the high heterogeneity observed
showed varying levels of importance. Although life stage explained a non-negligible amount of
heteregenity across effect sizes, the magnitude of IGEs did not clearly differ between early and late
life stages, which could be because our meta-analysis did not include maternal genetic effects, a
type of IGEs expected to be strong for juvenile traits (Moore et al., 2019). The sex of the studied
individuals explained a high amount of heterogeneity across effect sizes, with IGEs being most
important in female only estimates rather than in those including both sexes. However, we note that
we did not have enough male-only estimates for a meaningful comparison (7 effect sizes from only 2
studies) and that 30 out of the 40 female-only estimates corresponded to captive chickens, hence
more diversity in study systems is needed to directly compare sexes and confirm this trend. Indeed,
considering sex differences in the magnitude of IGEs in future studies will be useful as the sexes
often differ in their patterns of social interaction and competition, which is associated with sexually
dimorphic traits (Fairbairn et al., 2007; Shine, 1989).

The scarcity of wild studies hampered our assessment of whether IGEs estimates differ
compared to studies on captive animals. Fortunately, logistical challenges such as capturing
individuals and monitoring their social interactions, along with large sample sizes required by
quantitative genetics models (Charmantier et al., 2014), are gradually being met in recent years. The
development of large collaborative datasets (e.g. Culina et al. 2021), refined tools for spatial and
social network analyses (e.g. Fisher et al. 2019; Radersma 2021), tracking technology (e.g. Nathan et
al. 2022), as well as the ability to estimate genetic variances from genomic rather than pedigree data
(e.g. Gienapp et al. 2017, Johnston et al. 2022), are expected to lead to more studies estimating IGEs
in the wild. These findings mirror results of a meta-analysis estimating DGEs, where there was no
support for heritabilities differing between laboratory and field conditions, nor among wild,
domestic, or semi-domestic species (Dochtermann et al., 2019), although previous studies show that
behaviours measured in the field tended to exhibit higher heritabilities (Mousseau & Roff, 1987; van
Oers & Sinn, 2014; Stirling et al., 2002).

*Conclusions*

Our meta-analysis across 21 animal species highlights the small but statistically significant relative

magnitude of IGEs, yet also highlights their potentially large contribution to adaptive potential. IGEs

were most prominent for behavioural and reproduction traits, possibly suggesting a different

evolutionary route for these traits, despite the considerable variation across studies. Not accounting

for IGEs underestimates the total amount of genetic variance available to selection in most contexts.

IGEs can instead enhance the evolutionary potential of traits and speed up evolutionary rates by

facilitating a response to selection, possibly promoting the adaptation of populations to rapidly

changing environmental conditions. The increase in adaptive potential was mostly due to the

relationship between the contribution of IGEs to $\tau^2$ and group size, but also due to the correlation

between IGEs and DGEs tending to be, on average, positive across studies. However, it may take the

action of multilevel selection to realise a population's full potential to respond to selection in the

presence of IGEs, and therefore understanding the interplay between IGEs, multilevel selection, and

relatedness will be key for accurately predicting evolutionary change. Additionally, future studies

should expand our knowledge to underrepresented taxonomic groups, particularly focusing on wild

populations, which still remain severely understudied.

## References

Adams, M. J., Robinson, M. R., Mannarelli, M., & Hatchwell, B. J. (2015). Social genetic and social environment effects on parental and helper care in a cooperatively breeding bird. *Proceedings of the Royal Society B: Biological Sciences*, *282*(1810), 20150689. https://doi.org/10.1098/rspb.2015.0689

Alemu, S. W., Berg, P., Janss, L., & Bijma, P. (2016). Estimation of indirect genetic effects in group-housed mink (Neovison vison) should account for systematic interactions either due to kin or sex. *Journal of Animal Breeding and Genetics*, *133*(1), 43–50. https://doi.org/10.1111/jbg.12163

Alemu, S. W., Bijma, P., Møller, S. H., Janss, L., & Berg, P. (2014). Indirect genetic effects contribute substantially to heritable variation in aggression-related traits in group-housed mink (Neovison vison). *Genetics Selection Evolution*, *46*, 30. https://doi.org/10.1186/1297-9686-46-30

Arango, J., Misztal, I., Tsuruta, S., Culbertson, M., & Herring, W. (2005a). Estimation of variance components including competitive effects of Large White growing gilts. *Journal of Animal Science*, *83*(6). https://doi.org/10.2527/2005.8361241x

Arango, J., Misztal, I., Tsuruta, S., Culbertson, M., & Herring, W. (2005b). Threshold-linear estimation of genetic parameters for farrowing mortality, litter size, and test performance of Large White sows. *Journal of Animal Science*, *83*(3). https://doi.org/10.2527/2005.833499x

Araya-Ajoy, Y. G., Westneat, D. F., & Wright, J. (2020). Pathways to social evolution and their evolutionary feedbacks. *Evolution*, *74*(9), 1894–1907. https://doi.org/10.1111/EVO.14054

Bailey, N. W., & Desjonquères, C. (2022). The Indirect Genetic Effect Interaction Coefficient ψ: Theoretically Essential and Empirically Neglected. *Journal of Heredity*, *113*(1). https://doi.org/10.1093/jhered/esab056

Bailey, N. W., & Kölliker, M. (2019). Social runaway: Fisherian elaboration (or reduction) of socially selected traits via indirect genetic effects. *Evolution*, *73*(8). https://doi.org/10.1111/evo.13791

Bailey, N. W., Marie-Orleach, L., & Moore, A. J. (2018). Indirect genetic effects in behavioral ecology: Does behavior play a special role in evolution? In *Behavioral Ecology*. https://doi.org/10.1093/beheco/arx127

Baud, A., Mulligan, M. K., Casale, F. P., Ingels, J. F., Bohl, C. J., Callebert, J., ... & Stegle, O. (2017). Genetic variation in the social environment contributes to health and disease. *PLoS genetics*, *13*(1), e1006498.

Bergsma, R., Kanis, E., Knol, E. F., & Bijma, P. (2008). The contribution of social effects to heritable variation in finishing traits of domestic pigs (Sus scrofa). *Genetics*, *178*(3). https://doi.org/10.1534/genetics.107.084236

Bergsma, R., Mathur, P. K., Kanis, E., Verstegen, M. W. A., Knol, E. F., & Van Arendonk, J. A. M. (2013). Genetic correlations between lactation performance and growing-finishing traits in pigs. *Journal of Animal Science*, *91*(8). https://doi.org/10.2527/jas.2012-6200

Bijma, P. (2010). Estimating indirect genetic effects: Precision of estimates and optimum designs. *Genetics*, *186*(3). https://doi.org/10.1534/genetics.110.120493

Bijma, P., Muir, W. M., Ellen, E. D., Wolf, J. B., & Van Arendonk, J. A. M. (2007). Multilevel selection 2: Estimating the genetic parameters determining inheritance and response to selection. *Genetics*, *175*(1), 289–299. https://doi.org/10.1534/genetics.106.062729

Bijma, P., Muir, W. M., & Van Arendonk, J. A. M. (2007). Multilevel selection 1: Quantitative
genetics of inheritance and response to selection. *Genetics*, *175*(1), 277–288.
https://doi.org/10.1534/genetics.106.062711

Bijma, P., & Wade, M. J. (2008). The joint effects of kin, multilevel selection and indirect genetic
effects on response to genetic selection. *Journal of Evolutionary Biology*, *21*, 1175–1188.
https://doi.org/10.1111/j.1420-9101.2008.01550.x

Bourke, P.M., Evers, J.B., Bijma, P., van Apeldoorn, D.F., Smulders, M.J., Kuyper, T.W., Mommer, L.
and Bonnema, G., 2021. Breeding beyond monoculture: Putting the "intercrop" into
crops. *Frontiers in Plant Science*, *12*, p.734167. https://doi.org/10.3389/fpls.2021.734167

Bouwman, A. C., Bergsma, R., Duijvesteijn, N., & Bijma, P. (2010). Maternal and social genetic
effects on average daily gain of piglets from birth until weaning. *Journal of Animal Science*,
*88*(9). https://doi.org/10.2527/jas.2009-2494

Brinker, T., Bijma, P., Visscher, J., Rodenburg, T. B., & Ellen, E. D. (2014). Plumage condition in laying
hens: Genetic parameters for direct and indirect effects in two purebred layer lines. *Genetics
Selection Evolution*, *46*(1). https://doi.org/10.1186/1297-9686-46-33

Brinker, T., Ellen, E. D., Veerkamp, R. F., & Bijma, P. (2015). Predicting direct and indirect breeding
values for survival time in laying hens using repeated measures. *Genetics Selection Evolution*,
*47*(1). https://doi.org/10.1186/s12711-015-0152-2

Brommer, J. E., & Rattiste, K. (2008). "Hidden" reproductive conflict between mates in a wild bird
population. *Evolution*, *62*, 2326–2333. https://doi.org/10.1111/j.1558-5646.2008.00451.x

Canario, L., Lundeheim, N., & Bijma, P. (2017). The early-life environment of a pig shapes the
phenotypes of its social partners in adulthood. *Heredity*, *118*(6).
https://doi.org/10.1038/hdy.2017.3

Chakrabarty, A., van Kronenberg, P., Toliopoulos, N., & Schielzeth, H. (2019). Direct and indirect
genetic effects on reproductive investment in a grasshopper. *Journal of Evolutionary Biology*,
*32*(4). https://doi.org/10.1111/jeb.13417

Charmantier, A., Garant, D., & Kruuk, L. E. B. (2014). *Quantitative Genetics in the Wild*. Oxford
University Press.

Chen, C. Y., Kachman, S. D., Johnson, R. K., Newman, S., & Van Vleck, L. D. (2008). Estimation of
genetic parameters for average daily gain using models with competition effects. *Journal of
Animal Science*, *86*(10). https://doi.org/10.2527/jas.2007-0660

Cinar, O., Nakagawa, S., & Viechtbauer, W. (2022). Phylogenetic multilevel meta-analysis: A
simulation study on the importance of modelling the phylogeny. *Methods in Ecology and
Evolution*, *13*(2). https://doi.org/10.1111/2041-210X.13760

Costa E Silva, J., Potts, B. M., Bijma, P., Kerr, R. J., & Pilbeam, D. J. (2013). Genetic control of
interactions among individuals: Contrasting outcomes of indirect genetic effects arising from
neighbour disease infection and competition in a forest tree. *New Phytologist*.
https://doi.org/10.1111/nph.12035

Craig, J. V., & Muir, W. M. (1996). Group selection for adaptation to multiple-hen cages: Beak-
related mortality, feathering, and body weight responses. *Poultry Science*, *75*(3).
https://doi.org/10.3382/ps.0750294

David, I., Sánchez, J. P., & Piles, M. (2018). Longitudinal analysis of direct and indirect effects on average daily gain in rabbits using a structured antedependence model. *Genetics Selection Evolution*, *50*(1). https://doi.org/10.1186/s12711-018-0395-9

de Groot, C., Wijnhorst, R. E., Ratz, T., Murray, M., Araya-Ajoy, Y. G., Wright, J., & Dingemanse, N. J. (2023). The importance of distinguishing individual differences in 'social impact' versus 'social responsiveness' when quantifying indirect genetic effects on the evolution of social plasticity. *Neuroscience and Biobehavioral Reviews*, *144*. https://doi.org/10.1016/J.NEUBIOREV.2022.104996

Dewan, I., Garland, T., Hiramatsu, L., & Careau, V. (2019). I Smell a Mouse: Indirect Genetic Effects on Voluntary Wheel-Running Distance, Duration and Speed. *Behavior Genetics*, *49*(1). https://doi.org/10.1007/s10519-018-9930-2

Dickerson, G. (1947). Composition of hog carcasses as influenced by heritable differences in rate and economy of gain. *Iowa Agric Exp Stn Res Bull*, *354*, 492–524.

Dochtermann, N. A., Schwab, T., Anderson Berdal, M., Dalos, J., & Royauté, R. (2019). The Heritability of Behavior: A Meta-analysis. *Journal of Heredity*, *110*(4). https://doi.org/10.1093/jhered/esz023

Dochtermann, N. A., Schwab, T., & Sih, A. (2015). The contribution of additive genetic variation to personality variation⬚: heritability of personality. *Proceedings of the Royal Society B: Biological Sciences*, *282*(1798), 20142201.

Duijvesteijn, N., Knol, E. F., & Bijma, P. (2012). Direct and associative effects for androstenone and genetic correlations with backfat and growth in entire male pigs. *Journal of Animal Science*, *90*(8). https://doi.org/10.2527/jas.2011-4625

Edward, D. A., Poissant, J., Wilson, A. J., & Chapman, T. (2014). Sexual conflict and interacting phenotypes: A quantitative genetic analysis of fecundity and copula duration in drosophila melanogaster. *Evolution*, *68*(6). https://doi.org/10.1111/evo.12376

Ellen, E. D., Bas Rodenburg, T., Albers, G. A. A., Elizabeth Bolhuis, J., Camerlink, I., Duijvesteijn, N., Knol, E. F., Muir, W. M., Peeters, K., Reimert, I., Sell-Kubiak, E., van Arendonk, J. A. M., Visscher, J., & Bijma, P. (2014). The prospects of selection for social genetic effects to improve welfare and productivity in livestock. In *Frontiers in Genetics,* *5*(377). https://doi.org/10.3389/fgene.2014.00377

Ellen, E. D., Peeters, K., Verhoeven, M., Gols, R., Harvey, J. A., Wade, M. J., Dicke, M., & Bijma, P. (2016). Direct and indirect genetic effects in life-history traits of flour beetles (Tribolium castaneum). *Evolution*, *70*(1), 207–217. https://doi.org/10.1111/evo.12835

Ellen, E. D., Visscher, J., Van Arendonk, J. A. M., & Bijma, P. (2008). Survival of laying hens: Genetic parameters for direct and associative effects in three purebred layer lines. *Poultry Science*, *87*(2). https://doi.org/10.3382/ps.2007-00374

Evans, S. R., Postma, E., & Sheldon, B. C. (2020). It takes two: Heritable male effects on reproductive timing but not clutch size in a wild bird population. *Evolution*. https://doi.org/10.1111/evo.13980

Fairbairn, D. J., Blanckenhorn, W. U., & Székely, T. (2007). Sex, Size and Gender Roles: Evolutionary Studies of Sexual Size Dimorphism. In *Sex, Size and Gender Roles: Evolutionary Studies of Sexual Size Dimorphism*. https://doi.org/10.1093/acprof:oso/9780199208784.001.0001

Falconer, D. (1981). *Introduction to Quantitative Genetics* (2nd ed.). Longman.

Falconer, D. S., & Mackay, T. F. C. (1996). *Introduction to quantitative genetics* (Longman, Ed.).

Fisher, D. N., Haines, J. A., Boutin, S., Dantzer, B., Lane, J. E., Coltman, D. W., & McAdam, A. G.
(2019). Indirect effects on fitness between individuals that have never met via an extended
phenotype. In *Ecology Letters* (Vol. 22, Issue 4). https://doi.org/10.1111/ele.13230

Fisher, D. N., & McAdam, A. G. (2019). Indirect genetic effects clarify how traits can evolve even
when fitness does not. *Evolution Letters*. https://doi.org/10.1002/evl3.98

Fisher, D. N., Wilson, A. J., Boutin, S., Dantzer, B., Lane, J. E., Coltman, D. W., Gorrell, J. C., &
McAdam, A. G. (2019). Social effects of territorial neighbours on the timing of spring breeding
in North American red squirrels. *Journal of Evolutionary Biology*, *32*(6), 559–571.
https://doi.org/10.1111/jeb.13437

Fisher, R. (1930). *The genetical theory of natural selection*. Clarendon.

Frank,    S.    A.    (2007).    All    of    life    is    social.    *Current    Biology,*    17(16).
https://doi.org/10.1016/j.cub.2007.06.005

Germain, R. R., Wolak, M. E., Arcese, P., Losdat, S., & Reid, J. M. (2016). Direct and indirect genetic
and fine-scale location effects on breeding date in song sparrows. *Journal of Animal Ecology*,
*85*(6). https://doi.org/10.1111/1365-2656.12575

Gienapp, P., Fior, S., Guillaume, F., Lasky, J. R., Sork, V. L., & Csilléry, K. (2017). Genomic quantitative
genetics to study evolution in the wild. *Trends in Ecology & Evolution*, 32(12), 897-908.

Grafen, A. (1989). The phylogenetic regression. *Philosophical Transactions of the Royal Society of
London. Series B, Biological Sciences*, *326*(1233). https://doi.org/10.1098/rstb.1989.0106

Griffing, B. (1967). Selection in reference to biological groups. I. Individual and group selection
applied to populations of unordered groups. *Australian Journal of Biological Sciences*, *20*, 127–
139. https://doi.org/10.1071/BI9670127

Hadfield, J. D., & Wilson, A. J. (2007). Multilevel selection 3: Modeling the effects of interacting
individuals    as    a    function    of    group    size.    *Genetics*,    *177*(1),    667–668.
https://doi.org/10.1534/genetics.107.075622

Holtmann, B., Lagisz, M., & Nakagawa, S. (2017). Metabolic rates, and not hormone levels, are a
likely mediator of between-individual differences in behaviour: a meta-analysis. Functional
Ecology, 31(3), 685-696.

Han, C. S., Tuni, C., Ulcik, J., & Dingemanse, N. J. (2018). Increased developmental density decreases
the magnitude of indirect genetic effects expressed during agonistic interactions in an insect.
*Evolution*, *72*(11). https://doi.org/10.1111/evo.13600

Hansen, T. F., Pélabon, C., & Houle, D. (2011). Heritability is not Evolvability. *Evolutionary Biology*.
https://doi.org/10.1007/s11692-011-9127-6

Hong, J. K., Kim, Y. M., Cho, K. H., Cho, E. S., Lee, D. H., & Choi, T. J. (2019). Genetic association
between sow longevity and social genetic effects on growth in pigs. *Asian-Australasian Journal
of Animal Sciences*, *32*(8). https://doi.org/10.5713/ajas.18.0789

Hong, J. K., Kim, Y. M., Cho, K. H., Park, J. C., & Lee, D. H. (2017). Reproductive performance of sows
selected    for    divergent    social    genetic    effects    for    growth.    *Animal    Reproduction*,    *14*.
https://doi.org/10.21451/1984-3143-AR946

Houle, D. (1992). Comparing evolvability and variability of quantitative traits. In *Genetics*.

Hsu, W. L., Johnson, R. K., & van Vleck, L. D. (2010). Effect of pen mates on growth, backfat depth,
and longissimus muscle area of swine. *Journal of Animal Science*, *88*(3).
https://doi.org/10.2527/jas.2009-1879

Hunt, J., Rapkin, J., House, C. M., & Wilson, A. J. (2019). Beyond genes and environments: Indirect
genetic effects and the evolution of behaviour. In *Genes and Behaviour: Beyond Nature-*
*Nurture*. https://doi.org/10.1002/9781119313663.ch4

Johnston, S. E., Chen, N., & Josephs, E. B. (2022). Taking quantitative genomics into the wild.
*Proceedings of the Royal Society B*, 289(1989), 20221930.

Khaw, H. L., Ponzoni, R. W., Yee, H. Y., Aziz, M. A. bin, & Bijma, P. (2016). Genetic and non-genetic
indirect effects for harvest weight in the GIFT strain of Nile tilapia (Oreochromis niloticus).
*Aquaculture*, *450*. https://doi.org/10.1016/j.aquaculture.2015.07.033

Kjaer, J. B., & Sørensen, P. (1997). Feather pecking behaviour in White Leghorns, a genetic study.
*British Poultry Science*, *38*(4). https://doi.org/10.1080/00071669708417999

Kruuk, L. E. B., Slate, J., & Wilson, A. J. (2008). New answers for old questions: The evolutionary
quantitative genetics of wild animal populations. In *Annual Review of Ecology, Evolution, and*
*Systematics* (Vol. 39). https://doi.org/10.1146/annurev.ecolsys.39.110707.173542

Liu, J., & Tang, G. (2016). Investigating the contribution of social genetic effect to longer selection
response in a ten generations breeding programme simulated. *Italian Journal of Animal*
*Science*, *15*(4). https://doi.org/10.1080/1828051X.2016.1248868

Luan, S., Luo, K., Chai, Z., Cao, B., Meng, X., Lu, X., Liu, N., Xu, S., & Kong, J. (2015). An analysis of
indirect genetic effects on adult body weight of the Pacific white shrimp Litopenaeus
vannamei at low rearing density. *Genetics Selection Evolution*, *47*(1).
https://doi.org/10.1186/s12711-015-0164-y

Lynch, M., & Walsch, B. (1998). *Genetics and analysis of quantitative traits*. Sinauer.

Mcadam, A. G., Garant, D., & Wilson, A. J. (2014). *Chapter 6. The effects of others' genes: maternal*
*and other indirect genetic effects*. 1–33.

McGlothlin, J. W., Moore, A. J., Wolf, J. B., & Brodie, E. D. (2010). Interacting phenotypes and the
evolutionary process. III. Social evolution. *Evolution*, *64*(9), 2558–2574.
https://doi.org/10.1111/j.1558-5646.2010.01012.x

Merilä, J., Sheldon, B. C., & Kruuk, L. E. B. (2001). Explaining stasis: Microevolutionary studies in
natural populations. *Genetica*. https://doi.org/10.1023/A:1013391806317

Michonneau, F., Brown, J. W., & Winter, D. J. (2016). rotl: an R package to interact with the Open
Tree of Life data. *Methods in Ecology and Evolution*, *7*(12). https://doi.org/10.1111/2041-
210X.12593

Modlmeier, A. P., Keiser, C. N., Watters, J. V., Sih, A., & Pruitt, J. N. (2014). The keystone individual
concept: An ecological and evolutionary overview. In *Animal Behaviour* (Vol. 89).
https://doi.org/10.1016/j.anbehav.2013.12.020

Moher, D., Liberati, A., Tetzlaff, J., Altman, D. G., Antes, G., Atkins, D., Barbour, V., Barrowman, N.,
Berlin, J. A., Clark, J., Clarke, M., Cook, D., D'Amico, R., Deeks, J. J., Devereaux, P. J., Dickersin,
K., Egger, M., Ernst, E., Gøtzsche, P. C., … Tugwell, P. (2009). Preferred reporting items for
systematic reviews and meta-analyses: The PRISMA statement. In *PLoS Medicine* (Vol. 6, Issue
7). https://doi.org/10.1371/journal.pmed.1000097

Moore, A. J., Brodie, E. D. I., & Wolf, J. B. (1997). Interacting Phenotypes and the Evolutionary
Process : I . Direct and Indirect Genetic Effects of Social Interactions. *Evolution*, *51*(5), 1352–
1362.

Moore, M. P., Whiteman, H. H., & Martin, R. A. (2019). A mother's legacy: the strength of maternal
effects in animal populations. In *Ecology Letters* (Vol. 22, Issue 10).
https://doi.org/10.1111/ele.13351

Mousseau, T. A., & Fox, C. W. (1998). The adaptive significance of maternal effects. In *Trends in*
*Ecology and Evolution* (Vol. 13, Issue 10). https://doi.org/10.1016/S0169-5347(98)01472-4

Mousseau, T. A., & Roff, D. A. (1987). Natural selection and the heritability of fitness components.
*Heredity*. https://doi.org/10.1038/hdy.1987.113

Muir, W. M. (2005). Incorporation of competitive effects in forest tree or animal breeding
programs. *Genetics*. https://doi.org/10.1534/genetics.104.035956

Muir, W. M., Bijma, P., & Schinckel, A. (2013). Multilevel selection with kin and non-kin groups,
experimental results with japanese quail (coturnix japonica). *Evolution*, *67*(6).
https://doi.org/10.1111/evo.12062

Nakagawa, S., Lagisz, M., Jennions, M. D., Koricheva, J., Noble, D. W. A., Parker, T. H., Sánchez-Tójar,
A., Yang, Y., & O'Dea, R. E. (2022). Methods for testing publication bias in ecological and
evolutionary meta-analyses. In *Methods in Ecology and Evolution* (Vol. 13, Issue 1).
https://doi.org/10.1111/2041-210X.13724

Nakagawa, S., Lagisz, M., O'Dea, R., Pottier, P., Rutkowska, J., Senior, A., Yang, Y., & Noble, D.
(2023). orchaRd 2.0: an R package for visualizing meta-analyses with orchard plots.
*EcoEvoRxiv*.

Nakagawa, S., & Schielzeth, H. (2013). A general and simple method for obtaining R2 from
generalized linear mixed-effects models. *Methods in Ecology and Evolution*, *4*(2).
https://doi.org/10.1111/j.2041-210x.2012.00261.x

Nathan, R., Monk, C. T., Arlinghaus, R., Adam, T., Alós, J., Assaf, M., … & Jarić, I. (2022). Big-data
approaches lead to an increased understanding of the ecology of animal movement. *Science*,
375(6582), eabg1780.

Nielsen, H. M., Ask, B., & Madsen, P. (2018). Social genetic effects for growth in pigs differ between
boars and gilts. *Genetics Selection Evolution*, *50*(1). https://doi.org/10.1186/s12711-018-0375-
0

Nielsen, H. M., Monsen, B. B., Ødegård, J., Bijma, P., Damsgård, B., Toften, H., & Olesen, I. (2014).
Direct and social genetic parameters for growth and fin damage traits in Atlantic cod (Gadus
morhua). *Genetics Selection Evolution*, *46*(1). https://doi.org/10.1186/1297-9686-46-5

Oers, K. Van, & Sinn, D. L. (2014). Quantitative and Molecular Genetics of Animal Personality. In
*Animal Personalities*. https://doi.org/10.7208/chicago/9780226922065.003.0007

Ouzzani, M., Hammady, H., Fedorowicz, Z., & Elmagarmid, A. (2016). Rayyan-a web and mobile app
for systematic reviews. *Systematic Reviews*, *5*(1). https://doi.org/10.1186/s13643-016-0384-4

Paradis, E., & Schliep, K. (2019). Ape 5.0: An environment for modern phylogenetics and
evolutionary analyses in R. *Bioinformatics*, *35*(3).
https://doi.org/10.1093/bioinformatics/bty633

Peeters, K., Ellen, E. D., & Bijma, P. (2013). Using pooled data to estimate variance components and
breeding values for traits affected by social interactions. *Genetics Selection Evolution*, *45*(1).
https://doi.org/10.1186/1297-9686-45-27

Peeters, K., Eppink, T. T., Ellen, E. D., Visscher, J., & Bijma, P. (2012). Indirect genetic effects for
survival in domestic chickens (Gallus gallus) are Magnified in crossbred genotypes and show a
parent-of-origin effect. *Genetics*, *192*(2). https://doi.org/10.1534/genetics.112.142554

Petfield, D., Chenoweth, S. F., Rundle, H. D., & Blows, M. W. (2005). Genetic variance in female
condition predicts indirect genetic variance in male sexual display traits. *Proceedings of the
National Academy of Sciences of the United States of America*, *102*(17).
https://doi.org/10.1073/pnas.0409378102

Pick, J.L., Kasper, C., Allegue, H., Dingemanse, N.J., Dochtermann, N.A., Laskowski, K.L., Lima, M.R.,
Schielzeth, H., Westneat, D.F., Wright, J. and Araya-Ajoy, Y.G. (2023). Describing posterior
distributions of variance components: Problems and the use of null distributions to aid
interpretation. Methods in Ecology and Evolution, 14(10), 2557-2574.
https://doi.org/10.1111/2041-210X.14200

Piles, M., David, I., Ramon, J., Canario, L., Rafel, O., Pascual, M., Ragab, M., & Sánchez, J. P. (2017).
Interaction of direct and social genetic effects with feeding regime in growing rabbits. *Genetics
Selection Evolution*, *49*(1). https://doi.org/10.1186/s12711-017-0333-2

Postma, E. (2014). Four decades of estimating heritabilities in wild vertebrate populations:
improved methods, more data, better estimates? In *Quantitative Genetics in the Wild*.
https://doi.org/10.1093/acprof:oso/9780199674237.003.0002

Pujol, B., Blanchet, S., Charmantier, A., Danchin, E., Facon, B., Marrot, P., Roux, F., Scotti, I.,
Teplitsky, C., Thomson, C. E., & Winney, I. (2018). The Missing Response to Selection in the
Wild. In *Trends in Ecology and Evolution* (Vol. 33, Issue 5).
https://doi.org/10.1016/j.tree.2018.02.007

R Core Team. (2021). R core team (2021). In *R: A language and environment for statistical
computing. R Foundation for Statistical Computing, Vienna, Austria. URL http://www. R-
project. org*.

Radersma, R. (2021). Estimating heritability of social phenotypes from social networks. *Methods in
Ecology and Evolution*, 12(1), 42-53.

Ragab, M., Piles, M., Quintanilla, R., & Sánchez, J. P. (2019). Indirect genetic effect model using
feeding behaviour traits to define the degree of interaction between mates: An
implementation in pigs growth rate. *Animal*, *13*(2).
https://doi.org/10.1017/S1751731118001192

Rees, J. A., & Cranston, K. (2017). Automated assembly of a reference taxonomy for phylogenetic
data synthesis. *Biodiversity Data Journal*, *5*(1). https://doi.org/10.3897/BDJ.5.e12581

Rostellato, R., Sartori, C., Bonfatti, V., Chiarot, G., & Carnier, P. (2013). Comparison between direct
and competitive models to investigate variation of carcass and ham quality traits in heavy pigs.
*Agriculturae Conspectus Scientificus*, *78*(3).

Rostellato, R., Sartori, C., Bonfatti, V., Chiarot, G., & Carnier, P. (2015). Direct and social genetic
effects on body weight at 270 days and carcass and ham quality traits in heavy pigs. *Journal of
Animal Science*, *93*(1). https://doi.org/10.2527/jas.2014-8246

Sánchez-Tójar, A., Nakagawa, S., Sánchez-Fortún, M., Martin, D. A., Ramani, S., Girndt, A., Bókony,
V., Kempenaers, B., Liker, A., Westneat, D. F., Burke, T., & Schroeder, J. (2018). Meta-analysis

challenges a textbook example of status signalling and demonstrates publication bias. *ELife, 7*.
https://doi.org/10.7554/eLife.37385

Santostefano, F., Allegue, H., Garant, D., Bergeron, P., & Réale, D. (2021). Indirect genetic and
environmental effects on behaviors, morphology, and life-history traits in a wild Eastern
chipmunk population. *Evolution, 75*(6). https://doi.org/10.1111/evo.14232

Santostefano, F., Wilson, A. J., Niemelä, P. T., & Dingemanse, N. J. (2017). Indirect genetic effects: A
key component of the genetic architecture of behaviour. *Scientific Reports, 7*(1).
https://doi.org/10.1038/s41598-017-08258-6

Sartori, C., & Mantovani, R. (2012). Effects of inbreeding on fighting ability measured in Aosta
chestnut and Aosta black pied cattle. *Journal of Animal Science, 90*(9).
https://doi.org/10.2527/jas.2011-4010

Scott, J., & Fuller, J. (1965). *Genetics and the Social Behavior of the Dog*. University of Chicago Press.

Shine, R. (1989). Ecological causes for the evolution of sexual dimorphism: a review of the evidence.
*Quarterly Review of Biology, 64*(4). https://doi.org/10.1086/416458

Stirling, D. G., Réale, D., & Roff, D. A. (2002). Selection, structure and the heritability of behaviour.
*Journal of Evolutionary Biology, 15*(2), 277–289.

Sun, Y., Ellen, E. D., Van Der Poel, J. J., Parmentier, H. K., & Bijma, P. (2014). Modelling of feather
pecking behavior in beak-trimmed and non-beak-trimmed crossbred laying hens: Variance
component and trait-based approach. *Poultry Science, 93*(4). https://doi.org/10.3382/ps.2013-
03367

Székely, T., Moore, A. J., & Komdeur, J. (2010). *Social behaviour: genes, ecology and evolution*.
Cambridge University Press.

Teplitsky, C., Mills, J. A., Yarrall, J. W., & Merilä, J. (2010). Indirect genetic effects in a sex-limited
trait: The case of breeding time in red-billed gulls. *Journal of Evolutionary Biology*.
https://doi.org/10.1111/j.1420-9101.2010.01959.x

Trubenová, B., & Hager, R. (2012). Phenotypic and Evolutionary Consequences of Social Behaviours:
Interactions among Individuals Affect Direct Genetic Effects. *PLoS ONE, 7*(11).
https://doi.org/10.1371/journal.pone.0046273

Trubenová, B., Novak, S., & Hager, R. (2015). Indirect genetic effects and the dynamics of social
interactions. *PLoS ONE, 10*(5). https://doi.org/10.1371/journal.pone.0126907

Van Vleck, L. D., Cundiff, L. V., & Kocht, R. M. (2007). Effect of competition on gain in feedlot bulls
from Hereford selection lines. *Journal of Animal Science, 85*(7).
https://doi.org/10.2527/jas.2007-0067

Viechtbauer, W. (2010). Conducting Meta-analysis in R with the metafor package. *Journal of
Statistical Software, 36*(3).

Wade, M. J. (1976). Group selection among laboratory populations of Tribolium. *Proceedings of the
National Academy of Sciences of the United States of America, 73*(12).
https://doi.org/10.1073/pnas.73.12.4604

Wade, M. J. (1977). An Experimental Study of Group Selection. *Evolution, 31*(1).
https://doi.org/10.2307/2407552

Wade, M. J., Bijma, P., Ellen, E. D., & Muir, W. (2010). Group selection and social evolution in
domesticated animals. *Evolutionary Applications*, *3*, 453–465. https://doi.org/10.1111/j.1752-
4571.2010.00147.x

West-Eberhard, M. J. (2003). *Developmental plasticity and evolution*. Oxford University Press.

Westgate, M. J. (2019). revtools: An R package to support article screening for evidence synthesis.
In *Research Synthesis Methods* (Vol. 10, Issue 4). https://doi.org/10.1002/jrsm.1374

Westneat, D. F., & Fox, C. (2010). *Evolutionary behavioral ecology*. Oxford University Press.

Wilson, A. J. (2014). Competition as a source of constraint on life history evolution in natural
populations. *Heredity*, *112*(1), 70–78. https://doi.org/10.1038/hdy.2013.7

Wilson, A. J., Gelin, U., Perron, M., & Réale, D. (2009). Indirect genetic effects and the evolution of
aggression in a vertebrate system Indirect genetic effects and the evolution of aggression in a
vertebrate system. *Proceedings of the Royal Society B: Biological Sciences*, *276*, 533–541.
https://doi.org/10.1098/rspb.2008.1193

Wolf, J. B., Brodie, Cheverud, J. M., Moore, A. J., & Wade, M. J. (1998). Evolutionary consequences
of indirect genetic effects. *Trends in Ecology and Evolution*, *13*(2), 64–69.

Wood, J. L. A., Yates, M. C., & Fraser, D. J. (2016). Are heritability and selection related to
population size in nature? Meta-analysis and conservation implications. In *Evolutionary*
*Applications* (Vol. 9, Issue 5). https://doi.org/10.1111/eva.12375

Wright, A. J. (1985). Selection for improved yield in inter-specific mixtures or intercrops. Theoretical
and applied genetics, 69, 399-407.

Wu, P., Wang, K., Yang, Q., Zhou, J., Chen, D., Liu, Y., Ma, J., Tang, Q., Jin, L., Xiao, W., Lou, P., Jiang,
A., Jiang, Y., Zhu, L., Li, M., Li, X., & Tang, G. (2019). Whole-genome re-sequencing association
study for direct genetic effects and social genetic effects of six growth traits in Large White
pigs. *Scientific Reports*, *9*(1). https://doi.org/10.1038/s41598-019-45919-0

Yang, Y., Noble, D. W., Spake, R., Senior, A. M., Lagisz, M., & Nakagawa, S. (2023). Measuring
biological generality in meta-analysis: a pluralistic approach to heterogeneity quantification
and stratification. *EcoEvoRxiv*. Preprint. https://doi.org/10.32942/X2RG7X

## Author contributions

FS: Conceptualization; Methodology; Investigation; Data Curation; Writing – original draft preparation; Writing – review and editing; Project administration

MM: Conceptualization; Methodology; Validation; Formal analysis; Investigation; Writing – original draft preparation; Writing – review and editing; Visualization

AST: Conceptualization; Methodology; Formal analysis; Investigation; Data Curation; Writing – original draft preparation; Writing – review and editing; Visualization

DF: Conceptualization; Methodology; Validation; Formal analysis; Investigation; Writing – original draft preparation; Writing – review and editing; Visualization

## Acknowledgements

We would like to thank Denis Reale, Clint Kelly, and Pierre-Olivier Montiglio for their feedback in the earlier stages of the project, and Daniel W.A. Noble for statistical advice. We thank the participants of the 2023 WAMBAM conference and Alastair Wilson for useful discussion. We also thank Allen Moore, Joel Pick, and one anonymous reviewer for their constructive feedback, which improved the manuscript. We are grateful to the following authors who provided additional data for the meta-analysis when we contacted them: Alastair Wilson, Anasuya Chakrabarty, Cristina Sartori, Esther Ellen, Chang Han, Jane Reid, João Costa e Silva, Jennifer Morinay, Mark Adams, Miriam Piles, Moha Ragab, Ingrid David, Paolo Carnier, Ryan Germain, Setegn Alemu, Simon Evans, and Céline Teplitsky.

## Funding

FS was supported by the European Union's Horizon 2020 research and innovation programme under the Marie Sklodowska-Curie Individual Fellowship (INTERACTIVE, Grant Agreement Number 101023262). MM was funded by a Marie Skłodowska-Curie Individual Fellowship (PLASTIC TERN, Grant Agreement Number 793550) and an Alexander von Humboldt Research Fellowship for Postdoctoral Researchers. AST was partially funded by the German Research Foundation (DFG: Deutsche Forschungsgemeinschaft) as part of the SFB TRR 212 (NC3)—Project no. 316099922 and 396782608.

## Data and code availability statement

All data and code are available at the following GitHub repository (https://github.com/ASanchez-Tojar/meta-analysis_IGEs). Upon acceptance, all data and code will be provided with a doi via Zenodo.

## Conflict of interest statement

The authors declare no competing interests.

# Tables

935 **Table 1.** Glossary of notation used in the manuscript with associated names and formulae.

| Notation | Name | Formula |
|---|---|---|
| DGEs | Direct genetic effects | - |
| IGEs | Indirect genetic effects | - |
| $V_A$ | Direct additive genetic variance | - |
| $V_{IGE}$ | Indirect genetic effects variance | - |
| $V_P$ | Total phenotypic variance | - |
| $h^2$ | Heritability (narrow sense) | $V_A / V_P$ |
| *social $h^2$* | Social heritability (narrow sense) | $V_{IGE} / V_P$ |
| $I_{DGE}$ | Evolvability | $V_A/\bar{x}^2$ |
| $I_{IGE}$ | Social evolvability | $V_{IGE}/\bar{x}^2$ |
| $COV_{DGE-IGE}$ | DGE-IGE covariance | - |
| $r_{DGE-IGE}$ | DGE-IGE correlation | $COV_{DGE-IGE}/\sqrt{(V_A * V_{IGE})}$ |
| $\tau^2$ | Total heritable variance | $[V_A + Cov_{A-IGE}*2*(n-1) + V_{IGE}*(n-1)^2] / V_P$ |
| - | "Covariance component" of $\tau^2$ | $(Cov_{A-IGE}*2*[n-1])$ |
| - | "IGE variance component" of $\tau^2$ | $(V_{IGE}*[n-1]^2)$ |
| - | Response to individual selection among unrelated individuals | $1*[V_A + (n-1) Cov_{A-IGE}]$ |
| *n* | Group size | - |

 **Table 2.** List of studies (n = 47) from which data was extracted and included in our meta-analysis on indirect genetic effects.

| Study ID | Year | Journal | Number of effect sizes (k) | Species | Reference |
|---|---|---|---|---|---|
| IGE0057 | 2015 | Genetics Selection Evolution | 1 | *Penaeus vannamei* | (Luan et al., 2015) |
| IGE0144 | 2013 | Agriculturae Conspectus Scientificus | 3 | *Sus scrofa* | (Rostellato et al., 2013) |
| IGE0195 | 2012 | Journal Of Animal Science | 3 | *Sus scrofa* | (Duijvesteijn et al., 2012)* |
| IGE0196 | 2016 | Journal Of Animal Ecology | 1 | *Melospiza melodia* | (Germain et al., 2016) |
| IGE0197 | 2016 | Evolution | 1 | *Tribolium castaneum* | (Ellen et al., 2016) |
| IGE0199 | 2019 | Journal Of Evolutionary Biology | 3 | *Chorthippus biguttulus* | (Chakrabarty et al., 2019) |
| IGE0202 | 2015 | Journal Of Animal Science | 10 | *Sus scrofa* | (Rostellato et al., 2015) |
| IGE0203 | 2014 | Genetics Selection Evolution | 19 | *Gadus morhua* | (Nielsen et al., 2014)* |
| IGE0242 | 2007 | Journal Of Animal Science | 8 | *Bos taurus* | (Van Vleck et al., 2007)* |
| IGE0243 | 2010 | Journal Of Animal Science | 3 | *Sus scrofa* | (Hsu et al., 2010)* |
| IGE0253 | 2012 | Journal Of Animal Science | 1 | *Bos taurus* | (Sartori & Mantovani, 2012) |
| IGE0291 | 2008 | Journal Of Animal Science | 1 | *Sus scrofa* | (Chen et al., 2008)* |
| IGE0292 | 2016 | Journal Of Animal Breeding And Genetics | 3 | *Neovison vison* | (Alemu et al., 2016) |
| IGE0294 | 2005 | Journal Of Animal Science | 1 | *Sus scrofa* | (Arango et al., 2005) |
| IGE0347 | 1997 | British Poultry Science | 6 | *Gallus gallus* | (Kjaer & Sørensen, 1997) |
| IGE0380 | 2016 | Aquaculture | 1 | *Oreochromis niloticus* | (Khaw et al., 2016) |
| IGE0385 | 2019 | Asian-Australasian Journal Of Animal Sciences | 1 | *Sus scrofa* | (Hong et al., 2019) |
| IGE0391 | 2013 | Journal Of Animal Science | 3 | *Sus scrofa* | (Bergsma et al., 2013)* |
| IGE0406 | 2005 | Proceedings Of The National Academy Of Sciences Of The United States Of America | 16 | *Drosophila serrata* | (Petfield et al., 2005) |

| IGE0468 | 2019 | Behavior Genetics | 24 | *Mus musculus* | (Dewan et al., 2019) |
|---------|------|-------------------|-----|----------------|----------------------|
| IGE0483 | 2018 | Evolution | 6 | *Gryllus bimaculatus* | (Han et al., 2018) |
| IGE0493 | 2019 | Ecology Letters | 1 | *Tamiasciurus hudsonicus* | (Fisher, Haines, et al., 2019) |
| IGE0494 | 2019 | Animal | 1 | *Sus scrofa* | (Ragab et al., 2019) |
| IGE0501 | 2009 | Proceedings Of The Royal Society B-Biological Sciences | 5 | *Peromyscus maniculatus sonoriensis* | (Wilson et al., 2009)* |
| IGE0505 | 2014 | Genetics Selection Evolution | 3 | *Neovison vison* | (Alemu et al., 2014)* |
| IGE0507 | 2012 | Genetics | 4 | *Gallus gallus* | (Peeters et al., 2012)* |
| IGE0509 | 2010 | Journal Of Evolutionary Biology | 1 | *Chroicocephalus scopulinus* | (Teplitsky et al., 2010) |
| IGE0515 | 2017 | Scientific Reports | 1 | *Gryllus bimaculatus* | (Santostefano et al., 2017) |
| IGE0537 | 2017 | Genetics Selection Evolution | 2 | *Oryctolagus cuniculus* | (Piles et al., 2017) |
| IGE0572 | 2018 | Genetics Selection Evolution | 5 | *Oryctolagus cuniculus* | (David et al., 2018) |
| IGE0598 | 2010 | Journal Of Animal Science | 1 | *Sus scrofa* | (Bouwman et al., 2010)* |
| IGE0640 | 2014 | Poultry Science | 2 | *Gallus gallus* | (Sun et al., 2014) |
| IGE0654 | 2007 | Genetics | 1 | *Gallus gallus* | (Bijma et al., 2007) |
| IGE0657* | 2013 | Evolution | 1 | *Coturnix japonica* | (Muir et al., 2013)* |
| IGE0746 | 2014 | Genetics Selection Evolution | 8 | *Gallus gallus* | (Brinker et al., 2014)* |
| IGE0752 | 2015 | Genetics Selection Evolution | 4 | *Gallus gallus* | (Brinker et al., 2015) |
| IGE0786 | 2017 | Animal Reproduction | 1 | *Sus scrofa* | (Hong et al., 2017) |
| IGE0824 | 2014 | Evolution | 2 | *Drosophila melanogaster* | (Edward et al., 2014) |
| IGE0852 | 2019 | Journal Of Evolutionary Biology | 1 | *Tamiasciurus hudsonicus* | (Fisher, Wilson, et al., 2019) |
| IGE0857 | 2015 | Proceedings Of The Royal Society B: Biological Sciences | 1 | *Aegithalos caudatus* | (Adams et al., 2015) |

| IGE0858 | 2018 | Genetics Selection Evolution | 2 | *Sus scrofa* | (Nielsen et al., 2018) |
|---------|------|------------------------------|---|--------------|------------------------|
| IGE0889* | 2008 | Poultry Science | 3 | *Gallus gallus* | (Ellen et al., 2008)* |
| IGE0920 | 2008 | Genetics | 5 | *Sus scrofa* | (Bergsma et al., 2008) |
| IGE0927 | 2017 | Heredity | 1 | *Sus scrofa* | (Canario et al., 2017) |
| IGE1023 | 2013 | Genetics Selection Evolution | 2 | *Gallus gallus* | (Peeters et al., 2013) |
| IGE1038 | 2019 | Scientific Reports | 6 | *Sus scrofa* | (Wu et al., 2019) |
| IGE1050 | 2008 | Evolution | 1 | *Larus canus* | (Brommer & Rattiste, 2008) |

939      * References included in the qualitative review performed by Ellen et al. (2014: Table 4 in that review).

## Figures

**Figure 1.** Indirect genetic effects occur when the genotype of one individual influences the expression of a phenotype in a conspecific, mediated by social interactions.

**Figure 2**. Social heritability (*social h$^2$*) from a phylogenetic multilevel meta-analysis with 40 studies

and 146 effect sizes (Aim 1). Orchard plot showing the back-transformed meta-analytic mean as *r*,

95% confidence intervals (thick whisker), 95% prediction intervals (thin whisker) and individual effect

sizes (*r*) scaled by their precision (semitransparent circles). *k* corresponds to the number of effect

sizes, the number of studies is shown in brackets. Note that uncertainty measures (i.e., 95% CI and

95% PI) are generated assuming a normal distribution around the meta-analytic mean, explaining

why some intervals may overlap zero substantially despite the data analyzed being bounded

between 0 and 1.

**Figure 3.** Social heritability (*social $h^2$*) meta-analytic estimates from phylogenetic multilevel uni-moderator meta-regressions testing the following

moderators (Aim 2): (a) Trait category, (b) Age, (c) Sex, (d) Population type. Orchard plot showing the back-transformed meta-analytic mean as *r*, 95%

confidence intervals (thick whisker), 95% prediction intervals (thin whisker) and individual effect sizes (*r*) scaled by their precision (semitransparent circles).

*k* corresponds to the number of effect sizes, the number of studies is shown in brackets. Note that uncertainty measures (i.e., 95% CI and 95% PI) are

generated assuming a normal distribution around the meta-analytic mean, explaining why some intervals may overlap zero substantially despite the data

analyzed being bounded between 0 and 1.

# Social h$^2$

**Figure 4.** IGEs and DGEs meta-analytic estimates from phylogenetic multilevel uni-moderator meta-regressions (Aim 3) comparing: (a) $V_A$ vs. $V_{IGE}$, (b)

narrow-sense $h^2$ vs. *social $h^2$*, and (c) $I_{DGE}$ vs. $I_{IGE}$ (evolvability). Orchard plot showing the back-transformed meta-analytic mean as $r$, 95% confidence intervals

(thick whisker), 95% prediction intervals (thin whisker) and individual effect sizes ($r$) scaled by their precision (semitransparent circles). $k$ corresponds to the

number of effect sizes, the number of studies is shown in brackets. Note that for (b) and (c) uncertainty measures (i.e., 95% CI and 95% PI) are generated

assuming a normal distribution around the meta-analytic mean, explaining why some intervals may overlap zero substantially despite the data analyzed

being bounded between 0 and 1.

**Figure 5.** Meta-analytic estimates for Aim 4: (a) phylogenetic multilevel meta-regression comparing narrow-sense heritability ($h^2$) to total heritable variance

($\tau^2$); (b) phylogenetic multilevel meta-analysis of $r_{DGE\text{-}IGE}$. Orchard plot showing the original (a) and back-transformed meta-analytic mean as $r$, 95%

confidence intervals (thick whisker), 95% prediction intervals (thin whisker) and individual effect sizes ($r$) scaled by their precision (semitransparent circles).

$k$ corresponds to the number of effect sizes, the number of studies is shown in brackets. Note that for (a) uncertainty measures (i.e., 95% CI and 95% PI) are

generated assuming a normal distribution around the meta-analytic mean, explaining why some intervals may overlap zero substantially despite the data

analyzed being bounded between 0 and 1.