Kinetic properties of optogenetic DNA editing by LiCre-*loxP*

Alice Dufour[1,*], Hélène Duplus-Bottin[1,*], Thomas Boukéké-Lesplulier[1], Eliane Casassa[1], Gérard Triqueneaux[1], Camille Darthenay-Kiennemann[1], Agnès Dumont[1], Catherine Moali[2], Franck Vittoz[3], Daniel Jost[1,#] and Gaël Yvert[1,#,§]

1) Laboratory of Biology and Modeling of the Cell, Ecole Normale Superieure de Lyon, CNRS, UMR 5239, Inserm, U1293, Universite Claude Bernard Lyon 1, 46 allee d'Italie F-69364 Lyon, France

2) Tissue Biology and Therapeutic Engineering Laboratory, Universite Claude Bernard Lyon 1, CNRS, UMR 5305, 7 Passage du Vercors F-69367 Lyon, France

3) Laboratory of Physics, Ecole Normale Superieure de Lyon, CNRS, UMR 5672, Universite Claude Bernard Lyon 1, 46 allee d'Italie F-69364 Lyon, France.

*) equal contribution

#) Corresponding authors. Experiments: gael.yvert@ens-lyon.fr ; modeling: daniel.jost@ens-lyon.fr

§) Lead contact: gael.yvert@ens-lyon.fr

ABSTRACT

Previously, we developed an optogenetic tool made of a single chimeric protein called LiCre

that enables the edition of specific changes in the genome of live cells with blue light via DNA

recombination between *loxP* sites (Duplus-Bottin et al., 2021). Here, we used *in vitro* and *in vivo*

experiments combined with kinetic modeling to provide a deeper characterization of the photo-

activated LiCre-*loxP* recombination reaction. We find that LiCre binds DNA with high affinity in

absence of light stimulus, that this binding is cooperative although not as much as for the Cre

recombinase from which LiCre was derived and that increasing temperature from 20°C to 37°C

gradually increased LiCre efficiency. The recombination kinetics in live cells can be explained by a

model where photo-activation of two or more DNA-bound LiCre units (happening in seconds) can

produce (in several minutes) a functional recombination synapse. Our conclusions provide helpful

guidelines to induce specific genetic changes in live cells using light.

INTRODUCTION

Experimental biologists usually make discoveries by perturbing living organisms and
observing the consequences of these perturbations. Precision in these manipulations is often crucial
because biological systems are spatially complex and highly dynamic. Optogenetics offers the
possibility to alter cellular activities with light and provides unprecedented spatio-temporal
resolution. For this reason, a wide set of light-inducible biotechnologies have been developed in the
past years. These tools include photo-activatable ion channels that trigger neuronal activities when
illuminated (see (Altahini et al., 2024) for a review on neuroscience applications), photo-controlled
kinases activating signaling pathways, as well as systems controlling protein localization, protein-
protein interactions, gene expression or DNA editing (see (Armbruster et al., 2024) for a review on
molecular tools). Many of these systems are genetically encoded, thereby alleviating the need for
chemicals. They are all based on conformational changes of a macromolecule leading to its
activation or inactivation in response to light. Yet, each tool has its own specificity in terms of light
sensitivity, efficiency and response dynamics; optimal usage of a given tool therefore requires
quantitative knowledge of its biochemical properties.

We recently developed an optogenetic site-specific DNA recombinase called LiCre (Duplus-
Bottin et al., 2021). As for the Cre recombinase from which it derives, LiCre recognizes 34-bp DNA
sequences called *loxP* and catalyzes DNA recombination between two such sites. This allows
experimentalists to edit the genome of live cells in real time with blue light. For example, if a gene
is flanked by two *loxP* sites in LiCre-expressing cells, shining light on these cells can delete the
gene. Compared to other photo-activatable recombinases which are based on two protein halves
(splitCre) that complement in response to light (Hochrein et al., 2018; Kawano et al., 2016;
Kuwasaki et al., 2022; Taslimi et al., 2016), LiCre is unique as it consists in a single chimeric
protein comprising the Light-Oxygen Voltage domain from *Avena sativa* (asLOV2) fused to a
mutant version of Cre. It was designed such that the unfolding of a critical α-helix in response to
light enables the assembly of a functional recombination synapse (Duplus-Bottin et al., 2021).
Although LiCre outperformed split-based systems in terms of speed and efficiency, its mode of
action, represented in Fig. 1a, could only be speculated from the abundant knowledge previously
acquired on the Cre-*loxP* synapse. Formation of this complex involves the cooperative binding of
two Cre units on each *loxP* site and the subsequent assembly of the four Cre proteins complexed

with their DNA substrates (Duyne, 2015). The resulting intasome is stabilized by interactions between adjacent protomers: on one side of the DNA substrate, the four C-terminal domains are locked together in a cyclic manner, each unit hosting the αN helix of its neighbor in a nest; on the other side of DNA (depicted in Fig. 1a for LiCre), N-terminal domains also form a cycle of binding interfaces but in the reverse orientation (Guo et al., 1997). This way, each protomer is stabilized by its two neighbors. For LiCre, the strength of interactions between C-terminal domains was reduced by mutating two critical residues of the αN helix. In addition, the association of N-terminal domains was also perturbed: the αA helix interacting with the neighboring unit was truncated and fused to asLOV2, generating a chimeric α-helix comprising Jα from asLOV2 and the remaining part of αA from Cre. Given the known properties of the asLOV2 photo-switch (Zayner et al., 2012), LiCre is presumed to be inactive in the dark because i) docking with the adjacent N-terminal domain is probably hindered by the sterically-proximal asLOV2 domain, and ii) the chimeric helix is likely to be sequestered by asLOV2 and unavailable for interaction with other LiCre units. The unfolding of Jα in response to light may have two effects: clearance of the interface by the release of asLOV2, and the possibility that αA refolds and makes contacts with the other unit. However, the exact mechanism by which a functional LiCre-*loxP* complex assembles, and in particular the critical step that is unlocked by light, remains to be identified.

We report here a deeper analysis of LiCre that combined *in vitro* and *in vivo* experiments with kinetic modeling. We found that LiCre units bind to DNA with high affinity even if they are not photo-activated, that they do so in a cooperative manner although this cooperativity is weaker than for Cre, that temperature greatly affects LiCre efficiency, and that the kinetics of recombination in live cells can be explained by a model where photo-activation of two or more DNA-bound LiCre units happens in seconds and enables the formation, in several minutes, of a functional recombination synapse. This deeper understanding of the LiCre-*loxP* system provides helpful guidelines to conduct genetic perturbations of live cells with high spatio-temporal precision using light.

RESULTS

LiCre binds its target DNA in absence of photo-stimulation

LiCre photo-activation is based on a conformational change that enables protein-protein
interactions assembling the recombinogenic intasome (Duplus-Bottin et al., 2021). We previously
proposed two possible models for this activation. In the first model, LiCre monomers are unable to
bind DNA *loxP* sites in their 'dark' – inactive - state and the photo-induced conformational change
restores this ability. In the alternative model, LiCre binds to *loxP* sites even in the dark, and the
conformational change of photo-stimulated DNA-bound LiCre molecules enables the assembly of
the recombination synapse. To distinguish between these two models, we evaluated the ability of
LiCre to bind DNA in conditions where it was not photo-activated.

We first tested this DNA-binding ability *in vivo* by using chromatin immuno-precipitation
(ChIP). We constructed yeast strains harboring a single *loxP* site in the genome and expressing
LiCre or Cre tagged with repeated V5 epitopes (Fig. 1b). We cultured cells overnight and then either
let them in dark conditions or, for LiCre-V5 samples, illuminated them in conditions known to
trigger LiCre activity (Duplus-Bottin et al., 2021). Following illumination, we processed half of the
cells for fixation and ChIP and the other half for Western blot. We quantified ChIP by qPCR at three
positions near the *loxP* site (Fig. 1c) and, for normalization, at three unrelated positions on the
genome. As shown on Fig. 1d, we observed a strong ChIP signal for all three probes of the *loxP*
locus for Cre, consistent with the known high-affinity of Cre for its *loxP* target (Duyne, 2015).
Regarding LiCre, we observed a much weaker ChIP signal, which reached statistical significance
only for the probe located very close to *loxP* (probe P2 on Fig. 1c-d). When verifying the number of
V5 epitopes by Sanger sequencing, we found that they had been amplified for Cre-V5 (12 copies)
but not for LiCre-V5 (9 copies as expected). The fact that LiCre-V5 possessed fewer epitopes
certainly contributed to the weaker ChIP signal. However, given the magnitude of the difference, it
is also possible that LiCre binds chromatin with a weaker affinity than Cre. Importantly,
illumination prior to chromatin immuno-precipitation had clearly no effect on LiCre ChIP signals
(Fig. 1d). Finally, on Western-blots, the immuno-detection of LiCre-V5 on cell extracts was not
weaker than the detection of Cre-V5 (Fig. 1e-f). We conclude that, *in vivo*, LiCre binds *loxP*-
containing chromatin in the dark, that this binding has a similar affinity in light conditions, and that
this affinity is possibly lower than Cre affinity.

Figure 1. Analysis of LiCre:DNA binding *in vivo*. **a**) Hypothetical principle of LiCre photo-activation. The chimeric α-helix is presented as a rectangle; in blue when folded as the Jα helix of LOV2, in green when folded as the αA helix of Cre. Modified from (Duplus-Bottin et al., 2021). **b**) Chromatin immunoprecipitation (ChIP) experimental setup. **c**) Position of qPCR probes relative to *loxP*. **d**) ChIP-qPCR quantifications. Every dot corresponds to one biological replicate (independent culture inoculated from an independent transformant of the Cre-V5 or LiCre-V5 expression plasmid). Values correspond to enrichment by ChIP (relative to its input), normalized by the signals obtained on three unrelated loci of the genome (see Methods). *P* values: Welch *t*-test. **e**) Western blot analysis. Proteins were extracted from the same population of cells as for ChIP. **f**) Quantification of LiCre, Cre and GAPDH immunoblot shown in e.

123   Given the weakness of the ChIP signal supporting LiCre:DNA binding, we decided to

124 analyse this affinity *in vitro* by surface plasmon resonance (SPR). This method offers high

125 sensitivity to monitor macromolecular associations and dissociations; it was previously used to

126 characterize the binding affinity of Cre to *loxP* (Rüfer et al., 2002). We produced and purified

127 recombinant Cre and LiCre proteins from *E. coli* (Supplementary Fig. S1) and injected these

128 proteins into microfluidic chips containing immobilized biotinylated DNA substrates (see Methods).

129 We used DNA substrates containing either a full *loxP* site that can associate with two units of Cre or

130 LiCre, or a half site allowing the binding of only one unit. We also used a randomized DNA

131 sequence to quantify non-specific binding. We did not apply any blue light illumination to the

132 samples, neither before injection of the protein solution in the chip nor during acquisitions.

133   As shown in Fig. 2a, Cre association with the full *loxP* site was very rapid and its

134 dissociation was extremely slow. This observation is in perfect agreement with previous

135 experiments (Rüfer et al., 2002) based on the same DNA substrates as here. For LiCre, association

136 with the full loxP site was slightly slower than for Cre but still very fast while the dissociation was

137 slightly faster (Fig. 2b). This demonstrates that LiCre is able to bind its DNA target site even if it is

138 not photo-activated, which may explain why the blue-light treatment did not increase the detection

139 of LiCre chromatin-binding *in vivo* (Fig. 1). However, quantitatively, the high affinity of LiCre for

140 *loxP* DNA *in vitro* contrasted with the weak ChIP enrichment observed *in vivo*.

141   The profiles obtained with half *loxP* sites are also very informative. In accordance with

142 (Rüfer et al., 2002), Cre dissociation from the half site is much faster than its dissociation from the

143 full site (Fig. 2c). Similarly, for LiCre, dissociation from the half site is also faster, but the

144 difference is not as pronounced as for Cre (Fig. 2d). This suggests cooperativity between the two

145 Cre or LiCre units when binding to the full *loxP* site, and that cooperativity of LiCre in the dark

146 may be lower than for Cre.

Figure 2. Analysis of LiCre:DNA interaction *in vitro* by surface plasmon resonance (SPR). Net sensorgrams showing Cre (**a,c**) or LiCre (**b,d**) interaction with full (**a,b**) or half (**c,d**) *loxP* sites. At time zero, recombinant proteins were injected at the indicated concentrations over a BIACORE sensor chip containing immobilized *loxP* sites (a,b), *loxP* half sites (c,d) or unrelated DNA as negative control blank. After 90 seconds, the sensor chip was washed by injecting running buffer lacking the protein (dissociation step). Net sensorgrams were generated from the raw data by subtracting the negative blank control to correct injection-related fluctuations.

     LiCre binding to DNA is strong and cooperative

To rationalize our in vitro measurements in terms of binding affinities, we developed a

kinetic model of the association and dissociation of LiCre and Cre molecules to DNA, accounting

for the SPR protocol (see Methods). Briefly, based on (Rüfer et al., 2002), the binding of one

molecule to a half-site is described as a simple one-step assembly reaction; while the binding of two

molecules to a full site is assumed to happen sequentially: binding of the first molecule is assumed

to follow the same kinetics as for a half-site while the second binding event may have different

kinetics due to cooperativity mediated by the presence of the already-bound molecule (Fig. 3a). In

addition to the association (k_1, k_2) and dissociation (k_{-1}, k_{-2}) rates of each step, the model comprises

the following parameters that enable its direct confrontation to experimental acquisitions: a

coefficient of mass transport in the biosensor chip (k_t), a linear coefficient (α) linking the Response

Unit (RU) of the SPR experiments to the amount of bound proteins, and maximal RU values

($RUmax_h$ and $RUmax_f$) when all of the half or full *loxP* sites are fully occupied, respectively. We

fitted these parameters by least-squared minimization between predicted and experimental

sensorgrams (Fig. 3b, Figure 3 - figure supplement 1, see Methods). To study the identifiability of

model parameters, we ran the procedure 1,000 times starting from different initial parameter values

picked randomly within acceptable ranges. Among the resulting 1,000 trials, 541 and 859 led to

high-quality fits for Cre and LiCre, respectively (Figure 3 - figure supplement 2). The

corresponding sets of parameters are shown in Fig. 3c-e. Fitted values of $RUmax_h$ and $RUmax_f$ had

very low standard deviations and were therefore very well defined, both for Cre and LiCre (Fig. 3e).

Parameters k_t and α were each well estimated for LiCre (Fig. 3d). For Cre, their individual

estimation was less precise but, as anticipated by their definition, they co-varied and their product

was robustly estimated (Fig 3c,e). Similarly, association and dissociation rates were not individually

identifiable, but their ratio ($K_{D1} = k_{-1}/k_1$ and $K_{D2} = k_{-2}/k_2$, *aka* dissociation constants) were (Fig. 3c-

e). For Cre, these values ($K_{D1} = 1.15$ nM and $K_{D2} = 0.001$ nM) highlight a very strong cooperativity

($K_{D2} << K_{D1}$) and are consistent with those previously reported (Ringrose et al., 1998; Rüfer et al.,

2002) (see Discussion). Importantly, the cooperativity for LiCre is also strong (cooperative moment

$K_{D1}/K_{D2} = 81$) but much weaker than for Cre ($K_{D1}/K_{D2} = 1330$). This is consistent with the way LiCre

was derived from Cre, which involved the destabilization of protein:protein binding both by

introducing point mutations in the αN helix required to lock C-terminal domains together, and by

truncating and hindering the αA helix engaged in the interaction between N-terminal domains

(Duplus-Bottin et al., 2021) (see Discussion).

Figure 3. Kinetic modeling of DNA binding. a) Model for association of Cre or LiCre molecules (*C*) to DNA *loxP* site (*L*). L_0, L_1 and L_2 denote DNA sites occupied by zero, one and two molecules, respectively. **b**) Experimental (dots) versus model-predicted (lines) sensorgrams of LiCre binding to a full *loxP* site, using parameter values $\alpha = 1.7$, $k_t = 0.1$, $RUmax_h = 53.2$, $RUmax_f = 38.1$, $K_{D1} = 0.5$ nM, $K_{D2} = 0.00615$ nM. **c**) Best estimates of model parameters obtained from 541 optimizations for Cre:DNA binding (see Methods). Every dot corresponds to one optimization. Histograms show marginal distributions of x and y values. Red line, LOWESS local regression. **d**) Same representation as in (c), showing results from 859 optimizations for LiCre:DNA binding. **e**) Table of estimated parameter values (mean +/- sd).

Experimental quantification of LiCre-loxP recombination following various regimes of photo-stimulation

197   To quantify recombination in live cells, we used a yeast strain harboring a previously-
198 described quantitative reporter (Duplus-Bottin et al., 2021). This system is based on the LiCre-
199 mediated excision of a transcriptional terminator preventing the expression of a fluorescent protein
200 (Fig. 4a). This excision occurs by recombination between *loxP* sites located 1.3-Kb from one
201 another. In a typical experiment, a population of yeast cells is illuminated with chosen doses and
202 dynamics of blue light; cells are then re-suspended in fresh medium in the dark to let them express
203 the fluorescent protein if recombination occurred; the population is finally analyzed by flow
204 cytometry to determine the fraction of fluorescent cells. This fraction corresponds to the proportion
205 of stimulated cells that achieved recombination, therefore providing a clear estimate of
206 recombination efficiency. To avoid any potential bias associated with the chromosomal locus where
207 recombination takes place, every cell carried two versions of the reporter, one on chromosome IV
208 coding for mCherry (red fluorescence), and one on chromosome II coding for GFP (green
209 fluorescence). LiCre expression was conferred by a plasmid introduced in the resulting strain.

210   Figure 4b shows how recombination efficiency increased with the duration of a defined
211 pulsed stimulation. In agreement with previous observations (Duplus-Bottin et al., 2021), ten
212 minutes were enough to induce recombination in a significant proportion (~10%) of cells. In
213 general, we counted slightly more red cells than green cells. This could indicate that the reporter
214 locus on chromosome IV is more favorable to recombination than its counterpart on chromosome
215 II. Alternatively, the difference could result from different measurement accuracy between the green
216 and the red channel. Importantly, we never observed recombination in the entire cell population.
217 After prolonged (>3 hours) photo-stimulation, the proportion of positive cells reached a plateau of
218 about 85%. Most of these cells displayed both red and green fluorescence, leaving a remaining
219 ~15% cells negative for both reporters. This suggested that the upper limit of recombination
220 efficiency was caused by a cell-specific factor rather than a locus-specific factor. We hypothesized
221 that this upper limit could be linked to the occasional loss of the LiCre-encoding plasmid.

222   To test this possibility, we repeated the experiment but instead of illuminating the population
223 of cells, we plated diluted samples of it on both selective and non-selective solid media
224 (Supplementary Table S1). This revealed that, in our experimental conditions, the proportion of
225 cells having lost the LiCre plasmid at the time of illumination was between 10% and 30%. This

observation is remarkably consistent with the presence of ~15% of recombination-negative cells in our optogenetic assay with prolonged photostimulation. We therefore propose that, for cells having kept the plasmid, LiCre-mediated recombination efficiency after prolonged illumination is close to 100%.

We also observed substantial variability between experiments. For example, a pulsed stimulation of 60 minutes generated nearly 50% of positive cells in the experiment of Fig. 4b, performed in July 2019, whereas the same stimulation generated only ~30% of positive cells in another experiment (Fig. 3g of (Duplus-Bottin et al., 2021), performed in November 2020). For these two experiments, illumination was conducted at room temperature, which was not stable over the year: it exceeded 37°C in July 2019 and was below 25°C in November 2020. We therefore directly tested the effect of temperature on recombination efficiency by illuminating cells in thermo-controlled conditions. We observed that the same pulsed stimulation generated between ~15% and ~45% positive cells when temperature varied from 20°C to 37°C, respectively (Fig. 4c), higher temperature leading to more efficient recombination, thus explaining most of the variability mentioned above. We therefore conducted subsequent experiments under strict temperature-controlled conditions (30°C).

To further explore the dynamic properties of the LiCre-*loxP* reaction *in vivo*, we applied pulsed photo-stimulation at various periods, with two different duty cycles (fraction of the period when light is on). The list of experiments is summarized in Supplementary Table S2 and only a subset of the data is shown in Fig. 4d. For a fixed total cumulative time of light exposure, recombination efficiency declined when the period increased. For example, applying a total illumination of 6 minutes (dark red in Fig. 4d) led to an efficiency above 40% when the time between consecutive pulses of light was short (4.5 seconds) but below 25% when this period was longer (54 seconds). This may reflect the existence of a lag time, in the order of tens of seconds, before LiCre returns to its dark, inactive, state in absence of light.

In order to quantify the photo-sensitivity of the reaction, we illuminated yeast cells at various light intensities. We chose a regime where light pulses were long enough to ensure efficient activation (2 seconds) and frequent enough to prevent the system from reverting to its inactive state (every 7 seconds). As expected, we observed an overall increase of recombination efficiency with light intensity until maximal activation was reached at $12mW/cm^2$ (Fig. 4e). Intriguingly, the fraction of positive cells was slightly lower when we applied maximal intensity (35.6mW/cm2). This observation was reproduced when we repeated the experiment (Figure 4 - figure supplement 1,

and Dataset 11 listed in Supplementary Table S2), possibly pointing to photo-toxic effects under

prolonged or intense illumination.

Overall, the experimental data generated here describe the efficiency of LiCre following

illumination conditions varying in duration, in dynamics or in light intensity. It therefore provides a

useful resource to evaluate the kinetics of the recombination reaction.

Figure 4. Experimental quantification of *in vivo* LiCre-*loxP* recombination following
various photo-stimulation conditions. a) Reporter system. Yeast cells harbour two reporter
constructs integrated in their genome and express LiCre from a plasmid. A transcriptional
terminator (T_{ADH1}), flanked by *loxP* sites, prevents expression of mCherry or GFP fluorescent
proteins. LiCre photo-stimulation triggers recombination, which removes T_{ADH1} and releases
expression. The rate of recombination events can be estimated by counting fluorescent cells in a
flow cytometer. **b)** Recombination rate following pulsed stimulation of various duration. Cells were
subjected to pulses of blue light (450 nm, period of 10s, duty cycle of 5%, intensity of 35mW/cm2)
for the indicated amount of time, then transferred to fresh medium for 4 hours to enable expression
of the fluorescent proteins. Green, red and non-fluorescent cells were then counted by flow
cytometry. **c)** Same as in *b* but for a single duration and various temperatures. **d)** Same as in *b* but
with various periods and a duty cycle of 10%. Intensity of 35 mW/cm2. **e)** Same as in *b* and *d* but
with various light intensities, a period of 7 s, a duty cycle of 29%, applied for 60 minutes. In *b-e*,
replicates (dots) correspond to independent cultures, each inoculated with an independent
transformant of the LiCre plasmid. Lines connect mean values.

    Kinetic modeling of LiCre-loxP recombination

To describe the light-induced recombination catalyzed by LiCre, we built a mathematical
model based on three steps (Fig. 5a, see Methods).

First, association of LiCre to DNA occurs in the dark and is at equilibrium. Using the
dissociation constants inferred from the SPR experiments (see above), we plotted the probabilities
of occupancy of the four LiCre binding sites located on two *loxP* sequences assuming independent
binding at the two *loxP* loci (Fig. 5b). At LiCre concentrations above 1nM, over 97% of cells have
both their *loxP* sites fully occupied (four bound LiCre units in total).

Second, during stimulation by light, bound LiCre units are activated at a rate k_{ON}. In dark
condition, they deactivate at a rate k_{OFF}. Given these rates and the dynamics of illumination, we can
compute the probability $P_{act}(t)$ that a given LiCre unit is active at time t (Fig. 5c, top). Since LiCre is
activated by the asLOV2 switch, we chose k_{ON} as previously reported for the LOV2 domain
(Renicke et al., 2013). This rate increases linearly with the incoming flux of photons and is in the
order of second^{-1} (see Methods).

Third, if the two *loxP* sites are both fully-occupied and if at least x among the four LiCre
units are active, with x ranging from 1 to 4, recombination can occur at a constant rate R_0 that
encompasses the loop probability between the sites, the synapse formation and the actual catalytic
reaction of recombination. All this leads to an overall time-dependent recombination rate $R(t)$ that is
proportional to R_0, to the probability of having 4 LiCre bound to the *loxP* sites and to the probability
to have at least x light-activated LiCre molecules, which is a function of $P_{act}(t)$ (Fig.5a).

The well-known Cre-*loxP* reaction is reversible and this is presumably also the case for the
LiCre-*loxP* reaction. As described above, we concluded from our experimental data that all cells
having kept the LiCre plasmid would eventually achieve recombination after prolonged photo-
activation. Thus, at equilibrium, the backward reaction (re-insertion of the excised circular DNA) is
unlikely and out-competed by the forward reaction and we therefore neglected it in our model.

A reaction simulated with this model is shown in Fig. 5c. In this example, periodic pulses of
light were applied and the fraction of activated LiCre units reached a periodic steady-state after only
60 seconds, oscillating around ~80% (Fig. 5c, top) with the light stimulus. The overall time-
dependent reaction rate $R(t)$ also fluctuates, the speed of recombination being highly affected by the
number (x) of active LiCre units needed to form a functional complex (Fig. 5c, center). After

plotting the cumulative fraction of recombined molecules over time (Fig. 5c bottom) we concluded

that this three-steps model provides realistic outcomes that can be compared to experimental data.

Figure 5. Kinetic model of LiCre-*loxP* recombination. a) Model description. Purple and
cyan beads represent inactive and active LiCre units, respectively. Occurrence of the reaction
assumes the binding of four LiCre units (which happens with probability $P_{(2,2)}$), the photoactivation
of at least *x* of them, with $x \in \{1,2,3,4\}$, and a maximal reaction rate R_0 once the complex can form.
Reaction rate *R(t)* summarizes the triggering of the reaction by light. See text and Methods for
details. **b)** Effect of [LiCre] concentration on the probability of occupancy at two *loxP* sites. (*n,m*)
indicates *n* and *m* LiCre molecules bound to the first and second *loxP* site, respectively. **c)**
Simulations under periodic pulses of light. Light was ON during the vertical blue stripes, each pulse
occurred every 10s and lasted 0.5s. [LiCre] = 1nM, k_{ON} = 1s^{-1}, k_{OFF} = 0.007s^{-1}, R_0 = 10^{-3}s^{-1}. From up
to bottom: probability that one LiCre unit is active (P_{act}); reaction rate at each possible value of *x*
(required number of activated units); cumulative fraction of recombined molecules over time at
each possible value of *x*. **d-g)** Best model prediction using *x* = 2 and parameters simultaneously
fitted to all experimental data. Lines, model simulations. Dots and bars, experimental data (mean ±
2 standard deviations). Each panel corresponds to one consistent series of acquisitions, where the
fraction of mCherry-positive yeast cells which underwent recombination was measured by flow-
cytometry (datasets 1-10 listed in Supplementary Tables S2). Value k_{OFF} inferred from all data: 50 s$^-$
^{1}. **d)** Effect of the duration of illumination (same data as in Fig. 4b). Inferred R_0 value: 3.5.10^{-4} s^{-1}. **e)**
Effect of illumination dynamics and intensity (duration of 1 hour), same data as in Fig. 3f-g , from
ref. (Duplus-Bottin et al., 2021). Inferred R_0 value: 0.000125 s^{-1}. **f)** Effect of illumination dynamics
and duration. Inferred R_0 value: 0.00014 s^{-1}. **g)** Effect of illumination intensity (same data as in Fig.
4e). One unit = 35.6 mW/cm^2. Inferred R_0 value: 0.00022 s^{-1}.

    Model fitting to experimental data suggests that photo-activation of two LiCre protomers is

    sufficient to trigger recombination

We investigated if the above model could explain the efficiencies of DNA recombination

that we observed on yeast cells stimulated under various conditions. In this yeast assay, the fraction

of cells that switched ON fluorescence directly reports on the fraction of DNA molecules (among

the total cell population) that underwent recombination. The corresponding experimental data can

therefore be compared with model predictions.

Although we did not quantify LiCre concentration in yeast cells, we estimated from the

literature (see Methods) that it was probably above ~1nM, conferring near-complete occupancy at

the two *loxP* sites *in vitro* (Fig. 5b). We decided to set LiCre concentration in the model at 1nM.

This ensures near-complete occupancy of the two *loxP* sites while still tolerating the possibility that

LiCre:DNA affinity may not be as high *in vivo* as it is *in vitro*. Note that this choice does not impact

the quality of the fit nor the order of magnitudes of the inferred parameters (see Methods).

In total, three parameters of the model are unknown and can be optimized to fit experimental

data: k_{OFF}, the speed of LiCre inactivation in the dark; x, the minimal number of active LiCre

protomers in the complex required for the recombination reaction to occur; and R_0, the reaction rate

of the enabled synapse. We ran the optimization of k_{OFF} and R_0 for each value of x among {1, 2, 3,

4} over a rich set of experiments performed under various conditions of illumination (different

durations, periods, duty cycles and light intensities, data sets 1-10 listed in Supplementary Table

S2). Given the batch-to-batch experimental variability mentioned above, we optimized R_0

specifically for each series of experiments. Finally, to account for the frequent loss of the LiCre

plasmid (see above), we normalized experimental data so that the highest observed efficiency was

100% (see Methods).

Results from parameter optimization are summarized in Table 1. Inferred R_0 values ranged

from 1.25 to 4.10^{-4} s^{-1} (characteristic time from 42 min to 2.2 hrs) independently of x and this

variation primarily corrected for the above-mentioned suspected effect of temperature in one of the

data sets. When x was set to either 1, 2, 3 or 4, the inferred k_{OFF} rates corresponded to characteristic

LiCre deactivation times of about 8s, 20s, 50s and 140s, respectively. It has long been known that,

once photo-activated, the LOV2 domain needs tens of seconds to return to its dark state (Swartz et

al., 2001). It is therefore unlikely that LiCre can be deactivated after only 8 seconds in the dark, thus

excluding that a functional recombination complex is formed if only one LiCre unit is active ($x =$

1). Notably, the adequacy of the model predictions to the data was maximized for $x = 2$ (Table 1)

with an overall very good fit (Fig. 5d-g). This suggests that the complex may not need four active

LiCre units to be functional, but that photo-activation of only two units may be sufficient.

In conclusion, our three-steps model can recapitulate the recombination rates observed under

various dynamics and intensities of light stimulation, and it provides a better understanding of the

LiCre-*loxP* optogenetic system.

Table 1: Best-fit parameters of the recombination reaction model.

Parameter	Data set specificity	$x = 1$	$x = 2$	$x = 3$	$x = 4$
k_{OFF} (10^{-3} s^{-1})	All	125.6	50	19.9	7.1
R_0 (10^{-4} s^{-1})	Duration, data set 7 (Fig. 5d)	3.5	3.5	3.5	4.0
	Dynamics, data sets 1&2 (Fig. 5e)	1.25	1.25	1.25	1.4
	Dynamics, data sets 3,4,5&6 (Fig. 5f)	1.25	1.4	1.4	1.6
	Intensity (Fig. 5g)	2.2	2.2	2.0	2.2
Fit score[a]	All	0.034	0.030	0.033	0.041

(a) Chi-square-like score quantifying distance to observations, see Methods.

DISCUSSION

By coupling modeling with experiments conducted both *in vitro* and *in vivo*, this work
provides knowledge on how blue light stimulates site-specific DNA recombination of the LiCre-
loxP system. We found that LiCre has high affinity for its target *loxP* substrate even if it is not
photo-activated, that this binding is cooperative – although less than for Cre - and that the observed
data are best explained by a model where the photo-activation of at least two LiCre protomers
triggers recombination. We quantified LiCre DNA-binding affinity and we estimated characteristic
times involved in the photo-induced reaction.

The fact that LiCre binds its DNA target already before illumination probably explains why
it is faster and more efficient than other photo-activatable Cre-*loxP* systems (Duplus-Bottin et al.,
2021). These other systems were obtained by splitting Cre in two halves which were fused to
optogenetic dimerizers enabling their assembly in response to light (Hochrein et al., 2018; Kawano
et al., 2016; Kuwasaki et al., 2022; Taslimi et al., 2016). Given that Cre binds DNA by capturing it
between its N- and C-terminal domains (Duyne, 2015), it is unlikely that these split subunits can
bind DNA before they are assembled together. In contrast, LiCre is made of a single peptide chain
comprising both N- and C-terminal domains, which makes it compatible with DNA binding prior to
illumination.

Our SPR measurements allowed us to compare the DNA-binding affinities of LiCre and Cre.
For Cre, we estimated K_{D1} at 1.15 nM, which is concordant with previous reports where it was
estimated at 1.7 nM by electrophoretic mobility shift assays (EMSA) (Ringrose et al., 1998) and at
1.8 nM by SPR (Rüfer et al., 2002). For K_{D2}, however, these previous studies reported values of
0.0135 nM for EMSA and 0.02 nM for SPR, which are much higher than 0.001 nM found in the
present work. This difference, which results in different estimates of DNA-binding cooperativity,
can be explained by differences in Mg^{2+} concentration between the different studies. Unlike
previous reaction conditions which did not contain any Mg^{2+} cations (Rüfer et al., 2002) or
contained 2 mM Mg^{2+} fully chelated by 1 mM EDTA (Ringrose et al., 1998), our running buffer
contained 10 mM $MgCl_2$ and no EDTA. Magnesium as well as spermidine have long been known to
increase the activity of Cre *in vitro* (Abremski and Hoess, 1984), and the previous SPR study
reported a marked increase in cooperativity when one of these ingredients was added to the

reaction. For example, the DNA-binding cooperativity moment of Cre increased from 86 to 1700 when 5 mM spermidine was added, and the authors reported that the effect of Mg^{2+} was similar (Rüfer et al., 2002). Thus, the high cooperativity previously observed in the presence of Mg^{2+} is comparable to the one we found here (cooperativity moment of 1300). We conclude that our measure of the DNA-binding affinity of Cre by SPR is consistent with previous estimates, which provides confidence in the novel measures obtained here on LiCre.

Similarly to Cre, LiCre showed very high affinity for *loxP*, and LiCre proteins bind cooperatively to DNA. However, when compared to Cre, we observed a 2-fold stronger binding of the first unit to DNA and a 7-fold weaker binding of the second unit, resulting in a much less pronounced cooperativity (Fig. 3e). Given that cooperativity results from physical interactions between protomers, this observation is consistent with the way LiCre was engineered. Indeed, the original protein-protein interactions known to take place between Cre protomers were destabilized in three ways (Duplus-Bottin et al., 2021). First, the αA helix of the N-terminal domain was truncated. Second, two point mutations destabilizing protein-protein interactions were introduced in the αN helix of the C-terminal domain. These point mutations likely affect the mechanism by which this helix was recently shown to participate to the DNA-binding cooperativity of Cre: in the absence of DNA, soluble Cre is monomeric and retains its own αN helix in *cis*, but in the presence of *loxP* DNA, the αN helix is released and becomes an anchor that stabilizes the association of the second unit (Unnikrishnan et al., 2020). Finally, the remaining segment of the αA helix was directly fused to the Jα helix of asLOV2. In the dark, the resulting chimeric helix is probably unable to interact with the partner protomer because it is presumably well folded and sequestered by the asLOV2 domain. Considering these multiple changes that destabilize protein-protein interactions, it is not surprising to observe a weaker DNA-binding cooperativity for LiCre than for Cre. It would be interesting to measure the DNA-binding affinity of light-activated LiCre. We may expect an increase in cooperativity if the chimeric helix refolds in a conformation suitable for protein-protein interaction, but probably not to the extent of Cre given the C-terminal mutations. Such measurements require dedicated equipment that can simultaneously quantify molecular associations and illuminate samples in appropriate conditions, which was not possible on our SPR station.

Unlike classical genetics or drug-based treatments, optogenetics offers the possibility to induce an intracellular reaction dynamically with high temporal precision. In the present study, applying pulsed stimuli with various dynamics and feeding the resulting data into a kinetic model allowed us to produce fine-scale information from a rather simple yeast-based assay. We could

conclude that LiCre units are photo-activated in seconds, deactivated in tens of seconds, that a
functional LiCre-*loxP* synapse can form if at least two of its four LiCre units are active and that,
following this activation, it achieves DNA recombination quite slowly with a characteristic time of
tens to hundreds of minutes. It is interesting to compare these conclusions with the kinetic
properties of Cre-mediated recombination, which were previously studied by tracking the motion of
single tethered DNA molecules *in vitro* (Fan, 2012; Pinkney et al., 2012; Shoura et al., 2012). These
studies showed that, with a constitutively-active Cre protein, limiting steps of the reaction were the
assembly of a functional synapse and the catalytic activity of the complex. For example, for intra-
molecular recombination reactions between *loxP* sites separated by 3044 and 870 bp, transposing
the estimates of reaction rates found in (Shoura et al., 2012) to equivalents of the R_0 parameter of
our model (see Methods) leads to values of 0.2-0.55 10^{-4} s^{-1}, respectively, corresponding to
characteristic times reaching several hundreds of minutes, much slower than our *in vivo* estimation
for two *loxP* sites separated by 1300 bp. Since, for LiCre, the steps of synapse assembly and
catalysis are not expected to be faster (LiCre was derived from Cre by reducing, rather than
increasing, its propensity to assemble the complex and achieve the reaction), the main difference
between *in vitro* (Shoura et al., 2012) and *in vivo* (this study) R_0 values likely lies into the
probability of forming a DNA loop bringing the two *loxP* sites together (a pre-requisite to synapse
formation). This probability depends on the mechanical properties of the inter-*loxP* DNA segment
(Rosa et al., 2010; Shoura et al., 2020); it may thus differ strongly between a stiff inter-*loxP*
segment of naked dsDNA *in vitro* and a softer, chromatinized one *in vivo* (Hajjoul et al., 2013;
Ringrose et al., 1999).

Finally, our results have practical implications on experimental protocols employing LiCre.
First, given its high affinity for *loxP* (Fig. 5b), over-expressing LiCre at high levels will probably
not increase its efficiency. On the contrary, using it at relatively low levels may maximize the
proportion of LiCre molecules containing the flavin mono-nucleotide (FMN) co-factor needed for
photo-activation. FMN derives from riboflavin and is produced by the cell. Its abundance was
previously quantified at about 1 amol per cultured mammalian cell (Hühner et al., 2015). Assuming
a cell volume of 1 pL for mammals and 0.05 pL for yeast, this suggests that a yeast cell contains
about 1,000 FMN molecules, equivalent to ~3 nM. Thus, it may be judicious to express LiCre at
low levels (a few nM) to ensure stoichiometry with FMN while still maximizing its binding to
loxP. Second, LiCre was three times more efficient at 37°C than at 20°C, with no additional
residual activity in the dark (Fig. 4c). This observation is concordant with previous *in vitro*

measurements for Cre (Buchholz et al., 1996). We therefore invite LiCre users to conduct
experiments at 37°C whenever possible. Third, our data indicate how to minimize the amount of
blue light shone on cells – and its potential damaging effects thereof - while keeping LiCre
induction maximal. With a 450nm-wavelength, we recommend to use a photon flux of 12 mW/cm^2
and periodic pulses of 1 second spaced every 10 seconds.

In conclusion, by precisely characterizing the kinetic properties of the optogenetic LiCre-
loxP recombination, this study provides useful insights for conducting targeted genetic
manipulations of live cells using light.

     METHODS

Strains and plasmids. Plasmids, strains and oligonucleotides used in this study are listed in

Supplementary Tables S3-5. Laboratory stocks were traced using MyLabStocks (Chuffart and

Yvert, 2014).

To construct LiCre-V5, we placed a Gly-Ser flexible linker at the C-terminal junction in

order to avoid interference between these epitopes and the C-terminal αN helix known to be critical

for LiCre activity (Duplus-Bottin et al., 2021). Nine repeats of the V5 epitope were amplified, using

primers 1Q46 and 1Q47, from a plasmid kindly provided by Pascal Bernard. The resulting amplicon

was co-transformed in yeast with plasmid pGY466 previously linearized by NdeI. The plasmid

(pGY605) resulting from subsequent homologous recombination was recovered from yeast colonies

and amplified in *E. coli*. We constructed Cre-V5 similarly by co-transforming in yeast the same

amplicon with pGY502 linearized at NdeI. Subsequent rescue from yeast provided plasmid

pGY607. Both plasmids were verified by Sanger sequencing, which revealed that pGY605

contained 9 V5 repeats as expected and pGY607 contained 12 V5 repeats.

For recombinant protein production, the LiCre and Cre coding sequences were codon-

optimized for *E. coli* and the corresponding sequences were synthesized and cloned in pET-19 by

Genecust to produce plasmids pGY611 (pET-LiCre) and pGY709 (pET-Cre), respectively. An

mCherry reporter targeting the *LYS2* locus was obtained by cloning a KpnI-AvrII fragment from

pGY472 into the KpnI-AvrII sites of pGY537, producing plasmid pGY618. Finally, for integration

of a single *loxP* site at the *LYS2* locus, we built plasmid pGY621 by digesting pGY618 with BamHI

and AgeI, and applying Klenow fill-in and re-ligation.

Reporter yeast strains were either GY2214 (Duplus-Bottin et al., 2021) carrying an mCherry

reporter at the *HO* locus and a GFP reporter at the *LYS2* locus, or strain GY2517 carrying the same

mCherry and GFP reporters, but at the *LYS2* and *HO* locus, respectively, or strain GY2753 carrying

a single reporter. To obtain GY2517, we transformed GY855 with pGY618 linearized by NruI and

selected Lys-, Ura+ clones (pop-in) and then 5-FoA resistant clones that had become Lys-, Ura-

(pop-out). This produced strain GY2450 which we crossed with GY1761 to obtain GY2517. To

obtain GY2753, we first cured the ade2-1 mutation of strain OAy470 (Aparicio et al., 1997) by

transforming it with a PCR amplicon containing the wild-type ADE2 sequence. The resulting strain

(GY2752) was then transformed with a 4Kb NotI fragment from pGY472.

For ChIP, strain GY2416 containing a single *loxP* site at the *LYS2* locus on the right arm of
chromosome 2 was obtained by transforming strain GY855 with plasmid pGY621 linearized at
NruI followed by pop-in and pop-out selections as above.

Purification of recombinant Cre and LiCre from bacteria. The coding sequences of Cre
and LiCre fused to a N-terminal 6xHIS tag were codon-optimized for production in *E. coli*. They
were synthesized by GeneCust and cloned in a pET-19 vector. The resulting plasmids (pGY611 and
pGY709) were transformed in Rosetta2 bacteria. Bacterial cells were cultured in LB medium
containing ampicillin and chloramphenicol at 37°C. A 40 mL starter culture was used to inoculate a
1 L culture. When reaching $OD_{600} = 0.6$, these cultures were induced for 3 h with 0.1 mM IPTG.
Cells were harvested by centrifugation and resuspended in 50 mL of 50 mM Tris pH8, 500 mM
NaCl, 10% glycerol, 0.1% Triton, 0.5 M glucose, 1 mM DTT, 1 g/L lysosyme, 2X mix of protease
inhibitors, 5 µg/mL DNAse and 2.5 mM MgCl2. After 30 minutes on ice, the suspension was
sonicated 4 times for 1 min and then centrifuged at 14,000 g for 20 min at 4°C. The resulting
supernatant was filtered on a GD/X membrane (INI) and then loaded on a HisTrap 1 mL column
(GEHealthcare) using an AKTA PURE chromatography system and a varying ratio of two buffers A
and B. Buffer A contained 50 mM Tris pH8, 500 mM NaCl, 10% Glycerol. Buffer B had the same
composition of A with imidazole added at 300 mM. Three series of injections and washes were
performed using A:B mixtures of different proportions : (100% : 0%), (95% : 5%) and (90% : 10%).
Elution was done by applying a gradient of A:B mixtures from (90% : 10%) to (50% : 50%) which
lasted 10 CV (column volume taking into account the packing and porosity of the material),
followed by a (0% : 100%) mixture for 5 CV. Fractions were analyzed by SDS-PAGE. We noted
that the eluted fractions of interest had a yellow color (Supplementary Fig. S1a), indicating that
LiCre was co-purified with its co-factor flavin mononucleotide (FMN). Fractions of interest were
concentrated on a 10 kDa membrane (Vivaspin, Millipore). The resulting products were purified
through two columns: a GF Superdex75 10/300 Increase column (Cytiva) and a Zeba column which
was eluted in the final SPR buffer 20 mM Tris pH7.4, KCl 50 mM, NaCl 150 mM, $MgCl_2$ 10 mM,
DTT 1 mM, P20 surfactant 0.05%. The quality of the production was assessed by light diffusion on
a NanosizerS (MALVERN) and SDS-PAGE (Supplementary Fig. S1b).

Chromatin immunoprecipitation (ChIP). Strain GY2416 was transformed with plasmid
pGY44, pGY605 or pGY607. Three independent transformants were used as biological replicates (9

strains in total). For each replicate, the strain was cultured overnight in 120ml liquid flasks of
synthetic SD-MW medium containing 2% glucose. Composition of this medium was the same as
for the SD-Met medium previously described (Ansel et al., 2008), except that tryptophan was
missing in the Dropout Mix. The next day, 54 mL of each culture was loaded on a 6-well plate (9
mL per well) and illuminated in a DMX-controlled 450 nm-LED box placed in a 30°C incubator.
Light intensity was set to 35 mW/cm^2. Illumination consisted of periodic pulses of 1 s applied every
10 s for a total duration of 60 minutes. In parallel, 54 mL of the same culture was kept in the dark.
For each sample (illuminated or not), 25 mL was processed for Western Blot (WB) and 25 mL was
processed for ChIP. For WB, cells were washed twice with TBS1X, aliquoted in microtubes at 10^8
cells per tube and pelleted by centrifugation. For ChIP, 700 µL of 37% Formaldehyde (Sigma
F8775) was added to the 25 mL cell suspension. After 15 min at room temperature, 1250 µL of 5M
Glycine was added and tubes were placed on a rotating wheel for 5 min at room temperature. Cells
were then washed twice with TBS1X, aliquoted in microtubes at 10^8 cells per tube and pelleted by
centrifugation. All cell pellets (WB and ChIP) were frozen in liquid nitrogen and stored at -80°C.

For ChIP, aliquoted fixed cells were thawed on ice and resuspended in 300 µL lysis buffer
composed of 0.1% w/v sodium deoxycholate, 1 mM EDTA pH8, 50 mM HEPES-KOH pH7.5, 140
mM NaCl, 1% w/v Triton X-100 and proteinase inhibitors (Sigma P2815). Tubes were mixed by
pipeting. To disrupt cells, we added 500 µL of cold acid-washed glass beads (Sigma G8772) and
placed the tubes on a PreCellys apparatus. Tubes were shaken at 6800 rpm through 2 cycles, each
cycle consisting of 10 s ON, 10 s OFF, 10 s ON. Tubes were placed for 1 min on ice between the
two cycles. Cell lysates were recovered and sonicated on a Covaris apparatus with the following
settings: P 140 W, duty factor 5%, 200 bursts per cycle, 15 min. Debris were eliminated by
centrifugation at 14,000 rpm 4°C. We then added lysis buffer to reach a volume of 1 mL. After
vortexing, 200 µL were used to control sonication efficiency on agarose gels, 750 µL were used for
immunoprecipitation (IP) and 50 µL were kept as "inputs". For IP, anti-V5 antibody (BioRad
MCA1360) was captured on Dynabeads pA (Life Technologies 10002D) and 35 µL of this
preparation were added to the sample, tubes were placed overnight at 4°C on a rotating wheel. In
parallel, the "input" samples were also placed overnight at 4°C. The following day, 50 µL of lysis
buffer were added to the "input" samples, followed by 35 µL of naked (i.e. without antibody) pA
Dynabeads, and this mixture was incubated for 1 hour at 4°C on a rotating wheel. For IP and input
samples, beads were washed three times in 0.5 mL of washing buffer I (20 mM TrisHCl pH8, 150
mM NaCl, 2 mM EDTA, 1% Triton X-100, 0.1% SDS), once in 0.5 mL of washing buffer II (20
mM TrisHCl pH8, 500 mM NaCl, 2 mM EDTA, 1% Triton X-100, 0.1% SDS), once in 0.5 mL of

washing buffer III (10 mM TrisHCl pH8, 1 mM EDTA, 0.5% deoxycholate, 1% Igepal, 25 mM

LiCl) and twice in 0.5 mL of TE8 buffer (10 mM TrisHCl, 1 mM EDTA pH8). For elution, beads

were resuspended in 130 µL of Elution Buffer (30 mM TrisHCl pH8, 2 mM EDTA, 1% SDS, 0.35

mg/mL proteinase K), incubated at 65°C for 5 to 6 hours and briefly vortexed every hour. Eluates

were transferred to fresh tubes, beads were further incubated in 20 µL of 1 M Tris pH5.5 to generate

a second eluate that was also transferred and added to the first. DNA was further purified on

NucleoSpin® Gel and PCR Clean-up Columns (Macherey Nagel) and finally eluted in 23 µL ultra-

pure water. Real-time qPCR were run on a Rotor-Gene Q thermocycler from QIAGEN using primer

pairs (1R69,1R70), (1R71,1R72), and (1R73,1R74) corresponding to probes mentioned in Fig. 1a

as P1, P2 and P3, respectively. Unrelated control loci were quantified using primer pairs

(1H97,1H98) located in *HIS5*, (1E10,1E11) located in *AHA1* and (1B36,1B37) located in *iSWH1*.

We estimated relative quantification (ChIP/INPUT) by computing the NRQ ratio described in

(Hellemans et al., 2007). This ratio follows $NRQ = (E^{\Delta Ct})/R$, where E is the amplification efficiency

of the target amplicon of interest (for example P1), ΔCt is the difference in Ct values for this

amplicon between ChIP and INPUT, and R is the geometric mean of three reference $E_i^{\Delta Cti}$ values,

with E_i being the amplification efficiency of reference i and ΔCti being the difference in Ct values

for reference i between ChIP and INPUT. This way, every ratio was normalized by all three

references $i \in \{HIS5, AHA1, iSWH1\}$.

Analysis by surface plasmon resonance (SPR). We conducted experiments on a Biacore™

T200 (Cytiva) apparatus equipped with Biacore T200 control software v3.2.1, using streptavidin-

coated Series S Sensor Chips SA (Cytiva). We conditioned the chip according to manufacturer's

recommendations (three consecutive 1 min injections of 1 M NaCl in 50 mM NaOH). After

conditioning, we used, for all experiments, a running buffer composed of 20 mM Tris pH7.4, 50

mM KCl, 150 mM NaCl, 10 mM MgCl₂, 1 mM DTT, 0.05% P20 surfactant (Cytiva). The buffer

was filtered on a 0.22 µm MCE membrane (Millipore) before use. Recorded signals were expressed

in Response Units (RU) corresponding to a rise in SPR signal resulting from protein/DNA

interactions. We prepared ligands by annealing complementary oligonucleotides corresponding to

full *loxP* (1Y68/1Y69), or half *loxP* (1Y70/1Y71) or a random sequence (1Y72/1Y73). For each

pair, one of the oligonucleotides contained a biotin group at its 5' end (Supplementary Table S5).

Each sensor chip contained 4 flow cells (channels). We first used these channels independently to

capture DNA ligands: 32.1 RU of control DNA in channel 1 of the chip, 31.6 RU of half *loxP* DNA

in channel 2, 10.6 RU of control DNA in channel 3 and 10.8 RU of *loxP* DNA in channel 4. For

kinetic measurements, we then used series of two consecutive channels: FC2-1 (half *loxP)*, and

FC4-3 (full *loxP*). Temperature of the chip was set at 25°C. Samples were kept in the Biacore

chamber at 10°C. Flow rate was set at 100 µL/min. To measure association, we injected Cre or

LiCre for 90 seconds at concentrations varying from 0.3125 to 10 nM prepared as serial two-fold

dilutions. We then monitored dissociation for 400 seconds. We applied a regeneration step between

each sample to ensure that all detectable Cre or LiCre molecules were dissociated from DNA. These

steps consisted of two 30 s injections of 2 M MgCl$_2$ in 0.5 M NaCl for full *loxP* kinetics and one 60

s injection of 2 M NaCl for half *loxP* kinetics. Sensorgram artefacts were corrected by subtracting a

control injection of running buffer.

Kinetic model of protein:DNA binding. We based this model on the previous work of

(Rüfer et al., 2002). For binding to a half-*loxP* site, we assume that the binding of Cre or LiCre is

defined by the association (k_1) and dissociation (k_{-1}) rates and follows the mass-action model (Fig.

3a):

$$\frac{d[P_1]}{dt} = k_1[C]([P_{tot}]-[P_1])-k_{-1}[P_1] \quad (1)$$

with $[P_{tot}]$ the total concentration of half-*loxP* sites, $[P_1]$ the concentration of occupied half-

loxP sites and $[C]$ the concentration of Cre/LiCre units. In the SPR protocol, Cre or LiCre

proteins are injected at a controlled concentration $[C_i]$. They diffuse within the biosensor chip

and associate with half-*loxP* sites attached to the surface (Fig. 2). This leads to the following

equation of evolution for $[C]$:

$$\frac{d[C]}{dt} = k_t([C_i]-[C])-k_1[C]([P_{tot}]-[P_1])+k_{-1}[P_1] \quad (2)$$

with k_t the coefficient of mass transport within the device.

For binding to a full *loxP* site, we assume a two-step process (Fig. 3a): one unit first binds

with the same kinetic rates as for the half-*loxP* site (k_1, k_{-1}), the second unit then associates with

possibly different kinetics (k_2, k_{-2}). This way, putative cooperative effects can be captured. This

leads to the following mass-action model:

$$\frac{d[C]}{dt} = k_t([C_i]-[C])-2k_1[C]([P_{tot}]-[P_1])+k_{-1}[P_1]-k_2[C][P_1]+2k_{-2}[P_2] \quad (3)$$

$$\frac{d[P_1]}{dt} = 2k_1[C]([P_{tot}]-[P_1])-k_{-1}[P_1]-k_2[C][P_1]+2k_{-2}[P_2] \quad (4)$$

$$\frac{d[P_2]}{dt} = k_2[C][P_1] - 2\,k_{-2}[P_2] \quad (5)$$

with $[P_1]$ the concentration of half-occupied *loxP* sites and $[P_2]$ the concentration of fully-
occupied sites.

The output signal $[B](t)$ of SPR experiments, given in Response Units (RU), effectively
measures the total mass of Cre/LiCre units bound to DNA sites at a given time *t* and is thus
proportional to $[P_1]$ in the half-*loxP* experiments, noted ($[B]^h(t) \equiv \alpha[P_1]^h$) with superindex *h*,
and to $[P_1] + 2[P_2]$ in the full-*loxP* experiments, noted ($[B]^f(t) \equiv \alpha([P_1]^f + 2[P_2]^f)$) with
superindex *f*. The maximal values for $[B](t)$ are termed RU_{max} and are defined as
$RU_{max}^h = \alpha[P_{tot}]^h$ and $RU_{max}^f = 2\alpha[P_{tot}]^f$. Each experiment consists in two successive steps: an
association phase (duration t_a) where the protein is injected ($[C_i] \equiv [C_0] > 0$ for $0 \le t \le t_a$) and a
dissociation phase in the absence of new injected material ($[C_i] = 0$ for $t \ge t_a$). To compare
quantitatively model predictions with experiments, for a given experimental condition (Cre or
LiCre, half- or full-*loxP* site, $[C_0]$) and a given set of model parameters (k_1, k_{-1}, k_2, k_{-2}, k_t, α,
[P$_{tot}$]), we numerically solved the systems of ordinary differential equations Eqs. (1-2) or Eqs. (3-5)
with initial conditions $\{[C], [P_1], [P_2]\} = \{0,0,0\}$ and then computed the corresponding SPR-like
output signal (see above).

We fitted model parameters as follows. For each type of molecule (Cre or LiCre) and site
(half- or full-*loxP*), experimental data is available for seven values of $[C_0]$ and a fixed t_a value of
90 seconds (Fig. 2). For a given set of parameters (k_1, k_{-1}, k_2, k_{-2}, k_t, α, *[P$_{tot}$]*), we simulated all
experimental conditions (*n*=14) and computed a chi-square-like score defined as:

$$\chi_2 = \sum_{e,d} \left[\frac{y_{d,e} - f(t_d;\,param)}{y_{d,e} + \varepsilon}\right]^2 \quad (6)$$

with $y_{d,e}$ the SPR signal observed at time t_d ($\in [0:0.1:340]\,sec$) during experiment *e*, *f(t$_d$;param)*
the corresponding model prediction and ε=10 a meta-parameter that grossly estimates the data
uncertainty. Note that one score was defined for the Cre dataset and another for LiCre. The
parameters for each type of molecule were fitted independently by minimizing the corresponding
χ_2 using a trust region reflective algorithm (function *least_squares* in scipy) starting from initial
parameter values randomly sampled within acceptable ranges (Supplementary Table S6). We
repeated the minimization 1,000 times per type of molecule and stored, for each fitting round, the
optimal set of parameters and the minimum score. The distribution of minimum scores showed that

some rounds of minimization only reached a local minimum and did not fully optimize the fit (Figure 3 - figure supplements 1 and 2). We discarded them from the analysis by setting an arbitrary threshold above the first mode of the distribution, keeping only 541 and 859 sets of fitted parameters for Cre and LiCre, respectively.

Quantification of LiCre-*loxP* recombination in yeast. The yeast assay presented here was performed as previously described (Duplus-Bottin et al., 2021). Yeast reporter strains were transformed with the P_{met17}-LiCre plasmid pGY466, or plasmid pGY44 (empty vector) or P_{met17}-Cre plasmid pGY502 (positive control). For each strain-plasmid combination, three independent transformants were used as biological replicates. For each replicate, a fresh colony was used to inoculate 4 mL of synthetic selective medium lacking methionine (in order to induce LiCre expression) with no particular protection against ambient light. After 18 hours, the saturated culture was transferred to two 96-well polystyrene flat-bottom sterile plates (100 µL per well); one plate was illuminated with the indicated conditions while the other plate was kept in the dark at the same temperature. We used two devices for photoactivation. The first device was a LED spot (450 nm) equipped with a DMX controller (Neptune-LED, Grenoble, France), which we placed above a thermostated platform. The second device was a homemade aluminium box which was equipped (by Neptune-LED, Grenoble, France) with a ceiling of LED strips (450 nm) connected to a DMX controller. This box was placed into an incubator at 30°C. Both devices were controlled from a laptop computer via the QLC+ software. After illumination, cells from the two plates were diluted into fresh synthetic medium in deep-well plates and incubated at 30˚C for 4 hours to allow expression of the fluorescent protein. We then analyzed cells by flow cytometry either immediately, or the day after. In this latter case, cells were either kept in PBS + 1 mM sodium azide or fixed with 2% PFA before they were analyzed the following day. We then analyzed the raw flow-cytometry data using custom R scripts as previously (Duplus-Bottin et al., 2021). No statistical power analysis was done prior to the experiment, no masking of samples IDs was applied during the experiments, and no data randomization was used.

Kinetic model of recombination. For recombination to happen, we assume that (i) both *loxP* sites should be fully occupied by two LiCre units each and that (ii) at least *x* among 4 LiCre units should be currently in the activated state. If we neglect the backward, re-insertion reaction (see main text), the probability *p* for a cell to have not recombined yet follows the evolution equation:

$$\frac{dp(t)}{dt} = -R(t)p(t) \quad \text{with} \quad R(t) = R_0 P_{2,2}(t) A_x(t) \quad (7)$$

where $P_{2,2}(t)$ is the probability at a given time t that condition (i) is effective, $A_x(t)$ is the

probability that condition (ii) is realized and R_0 is the recombination rate when both conditions

(i) and (ii) are fulfilled. Note that R_0 encompasses the rate of DNA looping between the two

loxP sites, the rate of synapse formation and the actual rate of recombination when the synapse is

formed and functional. From Eq. 7, the fraction $\phi(t)$ of cells having recombined at time t is given by

$$\Phi(t) \equiv 1 - p(t) = 1 - \exp\left(-\int_0^t dt' R(t')\right) \quad (8)$$

where we assumed that initially ($t=0$) no cells have recombined ($p(0)=1$).

Occupation of loxP sites

We assume that binding of LiCre at one *loxP* is independent from binding events at the other *loxP*

site, which we write as:

$$P_{2,2}(t) = P_2(t)^2 \quad (9)$$

with $P_2(t)$ the probability that one site is fully occupied, which is given by Eqs. (4-5) by

replacing $[P_1]$ and $[P_2]$ by $P_1(t)$ and $P_2(t)$ (the probabilities that the site is occupied by

one and two LiCre units, respectively), and by setting $[C]$ as a constant equal to the average

nuclear concentration of LiCre.

Since light does not impact LiCre binding (Fig. 1) and binding/unbinding of LiCre is much faster (~

seconds, Fig. 3e) than the recombination step (minutes), we assume that occupation of *loxP* sites is

at steady-state. Finding the stationary solutions of Eq. (4-5) and using Eq. (9) leads to

$$P_{2,2} = \left(\frac{[C]^2}{[C]^2 + 2K_{d2}[C] + K_{d1}K_{d2}}\right)^2 \quad (10)$$

Photo-activation of individual LiCre units

Assuming that the activation of one LiCre unit is independent from the state of the 3 other units

bound to the two *loxP* sites,

$$705 \quad A_x(t) = \begin{cases} \left(1 - (1 - P_{act}(t))^4\right) \text{ for } x=1 \\ P_{act}^4(t) + 4P_{act}^3(t)(1 - P_{act}(t)) + 6P_{act}^2(t)(1 - P_{act}(t))^2 \text{ for } x=2 \\ P_{act}^4(t) + 4P_{act}^3(t)(1 - P_{act}(t)) \text{ for } x=3 \\ P_{act}^4(t) \text{ for } x=4 \end{cases} \quad (11)$$

706   with $P_{act}(t)$ being the probability that one LiCre unit is activated. The kinetics of

707   activation/inactivation is supposed to follow a simple two-state model:

$$708 \quad \frac{dP_{act}}{dt} = k_{ON} \Delta(t)(1 - P_{act}) - k_{OFF} P_{act} \quad (12)$$

709   where k_{ON} is the activation rate when light is ON and depends on the lighting intensity (see below),

710    $\Delta(t)$ is a time-dependent binary variable that is equal to 1 when the light is ON and 0 otherwise, k_{OFF}

711   is the deactivation rate. For a periodic illumination defined by a period T and a duty cycle dc, Eq.

712   (12) can be solved analytically:

$$713 \quad \begin{aligned} P_{act}(t) = &\left(\frac{k_{ON}}{k_{ON}+k_{OFF}}\right)\Bigg[\Theta(dc - (t/T - n))\Big((1 - e^{-(k_{ON}+k_{OFF})(t-nT)}) \\ &+ (1 - e^{-(k_{ON}+k_{OFF})dcT})e^{-k_{OFF}(1-dc)T}\left(\frac{1 - e^{-nT(k_{ON}dc+k_{OFF})}}{1 - e^{-T(k_{ON}dc+k_{OFF})}}\right)e^{-(k_{ON}+k_{OFF})(t-nT)}\Big) \\ &+ \Theta((t/T - n) - dc)(1 - e^{-(k_{ON}+k_{OFF})dcT})\left(\frac{1 - e^{-(n+1)T(k_{ON}dc+k_{OFF})}}{1 - e^{-T(k_{ON}dc+k_{OFF})}}\right)e^{-k_{OFF}(t-nT-dcT)}\Bigg] \end{aligned} \quad (13)$$

with $n = \lfloor t/T \rfloor$ and $\Theta(u)$ being the Heaviside function.

Choice of parameters, integration and fitting

For a given set of parameters $(R_0, K_{d1}, K_{d2}, [C], k_{ON}, k_{OFF}, T, dc)$, the time evolution of $\phi(t)$ can be

computed using Eqs. (7,8,10,11,13). To compare it to experimental data, we fix parameters for

which a value was known. We chose the K_{d1}, K_{d2} values that were inferred from SPR experiments on

LiCre (Fig. 3e). For k_{ON}, we used the estimate of (Renicke et al., 2013) who computed the activation

rate of a LOV2 domain as $k_{ON} = 0.26\,SF$ with S=4.3 10^{-17} cm^2 the surface of LOV2, F=(E λ

10^{13})/2 $cm^{-2}s^{-1}$ the flux of incoming photons, λ the light wavelength (in *nm*), and E the power of the

light in *mW.cm⁻²*. In our case, this led to $k_{ON} \approx (E/35)\,s^{-1}$. The concentration of LiCre in the yeast

nucleus is not known. For simplicity, we fixed it at *[C]=1 nM* which represents about 100

molecules per nucleus. *E, T* and *dc* are defined by the illumination conditions of each experiment.

To fit the remaining parameters (x, R_0 and k_{OFF}), we considered 4 sets of experiments listed in

Supplementary Table S2, each set corresponding to a coherent series (proximal dates, same operator

and comparable conditions): experiment 8 (Fig. 5d), experiments 1 and 2 (Fig. 5e), experiments 3 to

7 (Fig. 5f) and experiment 9 (Fig. 5g). For each set of experiments, we systematically varied x from

1 to 4, R_0 between 5.10^{-5} and $1.5 \cdot 10^{-3}$ s^{-1} and k_{OFF} between 8.10^{-4} and $2.5 \cdot 10^{-2}$ s^{-1}, we computed the

corresponding model prediction for $\phi(t)$ and we estimated a chi-square-like score (similar to Eq. 6)

where experimentally-measured recombination efficiencies were normalized by a factor 0.87 to

account for the frequency of LiCre-plasmid loss (see main text). For each set e of experiments and

each x and k_{OFF} values, we estimated the minimum chi-square-like score $\chi_2^{min}(e\,;k_{OFF},x)$ and the

corresponding $R_0(e)$ value. We then estimated the value of x and k_{OFF} that minimized

$\sum_e \chi_2^{min}(e\,;k_{OFF},x)$. For each x value, optimal $\{R_0(e;x)\}$ and $k_{OFF}(x)$ values are shown in Table 1.

Note that we fitted the model by imposing the same values of x and k_{OFF} for every experiments. In

contrast, R_0 was estimated for each experiment in order to capture the observed batch-to-batch

experimental variability (which was presumably due to changing environmental conditions such as

temperature, see main text). Note also that our arbitrary choice for *[C]* does not impact the quality

of the fit nor the best values for x and k_{OFF}, because changing its value only changes $P_{2,2}$ (Eq.10)

which acts as a normalizing factor in Eq.(7). Nonetheless, the choice of *[C]* may impact on the

fitted values of $R_0(e;x)$ as follows: $R_0(e\,;x\,;[C])=R_0(e\,;x\,;[C]=1\,nM)\dfrac{P_{2,2}([C]=1\,nM)}{P_{2,2}([C])}$.

Relation between R_0 and kinetic parameters previously-estimated for Cre. In (Shoura et al.,

2012), authors fitted *in-vitro* FRET measurements by a three-state model of intra-molecular Cre-

loxP recombination (Fig. 6A of their publication). They estimated the rate of synapse formation

from the unlooped state k_3, the rate of synapse unfolding k_{-3} and the rate of recombination catalysis

k_{-4}. They found that k_{-4} was the rate-limiting step in the two cases investigated (inter-*loxP* distances

of 3044 and 870 bp). In our model, R_0 encompasses all these different reactions. In the limit $k_{-4}<<$

k_3, k_{-3}, we can approximate R_0 by the probability for a non-edited DNA molecule to be in the

synaptic state, multiplied by the catalysis rate, giving $R_0 = k_{-4}(k_3)/(k_3+k_{-3})$. We used this

formula to compare our R_0 values with corresponding estimates from Table 2 of (Shoura et al.,

2012) (see Discussion).

Intracellular LiCre concentration. Although we did not quantify the concentration of LiCre in yeast cells, we anticipate that it is above 1 nM for the following reasons. In our conditions, LiCre expression was driven by the P_{MET17} promoter and its level is therefore likely comparable to the level of Met17p. In a previous study based on LC-MS, this protein was estimated at ~120,000 copies per yeast cell (equivalent to 4 µM if we assume a cell volume of 50 µm^3), an abundance 1.6 times higher than the average abundance of the proteins quantified in this study (Data S1B of (Martin-Perez and Villén, 2017)). Another study based on fluorescently-tagged proteins (Ghaemmaghami et al., 2003) estimated an average concentration of 12,100 copies per cell (0.4 µM) per protein. So, unless LiCre is particularly unstable in yeast, its concentration should be 2 or 3 orders of magnitude above 1 nM.

MATERIAL AND CODE AVAILABILITY

Plasmids used for the production of recombinant LiCre (pET-LiCre = pGY611) or Cre (pET-Cre = pGY709) are made available from Addgene (https://addgene.org) under accession ID 219797 and 219798, respectively. The computational code implementing the models of LiCre:DNA binding and recombination reaction are made available at https://github.com/physical-biology-of-chromatin/licre-optogenetic-kinetics/.

ACKNOWLEDGEMENTS

This work has benefited from the expertise of the Protein Science (Virginie Gueguen-Chaignon and Aurélie Thibaut) and Flow Cytometry facilities of the SFR Biosciences Lyon. We thank Fabien Duveau, Mirko Francesconi and Aurèle Piazza for fruitful discussions, Fabien Duveau, Mirko Francesconi and Valérie Robert for critical reading of the manuscript, Pascal Bernard for a plasmid carrying V5-epitope repeats, Jean-Philippe Robin for advice on recombinant proteins and developers of R, Bioconductor, QLC+, numpy, git and Ubuntu for their software. This

work was supported by CNRS under the "MITI 80 Prime" program grant READGEN, ANR Grant

Opto4D ANR-23-CE12-0039, a funding for interdisciplinary innovations from the Institut

RhôneAlpin des Systèmes Complexes (IXXI) and the Federation for Systems Biology in Lyon

(BioSyL) and a funding for technological developments from the SFR Biosciences Lyon.

AUTHORS CONTRIBUTION

Designed and supervised the study: D. J. for modeling, G. Y. for experiments (except SPR).

Designed and performed the SPR experiment: A. Duf, C.M. and G. T.

Constructed plasmids and strains: A. Dum, H. D-B., G. T.

Generated datasets 1-8 and 10-11: H. D-B.

Generated dataset 9: C. D-K.

Performed ChIP and WB experiments: H. D-B.

Constructed the DNA-binding model: T. B-L., E. C. and D. J.

Fitted parameters of the DNA-binding model: D. J. and G. Y.

Constructed and analyzed the recombination model: T. B-L., E. C. and D. J.

Fitted parameters of the recombination model: D. J.

Analyzed flow-cytometry data: G. Y.

Designed and constructed LED box chassis: F. V.

Interpreted results: A. Duf., D. J., C. M., G. T., G. Y.

Made the figures: G. Y.

Wrote the paper: D. J., G. Y. with corrections from C. M. and inputs from all the authors.

Read and approved the manuscript: all authors

REFERENCES

Abremski K, Hoess R. 1984. Bacteriophage P1 site-specific recombination. Purification and properties of the Cre recombinase protein. *J Biol Chem* **259**:1509–1514. doi:10.1016/S0021-9258(17)43437-5

Altahini S, Arnoux I, Stroh A. 2024. Optogenetics 2.0: challenges and solutions towards a quantitative probing of neural circuits. *Biol Chem* **405**:43–54. doi:10.1515/hsz-2023-0194

Ansel J, Bottin H, Rodriguez-Beltran C, Damon C, Nagarajan M, Fehrmann S, Francois J, Yvert G. 2008. Cell-to-cell stochastic variation in gene expression is a complex genetic trait. *PLoS Genet* **4**:e1000049.

Aparicio OM, Weinstein DM, Bell SP. 1997. Components and Dynamics of DNA Replication Complexes in S. cerevisiae: Redistribution of MCM Proteins and Cdc45p during S Phase. *Cell* **91**:59–69. doi:10.1016/S0092-8674(01)80009-X

Armbruster A, Mohamed AM, Phan HT, Weber W. 2024. Lighting the way: recent developments and applications in molecular optogenetics. *Curr Opin Biotechnol* **87**:103126. doi:10.1016/j.copbio.2024.103126

Buchholz F, Ringrose L, Angrand P-O, Rossi F, Stewart AF. 1996. Different Thermostabilities of FLP and Cre Recombinases: Implications for Applied Site-Specific Recombination. *Nucleic Acids Res* **24**:4256–4262. doi:10.1093/nar/24.21.4256

Chuffart F, Yvert G. 2014. MyLabStocks: a web-application to manage molecular biology materials. *Yeast* **31**:179–184. doi:10.1002/yea.3008

Duplus-Bottin H, Spichty M, Triqueneaux G, Place C, Mangeot PE, Ohlmann T, Vittoz F, Yvert G. 2021. A single-chain and fast-responding light-inducible Cre recombinase as a novel optogenetic switch. *eLife* **10**:e61268. doi:10.7554/eLife.61268

Duyne GDV. 2015. Cre Recombinase. *Microbiol Spectr* **3**. doi:10.1128/microbiolspec.MDNA3-0014-2014

Fan H-F. 2012. Real-time single-molecule tethered particle motion experiments reveal the kinetics and mechanisms of Cre-mediated site-specific recombination. *Nucleic Acids Res* **40**:6208–6222. doi:10.1093/nar/gks274

Ghaemmaghami S, Huh W-K, Bower K, Howson RW, Belle A, Dephoure N, O'Shea EK, Weissman JS. 2003. Global analysis of protein expression in yeast. *Nature* **425**:737–741. doi:10.1038/nature02046

Guo F, Gopaul DN, van Duyne GD. 1997. Structure of Cre recombinase complexed with DNA in a site-specific recombination synapse. *Nature* **389**:40–6.

Hajjoul H, Mathon J, Ranchon H, Goiffon I, Mozziconacci J, Albert B, Carrivain P, Victor J-M, Gadal O, Bystricky K, Bancaud A. 2013. High-throughput chromatin motion tracking in living yeast reveals the flexibility of the fiber throughout the genome. *Genome Res* **23**:1829–1838. doi:10.1101/gr.157008.113

Hellemans J, Mortier G, De Paepe A, Speleman F, Vandesompele J. 2007. qBase relative quantification framework and software for management and automated analysis of real-time quantitative PCR data. *Genome Biol* **8**:R19. doi:10.1186/gb-2007-8-2-r19

Hochrein L, Mitchell LA, Schulz K, Messerschmidt K, Mueller-Roeber B. 2018. L-SCRaMbLE as a tool for light-controlled Cre-mediated recombination in yeast. *Nat Commun* **9**:1931. doi:10.1038/s41467-017-02208-6

Hühner J, Ingles-Prieto Á, Neusüß C, Lämmerhofer M, Janovjak H. 2015. Quantification of riboflavin, flavin mononucleotide, and flavin adenine dinucleotide in mammalian model cells by CE with LED-induced fluorescence detection. *ELECTROPHORESIS* **36**:518–525. doi:10.1002/elps.201400451

Kawano F, Okazaki R, Yazawa M, Sato M. 2016. A photoactivatable Cre-loxP recombination system for optogenetic genome engineering. *Nat Chem Biol* **12**:1059–1064. doi:10.1038/nchembio.2205

Kuwasaki Y, Suzuki K, Yu G, Yamamoto S, Otabe T, Kakihara Y, Nishiwaki M, Miyake K, Fushimi K, Bekdash R, Shimizu Y, Narikawa R, Nakajima T, Yazawa M, Sato M. 2022. A red light–responsive photoswitch for deep tissue optogenetics. *Nat Biotechnol* **40**:1672–1679. doi:10.1038/s41587-022-01351-w

Martin-Perez M, Villén J. 2017. Determinants and Regulation of Protein Turnover in Yeast. *Cell Syst* **5**:283-294.e5. doi:10.1016/j.cels.2017.08.008

Pinkney JNM, Zawadzki P, Mazuryk J, Arciszewska LK, Sherratt DJ, Kapanidis AN. 2012. Capturing reaction paths and intermediates in Cre-loxP recombination using single-molecule fluorescence. *Proc Natl Acad Sci* **109**:20871–20876. doi:10.1073/pnas.1211922109

Renicke C, Schuster D, Usherenko S, Essen L-O, Taxis C. 2013. A LOV2 Domain-Based Optogenetic Tool to Control Protein Degradation and Cellular Function. *Chem Biol* **20**:619–626. doi:10.1016/j.chembiol.2013.03.005

Ringrose L, Chabanis S, Angrand P-O, Woodroofe C, Stewart AF. 1999. Quantitative comparison of DNA looping in vitro and in vivo: chromatin increases effective DNA flexibility at short distances. *EMBO J* **18**:6630–6641. doi:10.1093/emboj/18.23.6630

Ringrose L, Lounnas V, Ehrlich L, Buchholz F, Wade R, Stewart AF. 1998. Comparative kinetic analysis of FLP and cre recombinases: mathematical models for DNA binding and recombination. *J Mol Biol* **284**:363–384. doi:10.1006/jmbi.1998.2149

Rosa A, Becker NB, Everaers R. 2010. Looping Probabilities in Model Interphase Chromosomes. *Biophys J* **98**:2410–2419. doi:10.1016/j.bpj.2010.01.054

Rüfer A, Neuenschwander PF, Sauer B. 2002. Analysis of Cre–loxP interaction by surface plasmon resonance: Influence of spermidine on cooperativity. *Anal Biochem* **308**:90–99. doi:10.1016/S0003-2697(02)00247-6

Shoura MJ, Giovan SM, Vetcher AA, Ziraldo R, Hanke A, Levene SD. 2020. Loop-closure kinetics reveal a stable, right-handed DNA intermediate in Cre recombination. *Nucleic Acids Res* **48**:4371–4381. doi:10.1093/nar/gkaa153

Shoura MJ, Vetcher AA, Giovan SM, Bardai F, Bharadwaj A, Kesinger MR, Levene SD. 2012. Measurements of DNA-loop formation via Cre-mediated recombination. *Nucleic Acids Res* **40**:7452–7464. doi:10.1093/nar/gks430

Swartz TE, Corchnoy SB, Christie JM, Lewis JW, Szundi I, Briggs WR, Bogomolni RA. 2001. The Photocycle of a Flavin-binding Domain of the Blue Light Photoreceptor Phototropin. *J Biol Chem* **276**:36493–36500. doi:10.1074/jbc.M103114200

Taslimi A, Zoltowski B, Miranda JG, Pathak GP, Hughes RM, Tucker CL. 2016. Optimized second-generation CRY2-CIB dimerizers and photoactivatable Cre recombinase. *Nat Chem Biol* **12**:425–430. doi:10.1038/nchembio.2063

Unnikrishnan A, Amero C, Yadav DK, Stachowski K, Potter D, Foster MP. 2020. DNA binding induces a cis-to-trans switch in Cre recombinase to enable intasome assembly. *Proc Natl Acad Sci* **117**:24849–24858. doi:10.1073/pnas.2011448117

Zayner JP, Antoniou C, Sosnick TR. 2012. The Amino-Terminal Helix Modulates Light-Activated Conformational Changes in AsLOV2. *J Mol Biol* **419**:61–74. doi:10.1016/j.jmb.2012.02.037

801   **Figure 3 - figure supplement 1.** Observed (dots) and predicted (lines) values for all SPR
802   sensorgrams. Parameters for Cre (panels a and c) were: α = 0.87, $RUmax_h$ = 38.2, $RUmax_f$ = 28.8, k_t
803   = 0.51, k_1 = 0.962, k_{-1} = 1.106, k_2 = 0.498, k_{-2} = 0.000452, which gave a fit score of 0.0011.
804   Parameters for LiCre (panels b and d) were: α = 1.71, $RUmax_h$ = 53.2, $RUmax_f$ = 38.1, k_t = 0.099, k_1
805   = 0.98, k_{-1} = 0.481, k_2 = 4.655, k_{-2} = 0.00286, which gave a fit score of 0.00193.

Figure 3 - figure supplement 2. Distribution of fit scores across 1,000 optimizations of the DNA-binding model for Cre (panel a) and LiCre (panel b). The lower the score, the better the fit (see Methods). Dashed blue line, arbitrary threshold used to select high-quality fits.

809   **Figure 4 - figure supplement 1.** Recombination efficiency as a function of light intensity.
810   The experiment shown in Fig. 4e (period of 7s, duty cycle of 29%, duration of 60 minutes) was
811   replicated by another operator and at a different date (dataset 11 of Supplementary Table S2).