Ecosystem's Oxygen Deficiency and Selection of Antimicrobial Resistance Genes in a One Health Perspective

Laura Ortiz-Miravalles [†,1,2], Amalia Prieto[†,1,2], Nicolas Kieffer[1,2], Ester Vergara[1,2], Rafael Cantón[3,4], Álvaro San Millán[5], Fernando Baquero[3,6], Alberto Hipólito*[1,2], **José Antonio Escudero***[1,2].

[1]Molecular Basis of Adaptation. Departamento de Sanidad Animal. Facultad de Veterinaria de la Universidad Complutense de Madrid. Spain.

[2]VISAVET Health Surveillance Centre, Universidad Complutense de Madrid.

[3]Servicio de Microbiología, Hospital Universitario Ramón y Cajal, Instituto Ramón Cajal de Investigación Sanitaria (IRYCIS), Madrid, Spain.

[4]CIBER en Enfermedades Infecciosas (CIBERINFEC), Instituto de Salud Carlos III, Madrid, Spain.

[5]Department of Microbial Biotechnology, Centro Nacional de Biotecnología - CSIC, Madrid, Spain.

[6]Network Center for Biomedical Research in Epidemiology and Public Health (CIBERESP), Madrid, Spain.

† These authors have contributed equally.

* Correspondence should be addressed to Jose Antonio Escudero (jaescudero@ucm.es) and Alberto Hipólito (alberhip@ucm.es)

Abstract

Bacteria must face and adapt to a variety of physicochemical conditions in the environment and during infection. A key condition is the concentration of dissolved oxygen, proportional to the partial pressure of oxygen (PO_2), which is extremely variable among environmental biogeographical areas and also compartments of the human and animal body. Here, we sought to understand if the phenotype of resistance determinants commonly found in Enterobacterales can be influenced by oxygen pressure. To do so, we have compared the MIC in aerobic and anaerobic conditions of isogenic *Escherichia coli* strains containing 136 different resistance genes against 9 antibiotic families. Our results show a complex landscape of changes in the performance of resistance genes in anaerobiosis. Certain changes are especially relevant for their intensity and the importance of the antibiotic family, like the large decreases in resistance observed against ertapenem and fosfomycin among *bla*$_{VIM}$ ß-lactamases and certain *fos* genes, respectively; however, the *bla*$_{OXA-48}$ ß-lactamase from the clinically relevant pOXA-48 plasmid conferred 4-fold higher ertapenem resistance in anaerobiosis. Strong changes in resistance patterns in anaerobiosis were also conserved in *Klebsiella pneumoniae*. Our results suggest that anaerobiosis is a relevant aspect that can affect the action and selective power of antibiotics for specific AMRs in different environments.

Keywords: Dissolved oxygen, anaerobiosis, antimicrobial resistance, antimicrobial resistance genes, environmental pharmacokinetics, integrons, antibiotics.

Antimicrobial resistance (AMR) is one of the major health concerns of the 21st century

globally, according to the World Health Organization (WHO)[1,2]. The dissemination of AMR

occurs in the global human-animal-environmental continuum, as considered in the One

Health perspective[3,4], yet the physicochemical conditions of these different compartments are

diverse. In humans and animals AMR frequency is mostly determined by antibiotic

consumption, following selection of bacteria harboring antibiotic resistance genes (ARGs),

frequently harbored in mobile genetic platforms such as plasmids and integron cassettes

(ARCs)[5,6]. Additionally, growing antibiotic pollution in the environment also exerts a

significant selective pressure on AMR bacteria[7]. The environmental pharmacokinetics of

antibiotics remain poorly explored, and the laboratory conditions in which the susceptibility

of pathogenic bacteria is tested (based on MIC, minimal inhibitory concentration), strongly

differ from the variable ecological compartments, in animal body sites, or in the environment

at large[8].

One of these unmet conditions in laboratory tests is the variability of oxygen

concentrations. Here, we sought to understand if the detoxifying performance of ARGs

commonly found in Enterobacterales can be influenced by dissolved oxygen (directly

proportional to the PO2, the partial pressure of oxygen) in such a way that it contributes in

various degrees to the selection of AMR in aerobic/anaerobic conditions.

To investigate the impact of anaerobiosis on resistance genes, we used mainly our pMBA

collection[9] that is composed of 136 isogenic strains of *E. coli* MG1655*,* each containing a

particular ARC as the first cassette of a class 1 integron borne by the pMBA plasmid vector[10].

The collection contains resistance genes against 9 antimicrobial families covering a broad

variety of mechanisms. We have also included other clinically relevant AMR plasmids and

genes. We have determined the MIC levels of every member in the collection in the presence

and absence of oxygen (Figs. 1B and 1C). As a control, a strain containing the empty pMBA

vector was included. To rid our analysis of the potential effects of anaerobiosis on the

bacterium's resistance -and not in the gene's function- the *performance* of an AMR gene is

measured as the fold-increase in resistance it confers compared to the empty pMBA in a given

environmental condition. Additionally, the strain containing pMBA reproduced the effect of

anaerobiosis found in *E. coli* without pMBA, allowing to rule out confounding effects of the

vector (Fig. 1A). Our data show that several ARCs varied their antimicrobial resistance effect

according to oxygen availability. For example, the aminoglycoside resistance cassette

aacA56 (Figs. 1B and 1C) increases 8-fold the MIC to amikacin in the absence of oxygen,

enhancing the effect of energy-dependent antibiotic reduced uptake. In addition, some

families of ARCs showed opposite trends in resistance in anaerobiosis: different variants of

the metallo-ß-lactamase bla_{VIM} and bla_{IMP-2} were less efficient in anaerobiosis, reducing their

performance against ertapenem between 8 and 64-fold on average (Figs. 1B and 1C). A

similar decrease in resistance was observed for several fosfomycin resistance genes (*fos*) in

anaerobiosis -with the exception of *fosE,* that showed a 4-fold increase (Figs. 1B). Because

both bla_{VIM} and *fos* encode metalloenzymes (that use respectively Zn^{2+} and Mn^{2+} as

cofactors) it is tempting to speculate that metal coordination is being affected in anaerobiosis.

Figure 1. Anaerobiosis affects ARC's function in *E. coli*. A) Heatmap showing the fold-change in resistance of *E. coli* MG1655 -with and without pMBA- between anaerobic and aerobic conditions against 18 antimicrobials (KAN: kanamycin, TOB: tobramycin, AMK: amikacin, GEN: gentamicin, STR: streptomycin, APR: apramycin, AMX: amoxicillin, CEC: cefaclor, CAZ: ceftazidime, ETP: ertapenem, ATM: aztreonam, CHL: chloramphenicol, TMP: trimethoprim, ERY: erythromycin, AZM: azithromycin, FOF: fosfomycin, CHX:

chlorhexidine, RIF: rifampicin). Numbers in each square indicate the fold-change in resistance between anaerobic and aerobic conditions. **B)** Heatmap showing the resistance profile of each ARC in anaerobiosis, normalized by its effect in aerobic conditions. Purple indicates that an ARC confers a higher resistance increase (compared to pMBA) in anaerobic conditions than in aerobic conditions, while green represents a lower resistance increase (compared to pMBA) in anaerobiosis compared to aerobiosis. Red squares depict cases for which a precise MIC could not be determined (strains showing growth at the highest concentration of antibiotic tested) in any of both conditions. Symbols > and < are used to indicate strains that grew at the highest concentration of antibiotic tested in anaerobic or aerobic conditions respectively. A hierarchical clustering tree showing protein sequence similarity is shown in the left side of each panel for (from left to right) aminoglycoside resistance cassettes, ß-lactam resistance cassettes, other ARCs. **C)** MIC fold change (compared to pMBA) in aerobiosis and anaerobiosis of the subset of ARCs, showing at least an 8-fold difference between conditions. Points represent the mode of at least three biological replicates.

At least 8 ARCs showed strong changes in their performance in anaerobiosis (defined as fold changes $\geq$ 8-fold between conditions) (Fig. 1C). This can be of clinical relevance, as carbapenemases inactivating last resource antimicrobial agents, such as bla_{VIMs} and bla_{IMP-2}, confer lower resistance levels in anaerobiosis. Our data suggests that bacteria carrying these metallo-ß-lactamase genes could become susceptible to drugs such as ertapenem in anaerobic compartments of the animal or human body (such as the large intestine), including infection sites or in particular environmental spaces. Yet, the conventional antibiogram would yield an ertapenem resistance phenotype, wrongly preventing clinicians from using a possibly successful drug. The opposite trend is observed for other ß-lactamases, such as OXA-9 and OXA-48 ß-lactamases. Indeed, plasmid pOXA-48 confers resistance to ertapenem in *E. coli* under anaerobic conditions but the strain is considered susceptible to this antibiotic under aerobic conditions. A similar phenomenon is observed for plasmid R388[11] containing bla_{VEB-}

$_1$, where resistance to ß-lactam antibiotic ceftazidime increases in anaerobiosis (Fig. 2A). The

resistance profile of an *mcr-1* did not show relevant changes (≥ 4-fold change) in anaerobic

conditions (Fig 2A).

To test if these results are conserved in other Enterobacterales, we selected the ARCs showing

≥ 8-fold change between conditions in *E. coli* and introduced them in strain ATCC23357 of

Klebsiella pneumoniae. The trends in resistance changes were similar between both species

(Fig. 2B).

Figure 2. Anaerobiosis modulates antimicrobial susceptibility across different species. A) MIC of *E. coli*

MG1655 harboring clinically relevant resistance genes and plasmids, against several antimicrobials. Bars

represent the mode of three biological replicates (black dots). Light and dark colored bars represent aerobic and

anaerobic measures respectively. S and R letters stand for susceptible and resistant according to EUCAST

clinical breakpoints. A scheme illustrating the genetic background of each ARG is displayed above each graph.

B) Comparison between anaerobic MIC fold change normalized to aerobic MIC fold change of pMBA$_{ARC}$-

containing *E. coli* and *K. pneumoniae*. The effect of oxygen concentration on ARCs is similar in both species.

MIC fold change is calculated as the mode of at least six biological replicates. Non-patterned and patterned

columns represent data retrieved from *E. coli* and *K. pneumoniae* respectively.

In this work we show how oxygen availability, as an example of physico-chemical

conditions of ecosystems can influence AMR. Our results have implications in clinical

treatments but also in environmental settings. Indeed, in natural water bodies the amount of

dissolved oxygen depends on the depth of the water column and the action of winds, waves,

photosynthetic organisms, temperature, and salinity. Water bodies are anthropogenically

polluted with antibiotics and antibiotic-resistant bacteria, many of them originating from

human and animal microbiota[12]. Hence, oxygen availability can influence selection of AMR

organisms in the environment, in particular in sewage and sludge water treatment plants, that

are frequently based on treatments decreasing dissolved oxygen[13]. Altogether, this means that

the possibility of modifying the effect on the AMR genes is ecologically and geographically

dependent, and AMR surveillance research on the relation of oxygen and the abundance of

AMRs should probably be developed in the near future[14]. Finally, we should mention the

consequences of global warming, increasing bacterial growth but reducing the availability of

dissolved oxygen in different biogeographical scenarios, thus altering the environmental

selection of AMR[15].

Conflict of Interest

The authors have no conflicts of interest to declare.

Funding

The work in the MBA laboratory is supported by the European Research Council (ERC) through a Starting Grant [ERC grant no. 803375-KRYPTONINT] and the Ministerio de Ciencia e Innovación [PID2020-117499RB-100 and CNS2022-135857]; and the EU HARISSA JPI-AMR program [PCI2021-122024-2A]. J.A.E. has been supported by the Atracción de Talento Program of the Comunidad de Madrid [2020-5A/BIO19726]; L.O. is supported by a grant from the Ministerio de Universidades (FPU21/03268). N.K. is supported by EU's Horizon 2020 Marie Skłodowska-Curie – UNA4CAREER grant (Nº: 847635). A.H. is supported by the PhD program at UCM;

Data Availability

Data sharing not applicable to this article as no datasets other than what is shown in figures were generated or analysed during the current study.

Acknowledgements

We would like to thank the members of MBA lab for their helpful comments and the funding agencies for their support.

References

1. WHO. Antimicrobial resistance. 2023 https://www.who.int/news-room/fact-sheets/detail/antimicrobial-resistance (accessed April 8, 2024).

2. Naghavi, M. *et al.* Global burden of bacterial antimicrobial resistance 1990–2021: a systematic analysis with forecasts to 2050. *The Lancet* 404, 1199–1226 (2024).

3. Hernando-Amado, S., Coque, T. M., Baquero, F. & Martínez, J. L. Defining and combating antibiotic resistance from One Health and Global Health perspectives. *Nat Microbiol* 4, 1432–1442 (2019).

4. Zhu, Y.-G. *et al.* Microbial mass movements. *Science (1979)* 357, 1099–1100 (2017).

5. Escudero*, J. A., Loot*, C., Nivina, A. & Mazel, D. The Integron: Adaptation On Demand. *Microbiol Spectr* 3, (2015).

6. MacLean, R. C. & Millan, A. S. The evolution of antibiotic resistance. *Science (1979)* 365, 1082–1083 (2019).

7. Berendonk, T. U. *et al.* Tackling antibiotic resistance: the environmental framework. *Nat Rev Microbiol* 13, 310–317 (2015).

8. Berryhill, B. A. *et al.* What's the Matter with MICs: Bacterial Nutrition, Limiting Resources, and Antibiotic Pharmacodynamics. *Microbiol Spectr* 11, (2023).

9. Hipólito, A., García-Pastor, L., Vergara, E., Jové, T. & Escudero, J. A. Profile and resistance levels of 136 integron resistance genes. *npj Antimicrobials and Resistance* 1, 13 (2023).

10. Hipólito, A. *et al.* The expression of aminoglycoside resistance genes in integron cassettes is not controlled by riboswitches. *Nucleic Acids Res* (2022) doi:10.1093/nar/gkac662.

11. Avila, P. & de la Cruz, F. Physical and genetic map of the IncW plasmid R388. *Plasmid* 20, 155–157 (1988).

12. Baquero, F., Martínez, J.-L. & Cantón, R. Antibiotics and antibiotic resistance in water environments. *Curr Opin Biotechnol* 19, 260–265 (2008).

196   13.   Radetic, B. The Anaerobic Process in Used Water Purification and Sludge
197   Treatment, Basics. in *Handbook of Water and Used Water Purification* 1–18
198   (Springer International Publishing, Cham, 2020). doi:10.1007/978-3-319-66382-
199   1_94-1.

200   14.   Sperling, E. A., Knoll, A. H. & Girguis, P. R. The Ecological Physiology of Earth's
201   Second Oxygen Revolution. *Annu Rev Ecol Evol Syst* 46, 215–235 (2015).

202   15.   Larsson, D. G. J. & Flach, C.-F. Antibiotic resistance in the environment. *Nat Rev*
203   *Microbiol* 20, 257–269 (2022).

204