Title:

Proteomic analysis to classify extracellular vesicles and plasmodesmata in

Arabidopsis cell culture supernatants

Takehito Takei[1], Teppei Asou[1], Nobuteru Hamakawa[1], Ryo Noguchi[1], Kohei Sumiyoshi[2],

Kenji Sueyoshi[3], Koshi Imami[4], Takahiro Hamada[1]

1. Department of Bioscience, Faculty of Life Science, Okayama University of Science,

Ridaicho 1-1, Okayama, 700-0005, Japan

2. Department of Applied Chemistry, Graduate School of Engineering, Osaka

Metropolitan University, Osaka, 599-8531, Japan

3. Department of Chemistry, School of Science, Kitasato University, Sagamihara,

Kanagawa, 252–0373, Japan

4. Proteome Homeostasis Research Unit, RIKEN Center for Integrative Medical Sciences,

Tsurumi-ku, Yokohama, Kanagawa 230-0045, Japan

Corresponding author e-mail:

hama.micro@ous.ac.jp

Short title: Arabidopsis PD and EV proteomes

Abstract

The plant apoplast contains diverse proteins. In this study, we analyzed the extracellular

vesicle (EV) proteome using cell culture supernatants. Unexpectedly, many

plasmodesmata proteins were identified. The comparison between the cell culture EV

proteome and the plasmodesmata proteome revealed that EV and plasmodesmata

proteins were indistinguishable. Highly sensitive imaging analyses showed that PDLP3,

a representative plasmodesmata marker, was localized to plasmodesmata, but it was also

present as small granules in apoplast regions. The small PDLP3 granules were similar in

size to the EVs, implying that the PDLPs identified during the proteomic analysis may

be small PDLP granules secreted into apoplast regions, rather than proteins derived

from the plasmodesmata itself. Next, we attempted to distinguish EV proteins from

plasmodesmata proteins. The step-wise density gradient ultracentrifugation and

clustering analysis separated EV protein clusters (represented by tetraspanins and

membrane trafficking proteins) from plasmodesmata clusters (represented by PDLPs

and PDCBs). These results provide the basis for future explorations of plant

plasmodesmata and EVs.

Introduction

Plant extracellular vesicles (EVs) are mainly secreted from multi-vesicular bodies (MVBs) generated in the endoplasmic reticulum (ER) via Golgi bodies and the trans-Golgi network (TGN) pathway (Gao et al., 2017a; Otegui, 2018). Multi-vesicular bodies also affect degradation pathways by transporting proteins and lipids to vacuoles. For example, proteins ubiquitinated in the plasma membrane are transported to MVBs via endocytosis, after which the MVBs fuse to the vacuolar membrane (Mosesso et al., 2019). Different SNAREs and Rab GTPases contribute to the fusion of MVBs to the plasma and vacuolar membranes (Ito and Uemura, 2022). For example, ARA6, VAMP721/VAMP722, and SYP121/SYP122 mediate the fusion of MVBs to the plasma membrane, whereas ARA7/RHA1 and VAMP727 participate in the fusion of MVBs to the vacuolar membrane.

However, EVs are not only derived from MVBs, suggestive of a relatively complex process. The secretion of EVs is activated in plants infected by pathogenic fungi and bacteria (Wang et al., 2017; Hansen and Nielsen, 2017). During an infection by pathogenic fungi, vacuolar and plasma membranes fuse together, which leads to the extracellular secretion of vacuolar components, including MVB-derived vesicles and autophagosomes (Cui et al., 2020). Although *Arabidopsis thaliana* (Arabidopsis) plants are normally resistant to *Golovinomyces orontii*, a mutation to *PEN1* (*SYP121*) results in plants that are susceptible to this phytopathogenic fungus that causes powdery mildew (Collins et al., 2003; Assaad et al., 2004; Lipka et al., 2005). The *SYP121* gene encodes a SNARE protein involved in fusing membranes, suggesting it contributes to the secretion of EVs. However, SYP121 was also detected in EVs, implying it may not be involved in the fusion of MVBs to the plasma membrane (Meyer et al., 2009; Nielsen and Thordal-Christensen, 2012). These findings reflect the importance of membrane trafficking factors

for plant responses to disease, but the precise functions of SYP121 on plant resistance to

phytopathogenic fungi remain unclear.

Identifying the contents of purified EVs is critical for elucidating EV origins and

functions. The purification of EVs requires the recovery of apoplasts that are not

contaminated with the inner cellular components and vascular components. Obtaining

apoplasts from the imbibed leaf lumen via centrifugation is reportedly the optimal method

(Rutter et al., 2017). In this method, the apoplast recovery solution is infiltrated into the

leaf lumen and then gently recovered by a slow-speed centrifugation. The apoplasts are

subsequently collected following a density gradient centrifugation. The use of this method

resulted in the identification of 598 proteins in the EV fraction of Arabidopsis (Rutter and

Innes, 2017). Among these proteins, several EV markers, including tetraspanins and

SYP121, were identified, indicative of the successful recovery of EVs. Additionally,

biotic and abiotic stress-responsive proteins, membrane trafficking proteins, plasma

membrane-localized proteins, and vacuolar proteins were enriched. Moreover, several

ribosomal proteins, cytoplasmic enzymes, and cell wall-associated proteins were also

identified. Similarly, 237 proteins were identified in the EV fraction of sunflower

seedlings (Regente et al., 2017). The protein composition did not differ significantly from

that of the Arabidopsis EV fraction. The increase in EV secretion following an infection

by pathogenic fungi and bacteria may be associated with a change in the EV protein

composition (e.g., increase in the abundance of resistance-related proteins). However, a

previous study on Arabidopsis showed the EV contents are not significantly affected by

a *Pseudomonas syringae* infection (Rutter and Innes, 2017). An in depth analysis of the

EV proteome of *Botrytis cinerea*-infected Arabidopsis identified 981 proteins including

several RNA-binding proteins (e.g., AGO1 and the DEAD-box RNA helicases RH11 and

RH37) (He et al., 2021).

In this study, we conducted a proteomic analysis of Arabidopsis EVs obtained from

cell culture supernatants. The protein composition of the EV proteome was similar to that

of plasmodesmata (PD) proteomes (Fernandez-Calvino et al., 2011; Brault et al., 2019).

Confocal microscopy images revealed that a typical PD marker, PDLP3-GFP, was

localized to small particles in the apoplast region where PDs are absent. The optimized

iodixanol density gradient ultracentrifugation proteomic analysis and the in depth

clustering analysis enabled the separation of EV and PD proteins. The apoplast-secreted

EV- and PD-related proteins newly identified in this study will be useful for future

research on plant apoplasts as well as investigations involving EV and PD.

Results

Extracellular vesicles were detected in the plant cell culture medium

The supernatant of a plant cell culture is ideal biochemical materials for plant apoplast studies. More specifically, it is highly homogeneous and can be obtained from large cell cultures relatively easily. Thus, to study plant EVs, we focused on cell culture supernatants. First, to verify the presence of EVs in plant cell culture supernatants, the cell culture supernatant of tobacco BY-2 cells or Arabidopsis MM2d cells were prepared via centrifugation for the subsequent analysis (Fig. 1A). The sedimented fraction on the Percoll cushion was stained with BODIPY TR Ceramide to detect sphingolipids, which are abundant in animal EVs. Many particles were detected only in the sedimented fraction on the Percoll cushion (Fig. 1B, C and D, Supplemental Movie 1). To analyze these sedimented particles, the size and number of particles were measured using qNano. The results showed that the size of these particles ranged from 70 to130 nm with a peak at 87.5 nm (Fig. 1E). The size distribution of these particles is similar to that of typical exosome like EVs (Meldolesi 2018). Furthermore, to check whether EVs were collected in the sedimented fraction, we used green fluorescent protein (GFP)-tagged SYP121, which is a plant EV marker (He et al., 2021; Rutter and Innes, 2017). The SYP121-GFP signal was detected in the sedimented fraction on the Percoll cushion and overlapped with BODIPY TR Ceramide staining (Fig. 1F). These results indicated that EVs obtained from plant cell culture supernatants.

The Arabidopsis cell culture EVs proteome revealed the EV protein composition more accurately than the earlier seedling apoplast EV proteomes

To characterize the cell culture EVs, we collected EVs from the supernatant of a

healthy 7-day old MM2d cell culture (Supplemental Figure 1) and analyzed the protein

composition of the Arabidopsis MM2d EV fraction by mass spectrometry. A total of 1,356

proteins were identified in the cell culture EV proteome (Supplemental Table 1),

including 8.2% and 24.1% of the proteins in the seedling apoplast EV proteome identified

by He et al. (2021) and Rutter and Innes (2017), respectively (Fig. 2A). The major

proteins in the cell culture EV proteome were apoplast-related proteins, membrane

trafficking proteins, cell wall-associated proteins, plasma membrane proteins, membrane

transporters, and PD-related proteins (Fig. 2B).

Compared with the earlier analyses of seedling apoplast EV proteomes, our

examination of the cell culture EV proteome identified more proteins, including specific

protein subgroups predicted to have overlapping functions. This indicates that cell culture

EVs may provide a more comprehensive overview of the EV proteome than seedling

apoplast EVs. For example, of the 17 tetraspanin proteins in Arabidopsis (Cnops et al.,

2006), only one (TET8) was identified in the seedling apoplast EV proteome (Rutter and

Innes, 2017; He et al., 2021) (Fig. 2C). In contrast, eight of the 10 highly conserved plant

tetraspanins (TET1–TET10) were identified in the cell culture EVs (Fig. 2D). TET11–

TET17, which are less conserved tetraspanin family members (Fig. 2D) and lack

homologs in other plants (Cnops et al., 2006), were not identified. Accordingly, TET1–

TET10 are likely the functional tetraspanins in plant EVs.

Membrane trafficking proteins were also identified as EV proteins (Fig. 2E). For

example, SYP121 and its functional ortholog SYP122, were identified in both the cell

culture EV and seedling apoplast EV proteomes. In addition, VAMP721/VAMP722,

which participate in the fusion of MVBs to the plasma membrane together with

SYP121/SYP122 (Kwon et al., 2008), were also identified in both the cell culture EV and seedling apoplast EV proteomes. Unlike seedling apoplast EV proteomes, the cell culture EV proteome was enriched with specific subgroups of MVBs, such as ESCRT-0 and ESCRT-3 complexes and the VPS4 complex, but not ESCRT-1 and ESCRT-2 (Fig. 2E). Thus, compared with the previously characterized EV proteomes, the cell culture EV proteome may provide additional insights into plant EV protein compositions.

In addition to the typical EV proteins, the cell culture EV proteome contained many membrane trafficking proteins, including specific subgroups of small GTPases (ARF, RAB, and ROP) (Supplemental Table 2). Earlier research showed ARA6/AtRABF1 helps mediate the fusion of MVBs to plasma membranes (Ebine et al., 2012). Additionally, AtRABA1a is colocalized with VAMP721/722 and facilitates the transport between the TGN and the plasma membrane (Asaoka et al., 2013). Both AtRABD2a and AtRABD2b are also involved in the transport from the TGN to the plasma membrane (Pinheiro et al., 2009; Drakakaki et al., 2012). The SCAMP family members, which help transport vesicles to the plasma membrane (Toyooka et al., 2009), were also enriched in the cell culture EV proteome. There were some differences between the cell culture EV proteome and the seedling apoplast EV proteome. More specifically, all clathrin heavy and light chains and many clathrin-related proteins were detected in the seedling apoplast EV proteome (He et al., 2021). Conversely, most of these proteins were not identified in the cell culture EV proteome (Supplemental Table 2). Similarly, there were differences in the identified EXOCYST-related components (Supplemental Table 2). The membrane trafficking proteins detected in the cell culture EV proteome mediate various types of transport to the plasma membrane, suggesting the cell culture EV fraction contained various types of EVs.

Transporters, proton pumps, and auxin carriers, which are plasma membrane proteins,

were also detected in the cell culture EV proteome (Supplemental Table 3). These proteins

are ubiquitinated and transported to vacuoles for degradation via the MVB pathway

(Mosesso et al., 2019). If MVBs are transported to the plasma membrane rather than to

the vacuolar membrane, the vesicles within the MVBs are considered to be EVs.

Therefore, it is not surprising that these proteins were identified in EVs.

O-Glycosyl hydrolases that degrade β-1,3-glucan were relatively abundant in the cell

culture EV fraction. Among the 51 O-glycosyl hydrolase family 17 members in

Arabidopsis, 25 were previously predicted to contain the GPI-anchor motif that binds to

membranes (Doxey et al., 2007). In the cell culture EV proteome, 23 O-glycosyl

hydrolases were identified, of which 18 had the GPI-anchor motif, suggesting that

membrane-bound O-glycosyl hydrolases are selectively concentrated in the cell culture

EV proteome (Supplemental Table 4). Some O-glycosyl hydrolases were detected in

seedling apoplast EV proteomes, but they have not been thoroughly characterized.

The cell culture EV proteome also included several enzymes, such as strictosidine

synthase, eukaryotic aspartyl protease, purple acid phosphatase, and rotamase CYP,

which were previously identified in the apoplast proteome (Supplemental Table 1)

(Uemura et al., 2019). Although their exact functions are unknown, these enzymes might

modulate secondary metabolite metabolism or cell wall regulation in apoplasts.

The EV and PD proteomes had similar compositions

Rutter and Innes (2017) reported that the identified proteins in the seedling apoplast

EV proteome overlapped approximately 50% of the identified proteins in the PD

proteome (Fernandez-Calvino et al., 2011) and the membrane trafficking-related

proteome (Heard et al., 2015). In the current study, the identified proteins in the cell

culture EV proteome included many PD proteins. In terms of the top 30 high-intensity

proteins in the cell culture EV proteome, major plant EV markers, such as tetraspanins

(TET8 and TET1) and SNAREs (SYP121, SYP122, and SYP132), were identified in both

EV and PD proteomes (Fig. 3A) (Fernandez-Calvino et al., 2011; Brault et al., 2019).

Moreover, major PD markers, such as PDLP and PCBP family members, were also

identified in the cell culture EV proteome (Fig. 3A). Several specific proteins, including

the disease resistance-related protein NDR1, late embryogenesis abundant (LEA)

proteins, O-glycosyl hydrolases, and oxidases (including SKU5 family members) were

common to the EV and PD proteomes (Fig. 3A).

Of the 115 proteins identified in an earlier analysis of a PD proteome (Brault et al.,

2019), 77 were detected in the cell culture EV proteome (Fig. 3B). Among the 50 proteins

with the highest scores in the PD proteome (Brault et al., 2019), 43 were also identified

in the cell culture EVs. A similar correlation was also observed between the seedling

apoplast EV proteome (Rutter and Innes, 2017) and the PD proteome (Fernandez-Calvino

et al., 2011); 113 proteins in the PD proteome were also among the 204 proteins in the

seedling EV proteome (Fig. 3B). In the previous proteomic analyses, PDs were isolated

from the supernatant of cell wall-digested cultures by differential centrifugation. Isolated

PDs are reported as membranous components containing 50–100 nm vesicle-like

structures (Fernandez-Calvino et al., 2011). Hence, they are similar in size to EVs. Due

to the similarities in the purification protocols and vesicle sizes, it seems that the

separation of EVs from PDs was difficult.

PDLP3 localized to small particles in the apoplast and was distinguishable from

EVs

This study identified the amount of PD-related factors in the cultured cell supernatant

fraction (Fig 3). However, large membrane structures, as seen in the purified fraction in

the PD proteome analysis (Fernandez-Calvino et al., 2011), were not observed in the

cultured cell supernatant fraction. Therefore, we considered the possibility that PD-

related factors are secreted into the extracellular space as small particles.

To assess this possibility, we used an established PD marker line, pPDLP3::PDLP3-

GFP transgenic seedling (Thomas et al., 2008) and checked the localization by a spinning-

disc confocal laser microscopy equipped with hyper-sensitive sCMOS cameras (Fig. 4A).

PDLP3-GFP clearly localized to PDs (Fig. 4A). In addition, observations with a hyper-

sensitive camera showed that PDLP3-GFP localized to particles in cells such as

membrane trafficking vesicles (Fig. 4B) as previously reported (Thomas et al., 2008).

Observations with a hyper-sensitive camera also revealed that PDLP3-GFP particles

localized to both the plasma membrane and the extracellular region of hypocotyl

epidermal cells (Fig. 4C, D). Timelapse analyses revealed that signals of extracellular

PDLP3-GFP particles were immobile, sticking one spot for a long time (Fig. 4E,

Supplemental Movie 2), whereas signals of plasma membrane-bound PDLP3-GFP

particles showed intense two-dimensional motility (Fig. 4F, Supplemental Movie 2).

These results showed that PDLP3-GFP secreted into apoplastic regions where PDs are

absent.

Next, we checked whether the extracellular PDLP3-GFP particles were identical to

EVs. To assess this possibility, we selected TagRFP-SYP121 (Ebine et al. 2011) as an EV

marker. Observation of the sample mounting water of stable transgenic plants expressing

PDLP3-GFP and TagRFP-SYP121 showed that the particles of PDLP3-GFP and

TagRFP-SYP121 were independent (Fig. 4G), indicating that the extracellular PDLP3-

GFP particles are not identical to EVs.

The EV and PD proteins were distinguished by a density gradient

ultracentrifugation

The differences in the EV and PD protein localization implied these proteins may be

separated experimentally. To generate more pure EV fractions, we separated the crude EV

fractions by density gradient ultracentrifugation. The crude EV fractions were collected

from the cell culture supernatants by ultracentrifugation and then used for the 24%–3%

step-wise iodixanol density gradient ultracentrifugation, which resulted in six fractions

(Fig. 5A) that were subjected to the size distribution analysis and the proteome analyses.

Although the size distribution analysis did not show significant differences between

fractions (Supplemental Figure 2), the proteome analyses identified 524 proteins (Fig. 5B,

Supplemental Table 5).

The 524 identified proteins were classified into 18 clusters according to their

distribution (Fig. 5B, Supplemental Table 5). Tetraspanin and SNARE proteins were

distributed between 6% and 12% (fractions 2, 3, and 4), suggesting that typical EVs in

plant cell cultures are distributed between 6% and 12% (Fig. 6A). The PDLP (PDLP1, 3,

and 6) and PDCB (PDCB1 and 4) proteins, which are representative PD proteins, were

also distributed between 6% and 12% (fractions 2, 3, and 4) (Fig. 6B). The EV markers

were often most concentrated in fraction 4, followed by fractions 2 and then 3 or they

were exclusively concentrated in fraction 4 (Fig. 6A), whereas the PD markers were most

enriched in fraction 2, followed by fractions 3 and then 4 (Fig. 6B). Accordingly, although

it is likely difficult to completely separate EVs and PD, the EV markers can be

distinguished from the PD markers by iodixanol density gradient ultracentrifugation and

clustering analyses.

We subsequently classified additional proteins in the EV, PD, and Other categories.

Among the membrane trafficking markers, the SNAREs VAMP721, VAMP724, and

YKT61 as well as SYP121 and SYP122 were classified in the EV category (Fig. 6A).

Among the ESCRT proteins involved in MVB formation, CHMP2A and VPS24A were

classified in the EV category, whereas CHMP2B, VPS20B, and SNF7A were classified

in the PD category (Fig. 6C). Moreover, SCAMP and the RAB GTPases were also divided

into the EV and PD categories (Fig. 6C). The crude EV fraction contained many

transporters belonging to the PD, EV, and Other categories (Fig. 6D). In the Other

category, NRT1:2, proton transporters, and auxin transporters were concentrated in the

heaviest fraction (i.e., fraction 6). The NDR1/HIN1 family, which is involved in disease

resistance, was clearly concentrated in the EV category (Fig. 6E). The LEA family, which

is phylogenetically closely related to the NDR1/HIN1 family, was classified in either the

EV or PD category (Supplemental Table 5). In addition, the PTI1 family, which comprises

key signaling proteins related to disease resistance (Sun et al., 2022), was concentrated in

the EV category (Fig. 6E).

A number of transporters were classified in the PD category, among which CHER1

(choline transporter) is localized in TGNs, cell plates, and sieve pores after cell division;

it is important for the development of sieve pores and PD (Dettmer et al., 2014; Kraner

et al., 2017; Gao et al., 2017b). Although CHER1 has not been confirmed as a PD-

localized protein, it was classified in the PD category in this study. Other transporters in

the PD category may also affect PD development and functions.

Oxidases and peroxidases, which regulate the apoplast redox state and pH homeostasis,

were relatively abundant in the crude EV fraction. The apoplast redox state and pH

homeostasis are crucial factors influencing cell expansion, development, and defenses

against pathogens (Smirnoff and Arnaud, 2019). Both SKU5 and SKS1 contribute to cell

wall formation (Chen et al., 2023; Sedbrook et al., 2002). In the present study, SKS3 and

SKS1 were classified in the PD and EV categories, respectively (Fig. 6F). Furthermore,

SKU5, SKS5, and SKS12 were concentrated in fractions 4 and 5, which were slightly

heavier than the EV fraction. Interestingly, most of the other oxidases and peroxidases

were also concentrated in fractions 4 and 5. Beta-glucosidases were also enriched in

fractions 4 and 5, one of which (BGLU15) is an apoplastic enzyme that mediates the

degradation of flavonol bisglycosides (Roepke et al., 2017). These results suggest that

oxidases, peroxidases, and beta-glucosidases are localized to some apoplastic structures

other than EVs and PD.

The beta-1,3-glucanases, which are O-glycosyl hydrolases involved in degrading

callose, were widely distributed in each fraction. Earlier research showed PdBG1, PdBG2,

and PdBG3 are localized to PD (Levy et al., 2007; Benitez-Alfonso et al., 2013). In the

current study, PdBG2 was classified in the PD category, but PdBG1 and PdBG3 were

concentrated in the lightest fraction (i.e., fraction 1) (Fig. 6G). Moreover, BG_pap is a

PD-localized beta-1,3-glucanase (Levy et al., 2007; Zavaliev et al., 2016). However,

BG_pap was abundant in the heaviest fraction (i.e., fraction 6) (Fig. 6G). The EV fractions

contained ZERZAUST, which is an important O-glycosyl hydrolase for intercellular

communication (Vaddepalli et al., 2017). In addition, several O-glycosyl hydrolases were

enriched in the EV fraction (Fig. 6G). These findings suggest that O-glycosyl hydrolases

are localized to various structures in the apoplast.

Discussion

Extracellular vesicles are membrane-coated vesicles in apoplasts. These vesicles are transported via various membrane trafficking pathways and degradation pathways. Plasmodesmata connect neighboring cells as cell wall-embedded structures consisting of the plasma membrane, cytoplasm, and ER. The apoplast side of PD accumulates callose and some specific components, including callose-binding proteins, callose synthases, and callose degrading enzymes (β-1,3-glucanases). On the basis of these distinct structural differences, EVs and PD are generally considered to be different structures.

The EV proteomes analyzed in a previous study (Rutter and Innes, 2017) and in the current study were revealed to contain a considerable abundance of PD proteins. Similarly, earlier examinations of PD proteomes identified EV proteins (Fernandez-Calvino et al., 2011; Brault et al., 2019). Hence, EV and PD proteins were not separated in previous proteome-level investigations, possibly because of the similarities in the purification methods. More specifically, EVs were purified from the apoplast fraction by centrifugation, while PD were purified via the centrifugation of solutions comprising digested cell walls. The reported size of EVs (50–300 nm) (Rutter and Innes, 2017) is similar to the size of vesicle-like PD structures purified from the cellulase-digested cell wall fraction (50–100 nm) (Fernandez-Calvino et al., 2011).

In the present study, we distinguished EVs from PD on the basis of a density gradient centrifugation and fine-tuned clustering. We revealed the differences in the clustering of typical EV markers, such as tetraspanins and specific SNARE family members, and typical PD markers, such as PDLP and PDCB proteins. Recent analyses of the *Physcomitrium patens* PD proteome successfully identified core PD proteins (Johnston et al., 2023; Gombos et al., 2023). The PDLP and PDCB families are apparently not

conserved in *P. patens* or in *Marchantia polymorpha* and *Klebsormidium nitens* (Bowman

et al., 2017), implying that the fraction containing the PDLP and PDCB proteins in this

study may not be the PD fraction that includes the desmotubule. A number of unclassified

clusters were identified in this study (Figure 5 and Supplemental Table 5). Of these,

cluster 18, which was enriched in the heaviest fraction (i.e., fraction 6), is notable because

it contained MCTP4, but lacked typical PD markers. The MCTP4 protein belongs to the

MCTP family, which may be involved in linking the ER membrane and the plasma

membrane within PD (Brault et al., 2019). Additionally, among the SNARE proteins, only

SYP71 was classified in cluster 18. This protein is localized to the plasma membrane, ER,

and cell plate (Suwastika et al., 2008; El Kasmi et al., 2013). These results suggest that

cluster 18 may be enriched with proteins linking the desmotubule to the plasma

membrane in PD. In the current study, the PD fraction including the PDLP and PDCB

proteins was slightly lighter than the EV fraction, suggesting that the PD fraction may

consist of only small particles (e.g., vesicles) and proteins that aggregate on the apoplast

side of PD. The heavy fraction (i.e., cluster 18) may contain PD. Clarifying whether the

proteins in cluster 18 actually belong to desmotubule–plasma membrane structures in PD

may require additional electron microscopy analyses.

In contrast to plant seedlings, cultured cells are undifferentiated and homogeneous. In

this study, we originally planned to recover pure homogeneous EVs via the

ultracentrifugation of a large volume of cell culture supernatant. However, the recovered

fraction contained various apoplast granules, including PD-associated proteins and

apoplast proteins as well as EV proteins. The density gradient centrifugation and

clustering categorized these proteins into 18 clusters, but it is unclear whether apoplast

granules and structures remained. The study findings indicate that the secretion of

apoplast materials is a complex process, even in homogeneous cell cultures. Plant

seedlings consisting of various differentiated cells secrete more complex apoplast

components. Furthermore, the types and amount of secreted apoplast components are

influenced by biotic and abiotic stimuli. Future analyses of the clusters identified in this

study will further characterize apoplast structures and physiological responses in plant

seedlings.

Materials and Methods

Growth conditions of Arabidopsis cultured cells

Arabidopsis MM2d cells (Menges and Murray, 2002) and tobacco BY-2 cells were cultured in modified LS medium (Nagata et al., 1981) on a rotary shaker (130 rpm) at 27 °C in darkness. To maintain the cell culture, 3 mL 7-day-old MM2d cultured cells were transferred to 95 mL fresh medium in 300 mL flasks. Arabidopsis plants (Col-0) were grown on 1.5% agar medium containing 1% sucrose and 1% MES-KOH (pH 5.7) in plates and in half-strength Murashige and Skoog salt medium at 22 °C for 5 days.

Isolation of EVs from the culture medium

Seven-day-old MM2d cell cultures were centrifuged twice (1,500 × g for 5 min). The supernatants were collected for the high-speed centrifugation (10,000 × g for 1 h), which was completed to eliminate any remaining debris. The purified cell culture supernatants were placed on a Percoll cushion (37% Percoll and 0.6 M sorbitol) prepared in a centrifuge tube for the subsequent ultracentrifugation (100,000 × g for 1 h). The white layer on the Percoll cushion was recovered. The EVs were further purified using exoEasy Maxi kit (Qiagen). The resulting samples, which were designated as EV fractions, were used for the mass spectrometry analysis.

For the density gradient ultracentrifugation, the supernatants of 5-day-old MM2d cell cultures were prepared via the centrifugations described above and collected from the top of the 24% iodixanol solution containing 0.25 M sucrose and 20 mM HEPES-KOH (pH 7.4) after the ultracentrifugation (100,000 × g for 1 h). The collected crude EV fractions were diluted with 10 volumes of buffer (0.25 M sucrose and 20 mM HEPES-KOH, pH 7.4) before the density gradient ultracentrifugation (100,000 × g for 1 h) involving the

24%, 12%, 6%, and 3% step-wise iodixanol density gradient. Each layer was recovered

and used for the mass spectrometry analysis.

Size distribution and particle concentration

The size and number of particles in each fraction were analyzed using qNano (Izon,

Christchurch, New Zealand). The NP200 nanopore and CPC100 calibration sample were

used for particle detection and evaluation.

Mass spectrometry

The mass spectrometry analysis was performed as described by Imami et al. (2018),

with slight modifications. Proteins were precipitated in 20% trichloroacetic acid (TFA).

The pelleted proteins were washed three times in ice-cold 100% acetone and then

resuspended in 50 μL resuspension buffer (8 M urea and 0.1 M Tris-HCl, pH 8). After

adding 10 mM dithiothreitol, the proteins were reduced at room temperature for 30 min

and then alkylated using 50 mM iodoacetamide at room temperature for 30 min in

darkness. Proteins were digested with lysyl endopeptidase (LysC) (Wako) at room

temperature for 3 h. The sample solutions were diluted to a final concentration of 2 M

urea and 50 mM ammonium bicarbonate prior to the trypsin (Promega) digestion, which

was performed with constant agitation at room temperature for 16 h. The resulting peptide

samples were desalted using SDB-XC (upper)/SCX (bottom) (GL Sciences) StageTips

(Adachi et al., 2016). The peptides were sequentially eluted using elution buffer 1 (4%

TFA, 500 mM ammonium acetate, and 30% acetonitrile) and elution buffer 2 (500 mM

ammonium acetate and 30% acetonitrile).

The nano-scale reversed-phase liquid chromatography and tandem mass spectrometry

(nanoLC/MS/MS) analysis was completed using the Orbitrap Fusion Lumos mass

spectrometer (Thermo Fisher Scientific) connected to a Thermo Ultimate 3000

RSLCnano pump and an HTC-PAL autosampler (CTC Analytics, Zwingen, Switzerland)

equipped with a self-packing analytical column (150 mm long with 100 μm internal

diameter) (Ishihama et al., 2002) packed with ReproSil-Pur C18-AQ (3 μm; Dr. Maisch

GmbH). Mobile phase A consisted of 0.5% acetic acid, whereas mobile phase B

comprised 0.5% acetic acid and 80% acetonitrile. Peptides were eluted from the analytical

column at a flow rate of 500 nL/min with the following gradient: 5%–10% B in 5 min,

10%–40% B in 20 min, 40%–99% B in 1 min, and 99% for 5 min. The Orbitrap Fusion

Lumos instrument was operated in the data-dependent mode with a full scan in the

Orbitrap followed by MS/MS scans for 3 s for the higher-energy collisional dissociation

(HCD). The applied voltage for ionization was 2.4 kV. The full scans were performed

with a resolution of 120,000, a target value of 4×10^5 ions, and a maximum injection time

of 50 ms. The MS scan range was m/z 300–1,500. The MS/MS scans were performed

with a resolution of 15,000, a target value of 5×10^4 ions, and a maximum injection time

of 50 ms. The isolation window was set to 1.6. The normalized HCD collision energy was

30. Dynamic exclusion was applied for 20 s.

All raw data files were analyzed and processed using MaxQuant (v1.6.15.0) (Cox and

Mann, 2008). Andromeda (Cox et al., 2011) was used to search the

Araport11_genes.201606.pep (https://www.arabidopsis.org/) database spiked with

common contaminants and enzyme sequences. The search parameters included two

missed cleavage sites and variable modifications, including methionine oxidation,

deamidation of glutamine and asparagine and protein N-terminal acetylation. Cysteine

carbamidomethylation was set as a fixed modification. The peptide mass tolerance was

4.5 ppm, whereas the MS/MS tolerance was 20 ppm. The false discovery rate at the

peptide spectrum match and protein levels was set to 1%.

Microscopy analyses

The confocal microscopy analysis was performed using the ECLIPSE Ti microscope

(Nikon) equipped with a confocal spinning disc unit (CSU-W1, Yokogawa) and the

sCMOS camera (Sona, Andor). The isolated EVs were stained with 1 μM BODIPY TR

Ceramide (ThermoFisher Scientific). For the optical sectioning of Arabidopsis seedlings,

transgenic plants expressing both PDLP3-GFP and TagRFP-SYP121 were mounted

between 60×24 and 18×18 coverslips (Iwanami) with a 0.15 mm spacer. Images were

captured using NIS-Element AR (Nikon) and analyzed using Fiji (Schindelin et al., 2012).

Acknowledgements and fundings

We thank Tsuyoshi Nakagawa (Shimane University) for providing Gateway binary

vectors containing bar gene identified by Meiji Seika Kaisha, Ltd., Andrew J. Maule

(John Innes Centre) for providing the PDLP3-GFP transgenic plant, Masa H. Sato and

Tomoko Hirano (Kyoto Pref. Univ.) for providing the pGWB1-GFP-SYP121 plasmid,

Kazuo Ebine and Takashi Ueda (NIBB) for providing the TagRFP-SYP121 seedling,

Akihiko Nakano (Riken) and Tomohiro Uemura (Ochanimizu Univ) for helpful

discussion. This study was supported by JST-PRESTO [JPMJPR18H7 to T. H.,

JPMJPR19H7 to K. Sueyoshi, and JPMJPR18H2 to K. I.] and JST-CREST

[JPMJCR18H4 to T. H. and JPMJCR19H1 to K. Sueyoshi].

 Figure legends

Figure 1. Isolation of EVs from a plant cell culture suspension

A) Protocol for the isolation of EVs from plant cell cultures. **B–D)** BODIPY TR Ceramide

staining of the ultracentrifugation-derived sedimented fraction (B) and supernatant

fraction (C) for the tobacco BY-2 culture medium and the ultracentrifugation-derived

sedimented fraction for the medium before culturing (D). The panels on the right present

the enlarged images of the area marked by a white square in the panels on the left. These

images correspond to the optical section presented in Supplemental Movie 1. Bar, 2 μm.

E) Size distribution of sedimented particles by ultracentrifugation from the supernatant

of Arabidopsis MM2d cells. **F)** Purified EVs from SYP121-GFP transgenic tobacco BY-

2 cell cultures. Bar, 2 μm.

Figure 2. Comparison between the cell culture EV proteome and the seedling

apoplast EV proteome

A) Venn diagram of the identified proteins in the cell culture EV proteome and the

seedling apoplast EV proteome. **B)** Classification of the top 100 high-intensity proteins

in the cell culture EV proteome. **C)** Identified tetraspanin family members in the cell

culture EV proteome and the seedling apoplast EV proteome. **D)** Phylogenetic tree of the

tetraspanin family in *Arabidopsis thaliana*. Identified tetraspanin family members in the

cell culture EV proteome were colored in gray. **E)** Identified membrane trafficking

proteins in the cell culture EV proteome and the seedling apoplast EV proteome.

Figure 3. Comparison between the EV proteome and the plasmodesmata

proteome

A) Comparison of the EV proteomes and the plasmodesmata proteomes in terms of the

top 30 high-intensity proteins in the cell culture EV proteomes. **B)** Overlap percentages

between the three EV proteomes and the two plasmodesmata proteomes. The EV and

plasmodesmata fractions were indistinguishable in these analyses.

Figure 4. Localization of PDLP3-GFP in the apoplast region

A-D) Confocal microscopy images of PDLP3-GFP. Plasma membrane was labeled by

Plasmembrane Red. PDLP3-GFP clearly localized to plasmodesmata (A, white

arrowheads). At high contrast images, the PDLP3-GFP signals were also observed as

particles at inside (B), plasma membrane (C), and surface (D). Some PDLP3-GFP signals

were detected as small particles in the apoplast region (D, white arrowheads). Bar, 10 μm.

E, F) Projection images from PDLP3-GFP time-lapse analyses. The 15 time-lapse images

were acquired every 5 seconds and grouped into early (1-5), middle (6-10), and late (11-

15) phases. These images were projected at maximum intensity, and superimposed into

one merge image using blue, green and red colour, respectively. Bars, 10 μm. **G)**

Comparison of PDLP3-GFP and TagRFP-SYP121 at the apoplast region. TagRFP-

SYP121 (white arrowheads) were not colocalized with PDLP3-GFP (yellow arrowheads).

Bar, 5 μm.

Figure 5. Iodixanol density gradient ultracentrifugation of the crude EV fraction

A) Separation of the crude EV fraction by an iodixanol density gradient

ultracentrifugation. The crude EV fraction was mounted on the top of the iodixanol

solution during the density gradient ultracentrifugation (24%, 12%, 6%, and 3% iodixanol

solutions). After the ultracentrifugation, six fractions were recovered and used for

proteome analyses. B) Classification of the 524 identified proteins on the basis of their

distribution. The presented data correspond to the information provided in Supplemental

Table 5.

Figure 6. Representative proteins in each gene ontology category

A) Representative EV proteins. B) Representative PD proteins. C) Membrane trafficking-

related proteins. D) Membrane transporters and pumps. E) Disease resistance-related

proteins. F) Oxidases and peroxidases. G) O-glycosyl hydrolases. The presented data

correspond to the information provided in Supplemental Table 5. cl. indicates the cluster

number in Figure 5B; cat. indicates category.

Supplemental Figure 1. Growth condition of MM2d Arabidopsis cultured cells for

this study

A) Growth curve of MM2d Arabidopsis cultured cells in this study. Supernatant of 5 days-

old and 7 days-old cultured cells was applied for EV purification. B) DNA degradation

assay of MM2d cells. In 5 days-old and 7 days-old cultured cells, DNA degradation did

not occur at all.

Supplemental Figure 2. Size distribution of EVs in each fraction after an iodixanol

density gradient ultracentrifugation

Six fractions were recovered after an iodixanol density gradient ultracentrifugation and

used for size distribution analyses. These fractions correspond to the fractions in Figure

5A.

Supplemental Table 1　List of 1356 proteins identified in the cultured cell EV

proteomics

Supplemental Table 2 Comparison of membrane traffic proteins between the cultured

cell and the leaf apolast EV proteomics

Supplemental Table 3　Comparison of transpoters and ubiquitin-related proteins

between the cultured cell and the leaf apolast EV proteomics

Supplemental Table 4　Comparison of O-Glycosyl hydrolases family 17 proteins

between the cultured cell and the leaf apolast EV proteomics

Supplemental Table 5　List of 524 proteins Identified in the density gradient ultra-

centrifugation proteome

Supplemental Movie 1 Optical sections of isolated EVs.

EVs were stained by BODIPY TR Ceramide. Images were taken from the surface of

coverslip toward to the top of a drop.

Supplemental Movie 2　Time-lapse analyses of PDLP3-GFP.

Images were taken from the same seedling at different depths. These images were used

to generate the projection images of Figure 4E and F. Bars, 10 μm.

References

Adachi, J., Hashiguchi, K., Nagano, M., Sato, M., Sato, A., Fukamizu, K.,

Ishihama, Y., and Tomonaga, T.. Improved proteome and phosphoproteome

analysis on a cation exchanger by a combined acid and salt gradient. Anal. Chem.

2016:**88**: 7899–7903.

Asaoka, R., Uemura, T., Ito, J., Fujimoto, M., Ito, E., Ueda, T., and Nakano, A..

Arabidopsis RABA1 GTPases are involved in transport between the trans-Golgi

network and the plasma membrane, and are required for salinity stress tolerance.

Plant J. 2013:**73**: 240–249.

Assaad, F.F., Qiu, J.-L., Youngs, H., David, E., Zimmerli, L., Kalde, M., Wanner,

G., Peck, S.C., Edwards, H., Ramonell, K., Somerville, C.R., and Thordal-

Christensen, H. The PEN1 Syntaxin Defines a Novel Cellular Compartment upon

Fungal Attack and Is Required for the Timely Assembly of Papillae. Mol Biol Cell

2004:**15**: 5318–5328.

Benitez-Alfonso, Y., Faulkner, C., Pendle, A., Miyashima, S., Helariutta, Y., and

Maule, A. Symplastic Intercellular Connectivity Regulates Lateral Root

Patterning. Dev. Cell 2013:**26**: 136–147.

Bowman, J.L. et al. Insights into Land Plant Evolution Garnered from the Marchantia

polymorpha Genome. Cell 2017:**171**: 287-304.e15.

Brault, M.L. et al. Multiple C2 domains and transmembrane region proteins (MCTP s)

tether membranes at plasmodesmata . EMBO Rep. 2019:**20**: 1–26.

Chen, C. et al. Multi-copper oxidases SKU5 and SKS1 coordinate cell wall formation

using apoplastic redox-based reactions in roots. Plant Physiol. 2013:**192**: 2243–

2260.

Clough, S.J. and Bent, A.F. Floral dip: A simplified method for Agrobacterium-

mediated transformation of Arabidopsis thaliana. Plant J. 1998:**16**: 735–743.

Cnops, G. et al. The TORNADO1 and TORNADO2 genes function in several

patterning processes during early leaf development in Arabidopsis thaliana. Plant

Cell 2006:**18**: 852–866.

Collins, N.C., Thordal-christensen, H., and Lipka, V. SNARE-protein-mediated

disease resistance at the plant cell wall. Nature 2003:**312**: 310–312.

Cox, J. and Mann, M. MaxQuant enables high peptide identification rates,

individualized p.p.b.-range mass accuracies and proteome-wide protein

quantification. Nat. Biotechnol. 2008:**26**: 1367–1372.

Cox, J., Neuhauser, N., Michalski, A., Scheltema, R.A., Olsen, J. V, and Mann, M.

Andromeda : A Peptide Search Engine Integrated into the MaxQuant

Environment.: 2011:**10(4)**:1794–1805.

Cui, Y., Gao, J., He, Y., and Jiang, L. Plant extracellular vesicles. Protoplasma

2020:**257**: 3–12.

Dettmer, J. et al. CHOLINE TRANSPORTER-LIKE1 is required for sieve plate

development to mediate long-distance cell-to-cell communication. Nat. Commun.

2014:**4276**.

Doxey, A.C., Yaish, M.W.F., Moffatt, B.A., Griffith, M., and McConkey, B.J.

Functional divergence in the Arabidopsis β-1,3-glucanase gene family inferred by

phylogenetic reconstruction of expression states. Mol. Biol. Evol. 2007:**24**: 1045–

1055.

Drakakaki, G., Van De Ven, W., Pan, S., Miao, Y., Wang, J., Keinath, N.F.,

Weatherly, B., Jiang, L., Schumacher, K., Hicks, G., and Raikhel, N. Isolation

616 and proteomic analysis of the SYP61 compartment reveal its role in exocytic

617 trafficking in Arabidopsis. Cell Res. 2012:**22**: 413–424.

618 **Ebine, K., Miyakawa, N., Fujimoto, M., Uemura, T., Nakano, A., and Ueda, T.**

619 Endosomal trafficking pathway regulated by ARA6, a RAB5 GTPase unique to plants.

620 Small GTPases 2012:**3**: 23–27.

621 **Fernandez-Calvino, L., Faulkner, C., Walshaw, J., Saalbach, G., Bayer, E.,**

622 **Benitez-Alfonso, Y., and Maule, A.** Arabidopsis plasmodesmal proteome. PLoS

623 One 2011:**6**.

624 **Gao, C., Zhuang, X., Shen, J., and Jiang, L.** Plant ESCRT Complexes: Moving

625 Beyond Endosomal Sorting. Trends Plant Sci. 2017a:**22**: 986–998.

626 **Gao, Y.Q., Chen, J.G., Chen, Z.R., An, D., Lv, Q.Y., Han, M.L., Wang, Y.L., Salt,**

627 **D.E., and Chao, D.Y.** A new vesicle trafficking regulator CTL1 plays a crucial

628 role in ion homeostasis. PLoS Biol. 2017b:**15**: 1–27.

629 **Gombos, S. et al.** A high-confidence Physcomitrium patens plasmodesmata proteome

630 by iterative scoring and validation reveals diversification of cell wall proteins

631 during evolution. New Phytol. 2023:**238**: 637–653.

632 **Hansen, L.L. and Nielsen, M.E.** Plant exosomes: Using an unconventional exit to

633 prevent pathogen entry? J. Exp. Bot. 2017:**69**: 59–68.

634 **He, B., Cai, Q., Qiao, L., Huang, C.Y., Wang, S., Miao, W., Ha, T., Wang, Y., and**

635 **Jin, H.** RNA-binding proteins contribute to small RNA loading in plant

636 extracellular vesicles. Nat. Plants 2021:**7**: 342–352.

637 **Heard, W., Sklenář, J., Tomé, D.F.A., Robatzek, S., and Jones, A.M.E.**

638 Identification of regulatory and cargo proteins of endosomal and secretory

639 pathways in arabidopsis thaliana by proteomic dissection. Mol. Cell. Proteomics

2015:**14**: 1796–1813.

Imami, K., Milek, M., Bogdanow, B., Yasuda, T., Kastelic, N., Zauber, H.,

Ishihama, Y., Landthaler, M., and Selbach, M. Phosphorylation of the

Ribosomal Protein RPL12/uL11 Affects Translation during Mitosis. Mol. Cell

2018:**72**: 84-98.e9.

Ishihama, Y., Rappsilber, J., Andersen, J.S., and Mann, M. Microcolumns with self-

assembled particle frits for proteomics. J. Chromatogr. A 2002:**979**: 233–239.

Ito, Y. and Uemura, T. Super resolution live imaging: The key for unveiling the true

dynamics of membrane traffic around the Golgi apparatus in plant cells. Front.

Plant Sci. 2022:**13**: 1–8.

Johnston, M.G., Breakspear, A., Samwald, S., Zhang, D., Papp, D., Faulkner, C.,

and de Keijzer, J. Comparative phyloproteomics identifies conserved

plasmodesmal proteins. J. Exp. Bot. 2023:**74**: 1821–1835.

El Kasmi, F., Krause, C., Hiller, U., Stierhof, Y.D., Mayer, U., Conner, L., Kong,

L., Reichardt, I., Sanderfoot, A.A., and Jürgens, G. SNARE complexes of

different composition jointly mediate membrane fusion in Arabidopsis cytokinesis.

Mol. Biol. Cell 2013:**24**: 1593–1601.

Kraner, M.E., Link, K., Melzer, M., Ekici, A.B., Uebe, S., Tarazona, P., Feussner,

I., Hofmann, J., and Sonnewald, U. Choline transporter-like1 (CHER1) is crucial

for plasmodesmata maturation in Arabidopsis thaliana. Plant J. 2017:**89**: 394–406.

Kwon, C. et al. Co-option of a default secretory pathway for plant immune responses.

Nature 2008:**451**: 835–840.

Lee, J.Y., Wang, X., Cui, W., Sager, R., Modla, S., Czymmek, K., Zybaliov, B.,

Van Wijk, K., Zhang, C., Lu, H., and Lakshmanana, V. A plasmodesmata-

664 localized protein mediates crosstalk between cell-to-cell communication and innate

665 immunity in arabidopsis. Plant Cell 2011:**23**: 3353–3373.

666 **Levy, A., Erlanger, M., Rosenthal, M., and Epel, B.L.** A plasmodesmata-associated

667 β-1,3-glucanase in Arabidopsis. Plant J. 2007:**49**: 669–682.

668 **Lipka, V. et al.** Plant science: Pre- and postinvasion defenses both contribute to

669 nonhost resistance in Arabidopsis. Science. 2005:**310**: 1180–1183.

670 **Matsui, H., Matsui, H., Fujiwara, M., Hamada, S., Shimamoto, K., Nomura, Y.,**

671 **Nakagami, H., Takahashi, A., and Hirochika, H.** Plasma membrane localization

672 is essential for oryza sativa pto-interacting protein 1a-mediated negative regulation

673 of immune signaling in rice. Plant Physiol. 2014:**166**: 327–336.

674 **Menges, M. and Murray, J.A.H.** Synchronous Arabidopsis suspension cultures for

675 analysis of cell-cycle gene activity. Plant J. 2002:**30**: 203–212.

676 **Meyer, D., Pajonk, S., Micali, C., O'Connell, R., and Schulze-Lefert, P.**

677 Extracellular transport and integration of plant secretory proteins into pathogen-

678 induced cell wall compartments. Plant J. 2009:**57**: 986–999.

679 **Meldolesi, J.** Exosomes and ectosomes in intercellular communication. Curr. Biol.

680 2018:28: R435-R444.

681 **Mosesso, N., Nagel, M.K., and Isono, E.** Ubiquitin recognition in endocytic trafficking

682 - With or without ESCRT-0. J. Cell Sci. 2019:**132**.

683 **Nagata, T., Okada, K., Takebe, I., and Matsui, C.** Delivery of tobacco mosaic virus

684 RNA into plant protoplasts mediated by reverse-phase evaporation vesicles

685 (Liposomes). MGG Mol. Gen. Genet. 1981:**184**: 161–165.

686 **Nakamura, S., Mano, S., Tanaka, Y., Ohnishi, M., Nakamori, C., Araki, M., Niwa,**

687 **T., Nishimura, M., Kaminaka, H., Nakagawa, T., Sato, Y., and Ishiguro, S.**

Gateway binary vectors with the bialaphos resistance gene, bar, as a selection

marker for plant transformation. Biosci. Biotechnol. Biochem. 2010:**74**: 1315–

1319.

Nielsen, M.E. and Thordal-Christensen, H. Recycling of Arabidopsis plasma

membrane PEN1 syntaxin. Plant Signal. Behav. 2012:**7**: 1541–1543.

Otegui, M.S. ESCRT-mediated sorting and intralumenal vesicle concatenation in

plants. Biochem. Soc. Trans. 2018:**46**: 537–545.

Pinheiro, H., Samalova, M., Geidner, N., Chory, J., Martinez, A., and Moore, I.

Genetic evidence that the higher plant Rab-D1 and Rab-D2 GTPases exhibit

distinct but overlapping interactions in the early secretory pathway. J. Cell Sci.

2009:**122**: 3749–3758.

Regente, M., Pinedo, M., Clemente, H.S., Balliau, T., Jamet, E., and De La Canal,

L. Plant extracellular vesicles are incorporated by a fungal pathogen and inhibit its

growth. J. Exp. Bot. 2017:**68**: 5485–5495.

Roepke, J., Gordon, H.O.W., Neil, K.J.A., Gidda, S., Mullen, R.T., Freixas Coutin,

J.A., Bray-Stone, D., and Bozzo, G.G. An apoplastic β-glucosidase is essential

for the degradation of flavonol 3-O-β-Glucoside-7-O-α-rhamnosides in

arabidopsis. Plant Cell Physiol. 2017:**58**: 1030–1047.

Rutter, B., Rutter, K., and Innes, R. Isolation and Quantification of Plant

Extracellular Vesicles. Bio-Protocol 2017:**7**: 1–13.

Rutter, B.D. and Innes, R.W. Extracellular vesicles isolated from the leaf apoplast

carry stress-response proteins. Plant Physiol. 2017:**173**: 728–741.

Schindelin, J. et al. Fiji: An open-source platform for biological-image analysis. Nat.

Methods 2012:**9**: 676–682.

Sedbrook, J.C., Carroll, K.L., Hung, K.F., Masson, P.H., and Somerville, C.R. The

Arabidopsis SKU5 gene encodes an extracellular glycosyl phosphatidylinositol-

anchored glycoprotein involved in directional root growth. Plant Cell 2002:**14**:

1635–1648.

Smirnoff, N. and Arnaud, D. Hydrogen peroxide metabolism and functions in plants.

New Phytol. 2019:**221**: 1197–1214.

Sun, M., Qiu, L., Liu, Y., Zhang, H., Zhang, Y., Qin, Y., Mao, Y., Zhou, M., Du, X.,

Qin, Z., and Dai, S. Pto Interaction Proteins: Critical Regulators in Plant

Development and Stress Response. Front. Plant Sci. 2022:**13**: 1–11.

Suwastika, I.N., Uemura, T., Shiina, T., Sato, M.H., and Takeyasu, K. SYP71, a

plant-specific Qc-SNARE protein, reveals dual localization to the plasma

membrane and the endoplasmic reticulum in Arabidopsis. Cell Struct. Funct.

2008:**33**: 185–192.

Thomas, C.L., Bayer, E.M., Ritzenthaler, C., Fernandez-Calvino, L., and Maule,

A.J. Specific targeting of a plasmodesmal protein affecting cell-to-cell

communication. PLoS Biol. 2008:**6**: 0180–0190.

Toyooka, K., Goto, Y., Asatsuma, S., Koizumi, M., Mitsui, T., and Matsuoka, K. A

mobile secretory vesicle cluster involved in mass transport from the golgi to the

plant cell exterior. Plant Cell 2009:**21**: 1212–1229.

Uemura, T., Nakano, R.T., Takagi, J., Wang, Y., Kramer, K., Finkemeier, I.,

Nakagami, H., Tsuda, K., Ueda, T., Schulze-Lefert, P., and Nakano, A. A

golgi-released subpopulation of the trans-golgi network mediates protein secretion

in arabidopsis 1. Plant Physiol. 2019:**179**: 519–532.

Vaddepalli, P., Fulton, L., Wieland, J., Wassmer, K., Schaeffer, M., Ranf, S., and

Schneitz, K. The cell wall-localized atypical β-1,3 glucanase ZERZAUST controls

tissue morphogenesis in Arabidopsis thaliana. Dev. 2017:**144**: 2259–2269.

Wang, X., Chung, K.P., Lin, W., and Jiang, L. Protein secretion in plants:

Conventional and unconventional pathways and new techniques. J. Exp. Bot.

2017:**69**: 21–37.

Zavaliev, R., Dong, X., and Epel, B.L. Glycosylphosphatidylinositol (GPI)

modification serves as a primary plasmodesmal sorting signal. Plant Physiol.

2016:**172**: 1061–1073.

Figure 1. Isolation of EVs from a plant cell culture suspension

A) Protocol for the isolation of EVs from plant cell cultures. **B–D)** BODIPY TR Ceramide staining of the ultracentrifugation-derived sedimented fraction (B) and supernatant fraction (C) for the tobacco BY-2 culture medium and the ultracentrifugation-derived sedimented fraction for the medium before culturing (D). The panels on the right present the enlarged images of the area marked by a white square in the panels on the left. These images correspond to the optical section presented in Supplemental Movie 1. Bar, 2 μm. **E)** Size distribution of sedimented particles by ultracentrifugation from the supernatant of Arabidopsis MM2d cells. **F)** Purified EVs from SYP121-GFP transgenic tobacco BY-2 cell cultures. Bar, 2 μm.

Figure 2. Comparison between the cell culture EV proteome and the seedling apoplast EV proteome
A) Venn diagram of the identified proteins in the cell culture EV proteome and the seedling apoplast EV proteome. **B)** Classification of the top 100 high-intensity proteins in the cell culture EV proteome. **C)** Identified tetraspanin family members in the cell culture EV proteome and the seedling apoplast EV proteome. **D)** Phylogenetic tree of the tetraspanin family in Arabidopsis thaliana. Identified tetraspanin family members in the cell culture EV proteome were colored in gray. **E)** Identified membrane trafficking proteins in the cell culture EV proteome and the seedling apoplast EV proteome.

A

AGI code	Name	Protein ID	Cultured cell EV this study	Category	Seedling EV Rutter et al. 2017	Seedling EV He et al. 2021	PD Fernandez-Calvino et al. 2011	PD Brault et al. 2019	Apoplast Uemura et al. 2019
AT5G61130	plasmodesmata callose-binding protein 1	PDCB1	*	PD	N.D.	N.D.	N.D.	*	*
AT1G69295	plasmodesmata callose-binding protein 4	PDCB4	*	PD	N.D.	N.D.	N.D.	*	*
AT5G06320	NDR1/HIN1-like 3	NHL3	*	Cytoplasm	*	*	*	*	N.D.
AT1G64760	O-Glycosyl hydrolases family 17 protein	1,3-beta-glucosidase	*	apoplast	N.D.	N.D.	*	*	N.D.
AT2G23810	tetraspanin8	TET8	*	TET	*	*	*	*	N.D.
AT4G31140	O-Glycosyl hydrolases family 17 protein	1,3-beta-glucosidase	*	apoplast	*	*	*	*	N.D.
AT5G43980	plasmodesmata-located protein 1	PDLP1	*	PD	N.D.	N.D.	*	*	*
AT1G74020	strictosidine synthase 2	SSL12	*	apoplast	N.D.	N.D.	*	N.D.	*
AT5G58090	O-Glycosyl hydrolases family 17 protein	1,3-beta-glucosidase	*	apoplast	*	*	*	*	*
AT4G12420	Cupredoxin superfamily protein	SKU5	*	apoplast	*	*	*	N.D.	*
AT3G25290	Auxin-responsive family protein	Cytochrome b561	*	PM	N.D.	N.D.	*	N.D.	N.D.
AT5G26340	Major facilitator superfamily protein	STP13	*	PM	*	*	*	N.D.	N.D.
AT5G08080	syntaxin of plants 132	SYP132	*	membrane traffic	*	*	*	N.D.	N.D.
AT3G52400	syntaxin of plants 122	SYP122	*	membrane traffic	*	*	*	N.D.	N.D.
AT5G46700	Tetraspanin family protein	TET1, TRN2	*	TET	N.D.	N.D.	*	*	N.D.
AT1G31340	related to ubiquitin 1	RUB1	*	Cytoplasm	N.D.	N.D.	N.D.	N.D.	N.D.
AT2G27260	Late embryogenesis abundant (LEA)	LEA	*	Cytoplasm	N.D.	N.D.	*	N.D.	N.D.
AT2G18960	H[+]-ATPase 1	AHA1	*	PM	*	*	*	N.D.	*
AT2G47170	Ras-related small GTP-binding family protein	ARF1	*	membrane traffic	N.D.	N.D.	*	N.D.	N.D.
AT4G25240	SKU5 similar 1	SKS1	*	apoplast	N.D.	N.D.	*	*	*
AT5G25460	transmembrane protein 2C putative	DGR2	*	PM	N.D.	*	*	N.D.	*
AT2G16600	rotamase CYP 3	CYP19-1	*	Cytoplasm	*	N.D.	*	N.D.	N.D.
AT3G52470	Late embryogenesis abundant (LEA)	LEA	*	Cytoplasm	N.D.	N.D.	*	*	N.D.
AT5G42980	thioredoxin 3	TRX3	*	enzyme	*	N.D.	*	N.D.	N.D.
AT2G44790	uclacyanin 2	uclacyanin 2	*	apoplast	*	N.D.	*	N.D.	*
AT1G74010	Calcium-dependent phosphotriesterase	phosphotriesterase	*	enzyme	N.D.	N.D.	N.D.	*	N.D.
AT2G16060	hemoglobin 1	AHB1	*	apoplast	N.D.	N.D.	*	N.D.	N.D.
AT5G19390	Rho GTPase activation protein (RhoGAP)	ROPGAP7	*	membrane traffic	N.D.	N.D.	N.D.	N.D.	N.D.
AT3G11820	syntaxin of plants 121	SYP121	*	membrane traffic	*	*	*	N.D.	N.D.
AT3G02740	Eukaryotic aspartyl protease family protein	aspartyl protease	*	apoplast	*	N.D.	N.D.	N.D.	*

B

	Number of Identified proteins	Cultured cell EV	Seedling EV Rutter et al. 2017	Seedling EV He et al. 2021	Plasmodesmata Fernandez et al. 2011	Plasmodesmata Brault et al. 2019	Apoplast Uemura et al. 2019
Cultured cell EV	1356	N. A.	8.2% (111/1356)	24.1% (326/1356)	23.5% (318/1356)	5.7% (77/1356)	44.6% (604/1356)
Seedling EV Rutter et al. 2017	204	54.5% (111/204)	N. A.	75% (153/204)	55.4% (113/204)	3% (6/204)	66.7% (136/204)
Seedling EV He et al. 2021	981	33.3% (326/981)	15.6% (153/981)	N. A.	37.7% (370/981)	1.8% (17/981)	55% (539/981)
Plasmodesmata Fernandez-Calvino et al. 2011	1260	25.2% (318/1260)	9% (113/1260)	29.4% (370/1260)	N. A.	4.8% (61/1260)	41.0% (517/1260)
Plasmodesmata Brault et al. 2019	115	67% (77/115)	5.3% (6/115)	14.8% (17/115)	53% (61/115)	N. A.	30.5% (35/115)
Apoplast Uemura et al. 2019	3219	18.8% (604/3219)	4.3% (136/3219)	16.8% (539/3219)	16.1% (517/3219)	1.1% (35/3219)	N. A.

80　60　40　20　0 (%)

Figure 3. Comparison between the EV proteome and the plasmodesmata proteome
A) Comparison of the EV proteomes and the plasmodesmata proteomes in terms of the top 30 high-intensity proteins in the cell culture EV proteomes. **B)** Overlap percentages between the three EV proteomes and the two plasmodesmata proteomes. The EV and plasmodesmata fractions were indistinguishable in these analyses.

Figure 4. Localization of PDLP3-GFP in the apoplast region
A-D) Confocal microscopy images of PDLP3-GFP. Plasma membrane was labeled by Plasmembrane Red. PDLP3-GFP clearly localized to plasmodesmata (**A**, white arrowheads). At high contrast images, the PDLP3-GFP signals were also observed as particles at inside (**B**), plasma membrane (**C**), and surface (**D**). Some PDLP3-GFP signals were detected as small particles in the apoplast region (**D**, white arrowheads). Bar, 10 μm.
E, F) Projection images from PDLP3-GFP time-lapse analyses. The 15 time-lapse images were acquired every 5 seconds and grouped into early (1-5), middle (6-10), and late (11-15) phases. These images were projected at maximum intensity, and superimposed into one merge image using blue, green and red colour, respectively. Bars, 10 μm. **G)** Comparison of PDLP3-GFP and TagRFP-SYP121 at the apoplast region. TagRFP-SYP121 (white arrowheads) were not colocalized with PDLP3-GFP (yellow arrowheads). Bar, 5 μm.

Figure 5. Iodixanol density gradient ultracentrifugation of the crude EV fraction

A) Separation of the crude EV fraction by an iodixanol density gradient ultracentrifugation. The crude EV fraction was mounted on the top of the iodixanol solution during the density gradient ultracentrifugation (24%, 12%, 6%, and 3% iodixanol solutions). After the ultracentrifugation, six fractions were recovered and used for proteome analyses. **B)** Classification of the 524 identified proteins on the basis of their distribution. The presented data correspond to the information provided in Supplemental Table 5.

Figure 6. Representative proteins in each gene ontology category

A) Representative EV proteins. B) Representative PD proteins. C) Membrane trafficking-related proteins. D) Membrane transporters and pumps. E) Disease resistance-related proteins. F) Oxidases and peroxidases. G) O-glycosyl hydrolases. The presented data correspond to the information provided in Supplemental Table 5. cl. indicates the cluster number in Figure 5B; cat. indicates category.

Supplemental Figure 1. Growth condition of MM2d Arabidopsis cultured cells for this study

A) Growth curve of MM2d Arabidopsis cultured cells in this study. Supernatant of 5 days-old and 7 days-old cultured cells was applied for EV purification. **B)** DNA degradation assay of MM2d cells. In 5 days-old and 7 days-old cultured cells, DNA degradation did not occur at all.

Supplemental Figure 2. Size distribution of EVs in each fraction after an iodixanol density gradient ultracentrifugation.

Six fractions were recovered after an iodixanol density gradient ultracentrifugation and used for size distribution analyses. These fractions correspond to the fractions in Figure 5A.