SlimVar: rapid *in vivo* single-molecule tracking of chromatin
regulators in plants

Alex L. Payne-Dwyer[1,2], Geng-Jen Jang[3], Caroline Dean[3], and Mark C. Leake[1,2]*

[1] School of Physics, Engineering and Technology, University of York, YO10 5DD, UK.

[2] Department of Biology, University of York, YO10 5DD, UK.

[3] Cell and Developmental Biology, John Innes Centre, Norwich Research Park, NR4 7UH, UK.

*Correspondence: mark.leake@york.ac.uk

Abstract

Epigenetic regulation occurs over many rounds of cell division in higher organisms.
However, visualisation of the regulators *in vivo* is limited by imaging dynamic molecules
deep in tissue. We report new technology—Variable-angle Slimfield microscopy (SlimVar)—
that enables tracking of single fluorescent reporters to 30 µm depth through multiple
Arabidopsis thaliana root tip cell layers. SlimVar uses rapid photobleaching to resolve
tracked particles to molecular steps in intensity. By modifying widefield microscopy to
minimise optical aberrations and robustly post-process few-photon signals, SlimVar
mitigates performance losses at depth. We use SlimVar to quantify chromatin-protein
assemblies in nuclei, finding that two homologous proteins key to epigenetic switching at
FLOWERING LOCUS C (*FLC*) —cold-induced VERNALISATION INSENSITIVE3 (VIN3) and
constitutively expressed VERNALISATION 5 (VRN5)—exhibit dynamic assemblies during *FLC*
silencing. Upon cold exposure, the number of assembly molecules increases up to 100% to a
median of ~20 molecules. Larger VRN5 assemblies preferentially colocalise with an *FLC lacO*
transgenic reporter during prolonged cold and persist after return to warmth. Our findings
support a hybrid model of epigenetic memory in which nucleation of histone trimethylation
is assisted by dynamic protein assemblies over extended durations. SlimVar offers molecular
insights into proteins expressed at physiological levels in tissues.

Introduction

Understanding the basis of epigenetic silencing remains a major question, with potential
implications for both novel medical therapeutics [1] and agricultural biotechnology [2]. It is not
yet known how cells process delocalised, long-term sensory information into genetic states
that are stable enough to facilitate not only cellular differentiation but also determine key
organism-level transitions, including those relevant to medicine such as ageing and
disease [3]. Similarly, mysteries remain regarding long-term epigenetic responses to key
environmental cues, including seasonal temperature fluctuations, govern the productivity
and resilience of crops to challenges such as climate change [4].

A key conserved epigenetic mechanism across eukaryotes - including humans and plants - is
Polycomb-mediated silencing [5], which involves modification of trimethylated histone H3
lysine 27 (H3K27me3). H3K27me3 initially accumulates at a nucleation site then spreads

across a locus to stably maintain a silenced state through many rounds of cell division [6,7].
Polycomb Repressive Complex 2 (PRC2) silencing has been well studied in *Arabidopsis* gene
FLOWERING LOCUS C (*FLC*) [8]. *FLC* encodes a repressor of flowering and is epigenetically
silenced during a process called vernalisation by the prolonged cold of winter; lowering
repressor levels enables flowering in spring [9–11]. Cold exposure increases the probability that
each *FLC* locus will epigenetically switch from ON to OFF states through nucleation of
H3K27me3 at an intragenic site [12,13]. Upon return to warm conditions, H3K27me3 spreads
across the locus to give long-term stable silencing [14]. What has been less clear is how the
relative stability of the nucleated state is inherited following cell division. Each nucleation
event involves only three nucleosomes, too few to survive random replicative dilution
through the classic Polycomb 'read-write' mechanism [11,15,16]. Metastable protein assemblies
have been proposed to explain the inheritance of the nucleated state [17]. PRC2 accessory
proteins are thought to restore the stochastic loss of histone marks by stimulating Polycomb
Repressive Complex 2 (PRC2) activity [18]. While individual proteins may only interact
transiently with nucleation factors and the locus, a recent model predicts that an assembly -
with the appropriate positive cooperativity – can become dynamically self-sustaining above
a threshold number of recruited proteins [17]. In this framework, a sufficiently large assembly
of proteins could act as a binary memory element working together with established
H3K27me3 machinery at a given locus. This model highlights molecule number as a key
factor.
61
Two PRC2 accessory proteins required for stable cold-induced silencing at *FLC*,
VERNALISATION INSENSITIVE3 (VIN3) and VERNALISATION 5 (VRN5, also known as VIN3-LIKE
1 / VIL1), expressed in shoot and root tips are clear candidates for this form of memory
storage [19]. VIN3 and VRN5 (collectively called VEL proteins) associate with the PRC2 complex
and with the *FLC* nucleation region that accumulates H3K27me3 specifically during cold
conditions [11,20]. At <15°C, *VIN3* expression gradually rises over several weeks, while in warm
conditions >15°C, expression decays rapidly within approximately 4 h [9,10,20]. VIN3 and its
assemblies could therefore in principle report the duration of cold conditions during winter
to promote an epigenetic switch. However, due to its instability in the warm, one or more
additional factors—a promising candidate being VRN5—are required at the *FLC* nucleation
region to explain persistent memory of silencing following a change to warm conditions.
Both VIN3 and VRN5 contain complex plant homeodomains (PHDs) [19] which do not interact
directly with histone tails [21], FNIII domains and C-terminal VEL domains [22,23]. Of these three
domains common to VIN3 and VRN5, the VEL domain mediates head-to-tail interactions.
VEL proteins that oligomerise and even form phase-separated droplets under transient
overexpression are directly associated with stable silencing at *FLC* [23]. A key question is then:
do physiological levels of VIN3 and VRN5 oligomerise sufficiently in the vicinity of *FLC* during
vernalisation, to fulfil this model?
80
Here, we describe development of an efficient new technology, Slimfield Variable Angle
(SlimVar), and apply it to rapidly track and quantify VEL proteins in live plant tissue. Prior
investigation of nuclear oligomers has been constrained by an inability to detect individual,

rapidly diffusing protein molecules deep within plant tissue. Traditional confocal [24],
structured illumination [25] and lightsheet microscopy [26] lack the required combination of
sensitivity and speed for single-molecule tracking in live plants, which remains a challenge
beyond the first cell layer [27–34]. Although complex, expensive super-resolution methods
including lattice lightsheet [35,36] and MINFLUX [37,38] are in principle capable of deeper imaging,
to date neither has been successfully applied at a molecular scale in plants. In addition,
these require both specialised hardware and/or photoactivatable/photoswitchable
fluorescent proteins [39,40] or dyes [41,42].

SlimVar, instead uses common fluorescent protein fusions in existing transgenic plants
without requiring overexpression, as well as a relatively accessible microscope platform
compared to other single molecule techniques. Adapted from Slimfield microscopy,[43–45]
SlimVar uses a grazing angle of incidence optimised for observing single nuclei up to
approximately 30 μm deep in root tissue, while tracking single fluorescent protein
molecules at up to 40 nm lateral spatial precision. Whereas Slimfield has been limited to
simple unicellular samples or the surfaces of *ex vivo* tissue sections [46], SlimVar enhances the
image contrast and mitigates optical aberrations at depth to enable dynamic spatial
localisation in complex multicellular samples. The use of root tips confers optical advantages
of lower autofluorescence and more regular, less refractive layers than other plant tissues.
Single-molecule sensitivity is combined with stepwise photobleaching analysis [43] to quantify
the number of molecules in any observed oligomeric assemblies. This sensitivity simplifies
the use of transgenic plant lines expressing from as few as a single gene copy and enables
imaging of physiological states of low-abundance nuclear proteins.

We first introduce the principles and operational procedures of SlimVar, and demonstrate
its measurement capabilities. Applying to it root tips, we find that both VIN3 and VRN5
proteins form assemblies in cell nuclei, composed of consistent dimeric subunits. In lines
expressing from single-digit exogenous copies of VIN3 and VRN5, the median assembly
comprises up to ~20 molecules of each protein, in agreement with that required for protein
memory elements predicted from modelling [17]. We also use an *FLC-lacO*/LacI-YFP transgenic
reporter to localise the *FLC* locus and characterise its mobility relative to VEL proteins.
Finally, we demonstrate dual-colour SlimVar, which directly shows VRN5 assemblies present
at *FLC*. We find that larger VRN5 assemblies preferentially colocalise with *FLC* after long cold
exposure and after return to warm conditions, and that this interaction between individual
larger oligomers of VRN5 and *FLC* is dynamic on a sub-second timescale.

Results

SlimVar enables molecular quantification of diffusing particles deep in tissue

SlimVar is a 2D + time imaging technique which identifies fluorescent **foci** from local
intensity maxima in each video frame (Fig. 1, Table 1). Foci correspond to at least one, or
more generally, of a localised group of labelled molecules much smaller than the widefield
resolution limit; output parameters include 2D spatial location, total intensity in photons,
and **signal-to-noise ratio** (SNR). If the frame rate is rapid enough to overcome motion blur,
foci from sequential frames can be linked into **tracks**. Since the molecules in a track are
spatially correlated we can infer that they form an intermolecular **assembly**, and that each
track corresponds to one assembly.

Rapid tracking of single diffusive proteins requires minimal background (equivalent to $<10^7$
photons per μm^2/s), and a photon-sensitive, low-noise detector (<2 electrons readout per
pixel) with millisecond or less sampling. Maximising the emissive output of fluorescent
protein tags in these short exposures requires high excitation intensity (~kW/cm^2).

Photobleaching dominates under these conditions, which SlimVar uses to capture the
emission intensities of both

i)   the initial unbleached state of each assembly, and
ii)   its constituent single fluorophores,

in the same short (~10 s) acquisition.

The ratio of these—**stoichiometry**—is an estimate of the number of molecules in each
assembly. The numerator in this ratio is the track's initial intensity; its relative uncertainty is
low since the first of its foci contains many photons. The uncertainty can be reduced further
by interpolation from fitting the photobleach trend in time.

The denominator in the ratio relates to single molecule detection events, each containing
few photons; there may be very few or no such events that can be associated with a
particular assembly, for example if it leaves the detection volume. Instead, we consider the
population-average number of photons associated with each fluorophore—the
characteristic molecular brightness. It can be determined by averaging the height of
individual photobleaching steps [43] (Supplementary Fig. 1). As an average, it is only a reliable
estimate for individual assemblies if the intensities are narrowly distributed for fluorophores
of the same type in the same environment, and independent of concentration. These
conditions are well satisfied by intracellular fluorescent protein fusion constructs, which do
not easily self-quench [47,48]. These intensities represent photons per frame and, if not in a
saturating regime for emission, scale with the exposure time, excitation power and the
photon collection efficiency of the microscope. However, the characteristic molecular
brightness can be internally calibrated within each sample or dataset for a ~20 μm range of
working depths. For this reason, it is best practice to acquire additional frames until an
excess of independent single-molecule events has been accrued.

It is possible to estimate stoichiometry by counting photobleaching steps directly,[49] but this
strategy is typically limited to roughly <10 steps per track due to increasing likelihood of
stochastic missing/overlapping steps for higher stoichiometry assemblies. Our ratiometric
method instead estimates the stoichiometry of tracks which begin close to the start of the
acquisition, typically within a quarter of a photobleaching decay time or less, thus

containing minimal prior photobleached content. This approach enables accurate counting
over a broader range of stoichiometries than direct step detection.

Independent of the tracking pipeline, SlimVar can quantify the **total protein number** in the
detection volume, or sub-region such as a section of the cell nucleus. We sum all the initial
pixel intensities (measured as number of photons) in segmented region of each image and
normalise these values by the characteristic brightness corresponding to a single fluorescent
dye tag. We define the total protein number as the difference in mean integrated intensity
between the test population and that of the unlabelled negative control to corrected for any
autofluorescence. We interpret this as the average number of labelled molecules in the
volume. It is a reliable population-level measure of protein content and concentration.[45]

Since each assembly is associated with a sequence of discontinuous steps along its track, we
also quantify the assembly's mean **diffusivity** using mean-square displacement analysis
(Methods). While photobleaching shortens the average track, it transiently improves the
optical contrast which is ideal for rapid, high-content tracking. The diffusivity can be used to
determine if the assembly is immobile on the timescale of the experiment and therefore
likely to be bound, for example to chromatin.

In multicolour SlimVar, multiple channels are captured and tracked independently from the
same acquisition. Once the two channels are spatially coaligned, the tracks' spatial and
temporal coordinates are compared. This enables measurement of dynamic **colocalisation**
of pairs of tracked particles such as different types of protein that are labelled with different
colour dyes. This is a powerful correlative approach which can test the dependence of
colocalisation on metrics associated with each assembly, such as stoichiometry and
diffusivity.

Fig. 1. Correlative quantification of diffusing assemblies using SlimVar. Terms are defined

in Table 1. SlimVar delivers a) rapid photobleaching in image sequences at high framerate to

outpace molecular diffusion, followed by b) robust postprocessing and quality control steps

to identify foci in individual frames, and tracks across multiple (up to 20) frames, which

correspond in general to assemblies of labelled molecules. c) The full extent of

photobleaching enables estimates of the characteristic molecular brightness, which is

narrowly distributed for a fluorescent protein. The characteristic molecular brightness is

used to determine d) total protein number for each region of interest and e) stoichiometry

for each tracked assembly near the start of the image sequence, as number of molecules.

These metrics are corrected for autofluorescence using unlabelled wild type then collated

over a population, enabling robust estimation for average total protein number. f)

Periodicity analysis extracts patterns from the stoichiometry distribution to infer consistent

repeat units of assemblies. g) Rapid tracking facilitates mean-square displacement analysis

to estimate individual assembly mobility. h) Multicolour SlimVar assesses whether

stoichiometry and diffusivity are dependent on colocalisation between different pairs of

assemblies.

Fig. 2. SlimVar enhances optical contrast at greater working depths. The optical scheme for SlimVar adapts widefield or objective-based TIRF (total internal reflection fluorescence) microscopy capable of detecting single molecules at a coverslip surface, and extends this to greater working depths. A narrow, collimated excitation beam is delivered at a steep but subcritical angle by (1) adjusting the position of a steering lens. The intersection of the focal plane and excitation beam defines a sub-micron high detection volume at the set working depth. The lateral size of this volume can be (2) adjusted using an iris or beam stop to match sample dimensions and reduce background. Aberrations, inherent to oil immersion lenses at depth, are mitigated at the set working depth using a calibration procedure. Either a test sample or an *in vitro* beads-in-agarose phantom may be used. This comprises a combination of adjustments to (3) an objective lens correction collar and, where necessary, (4) shifting the tube lens towards the objective. The microscope uses a single detector with two colour channel splitter (Cairn OptoSplit); note, the beam is not stopped down after entering the splitter and is shown here with a narrowed diameter only for clarity. In multicolour experiments, contrast is protected from channel crosstalk by (5) alternating excitation wavelengths between subsequent frames. The second pair of lenses in the detection path provides (6) additional magnification (1.2–2.2× depending on physical sensor pixel size) to ensure the point spread function (PSF) is spatially oversampled for super-resolved localisations.

Implementation of single-molecule photobleaching analysis requires the maximum optical contrast, while maintaining both sampling rate and signal-to-noise sufficient to detect foci. The widefield configuration (Fig. 2) maximises fluorescence emission collection from foci within each detection volume. It uses the full back aperture of the objective lens and a minimal number of optical components in the detection path. SlimVar adjusts these

elements to mitigate loss of optical contrast at high working depths. These adjustments
address either the fluorescence excitation, or the aberrations (blurring of static objects) in
the emission path.

The optimal excitation intensity is a few kW/cm^2 (~10 mW total power within a beam of
approximately 25 µm diameter) which maximises fluorescence emission, whilst being sub-
saturating for the dyes and ensuring photobleaching is not too rapid relative to the sampling
time. The contrast can be further improved by minimising out-of-focus background. SlimVar
avoids exciting the sample outside the detection volume by employing a narrow beam of
near-collimated illumination delivered at an oblique-angle similar to variable angle
epifluorescence microscopy (VAEM) [50] and highly inclined illumination (HILO) [51,52]. The
excitation laser(s) are coaligned and focused into the back focal plane of the high numerical
aperture objective lens. To progressively tilt the beam in the sample, this focus is moved to
a precise lateral displacement from the optic axis using a steering lens (Fig. 2, point 1)
placed conjugate to the back focal plane. The maximum shift of ~3 mm radius would
generate a large tilt for TIRF) excitation (Fig. 2, inset); for SlimVar, the beam is instead
shifted by 2.3 ± 0.2 mm corresponding to a free beam tilted to 60 ± 5 $^{\circ}$ at the sample,
subcritical for TIRF. The beam is then stopped down to match the sample dimensions (Fig.
2, point 2) which eliminates further background unrelated to the region of interest.

At low working depths, aberrations in the emission path are corrected by design for most
high numerical aperture lenses. However, scattering and aberrations emerge at higher
working depths; of these, the most detrimental is spherical aberration caused by differences
in refractive index between the immersion medium and the aqueous sample, that
compounds rapidly with additional working depth. While in principle SlimVar admits the use
of water immersion lenses (NA < 1.3) that minimise this effect, here we adapt the widefield
microscope to make more efficient use of a higher numerical aperture oil immersion
objective lens with a correction collar (NA = 1.49). These adaptations compensate the
optical aberrations at a desired working depth. Using a precision microscope stage, the
working depth is set by difference from the axial position where the coverslip surface
appears in focus. We present a **calibration procedure**, inspired by previous works
improving optical trapping[53–55] or imaging [56,57] using high numerical aperture in aqueous
samples. It adjusts the optical pathway—including beam collimation, objective correction
collar setting and tube lens position—to minimise aberrations at a representative depth of
25 µm (Fig. 2, points 3-4). This calibration improves the optical contrast sufficiently for
imaging and tracking across the range of working depths up to 30 µm (Supplementary Figs.
2 and 3). This procedure can be performed either with a live root sample resting on the
coverslip, or a phantom using beads suspended in agarose (Methods). We assessed the
improved optical contrast using the width of the point spread functions (PSFs)
(Supplementary Fig. 3) and quantified an effective numerical aperture of 1.38 ± 0.02, which
though reduced from the diffraction-limited performance, still exceeds that of a water
immersion objective lens. Both axial and lateral resolution are improved for beads, and the
severe loss of axial performance associated with depth is mitigated in plant roots.

For multicolour experiments, multiple continuous wave lasers are spatially filtered and
expanded to the same dimensions, then coaligned using dichroic mirror beamsplitters.

Crosstalk and bleed-through effects tend to decrease the contrast or introduce ambiguous signals; to avoid this we excite and subsequently analyse each channel in alternating interleaved frames (Fig. 2, point 5). This proportionally decreases the effective sampling rate but otherwise maintains the tracking performance of single-colour imaging.

Spatial oversampling—a small pixel size relative to the widefield spatial resolution—is necessary for postprocessing. It is introduced by additional magnification in the detection path (Fig. 2, point 6). Although each signal is spread more thinly over a greater number of pixels, the lower chance of spurious correlations makes detection more robust to variations in foci shape due to defocus or motion. Oversampling also ensures the localisation precision is not limited by pixel size.

To mitigate detrimental effects of background noise and spatial overlap on the final metrics, **sifting** is performed to only accept tracks above a minimum set signal-to-noise ratio and track length. Using a dedicated single molecule assay *in vitro* (Supplementary Fig. 2a-b) and our tracking data from live plants, we found appropriate sifting thresholds determined by two factors: detector noise (Supplementary Fig. 2c) and autofluorescence relative to the probe's characteristic molecular brightness (Supplementary Fig. 4). Sifted tracks have a positive predictive value >95% for single YFPs *in vitro* (Supplementary Fig. 2d) and >90% for YFP-labelled assemblies in plant roots (Supplementary Fig. 2e).

We assessed the combined improvements to optical and sifting performance using the characteristic molecular brightness of the fluorophore and the total number of photons collected per track. The increase in apparent characteristic molecular brightness with excitation power (Supplementary Fig. 2f), indicates that the detected photon flux associated with single molecule detection at 25 μm depth *in vitro* is only >50 photons per frame, and therefore >150 photons per single molecule track. Our practical values for characteristic molecular brightness fall in the range 70-200 photons per frame. The total flux emitted is much larger (~1,000 photons/ms), with many photons producing the raw image but fewer photons captured in tracks. This is further illustrated by comparing the numbers of photons per track for *in vitro* control and *in vivo* data (Supplementary Fig. 2g-h). Modes of 190–400 photons per single molecule track are observed, up to a maximum of ~5,000 photons per single molecule. This accounts for ~5% of a typical single fluorescent protein molecule budget of 10^5 photons [58,59]. This net transmittance corresponds reasonably well to the expected 10%, accumulated from transmittance of the sample (~80% in agarose, ~60% in root tips), effective numerical aperture (29% solid angle), detection optics (~60%) and detector (fill factor >99%, quantum efficiency ~92%). Most tracks have a photon total an order of magnitude lower, primarily due to the truncating effect of photobleaching (which is accelerated at the high excitation irradiance required), or simply by diffusion out of the detection volume. So, while the observed photon counts *in vivo* are low compared to theoretical limits of photon budgets for bright fluorescent proteins, they are reasonable considering a realistic budget subject to diffusive, photobleaching and scattering losses.

The fastest diffusivity that SlimVar can detect lies in the range 5–30 μm/s^2 depending on sampling rate. Only single proteins, if any, exhibit this diffusivity in an intracellular context, and these can still be inferred from the total protein number. Importantly, this suggests that oligomeric assemblies and associated metrics remain representative of the underlying

population through sifting, even in cases where the direct detection rate of single molecules is reduced, or where the tracks are truncated. Subject to these limitations, SlimVar is therefore suitable not only for detecting single molecules in plant tissues but also counting molecules within assemblies (stoichiometry and periodicity), and measuring their mobility (diffusivity) and interactions (colocalisation) simultaneously.

SlimVar resolves dynamic single assemblies of VIN3 and VRN5 in plant nuclei

For examining VIN3 and VRN5 protein localisation and self-assembly behaviours, during and after cold treatment, we utilised lines with VIN3-EGFP/*vin3-4 FRI* [11,60] or VRN5-EYFP/*vrn5-8 JU223* [11,19] (referred to as VIN3-GFP and VRN5-YFP respectively) with variable transgene copy numbers among progeny due to genetic segregation. To investigate the effects of protein expression and to exploit greater imaging contrast and lower autofluorescence associated with yellow and red fluorophores, we later also characterised new lines containing VIN3-SYFP2/Col*FRI* (Supplementary Fig. 5) or VRN5-mScarlet-I/*vrn5-8 FRI* (Supplementary Fig. 6) with different numbers of transgenes. These lines include active *FRI* alleles so require effective vernalisation for flowering [20].

As a benchmark, we performed traditional confocal microscopy imaging on whole roots (Fig. 3a) with typical sampling at 35 s per frame (Fig. 3b and Supplementary Fig. 7). After identifying nuclei from transmitted light images, we found that both VIN3 and VRN5 exhibited bright but largely diffuse fluorescence localised to the nucleoplasm. We determined the qualitative autofluorescence from the Col*FRI* negative control; the autofluorescence is greater under 488 nm wavelength excitation but this line shows no signal localisation in either channel (Supplementary Fig. 7). While VRN5 was detectable above this unlabelled background in all nuclei (N=241) at all timepoints, VIN3 is only discernible during the cold period itself. Its total brightness in nuclei decreased at subsequent timepoints after this, being undetectable within one week after return to warm conditions, as reported previously [10]. During cold, the VIN3 signal per cell was initially greatest in the vicinity of the meristem and epidermis, before becoming brighter in all cells after further cold exposure (Supplementary Fig. 7).

We then performed Airyscan, an enhanced form of confocal laser scanning microscopy which uses a point detector array; after optimising the point dwell time, field of view and laser irradiance, we obtained faster frame sampling times down to 60 ms for individual root tip nuclei enabling attempts at video tracking [61]. These image sequences showed a marginally more granular spatial patterning than standard confocal microscopy, hinting at the presence of distinct foci within the diffusive fluorescence (Fig. 3c), with residence times in a similar range to the sampling time (Fig. 3d).

Fig. 3. SlimVar resolves dynamics of VIN3 and VRN5 assemblies during cold exposure of root tips. a) Whole roots mounted horizontally in aqueous media between agarose and coverslip, suitable for confocal and SlimVar microscopy. b) Projections of confocal z-stacks of root tips after 6 weeks of cold with VRN5-YFP reporter; acquisition time 35 s. Insets (interpolated) show labelled protein signals consistently localised to the nucleoplasm but not the nucleolus. Nuclear patterning of VEL proteins (c.f. confocal images in Supplementary Fig. 7) was typically round or lens-shaped, with a median length 7.8 µm (interquartile range IQR: 5.7–10.3 µm, N=571), and aspect ratio of 1.16 (IQR: 1.06–2.10), comparable to other nuclear reporters [62]. c,d) Airyscan images of VRN5-YFP after 2 weeks' cold indicating heterogeneous spatial distribution; c) maximum intensity projection of three *z*-slices, averaged over three consecutive timepoints; d) residence times over the same timepoints estimated from ratio between median and standard deviation of pixel values across the three frames. Low standard deviation (cyan) indicates low foci displacement over the three frames (200 ms) equivalent to diffusivity <0.1 µm^2/s, while high standard deviation (magenta) indicates high displacement over 70 ms, or diffusivity >0.3 µm^2/s. e) Schematic indicating illumination and detection volumes and working depth specific to SlimVar of root tip nuclei. f-k) SlimVar images of a root tip labelled at VRN5-YFP before vernalisation. f) Brightfield, used to identify nuclei in the field of view; g) initial frame in YFP channel prior to photobleaching, with nucleolus indicated (white dashes) showing background of overlapping signals; h) early stage photobleaching revealing distinct assemblies due to transient increase in background contrast (mean projection of frames 4–6); i) SlimVar resolves assemblies of differential mobility on ms timescales shown here as distinct slow-(cyan, >60 ms residence time, diffusivity <0.4 µm^2/s) and fast-moving (magenta, <20 ms residence time, diffusivity >1.4 µm^2/s) objects, represented by pixelwise ratio of median and standard deviation projections over three frames. j) Foci are detected from local maxima to super-resolved localisation precision. All sifted foci (Methods) for full sequence shown superimposed (white circles) on image from panel h (greyscale); k) tracks, generated by linking similar nearby foci, indicate individual assemblies with independent estimates of

stoichiometry and diffusivity. All sifted tracks from entire sequence shown superimposed on image from panel with one vertex per timepoint (white arrows).

We then implemented SlimVar (Fig. 3e-k) capable of single-molecule fluorescent protein detection at sub-ms levels, by first identifying nuclei in brightfield (Fig. 3f) to avoid premature photobleaching. We found qualitatively similar nucleoplasmic morphology and localisation (Fig. 3g), but instead of diffuse fluorescence we observed multiple, highly mobile, distinct particles (Fig. 3h) with residence times longer than the exposures, consistent with a sensitivity and sampling speed sufficient to overcome motion blur (Fig. 3i). We thereby detected protein assemblies as distinct foci (Fig. 3j) and connected them into tracks (Fig. 3k).

The excitation beam encapsulates individual cell nuclei between 4–16 µm wide but with minimal excitation of the remaining >70% of the cell volume. It is ideally aligned for the target root tip cells in at least three surface cell layers overlaying the stem cell niche (Fig. 3a and Supplementary Fig. 8), with associated reduction in aberration, backscatter and out-of-focus fluorescence excitation of intermediate cell layers. The contrast available for imaging and tracking indicates that, in principle, cells of different types can be quantitatively discriminated. While SlimVar does not fully recover diffraction-limited resolution in images (Supplementary Fig. 3), the net result is an improvement in our signal-to-noise metric for tracking VRN5 fluorescent reporters in assemblies by a factor of ~2.6 relative to epifluorescence microscopy, including adjustment for the faster exposures (Supplementary Fig. 2i, Methods). The capability for single-molecule detection is comparable with control samples of purified fluorescent protein (Supplementary Fig. 2). The characteristic molecular brightness is sufficiently consistent across the range of working depths to collate acquisitions for each line.

Vernalisation induces upregulation and self-assembly of VIN3 and VRN5

The visual changes in VIN3 and VRN5 fluorescence during cold exposure suggest corresponding changes in nuclear expression. To test this quantitatively, we acquired SlimVar datasets from nuclei in VRN5-YFP and VIN3-GFP lines. From these, we first determined the characteristic molecular brightness for the GFP and YFP tags (Methods, Supplementary Fig. 1). As a negative control, we imaged nuclei in the Col*FRI* line using the same 488 nm and 514 nm wavelength excitation modes (Supplementary Fig. 4). We then estimated the total protein numbers in each nucleus of VRN5-YFP and VIN3-GFP (Fig. 4a).

VRN5 was highly abundant at all timepoints, with levels an order of magnitude greater than VIN3 in the VIN3-labelled lines (Fig. 4a). Total VRN5 approximately doubles in response to full vernalisation and persists after return to warm conditions. Total protein numbers translate to nucleoplasmic concentrations of ~100 nM–1 µM for VIN3 and 1–10 µM for VRN5 (Methods). When applied to the cell cytoplasm, the high sensitivity of SlimVar was also able to establish that the fluorescence signals for both VIN3 and VRN5 were marginally above Col*FRI* negative control levels, equivalent to a concentration at least 10,000-fold less than those measured in the nucleus.

431
432

Fig. 4. Cold exposure causes VIN3 and VRN5 to form higher stoichiometry assemblies, but only VRN5 assemblies become more numerous. a) Distributions of integrated nuclear intensity (total number of labelled molecules per nucleus prior to correcting for autofluorescence) of individual nuclei collated from cells imaged at working depths of 20 μm ± 10 μm at different timepoints before, during and after vernalisation, for VIN3-GFP and VRN5-YFP: NV = not vernalised; V2W = two weeks of cold; V6W = six weeks of cold; V6W+T7 = six weeks of cold followed by one week of warm conditions; V6W+T14 = six weeks of cold followed by two weeks of warm. The total protein number is calculated as the excess of the integrated nuclear intensity above the mean level of the autofluorescence background in the negative control line, Col*FRI* (horizontal line). Bar, box and whisker denote median, interquartile range (IQR) and 1.5× IQR respectively; cross: mean ± sem. The abundance of VIN3 protein was insignificant prior to vernalisation (Brunner-Munzel (BM) test vs Col*FRI*, N=66, p=0.11 |ns: not significant at adjusted p<0.01). However, the VIN3 total protein number increases sharply to ~28,000 ± 3,700 molecules above control after 2 weeks cold (N=64, p=0.0031 |*), and peaks at ~44,000 ± 4,700 after 6 weeks cold (N=83, p<0.001 |**). Following transfer to warm conditions, VIN3-GFP levels reduce to ~3,200 ± 1,600 molecules within 7 days (N=74, p=0.04 |ns).   VRN5 levels increase between non-vernalised and post-vernalised timepoints from ~110,000 ± 23,000 to ~190,000 ± 37,000 molecules (N=94, p=0.0089|*). b) Distributions of number of tracks per nucleus (binned in columns of width 2 for clarity). The number of VRN5 tracks increases initially (NV: 20.8 ± 1.9 up to 26.8 ± 1.6 tracks per nucleus at 2 weeks cold; N=86, p=0.0054|*) and this increase is retained after 6 weeks' cold (27.0 ± 1.5 and 26.2 ± 2.6 tracks per nucleus at 6 weeks cold and 14 days post-cold respectively; N=94, p=0.80|ns); c) Collated distributions of stoichiometry (number of labelled molecules per nuclear assembly) of individual tracks at different timepoints before, during and after vernalisation; *nt*: no tracks detected.

We also explored the effect of transgene copy number on the abundance of VEL proteins
during cold exposure. We generated a homozygous single transgene copy line of VIN3-
SYFP2, though not in a deletion background, meaning endogenous VIN3 is also present at a
similar level (Supplementary Fig. 5a). The expression of exogenous VIN3 still follows the
expected pattern (Supplementary Fig. 7) but at much lower levels than in VIN3-GFP,
reflecting the reduction to a single transgene copy of *VIN3*. We used SlimVar to estimate
the characteristic molecular brightness for SYFP2 to quantify the total protein number in the
VIN3-SYFP2 line (Supplementary Fig. 11). Considering only the labelled VIN3 in the SYFP2
line, the total protein number is 31 ± 4 % of that of the VIN3-GFP line at both of the two
timepoints. Accounting for the unlabelled copy, this rises to 62 ± 11 %. This is consistent
with the VIN3-GFP line having three transgene copies, with each copy generating the same
amount of protein as the endogenous *VIN3* gene independent of tag.

In both VIN3 fusions, the total protein number exhibits an increase from two to six weeks of
cold (Fig. 4a and Supplementary Fig. 11a.), although the distributions partially overlap
between the timepoints. The two-week level is between half to two-thirds of the six-week
level that approaches full vernalisation.   For these lines containing 1–3 copies of VIN3, this
steady upregulation does not appear to change from that of a single functional copy.

We then asked whether these cold-dependent increases in protein abundance are
evidenced as higher stoichiometry assemblies, or a greater number of assemblies.
Identifying these with stoichiometry and number of tracks detected per nucleus
respectively, we saw SlimVar as a uniquely powerful tool to address this question *in vivo*.

Highly mobile fluorescent foci for both VIN3 and VRN5 could be tracked for up to ~20
consecutive image frames before photobleaching of nuclear contents occurred
(Supplementary Videos 1-3). Using bespoke ADEMScode tracking software (Methods) we
found that these nucleoplasm-localised protein assemblies were largely excluded from the
nucleolus, evident as dark regions 3–6 µm in diameter (Fig. 1k and Supplementary Figure 8).
In larger nuclei, the centres of nucleoli also appeared to exhibit weak VIN3 and VRN5
localisation (Fig. 3b-c and Supplementary Figure 8). About 1–2% of the total protein number
was detected in tracks, which compares well with our estimate, based on the depth-of-field
(Supplementary Fig. 2a), that 4–6% of the mean nuclear volume is in sharp focus in each
frame.

We detected 10–40 tracks per nucleus in the VIN3 line during cold exposure, and for VRN5
at all timepoints (Fig. 4b). No tracks were detected either in the Col*FRI* negative control or
for VIN3 at pre- or post-vernalised timepoints. A key finding was that the mean number of
VRN5 tracks per nucleus increased by ~30% after the onset of vernalisation and this increase
was maintained after return to warm (Fig. 4b). We also considered the number density of
these tracks (the number of tracks normalised by the nuclear cross-sectional area). The
number density of VRN5 tracks similarly increased by about 30% over the vernalisation time
course (0.55 ± 0.05 and 0.71 ± 0.05 μm^{-2} before cold and 14 days post-cold respectively:
N=62, p=0.0042|*).

We found that both VIN3 and VRN5 exhibited broad stoichiometry distributions from a few
molecules up to several tens of molecules for individual assemblies (Fig. 4c) and that the
average stoichiometry increased with time duration spent in the cold during vernalisation.

For VIN3, the mean stoichiometry was 12.0 ± 0.4 molecules at V2W, increasing to 18.6 ± 0.5 molecules at V6W (N=1,988, p<0.001|**). For VRN5, assemblies were found to be well developed prior to vernalisation (mean 18.5 ± 0.6 molecules at NV). However, there was an increase in stoichiometry during vernalisation which persisted after the return to warm conditions (mean of 24.4 ± 0.9 molecules at V6W+T14; N=1,626, p<0.001|**). The greatest change occurred during the intermediate stages of vernalisation between V2W and V6W (17.4 ± 0.7 to 23.4 ± 0.6 molecules, N=1,928, p<0.001|**).

In summary, the mean number of tracked VRN5 assemblies in each nucleus, and mean stoichiometry of assemblies, each increased over the full course of vernalisation by the same proportion: approximately 30–35%. Thus, the additional VRN5 protein is divided equally into new vernalisation-induced assemblies, as well as into enlarging assemblies that resembled the pre-vernalised state.

Conversely, VIN3-GFP increased in total protein number by 58% between two-weeks and six-weeks of cold, which matched the proportional 55% increase in its stoichiometry during the same interval. The number of tracks per nucleus did not increase significantly in either of the VIN3 lines (Fig. 4b and Supplementary Fig. 11). All the additional VIN3 protein generated during vernalisation was incorporated into the existing number of assemblies.

Multimolecular assemblies of VIN3 and VRN5 contain multiples of two molecules

The stoichiometry distributions show a series of periodic peaks (Fig. 5) that are revealed when represented as a kernel density estimate, a method which objectifies the equivalent histogram bin width used.[63] If the assemblies represented have a common oligomeric structure, the characteristic peak-to-peak interval—the periodicity—is equivalent to the number of molecules associated with a physical subunit of the assembly.[64] We developed a new analysis method to discriminate this periodicity using the most common nearest-neighbour peak intervals, verified using realistic statistical simulations and experimental data from standard LacI tetramers *in vivo* [65] (Methods, Supplementary Figs. 9 and 10). Both VIN3 and VRN5 exhibit neighbouring peaks in their stoichiometry distributions separated by two molecules (Fig. 5 insets).

We performed SlimVar on the VIN3-SYFP2 line (Supplementary Fig. 11). Our analysis indicated VIN3-SYFP2 has averages of total protein number, stoichiometry and periodicity consistent with the VIN3-GFP line, but only when corrected for the proportion of unlabelled VIN3 present, as estimated from qPCR (Supplementary Fig. 5). We estimated the correction factor by taking the ratio of mRNA expression levels of VIN3 to SYFP2, then normalising by the ratio of VIN3 to GFP expression in the green line, which lacks endogenous VIN3. While inference of quantitative protein levels from mRNA levels is limited, the dependence of the observed periodicity on labelling provides further confidence: if the observed oligomeric species were artifacts mediated by self-interactions of the fluorescent protein tags,[66] the periodicity of visible oligomers would not vary in proportion to unlabelled VIN3. Together, these observations are consistent with VEL proteins dimerising within higher-order oligomeric assemblies *in vivo*. It is intuitive to think of these oligomers growing with the addition of dimeric units, though we cannot test this directly; we do not probe the

544 molecular kinetics or structure that would indicate the pathways of scaffold assembly and
545 disassembly.

547
548

Fig. 5. VIN3 and VRN5 assemblies exhibit a two-molecule spacing in their stoichiometry
distributions. The number of labelled molecules in each assembly (stoichiometry) shows
consistent peak-to-peak spacing via periodicity analysis of a) VRN5-YFP and b) VIN3-GFP
across different vernalisation timepoints: NV = not vernalised; V2W = two weeks of cold;
V6W = six weeks of cold; V6WT14 = six weeks of cold followed by two weeks of warm
conditions.   A kernel width (curve smoothing parameter) of 0.6 molecules was used
corresponding to the standard deviation in the observed intensity of a single molecule at
the sifting signal-to-noise threshold. Insets: Periodicity analysis - the number of molecules in
this subunit can be estimated from the most common spacing between neighbouring peaks
in each stoichiometry distribution. The threshold above which a null (aperiodic) distribution
can be rejected is the 95[th] percentile fraction of intervals (grey trace) output from simulated
random stoichiometry (Methods). The most common interval is given by the modal kernel
density estimate ± s.e.m. above the null threshold (VIN3-GFP: V2W, 1.9 ± 0.3; V6W, 2.2 ±
0.3. VRN5-YFP: NV, 1.9 ± 0.4; V2W, 2.2 ± 0.4; V6W, 2.0 ± 0.3; V6W+T14, 2.0 ± 0.4). The
periodic unit in each of these cases is consistent only with an assembly subunit of 2
molecules of either VIN3-GFP or VRN5-YFP.

Mobility of larger VIN3 and VRN5 assemblies matches that of *FLC* during cold exposure

We estimated microscopic diffusivity *D* for each detected track by calculating the gradient to
the initial portion of its corresponding mean square displacement (Methods).

Fig. 6. Microscopic diffusivity of VIN3 and VRN5 assemblies decrease towards that of
FLC **loci during vernalisation.** a-d) Diffusivity *D* of individual tracks estimated from mean-
square displacement analysis at different vernalisation timepoints: NV = not vernalised;
V2W = two weeks of cold; V6W = six weeks of cold; V6WT14 = six weeks of cold followed by
two weeks of warm conditions. a) LacI-YFP tracks of fewer than 12 molecules, detected from
nuclei without pre-bleaching and identified as unbound LacI, and of LacI-YFP tracks of more
than 12 molecules after pre-bleaching, identified as *FLC* candidates; b) Diffusivity of VIN3-
GFP and c) VRN5-YFP before, during and after vernalisation. VIN3-GFP and each exhibit a
decrease in mobility during the latter part of vernalisation, persisting in VRN5 following
return to warm conditions: (VIN3-GFP: D = 0.52 ± 0.03 to 0.41 ± 0.01 $\mu m^2 s^{-1}$; mean ± sem;
N=672, p=0.0011|*; VIN3-SYFP2: D = 0.58 ± 0.02 to 0.49 ± 0.02 $\mu m^2 s^{-1}$; N=1,552,
p<0.001|**; VRN5-YFP: 0.47 ± 0.02 / 0.48 ± 0.02 $\mu m^2 s^{-1}$ at NV / V2W to 0.38 ± 0.01 / 0.40 ±
0.02 $\mu m^2 s^{-1}$ at V6W / V6W+T14; N≥982, p=0.0072 |*). Horizontal lines denote diffusivity of
FLC-lacO/LacI-YFP foci under the same conditions (solid line: mean value; grey area,
agreement within error).

For comparison with the diffusivity of *FLC* loci, we performed SlimVar imaging of a *FLC-*
lacO/LacI-YFP line with 120 *lacO* copies integrated downstream of the *FLC* transgene [67]. To
obtain a qualitative indication of the proportion of VIN3 and VRN5 assemblies which might
be bound to *FLC* and their stoichiometry, we analysed just tracks whose diffusivity was
comparable to that of a typical *FLC* locus (D_{FLC} = 0.20 $\mu m^2 s^{-1}$, Supplementary Fig. 6) within
individual track measurement error (± 0.07 $\mu m^2 s^{-1}$) such that D lies in the range
0.13 – 0.27 $\mu m^2 s^{-1}$ (Fig. 6a). This simple method of diffusivity matching [68] is particularly
helpful when only single-label lines are available. The subset of VIN3 or VRN5 assemblies
with diffusivity consistent with *FLC* (Supplementary Table 1) have stoichiometries
distributed similarly to the full cohort of VIN3 or VRN5 tracks (Fig. 5c) and show similar
increases in stoichiometry over the course of vernalisation for both VIN3 and VRN5. The
median stoichiometry of each protein using this diffusivity matching was in the range 10–20
molecules per assembly, increasing with vernalisation.

We found that VIN3 assemblies became significantly less mobile between two and six weeks
of cold exposure (Fig. 6b, Supplementary Fig. 11e). VRN5 assemblies exhibited a similar
~20% decrease in diffusivity during the same central stage of vernalisation (Fig. 6c). In

keeping with the increase in stoichiometry and lower diffusivity, the proportion of VIN3 assemblies that show slow *FLC*-like diffusion increases from around 11–15% during early stages of vernalisation up to around 17–18% (Supplementary Table 1). The fraction of *FLC*-like VRN5 assemblies present in the nucleus is already ~18% prior to vernalisation and maintains this level throughout. However, given the greater concentration of VEL protein assemblies compared to *FLC* loci, this proportion is unlikely to be representative of VEL localisations at *FLC*, which if present would be a minority observable only by direct colocalisation.

VRN5 assemblies at *FLC* have higher stoichiometry during and after vernalisation

To directly track VRN5 at the *FLC* locus, we generated transgenic plants co-expressing VRN5 (fused to mScarlet-I [69], mScI) and *FLC-lacO* (via LacI-YFP) for dual-colour SlimVar. Our test for colocalisation of VRN5 at *FLC* first required a reliable method to identify *FLC* without perturbing the mScarlet-I reporter for VRN5. For each nucleus, we first performed a rapid z-stack using 514 nm wavelength laser excitation to screen for and localise bright, low mobility LacI-YFP foci consistent with *FLC* genomic loci (Fig. 7a). At a chosen z-position containing one or more of these *FLC* candidates, we then tracked VRN5-mScI and LacI-YFP using alternating laser excitation (Methods); most images were dominated by the presence of unbound LacI-YFP foci, however, we measured a distinct subset bound to *FLC* (Fig. 7b) with a frequency of 2.3 ± 1.4 (mean ± s.d.) per nucleus in the meristem, matching the expectation of 2 *FLC* loci per nucleus within experimental error [67]. This rose to 3.3 ± 1.7 in cells toward the transition zone, consistent with more nuclei exhibiting additional pairs of *FLC* loci under genomic endoreduplication. Approximately 40% of detected *FLC* loci were colocalised with VRN5-mScI assemblies, though this proportion was constant across all vernalisation times. This led us to question: might a putative memory element be conditional on the properties of the colocalised VRN5 assemblies, such as stoichiometry?

634
635

Fig. 7. A dynamic subset of enlarged VRN5 assemblies is present at *FLC* loci after long cold exposure and after return to warm.

a) Representative example of a z-stack used to screen for *FLC* loci in the LacI-YFP channel, followed by b) two-colour alternating excitation at a single target z-plane to capture *FLC* loci (circles) and colocalisation events (solid circles) between *FLC* (yellow) and VRN5-mScI assemblies (magenta) as determined by tracking (Methods). The example shown is at six weeks of cold exposure. Note that the genomic loci *FLC-lacO*, rather than individual nucleosomes, are localised experimentally by LacI-YFP in yellow tracks. c) The mean stoichiometry of VRN5 colocalised at *FLC* exceeds that of uncolocalised VRN5 because of vernalisation. The timepoints are: NV = not vernalised; V2W = two weeks of cold; V6W = six weeks of cold; V6W+T14 = six weeks of cold followed by two weeks of warm conditions. The difference is initially negligible before vernalisation (3.5 ± 0.3 vs 3.5 ± 0.1, NS, p=0.39, N=365) but appears clearly at V6W (7.7 ± 0.6 vs 5.0 ± 0.2, p<0.0001***, N=2867) and is sustained for at least two weeks after return to warm (7.8 ± 0.5 vs 5.2 ± 0.2, p<0.0001***, N=1416).   d) Vernalisation increases the fraction of colocalised *FLC* with VRN5 above a

stoichiometry threshold. Total number of detected *FLC* (Supplementary Table 2) found to be
colocalised with VRN5 are: (NV) 11, (V2W) 43, (V6W) 83, (V6W+T14) 29. Thresholds are set
at which dependence on vernalisation becomes statistically significant (>10 molecules) or
maximal (>6 molecules). The proportion of colocalised *FLC* associated with >6 VRN5
molecules are (NV) 1/11 and (V2W) 3/43, while this frequency rises with vernalisation to
(V6W) 35/83 and (V6W+T14) 12/29 colocalised *FLC.* The increases between two and six
weeks, and between two weeks and return to warm are significant (Fisher's exact test two-
tailed, p=7 × 10^{-4} and p=3 × 10^{-5} respectively). e) The diffusivity of VRN5-mScI tracks either
colocalised to *FLC* (decreasing with vernalisation) or independent of *FLC.* f) An illustration of
the model supported by the imaging results. The model uses *FLC* as the locus, showing VEL
protein dynamic self-assembly and function in vernalization-driven epigenetic silencing;
prior to vernalisation, VEL proteins interact with each other and PRC2, but the assemblies
are modest in size and the majority diffuse too quickly to be dependent on interaction with
chromatin. During vernalisation, the VEL assemblies become larger and preferentially
localize at *FLC* compared to their status prior to vernalization, consistent with facilitating
greater PRC2-mediated nucleation of silencing marks. Large VRN5 assemblies persist at *FLC*
after the cold stimulus is removed, which may assist the sustained spreading of marks and
maintenance of vernalisation memory.

Both number and stoichiometry of VRN5 assemblies were considerably lower in this single-
copy VRN5-mScI line than the multiple-copy VRN5-YFP line (Fig. 4c). Like VRN5-YFP,
however, these VRN5 assemblies showed stoichiometries that increased with vernalisation
independent of colocalisation at *FLC* (Fig. 7c, grey). There was a proportionally far greater
increase in stoichiometry of VRN5 assemblies colocalised at *FLC,* particularly after long cold
exposure and after the return to warm (Fig. 7c, magenta). Correspondingly, the fraction of
colocalised *FLC* sites associated with assemblies of more than 6 VRN5 molecules increased
around fourfold to approximately 40% between two and six weeks of cold (Fig. 7d). At six
weeks' cold, the frequency of *FLC* loci colocalised with >6 VRN5 molecules exceeded an
average of one site per nucleus. This preferential colocalisation of FLC with larger VRN5
assemblies after six weeks of cold and on return to warm was detectable for thresholds
between 5–10 VRN5 molecules.

To establish the timescale of interaction between VRN5 and *FLC*, we then compared their
mobility (Fig. 7e). Before cold and at two weeks of cold exposure, the mean diffusivity of
the colocalised VRN5 assemblies exceeded that of *FLC* loci by a factor of ~3, indicating short-
lived binding relative to the ~10 ms sampling timescale. After six weeks of cold and after
return to warm, the diffusivity of colocalised VRN5 ever more closely matched the typical
diffusivity of *FLC* loci, potentially pointing to a tighter interaction. Notably, the fraction of
VRN5 not colocalised at *FLC* remained at the higher diffusivity independent of vernalisation.
An unexpected observation was the rapid rate of VRN5 turnover at *FLC* that is sensitive to
assembly size. The mean apparent residence time [70] at *FLC* of colocalising assemblies
smaller than 6 VRN5 was 26 ms before, and 34 ms after, vernalisation, distinctly shorter
than the typical photobleaching time of 63 ± 2 ms. For assemblies larger than 6 VRN5 after
vernalisation, the residence time was 64 ± 6 ms, indicating minimal dissociative loss on this

timescale relative to photobleaching. This difference indicates turnover of smaller
assemblies at *FLC* on subsecond timescales, in contrast to less transient binding of larger
assemblies.
697
In Lövkvist et al., 2021 [17], an assembly size of 17 was predicted as the best fit to
observations of nucleating silencing marks, potentially reducing further to 10 in the case of
maximal positive cooperativity. Although we do not know whether VIN3 assemblies
colocalise specifically with *FLC*, and if so, the size of such an assembly, a large minority of
observed VIN3 and VRN5 assemblies with sufficiently low mobility to interact with *FLC*
clearly do exceed this size threshold during and after vernalisation. In the case of VRN5, we
show this is exceeded by a small minority of assemblies directly colocalised with *FLC* only
after vernalisation, though at a single gene copy, the maximal distinction with uncolocalised
assemblies occurs at even lower thresholds of as low as 6–10 molecules. This finding
suggests that if any lower bound size is required for positive feedback to become active in
these colocalised assemblies, it must be very low.
709

Discussion

Here, we have developed a new optical microscopy pipeline—SlimVar—and applied it to
image cell nuclei in *Arabidopsis* root tips. SlimVar is optimised to detect single fluorescent
protein fusions by sampling faster than their typical motion in cells. It therefore enables
tracking of molecular assemblies diffusing in comparatively deep, multicellular samples,
without requiring complex protocols for chemically conjugating target biomolecules.

In animal tissues, studies using HILO [61 71], lightsheet [72 73] or lattice lightsheet [35] have
previously demonstrated single-molecule tracking to a standard of 50 ms sampling at ~300
μm depth, or 10 ms at ~30 μm depth using dyes, which is comparable to SlimVar with
fluorescent proteins. However, in plant tissues, single particle tracking at molecular
sensitivity has been demonstrated with TIRF [74,75], or VAEM [31,33,76–78] only in the vicinity of a
surface cell layer. SlimVar therefore advances the ability to track and count single-molecular
assemblies in plants, and potentially in a range of tissues, to that of more complex existing
microscopy technologies.

In achieving this speed, SlimVar trades off some of its 3D capability; the detection of foci is
restricted to a limited depth of field much smaller than the nucleus. Nonetheless, it is
capable of z-stacks as used here for systematic *FLC* detection, and it should be possible to
extend oblique angle [51] or lightsheet-based approaches to achieve rapid volumetric scans [79]
and extended depth of field [80]. Within the depth of field itself, astigmatic 3D imaging has
been demonstrated with Slimfield [81]. The implementation of better index-matching and
photon-efficient adaptive optics [36,82] could also improve the range of accessible working
depths currently limited by scattering losses, while also mitigating the required adjustments
at each depth. Single-molecule experimental schemes related to SlimVar such as multiple-
colour or photoconvertible labelling on the same target [83], may provide promising future
avenues to probe the turnover dynamics over longitudinal experiments.

As presented here, SlimVar is implemented on a custom microscope with free-space components. However, the optical components and controls are essentially similar to commercial widefield/TIRF microscopes used for single-molecule localisation microscopy. Commercial instruments do not typically provide the intense excitation or detector sensitivity sufficient to track single fluorescent proteins at depth (Supplementary Fig. 2). As such, we use the new platform to explore newly accessible biological insight. In future work, we aim to democratise and enhance SlimVar accessibility further by minimally adapting a representative, modern commercial microscope that meets these requirements. Although optimised for applications in *Arabidopsis* root tips, SlimVar is expected to be adaptable for functional bioimaging research at the molecular scale inside a range of living tissues.

Using SlimVar, we demonstrated oligomeric assemblies of the PRC2 accessory proteins VIN3 and VRN5. VIN3 and VRN5 assemblies increase in stoichiometry above a demonstrable threshold during prolonged cold. Both sets of assemblies exhibit mobility signatures similar to *FLC* gene loci, with a subset of larger VRN5 assemblies clearly demonstrating *FLC* colocalisation. The higher stoichiometry assemblies are therefore prime candidates to contribute to memory element function predicted in a hybrid model coupling protein self-assembly and histone modification [17]. A major advantage of SlimVar over ensemble techniques is that it directly probes not only whether proteins become abundant (total protein number), but whether such change take effect as higher/lower assembly concentration (tracks per nucleus) or as larger/smaller assemblies (stoichiometry).   Our observation that adding more protein subunits preferentially results in a larger average assembly stoichiometry, rather than a greater number of assemblies, is itself consistent with a model of positive cooperativity. Taken together, these findings support the view that VIN3 and VRN5 assemblies mediate epigenetic memory over the extended vernalisation cycle of several weeks (Fig. 7f). However, our current imaging results do not make a direct claim about nucleosomes, only about the *FLC* locus reporter, for which the endogenous locus contains at least three nucleosomes in the nucleation region relevant to VEL function.

The model [17] makes no predictions as to the underpinning factors and mechanisms - neither for protein self-assembly, nor dynamic exchange with the surrounding nucleoplasm. The head-to-tail polymerization via the VEL domain [23] is likely involved, but the predominantly transient interactions between most individual VEL protein assemblies and *FLC* may suggest that the physical feedback processes are more complex than currently understood. Our current interpretation is that the VRN5 assemblies enriched at *FLC*, of sufficient size to satisfy the model, are most likely simple oligomers. Yet, we contemplate whether the very largest of these (~100 VRN5) are instead small, dynamic, phase-separated condensates, related to those observed during transient overexpression of VEL proteins [19,23,84]. If so, the collective, multivalent interactions characteristic of condensates might offer a longer, or otherwise more effective, residence time at *FLC* than we observed for typical molecular assemblies, and therefore a disproportionate contribution to epigenetic memory. SlimVar is an excellent tool capable and primed to further investigate these rare mechanistic events *in vivo*. Further work will investigate the phenotypic and molecular interactions between VRN5 and VIN3 mutants [84].

While pioneered for the *Arabidopsis FLC* system, these protein-mediated feedback
processes may underpin Polycomb-based epigenetic memory common to all eukaryotic
systems. Our study demonstrates the interdisciplinary value at the interface of the physical
and life sciences of developing SlimVar, and other bioimaging technologies at single
molecule precision, to tackle outstanding biological questions in epigenetic processing and
memory.

Methods

Plant material and growth conditions

VIN3-GFP and VRN5-YFP lines have previously been described [11,19]. To generate the VIN3-
SYFP2 line, GFP of the pENTR pVIN3::VIN3-GFP construct [85] was replaced by SYFP2 by
seamless mega-primer cloning. VIN3-SYFP2 was cloned to the SLJ destination vector, a
derivative of SLJ755I5 [86] and transformed to *Agrobacterium tumefaciens* C58 by triparental
mating. The transgenic VIN3-SYFP2 plant in Col*FRI* background was generated by floral
dipping with *Agrobacterium*. To generate pENTR pVRN5::VRN5-mScarlet-I, the SYFP2 of
VRN5-SYFP2 [84] was replaced by mScarlet-I by seamless mega-primer cloning. The VRN5-
mScarlet-I was cloned to the SLJ destination vector, a derivative of SLJ6991 [86], transformed
to *Agrobacterium tumefaciens* C58 and subsequently transferred into *vrn5-8 FRI* mutants as
described above. All primers used are listed in Supplementary Table 3. Transgene copy
number was determined in T1 or T2 transformants by IDna Genetics (Norwich Research
Park). To generate the plant co-expressing VRN5-mScarlet-I and *FLC-lacO*/LacI-YFP, the
VRN5-mScarlet-I line was crossed into the *FLC-lacO/lacI-YFP* line [67] and was selected by
antibiotics.

All seeds were surface-sterilised and sown on 100 mm growth plates containing Murashige
and Skoog (MS) medium (Duchefa) with 1 wt. % agar (Difco Bacto) without sucrose. The
plates were sealed with Micropore tape (3M) and kept at 4°C in the dark for 2–3 days to
stratify the seeds. Plates were racked vertically in the growth chamber in warm conditions
(16 h light/ 8 h dark with constant 22 ± 2°C, 60 ± 10% Relative Humidity (RH)) for 7 days.
Non-vernalised timepoints (NV) were then imaged on the final day. For all other timepoints,
plants were grown in warm conditions for 7 days as above and then were transferred to
cold conditions (8h light / 16h dark, 5 ± 1°C, 50 ± 30 % RH) to vernalise for either 2 or 6
weeks (V2W and V6W respectively). Following vernalisation, a subset of plates was
returned to warm conditions for an additional period of either 7 or 14 days (V6W+T7 or
V6W+T14). Plates for warm timepoints (NV, V6W+T7, V6W+T14) were handled at room
temperature, while those imaged at cold timepoints (V2W and V6W) were transferred on
ice to a 4°C cold room for slide preparation to avoid temperature spikes affecting the fragile
vernalised state. VIN3 expression is known to be modulated by the cellular circadian clock
[10]; to isolate the long-term trends in expression relating to cold exposure, imaging was
performed in daylight hours 4–8 to align with the diurnal maximum in VIN3 expression.   At
least 3 independent vernalisation courses were grown for each line and timepoint
(Supplementary Table 2). Immediately prior to imaging the *FLC-lacO*/LacI-YFP lines (single
colour or dually labelled with VRN5-mScarlet-I), LacI-YFP was induced by placing opened

growth plates next to a bath of 0.5% EtOH at 25°C in an airtight container for 2.5 h. This
resulted in an optimal amount of LacI-YFP expression in the meristem for SlimVar imaging
without spot-bleaching, below the near-saturated induction level (1.5–2% EtOH for 1.5 h [67])
used for confocal study (Supplementary Fig. 9a-d).

RNA expression analysis

Total RNA was extracted using the phenol method as previously described [11,87]. Genomic
DNA was removed from the TURBO DNA-free kit (Invitrogen, AM1907) before reverse
transcription with SuperScript IV reverse transcriptase (Invitrogen, 18090050) and gene-
specific primers. Quantitative PCR analysis was performed on a LightCycler480 II (Roche).
Target gene expression was normalised by PP2A (AT1G13320) and UBC (AT5G25760). All
primers used in this study are listed in Supplementary Table 3.

Preparation of samples for imaging

Identical slides were prepared for confocal, Airyscan or SlimVar imaging [61]. Briefly,
GeneFrames (Thermo Scientific, AB0578) were fixed to standard slides (VWR) and filled with
MS medium plus 1 wt.% agarose to produce agar pads. Where necessary due to seedling
size, the terminal >10mm of the primary root of each plant was excised using a razor. Root
tips were laid on each agar pad with tweezers. Liquid MS media was applied to exclude air
and each slide was sealed with a plasma-cleaned #1.5 coverslip (VWR). Each slide was
imaged within <1 h.

Confocal and Airyscan imaging

Confocal imaging was performed on a Zeiss LSM880 microscope equipped with argon ion
laser and Plan-Apochromat 63×/1.40 NA oil objective lens (Zeiss). Samples were illuminated
at 488 or 514 nm wavelengths (GFP or YFP/SYFP2 channels) respectively, and the emission
detected at 490–550 nm or 518–550 nm respectively [61]. The root tip confocal images (Fig.
3b, Supplementary Fig. 7, Supplementary Fig. 9a-d) were acquired as z-stacks over ≤3 z-
slices of 1,000 × 1,000 pixels at 0.6× zoom factor, at 1.5 µm intervals using 20 mW excitation
power. Slices were postprocessed in FIJI/ImageJ [88] with a 2D median filter (0.2 µm filter
size) to suppress noise before performing a mean z-projection.

Timelapse z-stacks of nuclei (Fig. 3c) were acquired in Airyscan RS mode, after aligning the
detector with immobilised TetraSpeck microspheres (Invitrogen, 0.1 µm diameter). Each
sequence contained ≤25 volumetric cycles, each taking 0.84 s (12 slices × 56 ms/slice). The
axial step size was 0.65 µm with 112 × 112 pixels at 14× zoom factor and 20 mW excitation
power. The 3D z-stacks were Airyscan post-processed in Zen Black software (Zeiss) with a
user-optimised strength parameter of 5.0. Vernalised timepoints at two and six weeks of
cold (V2W, V6W) were imaged at 5 ± 1 °C on a water-recirculating Peltier-cooled stage
(Linkam PE100) calibrated using hydrocarbon melting points.

SlimVar imaging platform

857 SlimVar was adapted from a Slimfield microscope comprising objective-lens-based total
858 internal reflection fluorescence (OpenFrame, Cairn Research), custom-built from benchtop
859 optical components (Thorlabs) and a nanopositioning stage (Mad City Labs). A general
860 scheme of Slimfield microscopy is available [43,44,89 48] with key terms defined in Table 1.

Metric / Object	Definition
Foci	Spot-like local intensity maxima in a single frame, each corresponding to a localised group of labelled molecules. Associated properties include spatial/temporal location, total intensity, and signal-to-noise ratio.
Track	A set of foci in adjacent frames which are spatially close enough to form a contiguous trajectory. Associated properties include those of the set of foci, plus stoichiometry, diffusivity, and signal-to-noise ratio.
Characteristic molecular brightness	The average number of photon counts per frame associated with a single fluorescent reporter molecule (e.g. GFP), under a fixed imaging condition. Equivalent to the number of photons in the most common intensity step observed for tracks in the final stage of photobleaching.
Integrated nuclear intensity	The total fluorescence intensity, including autofluorescence, of an entire nuclear segment in photon counts, normalised by the characteristic molecular brightness. Dimensionless; described in fluorescent protein equivalents.
Total protein number	The average number of labelled molecules in a nucleus, not including autofluorescence, estimated from the difference in integrated nuclear intensity from that of an unlabelled negative control.
Stoichiometry	The number of labelled molecules in a track, as estimated by dividing the track's initial intensity by the characteristic single-molecule brightness.
Periodicity	The number of labelled molecules in a repeat unit within tracked objects, as estimated by the consistent stoichiometry intervals between nearest-neighbour peaks in the stoichiometry distribution.
Diffusivity	An average measure of the rate of random microscopic motion of a track based on the increase in its mean-squared displacement over time.
Signal-to-noise ratio (SNR)	A measure of the signal strength of foci or tracks compared to background noise. Higher SNR implies higher confidence of a true positive detection.
Sifting	Sifting is the postprocessing step which imposes a minimum SNR threshold on foci and tracks, and a minimum track length, to improve robustness of the track-wise metrics.

863 **Table 1. Glossary definitions of analysis metrics for single particle tracking.**

With increasing acquisition depth, refractive index mismatch between immersion oil and aqueous sample is a key challenge [56] due to spherical aberration and excitation beam deviations. This is usually avoided with expensive water or silicone immersion objective lenses, but we demonstrate equal or superior performance with an affordable oil immersion objective lens: NA 1.49 Apo TIRF 100× oil (Nikon). For Slimfield and SlimVar, as with other single-molecule techniques with widefield detection, objective lenses must have a large back aperture diameter, common in lenses specialised for TIRF. Single molecule sensitivity was afforded by a fast sCMOS camera (Teledyne Photometrics Prime95B, 12-bit 'Sensitivity', or BSI, 16-bit 'Sensitivity' i.e. gain of 0.6 photoelectrons per count with an offset of 100 counts). The turret dichroic mirror was either dual-band for GFP/mScarlet-I (Chroma ZT488/561rpc) or for YFP/mScarlet-I (Chroma ZT442/514/561rpc) as appropriate. A 580 nm wavelength longpass beamsplitter (OptoSplit, Cairn Research) after the tube lens (f=200 mm achromat, ThorLabs AC254-200-A-ML) enabled simultaneous detection of GFP (525/50 nm wavelength centre/bandwidth emission filter) or YFP (550/25 nm) in a green/yellow channel, and mScarlet-I (594/25 nm) in a red channel. Additional magnification (1.2 – 2.2×) was used to compensate for different camera pixel sizes to maintain an oversampled pixel width of 53 ± 5 nm in the images.

Continuous wave lasers (Coherent OBIS) delivered Gaussian beams (TEM_{00}) at 488 nm, 514 nm and 561 nm wavelengths with 1.9 mm FWHM beam diameter that were circularised by an achromatic quarter waveplate [90]. These were steered and focused at the objective back aperture using f=150 mm lenses to collimate them through the sample with a FWHM diameter of 25 µm.

To optimise background contrast for *Arabidopsis* root tips, the following adaptations were made to the Slimfield microscope:

i) the second convex lens in the expansion telescope was mounted on a lateral translation stage with a high-precision micrometer; this shift (up to 2.3 mm) generated an equivalent lateral displacement of the beam at the 6 mm diameter objective back aperture, thereby tilting the beam away from the optic axis at the sample. The beam delivery angle was calibrated for lens micrometer position following [49] and set to 50° ± 3° from normal incidence in oil, corresponding to 60° ± 5° in water by Snell's Law, with minimal coupling into the evanescent field at the coverslip, similar to HILO [51] but unlike PaTCH [91] or VAEM [50–52,92].

ii) a field stop was placed in the conjugate plane upstream of the first telescope lens. Rather than thinning the beam with rectangular slits to further suppress background,[52] we chose to maintain a circular beam; this illuminated the full depth of each nucleus for representative estimates of nuclear protein copy and efficient screening of *FLC* loci. The stop was tightened to crop the beam from 25 µm FWHM to between 4–9 µm cross-section. The field then approximated an ellipse of $1 \times 3^{1/2}$ the chosen beam cross-section, or 4–16 µm diameter, uniformly illuminated at a power density of 1–5 kW cm^{-2}, which is a few-fold less than the saturation excitation intensity of the fluorescent proteins.[58]

iii) a pair of mirrors were used to incline the beam ~5 mrad away from the optic axis at its intersection with the objective back aperture. This compensates the beam's mismatch-dependent lateral deflection of ~11 µm at the 25 µm calibration depth.

908   The aberrations were then corrected using the following protocol.

Optical calibration protocols

910   Two forms of calibration are presented: basic, relying only on high-signal fluorescence with
911   a correction collar, or an advanced protocol, also using single molecules and tube lens shift.
912   For best results, the advanced protocol requires independent z-positioning of the objective
913   lens and the sample stage, since this enables decoupled correction of both defocus and
914   spherical aberrations.

Basic aberration correction with objective collar only

1. Choose a nominal working depth (e.g. d = 25 µm).
2. Prepare a sample with sub-diffractive fluorescent features at this desired working
depth or deeper. This could comprise a suspension of fluorescent beads of diameter
<200 nm (Fluospheres, Invitrogen) in 1 wt.% agarose, or weakly autofluorescent
point-like features in a root tip.
3. Set the laser excitation to a modest level to reduce photobleaching, e.g. 0.2 mW
source power ~ 0.1 kW/cm^2 irradiance and a long exposure time ~100 ms.
4. Move the stage (or objective lens) to focus on the upper surface of the coverslip,
then move an additional distance corresponding to the nominal working depth.
5. Move the stage laterally to find a point-like sample feature close to best focus.
Centre it in the field of view without refocusing.
6. Acquire a z-stack at <0.2 µm spacing over a maximum range of ± 2 µm.
7. Repeat steps 4–6 for several such features.
8. Repeat steps 4–5 to find a new feature. Find best focus using the objective.
Iteratively decrease the collar setting to a thinner coverslip setting
(by $n_{water}/n_{oil} \times d = 30$ µm, i.e. to ~140 µm) and refocus the objective to achieve best
lateral focus (narrowest width of the feature).
9. Repeat steps 4–7 but with the new collar setting, to generate a second set of z-
stacks. Compare the differences in PSF (e.g. using MetroloJ QC).

Advanced aberration correction with single molecule sample and tube lens shift

1. First, mark the tube lens position. It is best to set any components such as filters
between the objective lens and the detector, before—rather than after—performing
the calibration.
2. Measure the total image magnification using a graticule.
3. Choose a nominal working depth (e.g. d = 25 µm). Prepare samples containing sub-
diffractive fluorescent features and slow (or immobilised) single molecules at this
desired working depth, such as a dilute (<1 nM) suspension of fluorescent proteins
and/or beads in 1 wt.% agarose, or on a tilted coverslip (Supplementary Fig. 2a).
4. Set the laser excitation to a high level, e.g 10 mW power or ~5 kW/cm^2 irradiance
with 10 ms exposure time. Adjust the camera to the corresponding imaging settings
above for the 'balanced mode'.
5. Focus on the coverslip and move to the nominal working depth using the sample
stage only (e.g. d = 25 µm)

6. Move the stage to a point-like sample feature near focus (this could be a single bead or a section of the volume or surface containing single molecules in focus). Centre it in the field of view.

7. Acquire several fields of view for reference: for beads, acquire z-stacks with <0.2 μm spacing over a range ± 2 μm. For fields of view containing visible single molecules, acquire multiple SlimVar sequences with >300 frames; a longer exposure time may be needed to distinguish single-molecule foci from background. Note this exposure time.

8. Iteratively decrease the collar setting to a thinner coverslip setting (by $n_{water}/n_{oil} \times d = 30$ μm, i.e. to ~140 μm) and move the stage (not the objective lens if possible) to achieve best lateral focus corresponding to the narrowest width of the feature.

9. Move the objective lens towards the sample (or if this is not possible, move the sample stage towards the objective) a further 20% of the working depth (e.g. by 4 μm to a new total working depth of 29 μm).

10. Compensate by moving the tube lens towards the objective (by up to 40 mm for a 200 mm focal length lens) until the plane of best focus is pulled back onto the chosen feature.

11. Using the stage, re-centre the feature in the field of view. If the region has bleached, pan the stage sideways to a new feature or field of view at the same depth.

12. Then, adjust the collar setting for best lateral resolution as above. Refocus onto the feature (preferably using the objective lens).

13. Repeat the acquisitions in the calibrated state: starting from 10 ms, adjust the exposure time until single molecules are visible. Acquire multiple fields of view in SlimVar sequences containing visible single molecules and track in ADEMScode. Compare the SNR of single molecules. If the exposure time was longer for the reference set, divide the reference SNRs by the square root of the ratio of exposure times. For beads, acquire z-stacks with <0.2 μm spacing. Compare the differences in axial and lateral resolution from the reference state (e.g. using MetroloJ QC).

14. Further iteration of focal position with either collar setting or tube lens position may be desirable for best results, as determined by minimal axial FWHM and maximum signal-to-noise metric.

15. Finally, measure the total image magnification using a graticule, as this may have changed from the nominal design magnification.

Evaluation of the advanced calibration

We implemented the latter protocol using a combination of *in vitro* YFP (Supplementary Fig. 2) and fluorescent beads (Supplementary Fig. 3). This introduced a correction collar setting of 140 μm (for a coverslip #1.5H of 170 μm thickness) and a shift of $d = -32$ mm in an $f = 200$ mm tube focal length (Supplementary Fig. 3). Despite this shift in the tube lens position, we found the optical pixel size remained consistent within the range 53 ± 5 nm, implying a change in magnification no larger than -8%. This is less than the drop of $d/f \sim -16\%$ expected from a formal change in tube length, i.e. using a tube lens with a different focal

length of (f + d) to restore the 4f imaging condition. Since the imaging is improved, the sine
condition between the sample and the intermediate image planes must be maintained,
which suggests that infinite conjugation breaks down: the emission reaching the camera is
no longer parallel to the optical axis at the back aperture of the objective. This also explains
the observed reduction in effective numerical aperture by a factor of approximately (1 -
$d/f)^{1/2}$ ~ 8% from 1.49 to ~1.38 (Supplementary Fig. 2). We suppose that relaxing the infinite
conjugation enables the combined calibration elements to deliver not only the required
amount of spherical aberration, but also the correct amount of wavefront defocus[55] for
deeper imaging.   This result suggests this strategy is workable for small defocus
compensations, without introducing a different tube lens or using e.g. spatial light
modulators that can provide separable, tuneable corrections to defocus and spherical
aberration. However, the beamsplitters and other elements between the objective and
tube lens would ideally need to be kept constant between calibrations.

The calibration yields a minimal FWHM of detected foci ~170 nm (Fig. 3d) and a localisation
precision[93] of 40–80 nm (Supplementary Fig. 12). In plants we see a net increase of 2.6 in
the median signal-to-noise ratio relative to epifluorescence microscopy, when accounting
for the decreased exposure time (normalised by the square root of exposure time ×
excitation power, n = 500 foci, Supplementary Fig. 2j).

Settings for SlimVar imaging in plants

Nuclei were identified in brightfield to find best focus at the nucleolar midbody without
photobleaching and were captured for manual segmentation. Fluorescence acquisition
settings in green, yellow and/or red channels were pre-optimised to avoid initial camera
saturation and to ensure detection of individual tracks of molecular brightness for GFP,
SYFP2 / YFP, and mScarlet-I respectively. Three exposure times were used: i) a
'representative' 20 ms providing a low-bias detection of all particles down to single
molecules, ii) 'fast' 2 ms to ensure robustness of tracking fidelity and mobility
measurements of only the brightest assemblies, and iii) a 'balanced' 10 ms, also capable of
single molecule detection. Fast acquisitions captured the subset of assemblies with fourfold
(upper quartile) mean stoichiometry at a tripled detection rate due to the faster sampling
bandwidth. Results shown derive from data collected with the representative (VRN5-YFP
and VIN3-GFP lines) or balanced mode (VIN3-SYFP2, *FLC-lacO*/LacI-YFP and VRN5-mScarlet-I
lines) unless otherwise stated. These settings were fixed to minimise systematic variation in
the characteristic molecular brightness (Supplementary Fig. 1).

The region of interest spanned ≤300 rows (16 μm) giving a readout time of 2.7 ms per
frame, and sampling rates of 44–217 fps. Lasers were triggered in each frame by the
camera in 'All Rows' mode to provide global shuttering without extraneous photobleaching.
To capture photoblinking, the number of frames in each sequence was set to >50× the
photobleaching decay constant.

Protein tracking, sifting and intensity analysis

SlimVar analysis used batches of OME TIFF image sequences as input to MATLAB-based
ADEMScode v2.2 software [94,45], whose key features are also available in open-source Python
package PySTACHIO [95].   The fluorescence sequences were cropped to specify individual

nuclei, using masks manually segmented from corresponding brightfield images (code -

Segmentation: *mask2seg*). All analysis was restricted to the image region of effectively

uniform laser illumination (80% ± 9% s.d. peak irradiance) no greater than 190 × 300 pixels

(10 × 16 μm). Each fluorescence sequence was then processed independently (code -

Tracking: *trackAllFields*). For each subsequent frame, the local fluorescence maxima—foci—

were identified, and their intensity estimated, by integrating the pixel value intensity within

5 pixels, and subtracting a background level averaged over the remainder of a 17 × 17-pixel

sliding window. Each of the foci was assigned a signal-to-noise ratio (SNR) equal to its

intensity divided by the standard deviation of the associated background region. Foci above

an initial, permissive SNR threshold of 0.2 were tentatively accepted. These foci were

refined to subpixel precision with an elliptical Gaussian masking algorithm. This returned

fitted estimates of their semi-axes, which reflected the widefield spatial resolution plus out-

of-plane defocus and any residual motion blur. These widefield dimensions (Supplementary

Fig. 12) are close to laterally isotropic and insensitive to stoichiometry, since all the

assemblies are expected to be much smaller than the diffraction limit and to rotate rapidly

relative to the exposure time. The centroids of the foci were also estimated to super-

resolved localisation precision [93] of typically 45–80 nm (Supplementary Fig. 12) but as low as

40 nm for bright assemblies, consistent with best Slimfield performance [96]. For each image

sequence, pairs of foci were then linked consecutively into tracks if their centroids lay within

8 pixels, with width and intensity ratios in the ranges 0.5–2 and 0.5–3 respectively. When

multiple links were possible, the nearest suitable neighbour was chosen.

We then performed sifting of foci and tracks (code - Analysis: *analyseAllTracks*), retaining

only those above a strict SNR threshold and minimum number of frames, respectively.   The

first sifting criterion - an SNR threshold - was robustly determined as the SNR value at which

true and false positives occur at equal frequency in a positive control that contains only

single molecules; the false positive rate is estimated from negative controls containing only

noise. For negative controls, we tracked acquisitions of dark noise (zero excitation

intensity), as well as image stacks of simulated noise based on the autofluorescence level

measured in wild-type root tips at the typical excitation intensity (Supplementary Fig. 4).

Prior works recommend a sifting SNR threshold of 0.4 [89,90,97]; here, we found the

appropriate SNR threshold was largely independent of excitation intensity in the kW/cm^2

range and instead determined by two factors: the autofluorescence relative to the

characteristic molecular brightness (Supplementary Fig. 4), and the detector noise. The

actual SNR threshold used ranged from 0.35 for the Prime 95B (12-bit Sensitivity) to 0.50 for

the Prime BSI 16-bit mode (Supplementary Fig. 2c).

Even with accurate distinction of foci from background noise, random overlap between

tracks increases at high emitter density, leading to spurious summation of their intensities.

Noting the temporal correlations present only in real signals, we further strengthened the

sifting procedure with a second criterion: we retained only tracks containing at least 3

consecutive foci. This second criterion reduced the true positive detection rate to 35–50%

(code – Analysis*: plotTrackFrequency*) but increased the positive predictive value (the mean

probability that sifted tracks were correctly identified) from <70% to >95% for single YFP

molecules *in vitro* (Supplementary Fig. 2d). Controls similarly indicate a positive predictive value of >90% for dimers and higher assemblies *in planta* (Supplementary Fig. 2g).

To understand the impact of higher particle densities, we estimated the theoretical probability of random overlap between tracks using a continuum model [89], which estimates the probability of nearest neighbour distances in a random Poisson process falling within the widefield localisation precision. According to this model, the expected fraction of randomly overlapping tracks retained after sifting was <10% for all cases in this study. Conversely for extremely low densities, the higher the SNR threshold is set, the brighter false positives will appear when they do eventually arise. This can be problematic when the true positive rate is very low, for example when estimating characteristic molecular brightness at very low emitter density.

The 2D diffusivity of each track was estimated according to a random walk model as a quarter of the rate of increase of the mean-squared displacement with lag time [98] (code – Analysis*: plotDiffusivity*).

The initial intensity of each track was determined by backward extrapolation of the intensities of its first 5 foci to a virgin timepoint prior to photobleaching.

The characteristic molecular brightness of each fluorescent reporter species was determined based on the Chung-Kennedy-filtered [99,100] terminal intensity of tracks in each acquisition (Supplementary Fig. 1, code – Characteristic Molecular Brightness: *overTrackAll*, *CKfilterBaseline, plotOverTracks*). After calibration, at 10 ms exposure time in plant nuclei at 20 μm working depth, the characteristic molecular brightness was:

VIN3-GFP:	103 ± 9 photons (172 ± 15 counts, mean ± sem),
VRN5-YFP and LacI-YFP:	76 ± 10 photons (126 ± 16 counts),
VIN3-SYFP2:	79 ± 15 photons (131 ± 25 counts) and
VRN5-mScarlet-I:	84 ± 11 photons (140 ± 18 counts).

Previous Slimfield work gives comparable values ranging from 60–250 photons per frame, albeit for fluorescent proteins within 1 μm from the coverslip surface [61,89,48]. For a quantitative comparison at greater working depths, *in vitro* values were obtained from recombinant YFP on a tilted surface under the same SlimVar imaging conditions (Supplementary Fig. 2f). The drops in characteristic molecular brightness imply photon collection losses of 25% over the working depth, compared to 30% in the root tissue itself. This corresponds to an average emission decay length in tissue of about 60 μm, which limits the range of working depths across which a given characteristic molecular brightness value can be used within error, to ± 10 μm.

The internally calibrated values listed above were then used to normalise intensity metrics into numbers of labelled molecules. First, the number of molecules associated with each tracked assembly – its **stoichiometry** – is the initial track intensity divided by the characteristic molecular brightness (code – Analysis*: plotStoichiometry*). The same normalisation factor is used for each dataset's stoichiometry **periodicity**, as well as its **total protein number**.

Autofluorescence and total protein number

Raw estimates of the total number of molecules in each nucleus were extracted with an *ImageJ* macro following [45] (code – Total Protein Number: *MonoCropper.ijm*); the pixel values were integrated within each nuclear segment, then normalised by the characteristic molecular brightness to give an integrated nuclear intensity in molecular equivalents (Fig. 4a). These values did not account for the additive contribution from autofluorescence background, which we estimated using the corresponding unlabelled control line, Col*FRI*. The total protein number of each labelled dataset was refined to exclude autofluorescence by taking the difference between mean integrated nuclear intensities of the labelled dataset and unlabelled control, adjusted in proportion to the ratio of mean areas of nuclear segments. The negative control was much brighter in green acquisitions (15,000 ± 1500 GFP equivalents, or 1.5×10^6 photons, vs. 3,800 ± 400 YFP/SYFP2, or 2.8×10^5 photons; mean ± sem, N=33, N=27 respectively). The nucleoplasmic concentrations were estimated by dividing each total protein number by the mean nuclear volume (assuming prolate spheroidal nuclei aligned in the image plane) and multiplying by Avogadro's number.

We quantified the mean autofluorescence intensity of nuclei in wild-type control (Col*FRI*) as 45 and 26 photons/pixel for 488 nm and 514 nm wavelength excitation (n=250 and 71 nuclei) respectively. We show that this is not prohibitive for SlimVar in these tissues by considering our SNR metric: if we divide the 103 or 76 photons from a single molecule (characteristic molecular brightness of GFP or YFP respectively) over the average area of 30 pixels per foci, then divide by the square root of this autofluorescence level, we find expected signal-to-noise ratios of 0.52 and 0.50, which correspond very well to the empirical SNR thresholds 0.35-0.50 used in sifting the single molecules from noise. At the 95th percentile of autofluorescence (109 and 114 photons/pixel), this typical single molecule SNR reduces to 0.24 and 0.33, reflecting the few adverse cases where only oligomers, and not single molecules, can be detected reliably. However, the stoichiometry of those oligomers can still be precisely counted with characteristic molecular brightness obtained elsewhere in regions of sufficiently low local autofluorescence.

Stoichiometry periodicity analysis

To calculate stoichiometry periodicity (code – Analysis: *plotNearestNeighbourPeriodicity*), first the stoichiometries of all tracks within each nucleus were aggregated across nuclei in a dataset (genotype and vernalisation status), then represented as a kernel density distribution. The choice of the kernel width is informed by the empirical variation in the characteristic molecular brightness. The observed range is typically ± 30%, i.e. a width of 0.6 molecules in each frame at the SNR threshold [43], but to avoid oversmoothing during the periodicity analysis, we used the standard error of ± 14%, i.e. 0.3 molecules. Peaks in this distribution were detected using the MATLAB *findpeaks* function, and the intervals between nearest neighbour peaks were calculated. The uncertainty in peak-to-peak interval was estimated as the single-molecule uncertainty of 0.6 molecules multiplied by the square root of the mean stoichiometry, divided by the square root of the number of interpolated intervals [91]. To suppress noise and spurious intra-peak sampling, all peak intervals smaller than the interval uncertainty were discarded. A second kernel density estimate was calculated over the remaining peak intervals, with the interval uncertainty as the kernel

width. This curve describes the distribution of peak intervals in the stoichiometry as shown
in Fig. 5 (insets) and Supplementary Fig. 11d. The modal value of this interval distribution
was reported as the predominant periodicity of assemblies in each dataset. This method of
estimating periodicity was verified as independent of the mean stoichiometry using
simulated positive control data drawn from noisy Poisson-distributed multiples of an
oligomeric ground truth [48]. This analysis reproduced the expectation that the minimum
number of tracks required for sufficient peak sampling, and therefore the limit of periodicity
detection, scales with the square root of the mean stoichiometry. To demonstrate a
negative control (code – Analysis: *simulateControlPeriodicity*), 100 aperiodic sets of 10^4
stoichiometry values, uniformly distributed at random between 1–30 molecules, were
generated using the *randperm* MATLAB function and processed to generate a set of 100
independent interval distribution curves each corresponding to null periodicity. The 95th
percentile fraction of peak intervals was calculated at each interval size to generate a null
curve, below which test data could no longer be considered periodic. The uncertainty in the
reported modal peak interval was estimated as the s.e.m. of the peak intervals falling within
the range above the null threshold line. To avoid undersampling the peaks, a minimum
number of tracks were needed: at least 14 multiplied by the mean stoichiometry. Where
necessary, stoichiometry lists from different replicates were aggregated to provide more
robust datasets for periodicity analysis. The variation of periodicity for each individual
replicate may be estimated by bootstrapping pairs of replicates together to meet the
threshold number of tracks. The periodicity analysis was validated *in vivo* using the
standard of tetrameric LacI-YFP [65] detected in the *FLC-lacO*/LacI-YFP lines (Supplementary
Fig. 10b).

Two-colour imaging and colocalisation

For each nucleus, a z-stack was performed, first with brightfield imaging to ensure
alignment, then with 514 nm wavelength SlimVar excitation at 10 ms/frame exposure to
track *FLC* loci via LacI-YFP. This z-range extended from the highest to lowest surface of each
nucleus with respect to the coverslip surface and was divided into equally spaced intervals
of 280–360 nm. During each stack, the z-position of the image frame (denoted I*)
containing the brightest LacI-YFP foci was noted and subsequently revisited (Fig. 7a). We
then performed a dual-colour SlimVar acquisition, 10 ms exposure time alternating between
561 nm and 514 nm excitation wavelengths to facilitate distinct signals for each reporter
free from bleed-through. The total duration of the fluorescent z-stack and alternating
excitation acquisitions for each nucleus was ≤15 s. We estimate that the maximum
displacement of *FLC* loci during the 15 s period was within 180 nm, less than our optical
resolution limit (Supplementary Fig. 12d) and consistent with previous observations,[101] thus
in effect immobile over this timescale.
During post-processing, the two colour channels were spatially aligned to sub-pixel precision
using affine transforms generated from SlimVar images of 200 nm diameter fluorescent
beads *in vitro* (code - Alignment: *generateBeadTransform*). Then both yellow and red
channel image sequences were tracked independently (Fig. 7b) to generate lists of LacI-YFP
and VRN5-mScI tracks respectively. To account for *FLC* candidates observed in the z-stack

but photobleached prior to the alternating acquisition, a copy of the alternating sequence was generated in which each yellow channel image simply comprised I*. This copy was also tracked, and its list of LacI-YFP tracks appended to the list from the original sequence. Collected LacI-YFP tracks are shown in Fig. 7b (yellow traces).

To exclude false positive detections of *FLC* due to free LacI-YFP, we then selected only slow and bright yellow tracks. Our earlier observations suggest a maximum of 8 *FLC* loci in the entire nucleus (Supplementary Fig. 9). On that basis, we selected the LacI-YFP tracks whose diffusivity was ≤ D_{FLC} (Fig. 6a) and from these, retained the 8 brightest (or all if fewer than 8) tracks in each nucleus. These selected *FLC* tracks, typically 1–4 per nucleus (Fig. 7b, white circles), were run through colocalisation analysis (code – Analysis: *analyseAllTracks*) with the corresponding VRN5-mScI tracks (Fig. 7b, magenta traces) using a previously reported algorithm [89]. Briefly, VRN5 and *FLC* tracks were deemed colocalised if they met an intensity overlap condition [102] of at least 50% (effectively a lateral distance of ~3 pixels or one widefield localisation precision) and remained within a distance of 7 pixels (twice the widefield localisation precision) for ≥3 frames. The high numerical aperture and short depth of field ensured an axial precision better than <220 nm FWHM for all colocalisations. The likelihood of false positive overlaps between VRN5 tracks, and the likelihood of false positive colocalisations (an *FLC* locus being colocalised with a VRN5 assembly by random chance) were both <5%, based on the average initial number density of tracks in each frame (5.2 VRN5 and 3.3 candidate *FLC*) distributed in the nucleoplasm under random point statistics [103]. The residence time of each colocalised track was determined from the number of adjacent colocalised frames [70] (code – Analysis: *plotResidenceTimes*).

Statistics and samples

Comparisons of the frequency of colocalised assemblies (Fig. 7d) used Fisher's exact test (two-tailed). All other pairwise comparisons used the non-parametric Brunner-Munzel test (code – Analysis: *BrunnerMunzelTest*). Sample size and significance are indicated alongside each result. Investigators were not blinded and each acquisition was independent. We predetermined a target sample size of >24 cells total per line per condition, sufficiently powered to detect changes of <1 s.d. in each of the five test variables (number of tracks, total protein number, stoichiometry, periodicity, diffusivity) at a Bonferroni-adjusted significance level of α = 0.05/5 = 0.01. We planned biological replicates of >3 independent cycles of growth and vernalisation, with >3 roots per cycle and >3 cells per root. In the dual line, >10 cells per condition sufficed for estimates of track number and stoichiometry disaggregated by colocalisation. Technical replicates were identified with tracks detected within each nuclear segment. Actual numbers of replicates analysed are detailed in Supplementary Table 2.

Data and Code Availability

All raw and processed imaging data and analysed tracks are available at the BioImage Archive: https://doi.org/10.6019/S-BIAD1217. ADEMScode v2.2 software for tracking analysis in MATLAB, and the corresponding documentation and exemplar data, can be found at https://github.com/alex-payne-dwyer/single-molecule-tools-alpd.

References

Johnstone SE, Gladyshev VN, Aryee MJ, Bernstein BE. Epigenetic clocks, aging, and cancer. *Science* 2022; **378**: 1276–1277.

Varotto S, Tani E, Abraham E, Krugman T, Kapazoglou A, Melzer R *et al.* Epigenetics: possible applications in climate-smart crop breeding. *Journal of Experimental Botany* 2020; **71**: 5223–5236.

Cavalli G, Heard E. Advances in epigenetics link genetics to the environment and disease. *Nature* 2019; **571**: 489–499.

Zetzsche J, Fallet M. To live or let die? Epigenetic adaptations to climate change—a review. *Environ Epigenet* 2024; **10**: dvae009.

Müller J, Kassis JA. Polycomb response elements and targeting of Polycomb group proteins in Drosophila. *Curr Opin Genet Dev* 2006; **16**: 476–484.

Coleman RT, Struhl G. Causal role for inheritance of H3K27me3 in maintaining the off state of a Drosophila HOX gene. *Science* 2017; **356**: eaai8236.

Blackledge NP, Klose RJ. The molecular principles of gene regulation by Polycomb repressive complexes. *Nat Rev Mol Cell Biol* 2021; **22**: 815–833.

Menon G, Schulten A, Dean C, Howard M. Digital paradigm for Polycomb epigenetic switching and memory. *Curr Opin Plant Biol* 2021; **61**: 102012.

Hepworth J, Antoniou-Kourounioti RL, Bloomer RH, Selga C, Berggren K, Cox D *et al.* Absence of warmth permits epigenetic memory of winter in Arabidopsis. *Nature Communications* 2018; **9**: 639.

Antoniou-Kourounioti RL, Hepworth J, Heckmann A, Duncan S, Qüesta J, Rosa S *et al.* Temperature Sensing Is Distributed throughout the Regulatory Network that Controls FLC Epigenetic Silencing in Vernalization. *Cell Systems* 2018; **7**: 643–655.

Yang H, Berry S, Olsson TSG, Hartley M, Howard M, Dean C. Distinct phases of Polycomb silencing to hold epigenetic memory of cold in Arabidopsis. *Science* 2017; **357**: 1142–1145.

Angel A, Song J, Yang H, Questa JI, Dean C, Howard M. Vernalizing cold is registered digitally at FLC. *Proc Natl Acad Sci U S A* 2015; **112**: 4146–4151.

Costa S, Dean C. Storing memories: The distinct phases of Polycomb-mediated silencing of Arabidopsis FLC. *Biochemical Society Transactions* 2019; **47**: 1187–1196.

Berry S, Hartley M, Olsson TSG, Dean C, Howard M. Local chromatin environment of a Polycomb target gene instructs its own epigenetic inheritance. *eLife* 2015; **4**: e07205.

Mylne JS, Barrett L, Tessadori F, Mesnage S, Johnson L, Bernatavichute YV *et al.* LHP1, the Arabidopsis homologue of HETEROCHROMATIN PROTEIN1, is required for epigenetic silencing of

FLC. *Proceedings of the National Academy of Sciences of the United States of America* 2006; **103**: 5012–7.

Saxton DS, Rine J. Epigenetic memory independent of symmetric histone inheritance. *eLife* 2019; **8**: 1–44.

Lövkvist C, Mikulski P, Reeck S, Hartley M, Dean C, Howard M. Hybrid protein assembly-histone modification mechanism for PRC2-based epigenetic switching and memory. *eLife* 2021; **10**: e66454.

Laugesen A, Højfeldt JW, Helin K. Molecular Mechanisms Directing PRC2 Recruitment and H3K27 Methylation. *Molecular Cell* 2019; **74**: 8–18.

Greb T, Mylne JS, Crevillen P, Geraldo N, An H, Gendall AR *et al.* The PHD Finger Protein VRN5 Functions in the Epigenetic Silencing of Arabidopsis FLC. *Current Biology* 2007; **17**: 73–78.

Sung S, Amasino RM. Vernalization in Arabidopsis thaliana is mediated by the PHD finger protein VIN3. *Nature* 2004; **427**: 159–64.

Franco-Echevarría E, Rutherford TJ, Fiedler M, Dean C, Bienz M. Plant vernalization proteins contain unusual PHD superdomains without histone H3 binding activity. *Journal of Biological Chemistry* 2022; **298**. doi:10.1016/j.jbc.2022.102540.

Kim DH, Sung S. Coordination of the vernalization response through a VIN3 and FLC gene family regulatory network in Arabidopsis. *Plant Cell* 2013; **25**: 454–69.

Fiedler M, Franco-Echevarría E, Schulten A, Nielsen M, Rutherford TJ, Yeates A *et al.* Head-to-tail polymerization by VEL proteins underpins cold-induced Polycomb silencing in flowering control. *Cell Rep* 2022; **41**: 111607.

Hepler PK, Gunning BES. Confocal fluorescence microscopy of plant cells. *Protoplasma* 1998; **201**: 121–157.

Shaw SL, Thoms D, Powers J. Structured illumination approaches for super-resolution in plant cells. *Microscopy (Oxf)* 2019; **68**: 37–44.

Ovecka M, Sojka J, Tichá M, Komis G, Basheer J, Marchetti C *et al.* Imaging plant cells and organs with light-sheet and super-resolution microscopy. *Plant Physiology* 2022; **188**: 683–702.

Guo AY, Zhang YM, Wang L, Bai D, Xu YP, Wu WQ. Single-molecule imaging in living plant cells: A methodological review. *International Journal of Molecular Sciences* 2021; **22**: 5071.

Dumur T, Duncan S, Graumann K, Desset S, Randall RS, Scheid M *et al.* Probing the 3D architecture of the plant nucleus with microscopy approaches: challenges and solutions. *Nucleus* 2019; **10**: 181–212.

Colin L, Martin-Arevalillo R, Bovio S, Bauer A, Vernoux T, Caillaud MC *et al.* Imaging the living plant cell: From probes to quantification. *Plant Cell* 2022; **34**: 247–272.

30 Wang X, Li X, Deng X, Luu DT, Maurel C, Lin J. Single-molecule fluorescence imaging to quantify
membrane protein dynamics and oligomerization in living plant cells. *Nature Protocols* 2015; **10**:
2054–63.

31 Cui Y, Yu M, Yao X, Xing J, Lin J, Li X. Single-Particle Tracking for the Quantification of Membrane
Protein Dynamics in Living Plant Cells. *Molecular Plant* 2018; **11**: 1315–1327.

32 Prunet N, Duncan K. Imaging flowers: a guide to current microscopy and tomography techniques
to study flower development. *Journal of Experimental Botany* 2020; **71**: 2898–2909.

33 Platre MP, Bayle V, Armengot L, Bareille J, Marquès-Bueno M del M, Creff A *et al.*
Developmental control of plant Rho GTPase nano-organization by the lipid phosphatidylserine.
Science 2019; **364**: 57–62.

34 Zhang X, Cui Y, Yu M, Lin J. Single-Molecule Techniques for Imaging Exo-Endocytosis Coupling in
Cells. *Trends Plant Sci* 2019; **24**: 879–880.

35 Chen B-C, Legant WR, Wang K, Shao L, Milkie DE, Davidson MW *et al.* Lattice light-sheet
microscopy: Imaging molecules to embryos at high spatiotemporal resolution. *Science* 2014;
346: 1257998.

36 Liu T-L, Upadhyayula S, Milkie DE, Singh V, Wang K, Swinburne IA *et al.* Observing the cell in its
native state: Imaging subcellular dynamics in multicellular organisms. *Science* 2018; **360**:
eaaq1392.

37 Schmidt R, Weihs T, Wurm CA, Jansen I, Rehman J, Sahl SJ *et al.* MINFLUX nanometer-scale 3D
imaging and microsecond-range tracking on a common fluorescence microscope. *Nat Commun*
2021; **12**: 1478.

38 Deguchi T, Iwanski MK, Schentarra E-M, Heidebrecht C, Schmidt L, Heck J *et al.* Direct
observation of motor protein stepping in living cells using MINFLUX. *Science* 2023; **379**: 1010–
1015.

39 Bayle V, Fiche JB, Burny C, Platre MP, Nollmann M, Martinière A *et al.* Single-particle tracking
photoactivated localization microscopy of membrane proteins in living plant tissues. *Nature*
Protocols 2021; **16**: 1600–1628.

40 Lummer M, Humpert F, Wiedenlübbert M, Sauer M, Schüttpelz M, Staiger D. A New Set of
Reversibly Photoswitchable Fluorescent Proteins for Use in Transgenic Plants. *Molecular Plant*
2013; **6**: 1518–1530.

41 Iwatate RJ, Yoshinari A, Yagi N, Grzybowski M, Ogasawara H, Kamiy M *et al.* Covalent Self-
labeling of tagged proteins with chemical fluorescent dyes in BY-2 cells and arabidopsis
seedlings. *Plant Cell* 2020; **32**: 3081–3094.

42 Yagi N, Yoshinari A, Iwatate RJ, Isoda R, Frommer WB, Nakamura M. Advances in Synthetic
Fluorescent Probe Labeling for Live-Cell Imaging in Plants. *Plant and Cell Physiology* 2021; **62**:
1259–1268.

43 Leake MC, Chandler JH, Wadhams GH, Bai F, Berry RM, Armitage JP. Stoichiometry and turnover
in single, functioning membrane protein complexes. *Nature* 2006; **443**: 355–358.

44 Plank M, Wadhams GH, Leake MC. Millisecond timescale slimfield imaging and automated
quantification of single fluorescent protein molecules for use in probing complex biological
processes. *Integrative Biology* 2009; **1**: 602–612.

45 Wollman AJM, Leake MC. Millisecond single-molecule localization microscopy combined with
convolution analysis and automated image segmentation to determine protein concentrations
in complexly structured, functional cells, one cell at a time. *Faraday Discussions* 2015; **184**: 401–
424.

46 Miller H, Cosgrove J, Wollman AJM, Taylor E, Zhou Z, O'Toole PJ *et al.* High-speed single-
molecule tracking of CXCL13 in the B-follicle. *Frontiers in Immunology* 2018; **9**: 1073.

47 Shashkova S, Wollman A, Hohmann S, Leake MC. Characterising Maturation of GFP and mCherry
of Genomically Integrated Fusions in Saccharomyces cerevisiae. *Bio-Protocol* 2018; **8**: e.2710.

48 Payne-Dwyer AL, Syeda AH, Shepherd JW, Frame L, Leake MC. RecA and RecB: probing
complexes of DNA repair proteins with mitomycin C in live Escherichia coli with single-molecule
sensitivity. *Journal of The Royal Society Interface* 2022; **19**: 20220437.

49 Dresser L, Hunter P, Yendybayeva F, Hargreaves AL, Howard JAL, Evans GJO *et al.* Amyloid-β
oligomerization monitored by single-molecule stepwise photobleaching. *Methods* 2021; **193**:
80–95.

50 Konopka CA, Bednarek SY. Variable-angle epifluorescence microscopy: A new way to look at
protein dynamics in the plant cell cortex. *Plant Journal* 2008; **53**: 186–96.

51 Tokunaga M, Imamoto N, Sakata-Sogawa K. Highly inclined thin illumination enables clear single-
molecule imaging in cells. *Nature Methods* 2008; **5**: 159–61.

52 Gardini L, Vignolini T, Curcio V, Pavone FS, Capitanio M. Optimization of highly inclined
illumination for diffraction-limited and super-resolution microscopy. *Opt Express* 2023; **31**:
26208–26225.

53 Reihani SNS, Charsooghi MA, Khalesifard HR, Golestanian R. Efficient in-depth trapping with an
oil-immersion objective lens. *Opt Lett* 2006; **31**: 766–768.

54 Reihani SNS, Mir SA, Richardson AC, Oddershede LB. Significant improvement of optical traps by
tuning standard water immersion objectives. *Journal of Optics* 2011; **13**. doi:10.1088/2040-
8978/13/10/105301.

55 Dienerowitz M, Gibson G, Bowman R, Padgett M. Holographic aberration correction: optimising
the stiffness of an optical trap deep in the sample. *Opt Express* 2011; **19**: 24589–24595.

56 Schwertner M, Booth MJ, Wilson T. Simple optimization procedure for objective lens correction
collar setting. *J Microsc* 2005; **217**: 184–187.

57 Ue Y, Monai H, Higuchi K, Nishiwaki D, Tajima T, Okazaki K *et al.* A spherical aberration-free
microscopy system for live brain imaging. *Biochem Biophys Res Commun* 2018; **500**: 236–241.

58 Kubitscheck U, Kückmann O, Kues T, Peters R. Imaging and Tracking of Single GFP Molecules in
Solution. *Biophysical Journal* 2000; **78**: 2170–2179.

59 Zhao Q, Young IT, de Jong JGS. Photon budget analysis for fluorescence lifetime imaging
microscopy. *J Biomed Opt* 2011; **16**: 086007.

60 Qüesta JI, Antoniou-Kourounioti RL, Rosa S, Li P, Duncan S, Whittaker C *et al.* Noncoding SNPs
influence a distinct phase of Polycomb silencing to destabilize long-term epigenetic memory at
Arabidopsis FLC. *Genes and Development* 2020; **34**: 446–461.

61 Payne-Dwyer AL, Leake MC. Single-molecular quantification of flowering control proteins within
nuclear condensates in live whole Arabidopsis root. In: Leake MC (ed). *Chromosome*
Architecture: Methods in Molecular Biology. Springer US, Humana: New York, NY, 2022, p 2476.

62 Dittmer TA, Stacey NJ, Sugimoto-Shirasu K, Richards EJ. LITTLE NUCLEI genes affecting nuclear
morphology in Arabidopsis thaliana. *Plant Cell* 2007; **19**: 2793–2803.

63 Leake MC. Analytical tools for single-molecule fluorescence imaging in cellulo. *Physical*
Chemistry Chemical Physics 2014; **16**: 12635–47.

64 Adler L, Lau CS, Shaikh KM, van Maldegem KA, Payne-Dwyer AL, Lefoulon C *et al.* The role of
BST4 in the pyrenoid of Chlamydomonas reinhardtii. *bioRxiv* 2023; : 2023.06.15.545204.

65 Lewis M, Chang G, Horton NC, Kercher MA, Pace HC, Schumacher MA *et al.* Crystal structure of
the lactose operon repressor and its complexes with DNA and inducer. *Science* 1996; **271**: 1247–
1254.

66 Wollman AJM, Syeda AH, Howard JAL, Payne-Dwyer A, Leech A, Warecka D *et al.* Tetrameric
UvrD Helicase Is Located at the E. Coli Replisome due to Frequent Replication Blocks. *J Mol Biol*
2024; **436**: 168369.

67 Rosa S, De Lucia F, Mylne JS, Zhu D, Ohmido N, Pendle A *et al.* Physical clustering of FLC alleles
during polycomb-mediated epigenetic silencing in vernalization. *Genes and Development* 2013;
27: 1845–50.

68 Ladouceur A-M, Parmar BS, Biedzinski S, Wall J, Tope SG, Cohn D *et al.* Clusters of bacterial RNA
polymerase are biomolecular condensates that assemble through liquid-liquid phase separation.
Proc Natl Acad Sci U S A 2020; **117**: 18540–18549.

69 Bindels DS, Haarbosch L, van Weeren L, Postma M, Wiese KE, Mastop M *et al.* mScarlet: a bright
monomeric red fluorescent protein for cellular imaging. *Nat Methods* 2017; **14**: 53–56.

70 Stracy M, Wollman AJM, Kaja E, Gapinski J, Lee JE, Leek VA *et al.* Single-molecule imaging of DNA
gyrase activity in living Escherichia coli. *Nucleic Acids Research* 2019; **47**: 210–220.

71 Zhan H, Stanciauskas R, Stigloher C, Keomanee-Dizon K, Jospin M, Bessereau J-L *et al.* In vivo
single-molecule imaging identifies altered dynamics of calcium channels in dystrophin-mutant C.
elegans. *Nat Commun* 2014; **5**: 4974.

72 Ritter JG, Veith R, Veenendaal A, Siebrasse JP, Kubitscheck U. Light Sheet Microscopy for Single
Molecule Tracking in Living Tissue. *PLOS ONE* 2010; **5**: e11639.

73 Reisser M, Palmer A, Popp AP, Jahn C, Weidinger G, Gebhardt JCM. Single-molecule imaging
correlates decreasing nuclear volume with increasing TF-chromatin associations during zebrafish
development. *Nat Commun* 2018; **9**: 5218.

74 Li X, Wang X, Yang Y, Li R, He Q, Fang X *et al.* Single-Molecule Analysis of PIP2;1 Dynamics and
Partitioning Reveals Multiple Modes of Arabidopsis Plasma Membrane Aquaporin Regulation.
Plant Cell 2011; **23**: 3780–3797.

75 Song K, Xue Y, Wang X, Wan Y, Deng X, Lin J. A modified GFP facilitates counting membrane
protein subunits by step-wise photobleaching in *Arabidopsis*. *Journal of Plant Physiology* 2017;
213: 129–133.

76 Pain C, Tynan C, Botchway SW, Kriechbaumer V. Variable-Angle Epifluorescence Microscopy for
Single-Particle Tracking in the Plant ER. *Methods Mol Biol* 2024; **2772**: 273–283.

77 von Arx M, Xhelilaj K, Schulz P, zur Oven-Krockhaus S, Gronnier J. Photochromic reversion
enables long-term tracking of single molecules in living plants. 2024; : 2024.04.10.585335.

78 Rohr L, Ehinger A, Rausch L, Glöckner Burmeister N, Meixner AJ, Gronnier J *et al.* OneFlowTraX: a
user-friendly software for super-resolution analysis of single-molecule dynamics and nanoscale
organization. *Front Plant Sci* 2024; **15**. doi:10.3389/fpls.2024.1358935.

79 Tang J, Han KY. Extended field-of-view single-molecule imaging by highly inclined swept
illumination. *Optica* 2018; **5**: 1063.

80 Ren J, Han KY. 2.5D Microscopy: Fast, High-Throughput Imaging via Volumetric Projection for
Quantitative Subcellular Analysis. *ACS Photonics* 2021; **8**: 933–942.

81 Wollman AJM, Hedlund EG, Shashkova S, Leake MC. Towards mapping the 3D genome through
high speed single-molecule tracking of functional transcription factors in single living cells.
Methods 2020; **170**: 82–89.

82 Miura N, Ashida Y, Matsuda Y, Shibuya T, Tamada Y, Hatsumi S *et al.* Adaptive Optics Microscopy
with Wavefront Sensing Based on Neighbor Correlation. *Plant and Cell Physiology* 2023; **64**:
1372–1382.

83 Jin X, Lee J-EJE, Schaefer C, Luo X, Wollman AJM, Payne-Dwyer AL *et al.* Membraneless
organelles formed by liquid-liquid phase separation increase bacterial fitness. *Science Advances*
2021; **7**: eabh2929.

84 Schulten A, Jang G-J, Payne-Dwyer A, Nielsen ML, Bienz M, Leake MC *et al.* Functional
specialization of Arabidopsis VEL polymerization domains in the switch to Polycomb silencing.
2024; : 2024.02.15.580496.

85   Qüesta JI, Song J, Geraldo N, An H, Dean C. Arabidopsis transcriptional repressor VAL1 triggers
Polycomb silencing at FLC during vernalization. *Science* 2016; **353**: 485–488.

86   Jones JDG, Shlumukov L, Carland F, English J, Scofield SR, Bishop GJ *et al.* Effective vectors for
transformation, expression of heterologous genes, and assaying transposon excision in
transgenic plants. *Transgenic Research* 1992; **1**: 285–297.

87   Box MS, Coustham V, Dean C, Mylne JS. Protocol: A simple phenol-based method for 96-well
extraction of high quality RNA from Arabidopsis. *Plant Methods* 2011; **7**: 7.

88   Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T *et al.* Fiji: an open-
source platform for biological-image analysis. *Nat Methods* 2012; **9**: 676–682.

89   Syeda AH, Wollman AJM, Hargreaves AL, Howard JAL, Brüning J-G, McGlynn P *et al.* Single-
molecule live cell imaging of Rep reveals the dynamic interplay between an accessory replicative
helicase and the replisome. *Nucleic Acids Research* 2019; **47**: 6287–6298.

90   Shepherd JW, Payne-Dwyer AL, Lee J-E, Syeda A, Leake MC. Combining single-molecule super-
resolved localization microscopy with fluorescence polarization imaging to study cellular
processes. *JPhys Photonics* 2021; **3**. doi:10.1088/2515-7647/ac015d.

91   Hunter P, Payne-Dwyer AL, Shaw M, Signoret N, Leake MC. Single-molecule and super-resolved
imaging deciphers membrane behavior of onco-immunogenic CCR5. *iScience* 2022; **25**: 105675.

92   Grossmann G, Krebs M, Maizel A, Stahl Y, Vermeer JEM, Ott T. Green light for quantitative live-
cell imaging in plants. *Journal of Cell Science* 2018; **131**: jcs209270.

93   Thompson RE, Larson DR, Webb WW. Precise nanometer localization analysis for individual
fluorescent probes. *Biophysical Journal* 2002; **82**: 2775–83.

94   Llorente-Garcia I, Lenn T, Erhardt H, Harriman OL, Liu L-N, Robson A *et al.* Single-molecule in vivo
imaging of bacterial respiratory complexes indicates delocalized oxidative phosphorylation.
Biochimica et Biophysica Acta (BBA) - Bioenergetics 2014; **1837**: 811–824.

95   Shepherd JW, Higgins EJ, Wollman AJM, Leake MC. PySTACHIO: Python Single-molecule TrAcking
stoiCHiometry Intensity and simulatiOn, a flexible, extensible, beginner-friendly and optimized
program for analysis of single-molecule microscopy data. *Computational and Structural
Biotechnology Journal* 2021; **19**: 4049–4058.

96   Miller H, Zhou Z, Wollman AJM, Leake MC. Superresolution imaging of single DNA molecules
using stochastic photoblinking of minor groove and intercalating dyes. *Methods* 2015; **88**: 81–
88.

97   Lecinski S, Shepherd JW, Bunting K, Dresser L, Quinn SD, MacDonald C *et al.* Correlating viscosity
and molecular crowding with fluorescent nanobeads and molecular probes: in vitro and in vivo.
Interface Focus 2022; **12**: 20220042.

98   Robson A, Burrage K, Leake MC. Inferring diffusion in single live cells at the single-molecule
level. *Philosophical Transactions of the Royal Society B: Biological Sciences* 2013; **368**: 20120029.

99 Chung SH, Kennedy RA. Forward-backward non-linear filtering technique for extracting small
biological signals from noise. *Journal of Neuroscience Methods* 1991; **40**: 71–86.

100 Leake MC, Wilson D, Bullard B, Simmons RM, Bubb MR. The elasticity of single kettin
molecules using a two-bead laser-tweezers assay. In: *FEBS Letters*. 2003, pp 55–60.

101 Meschichi A, Rosa S. Visualizing and Measuring Single Locus Dynamics in Arabidopsis
thaliana. In: *Arabidopsis Protocols: Methods in Molecular Biology*. 2021, pp 213–224.

102 Wollman AJM, Fournier C, Llorente-Garcia I, Harriman O, Payne-Dwyer AL, Shashkova S *et al.*
Critical roles for EGFR and EGFR-HER2 clusters in EGF binding of SW620 human carcinoma cells.
Journal of The Royal Society Interface 2022; **19**: 20220088.

103 Payne-Dwyer A, Kumar G, Barrett J, Gherman LK, Hodgkinson M, Plevin M *et al.* Predicting
Rubisco-Linker Condensation from Titration in the Dilute Phase. *Phys Rev Lett* 2024; **132**:
218401.

104 Faklaris O, Bancel-Vallée L, Dauphin A, Monterroso B, Frère P, Geny D *et al.* Quality
assessment in light microscopy for routine use through simple tools and robust metrics. *J Cell
Biol* 2022; **221**: e202107093.

105 Wilson T. Resolution and optical sectioning in the confocal microscope. *J Microsc* 2011; **244**:
113–121.

1507

1508

Acknowledgements

We would like to thank Ibrahim Cissé and Jamie Hobbs for preliminary discussions, Anna Schulten, Govind Menon, Martin Howard, Pan Zhu, Cecilia Lövkvist, Grant Calder and Aisha Syeda for training and discussions, Mathias Nielsen and Yaoxi Li for generating the VRN5-mScarlet-I and VIN3-SYFP2 constructs, and the Horticulture and Bioscience Technology Facility at the University of York for supporting growth chamber resources and confocal microscopy. Funded by EPSRC (grants EP/T00214X/1, EP/T002166/1 and EP/Y000501/1). For the purpose of open access, a Creative Commons Attribution (CC BY) licence is applied to any Author Accepted Manuscript version arising from this submission.

Author Contributions

Conceptualization, funding acquisition, administration, supervision; ML, CD. Methodology, software and validation: APD, ML; Investigation, data curation, analysis and visualization: APD, GJJ; Writing – original draft preparation: APD; Writing – review & editing: APD, GJJ, CD, ML.

Competing interests

The authors declare no competing interests.

 # Supplementary Information

1529

Supplementary Table 1. Average stoichiometry of VIN3-GFP, VIN3-SYFP2 and VRN5-YFP assemblies whose diffusivity matches that of *FLC* loci. The diffusivity-matched stoichiometry is distributed similarly to the population (VIN3-SYFP2 V6W, N=296, p=0.08 |ns), and similarly increases with vernalisation for both VIN3 (V2W to V6W, SYFP2: N=262, p<0.001 |**; GFP: N=302, p<0.001 |**) and VRN5 (NV to V6W+T14, N=318, p<0.001 |**). VIN3-SYFP2 values are corrected for the fraction of endogenous VIN3 (Supplementary Fig. 5). NV = not vernalised; V2W = two weeks of cold; V6W = six weeks of cold; V6WT14 = six weeks of cold followed by two weeks of warm conditions.

1538

Line	Timepoint	*FLC*-like tracks $D \approx D_{FLC}$		Stoichiometry (molecules)	
		Number	% of total (± s.e.m.)	Median	IQR
VIN3-GFP	V2W	151	15.2 (1.2)	9.9	5.9–16.3
	V6W	262	17.6 (1.1)	15.1	9.0–26.0
VIN3-SYFP2	V2W	131	11.3 (1.0)	3.2	2.0–6.0
				6.4	3.9–11.9 *corrected*
	V6W	148	16.5 (1.4)	5.2	3.8–8.4
				10.4	7.6–16.8 *corrected*
VRN5-YFP	NV	203	18.4 (1.3)	11.3	7.3–23.4
	V2W	168	17.4 (1.3)	17.7	11.1–29.9
	V6W	355	20.8 (1.1)	18.1	11.5–30.2
	V6W+T14	249	19.6 (1.2)	20.4	12.7–31.0

1539

1540

1541 **Supplementary Table 2. Sample statistics used for quantitative SlimVar analysis.** Numbers
1542 of nuclei and tracks quantified. Col*FRI* acquisitions were aggregated over all tested
1543 vernalisation conditions: NV = not vernalised; V2W = two weeks of cold; V6W = six weeks of
1544 cold; V6WT7 = six weeks of cold followed by one week of warm conditions; V6WT14 = six
1545 weeks of cold followed by two weeks of warm conditions.

1546

Line	Vernalisation / imaging condition	Independent growth cycles	Cells quantified	Nuclear tracks detected
VIN3-SYFP2	NV	3	22	0
	V2W	4	71	1,158
	V6W	3	46	896
	V6W T7	3	23	0
VIN3-GFP	NV	3	24	0
	V2W	5	57	994
	V6W	4	83	1,485
	V6W T7	4	36	0
VRN5-EYFP	NV	4	50	1,039
	V2W	3	36	964
	V6W	4	63	1,703
	V6W T14	4	54	1,285
VRN5-mScarletI	NV	5	44	552
	V2W	6	29	426
	V6W	2	16	256
	V6W T14	3	29	365
*FLC-lacO/*LacI-YFP	NV	3	38	166
VRN5-mScarletI × *FLC-lacO/*LacI-YFP	NV	3	10	365 × 23
	V2W	2	28	1,703 × 156
	V6W	3	25	2,867 × 192
	V6W T14	2	18	1,416 × 74
Col*FRI*	All (green / 488 nm)	3	79	0
	All (yellow / 514 nm)	3	38	0
	All (red / 561 nm)	3	79	0

1547
1548
1549

 Supplementary Table 3. Primers used in this study.

Primer name	Primer sequence 5' -> 3'
Primers for cloning	
Mega-pENTR VIN3-mTurq2 F	GCGGCAGCTTCAGCCGTGAGCAAGGGCGAGGAG
Mega-pENTR VIN3-mTurq2 R	GAGTTAATAAATAGACCGATTTCAAAAGCTTTTACTTGTACAGCTCGTCCATG
VRN5_mScarlet_T	GCATCCGCGGCAGCTTCAGCCGTGAGCAAGGGCGAGGCA
VRN5_mScarlet_B	GTTGAATCAATGGGAAGCTTGATTACTTGTACAGCTCGTCCATG
qRT-PCR primers	
FP-qPCR-F2	GGACGACGGCAACTACAAGA
FP-qPCR-R2	GTTGCCGTCCTCCTTGAAG
VIN3-qPCR-1F	TGCTTGTGGATCGTCTTGTCA
VIN3-qPCR-1R	TTCTCCAGCATCCGAGCAAG
VRN5_qPCR_F	GAGGCATTGTTAGTAGGCTTCC
VRN5_qPCR_R	CACGCAGCTGGAACTAAATCTC
FLC spliced F	AGCCAAGAAGACCGAACTCA
FLC spliced R	TTTGTCCAGCAGGTGACATC
UBC_qPCR_F	CTGCGACTCAGGGAATCTTCTAA
UBC_qPCR_R	TTGTGCCATTGAATTGAACCC
PP2A F2	ACTGCATCTAAAGACAGAGTTCC
PP2A R2	CCAAGCATGGCCGTATCATGT

1553

Supplementary Fig. 1. Determination of characteristic molecular brightness in live tissues from photobleaching. Intensity of foci (black circles) near the end of a single track in a post-vernalised VRN5-YFP image sequence, interpolated using a Chung-Kennedy edge-preserving filter (centred window of 3 points, blue line). With sufficient signal-to-noise, photobleaching enables counting of individual steps corresponding to the presence of individual labelled molecules in an assembly. The consistent photobleaching steps from fluorescent fusions provide enough resolution to count not only molecular steps in specific traces, but also to derive an ensemble estimate of the characteristic molecular brightness for working depths of 20 ± 10 μm. The steps yield an estimate of 76 ± 10 photons for the characteristic single-molecule brightness of the VRN5-YFP fusion *in planta*. Normalisation of the intensity trace by this value gives the absolute number of active fluorophore reporters over the entire course of photobleaching, which remains robust when individual steps would be missed or otherwise unidentifiable. A backward extrapolation of the first five points accounts for photobleaching and allows thee track stoichiometry to be estimated with better precision than the individual foci.

1569

Supplementary Fig. 2, **SlimVar detects and quantifies single fluorescent proteins in tracks**
above threshold levels of excitation power and signal-to-noise. a) A channel with a tilted
internal coverslip surface facilitated observation of single particles *in vitro* at a range of
working depths. Acquisition and tracking of a purified YFP sample were performed both at
zero depth in epifluorescence mode (10 ms exposures) and at a chosen calibration depth of
25 µm in SlimVar mode (10 ms exposures).

If the imaged surface is sufficiently tilted by an angle θ, the depth-of-field (d_{DOF}, the axial
extent of the detection volume) limits the width of the region in focus to $d_{DOF} \cot\theta$ (inset
scale bar 2 µm). At 25 µm working depth in SlimVar, we found the width to be
6,500 ± 300 nm, thus d_{DOF} = 530 ± 30 nm. At the emission wavelength of 528 nm, this result
indicated an effective numerical aperture [104] of 1.38 ± 0.02 (from the nominal NA 1.49),
thereby achieving better light collection than possible with any water immersion lens
(NA < 1.33).

To create the channel, a 3.2 × 22 mm section of #1.5 coverslip was placed between 5 ×
22 mm sections of adhesive spacer of 260 µm depth (125 µL GeneFrame, ThermoFisher)
mounted on a standard 25 × 75 mm slide. Additional coverslip sections were added to the

spacers to compensate for the thickness of the tilted slip, before sealing with a final

coverslip used for imaging. The formed channel was incubated in 1 µg/ml anti-YFP in

phosphate buffer saline (PBS) for 5 min, washed with PBS, followed by 50 nM mYFP in PBS

for 5 min, then washed with PBS. For imaging, the channel was placed so that the surface's

direction of tilt away from the optical axis was orthogonal to the plane of the beam delivery.

b) Examples of YFP foci observed within a 300 ms window (20 mW excitation, 10 ms

exposure per frame), showing similar performance in epifluorescence at zero depth, and in

SlimVar at the calibration depth. The temporal recolouring indicates how foci appear (bind),

diffuse in short, confined trajectories and disappear (photobleach or unbind); scale bar 2

µm.

c) The appropriate SNR threshold for sifting can be estimated as the SNR above which single

molecules are unlikely to be false positives, here at 0.5. This panel shows the frequency

distribution of the signal-to-noise metric for individual tracks detected above a nominal

capture SNR of 0.2, in 10 fields of view at different excitation powers and working depths

after SlimVar calibration: 20 mW power at the coverslip, or zero working depth (solid

black), at the calibration working depth of 25 µm at the working excitation powers of

20 mW (yellow) and lower powers of 16 mW (brown), 8 mW (orange) and 1 mW (red). The

distribution at zero power is derived from tracking simulations of Gaussian noise matched to

the experimental dark detector noise evaluated over three fields of view. This has a

standard deviation of 2.3 photons/pixel, which is equivalent to an average background of

about 15 photons/pixel.

d) After sifting above the SNR threshold of 0.5, the tracks have a distribution of lengths

(number of foci) which varies by modality and working depth. 'Epifluorescence' is

performed at 0^o beam incidence at 20 mW source power (blue), while 'SlimVar' refers to 60^o

beam incidence at 20 mW source power after the calibration protocol (yellow). The tracking

length for epifluorescence deteriorates with increasing working depth (dark blue vs light

blue), while those of SlimVar are less dependent on depth (dark yellow vs light yellow). The

tracks of length 3 or more have a much lower overlap with false positives from noise (black),

which motivates us to include a minimum track length of 3 in the sifting process.

e) Lengths of tracks detected in plants using SlimVar (LacI-YFP, brown; VRN5-YFP, orange;

VIN3-SYFP2, yellow; noise, black). The track lengths for assemblies are correspondingly

greater for species with higher mean stoichiometry and lower mean diffusivity.

f) The characteristic molecular brightness, estimated from the mean photons per frame in

detected tracks, as a function of excitation power. The detector noise generates a floor

value of 50 photons/frame, from the few false positives that inevitably pass the sifting step.

For epifluorescence at depth (blue bars), the high background immediately causes the

apparent value to increase with excitation power, but for isolated molecules, this is not well

correlated with true positive events (panel d). For SlimVar at depth (yellow bars), the

background is low and there is a clear onset above a critical power of ~8 mW (irradiance ~4

kW/cm^2) corresponding to the true number of photons detected from a single fluorescent

protein. The value is highest at zero working depth (rightmost bar), decreasing slightly at the

calibration working depth, and drops further due to scattering and aberrations in plants

(orange), yet remains above the floor level.

g) The distribution of the total number of photons associated with each track for optimal epifluorescence at the surface and 20 mW source power (blue), optimal SlimVar at 25 μm working depth and 20 mW source power (yellow), compared with those of SlimVar at 1 mW source power (red), below the requirement for single molecule detection, and the simulated detector noise (black).

h) The total number of photons per track in vivo, plotted on a logarithmic scale (LacI-YFP, brown; VRN5-YFP, orange). In vitro controls are also shown for comparison (single molecule YFP, yellow; noise, black). The total increases for states with high stoichiometry and low diffusivity. Distinct peaks can be seen at low values, corresponding to single molecules integrated over a small discrete number of frames.

i) The signal-to-noise improvement in vivo due to the SlimVar modality for tracks of VRN5-YFP in plants. Epifluorescence is performed at a longer exposure of 50 ms/frame, while SlimVar is at the balanced mode (10 ms/frame). The SNR values shown account for the additional exposure time required by dividing the raw values for epifluorescence by a factor of $(50 \text{ ms}/10 \text{ ms})^{0.5}$. SlimVar detects additional tracks that are lower stoichiometry and higher diffusivity including single molecules, that for epifluorescence would have raw SNR below the sifting threshold of 0.5 and are therefore invisible. Even with epifluorescence showing this bias towards brighter particles, the mean signal-to-noise in SlimVar track is still at least 2.6× greater.

Supplementary Fig. 3. SlimVar sufficiently mitigates aberrations associated with high-numerical aperture imaging at a specified calibration depth.

a) Individual PSFs shown in axial cross section, acquired as volume stacks from either subdiffractive fluorescent beads (24 nm diameter FluoSpheres) suspended in 1 wt.% agarose at 488 nm wavelength excitation at 20 nm z-step, or LacI-YFP foci in root tip nuclei with 514 nm excitation at 250 nm step. Scale bar 1 μm. All PSFs are spatially upsampled to 20 nm (linear interpolation), background-subtracted and normalised to the same total pixel intensity. Top and bottom rows correspond to independent replicate foci. The 'uncorrected state' corresponds to a standard epifluorescence configuration (beam delivery angle of 0° and correction collar set to the coverslip thickness). The 'corrected' state refers to SlimVar including optimal settings of beam delivery angle (60° in water), telescope lens (32 mm towards objective) and correction collar (coverslip thickness - 30 μm). The 'overcorrected' measurement retains the oblique beam delivery angle and standard tube lens position, but

the correction collar is adjusted by an equivalent amount in the opposite direction (coverslip
thickness +30 µm) to indicate the impact of the spherical aberration removed in SlimVar.
The examples *in planta* are of FLC foci identified in z-stacks (Fig. 7a) using a 'corrected'
SlimVar configuration. (b) Lateral and (c) axial FWHM dimensions determined from the PSF
volume stacks (n=2–4) using MetroloJ-QC (Image J plugin).[104] The axial FWHMs of beads
prior to correction at zero depth, and 'corrected state' at the calibration depth, agree
qualitatively with the theoretical diffraction limited resolution [105] of d_z = 0.89
$\lambda/(n - \sqrt{n^2 - NA^2})$ = 380 nm, and the empirical depth-of-field estimate of 530 nm
(Supplementary Fig. 2a) respectively. The axial improvement demonstrated in beads is
necessary and sufficient for feasible imaging at depths up to 30 µm inside plant tissue, in
which the PSF is subject to additional aberration and scattering.

1679
1680
1681
1682

488 nm   514 nm

1683
1684

Supplementary Fig. 4. Autofluorescence controls for SlimVar microscopy.

a) SlimVar images of non-vernalised nuclei in the Col*FRI* wild-type (unlabelled) line; top,
brightfield; bottom: SlimVar average projections over 100 frames, with contrast set to 120–
370 counts (12–162 photons/pixel). Plastids and cell walls have higher autofluorescence
than the nucleus, especially at 488 nm (bottom left), but do not contribute to false positives
as they are removed during segmentation of the nucleus with the brightfield image. Scale
bar: 5 µm. b) The mean number of background fluorescence photons per 53 nm pixel per
10 ms exposure in each nucleus for 488 nm and 514 nm excitation modes, aggregated over
all vernalisation timepoints. The mean ± s.e.m. (s.d.) autofluorescence is 45 ± 4 (34)
photons/pixel, or 26 ± 4 (33) photons/pixel under 488 nm and 514 nm (n=250 and 71 nuclei)
respectively.

1696

1697

Supplementary Fig. 5. RNA expression analysis and characterization of VIN3-SYFP2.

Expression level of *VIN3* (A), fluorescent protein (B) in VIN3-GFP, VIN3-SYFP2 and Col*FRI*
seedlings. (C) Expression level of spliced *FLC* in seedlings after 2 weeks (V2W) and 6 weeks
vernalised (V6W) cold seedlings relative to non-vernalised (NV) seedlings. (A-C) Data are
presented as mean ± s.e.m. (N=9) from three biologically independent experiments. (D)
Based on the assumption that protein levels scale proportionally with RNA levels, we derive
a correction factor to infer the total number of VIN3 proteins in the yellow line from the
detected number of VIN3-SYFP2 proteins. This factor can be estimated by taking the ratio of
mRNA expression levels of *VIN3* to *SYFP2*, then normalising this by the ratio of *VIN3* to *GFP*
expression in the green line, which lacks endogenous *VIN3*. The error in the correction
factor is propagated directly from the s.e.m. of the independent measurements in panels A-
B. Given the uniform rate of increase of VIN3 expression in all lines regardless of labelling,
the correction factor is expected to be constant over time. Taking an inverse-variance-
weighted average over the three timepoints, the correction factor used in this work is 2.0 ±
0.7, which is consistent with the expected genotype of the yellow line: a single exogenous
gene copy of *VIN3-SYFP2* and a single native gene copy of *VIN3*.

1714

1715

Supplementary Fig. 6. RNA expression analysis of *VRN5* and *FLC* in single-colour VRN5-mScarlet-I lines. Expression level of *VRN5* (A), of spliced *FLC* (B) in VRN5-mScarlet-I, *vrn5-8* and Col*FRI* seedlings before (NV) and after 2 weeks and 6 weeks cold treatments (V2W, V6W). V6T10 represents 10 days after a 6 week-cold treatment. Data are presented as mean ± s.d. (N=3) from 1 biological experiment. The samples of VRN5-mScarlet-I are from heterozygous T2 seedlings.

1722

1723

Supplementary Fig. 7. Confocal microscopy indicates VIN3 and VRN5 expression increase due to vernalisation.

Confocal microscopy projections of live whole root tips of VIN3-GFP and VRN5-YFP at different timepoints. NV is non-vernalised; V2W and V6W refers to 2 or 6 weeks of cold respectively; the post-vernalised timepoint V6W+T refers to 7 days warm (V6W+T7) for VIN3 and 14 days warm (V6W+T14) for VRN5. Bright particles outside the root envelope are remnants of the root cap. Also shown are insets at 13× scale to show detail of the labelled protein signals, which are consistently localised to the nucleoplasm, but not the nucleolus of each cell.

1736

1737

Supplementary Fig. 8. SlimVar detects molecular assemblies with high contrast across at least 3 cell layers in a range of cell morphologies.

SlimVar images of individual nuclei in the VRN5-YFP line, classified qualitatively by cell shape and location in root tip. Three independent fields of view (rows) are shown for each class at a mixture of timepoints before, during and after vernalisation. Left: brightfield, centre: initial fluorescence in SlimVar (average of first 3 frames); right: assemblies revealed by partial photobleaching during SlimVar sequence; visualisation generated in ImageJ from raw sequences by Gaussian smoothing at 1.0 pixel radius, gamma transform of 1.25, background subtraction at 10 pixel radius, followed by maximum projection over frames 30–100 in each sequence, then repeat background subtraction and Gaussian smoothing, then inverse gamma transform of 0.80. Exposure time: 10-20 ms. Scale bars: 10 μm.

1749

Supplementary Fig. 9. Confocal and SlimVar imaging of the FLC-*lacO*/LacI-YFP line. a-d)
Confocal: a) Transmission and b) confocal fluorescence imaging of an non-vernalised
nucleus in the epidermal meristem showing at least 2 loci, and c,d) a vernalised (six weeks of
cold) nucleus in the epidermal elongation zone, featuring additional clustered pairs
of *FLC* foci consistent with endoreduplication. Mobile LacI-YFP form a uniform background
due to motion blur. Scale bars 2 μm. e-h) SlimVar: e) a large nucleus (dotted outline) of a
non-vernalised cortical cell imaged in brightfield with superposed intensity profile of a
focused laser beam (cross) inside the nucleolus; an initial exposure with this beam, typically
30 s, removes excess unbound LacI-YFP that diffuses through the excitation volume, while
preserving any bound to the *lacO* array at the *FLC* transgene; f-h) the subsequent SlimVar
acquisition shown as f) mean average, g) median projection showing isolated and low
mobility *FLC* foci at the expected frequency of ≥2 loci per nucleus, whose diffusivity is
quantified and used to match the diffusivity of VIN3 and VRN5 proteins of interest under the
same acquisition conditions (Supplementary Table 1); and h) standard deviation projection
indicating residual fast-moving LacI-YFP. In spot-bleached nuclei, we find a mean of 4.4 ±
0.8 slow LacI-YFP tracks per nucleus (N=38 nuclei) reflecting the expected two *FLC*
transgenes plus a minority of endoreduplication loci. At the observed particle number
density, the likelihood of random overlap of more than two free LacI-YFP tetramers is
extremely low. We therefore identify all LacI-YFP tracks with stoichiometry greater than
three tetramers, or a threshold of 12 LacI-YFP, as potential *FLC* loci. These tracks exhibit a
diffusivity (Fig. 6a) of D_{FLC} = 0.20 ± 0.02 μm^{2}s^{-1} (N=129, mean ± s.e.m., 20 ms exposure).
Tracks in the upper quartile of LacI-YFP stoichiometry (>81 LacI-YFP molecules),
corresponding to the brightest foci per nucleus, yield an equivalent diffusivity estimate of
0.17 ± 0.04 μm^{2}s^{-1} (N=36; p= 0.097 |ns). With or without pre-bleaching, tracks with fewer
than 12 LacI-YFP molecules typically diffuse faster, around 0.54 ± 0.04 μm^{2}s^{-1} (N=153; p=
0.0094 |*), which suggests they may consist of free LacI tetramers.

1780

1781

1782

1783

1784

1785

Supplementary Fig. 10. Tetrameric spacing of LacI-YFP validates peak interval extraction

from stoichiometry distributions. a) Representative SlimVar images of neighbouring

epidermal nuclei before vernalisation in single-colour *FLC-lacO*/LacI-YFP line under strong

EtOH induction, shown as brightfield, initial SlimVar frame, and projections of mean

(yellow), and merged median (cyan) and standard deviation (magenta) respectively over 5

frames. b) The stoichiometry of detected LacI-YFP tracks (n=225) shows clear peaks at

approximate multiples of 4, with occasional interceding multiples of 2, consistent with the

tightly bound tetrameric (dimer-of-dimer) complex of LacI. The maximum stoichiometry

observed is 380 molecules (not shown for clarity). Inset: the most common spacing

between neighbouring peaks in the stoichiometry distribution as estimated by the new

method. The threshold above which the null distribution can be rejected is the 95[th]

percentile fraction of intervals (grey trace) output from simulated random stoichiometry

(Methods). The most common interval is given by the modal peak value above the null

threshold (3.8 ± 0.3 molecules), which is consistent only with integer 4 molecules, and thus

a tetrameric repeat unit as expected.

1801

1802

Supplementary Fig. 11. Collated SlimVar observations of VIN3-SYFP2. a) Distributions of integrated nuclear intensity of VIN3-SYFP2 in individual nuclei before, during and after vernalisation. NV = not vernalised; V2W = two weeks of cold; V6W = six weeks of cold; V6WT7 = six weeks of cold followed by one week of warm conditions. In the VIN3-SYFP2 line, the unlabelled, endogenous fraction of VIN3 protein can be accounted for by a factor of 2.0 ± 0.7 derived from the relative mRNA expression levels (Supplementary Fig. 5). Bar, box and whisker denote median, interquartile range (IQR) and 1.5× IQR respectively; cross: mean ± s.e.m. Total protein numbers of VIN3-SYFP2 range from negligible before cold (410 ± 560 molecules, BM test vs. Col*FRI*, N=52, p= 0.52 |ns), through 7,800 ± 1,600 molecules above background at 2 weeks cold (N=68, p<0.001|**) up to a peak of 15,000 ± 3,000 molecules at 6 weeks cold (N=51, p=0.0045|*). At 7 days after transfer to warm, the number of VIN3-SYFP2 again drops to a negligible amount (900 ± 700 molecules per nucleus, BM test vs. Col*FRI*, N=54, p=0.08|ns). b) The mean number of VIN3 tracks per nucleus cannot be distinguished between VIN3-SYFP2 and VIN3-GFP, nor does this change significantly between 2 and 6 weeks' vernalisation in either case (V2W: 16.3 ± 1.0 and 18.3 ± 1.9, N=120, p=0.42|ns; V6W: 17.4 ± 2.4 and 17.9 ± 1.8, N=140, p=0.22|ns respectively). c) After accounting for unlabelled VIN3, the mean labelled stoichiometry increases from 7.8 ± 2.7 molecules at 2 weeks' cold to 13.8 ± 3.9 at 6 weeks' cold (N=1792 tracks, p<0.001). d) The stoichiometry distribution for VIN3-SYFP2 has a smaller periodicity (dashed lines), which is incompatible with an integer number of molecules greater than one. However, after accounting for unlabelled VIN3, the resulting periodicity (2.2 ± 0.8, V2W; 2.2 ± 0.8, V6W, solid lines) is also consistent with dimeric subunits in VIN3 assemblies during vernalisation.

Supplementary Fig. 12. Localisation precision of tracked fluorescent foci in SlimVar acquisitions. Gaussian-fit widefield FWHM (major axis) and the corresponding super-resolved localisation precision estimated using the formulation of Thompson et al. [93] for foci within a) VRN5-YFP tracks (n= 5326), b) VIN3-GFP tracks (n=2479) at 2 ms exposure per frame; c) VRN5-mScI tracks (n=6395) and d) *FLC-lacO*/LacI-YFP loci (n=475) at 20 ms exposure. The localisation precision becomes finer with greater photon counts in each focus. However, most localisations occur during later stages of photobleaching; these are relatively dim compared to background counts, and therefore do not reach the theoretical limit scaling as the inverse square root of the photon count. For the brightest foci, we observe values smaller than the lateral spatial localisation precision for a single molecule (40 nm) but these should be interpreted with caution since the physical size of each assembly is unknown.

Supplementary Video legends

Supplementary Video 1. SlimVar resolves dynamic VRN5-YFP assemblies in live root tips.
A single nuclear acquisition of VRN5-YFP line in an epidermal cell (10 μm deep) at non-vernalised (NV) conditions showing distinct assemblies. Left: raw data; right: photobleach corrected and Gaussian smoothed by 1 pixel width to emphasise dynamic assemblies. Scale bar: 1 μm, frame duration 2 ms, frame rate 25 fps.

Supplementary Video 2. VRN5-YFP assemblies in live root tips after vernalisation.

A SlimVar acquisition in a large cortical nucleus (25 µm deep) in the VRN5-YFP line at post-

vernalised (V6WT14) conditions showing large assemblies. Left: raw data; right:

photobleach corrected and Gaussian smoothed by 1 pixel width. Scale bar: 2 µm, frame

duration 20 ms, frame rate 10 fps.

Supplementary Video 3. VIN3-GFP assemblies during vernalisation.

A representative SlimVar acquisition of a smaller epidermal nucleus in VIN3-GFP line at 2

weeks' cold. VIN3 has significantly lower copy number and stoichiometry compared to

VRN5, and background is higher in GFP channel relative to YFP. Scale bar 1 µm, frame

duration 2 ms, frame rate 25 fps, as for Video 1.