Transcriptomic analysis identifies injury-responsive fibroblast populations as potential mediators of Wnt-dependent spinal cord regeneration

Samuel R. Alper[†], Deeptha Vasudevan[†], Maya K. Wheeler, Samin Panahi, and Richard I. Dorsky*

Department of Neurobiology, University of Utah, Salt Lake City, UT 84112, USA.

† These authors contributed equally to this work.

*Author for Correspondence. Email: richard.dorsky@neuro.utah.edu

Keywords: Spinal cord injury, Wnt signaling, Regeneration, Fibroblasts, Meninges, Zebrafish

ABSTRACT

Background: In humans and other mammals, spinal cord injury (SCI) can lead to a permanent loss of sensory and motor function, due to the inability of damaged neurons and axons to regenerate. However, other vertebrate species including zebrafish exhibit complete spinal cord regeneration and functional recovery after SCI. Wnt signaling is required for neurogenesis and axon regrowth in a larval zebrafish SCI model, but the genes regulated by this pathway and the cell types that express them remain largely unknown. Results: In this study, we used bulk RNA-sequencing (RNAseq) to identify candidate genes regulated by Wnt signaling that are expressed after SCI. Using this unbiased screen, we identified multiple genes previously unassociated with SCI in larval zebrafish, and confirmed by in situ hybridization that their expression is both injury-responsive and Wnt-dependent. Conclusions: Together, our data suggest novel gene targets and cell types that may play important roles in spinal cord regeneration.

INTRODUCTION

Treatment options for human spinal cord injury (SCI) are limited and no current therapies promote full functional recovery. Permanent sensorimotor deficits following mammalian SCI are due to both a lack of neurogenesis and poor axon regrowth at the injury site. The absence of neural progenitor cells in the adult mammalian spinal cord explains the inability to replace lost neurons [1]. Poor axon regrowth results from intrinsic deficits in neuronal regenerative capacity [2], as well as a complex web of extrinsic factors. Data from mammalian models of SCI demonstrate significant inhibitory actions of non-neural, extraspinal cells . Several cell types including microglia, astrocytes, immune cells, fibroblasts, and meningeal cells infiltrate the injury site, induce inflammation, and secrete an array of growth-inhibiting extracellular matrix (ECM) components [3]. The result is the formation of a fibrotic scar that poses both a molecular and physical barrier for regrowing axons.

An increasing interest in the mechanisms that promote functional recovery in spontaneously regenerating species has led to the development of zebrafish SCI models. Zebrafish show regenerative neurogenesis, axon regrowth, and rewiring of spinal circuits following injury, leading to functional locomotor recovery [4-7]. The larval zebrafish SCI model is particularly advantageous due to its high throughput, rapid regeneration time, optical accessibility, and general conservation of regenerative mechanisms with adult fish [8]. Using this model, we aimed to better understand the mechanisms of spinal cord regeneration.

A wide array of cell types and signaling cascades are activated following zebrafish SCI. Of particular importance is the injury-induced reactivation of signaling pathways that play prominent

roles during embryonic development, including the Wnt/β-catenin pathway [9]. After SCI in larvae, Wnt signaling activity begins within 1 day post-injury (dpi), peaks by 2 dpi, and dissipates by around 3 dpi [10]. Prior data from our lab and others shows that Wnt signaling is required for both regenerative neurogenesis and axon regrowth [6,10-12]. After SCI at both larval and adult stages, resident neural progenitors proliferate and differentiate into newborn neurons. However, only a small fraction of spinal cord progenitors and no surviving neurons appear to respond directly to Wnt signaling after SCI [6,10].

Instead, most Wnt-responsive cells after injury are extraspinal and express general fibroblast markers, but more specific identification and characterization of these cells has remained elusive [10]. Wnt pathway activity in fibroblast-like cells after SCI is required for the expression of collagens and other ECM components. However, so far only two genes encoding Collagen XII orthologs have been shown to be induced by SCI, Wnt-dependent, and required for axon regrowth [10]. While valuable, these studies have led to an incomplete understanding of the Wnt target genes that mediate spinal cord regeneration and the identity of extraspinal Wnt-responsive cell populations that express them.

To address these questions, we performed bulk RNA sequencing (RNAseq) analyses on injured zebrafish larvae treated with pharmacological inhibitor of Wnt signaling. By comparing mRNA expression from these larvae with untreated controls, we have identified SCI- and Wnt-dependent genes that we have been able to functionally categorize based on gene ontology (GO) enrichment analysis. Of particular interest are genes encoding feedback inhibitors of Wnt signaling, and markers of injury-associated infiltrating and meningeal fibroblasts. We have validated the SCI- and Wnt-dependent expression patterns of transcripts from each of these categories in the

regenerating larval spinal cord by in situ hybridization. Together, our data identify novel Wnt target genes and cell populations that may play important roles in spinal cord regeneration.

EXPERIMENTAL PROCEDURES:

Zebrafish

The following zebrafish strains were used for this study: *AB, *mitfa*[w2] [13], and *Tg(7xTCF-Xla.Siam:GFP)[ia4]* [14]. Larvae used for experiments were obtained from incrosses and staged according to [15]. All experiments were approved by the University of Utah Institutional Animal Care and Use Committee.

Bulk RNA sequencing

*AB and *Tg(7xTCF-Xla.Siam:GFP)[ia4]* larvae were raised in E3 media until 5 dpf (days post fertilization), anesthetized in 0.016% Tricaine (Sigma), and injured by complete transection of the spinal cord at the level of the anal pore using a sharpened and sterilized FST injury tool. Larvae were allowed to recover for 6 hours in E3 before being transferred into media containing E3 with 35µM IWR-1 (Sigma) in 0.25% DMSO or 0.25% DMSO as a vehicle control. Larvae were fed rotifers starting at 1 dpi until fixation at 3 dpi (8 dpf). To confirm Wnt pathway inhibition, *Tg(7xTCF-Xla.Siam:GFP)[ia4]* larvae were imaged with a Nikon A1 HD25 confocal microscope using a 60X oil immersion lens and images were processed in Adobe Photoshop.

Larvae were fixed in 4% paraformaldehyde (PFA) and 10% sucrose in PBS at 3 dpi at 4°C overnight. For each biological replicate of 20 animals, spinal cord and adjacent tissues dorsal to the notochord were dissected 1-2 segments caudal and rostral to the injury site. This process was

repeated for three biological replicates per treatment condition. The Invitrogen Recover All Total Nucleic Acid Isolation Kit for FFPE was used to extract RNA from the tissue samples.

Purified RNA samples were analyzed by the High Throughput Genomics Core Facility at the Huntsman Cancer Institute using the NEBNext Ultra II Directional RNA Library Prep Kit with rRNA depletion. cDNA libraries were then sequenced using a NovaSeq 6000 instrument at 50X50 bp read length and 100M read-pair increments. Cutadapt was used for adapter trimming, FastQC for quality control, STAR for alignment and featureCounts for quantification with genome and annotation downloaded from Ensembl version Danio_rerio.GRCz11.94, and DESeq2 for differential expression analysis. Gene Ontology (GO) enrichment analysis for biological process and cellular component was performed for all genes with a negative log2 fold change and adjusted p values <0.05.

In situ hybridization and immunohistochemistry

mitfa^{w2} homozygous mutant larvae were raised in E3 media until 3 dpf, anesthetized in 0.016% Tricaine (Sigma), and injured by complete transection of the spinal cord at the level of the anal pore using a sterile 30G needle [16]. Injured larvae were allowed to recover in E3 for 6 hours before being transferred to media containing either 15uM IWR-1 in 0.25% DMSO or 0.25% DMSO as a vehicle control.

In situ hybridization and immunohistochemistry

mtfa^{w2} homozygous mutant larvae were raised in E3 media until 3 dpf, anesthetized in 0.016% Tricaine (Sigma), and injured by complete transection of the spinal cord at the level of the anal pore using a sterile 30G needle [16]. Injured larvae were allowed to recover in E3 for 6 hours before

being transferred to media containing either 15uM IWR-1 in 0.25% DMSO or 0.25% DMSO as a vehicle control [10]. In situ hybridization probes were made by a clone-free method as described previously[17]with DNA templates purified using Zymo Research DNA clean & concentrator™-5 kit. Primers were designed by Primer-BLAST (NCBI).

Injured fish were fixed at 2 dpi in 4% PFA containing 0.1% Tween 20 for 1 hour at room temperature and subsequently dehydrated in 100% Methanol and stored at -20°C. Whole mount *in situ* hybridization and IHC were performed according to [16] with a 10 minute pre-incubation in 2%H_2O_2 in MeOH. ISH-stained fish were imaged on an Olympus compound microscope using a 20X objective. Chicken anti-GFP (Aves #GFP-2010) and Goat anti-Chicken Alexa-488 (Invitrogen #A-11039) antibodies were used to label GFP+ cells in *Tg(7xTCF-Xla.Siam:GFP)[ia4]* larvae. 30um Z-stacks of antibody-labeled larvae were obtained using a Zeiss700 confocal microscope with a 20x objective. Linear adjustments to brightness and contrast were made using FIJI.

RESULTS AND DISCUSSION

<u>RNAseq analysis reveals known and novel Wnt-dependent genes following SCI</u>

Previous work has demonstrated that Wnt signaling is integral to both axon regrowth and neurogenesis after injury [6,10]. However, the transcriptional targets of Wnt signaling that mediate these regenerative processes remain incompletely characterized. To generate a list of candidate target genes, we performed bulk RNAseq on regenerating spinal cord and adjacent tissues dissected from larvae treated with the Wnt pathway inhibitor IWR-1 [18], and from vehicle treated controls (Fig. 1A-B). Tissue was collected at 3 dpi following spinal cord transections performed on 5 dpf larvae (Fig. 1A). Previous work from our lab found that at this timepoint axon regrowth

is observed at the injury site, but regenerated neurons are not yet present [6,7]. After DESeq2 analysis of IWR-1 treated vs. untreated samples, we were able to identify 1,791 differentially expressed annotated genes. Because we are primarily interested in genes with increased expression after SCI, and because most known direct Wnt/ß-catenin targets are transcriptionally activated by pathway activity [19] we focused specifically on the 872 genes with expression that was significantly decreased by Wnt inhibition (adjP<0.05). When we performed a gene ontology (GO) analysis of this list using the PANTHER overrepresentation test for biological process, we found high enrichment of terms such as "negative regulators of Wnt signaling pathway" (13.5-fold enrichment, adjP=3.24E-04), all of which included multiple genes that are known direct feedback targets of Wnt signaling. This result confirmed the effectiveness of our screen in detecting Wnt-dependent gene expression. We also found significant enrichment of GO terms related to tissue development and regeneration, suggesting that some novel Wnt targets identified by our screen could also be involved in the injury response. Finally, when we performed a GO analysis for cellular component, we found high enrichment of "collagen-containing extracellular matrix" (5.02-fold enrichment, adjP=4.57E-03), which was consistent with the requirement for Wnt-dependent collagen expression by infiltrating fibroblasts in axon regrowth after SCI [10] (Fig. 3A). Together, these data suggested that our RNAseq analysis could help identify novel Wnt-dependent molecular and cellular mediators of spinal cord regeneration.

Validation of candidate Wnt target gene expression in a 3 dpf injury model

While our RNAseq analysis was performed after SCI at 5 dpf, several recent studies have standardized an earlier 3 dpf SCI model (Fig 1A, E-F) [10,20,21]. At that timepoint the zebrafish larva is more accessible for whole-mount imaging and histological analysis, and the regenerative

response to injury largely recapitulates the responses in both 5 dpf larval and adult spinal cord injury paradigms [8](Alper & Dorsky 2022). Thus, we decided to adopt 3 dpf as our new injury paradigm. In order to confirm that Wnt signaling is both injury-responsive and IWR-1-dependent following SCI at this stage, we examined *Tg(7xTCF-Xla.Siam:GFP)[ia4]* larvae (henceforce TCF:GFP), which express a GFP reporter of Wnt/ß-catenin pathway activity [14]. By 2 dpi (5 dpf), we observed numerous GFP+ cells near the injury site in control larvae, while IWR-1 treated larvae showed a clear reduction in GFP expression (Fig. 1C-D). From these results we concluded that we could use this new paradigm to validate the injury-responsive and Wnt-dependent expression of candidate genes from our RNAseq analysis.

Injury-dependent expression of Wnt signaling feedback inhibitors

Our RNAseq analysis identified several known Wnt target genes encoding secreted inhibitors of pathway activity, including orthologs of *notum1*, *sfrp1*, *wif1*, *and dkk1* (Fig. 2A). We performed in situ hybridization for *notum1a* mRNA to determine whether this gene is expressed at the injury site following SCI. At 2 dpi, *notum1a* expression was present in close proximity to the injury site in vehicle treated controls, but was clearly diminished in IWR-1 treated larvae (Fig. 2B-C). This experiment confirms the injury and Wnt dependence of a candidate identified by our screen, the efficacy of our drug treatment, and suggests that *notum1a* may play a currently unexplored role in Wnt-dependent spinal cord regeneration. A requirement for the injury-induced activation of Wnt signaling in zebrafish spinal cord regeneration is well documented [6,10,12]. However, other roles of Wnt signaling also require the subsequent down-regulation of pathway activity in order for complete function to be achieved [22,23]. Studies have also shown that transient cell-type specific signals, including the Wnt pathway, allow cellular populations to execute pro-regenerative

functions [20,21,24-26]. Consistent with this possibility, the secreted inhibitors that we identified may act to restrict Wnt-signaling from cells near the injury site. Future studies will be required to determine the precise dynamics of Wnt signaling after SCI.

Wnt-dependent expression of infiltrating fibroblast markers

After SCI in larval zebrafish, most Wnt-responsive cells accumulate dorsal to the injury site [10]. These cells have been classified as "fibroblast-like" based on tissue localization, collagen expression, and occasional expression of the dermal fibroblast marker p63, but a more specific classification remains unknown [10]. Furthermore, CollagenXII expression by these cells is the only known Wnt-dependent gene shown to be required for axon regrowth and functional recovery [10]. Our RNAseq data show significant downregulation of several transcripts known to be expressed by injury-responsive infiltrating fibroblasts including orthologs of *fstl1*, *cthrc1*, and *ptx3* (Fig. 3A) [21,27,28]. These data also show that Wnt inhibition leads to significant downregulation of two paralogous genes encoding the homeodomain transcription factor Prrx1, which has previously been implicated in fibroblast-mediated tissue regeneration (Fig. 3A) (Currie et al., 2019; de Bakker et al., 2021; Leavitt et al., 2020; Yeo et al., 2018). In situ hybridization confirmed that IWR-1 treatment decreased the injury-induced expression of both paralogs, *prrx1a* and *prrx1b* (Fig. 3B-E). A role for Prrx1 in central nervous system repair has not been described previously, and the Wnt-dependent expression of *prrx1a/b* after SCI could indicate a function for the pathway in the switch of infiltrating fibroblasts from a fibrotic to a pro-regenerative state, by regulating the expression of signals that promote neurogenesis and axon regrowth.

Wnt-dependent expression of meningeal fibroblast markers

Meningeal cells are hypothesized to inhibit axon regrowth after mammalian SCI by secreting ECM components that contribute to chronic scarring and inflammation [29]. In axolotls, which show spontaneous spinal cord regeneration, putative meningeal fibroblasts aggregate on the leading edge of regrowing axons, consistent with a growth-promoting function [30]. However, a meningeal role in zebrafish spinal cord regeneration has not yet been demonstrated. A recent paper performed scRNAseq after injury to the mouse spinal cord and used two well-known meningeal cell markers, *decorin* and *col1a1*, to identify additional genes expressed in injury- responsive meningeal cells (Matson et al., 2022). We found that nine of these 20 meningeal genes had significant differential expression in our RNAseq data. In addition, six of these nine Wnt-dependent genes were also enriched in a scRNAseq dataset of embryonic mouse dural and arachnoid fibroblasts (DeSisto et al., 2020). Furthermore, Daniocell, an scRNAseq database of 0-5dpf zebrafish, shows enriched expression of these genes in meningeal fibroblasts and meningeal precursor cells: *fmoda*, *bambia*, *ogna*, *col1a1a*, *col1a1b*, and *aldh1a2* (Fig. 4A) [31]. These observations are consistent with a previous report of Wnt-activity in cells immediately adjacent to the spinal cord after zebrafish SCI [10] and thus suggest that Wnt-responsive meningeal cells may play a role in zebrafish spinal cord regeneration. While the cerebral meninges have begun to be characterized in zebrafish [32], the molecular and cellular composition of the spinal cord meninges and its roles in development, homeostasis, and regeneration remain poorly understood. We therefore sought to validate two of our RNAseq-identified candidate meningeal cell markers, *col1a1a* and *fmoda*. Expression was upregulated near the injury site at 2 dpi in controls, while treatment with IWR-1 after SCI decreased the expression of both transcripts (Fig. 4B-E). This suggests possible Wnt-dependent pro-regenerative roles of meningeal cells after spinal cord injury in zebrafish. However, the extent to which these cells contribute to regeneration requires further investigation.

CONCLUSIONS

In summary, our identification and validation of Wnt-dependent gene expression after SCI suggests that the subsequent inhibition of Wnt signaling may play a role in spinal cord regeneration, and identifies two novel fibroblast populations that could participate in the repair process. Future studies will characterize the functional relevance of these genes and cell types in axon regrowth, neurogenesis, and the recovery of sensorimotor function.

CONTRIBUTIONS
S.A.R. (Investigation, analysis, writing)
D.V. (Investigation, analysis, writing)
M.K.W. (Investigation, analysis, writing)
S.P. (Investigation)
R.I.D. (Funding acquisition, analysis, writing)

Figure 1. Workflow for bulk RNA sequencing and validation of Wnt inhibition by IWR-1.

(A) Schematic of SCI and IWR-1 treatment. (B) Schematic of RNAseq workflow. (C-D) SCI was performed on TCF:GFP fish at 3dpf and treated with IWR-1 or vehicle control (DMSO) until fixation at 2dpi. Wnt-responsive cells (GFP+) are present at the injury site both within and dorsal to the spinal cord in vehicle-treated controls. A decrease in Wnt-responsive cells in IWR-1 treated animals indicates successful pharmacological inhibition of Wnt signaling. (E-F) SCI was performed on 5dpf WT larvae and treated continuously with IWR-1 until sacrifice at 3dpi. Wnt-responsive (GFP+) cells are present at the injury site in DMSO controls. GFP expression is reduced in IWR-1 treated fish. At both timepoints, analysis of GFP expression in the transgenic Wnt reporter line shows successful IWR-1 mediated inhibition of Wnt signaling. Asterisk: Injury site.

Figure 2. Wnt target genes are expressed in response to SCI.

(A) Gene Ontology (GO) analysis of the 872 significantly downregulated genes from our RNAseq dataset shows significant enrichment of "Wnt protein binding". IWR-1 treatment after SCI leads to significant downregulation of known Wnt target genes encoding secreted feedback inhibitors. (B) *in situ* hybridization shows that *notum1a* is expressed in close proximity to the injury site at 2 dpi in control larvae. (C) Injury-induced expression of *notum1a* is reduced by IWR-1 treatment. Asterisk: injury site

Figure 3. Infiltrating fibroblast markers are SCI- and Wnt-dependent.

(A) GO analysis of the 872 significantly downregulated genes from our RNAseq dataset shows significant enrichment in "basement membrane" and "collagen-containing ECM". IWR-1 treatment after SCI leads to significant downregulation of transcripts expressed in infiltrating fibroblasts. (B,D) *in situ* hybridization shows that *prrx1a* and *prrx1b* are expressed in close proximity to the injury site at 2 dpi in vehicle treated controls. (C,E) IWR-1 treatment reduces *prrx1a/b* expression after SCI. Asterisks: Injury site.

Figure 4. Markers for meningeal fibroblasts are SCI- and Wnt-dependent.

(A) IWR-1 treatment after SCI leads to significant reduction of markers known to be expressed in meningeal fibroblasts. (B,D) Two of these markers, *col1a1a* and *fmoda*, are expressed near the injury site in control larvae at 2dpi. (C,E) IWR-1 treatment reduces the SCI-induced expression of *col1a1a* and *fmoda* at 2dpi. Asterisks: Injury site.

REFERENCES

1.   Tai W, Wu W, Wang LL, et al. In vivo reprogramming of NG2 glia enables adult neurogenesis and functional recovery following spinal cord injury. *Cell Stem Cell*. May 6 2021;28(5):923-937 e4. https://doi.org10.1016/j.stem.2021.02.009.

2.   Bray ER, Cheret J, Yosipovitch G, Paus R. Schwann cells as underestimated, major players in human skin physiology and pathology. *Exp Dermatol*. Jan 2020;29(1):93-101. https://doi.org10.1111/exd.14060.

3.   Shafqat A, Albalkhi I, Magableh HM, Saleh T, Alkattan K, Yaqinuddin A. Tackling the glial scar in spinal cord regeneration: new discoveries and future directions. *Front Cell Neurosci*. 2023;17:1180825. https://doi.org10.3389/fncel.2023.1180825.

4.   Becker T, Wullimann MF, Becker CG, Bernhardt RR, Schachner M. Axonal regrowth after spinal cord transection in adult zebrafish. *J Comp Neurol*. Jan 27 1997;377(4):577-95.

5.   Briona LK, Dorsky RI. Radial glial progenitors repair the zebrafish spinal cord following transection. *Exp Neurol*. Jun 2014;256:81-92. https://doi.org10.1016/j.expneurol.2014.03.017.

6.   Briona LK, Poulain FE, Mosimann C, Dorsky RI. Wnt/ß-catenin signaling is required for radial glial neurogenesis following spinal cord injury. *Dev Biol*. Apr 14 2015. https://doi.org10.1016/j.ydbio.2015.03.025.

7.   Vasudevan D, Liu YC, Barrios JP, Wheeler MK, Douglass AD, Dorsky RI. Regenerated interneurons integrate into locomotor circuitry following spinal cord injury. *Exp Neurol*. Aug 2021;342:113737. https://doi.org10.1016/j.expneurol.2021.113737.

8.   Alper SR, Dorsky RI. Unique advantages of zebrafish larvae as a model for spinal cord regeneration. *Front Mol Neurosci*. 2022;15:983336. https://doi.org10.3389/fnmol.2022.983336.

9.   Cardozo MJ, Mysiak KS, Becker T, Becker CG. Reduce, reuse, recycle - Developmental signals in spinal cord regeneration. *Dev Biol*. Dec 1 2017;432(1):53-62. https://doi.org10.1016/j.ydbio.2017.05.011.

10.   Wehner D, Tsarouchas TM, Michael A, et al. Wnt signaling controls pro-regenerative Collagen XII in functional spinal cord regeneration in zebrafish. *Nature communications*. Jul 25 2017;8(1):126. https://doi.org10.1038/s41467-017-00143-0.

11.   Becker CG, Becker T, Hugnot JP. The spinal ependymal zone as a source of endogenous repair cells across vertebrates. *Prog Neurobiol*. Nov 2018;170:67-80. https://doi.org10.1016/j.pneurobio.2018.04.002.

12.   Strand NS, Hoi KK, Phan TMT, Ray CA, Berndt JD, Moon RT. Wnt/beta-catenin signaling promotes regeneration after adult zebrafish spinal cord injury. *Biochem Biophys Res Commun*. Sep 2 2016;477(4):952-956. https://doi.org10.1016/j.bbrc.2016.07.006.

13.   Lister JA, Robertson CP, Lepage T, Johnson SL, Raible DW. nacre encodes a zebrafish microphthalmia-related protein that regulates neural-crest-derived pigment cell fate. *Development*. 1999;126(17):3757-3767.

14.   Moro E, Ozhan-Kizil G, Mongera A, et al. In vivo Wnt signaling tracing through a transgenic biosensor fish reveals novel activity domains. *Developmental biology*. Apr 21 2012. https://doi.org10.1016/j.ydbio.2012.03.023.

15.   Kimmel CB, Ballard WW, Kimmel SR, Ullmann B, Schilling TF. Stages of embryonic development of the zebrafish. *Dev Dyn*. Jul 1995;203(3):253-310.

16.   John N, Kolb J, Wehner D. Mechanical spinal cord transection in larval zebrafish and subsequent whole-mount histological processing. *STAR Protoc*. Mar 18 2022;3(1):101093. https://doi.org10.1016/j.xpro.2021.101093.

17.   Xie Y, Kaufmann D, Moulton MJ, et al. Lef1-dependent hypothalamic neurogenesis inhibits anxiety. *PLoS Biol*. Aug 2017;15(8):e2002257. https://doi.org10.1371/journal.pbio.2002257.

18.   Lu J, Ma Z, Hsieh JC, et al. Structure-activity relationship studies of small-molecule inhibitors of Wnt response [Research Support, N.I.H., Extramural

Research Support, Non-U.S. Gov't]. *Bioorg Med Chem Lett*. Jul 15 2009;19(14):3825-7. https://doi.org10.1016/j.bmcl.2009.04.040.

19.   Schuijers J, Mokry M, Hatzis P, Cuppen E, Clevers H. Wnt-induced transcriptional activation is exclusively mediated by TCF/LEF. *Embo J*. Jan 13 2014;33(2):146-56. https://doi.org10.1002/embj.201385358.

20.   Cavone L, McCann T, Drake LK, et al. A unique macrophage subpopulation signals directly to progenitor cells to promote regenerative neurogenesis in the zebrafish spinal cord. *Dev Cell*. Jun 7 2021;56(11):1617-1630 e6. https://doi.org10.1016/j.devcel.2021.04.031.

21.   Tsata V, Mollmert S, Schweitzer C, et al. A switch in pdgfrb(+) cell-derived ECM composition prevents inhibitory scarring and promotes axon regeneration in the zebrafish spinal cord. *Dev Cell*. Feb 22 2021;56(4):509-524 e9. https://doi.org10.1016/j.devcel.2020.12.009.

22.   Agathocleous M, Iordanova I, Willardsen MI, et al. A directional Wnt/beta-catenin-Sox2-proneural pathway regulates the transition from proliferation to differentiation in the Xenopus retina [Research Support, N.I.H., Extramural Research Support, Non-U.S. Gov't]. *Development*. Oct 2009;136(19):3289-99. https://doi.org10.1242/dev.040451.

23.   Wang X, Kopinke D, Lin J, et al. Wnt signaling regulates postembryonic hypothalamic progenitor differentiation [Research Support, N.I.H., Extramural]. *Developmental cell*. Sep 11 2012;23(3):624-36. https://doi.org10.1016/j.devcel.2012.07.012.

24.   Klatt Shaw D, Mokalled MH. Efficient CRISPR/Cas9 mutagenesis for neurobehavioral screening in adult zebrafish. *G3 (Bethesda)*. Mar 20 2021. https://doi.org10.1093/g3journal/jkab089.

25.   Wehner D, Cizelsky W, Vasudevaro MD, et al. Wnt/beta-catenin signaling defines organizing centers that orchestrate growth and differentiation of the regenerating zebrafish caudal fin. *Cell reports*. Feb 13 2014;6(3):467-81. https://doi.org10.1016/j.celrep.2013.12.036.

26. Gay D, Ghinatti G, Guerrero-Juarez CF, et al. Phagocytosis of Wnt inhibitor SFRP4 by late wound macrophages drives chronic Wnt activity for fibrotic skin healing. *Sci Adv*. Mar 2020;6(12):eaay3704. https://doi.org10.1126/sciadv.aay3704.

27. Maruyama T, Jeong J, Sheu TJ, Hsu W. Stem cells of the suture mesenchyme in craniofacial bone development, repair and regeneration. *Nature communications*. Feb 1 2016;7:10526. https://doi.org10.1038/ncomms10526.

28. Sun J, Peterson EA, Chen X, Wang J. ptx3a(+) fibroblast/epicardial cells provide a transient macrophage niche to promote heart regeneration. *Cell reports*. Apr 23 2024;43(4):114092. https://doi.org10.1016/j.celrep.2024.114092.

29. Dorrier CE, Jones HE, Pintaric L, Siegenthaler JA, Daneman R. Emerging roles for CNS fibroblasts in health, injury and disease. *Nat Rev Neurosci*. Jan 2022;23(1):23-34. https://doi.org10.1038/s41583-021-00525-w.

30. Zukor KA, Kent DT, Odelberg SJ. Meningeal cells and glia establish a permissive environment for axon regeneration after spinal cord injury in newts. *Neural Dev*. Jan 4 2011;6:1. https://doi.org10.1186/1749-8104-6-1.

31. Farrell JA, Wang Y, Riesenfeld SJ, Shekhar K, Regev A, Schier AF. Single-cell reconstruction of developmental trajectories during zebrafish embryogenesis. *Science*. Jun 1 2018;360(6392). https://doi.org10.1126/science.aar3131.

32. Venero Galanternik M, Castranova D, Gore AV, et al. A novel perivascular cell population in the zebrafish brain. *eLife*. Apr 11 2017;6. https://doi.org10.7554/eLife.24369.

A

Spinal cord transection

↓

Recovery in DMSO or IWR-1

↓

Tissue dissection

B

RNA extraction

↓

Library preparation

↓

Sequencing

↓

Identify up and down-regulated Wnt- dependent genes in SCI

C 2dpi, DMSO

GFP
Hoechst
20 µm

D 2dpi, IWR-1

E 3dpi, DMSO

GFP

F 3dpi, IWR-1

A

GO Molecular Function	Fold Enrichment	adjP
Wnt protein binding	8.57	9.54E-03

Gene	log2FoldChange	adjP
notum1a	-2.28	2.05E-19
notum1b	-2.18	2.85E-11
sfrp1a	-1.00	1.55E-06
wif1	-1.09	4.12E-05
dkk1b	-1.76	7.78E-04

A

GO Cellular Component	Fold Enrichment	adjP
Collagen-containing ECM	5.46	2.86E-11
Basement membrane	5.89	5.88E-04

Gene	log2FoldChange	adjP
fstl1a	-1.17	7.50E-07
prrx1a	-1.10	2.10E-07
prrx1b	-1.12	4.60E-06
cthrc1a	-0.73	3.60E-04
ptx3a	-0.72	5.60E-03

A

Gene	log2FoldChange	adjP
fmoda	-1.23	6.49E-14
bambia	-1.56	3.39E-09
ogna	-0.87	7.03E-07
col1a1a	-0.88	8.65E-06
col1a1b	-0.68	1.62E-04
aldh1a2	-0.67	2.67E-03