Accurate Inference of the Polyploid Continuum using Forward-time Simulations

Tamsen Dunn[1,2], Arun Sethuraman[1*]

[1] Department of Biology, San Diego State University

[2] Joint Doctoral Program in Evolutionary Biology, San Diego State University and
Department of Evolution, Ecology, and Organismal Biology, University of California
Riverside

[*]Corresponding author

Email: asethuraman@sdsu.edu

Abstract

Multiple rounds of whole-genome duplication (WGD) followed by diploidization have
occurred throughout the evolutionary history of angiosperms. To understand how these
cycles occur, much work has been done to model the genomic consequences and
evolutionary significance of WGD. The machinations of diploidization are strongly
influenced by the mode of speciation (allo or autopolyploidy). However, there is no
discrete boundary between allo and autopolyploidy, which is best described as a
continuum. Here we present a forward-time polyploid genome evolution simulator called
SpecKS. SpecKS models polyploid speciation as originating from a 2D continuum,
whose dimensions account for both the level of genetic differentiation between the
ancestral parental genomes, as well the time lag between ancestral speciation and their
subsequent reunion in the derived polyploid. Using extensive simulations, we
demonstrate that changes in initial conditions along either dimension of the 2D
continuum deterministically affect the shape of the *Ks* histogram. Our findings indicate
that the error in the common method of estimating WGD time from the *Ks* histogram
peak scales with the degree of allopolyploidy, and we present an alternative, accurate
estimation method that is independent of the degree of allopolyploidy. Lastly, we use
SpecKS to derive tests that infer both the lag time between parental divergence and
WGD time, and the diversity of the ancestral species, from an input *Ks* histogram. We
apply the latter test to transcriptomic data from over 200 species across the plant
kingdom, the results of which are concordant with the prevailing theory that the majority
of angiosperm lineages are derived from diverse parental genomes and may be of
allopolyploid origin.

Introduction

Multiple rounds of Whole-Genome Duplication (**WGD**) followed by diploidization
have occurred throughout the evolutionary history of angiosperms (Otto and Whitton
2000; Soltis and Soltis 2012; Wendel 2015). WGD is considered a major speciation
mechanism (Doyle and Egan 2010; Schranz et al. 2012; Wendel et al. 2016; Clark and
Donoghue 2018), presenting a massive "macromutation", potentially interfering with
sexual reproduction, releasing transposons, and unbalancing molecular signaling
pathways (McClintock 1929; Stebbins 1951; Mayer and Aguilera 1990; Comai et al.
2000; Ramsey and Schemske 2002; Le Comber et al. 2010; Arrigo and Barker 2012;
Yant and Bomblies 2015; Bomblies et al. 2016; Zhang et al. 2016; Baduel et al. 2018). It
has been theorized that the advantage of WGD lies not in the multiplicity of genomic
material itself, but the intense period of genomic reorganization and gene shedding that
follows, known as diploidization (Buggs et al. 2011; Madlung 2013; Soltis et al. 2014;
Tank et al. 2015; Dodsworth et al. 2016; Soltis et al. 2016; Robertson et al. 2017;
Baniaga et al. 2020; Carretero-Paulet and Van de Peer 2020; Nieto Feliner et al. 2020;
Van de Peer et al. 2020; Li et al. 2021).
The diploidization process depends on whether an ancient WGD event is due to
allopolyploidy (the multiplicity of genomes originating from two different species) or due
to autopolyploidy (the multiplicity of genomes originating from within the same species).
For example, allopolyploids are only capable of recombination across chromosomes
from the same parental lineage, while autopolyploids are capable of recombination
across all chromosomes (Blischak et al. 2023; Deb et al. 2023). Furthermore, the
existence of intermediate types of polyploids suggests that polyploidy is best modeled
as a continuum (Stebbins 1950; Meirmans and Van Tienderen 2013; Doyle and
Sherman-Broyles 2017; Mason and Wendel 2020; Blischak et al. 2023).
The estimation of the relative proportion of speciation by allopolyploidy or
autopolyploidy across the plant tree of life has been much debated (Stebbins 1947;
Soltis et al. 2007; Soltis et al. 2010; Barker et al. 2016). Recent assessments suggest
that the majority of sequenced plant genomes are derived from allopolyploidy
(Darlington 1937; Clausen et al. 1945; Parisod et al. 2010; Soltis et al. 2010; Barker et
al. 2016; Wang et al. 2019; Baniaga et al. 2020). Another critical parameter for
assessing the evolutionary significance of WGD is the timing of the WGD event. This is
critical for correlating WGD events with ecological and geological factors (Barba-
Montoya et al. 2018; L. Zhang et al. 2020; Clark and Donoghue 2023), as well as
understanding the timescale of the diploidization process (Hoegg et al. 2004; Doyle and
Egan 2010; Mable 2013; De La Torre et al. 2017; Laurent et al. 2017; David et al. 2020;
Li et al. 2021).
There are many methods to date WGD events. One method is to use genetic
data to place the WGD event on a phylogenetic tree, if enough is known about its
presence from multiple lineages (Bowers et al. 2003; Li et al. 2015; Li et al. 2018; Li and
Barker 2020; Parey et al. 2022), or its presumptive parental species (Lott et al. 2009;
Doyle and Egan 2010; Estep et al. 2014; Douglas et al. 2015; Thomas et al. 2017;
Mccann et al. 2018; Wen et al. 2018; Yan et al. 2022; Conant 2023). If only the
polyploid genome is available, a comparison of duplicated genes within the polyploid
genome may be made to estimate their likely time of divergence (Lynch and Conery
2000; Blanc and Wolfe 2004; Cui et al. 2006; Vanneste et al. 2013; Clark et al. 2019;
Chen and Zwaenepoel 2023). Syntenic relationships (self-synteny and with related

species) may also be used to corroborate WGD (Vandepoele et al. 2002; Hampson et
al. 2003; Wang et al. 2006; Parey et al. 2020).
86   Historically, the most common method to date a WGD event is to use a
87   molecular clock to calibrate the divergence times between paralogs in a polyploid
88   genome. The number of synonymous substitutions per synonymous site in protein-
89   coding genes (**Ks**) is calculated between all paralogous pairs, and if there is a peak at
90   $Ks>0,$ the position of this peak is used to infer the time of WGD (Blanc and Wolfe 2004;
91   Chen and Zwaenepoel 2023) and see figures 1 and 2.
92   Since paralogs can arise from a number of evolutionary processes, Ks histogram
shapes may be complex, and their interpretation may be challenging. Ks histograms
generally have a first peak at $Ks=0$, due to the constant birth and death of small-scale
duplications (SSDs). These SSD paralogs have recently arisen but have not yet been
shed, and their lifespan follows an exponential decay model (Lynch and Conery 2000).
If a species has undergone one or more WGD events, additional peaks will occur in the
histogram for $Ks>0$, and these peaks may even be overlapping. These additional peaks
are due to ohnologs (paralogs formed by WGD). The naive interpretation is that the
mode in ohnolog divergence times represents the time of the WGD event (Ohno 1970;
Blanc and Wolfe 2004). Univariate mixture models are used to empirically fit the peaks
(Schlueter et al. 2004; Cui et al. 2006; Vanneste et al. 2013; Tiley et al. 2018; Li and
Barker 2020; Chen and Zwaenepoel 2023).
104   However, the Ks-based approach has several pitfalls. The conversion between
Ks and time is not straightforward (Wolfe et al. 1987; Doyle and Egan 2010; Barba-
Montoya et al. 2018), Ks saturates for $Ks>2$, or roughly 200 million years (Cui et al.
2006; De La Torre et al. 2017; Li and Barker 2020), and multivariate fitting methods
have been shown to overfit distributions (Vanneste et al. 2013; Tiley et al. 2018;
Zwaenepoel and Van de Peer 2019). Restricting the Ks histogram to ohnologs may
improve resolution (Van de Peer 2004; Zwaenepoel and Van de Peer 2019; Sensalari et
al. 2022; Sutherland et al. 2024), but there remains significant methodological concerns.
112   While Ks-based methods can in theory correctly date the origin of a polyploid
lineage which traces back to a single individual (assuming the genomes began to
diverge at the same instant they duplicated (Doyle and Egan 2010)), the methods may
fail for allopolyploids (Thomas et al. 2017; Mccann et al. 2018; Wen et al. 2018;
Bouckaert et al. 2019; Conant 2023). This is because the peak of the Ks distribution
corresponds to the divergence time between the diploid parental species (hereon T_{DIV}),
not the time of origin of the polyploid (hereon T_{WGD}) (Doyle and Egan 2010; Thomas et
al. 2017; Chen and Zwaenepoel 2023), as shown in figure 1. Since plant genomes can
remain compatible for 10 mya or more after the last common ancestor (Senchina et al.
2003; Levin 2013) and up to 50 mya in one documented case (Rothfels et al. 2015), the
difference between T_{DIV} and T_{WGD} can be significant. Confounding the Ks peak with
T_{WGD} may also be problematic for autopolyploids, whose ohnologs may show complex
patterns of divergence post WGD (Gaeta and Chris Pires 2010; Parey et al. 2022; Lv et
al. 2024).
126   Several methods have been developed that can deal with the timing problems
unique to allopolyploidy. These methods rely on additional data from the parental
species or from broader sampling of the polyploid taxa (Thomas et al. 2017; Mccann et
al. 2018; Wen et al. 2018; Bouckaert et al. 2019; Conant 2023). Population-

demographic approaches have also been used with recent success to time WGD
(Gutenkunst et al. 2009; St Onge et al. 2012; Roux and Pannell 2015; Roux et al. 2017;
Blischak et al. 2023; Booker and Schrider 2023; Booker and Schrider 2023).
133   As part of the 1000 Plants (1KP) initiative (Leebens-Mack et al. 2019),
transcriptomic data from over 1000 plants spanning the plant kingdom was used to
compile *Ks* histograms. As in other works, Gaussian mixture models were fit and used
to detect WGD from *Ks* peaks (Clark et al. 2019; Qiao et al. 2019; Guo et al. 2020; Li
and Barker 2020). But a review of empirical *Ks* histogram shapes from the 1KP dataset
show great complexity beyond the Gaussian (Zwaenepoel and Van de Peer 2019;
Sensalari et al. 2022), which simulations have thus far have not been able to replicate
(Sutherland et al. 2024).
141   Here we present the *Ks* simulator SpecKS. SpecKS simulates the forward
evolution of polyploid genomes whereby the mode of speciation is not starkly allo or
autopolyploid, but instead a point in a 2D continuum model where the parental species'
divergence in time and genetic space may vary independently (fig. 3), which may better
align with our evolving definition of polyploidy (Parisod et al. 2010; Doyle and Sherman-
Broyles 2017). Thus, SpecKS is highly configurable (tables S2 and S3) and capable of
producing a rich array of *Ks* distribution shapes, modeling both the SSD and ohnolog
components. Here we use SpecKS to demonstrate the sensitivity of the *Ks* distribution
to several ancient speciation parameters, such as effective population sizes (N_e), gene
shedding rates, and the separation in time between T_{DIV} and T_{WGD}.
151

Results

Methods Overview

Ks histograms across the tree of life show a great diversity of distribution shapes, with
some showing a high degree of symmetry while others appearing more skewed (Li and
Barker 2020), suggesting the *Ks* histogram may bear signatures of evolutionary
parameters or events beyond the presence or absence of WGD. Thus, we sought to
build a simulation engine to investigate the potential effects of a variety of ancient
polyploid speciation parameters on the distribution of *Ks* histogram shapes.

We developed a novel simulation-engine, SpecKS, which models polyploid speciation
and evolution as a reticulate process. The simulation follows the evolution of an initial
ancestral genome, which diverges at a given time (T_{DIV}) into two sister diploid species.
These sister species later recombine at T_{WGD}, and the resulting polyploid continues to
evolve to the present day. A set of *i* gene-trees $\{G_i\}$ are embedded in this reticulate
topology with a configurable distribution (D_{TCOAL}) of coalescent times $\{T_{ci}\}$, which the
user might base on ancestral diversity (Π), effective population size (N_e), or some other
relationship. Genes (as a set of random strings of nucleotides, $\{N_i\}$) are evolved along
$\{G_i\}$ using PAML v4.10.7 (Yang 2007) under neutrality to the present time. *Ks* is then

calculated between the resulting paralogs, which are the leaves of the gene trees (fig.
4).
174
Implicit in this model is the concept that the polyploid continuum has more than one
dimension. We allow the degree of allopolyploidy to be a function of both length of the
time the parental species were separated (ΔT) and the ancestral diversity (Π) of the
subgenomes at T_{DIV}. This allows us to separate the effects of these critical speciation
parameters on the Ks histogram. We additionally disambiguate T_{DIV} and T_{WGD}, which
has confounded estimates of T_{WGD} in the past.
181
182   **SpecKS demonstrates that changes along either dimension of the 2D continuum**
183   **will deterministically affect the shape of the Ks histogram.**
184
The initial distribution of the divergence times for the gene trees ($D(T_c)$) is supplied by
the user as an input parameter. This is to allow the user as much flexibility as possible
with regard to modeling their system. To test if differences in these initial distribution
shapes can be theoretically detected even after WGD and thus potentially affect the Ks
histogram of extant polyploids, we simulated polyploids with modes of speciation from
all four corners of our 2D continuum (fig. 3). Specifically, we tested sets of allopolyploids
derived from (A) low-N_e ancestral species and a large ΔT between T_{DIV} and T_{WGD}, (B)
high-N_e ancestral species and a large ΔT, (C) low-N_e ancestral species and a small ΔT,
and (D) high-N_e ancestral species and a small ΔT. Set (B) comprises canonical
allopolyploids whose parental species were highly differentiated and have spent several
million years apart before hybridization. Set (C) represents polyploids whose parental
species were minimally differentiated and had no significant time between divergence
and conjunction. The off-diagonals (A & D) represent plausible, but less intuitive
polyploids. Specifically, set (A) are derived from parental species with little diversity
between them at speciation, but a great amount of time between parental divergence
and WGD. Set (D) are polyploids derived from parental species with a greater degree of
diversity between them at speciation (highly differentiated subpopulations leading to
separate species, but hybridization via polyploidy soon followed parental divergence).
For all four sets, we simulated a range of WGD times, from ancient to relatively recent
(80 - 0 MYA) (figure 5).
205
With regard to the diversity dimension of the 2D continuum, our results showed that the
polyploids derived from high-N_e ancestral species (A&B) show a more skew, fat-tailed
WGD component in the Ks distribution compared to the low-N_e polyploids (figure 5). We
also see that these differences persist for about 50 MY (figure 6, top). Along the ΔT
dimension, we see that differences in ΔT had little effect on the skew but will affect the

relative height differential between the SSD peak height and the WGD peak height
(figure 5, and figure 6 bottom).
213
214 **The error in the common method of estimating WGD time from the *Ks* histogram**
215 **peak scales with the degree of allopolyploidy.**
216
The "risk" of using the *Ks* peak to determine the timing of WGD, particularly for
allopolyploids, has been well described by (Thomas et al. 2017; Chen and Zwaenepoel
2023). However, the *Ks* peak continues to be used to determine the timing of WGD. We
were curious to see if SpecKS could be used to quantify the expected error in
estimating T_{WGD}, and potentially empirically relate the magnitude of T_{WGD} error to the
degree of allopolyploidy along a continuum. We were further interested to test if an
alternative method using the inferred start time of ohnolog-shedding would yield more
accurate estimates of T_{WGD}.
225
As expected, our simulations demonstrate that inferring T_{WGD} from the *Ks* peak may be
quite problematic (off by millions of years). Since the *Ks* peak gives the T_{DIV}, not the
T_{WGD}, it is no surprise that the error in estimates of T_{WGD} linearly scales with their
difference. Thus, the error in inferring T_{WGD} from the *Ks* peak is proportional to ΔT, the
degree of allopolyploidy as measured along the time axis of the 2D continuum (fig. 7,
top).
232
233 **An alternative, accurate T_{WGD} estimation method that is independent of the degree**
234 **of allopolyploidy.**
235
We tested the hypothesis that, since ohnolog-shedding can only begin after WGD
irrespective of the mode of polyploid speciation, using the proportion of duplicate genes
remaining may be a "less-risky" metric for inference, especially if the mode of speciation
is unclear. In practice, the rates of gene shedding would vary by lineage and gene
family, but for the purpose of this theoretical test, we assume the gene shedding rate is
constant and known *a priori* for our theoretical species. We thus simulated a range of *Ks*
histograms parameterized with a set gene shedding rate for a range of T_{DIV} and T_{WGD},
for polyploids across a 2D polyploid continuum, thus recreating the range of possible *Ks*
histograms for this hypothetical polyploid species, under a range of speciation
scenarios. The simulations revealed a clear logarithmic relationship between the
number of genes shed and T_{WGD}, invariant to the mode of speciation (fig. S2, left). We
were thus able to use the logarithmic relationship between T_{WGD} and the proportion of
ohnologs remaining to determine T_{WGD} from the *Ks* histogram. In fig. 7, bottom left, we
demonstrate the accuracy of this method, by comparing the input (true) T_{WGD} to the
recovered T_{WGD}, revealing a high accuracy, with r-value of 0.998 and a standard error of

0.004 MY. In fig. 7, bottom right, we show that the error (the difference between the true
and inferred T_{WGD}) does not increase with degree of allopolyploidy, as measured along
the ΔT dimension.
254
255 **SpecKS recreates the main features of empirical *Ks* histograms**
256
Next, we wanted to know if the evolutionary models incorporated in SpecKS would
sufficiently reproduce the exponential decay curves typical of the primary "SSD" peak of
the *Ks* histogram, as well as the secondary "WGD" peak, which contemporary works
have had difficulty to replicate (Tiley et al. 2018; Sutherland et al. 2024). We selected
three tetraploid species, *Coffea arabica, Zea mays* and *Populus trichocarpa* to
demonstrate this. *Coffea arabica* is a relatively recent allopolyploid (WGD ~10-600KYA).
Zea mays was formed by polyploidy ~14 MYA, and *Populus* ~56 MYA (Gaut and
Doebley 1997; Yu et al. 2011; Dai et al. 2014; Salojarvi et al. 2023). We show in figure 8
that SpecKS can well replicate both the SSD and ohnolog components of the *Ks*
histograms for all three species. It is particularly interesting to note that the simulated *Ks*
plots match the exponential-lognormal mixture models which have historically had the
greatest success fitting observed *Ks* distributions. We note that this is an emergent
property of our simulation, and no lognormal distributions were input. Furthermore, to
achieve the best fit, the SpecKS input parameters were optimized to minimize the
RMSE error (table S4), yielding T_{WGD} and T_{DIV} which correspond well with estimated
dates from other sources (table S5) (Gaut and Doebley 1997; Yu et al. 2011; Dai et al.
2014; Salojärvi et al. 2024). We also note that the main difference between the
simulated histograms and the true histograms is that the true histograms maintain a set
of paralogs whose numbers do not decay over time and whose *Ks* distribution appears
flat. One explanation for this phenomenon is that these paralogs are maintained by
selection, and thus not presently modeled by our system.
278
279
280 **Using SpecKS to estimate difference in time ($\Delta T = T_{DIV} - T_{WGD}$), and the diversity of**
281 **the ancestral species (Π), from an input *Ks* histogram.**
282
283 As we demonstrated previously, the changes along either dimension of the 2D
284 continuum deterministically affect the shape of the *Ks* histogram. Thus, we wanted to
285 know if, given the *Ks* histogram, could the initial parameters of polyploid speciation be
286 recovered? Since we have already demonstrated that T_{WGD} can be recovered (figure 7),
287 what remains is to demonstrate the recovery of T_{DIV} and Π.
288
The peak of the *Ks* histogram has long been attributed to T_{DIV}, albeit often confused
with T_{WGD} (Blanc and Wolfe 2004), but disambiguated in (Thomas et al. 2017; Chen and

Zwaenepoel 2023), thus we expected that the peak of the SpecKS-generated *Ks*

histogram, once converted from *Ks*-space to time, SpecKS, should given an accurate

recovery of the input T_{DIV}.

Indeed, SpecKS corroborates this expectation (figure S2, right), and we demonstrate

the accuracy of this method by comparing the input (true) T_{DIV} to the recovered T_{DIV},

revealing a high accuracy, with r-value of 0.995 and a standard error of 0.008 MY. We

note the caveat that this accuracy is contingent on the accuracy of the conversion factor

between time and *Ks*, which is a configurable parameter in our simulation. Since $\Delta T =$

$T_{DIV} - T_{WGD}$, the ability to estimate both T_{DIV} and T_{WGD} yields the estimated ΔT.

To test the recovery of ΔT, we generated a dataset of 160 simulations (dataset 1,

parameters described in table 3 for a variety of modes of speciation across the 2D

continuum, with a range of T_{DIV}, T_{WGD}, and Π. With these simulated datasets, using a ⅓

test, ⅔ training approach with the data, we trained a logistic regression classification

model to discriminate between large, medium or small ΔT. Within the context of our

simulated results, this approach gave 100% accuracy for WGD up to 80 MYA (figure 9).

To test the recovery of Π, we chose a model (described in Methods) that would relate Π

to the distribution of ortholog divergence times, since it is that distribution, which is input

to SpecKS, not Π directly. Based on this model, we were able to use a range of N_e to

generate a set of initial gene-tree divergence distributions as input to SpecKS. Using a

⅓ test, ⅔ training approach with the data, we trained a logistic regression classification

model on dataset 1 to discriminate between high, medium or low ancestral N_e. Within

the context of our simulated results, this approach gave 94% accuracy for WGD up to

80 MYA (figure 10).

Using SpecKS to infer the polyploid continuum with *Ks* distributions from

transcriptomic data from >200 angiosperms

Recent work has suggested that the majority of polyploid lineages may be derived from

allopolyploidy rather than autopolyploidy (Wang et al. 2019), although also see (Soltis et

al. 2007; Barker et al. 2016). Thus, we were interested to test if the logistical regression

model discussed previously, when applied to empirical *Ks* data from species across the

plant tree of life, would suggest high or low diversity between ancient subgenomes.

Thus we applied our low-vs-high N_e classifier to over 200 real *Ks* observations made

public by (Li and Barker 2020), derived from transcriptomic data from the 1KP study

(Leebens-Mack et al. 2019). We were able to classify 228 WGD events, resulting in 12,

38 and 178, low-N_e, medium-N_e and high-N_e determinations, respectively (fig. 11). Our

results suggest that >94.7% of the WGD in the 1KP dataset are moderately or strongly

allopolyploid, and the remaining 5.3% fall in an indeterminate category, with low
diversity between the initial orthologs. These results are concordant with (Wang et al.
2019), suggesting that the majority of ancient polyploidization events which contributed
to the genetic conduit may be derived from allopolyploidization with high ancestral N_e.
An alternative interpretation is that both our method and the method from (Wang et al.
2019) may under-report autopolyploids. One explanation for this might be that for
autopolyploids, the gene copies have not diverged sufficiently to form true paralogs,
thus the gene-pairs may not be detected as a multiplicity in transcriptomic analysis
(Mayfield-Jones et al. 2013; Garsmeur et al. 2014). Furthermore, our logistical
regression model was trained on simulations drawn from a continuum of allopolyploidy,
thus may not be applicable to autopolyploids whose speciation parameters lie outside
the scope of the SpecKS implementation. While much work exists describing the
complexity of post-WGD divergence of ohnologs in specific autopolyploid systems,
there currently exists no general, empirically-validated model (Parisod et al. 2010;
Spoelhof et al. 2017; Parey et al. 2022; Lallemand et al. 2023). While the derivation of
such a model is beyond the scope of this paper, the SpecKS architecture may prove to
be well-suited to its incorporation and assessment.
348
349

350 **Discussion**
351
352 Here we present the polyploid genome evolution simulator SpecKS and use it to
353 demonstrate the dependency of the shape of the Ks histogram on critical polyploid
354 speciation parameters.
355
Our simulations also show that the shape and skew of the *Ks* histogram is sensitive to
the evolutionary history of the ancestral polyploidization event: levels of ancestral
genetic divergence, as well as the relative timing of T_{WGD} and T_{DIV}, yielding
characteristically different *Ks* distributions. We also demonstrate that these differences
persist for 10s of millions of years (fig. 5). Furthermore, we have shown that the tail of
the *Ks* histogram depends on the divergence between the ancestral subgenomes of the
polyploid (fig. 6) and we demonstrate with simulations that the *Ks* distribution can be
used to test inferences linking N_e, Π, T_{WGD}, T_{DIV}, and distributions of ohnolog divergence
times (fig. 9 and 10). Specifically, we demonstrate that the commonly used method for
estimating T_{WGD} from the peak of the *Ks* histogram will be increasingly inaccurate for
greater differences in time between T_{WGD} and T_{DIV} (fig. 7), and we develop an alternative
method that works equally well for a variety of polyploid speciation scenarios. We
contend that the *Ks* distribution is therefore information rich, with the potential to aid in
the estimation of a variety of polyploid speciation parameters and scenarios.
370

We additionally show that SpecKS-generated histograms realistically capture the main
features of empirical *Ks* histograms. For instance, in figure 8, we parameterized
SpecKS to fit the observed *Ks* histograms of three well-studied species, coffee, maize
and poplar. In all cases the histogram features corresponding to the SSDs and ohnolog
peaks appear to be well replicated, with overall RMSE less than 2 in all cases (fig. 8).
This demonstrates that the parameters supplied may be reasonably estimable under the
SpecKS model. Interestingly, we note that the main difference between the simulated
histograms and the true histograms, is that the true histograms maintain very old
paralogous pairs ($Ks > T_{DIV}$) whose numbers do not decay over time. In contrast, in our
simulation, the number of paralogous pairs retained asymptotically approaches zero
over time. The latter behavior is expected under the SpecKS model, because both
SSDs and ohnologs are shed genes following a parameterized decay model (described
in Methods).
384
One explanation for this phenomenon is that the paralogous pairs that persist in the true
Ks histogram and not in the simulations may be duplicates that are maintained due to
selection or potentially under linked selection (hitchhiking) and thus not presently
modeled by our system. These persistent paralogs are more apparent in *Coffea arabica*
and *Zea mays* (fig. 8). Since these species have undergone recent domestication (Gaut
and Doebley 1997; Salojarvi et al. 2023; Yang et al. 2023), it may be possible that
domestication has hindered efficient gene shedding or promoted paralog retention
(Gaut et al. 2018). We also note that SpecKS currently only models ohnologs and
SSDs; some paralogs of a different origin, like tandem duplications, may be retained for
longer or shorter periods of time, and thus would be evident in the empirical histograms
but not in the modeled histograms.
396
While SpecKS internal 2D continuum model is currently best suited to model varying
degrees of allopolyploidy, it has not yet been extended to autopolyploidy. Furthermore,
despite its utility, SpecKS currently represents a simplistic model which does not include
selection, population dynamics, the effects of domestication, and other factors which
would reasonably impact the *Ks* histogram. SpecKS' internal model of evolution (GY94
with equal equilibrium codon frequencies) is simplistic, and the rate of *Ks* accumulation
is assumed to be constant over time. Furthermore, our ability to test SpecKS against
observed *Ks* histograms is limited to tetraploids whose WGD events are well-separated
in time. This is because SpecKS does not yet model multiple superimposed rounds of
WGD. An appropriate next step would be to iteratively include more evolutionary
complexity, thus making the simulation both more realistic and more testable.
408
We note that many of the parameters incorporated into SpecKS are lineage specific
(most significantly: rates of gene loss, the *Ks* distribution at the time of ancestral

divergence, the rate of the molecular clock) and are difficult to ascertain *a priori*.
Furthermore, the simplistic relationship between N_e and the gene tree coalescent
distribution we selected as input (described in more detail in the Methods section) might
not be appropriate for many use cases, including autopolyploids, or if complex
population structure existed within the ancestral species. For these situations, we
recommend the user consider the D_{TCOAL} most appropriate to their effort.
417
For simplicity, the discrimination models presented were drawn from simulation runs
where a small number of parameters were fixed (for example, fixed gene-birth-and-
death rate and molecular clock). These models are presented as proof of concept and
not meant to analyze any particular lineage. When making lineage-specific inferences,
we caution users to parameterize SpecKS with lineage-appropriate values, or if they are
unknown, to use SpecKS over a range of likely values, to model the impact of
uncertainty in their value.
425
426 **Materials and Methods**
427
428 **Data**
429
430 Transcriptomic data: *Ks* histograms in fig. 8 were generated from transcriptomic data for
431 *Coffea arabica*, *Zea mays* and *Populus trichocarpa*. Transcriptomic data were sourced
432 from NCBI (NCBI assembly GCF_003713225.1, GCF_902167145.1 and
433 GCF_000002775.5), and the software package KsRates (Sensalari, 2022) was used to
434 calculate the *Ks* histograms for each species.
435
1KP data: In results section 8, the low-vs-high N_e discrimination model is run on
histograms generated from *Ks* data from the 1KP dataset. The raw, empirical *Ks* data is
available at (https://gitlab.com/barker-lab/1KP/-/tree/master/1KP_ksplots). These *Ks*
data were generated by (Leebens-Mack et al. 2019; Li and Barker 2020). *Ks* histograms
from this data were plotted using matplotlib (Hunter 2007), and the skew of the *Ks*
histogram was measured using the same feature-extraction method used in the
derivation of the N_e discrimination model, and then input directly to the N_e discrimination
model.
444
445 **Logistic Regression Models**
446
An example use-case for SpecKS is the provision of simulated truth and training data
for the derivation of inference models. Here, we describe two discrimination models
used in this paper, one to infer N_e, and another to infer ΔT. For both of these models,
we used the logistic regression package from scikit-learn (Pedregosa et al. 2011).
SpecKS-simulated *Ks* histograms were used to derive features, and SpecKS input
parameters were used to derive target variables, which we organized into three
categories (low, medium, and high) for each classifier. With respect to target variable

categories, for the ΔT discrimination model, the target variable was based on whether
$\Delta T = T_{DIV} - T_{WGD} < 5, < 30,$ or $>= 30$ MY. For the N_e discrimination model, the target
variable was based on whether $N_e < 5, < 10,$ or $>= 10 * 10^6$. These levels were
selected to span realistic values (Gossmann et al. 2010). With respect to feature data,
for the ΔT discrimination model, we used the difference between estimated T_{WGD} and
T_{DIV} as our single feature, deriving both from the *Ks* histogram directly, using algorithms
for estimating T_{WGD} and T_{DIV} as discussed previously in the results section (figures 7 and
S2). For the N_e discrimination model, we used the natural logarithm of the skew of the
Ks histogram (measured as the difference between the x-value of the *Ks* peak and the
x-value of the *Ks* center of mass, in *Ks*-space) as the single key feature. The models
were trained with a ⅓ test, ⅔ training approach, splitting datasets as appropriate.
465
466
467 **Setting the gene-tree divergence time distribution "D_{TCOAL}"**
468
In the SpecKS simulator, the initial distribution of the divergence times for the gene
trees is supplied by the user as an input parameter. This allows the user as much
flexibility as possible with regard to modeling their system. Throughout this paper, a
simplistic input distribution D_{TCOAL} was used, which relates ancestral genetic diversity
(Π) to the distribution of ortholog divergence times, based on the assumption of some
level of allopolyploidy. We assumed that the ancestral diploid population was in
panmixia, and we used the Kingman coalescent (Kingman 1982) to derive a diversity of
initial gene-tree divergence times.
477
Under this model, the ancestral diploid species exists with some standing genetic
variation which scales with population size, such that N_e (effective population size) is
proportional Π. The ancestral species subsequently (at time T_{DIV}) diverges to give rise to
two diploid sister species, which evolve forward for a given amount of time ($\Delta T = T_{DIV} -$
T_{WGD}), before conjoining to form the diploid lineage at T_{WGD}. Prior to T_{DIV}, the
coalescence times for gene copies between two random individuals in the ancestral
population can be approximated by the Kingman coalescent, given Π. If these two
ancestral individuals found new species, their individual sets of gene copies are now
separated by speciation and are redefined as orthologs. The diversity of coalescent
times becomes the initial diversity in nodes for the bifurcating gene trees at T_{DIV}, which
follows an exponential distribution, under the Kingman coalescent (Kingman 1982).
489
490 Under this simple one-genome-one-species model, for more diverse ancestral species,
491 we see more initial diversity in node times, and a greater skew in the *Ks* distribution
492 towards the past. Mathematically, this is because the decay constant in the Kingman
493 coalescent is inversely proportional to N_e. Note that this model may not be appropriate
494 for autopolyploids, which is beyond the scope of this paper.
495

To obtain the Kingman coalescent from N_e, it is necessary to assume a generation time
(G_t). In all our simulations, for simplicity, we assumed at $G_t=1$. Since the *Ks* distributions
generated and the inferences made from them were done in *Ks* space, the exact G_t is
immaterial. It only matters that the Ks-to-My conversion factor, which is a configurable
input, properly factors in the G_t at the initiation of the simulation, and that this same
conversion factor is taken into account by the end-user when relating the output *Ks*
plots to chronological time.
503
504 **SpecKS Implementation**
505
SpecKS is implemented as a pipeline application in python. SpecKS takes as input an
XML configuration file listing the simulation parameters (Table 1, S2 & S3) and outputs
a text file (.csv) of all pairwise *Ks* accumulated between all gene pairs (ohnologs and
small-scale duplications) for each simulated genome. Architecturally, SpecKS is
designed as a pipeline with eight modules which are executed sequentially for each
polyploid in the simulation (fig. 4). The modules functions are (1) species tree
generation, (2) gene tree generation, (3) application of a gene birth and death model,
(4) application of a post WGD gene shedding model, (5) gene sequence evolution, (6)
the *Ks* calculation between gene pairs, (7) *Ks* histogrammer and (8) final results
assembly. We give details for each module below.
516
517 (1) Species tree generation: A newick-formatted species tree is generated for each
518 polyploid. This is aways in the format (O:T_{SIM}, (P1:T_{DIV}, P2:T_{DIV}): T_{SIM}-T_{DIV}), where T_{SIM}
519 specifies the full simulation time. P1 and P2 denote the subgenomes of the polyploid,
520 while O denotes the outgroup.
521
(2) Gene tree generation: Newick-formatted gene trees are generated from the
simulated species tree. Random variations in gene divergence times are introduced
according to the distribution specified in the configuration file, allowing for the
introduction of a range of divergence times for the ohnologs (genes duplicated by
whole-genome duplication). The distribution is configurable and might in theory be
selected based on the ancestral genetic diversity (Π), generation time, and evolutionary
model. The number of gene trees is also configurable, with a default set at 3000. The
distribution selected for our simulations was based on the Kingman coalescent, and we
give details in the methods section.
531
(3) Gene birth and death model: The gene birth and death model introduces small-scale
duplications ("SSDs") into the simulation. For each gene tree, genes are randomly born
at a rate specified in the configuration file, modeled as a Poisson process (Zhao et al.
2015). Genes are assigned a death-date at birth, with a life span drawn randomly from
an exponential decay distribution (Lynch and Conery 2000; Lynch et al. 2001). Gene
birth rate and mean life expectancy are configurable (Lynch and Conery 2003; Guo

2013). Genes born which will be dead before the simulation concludes are pruned at
this time for computational efficiency.
540
(4) Ohnolog gene shedding model: Genes duplicated by WGD also have a mean life
expectancy given by an exponential decay distribution (Ren et al. 2018). The proportion
of WGD duplicates which will be dead before the simulation concludes are calculated
based on WGD time and pruned for computational efficiency. For simplicity and
traceability, ohnolog duplicates are always removed from the P1 branch. Ohnolog life
expectancy is configurable and based on (Maere et al. 2005; Guo 2013).
547
(5) Gene sequence evolution: For each simulated gene tree, we simulated codon
sequences using the EVOLVER program within PAML v4.10.7 (Yang and Nielsen 2000:
200; Yang 2007). PAML was configured as in (Tiley et al. 2018) to simulate codon
evolution along each finalized gene tree using a Goldman–Yang (GY94) model of codon
evolution (Goldman and Yang 1994; Yang and Nielsen 1998) with equal equilibrium
codon frequencies, a transition/transversion rate ratio of 2 and a global dN/dS of 0.2 as
in (Tiley et al. 2018). The number of codons per alignment is set by default to 1000 as in
(Tiley et al. 2018), in line with reported plant transcript lengths (Luo et al. 2019; S.
Zhang et al. 2020; Al-Dossary et al. 2023), but is configurable.
557
558   (6) *Ks* calculation: *Ks* is calculated between all pairs of terminal genes in a given gene
559   tree, across all gene trees, using the CODEML program within PAML v4.10.7. Codon
560   frequencies, transition/transversion rate and *dN/dS* were set to match those used in
561   EVOLVER (Yang and Nielsen 2000; Yang 2007). *Ks* per million years is 0.01 by default
562   and is configurable (Blanc and Wolfe 2004; Tiley et al. 2018).
563
564   (7) *Ks* histogrammer: SpecKS thereon generates two default histograms (one with a
565   configurable maximum *Ks*, and one with no maximum *Ks*). The existence of these
566   histograms is only meant to give a confirmation of the run success. It is expected that
567   the end users will use their own visualization tools to generate more elegant histograms.
568
569   (8) Final results assembly: The final results assembler collates the CODEML results,
570   producing a single .csv file containing the *Ks* results for each polyploid. Additional
571   products of the simulation include *Kn* results (histogram of accumulated non-
572   synonymous mutations between paralogs) for the polyploid, and *Ks* and *Kn* results for
573   the outgroup.
574
575   Running SpecKS: SpecKS is run calling the "SpecKS.py" function from python and
576   passing it the config xml file. For example, "python3 SpecKS.py myconfig.xml"
577
578   SpecKS output: The SpecKS output file is a .csv file, with each line giving information
579   for a unique gene-pair. The data are organized into 4 columns, giving the gene-tree
580   name of the pair, the names of the two genes in the pair (the leaves of the tree), the *Ks*

581 value, and the path to the CODEML output file from which the value
582 originated,respectively.
583
Parallelization: For simplicity, SpecKS is not internally parallelized. A single polyploid of
about 3000 genes, 1000 codons long, simulated over 100 MY, runs on a typical cluster
(e.g. 16-node Dual-10 Xeon CPU, E5 -2630v4 2.20GHz, 12.8GB RAM per core) in
under 10 minutes. To simulate batches of polyploids, we recommend submitting each
polyploid as a separate batch job to a job scheduler such as SGE (Oracle) or SLURM
(SchedMD).
590
591 **Source code and data availability**
592
593 SpecKS is available from the github repository https://github.com/tamsen/SpecKS.
594 SpecKS input files used to drive the simulation and the scripts used to generate the
595 figures are at https://github.com/tamsen/SpecKS_paper_scripts.
596
597
598 **Acknowledgments**
599

We would like to thank members of the Barker and Sethuraman Labs for valuable inputs
during the conceptualization stage of this project. We would like to acknowledge
Matthew Hahn and two anonymous reviewers for valuable suggestions on the first
version of this manuscript. AS and TD were both supported by NSF CAREER: 2147812
and two CSUBIOTECH grants. All computations were performed on the *mesxuuyan*
cluster at San Diego State University, which was supported by startup funds and NSF
ABI: 1564659 to AS.

607
608
609 **References**
610

Al-Dossary O, Furtado A, KharabianMasouleh A, Alsubaie B, Al-Mssallem I, Henry RJ. 2023.
Long read sequencing to reveal the full complexity of a plant transcriptome by targeting
both standard and long workflows. *Plant Methods* 19:112.
Arrigo N, Barker MS. 2012. Rarely successful polyploids and their legacy in plant genomes.
Current Opinion in Plant Biology 15:140–146.
Baduel P, Bray S, Vallejo-Marin M, Kolář F, Yant L. 2018. The "polyploid hop": shifting
challenges and opportunities over the evolutionary lifespan of genome duplications.
Frontiers in Ecology and Evolution 6:117.
Baniaga AE, Marx HE, Arrigo N, Barker MS. 2020. Polyploid plants have faster rates of
multivariate niche differentiation than their diploid relatives. *Ecology Letters* 23:68–78.
Barba-Montoya J, Dos Reis M, Schneider H, Donoghue PCJ, Yang Z. 2018. Constraining
uncertainty in the timescale of angiosperm evolution and the veracity of a Cretaceous
Terrestrial Revolution. *New Phytol* 218:819–834.
Barker MS, Arrigo N, Baniaga AE, Li Z, Levin DA. 2016. On the relative abundance of
autopolyploids and allopolyploids. *New Phytologist* 210:391–398.
Blanc G, Wolfe KH. 2004. Widespread paleopolyploidy in model plant species inferred from age
distributions of duplicate genes. *Plant Cell* 16:1667.

Blischak PD, Sajan M, Barker MS, Gutenkunst RN. 2023. Demographic history inference and
the polyploid continuum. *Genetics* 224:iyad107.
Bomblies K, Jones G, Franklin C, Zickler D, Kleckner N. 2016. The challenge of evolving stable
polyploidy: could an increase in "crossover interference distance" play a central role?
Chromosoma 125:287–300.
Booker WW, Schrider DR. 2023. The genetic consequences of range expansion and its
influence on diploidization in polyploids. *bioRxiv* [Internet]. Available from:
https://www.biorxiv.org/content/early/2023/10/21/2023.10.18.562992
Bouckaert R, Vaughan TG, Barido-Sottani J, Duchêne S, Fourment M, Gavryushkina A, Heled
J, Jones G, Kühnert D, De Maio N, et al. 2019. BEAST 2.5: An advanced software
platform for Bayesian evolutionary analysis. *PLoS Comput Biol* 15:e1006650.
Bowers JE, Chapman BA, Rong J, Paterson AH. 2003. Unravelling angiosperm genome
evolution by phylogenetic analysis of chromosomal duplication events. *Nature* 422:433–
438.
Buggs RJA, Zhang L, Miles N, Tate JA, Gao L, Wei W, Schnable PS, Barbazuk WB, Soltis PS,
Soltis DE. 2011. Transcriptomic shock generates evolutionary novelty in a newly formed,
natural allopolyploid plant. *Curr Biol* 21:551–556.
Carretero-Paulet L, Van de Peer Y. 2020. The evolutionary conundrum of whole-genome
duplication. *Am J Bot* 107:1101–1105.
Chen H, Zwaenepoel A. 2023. Inference of ancient polyploidy from genomic data. *Methods Mol*
Biol 2545:3–18.
Clark JW, Donoghue PC. 2018. Whole-genome duplication and plant macroevolution. *Trends in*
Plant Science 23:933–945.
Clark JW, Donoghue PC. 2023. Constraining whole-genome duplication events in geological
time. In: Polyploidy: Methods and Protocols. Springer. p. 139–154.
Clark JW, Puttick MN, Donoghue PCJ. 2019. Origin of horsetails and the role of whole-genome
duplication in plant macroevolution. *Proc Biol Sci* 286:20191662.
Clausen J, Keck DD, Hiesey WM. 1945. Experimental studies on the nature of species. 2. Plant
evolution through amphiploidy and autoploidy, with examples from the Madiinae.
Experimental studies on the nature of species. 2. Plant evolution through amphiploidy
and autoploidy, with examples from the Madiinae.
Comai L, Tyagi AP, Winter K, Holmes-Davis R, Reynolds SH, Stevens Y, Byers B. 2000.
Phenotypic instability and rapid gene silencing in newly formed arabidopsis
allotetraploids. *Plant Cell* 12:1551.
Conant GC. 2023. POInT: Modeling polyploidy in the era of ubiquitous genomics. *Methods Mol*
Biol 2545:77–90.
Cui L, Wall PK, Leebens-Mack JH, Lindsay BG, Soltis DE, Doyle JJ, Soltis PS, Carlson JE,
Arumuganathan K, Barakat A, et al. 2006. Widespread genome duplications throughout
the history of flowering plants. *Genome Res* 16:738–749.
Dai X, Hu Q, Cai Q, Feng K, Ye N, Tuskan GA, Milne R, Chen Y, Wan Z, Wang Z, et al. 2014.
The willow genome and divergent evolution from poplar after the common genome
duplication. *Cell Research* 24:1274–1277.
Darlington CD. 1937. Recent advances in cytology. *Recent advances in cytology.*
David KT, Oaks JR, Halanych KM. 2020. Patterns of gene evolution following duplications and
speciations in vertebrates. *PeerJ* 8:e8813.
De La Torre AR, Li Z, Van de Peer Y, Ingvarsson PK. 2017. Contrasting rates of molecular
evolution and patterns of selection among gymnosperms and flowering plants. *Molecular*
Biology and Evolution 34:1363–1377.
Deb SK, Edger PP, Pires JC, McKain MR. 2023. Patterns, mechanisms, and consequences of
homoeologous exchange in allopolyploid angiosperms: a genomic and epigenomic
perspective. *New Phytologist*.

Dodsworth S, Chase MW, Leitch AR. 2016. Is post-polyploidization diploidization the key to the
evolutionary success of angiosperms? *Botanical Journal of the Linnean Society* 180:1–5.
Douglas GM, Gos G, Steige KA, Salcedo A, Holm K, Josephs EB, Arunkumar R, Ågren JA,
Hazzouri KM, Wang W. 2015. Hybrid origins and the earliest stages of diploidization in
the highly successful recent polyploid *Capsella bursa-pastoris*. *Proceedings of the
National Academy of Sciences* 112:2806–2811.
Doyle JJ, Egan AN. 2010. Dating the origins of polyploidy events. *New Phytol* 186:73–85.
Doyle JJ, Sherman-Broyles S. 2017. Double trouble: taxonomy and definitions of polyploidy.
New Phytologist 213:487–493.
Estep MC, McKain MR, Diaz DV, Zhong J, Hodge JG, Hodkinson TR, Layton DJ, Malcomber
ST, Pasquet R, Kellogg EA. 2014. Allopolyploidy, diversification, and the Miocene
grassland expansion. *Proceedings of the National Academy of Sciences* 111:15149–
15154.
Gaeta RT, Chris Pires J. 2010. Homoeologous recombination in allopolyploids: the polyploid
ratchet. *New Phytologist* 186:18–28.
Garsmeur O, Schnable JC, Almeida A, Jourda C, D'Hont A, Freeling M. 2014. Two
Evolutionarily Distinct Classes of Paleopolyploidy. *Molecular Biology and Evolution*
31:448–454.
Gaut BS, Doebley JF. 1997. DNA sequence evidence for the segmental allotetraploid origin of
maize. *Proceedings of the National Academy of Sciences* 94:6809–6814.
Gaut BS, Seymour DK, Liu Q, Zhou Y. 2018. Demography and its effects on genomic variation
in crop domestication. *Nature Plants* 4:512–520.
Goldman N, Yang Z. 1994. A codon-based model of nucleotide substitution for protein-coding
DNA sequences. *Mol Biol Evol* 11:725–736.
Gossmann TI, Song B-H, Windsor AJ, Mitchell-Olds T, Dixon CJ, Kapralov MV, Filatov DA,
Eyre-Walker A. 2010. Genome wide analyses reveal little evidence for adaptive
evolution in many plant species. *Mol Biol Evol* 27:1822–1832.
Guo X, Mandáková T, Trachtová K, Özüdoğru B, Liu J, Lysak MA. 2020. Linked by ancestral
bonds: Multiple whole-genome duplications and reticulate evolution in a Brassicaceae
tribe. *Molecular Biology and Evolution* 38:1695–1714.
Guo Y-L. 2013. Gene family evolution in green plants with emphasis on the origination and
evolution of *Arabidopsis thaliana* genes. *Plant J* 73:941–951.
Gutenkunst RN, Hernandez RD, Williamson SH, Bustamante CD. 2009. Inferring the joint
demographic history of multiple populations from multidimensional SNP frequency data.
PLoS genetics 5:e1000695.
Hampson S, McLysaght A, Gaut B, Baldi P. 2003. LineUp: statistical detection of chromosomal
homology with application to plant comparative genomics. *Genome Res* 13:999–1010.
Hoegg S, Brinkmann H, Taylor JS, Meyer A. 2004. Phylogenetic timing of the fish-specific
genome duplication correlates with the diversification of teleost fish. *Journal of Molecular
Evolution* 59:190–203.
Hunter JD. 2007. Matplotlib: A 2D graphics environment. *Computing in science & engineering*
9:90–95.
Kingman JFC. 1982. The coalescent. *Stochastic Processes and their Applications* 13:235–248.
Lallemand T, Leduc M, Desmazières A, Aubourg S, Rizzon C, Landès C, Celton J-M. 2023.
Insights into the evolution of ohnologous sequences and their epigenetic marks post-
WGD in *Malus domestica*. *Genome Biology and Evolution* 15:evad178.
Laurent S, Salamin N, Robinson-Rechavi M. 2017. No evidence for the radiation time lag model
after whole genome duplications in Teleostei. *PLoS One* 12:e0176384–e0176384.
Le Comber SC, Ainouche ML, Kovarik A, Leitch AR. 2010. Making a functional diploid: from
polysomic to disomic inheritance. *New Phytol* 186:113–122.
Leebens-Mack JH, Barker MS, Carpenter EJ, Deyholos MK, Gitzendanner MA, Graham SW,

Grosse I, Li Z, Melkonian M, Mirarab S, et al. 2019. One thousand plant transcriptomes and the phylogenomics of green plants. *Nature* 574:679–685.

Levin DA. 2013. The timetable for allopolyploidy in flowering plants. *Ann Bot* 112:1201–1208.

Li Z, Baniaga AE, Sessa EB, Scascitelli M, Graham SW, Rieseberg LH, Barker MS. 2015. Early genome duplications in conifers and other seed plants. *Science advances* 1:e1501084.

Li Z, Barker MS. 2020. Inferring putative ancient whole-genome duplications in the 1000 Plants (1KP) initiative: access to gene family phylogenies and age distributions. *GigaScience* 9:giaa004.

Li Z, McKibben MTW, Finch GS, Blischak PD, Sutherland BL, Barker MS. 2021. Patterns and Processes of Diploidization in Land Plants. *Annual Review of Plant Biology* 72:387–410.

Li Z, Tiley GP, Galuska SR, Reardon CR, Kidder TI, Rundell RJ, Barker MS. 2018. Multiple large-scale gene and genome duplications during the evolution of hexapods. *Proceedings of the National Academy of Sciences* 115:4713–4718.

Lott M, Spillner A, Huber KT, Moulton V. 2009. PADRE: a package for analyzing and displaying reticulate evolution. *Bioinformatics* 25:1199–1200.

Luo D, Zhou Q, Wu Y, Chai X, Liu W, Wang Y, Yang Q, Wang Z, Liu Z. 2019. Full-length transcript sequencing and comparative transcriptomic analysis to evaluate the contribution of osmotic and ionic stress components towards salinity tolerance in the roots of cultivated alfalfa (Medicago sativa L.). *BMC Plant Biology* 19:32.

Lv Z, Addo Nyarko C, Ramtekey V, Behn H, Mason AS. 2024. Defining autopolyploidy: Cytology, genetics, and taxonomy. *American Journal of Botany*:e16292.

Lynch M, Conery JS. 2000. The evolutionary fate and consequences of duplicate genes. *Science* 290:1151–1155.

Lynch M, Conery JS. 2003. The evolutionary demography of duplicate genes. *Genome Evolution: Gene and Genome Duplications and the Origin of Novel Gene Functions*:35–44.

Lynch M, O'Hely M, Walsh B, Force A. 2001. The probability of preservation of a newly arisen gene duplicate. *Genetics* 159:1789–1804.

Mable BK. 2013. Polyploids and hybrids in changing environments: winners or losers in the struggle for adaptation? *Heredity* 110:95–96.

Madlung A. 2013. Polyploidy and its effect on evolutionary success: old questions revisited with new tools. *Heredity* 110:99–104.

Maere S, De Bodt S, Raes J, Casneuf T, Van Montagu M, Kuiper M, Van de Peer Y. 2005. Modeling gene and genome duplications in eukaryotes. *Proc Natl Acad Sci U S A* 102:5454–5459.

Mason AS, Wendel JF. 2020. Homoeologous exchanges, segmental allopolyploidy, and polyploid genome evolution. *Frontiers in Genetics*:1014.

Mayer VW, Aguilera A. 1990. High levels of chromosome instability in polyploids of Saccharomyces cerevisiae. *Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis* 231:177–186.

Mayfield-Jones D, Washburn JD, Arias T, Edger PP, Pires JC, Conant GC. 2013. Watching the grin fade: tracing the effects of polyploidy on different evolutionary time scales. *Semin Cell Dev Biol* 24:320–331.

Mccann J, Jang T-S, Macas J, Schneeweiss GM, Matzke NJ, Novák P, Stuessy TF, Villaseñor JL, Weiss-Schneeweiss H. 2018. Dating the species network: Allopolyploidy and repetitive DNA evolution in American daisies (*Melampodium sect. Melampodium*, Asteraceae). *Syst Biol* 67:1010–1024.

McClintock B. 1929. A cytological and genetical study of triploid maize. *Genetics* 14:180.

Meirmans P, Van Tienderen P. 2013. The effects of inheritance in tetraploids on genetic diversity and population divergence. *Heredity* 110:131–137.

Nieto Feliner G, Casacuberta J, Wendel JF. 2020. Genomics of evolutionary novelty in hybrids

and polyploids. *Frontiers in Genetics* 11:792.
Ohno S. 1970. Evolution by gene duplication. Berlin, New York: Berlin, New York, Springer-
Verlag
Otto SP, Whitton J. 2000. Polyploid incidence and evolution. *Annu. Rev. Genet.* 34:401–437.
Parey E, Louis A, Cabau C, Guiguen Y, Roest Crollius H, Berthelot C. 2020. Synteny-Guided
Resolution of Gene Trees Clarifies the Functional Impact of Whole-Genome
Duplications. *Mol Biol Evol* 37:3324–3337.
Parey E, Louis A, Montfort J, Guiguen Y, Crollius HR, Berthelot C. 2022. An atlas of fish
genome evolution reveals delayed rediploidization following the teleost whole-genome
duplication. *Genome Research* 32:1685–1697.
Parisod C, Holderegger R, Brochmann C. 2010. Evolutionary consequences of autopolyploidy.
New Phytologist 186:5–17.
Pedregosa F, Varoquaux G, Gramfort A, Michel V, Thirion B, Grisel O, Blondel M, Prettenhofer
P, Weiss R, Dubourg V. 2011. Scikit-learn: Machine learning in Python. *the Journal of*
machine Learning research 12:2825–2830.
Qiao X, Li Q, Yin H, Qi K, Li L, Wang R, Zhang S, Paterson AH. 2019. Gene duplication and
evolution in recurring polyploidization–diploidization cycles in plants. *Genome Biology*
20:38.
Ramsey J, Schemske DW. 2002. Neopolyploidy in flowering plants. *Annu. Rev. Ecol. Syst.*
33:589–639.
Ren R, Wang H, Guo C, Zhang N, Zeng L, Chen Y, Ma H, Qi J. 2018. Widespread whole
genome duplications contribute to genome complexity and species diversity in
angiosperms. *Molecular Plant* 11:414–428.
Robertson FM, Gundappa MK, Grammes F, Hvidsten TR, Redmond AK, Lien S, Martin SAM,
Holland PWH, Sandve SR, Macqueen DJ. 2017. Lineage-specific rediploidization is a
mechanism to explain time-lags between genome duplication and evolutionary
diversification. *Genome Biology* 18:111.
Rothfels CJ, Johnson AK, Hovenkamp PH, Swofford DL, Roskam HC, Fraser-Jenkins CR,
Windham MD, Pryer KM. 2015. Natural hybridization between genera that diverged from
each other approximately 60 million years ago. *Am Nat* 185:433–442.
Roux C, Pannell JR. 2015. Inferring the mode of origin of polyploid species from next-generation
sequence data. *Mol Ecol* 24:1047–1059.
Roux J, Liu J, Robinson-Rechavi M. 2017. Selective constraints on coding sequences of
nervous system genes are a major determinant of duplicate gene retention in
vertebrates. *Molecular biology and evolution* 34:2773–2791.
Salojarvi J, Rambani A, Yu Z, Guyot R, Strickler S, Lepelley M, Wang C, Rajaraman S, Rastas
P, Zheng C. 2023. The genome and population genomics of allopolyploid *Coffea arabica*
reveal the diversification history of modern coffee cultivars. *bioRxiv*:2023.09. 06.556570.
Salojärvi J, Rambani A, Yu Z, Guyot R, Strickler S, Lepelley M, Wang C, Rajaraman S, Rastas
P, Zheng C. 2024. The genome and population genomics of allopolyploid Coffea arabica
reveal the diversification history of modern coffee cultivars. *Nature genetics* 56:721–731.
Schlueter JA, Dixon P, Granger C, Grant D, Clark L, Doyle JJ, Shoemaker RC. 2004. Mining
EST databases to resolve evolutionary events in major crop species. *Genome* 47:868–
876.
Schranz E, Mohammadin S, Edger PP. 2012. Ancient whole genome duplications, novelty and
diversification: the WGD Radiation Lag-Time Model. *Current Opinion in Plant Biology*
15:147–153.
Senchina DS, Alvarez I, Cronn RC, Liu B, Rong J, Noyes RD, Paterson AH, Wing RA, Wilkins
TA, Wendel JF. 2003. Rate variation among nuclear genes and the age of polyploidy in
Gossypium. *Mol Biol Evol* 20:633–643.
Sensalari C, Maere S, Lohaus R. 2022. ksrates: positioning whole-genome duplications relative

832   to speciation events in KS distributions. *Bioinformatics* 38:530–532.

833 Soltis DE, Buggs RJA, Doyle JJ, Soltis PS. 2010. What we still don't know about polyploidy.
834   *TAXON* 59:1387–1403.

835 Soltis DE, Soltis PS, Schemske DW, Hancock JF, Thompson JN, Husband BC, Judd WS. 2007.
836   Autopolyploidy in angiosperms: have we grossly underestimated the number of species?
837   *Taxon* 56:13–30.

838 Soltis DE, Visger CJ, Marchant DB, Soltis PS. 2016. Polyploidy: pitfalls and paths to a
839   paradigm. *American Journal of Botany* 103:1146–1166.

840 Soltis PS, Liu X, Marchant DB, Visger CJ, Soltis DE. 2014. Polyploidy and novelty: Gottlieb's
841   legacy. *Philos Trans R Soc Lond B Biol Sci* 369:20130351.

842 Soltis PS, Soltis DE. 2012. Polyploidy and genome evolution. Springer

843 Spoelhof JP, Soltis PS, Soltis DE. 2017. Pure polyploidy: closing the gaps in autopolyploid
844   research. *Journal of Systematics and Evolution* 55:340–352.

845 St Onge KR, Foxe JP, Li J, Li H, Holm K, Corcoran P, Slotte T, Lascoux M, Wright SI. 2012.
846   Coalescent-based analysis distinguishes between allo- and autopolyploid origin in
847   Shepherd's Purse (*Capsella bursa-pastoris*). *Mol Biol Evol* 29:1721–1733.

848 Stebbins CL. 1950. Variation and evolution in plants. *Variation and evolution in plants.*

849 Stebbins GL. 1947. Types of polyploids: their classification and significance. In: Advances in
850   genetics. Vol. 1. Elsevier. p. 403–429.

851 Stebbins GL. 1951. Cataclysmic Evolution. *Scientific American* 184:54–59.

852 Sutherland BL, Tiley GP, Li Z, McKibben MT, Barker MS. 2024. SLEDGe: Inference of ancient
853   whole genome duplications using machine learning. *bioRxiv*:2024.01.17.574559.

854 Tank DC, Eastman JM, Pennell MW, Soltis PS, Soltis DE, Hinchliff CE, Brown JW, Sessa EB,
855   Harmon LJ. 2015. Nested radiations and the pulse of angiosperm diversification:
856   increased diversification rates often follow whole genome duplications. *New Phytologist*
857   207:454–467.

858 Thomas GWC, Ather SH, Hahn MW. 2017. Gene-tree reconciliation with MUL-trees to resolve
859   polyploidy events. *Systematic Biology* 66:1007–1018.

860 Tiley GP, Barker MS, Burleigh JG. 2018. Assessing the performance of Ks plots for detecting
861   ancient whole genome duplications. *Genome Biology and Evolution* 10:2882–2898.

862 Van de Peer Y. 2004. Computational approaches to unveiling ancient genome duplications.
863   *Nature Reviews Genetics* 5:752–763.

864 Van de Peer Y, Ashman T-L, Soltis PS, Soltis DE. 2020. Polyploidy: an evolutionary and
865   ecological force in stressful times. *The Plant Cell* [Internet]. Available from:
866   https://doi.org/10.1093/plcell/koaa015

867 Vandepoele K, Saeys Y, Simillion C, Raes J, Van De Peer Y. 2002. The automatic detection of
868   homologous regions (ADHoRe) and its application to microcolinearity between
869   Arabidopsis and rice. *Genome Res* 12:1792–1801.

870 Vanneste K, Van de Peer Y, Maere S. 2013. Inference of genome duplications from age
871   distributions revisited. *Mol Biol Evol* 30:177–190.

872 Wang J, Qin J, Sun P, Ma X, Yu J, Li Y, Sun S, Lei T, Meng F, Wei C. 2019. Polyploidy index
873   and its implications for the evolution of polyploids. *Frontiers in genetics* 10:807.

874 Wang X, Shi X, Li Z, Zhu Q, Kong L, Tang W, Ge S, Luo J. 2006. Statistical inference of
875   chromosomal homology based on gene colinearity and applications to Arabidopsis and
876   rice. *BMC Bioinformatics* 7:447.

877 Wen D, Yu Y, Zhu J, Nakhleh L. 2018. Inferring phylogenetic networks using phyloNet.
878   *Systematic Biology* 67:735–740.

879 Wendel JF. 2015. The wondrous cycles of polyploidy in plants. *Am J Bot* 102:1753–1756.

880 Wendel JF, Jackson SA, Meyers BC, Wing RA. 2016. Evolution of plant genome architecture.
881   *Genome Biology* 17:37.

882 Wolfe KH, Li WH, Sharp PM. 1987. Rates of nucleotide substitution vary greatly among plant

mitochondrial, chloroplast, and nuclear DNAs. *Proc Natl Acad Sci U S A* 84:9054–9058.

Yan Z, Cao Z, Liu Y, Ogilvie HA, Nakhleh L. 2022. Maximum parsimony inference of
phylogenetic networks in the presence of polyploid complexes. *Systematic Biology*
71:706–720.

Yang N, Wang Y, Liu X, Jin M, Vallebueno-Estrada M, Calfee E, Chen L, Dilkes BP, Gui S, Fan
X, et al. 2023. Two teosintes made modern maize. *Science* 382:eadg8940.

Yang Z. 2007. PAML 4: phylogenetic analysis by maximum likelihood. *Molecular Biology and*
Evolution 24:1586–1591.

Yang Z, Nielsen R. 1998. Synonymous and nonsynonymous rate variation in nuclear genes of
mammals. *Journal of Molecular Evolution* 46:409–418.

Yang Z, Nielsen R. 2000. Estimating synonymous and nonsynonymous substitution rates under
realistic evolutionary models. *Mol Biol Evol* 17:32–43.

Yant L, Bomblies K. 2015. Genome management and mismanagement--cell-level opportunities
and challenges of whole-genome duplication. *Genes Dev* 29:2405–2419.

Yu Q, Guyot R, de Kochko A, Byers A, Navajas-Pérez R, Langston BJ, Dubreuil-Tranchant C,
Paterson AH, Poncet V, Nagai C, et al. 2011. Micro-collinearity and genome evolution in
the vicinity of an ethylene receptor gene of cultivated diploid and allotetraploid coffee
species (Coffea). *The Plant Journal* 67:305–317.

Zhang H, Gou X, Zhang A, Wang X, Zhao N, Dong Y, Li L, Liu B. 2016. Transcriptome shock
invokes disruption of parental expression-conserved genes in tetraploid wheat. *Scientific*
Reports 6:26363.

Zhang L, Wu S, Chang X, Wang X, Zhao Y, Xia Y, Trigiano RN, Jiao Y, Chen F. 2020. The
ancient wave of polyploidization events in flowering plants and their facilitated adaptation
to environmental stress. *Plant, Cell & Environment* 43:2847–2856.

Zhang S, Li R, Zhang L, Chen S, Xie M, Yang L, Xia Y, Foyer CH, Zhao Z, Lam H-M. 2020.
New insights into *Arabidopsis* transcriptome complexity revealed by direct sequencing of
native RNAs. *Nucleic Acids Res* 48:7700–7711.

Zhao J, Teufel AI, Liberles DA, Liu L. 2015. A generalized birth and death process for modeling
the fates of gene duplication. *BMC Evol Biol* 15:275.

Zwaenepoel A, Van de Peer Y. 2019. WGD—simple command line tools for the analysis of
ancient whole-genome duplications. *Bioinformatics* 35:2153–2155.

Figures and Tables

Figure 1. Relationship between the shape of the *Ks* histogram and the modes of auto versus allopolyploid speciation.
Top: Autopolyploid speciation. Bottom: Allopolyploid speciation. Outgroup (O) and parental species are indicated in light gray. Polyploid species are shown in dark gray with duplicated genomes denoted as P1 and P2. The arrow in the *Ks* histogram points backwards in time. The set of gene tree coalescents $\{T_{ci}\}$ are shown as nodes on the gene trees, and their distribution in time (D_{TCOAL}) is indicated by the gray shaded bands. For allopolyploids the offset between T_{WGD} and T_{DIV} represents the lag time between separation of the diploid parental species (T_{DIV}) and their later conjunction by polyploidization (T_{WGD}). For autopolyploids, there may be instances where the gene tree divergence may begin before T_{WGD} or after it. In both cases, T_{WGD} and T_{DIV} are not synonymous, and D_{TCOAL} may be complex. Figures at the far right and left are derived from (Chen and Zwaenepoel, 2023).

Paralogous genes modeled by SpecKS

Ohnologs

Small-scale duplications

Allopolyploid Autopolyploid Allopolyploid Autopolyploid

Figure 2. Ohnologs (paralogs generated by WGD) and small-scale duplications ("SSDs", paralogs generated as single or small-scale copies) have unique contributions to the *Ks* histogram. Ohnologs (left panel) are shown for both allo and autopolyploids. Green and pink ohnologs (far left) are attributed to allopolyploidy and are born as orthologs (genes duplicated by speciation, dashed lines). Gold and blue ohnologs (center left) are attributed to autopolyploidy and are born by WGD. SSDs (right panel) follow the same birth and death process for both allo and autopolyploids and are shown here in green and blue. Unless maintained by selection, SSDs tend to be shed rapidly, and are thus found at the far left of the *Ks* histograms. As in Fig. 1, the polyploid species is shown in dark gray while the parental and outgroups are shown in light gray. The arrow of time points into the past along the X axis of the *Ks* histogram.

Figure 3. The SpecKS 2-dimensional model parameterizing allopolyploid speciation.
A graphical representation of the 2D polyploid continuum modeled by SpecKS, showing Π,
ΔT, T_{DIV}, T_{WGD}, and the set of gene-tree coalescent times $\{T_{COAL\_i}\}$ in the SpecKS model.

Figure 4. The SpecKS pipeline.
A diagrammatic representation of the eight modules that make up the SpecKS polyploidy simulation pipeline.

Figure 5. SpecKS demonstrates how the shapes of the *Ks* distributions attenuate over time.
Set A: low Ne (1x10^5) ancestral species and a large ΔT (5 MY)
Set B: high Ne (5x10^6) ancestral species and a large ΔT (5 MY)
Set C: low Ne (1x10^5) ancestral species and a small ΔT (0 MY)
Set D: high Ne (5x10^6) ancestral species and a small ΔT (0 MY)

Figure 6. SpecKS demonstrates how the shapes of the *Ks* distribution vary with respect to ancestral diversity and ΔT. Top: Ancestral N_e varies from 10^5 to 20x10^6. T_{WGD} is fixed at 5 MYA and T_{DIV} is fixed at 10 MYA. Bottom: T_{WGD} varies from 0 to 40 MY, T_{DIV} is fixed at 40 MYA, and N_e is fixed at 1*10^6.

Figure 7. SpecKS shows that estimation of WGD time based number of genes remaining yields accurate results, irrespective of ΔT. (Left) T_{WGD} as estimated from the number of genes shed. (Top right) Error in the T_{WGD} as predicted from the number of genes shed (y axis) vs the degree of allopolyploidy (x-axis). (Bottom right) Error in T_{WGD} as estimated from the histogram mode (y axis) vs the degree of allopolyploidy (x-axis). Each datapoint represents results from simulations from dataset 1, described previously. Within dataset1, T_{DIV} times range from 10 to 80 MYA, by 10, with WGD offset (ΔT) by 0, 5, 10 and 50 MY. N_e*G_t is 0.1,1.0,5.0, 10 and 20 MY.

Figure 8. Using SpecKS to reproduce *Ks* empirical histograms from transcriptomic data.
SpecKS-simulated histogram in blue and the empirical *Ks* histogram from transcriptomic data in green.

Figure 9. The low, medium and high ΔT discrimination model.
(left) The ΔT discrimination threshold applied to the estimated ΔT. (right) Accuracy of inference of estimation ΔT, represented as an ROC plot. Each datapoint represents results from simulations from dataset 1, described previously. Lighter colors denote higher ΔT.

Figure 10. The low-vs-high Ne discrimination model.
(left) The low-vs-high diversity discrimination threshold applied to simulated polyploids. The x-axis gives the SPC time. The y-axis gives the value of the high-vs-low diversity discrimination metric. (right) The ROC plot to assess accuracy of inference into high versus low Ne, across threshold values. Darker colors denote higher N_e. Each datapoint represents results from a simulation from dataset 1, described previously. (Table 3).

1011

1012
1013
1014
1015

Figure 11. The Ne-classifier results for the 1KP dataset
(a) Violin plots indicating the number of species in each category. (b) Histogram giving the number of species in each category. Darker blue color represents higher-N_e ancestral species at SPC time, respectively. X-axis gives the numerical value of the discrimination metric. Y-axis gives the number of species per bin.

1016

Table 1. Subset of SpecKS configurable parameters		
Polyploid-specific. (Set for each simulated polyploid in the run):		
Parameter & default	**Default**	**Description**
DIV_time_MYA	0 MY	DIV time in MY. Time of subgenome divergence. For gradual speciation, this will be the mode of the gene tree divergence times.
WGD_time_MYA	0 MY	WGD time in MY. This will be the start time of ohnolog shedding, which will continue until the present time.
Gene_div_time_ distribution_ parameters	impulse,1,1	Sets the distribution of divergence times for gene trees at DIV time. For **autopolyploids**, use format: "impulse,1,1". For **allopolyploids**, use the format "expon,0,K", where K is the exponential decay constant. (We suggest K=Ne*Gt). For polyploids whose gene tree divergence might be a mix of distributions (ie, **segmental allopolyploids**), multiple distributions may be given, with the last parameter being the proportion of genes which belong in each distribution. Details in Table S2.
General. (The same for all polyploids in the run):		
Full_sim_time	100 MY	The length of the time period to simulate. Note that since speciation is a gradual process, it may be necessary to start the simulation well in advance of the SPC time, in order to fully capture the distribution of gene trees.
mean_gene_birth_ rate_GpMY	0.001359 genes per MY	Gene birth rate. Note this may be lineage specific. Our default is chosen from [Guo 2013]
SSD_half_life_MY	4 MY	Half-life of small-scale duplications. Our default is chosen from [Lynch and Conery 2003]
WGD_half_life_M Y	31 MY	Half-life of ohnologs (whole-genome duplications). Our default is based on Guo 2013 and Maere 2005
Num_gene_trees_ per_species_tree	3000	Default is set to give a well-supported histogram without taking too long to run. If set too low, the final histogram will look too sparse. Higher numbers may be more realistic [Sterk 2007] and result in smoother histograms but have a longer run time. Since gene trees are simulated independently, the number does not affect the

		general histogram shape, merely the number of samples in it, so high numbers are not always necessary.
Num_replicates_per_gene_tree	1	SpecKS can automatically run replicates for a given simulation, randomizing appropriately.
Num_codons	1000	The number of codons in each gene to be simulated. All genes in the sim have the same length. 1000 was chosen in agreement with [Tiley 2018].
Ks_per_Myr	0.01	*Ks* per million years. This number is lineage and gene family specific, and may need to change depending on user needs [Gaut 1996 and Koch 2000]. The default of 0.01 was chosen in agreement with [Tiley 2018], and in range with [Blanc and Wolfe 2004]
Per_site_evolutionary_distance Default: 0.01268182	0.01268182	The per-site-evoultionary_distance is used to calculate the total tree length per gene tree input to evolver. The default setting was derived by [Tiley 2018] for a Ks_per_Myr of 0.01 and the evolutionary GY94 model.

1018
1019

1020
1021
1022

Table 2. Summary of 1KP categorization results			
Ne categorization	Number in category	Percent of total	Metric mean value
Low	12	5.26	-5.05
Medium	38	16.67	-3.69
High	178	78.07	-2.45
Total	228	100	-2.80

1023
1024
1025

Table 3. SpecKS parameters used in dataset 1		
Polyploid	DIV_time_MYA	[80,70, 60, 50, 40, 30, 20,10]
Polyploid	WGD_time_MYA	[0,5,10,20,50]
Polyploid	gene_div_time_distribution_parameters	"Impulse,1,1" and "expo,0,K" Where K varied from [0.01,0.1,1.0,5.0,10.0,20.0] To span [Gossmann 2010]
Species Tree	full_sim_time	100 MY
Gene Tree	mean_gene_birth_rate_GpMY	0.001359 genes per MY
Gene Tree	SSD_half_life_MY	4 MY
Gene Tree	WGD_half_life_MY	31 MY
Gene Tree	num_gene_trees_per_species_tree	3000
Sequence Evolution	num_replicates_per_gene_tree	1
Sequence Evolution	num_codons	1000
Sequence Evolution	Ks_per_Myr	0.01
Sequence Evolution	per_site_evolutionary_distance	0.01268182
All other parameters are default.		