Dispersal, maturation and recruitment in a long-lived, trans-continentally migrating bird

Paweł Mirski[1], Wouter Vansteelant[2,3], Patrik Byholm[4,5]

[1] Faculty of Biology, University of Białystok, Ciołkowskiego 1J, 15-245 Białystok, Poland

[2] BirdEyes, Centre for Global Ecological Change at the Faculties of Science and Engineering and Campus Fryslân, University of Groningen, Zaailand 110, 8911 BN Leeuwarden, The Netherlands

[3] Theoretical and Computational Ecology, Institute for Biodiversity and Ecosystem Dynamics, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, The Netherlands

[4] Organismal and Evolutionary Biology, University of Helsinki, PO Box 65, 00014 Helsinki, Finland

[5] Novia University of Applied Sciences, Raseborgsvägen 9, FI-10600 Ekenäs, Finland

HIGHLIGHTS

- We used a unique GPS telemetry dataset on honey buzzards to track their natal dispersal and recruitment in this long-lived avian migrant.
- All individuals gradually decreased their ranging areas, moved closer to natal sites, and increased the time spent on the breeding grounds to synchronize with adults and begin recruitment.
- Mortality during return migration was high and acted as a bottleneck.
- We found both processes to be complex and lengthy, with the overall duration depending on individual decisions regarding the timing of return migration.

ABSTRACT

Natal dispersal is a multi-step process. It commences when a juvenile departs from its natal site and

concludes when it settles to breed for the first time. During this interim period, dispersing individuals

of long-lived species undergo a wandering phase, which may span years. This phase remains one of

the least studied aspects of species' life history. We utilized a unique GPS-telemetry dataset on a

slow-life history migratory bird of prey – the European honey buzzard, tracked for multiple years

after fledgling. Our aim was to assess how phenology, ranging behaviour and philopatry change as

individuals gain experience. Individuals exhibited variability in the age at which they first returned to

their breeding range, yet all survivors settled at sites proximate to their natal nests. In subsequent

returns, immatures increased the time spent within the breeding range and narrowed down their

ranging to smaller areas situated even closer to their natal nests. Our study unveils the complexity

and protraction inherent in natal dispersal, emphasizing the importance of individual experience and

migratory improvement for successful recruitment in long-lived species.

Keywords: floaters, homing, philopatry, immaturity, European honey buzzard, *Pernis apivorus*

INTRODUCTION

Many animals depart from their natal habitat upon attaining independence. The

phenomenon of permanent displacement from the natal area to the site of initial reproduction has

garnered significant attention among ecologists and is extensively deliberated under the term "natal

dispersal" (Greenwood and Harvey 1982). Natal dispersal strategy tend to vary greatly between and

within species, and seem to emerge through a trade-off between establishing in familiar areas and

habitats and conspecific attraction on the one hand, and avoiding intraspecies competition (Nilson

1989, Negro et al. 1997, Forero et al. 2002) and inbreeding (Alonso et al. 1998, Zedroesser et al.

2007, Ortego et al. 2008) on the other hand. Theprocess and outcome of natal dispersal not only

influences the life history and fitness of individual organisms (Forero et al. 2002, Nevoux et al. 2013,

Luna et al. 2020) but also shapes the genetic and social structure of populations (Ekman et al. 2001,

Ortego et al. 2008, 2011).

Dispersal constitutes a movement process, yet in many investigations, it is often distilled to a

single metric: dispersal distance (Benton and Bowler 2005). Undoubtedly, it is a much more intricate

phenomenon and challenging to track, especially among long-lived species that mature late. In such

species, dispersal unfolds as a multistep process. Between the initial departure from the natal site

and settling in a new breeding ground for the first time, dispersing individuals undergo a wandering

phase (Benton and Bowler 2005, Penteriani and Delgado 2009). During this stage, long-lived species,

even upon reaching maturity, frequently linger for one or a few seasons, or even their entire lives as

floaters, seeking a mate or suitable breeding territory (Penteriani and Delgado 2009), and this part of

life is attributed to recruitment process. Recruitment can be defined as the establishment of new

individuals in the breeding segment of a population (Reed et al. 2003), and the recruitment process is

considered complete once an individual breeds for the first time. The age of first breeding is a key

aspect of recruitment that significantly affects population demography (Gaillard et al. 1989, Stahl and

Oli 2006). Along with clutch size and adult survival, it is one of the main components of birds' life

history strategies (Sæther and Bakke 2000). Species that mature early typically exhibit high fertility

and low survival, characteristics of fast-life history species, while late maturation, low fertility, and

high survival represent the other end of the life history continuum (Sæther 1988). However, the age

of first breeding is also density-dependent, with younger individuals reproducing in low-density

populations (Wyllie and Newton 1991), thus indicating the stage of population development.

In migratory species, wherein the period between leaving the natal site and initial breeding is

preceded by one or more episodes of migration, dispersal is additionally complex. During this interim

period, immature individuals may traverse thousands of kilometres to non-breeding grounds before

returning to their breeding range. Inexperienced young bird migrants are often susceptible to

deviations during migration due to navigational errors or wind drifts, resulting in their positions often

shifting further and in a more random manner than those of resident species (Vickers et al. 2023).

Consequently, migration profoundly influences the dispersal process, resulting in relatively longer

dispersal distances (Paradis 1998) and relatively lower levels of philopatry (likelihood that individuals

breed at or near their place of origin; Weatherhead and Forbes 1994) in species that migrate longer

distances. At the same time, high philopatry (i.e. dispersal distances in the order of tens of meters to

kilometers) tends to be widespread even among migrants (Förschler and Bairlein 2010, Ceresa et al.

2016) and even in species that spend long periods outside the breeding range before their first

breeding attempt (Coulson 2016). In doing so, migratory birds, demonstrate remarkably long-term

spatial memory and homing abilities, as well as a strong bias towards natal environmental cues

during recruitment (Keiser 2003, Healy and Hurly 2004). And so the precise mechanisms through

which early-life movements outside the breeding range influence the natal dispersal process remain

obscure.

An additional challenge for migrants, especially those undertaking long-distance journeys, is

that they spend relatively little time in their breeding range prior to recruitment. While immature

individuals of resident species may spend the entire autumn, winter, and early spring exploring

resources and assessing competitors' distribution around potential breeding sites, migrants often

have only a few weeks from arrival to recruit into the breeding population. In long-lived species with

protracted offspring development and parental care, floaters, despite being physiologically mature,

may struggle to secure a territory and breed within the short window following their initial arrival

from wintering sites. In such species, the dispersal process is also protracted and typically

encompasses the floater stage for at least one season. Importantly, individuals that do not improve

their migration performance and arrival time might not even recruit into the breeding population

(Sergio et al. 2017). All considered, this underscores the importance of better studying the link

between migration timing and the natal dispersal and recruitment process.

The European honey buzzard (*Pernis apivorus*; therafter honey buzzard) serves as an example

of a long-lived, long-distance migratory species with potentially intricate dispersal and recruitment

strategies. Firstly, it is a specialist with a narrow diet, predominantly preying on Hymenoptera,

especially wasps and their larvae (Gamauf 1999, van Manen et al. 2011, van Manen 2013) within

tropical and temperate forests (van Manen et al. 2011, Howes et al. 2020a). This dietary

specialization significantly influences its annual cycle and appears to constrain the suitable habitats

where it can establish breeding populations. Secondly, migratory species of the honey buzzard genus

undertake transcontinental migrations, covering several thousand kilometres between their

temperate breeding and tropical non-breeding ranges (Higuchi et al. 2005, Strandberg et al. 2012),

with juveniles and adults exhibiting a large ontogenetic shift in migratory behaviour (Kjellen et al

1992, Schmid 2000, Hake et al 2003, Vansteelant et al 2020). Thirdly, it is a secretive species,

characterized by its brief residency in breeding areas, late spring arrival (during the full canopy

season), and relatively low population density. Consequently, it has been inadequately studied,

particularly in contrast to its relatively high conservation status (listed in Annex 1 of the Birds

Directive).

Here, we present longitudinal movement data of young honey buzzards tracked from their

natal areas in Finland for up to seven years after fledging, thus revealing their process of natal

dispersal and recruitment (cf. Roberts and Law 2014). We specifically aimed to understand the

interplay between maturity, natal dispersal, recruitment, and migratory behavior in this complex and

secretive species. It is well established from migration counts, ringing and tracking data (reviewed in

Vansteelant and Agostini 2021) that juvenile honey buzzards on their first outbound migration

migrate later and over a broader front than adults, with relatively few juveniles migrating through

traditional overland flyways (Kjellen et al 1992, Vansteelant et al 2020), and juveniles taking more

time to complete their first migration (Hake et al. 2003). Juvenile behaviour thus seems to be

consistent with an innate vector-based orientation strategy, making them highly susceptible to wind

drift (Thorup et al. 2003). Indeed, a previous study based on the same individuals that we study here

showed that stochastic wind influences greatly influenced the route choice of juveniles on their first

outbound migration, as well as the longitude at which they settled in tropical Africa (Vansteelant et

al. 2017). The remainder of the natal dispersal process, in particular the first return migration and

breeding range movements of young honey buzzards, have not been tracked until now.   We expect

most honey buzzards to initiate their first return migration upon reaching sexual maturity in their 3rd

calendar year, and to home back near to their natal areas to recruit as breeders. We also expect to

observe gradual changes in phenology and ranging behavior as the immatures mature, eventually

resembling the patterns observed in breeding adults once individuals recruit.

METHODS

GPS-tracking dataset

As a part of ongoing work on honey buzzard migration and movement ecology (Vansteelant et al.

2017, Nourani et al. 2020, Howes et al. 2020, Brønnvik et al. 2022), young honey buzzards (n = 29)

hatched in broods in southern Finland (latitude 61°14⃝–63°12⃝ N, longitude 21°16⃝–23°31⃝ E) were

equipped with solar-powered Argos-GPS platform terminal transmitters (PTTs) (Microwave

Telemetry Inc.) or Ecotone GSM-GPS-trackers (Ecotone) during 2011-2016 using body-loop harnesses

made of Teflon ribbon. Tags' weight (22–27 g) corresponded to ca. 3% of the birds' body mass at

deployment (908±83 g; avg ± s.d., n = 29). The amount of delivered GPS-fixes varied depending on

tracker model and programming. On average, immature individuals collected 1180 (± 764 SD) GPS

locations per individual per year. The sex of birds was determined from DNA as extracted from blood

samples using the salt extraction method. For more details on the field protocol see Vansteelant et

al. (2017). Additionally, a dataset on adult birds (n=9) was used for comparison of ranging areas on

wintering grounds. The details of this dataset are described in Brønnvik et al.(2022).

Ranging area

During consecutive wintering and breeding seasons, ranging areas were estimated to

compare how the area of exploration changes with age and experience. The dataset comprised

moderate-frequency GPS fixes, resulting in hundreds or occasionally dozens of fixes per season for

some individuals. To compare ranging areas rather than estimate space use requirements, a

geometric estimator – the 90% minimum convex polygon (mcp) – was employed. We considered

alternative utilization distribution methods, but in case on non-territorial, wandering individuals of

varying sample size and changing activity centres, utilization distribution methods overestimated the

ranging area more than mcp.

Calculations were conducted using the *adehabitatHR* package (Calenge 2006) in R 4.1.1.

Wintering ranging areas were calculated from the beginning of January until the end of March, when

individuals had reached their wintering quaters and ceased moving long distances, until spring

migration. Ranging areas upon returning to the breeding range were calculated from arrival until

departure in the potential breeding range. By the latter we assumed southern Finland, where the

tracked individuals originated and were expected to return, assuming the species is philopatric.

Phenological dates

To track how phenology of migration changed with age, departure from non-breeding range,

arrival to and departure from the breeding range were noted in the Julian calendar. Because

individuals differed in the latitude of their ranging in Africa and this affects the distance and

migration duration, the date of departure from non-breeding range was recorded as the date of

passing the 13[th] parallel (the approximate boundary of the desert zone). Arrival to the breeding range

was determined as the first instance where an individual ceased moving northward by more than 50

km in the next 5 days. Departure from the breeding range was identified when an individual began

moving southward by more than 50 km in the next 5 days.

Philopatry

To assess if immature honey buzzards tend to return and potentially settle close to their

natal sites, we calculated the mean and minimum distance at which honey buzzards resided from

their natal site in each consecutive breeding period. To do this we used the 'distance matrix' tool in

QGIS 3.22 for each breeding season.

Mortality

Confirmed and probable deaths of tracked individuals, especially during their first autumn

migration, wintering, and returning migration, were recorded to compare mortality among naïve

honey buzzards. Suspected mortality was investigated on-site when indicated by sensors showing

low temperatures, absence of movement over several days, or suspicious movements in urbanized

habitats. Mortality due to drowning during migration across the Mediterranean and Black Sea was

inferred from signal loss at the sea shore. Tag failure was suspected in some cases, especially in the

fourth-seventh year of tracking. Mortality ratios were estimated based on confirmed cases, excluding

suspected tag failures.

Analyses

Maps were done in QGIS 3.22 (QGIS.org 2024). All the analyses were carried out in R 4.1.1 (R

Core Team 2021). Due to sample size limitations, statistical approaches were constrained. Out of 29

individuals at the start, the sample size decreased to eleven individuals completing the first winter,

six at the first successful return to the breeding range, four at the second, and three at the third.

Wilcoxon tests were used to compare winter ranging areas between first and subsequent wintering

episodes as well as to test sex differences in the year of first return to breeding range and date of

arrival. Linear models were employed to test changes in dates of arrival and departure from breeding

ranges (Julian date), duration of stay and size of ranging areas in breeding and wintering grounds

(square km). Each of the above variables was tested in a separate model due to unequal and low

sample sizes and explained by individual experience, measured by the number of returns to the

breeding range or the number of wintering episodes in the case of winter ranging areas. Model

residuals were assessed for normality with the Shapiro test and variables were log-transformed in

the case of ranging areas.

RESULTS

Extended Immaturity Period in Wintering Range

Out of the 29 tracked juveniles, 22 (76%) successfully reached wintering sites in Africa (Tab.

1). None of the juveniles returned to the breeding range in their second calendar year. During their

first winter, honey buzzards ranged over extensive areas in Central-Western Africa, with some

individuals traversing distances exceeding 2000 km along longitudinal or latitudinal axes (Fig. 1A).

Subsequently, some individuals reduced their ranging areas considerably during the next winter,

while others maintained extensive ranges. However, the decrease in wintering area between the first

and next winter was statistically insignificant (W = 57.5, p = 0.11), and no significant changes were

observed in later life stages, although breeding adults tended to range over smaller areas (Fig. S1).

In the 3rd calendar year, six out of eleven tracked individuals attempted to return to the breeding

range, with only three succeeding. The individuals that skipped the returning journey all survived to

the next calendar year, and most of those (four out of five) attempted to return to the breeding

grounds in the following year, in their 4th calendar year. The last remaining individual, tried to return

in his 5th calendar year, but did not succeed in crossing the Mediterranean Sea and contined to

wander in non-breeding part of the range (Fig. 2). Finally, this male successfully returned to his natal

area in his 6th calendar year. In total, only six individuals (21%) reached the breeding range between

their 3rd and 6th calendar years. We did not confirm that establishing winter territories (represented

by the size of winter ranging area) impacted the age of returning migration (W = 47, p = 0.76).

Similarly, the change in winter ranging area between the first and second winters did not affect the

decision to undertake the returning migration in the 3rd calendar year (W = 10, p = 0.22).

Table 1. Characteristics of GPS-tracking dataset of twenty nine Honey Buzzards from Finland,
followed from fledging.

Name	Sex	Tracking start	Tracking end	Calendar year of return	Last data	Mortality	Mortality cause	Last signal site
Hans	M	03.08.2012	22.03.2014	-	1st wintering	Death	?	Cameroon
Ella	F	13.08.2013	16.12.2013	-	1st wintering	Death	?	Congo
Emma	F	06.08.2013	14.10.2013	-	1st autumn	Death	Shot	Sicily
Gilda	F	09.08.2012	08.11.2012	-	1st autumn	Death	Shot^	Niger
Ivar	M	07.08.2013	20.09.2013	-	1st autumn	Death	Predated	Estonia
Heidi	F	09.08.2013	28.11.2014	-	2nd wintering	Death	?	Gabon
Julia	F	17.08.2013	13.01.2014	-	1st wintering	Death	?	Congo
Kirsi	F	11.08.2012	13.11.2012	-	1st wintering	Death	Shot^	Nigeria
Lisa	F	03.08.2013	19.10.2013	-	1st autumn	Death	Predated	Deserts (Algeria)
Miikka	M	09.08.2013	04.04.2014	-	1st wintering	Tag failure	?	Nigeria
Per	M	02.09.2016	24.10.2016	-	1st autumn	Death	Exhaustion	Deserts (Algeria)
Piff	M	06.08.2011	17.02.2012	-	1st wintering	Death	?	Gabon
Roosa	F	29.08.2011	06.02.2012	-	1st wintering	Death	?	Congo
Rudolf	M	03.08.2012	02.02.2013	-	1st wintering	Death	?	Angola
Sven	M	07.08.2013	06.10.2013	-	1st autumn	Death	?	Algeria
Tor	M	29.07.2012	29.12.2012	-	1st wintering	Death	?	Angola
Ulla	F	06.08.2013	09.12.2013	-	1st wintering	Death	Collission	Congo
Viljo	M	03.08.2014	26.10.2014	-	1st autumn	Death	?	Deserts (Burkina Faso)
Venus	F	09.08.2012	11.05.2015	4*	1st return	Death	Exhaustion	Atlas mountains
Anni	F	06.08.2013	22.07.2016	4*	1st return	Unknown	?	Turkey
Edit	F	03.08.2013	18.06.2015	3*	1st return	Death	Drowned?	Medditeranean Sea
Matti	M	10.08.2013	29.05.2015	3*	1st return	Death	Predation?	Atlas mountains
Puff	F	07.08.2011	27.05.2013	3*	1st return	Death	Exhaustion?	Deserts (Mauritania)
Aida	F	03.08.2012	19.07.2014	3	breeding range	Death	Exhaustion^	Finland
Jaana	F	29.07.2011	20.05.2015	3	3rd return	Death	Exhaustion^	Deserts (Morocco)
Lars	M	10.08.2011	08.11.2015	3	5th wintering	Unknown	?	Cameroon
Mohammed	M	14.08.2013	25.09.2018	5* / 6	8th wintering	Tag failure?	?	Chad
Senta	F	09.08.2012	15.04.2018	4	3rd return	Unknown	?	Libya
Valentin	M	09.08.2012	09.10.2015	4	4th wintering	Tag failure	?	Nigeria

228   *- incomplete returning journey
229   ^- confirmed mortality
230

Figure 1. Differences in migration strategy and ranging behaviour with age (A) o or migratory experience (B-G) in GPS-tracked Honey Buzzards from Finland, followed from fledging. A - Size of ranging areas on wintering (January-March) and breeding (B) grounds, C - Date of departure from non-breeding range, D - Date of arrival to breeding range, E - Date of departure from breeding range (start of autumn migration), F - Duration of stay in the breeding range, G - Mean distance to natal nest during breeding season. Sample size decreased in time, from thirteen individuals followed through first winter, six at first return, four at second and three at third.

The First and Following Returns to the Breeding Range

All tracked individuals followed different paths during their first spring migration back to the

breeding range when compared to their first autumn migration, with most paths not intersecting

between autumn and spring (Fig. 2). Both clockwise and counterclockwise loop migrations occurred.

Honey buzzards advanced their departure from the non-breeding range with each following return

episode. The trend was visible (Fig. 1C), but only close to significant (Tab. 2). The date of arrival to the

breeding range varied significantly between individuals especially during the first returning migration

but tended to advance and narrow down slightly in subsequent returns (Fig. 1D, Tab. 2). The date of

departure from the breeding range tended to be delayed across returns (Fig. 1E, Tab. 2). The mean

departure date at first return was eight days earlier than the intial departure it their 1[st] calendar year

and about 2 weeks earlier at their second and third return. Immature honey buzzards spent more

time in the breeding range with each subsequent return (Fig. 1F, Tab. 2). At first return, it was 22.2

days on average (range: 5 – 54), at second – 51.0 days (31 – 79) and at third – 63.7 days (56 – 84).

Table 2. Results of linear models explaining change in migration strategy and ranging behaviour with
experience in GPS-tracked Honey Buzzards from Finland, followed from fledging (*p*-values are
rounded to two significant digits, significant values are bolded).

	Intercept	Beta coefficient	Standard error	*p*	Adjusted R^2
Departure from non-breeding range	143.05	-5.78	3.05	0.074	0.12
Arrival to breeding range	176.93	-7.4	6.76	0.30	0.02
Departure from breeding range	242.89	-1.3	2.12	0.55	-0.06
Duration of stay in the breeding range	32.18	16.51	6.83	**0.034**	0.29
Ranging area in breeding range (log)	13.15	-1.32	0.77	**0.0064**	0.46
Mean distance to natal site	376.23	-92.53	31.54	**0.014**	0.39

Figure 2. First autumn (continuous line) and first returning (dashed line) migrations of GPS-tracked
Honey Buzzards from Finland. The names of individuals provided in each map correspond to the
details given in Table 1. The year of the first autumn and returning migrations, respectively, are given
in brackets.

Recruitment and philopatry

The mean distance at which honey buzzards resided from their natal sites decreased

significantly with each return (see Table 2, Figure 1G), averaging 296 km at the first return, 162 km at

the second, and 135 km at the third (see Table S1). Similarly, the minimum distance to natal sites

decreased with each return, ranging from 5 to 177 km at the first return (mean 53 km) to 0 to 174 km

at the second return (mean 49 km). Five out of six returning individuals visited sites closer than 20

km from their natal sites, with four of them closer than 10 km. The area explored on the breeding

ground by individuals tracked, at least until the second return, showed a significant reduction (Fig.

1B, Fig. 3). Following the first return, honey buzzards explored hundreds of square kilometres,

whereas areas of only dozens of square kilometers were explored in the subsequent seasons. Some

individuals confined their ranging to a portion of the area explored during the first return, while

others moved to new, neighbouring areas that they had not explored before. Ranging areas showed

significant overlap between the second and third return.

Two males were tracked long enough to locate their first nests and observe a central place

foraging pattern around them (Tab. 1). However, we haven't confirmed successful breeding. Lars

built a nest in his 5th calendar year, during his third return to the breeding range, 16 km from the

natal site. Mohammed built his nest in his 6th calendar year, during his first return to the breeding

range, 189 km from the natal site. In his 7th calendar year, he refurbished the same nest, but in the

8th, he probably built another one. Mating and breeding, however, were not confirmed in either of

these males.

Figure 3. Change in summer ranging areas estimated with 90% minimum convex polygons of immature Honey Buzzards from Finland after first (white line), second (yellow line) and third return (red line). Plots A, B, C and D show ranging areas of Senta, Lars, Janna and Mohammed, respectively (see Tab. 1 for characteristics of individuals).

Mortality Across the Long Dispersal Period

Twenty three of twenty nine tracked individuals (79%) were found dead. In the remaining six individuals (21%), the signal was lost, usually after 4-8 years of tracking (see Tab. 1 for individuals' faith). In 4 cases (17%), we noted anthropogenic mortality, primarily due to shooting. Mortality on the first autumn migration reached 25% (7 out of 28 at this tage). All, except for one, managed to cross the Mediterranean Sea. Once they reached their wintering sites in the tropics, 32% of juveniles

perished at wintering sites during the first wintering season and one more individual (3.5%) in the

second season (altogether 10 out of 28). Finally, 39% of honey buzzards survived until their first

returning migration. Out of 11 individuals that survived long enough to undertake the returning

journey, as much as five (45%) perished trying to reach Europe. One individual turned back, so only

less than 50% of individuals that survived to this stage were able to reach the breeding range in the

first attempt and 55% in total. Once the honey buzzards successfully completed their dispersal cycle,

only two mortality cases were noted. Overall, only 22% of initially tracked individuals managed to

return to their breeding range, while only 15% lived up to their second return to the breeding range,

when potentially the first breeding attempt was possible.

DISCUSSION

Using GPS-GSM devices we were able to track the maturation and natal dispersal phase of

honey buzzards from fledging to recruitment. Overall we reveal a remarkably protracted natal

dispersal process, which sheds new light on the importance of individual experience and migratory

improvements for successful recruitment in this long-lived species.

Protracted immaturity period

Our study showed that the age of first breeding might vary between individuals, and that the

honey buzzard breeds at a later age as compared to similar-sized raptor species, including resident

species, and short-distance as well as long-distance migrants. While we could confirm nest-building

by two GPS-tracked Finnish males in the 5th and 6th calendar years, breeding attempts could not be

confirmed for other birds of similar age, and most individuals showed continuous changes in

breeding range movements for as long as they were tracked. These results agree with studies on

ringed British honey buzzards, which breed for the first time between the 4th and 7th calendar year

(Roberts and Law 2014). The very late average age of first breeding is rather unexpected for a meso-

predator like the honey buzzard. The similar-sized eurasian goshawk (*Accipiter gentilis*) can breed as

early as the 2nd calendar year, with as much as 42% of females of this resident species starting to

breed that early (Krüger 2005). Short-distance migrant eurasian buzzards (*Buteo buteo*) can also

breed early, but the average age of first breeding is 3.5 years (Davis and Davis 1992), and similarly

short-distance migrant red kites (*Milvus milvus*) breed for the first time between the 3rd and 8th

calendar year, with an average of 3.6 years (Newton et al. 1989). Long-distance migrant black kites

(*Milvus migrans*) start breeding between the 2nd and 8th calendar year (Sergio et al. 2017). Overall,

variability in the age of first breeding is expected in territorial, long-lived species that inhabit

relatively stable environments. Prior to breeding, they need to establish their territories over high-

quality sites, while non-territorial species inhabiting temporal habitats show much more limited

variation in recruitment age (Charlesworth 1994, Aubry et al. 2009, Acker et al. 2018).

In the case of the honey buzzard, the late age of first breeding seems to be associated with a

long delay of first return migrations, and an unexpectedly large variation in the age of returning

individuals in their first returns. Individuals that return to the breeding range in their 3rd calendar

year could likely attempt breeding in their 4th calendar year. By comparison, individuals like

Mohammed (Tab. 1) that attempt a first return as late as the 5th calendar year (or do not succeed to

return before the 6th calendar year) will recruit as breeders several years later. Such long delays of

first return migrations, far beyond the age of sexual maturation and the attainment of adult

plumage, were unexpected. The marked variation in the age of first return migrations, and the high

survival rates of birds 'oversummering' in Africa at late age, suggests that honey buzzards face a

trade-off between survival and attempting to return and recruit at early age. In this sense, it is

striking that Mohamed, the bird to return at latest age, was one of the only birds for which

attempted breeding could be confirmed in the field. It is possible that the age of first return is

constrained by other factors like body condition or environmental factors, as has been posited for

adult migrants that occasionally skip a northern summer by staying in sub-Saharan Africa (Sorensen

et al 2017). However, the fact that at least half of all birds delay the first return at least two years,

without engaging in failed or so-called 'reverse' migrations, suggest these between-individual

differences reflect the viability of both 'fast' and 'slow' life-history strategies in this species. It is

worth noting in this context that the longevity record in honey buzzards is 28 years, exceeding that of

similarly sized raptors, being more similar to that of long-lived seabirds and vultures that also recruit

at later age (Fransson et al 2023).

Natal dispersal and philopatry

We followed the long process of natal dispersal in Honey Buzzards, although only single birds

made it through to establish themselves as terrirtorial, nest-building individuals. Nevertheless, we

were able to document the complexity of the whole multistep process preceeding breeding in a slow-

life, transcontinental migrant species. All the tracked individuals found their way back to Finland and

located themselves relatively close from their natal sites, which agreed with our expectations on the

existance of philopatry in this species. The distance from natal nest and ranging areas decreased with

age, while duration of the stay increased, which also met our expectations on individual

improvements with age. This finding is in line with the results from the few studies documenting

individual improvements by the means of GPS telemetry in slow-life history species. Consequently, in

immature individuals of migratory Ferruginous hawks (*Buteo regalis*) returned significantly closer to

their natal sites in their third spring (when they may already breed), than a year before (Watson et

al. 2019). In partly migrating species, like the red kite, migration routes decresead in length along

with birds age (Literák et al. 2022), also showing adjustment to adult-like behaviour.

Relatively little is known on how philopatric birds actually find their way back home. Do they

repeat the autumn migration backwards up to familiar landscapes? Do they home by learned

gradients in environmental cues (Thorup and Holland 2009, Kishkinev et al 2021) to return to the

place they imprinted during the post-fledgling period? Our study adds insight to these uncertainities

by showing that most individuals performed a loop migration. First return routes crossed or got close

to first outbound routes in only two out of six individuals that completed their first return migration.

Yet all birds managed to home back to Finland, even if some first had to return southwards after

bypassing the Gulf of Bothnia through Sweden. To achieve such accurate navigation, birds may rely

on a variety of mechanisms, such as bi-coordinate gradient maps, whereby they could infer their

relative position by extrapolating from learned gradients in environmental cues (e.g. geomagnetic,

olfactory). Imprinting on "home" takes place prior to abandoning natal sites (Sokolov 2000), as is

evident from the fact that migratory birds that are displaced as eggs or chicks tend to return to the

areas they were displaced to (Cade et al. 2000). Such imprinting takes time and might be one of the

reasons why juveniles in many species, including honey buzzards, are migrating later than adults

(Kjellen et al 1992, Hake et al. 2003, Vansteelant and Agostini 2021). Tracking data confirmed that

juvenile Honey Buzzards were left by both adults at only 1 – 2.5 weeks after fledging (Zsiemer and

Meyburg 2015). The extra time spent by juveniles around natal sites might therefore be used for

further imprinting on the geographical location to which they will return via unknown routes, more

than 18 months after they left.

The process of recruitment to the breeding population

Tracking immature honey buzzards revealed that their recruitment to the breeding

population greatly depended on their migration performance. Whereas relatively few unexperience

juveniles died during their first autumn migration, survival was much lower during the returning

migration. As a result, only about 22% of tracked individuals survived the returning migration

bottleneck and were able to start the recruitment process.

We found that on their first return to the breeding range, immatures arrive much later than

the adults, usually in mid-June, but even as late as mid-July. This is weeks after the breeding adults

and certainly too late to breed. The time of arrival visibly advanced with each subsequent return, due

to a combination of advanced departures from Africa and reduced migration time, which in turn was

mostly due to reduced stop-over time in experienced migrants. Consequently the overall time spent

in the breeding grounds extended , while departure timing from Europe in autumn tended to remain

stable from the 2nd autumn migration onwards. Already on the third return, honey buzzards spent on

average 80 days on the breeding grounds, which is similar to adults (especially the ones that failed to

breed). Therefore, the purpose of the first return in such long-lived migrant species seems to serve

only for the exploration of the breeding grounds and possibly to initiate the annual routine and gain

necessary experience in it.

Especially following the first return, honey buzzards clearly ranged over much greater areas

in the breeding grounds than in following seasons. Wide-ranging behaviour is typical for floaters in

many bird species (Tanferna et al. 2013, Eeden et al. 2017, Mirski and Anderwald 2023). Floaters are

presumed to make such movements to obtain information on nest and foraging site distribution and

quality, which can be inferred from conspecific performance too (see review by Lenda et al. 2012).

The sharp decrease in ranging area and prolonged stay in the breeding grounds after the second

return migration suggests that already at that time, honey buzzards might recruit to the breeding

population, or at least have chosen the area of settlement. Indeed, two of the birds tracked here

were witnessed to build nests after the 2nd migration, in their 5th and 6th calendar year, respectively.

Though nest-building does not yet equate to recruitment as an actual breeding individual, we based

on our results find it is likely that most honey buzzards only engage in their first real breeding

attempt after having experienced multiple migrations at 6-8 years of age. The exact age of first

breeding in long-lived territorial species depends heavily on competitive skills (Becker and Bradley

2007) -which may include migratory performance (Sergio et al 2014, Flack et al 2024)- and habitat

quality (van de Pol et al. 2007), with additional variation in our study species due to the variable age

of first return.

Conclusions

We found that both natal dispersal and recruitment in long-lived trans-continental migrants, such as

honey buzzards, can be extremely long and complex processes. During the extended period of

immaturity, individuals must familiarize themselves with wintering and breeding grounds, adjust

their migration timing to match that of breeding adults, and navigate safely through the challenging

spring conditions of the Mediterranean Sea. Finally, the age of recruitment in long-lived migrants can

span greatly between individuals, largely due to the decision of when to return to the breeding

grounds, which appeared to be a trade-off between survival and early recruitment.

Ethics

Measuring and ringing of birds, was done as inherent of the normal ringing permit (permit 2604) as

issued by the Finnish Museum of Natural History. All fieldwork requiring special permits (taking blood

samples for DNA sexing, sampling and storing DNA-samples, attachment of PTTs/trackers) confirmed

to   five   separate   licences   as   issued   by   Finnish   authorities   (EPOELY/135/07.01.2013,

ESAVI/2195/04.10.07/2014, PIRELY/49/07.01.2013, VARELY/73/07.01.2013, VARELY/215/2015).

Data accessibility

The datasets generated and/or analyzed during the current study are available in the Movebank Data

Repository, https://doi.org/10.5441/001/1.335 (Byholm et al., [2024]).

Declaration of AI use

We have not used AI-assisted technologies in creating this article.

Authors' contributions

PM: conceptualization, data curation, formal analysis, investigation, methodology, visualization,

writing—original draft, writing—review and editing; WV: conceptualization, methodology, writing—

original draft, writing—review and editing; PB: conceptualization, data curation, funding acquisition,

project administration, investigation, methodology, writing—original draft, writing—review and

editing.

All authors gave final approval for publication and agreed to be held accountable for the work

performed therein.

Conflict of interest

We declare we have no competing interests.

Funding

This work was supported by Kone Foundation, Swedish Cultural Foundation in Finland, R.E. Serlachius

Foundation, Svensk-Österbottniska Samfundet and Aktiastiftelserna (to PB).

Acknowledgements

We wish to thank Mikko Honkiniemi, Ari Rantamäki, Jari Valkama, Risto Vilén, Jouko Kivelä, Annika

Rossi, Kari Palo, Hannu Vuoto, Daniel Burgas, Martti Peltola and Ismo Nousiainen for their help with

tagging European honey buzzards in Finland.

REFERENCES

Acker P, Francesiaz C, Béchet A, Sadoul N, Lessells CM, Pijl AS, Besnard A 2018. Insights on dispersal

and recruitment paradigms: sex- and age-dependent variations in a nomadic breeder.

Oecologia 186:85–97. https://doi.org/10.1007/s00442-017-3972-7

Agostini N, Lucia G, Mellone U, Panuccio M, Von Hardenberg J, Evangelidis A, Kominos T. 2012. Loop

migration of adult European Honey Buzzards (Pernis apivorus Linnaeus, 1758) through the

Central-Eastern Mediterranean, Ital J Zool 79:280-286.

https://doi.org/10.1080/11250003.2011.623245

Aikens E, Nourani E, Fiedler W, Wikelski M, Flack A 2024. Learning shapes the development of

migratory behavior. PNAS 121:e2306389121. https://doi.org/10.1073/pnas.230638912

Alonso JC, Martín E, Alonso JA, Morales MB 1998. Proximate and ultimate causes of natal dispersal in

the great bustard Otis tarda. Behav Ecol 9:243–252. https://doi.org/10.1093/beheco/9.3.243

Aubry LM, Cam E, Monnat J-Y 2009. Habitat selection, age-specific recruitment and reproductive

success in a long-lived seabird. In: Thomson DL, Cooch EG, Conroy MJ (eds) Modeling

demographic processes in marked populations. Springer, New York, pp 365–392

Becker PH, Bradley JS 2007. The role of intrinsic factors for the recruitment process in long-lived

birds. J Ornithol 148:377–384

Bensch S, Hasselquist D, Nielsen B, Hansson B 1998. Higher fitness for philopatric than for immigrant

males in a semi-isolated population of great reed warblers. Evolution 52: 877e883.

Bowler DE, Benton TG 2005. Causes and consequences of animal dispersal strategies: relating

individual behaviour to spatial dynamics. Biol Rev 80:205-225.

https://doi.org/10.1017/S1464793104006645

Brønnvik H, Safi K, Vansteelant WM, Byholm P, Nourani E 2022. Experience does not change the

importance of wind support for migratory route selection by a soaring bird. Roy Soc Open

Sci 9: 220746.

Cade TJ 2000. Progress in translocation of diurnal raptors. In: RD Chancellor and B-U Meyburg (EDS.),

Raptors at risk. WWGBP/Hancock House, London, U.K, pp. 343-372.

Calenge C 2006. The package adehabitat for the R software: a tool for the analysis of space and

habitat use by animals. Ecol Model 197:516-519.

Charlesworth B 1994. Evolution in age-structured populations. Cambridge University Press,

Cambridge

Ceresa F, Belda EJ, Monrós JS 2016. Similar dispersal patterns between two closely related birds with

contrasting migration strategies. Popul Ecol 58: 421–427. https://doi.org/10.1007/s10144-016-

0547-0

Coulson JC 2016. A Review of Philopatry in Seabirds and Comparisons with Other Waterbird Species.

Waterbirds 39:229-240. https://doi.org/10.1675/063.039.0302

Davis PE, Davis JE 1992. Dispersal and age of first breeding of Buzzards in central Wales. British Birds

85:578-587.

van Eeden R, Whitfield DP, Botha A, Amar A 2017. Ranging behaviour and habitat preferences of the

Martial Eagle: Implications for the conservation of a declining apex predator. PLOS ONE 12:

e0173956. https://doi.org/10.1371/journal.pone.0173956

Ekman J, Eggers S, Griesser M, Tegelström H 2001. Queuing for preferred territories: delayed

dispersal of Siberian jays. J Anim Ecol 70: 317-324.

Forero MG, Donázar JA, Hiraldo F 2002. Causes and fitness consequences of natal dispersal in a

population of black kites. Ecology 83:858-872. https://doi.org/10.1890/0012-9658

513 Förschler MI, del Val E, Bairlein F 2010. Extraordinary high natal philopatry in a migratory passerine. J

514   Ornithol 151:745–748. https://doi.org/10.1007/s10336-010-0495-y

515 Fransson T, Kolehmainen T, Moss D, Robinson R 2023. EURING list of longevity records for European

516   birds.

517 Gaillard JM, Pontier D, Allaine D, Lebreton JD, Trouvilliez J, Clobert J 1989. An analysis of

518   demographic tactics in birds and mammals. Oikos 56: 59-76.

519 Gamauf A 1999. Is the European honey buzzard (*Pernis apivorus*) a feeding specialist? The influence

520   of social hymenoptera on habitat selection and home range size. Egretta 42:57–85.

521 Gamauf A, Friedl C 2011. Zug- und Überwinterungsstrategien junger Wespenbussarde *Pernis*

522   *apivorus*. Vogelwarte 49:290–291.

523 Greenwood PJ, Harvey PH 1982. The Natal and Breeding Dispersal of Birds. Annual Review of Ecology

524   and Systematics 13:1–21. http://www.jstor.org/stable/2097060

525 Hake M, Kjellén N, Alerstam T 2003. Age-dependent migration strategy in honey buzzards *Pernis*

526   *apiorus* tracked by satellite. Oikos 103:385–396.

527 Healy SD, Hurly TA 2004. Spatial Learning and Memory in Birds. Brain Behav Evol 63: 211–220.

528   https://doi.org/10.1159/000076782

529 Higuchi H, Shiu HJ, Nakamura H, Uematsu A, Kuno K, Saeki M, Hotta M, Tokita KI, Moriya E, Morishita

530   E, Tamura M 2005. Migration of Honey-buzzards *Pernis apivorus* based on satellite tracking.

531   Ornithol Sci 4:109-115. 10.2326/osj.4.109.

532 Howes C, Byholm P, Symes CT 2020b. Forest availability and fragmentation drive movement

533   behaviour of wintering European Honey-buzzard *Pernis apivorus* in Africa. Ardea 108: 115-128.

534   https://doi.org/10.5253/arde.v108i2.a1

Howes C, Symes CT, Byholm P 2020a. Highly skewed sex and age ratios of European Honey Buzzard in

southern Africa. J Ornithol 161: 539–548. https://doi.org/10.1007/s10336-019-01736-w

Keiser JT 2003. Homing Success of Migrants Versus Nonmigrants: Do Differences in Spatial Memory

Generalize Across Spatial Scales?. Dissertations, Theses, and Masters Projects. William and

Mary. Paper 1539626402.

Kishkinev D, Packmor F, Zechmeister T, Winkler HC, Chernetsov N, Mouritsen H, Holland RA 2021.

Navigation by extrapolation of geomagnetic cues in a migratory songbird. Curr Biol

12;317:1563-1569.e4. https://doi.org/10.1016/j.cub.2021.01.051

Kjellén N 1992. Differential timing of autumn migration between sex and age groups in raptors at

Falsterbo, Sweden. Ornis Scand 23:420-434.

Lenda M, Maciusik B, Skórka P 2012.. The evolutionary, ecological and behavioural consequences of

the presence of floaters in bird populations. North-West J Zool 8: 394-408.

Literák, I., Raab, R., Škrábal, J., Vyhnal S, Dostál M, Matušík H, Makoň K, Maderič B, Spakovszky P

2022. Dispersal and philopatry in Central European Red Kites *Milvus milvus*. J Ornithol

163:469–479. https://doi.org/10.1007/s10336-021-01950-5

Luna Á, Palma A, Sanz-Aguilar A, Tella JL, Carrete M 2020. Sex, personality and conspecific density

influence natal dispersal with lifetime fitness consequences in urban and rural burrowing owls.

PLoS ONE 152: e0226089. https://doi.org/10.1371/journal.pone.0226089

Mirski P, Anderwald D 2023. Ranging behavior of non-breeding and breeding adult white-tailed

eagles. Diversity 15:1208.

Nilsson J-A 1989. Causes and Consequences of Natal Dispersal in the Marsh Tit, Parus palustris. J

Anim Ecol 582:619–636. https://doi.org/10.2307/4852

Negro JJ, Hiraldo F, Donazar JA 1997. Causes of Natal Dispersal in the Lesser Kestrel: Inbreeding

Avoidance or Resource Competition? J Anim Ecol 66:640–648. https://doi.org/10.2307/5917

Nevoux M, Arlt D, Nicoll M, Jones C, Norris K 2013. The short- and long-term fitness consequences of

natal dispersal in a wild bird population. Ecol Lett 16:438–445.

Newton I 1979. Population ecology of raptors. T and AD Poyser, Berkhamsted

Newton I, Davis PE, Davis JE 1989. Age of first breeding, dispersal and survival of Red Kites *Milvus*

milvus in Wales. Ibis 31:16-21.

Nourani E, Vansteelant WM, Byholm P, Safi K 2020. Dynamics of the energy seascape can explain

intra-specific variations in sea-crossing behaviour of soaring birds. Biol Lett 16: 20190797.

https://doi.org/10.1098/rsbl.2019.0797

Oppel S, Arkuma V, Bakari S et al 2021. Major threats to a migratory raptor vary geographically along

the eastern Mediterranean flyway. Biol Conserv 262: 109277

https://doi.org/10.1016/j.biocon.2021.109277

Ortego J, Calabuig G, Aparicio JM, Cordero PJ 2008. Genetic consequences of natal dispersal in the

colonial lesser kestrel. Mol Ecol 17:2051-2059. https://doi.org/10.1111/j.1365-

294X.2008.03719.x

Ortego J, García-Navas V, Ferrer E, Sanz J 2011. Genetic structure reflects natal dispersal movements

at different spatial scales in the Blue Tit, *Cyanistes caeruleus*. Animal Behav 82:131-137.

https://doi.org/10.1016/j.anbehav.2011.04.007

Paradis E, Baillie SR, Sutherland WJ, Gregory RD 1998. Patterns of natal and breeding dispersal in

birds. J Anim Ecol 67:518-536. https://doi.org/10.1046/j.1365-2656.1998.00215.x

Pärt T 1994. Male philopatry confers a mating advantage in the migratory collared flycatcher,

Ficedula albicollis. Anim Behav 48:401e409.

QGIS.org 2024. QGIS Geographic Information System. QGIS Association. http://www.qgis.org

R Core Team (2021). R: A language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria. https://www.R-project.org/.

Reed ET, Gauthier G, Pradel R, Lebreton JD 2003. Age and Environmental Conditions Affect

Recruitment in Greater Snow Geese. Ecology 84:219–230.

Roberts SJ, Law C 2014. Honey-buzzards in Britain. British Birds 107:668-691

Saurola P, Valkama J, Velmala W 2013. The Finnish Bird Ringing Atlas. Vol. I. Finnish Museum of

Natural History and Ministry of Environment, Helsinki.

Sæther B-E 1988. Pattern of covariation between life history traits of European birds. Nature

331:616–627.

Sæther B-E, Bakke Ø 2000. Avian life history variation and contribution of demographic traits to the

population growth rate. Ecology 81:642-653.

Schmid H 2000. Separate routes: autumn migration of juvenile and adult European Honey-Buzzards

Pernis apivorus – a synthesis. Der Ornithologische Beobachter 97:191-222.

Sergio F, Tanferna A, Stephanis R, Lopez-Jimenez L, Blas J, Hiraldo F2017.. Migration by breeders and

floaters of a long-lived raptor: implications for recruitment and territory quality. Anim Behav

131:59-72. https://doi.org/10.1016/j.anbehav.2017.07.011

Sokolov L 2000. Philopatry, dispersal and population structure of passerines on the Courish Spit of

the Baltic Sea. Vogelwarte 40:302-314.

Stahl JT, Oli MK 2006. Relative importance of avian life-history variables to population growth rate.

Ecol Model 198:23–39

Sørensen IH, Schlaich AE, Klaassen RHG, Heldbjerg H, Koks BJ 2017. Rare case of an adult male

Montagu's Harrier *Circus pygargus* over-summering in West Africa, as revealed by GPS

tracking. J Ornithol 158: 753–760. https://doi.org/10.1007/s10336-017-1445-8

Tanferna A, López-Jiménez L, Blas J, Hiraldo F, Sergio F 2013. Habitat selection by Black kite breeders

and floaters: Implications for conservation management of raptor floaters. Biol Conserv 160:1–

9.

Thorup K, Alerstam T, Hake M, Kjellén N 2003. Can vector summation describe the orientation

system of juvenile ospreys and honey buzzards? – An analysis of ring recoveries and satellite

tracking. – Oikos 103: 350–359.

Thorup K, Holland RA 2009. The bird GPS - long-range navigation in migrants. J Exp Biol 212:3597-

3604. https://doi.org/10.1242/jeb.021238

van de Pol M, Pen I, Heg D, Weissing FJ 2007. Variation in habitat choice and delayed reproduction:

adaptive queuing strategies or individual quality differences? Am Nat 170:530–541

van Manen W, van Diermen J, van Rijn S, van Geneijgen P 2011. Ecology of Honey Buzzard in the

Veluwe Natura 2000 site (central NL) during 2008-2010, population level, breeding biology,

habitat use and food (in Dutch). Natura 2000 report. Province of Gelderland, Arnhem NL/

Treetop foundation

van Manen W 2013. Biology of Honey Buzzards *Pernis apivorus* in the primaeval forest Białowieża,

Poland. De Takkeling 21:101–126

Vansteelant WMG, Kekkonen J, Byholm P 2017. Wind conditions and geography shape the first

outbound migration of juvenile honey buzzards and their distribution across sub-Saharan

Africa. Proc R Soc B 284: 20170387. http://dx.doi.org/10.1098/rspb.2017.0387

Vansteelant W.M.G., J. Wehrmann, D. Engelen, J. Jansen, B. Verhelst, R. Benjumea, S. Cavaillès, T.

Kaasiku, B. Hoekstra & F. de Boer. 2020. Accounting for differential migration strategies

between age groups to monitor raptor population dynamics in the eastern Black Sea flyway.

Ibis, 162: 356-372. doi: 10.1111/ibi.12773

Vansteelant WMG, Agostoni N 2021. Honey Buzzard *Pernis apivorus*. Pp. 38-45 in: Panuccio M.,

Mellone U. and Agostini A. [eds.]. Migration Strategies of Birds of Prey in Western Palearctic.

CRC Press, Boca Raton, Florida.

Vickers SH, Franco AMA, Gilroy JJ 2023. Non-reproductive dispersal: an important driver of migratory

range dynamics and connectivity. Ecography: e06201. https://doi.org/10.1111/ecog.06201

Watson JW, Banasch U, Byer T, Svingen DN, Mccready R, Hanni D, Gerhardt R 2019. First-Year

Migration and Natal Region Fidelity of Immature Ferruginous Hawks. J Raptor Res 53:266-275.

Weatherhead PJ, Forbes M 1994. Natal philopatry in passerine birds: genetic or ecological

influences? Behav Ecol 5:426–433. https://doi.org/10.1093/beheco/5.4.426

Wiltschko W, Wiltschko R 2017. Homing pigeons as a model for avian navigation? J Avian Biol 48: 66–

74.

Wright J, Cavaillès S, Vansteelant WMG 2019. Photographic evidence of a 2nd calendar year female

European Honey Buzzard *Pernis apivorus* on autumn migration in the Western Palearctic.

Sandgrouse 41:101-104.

Wyllie I, Newton I 1991. Demography of an Increasing Population of Sparrowhawks. J Anim Ecol 60:

749-766.

Zedrosser A, Støen O-G, Sæbø S, Swenson J 2007. Should I stay or should I go? Natal dispersal in

brown bear. Anim Behavi 74:369-376. https://doi.org/10.1016/j.anbehav.2006.09.015

Ziesemer F, Meyburg B-U 2015. Home range, habitat use and diet of Honey-buzzards during the

breeding season. British Birds 108:467– 481.