1 **Unraveling the impact of host genetics and factors on the urinary**
2 **microbiome in a young population**
3

4 Leying Zou[1,†], Zhe Zhang[2,†], Ruijin Guo[2], Xin Tong[2], Junhong Chen[1], Yanmei Ju[2,3],
5 Haorong Lu[4], Huanming Yang[2,5], Jian Wang[2,5], Yang Zong[2], Xun Xu[2], Xin Jin[2], Liang
6 Xiao[2,6], Tao Zhang[1,7*], Xiaomin Liu[2,7*]
7
1. BGI Research, Wuhan 430074, China;
2. BGI Research, Shenzhen 518083, China
3. College of Life Sciences, University of Chinese Academy of Sciences, Beijing
100049, China;
4. China National Genebank, BGI, Shenzhen 518120, China;
5. James D. Watson Institute of Genome Sciences, Hangzhou 310058, China;
6. Shenzhen Engineering Laboratory of Detection and Intervention of Human Intestinal
Microbiome, BGI-Shenzhen, Shenzhen 518083, China;
7. Shenzhen Key Laboratory of Human Commensal Microorganisms and Health
Research, BGI Research, Shenzhen 518083, China
18
19
20 [†] These authors contributed equally.
21
22 * To whom correspondence should be addressed: X.L., liuxiaomin@genomics.cn and
23 T.Z., tao.zhang@genomics.cn

24
25
Abstract

The significance of the urinary microbiome in maintaining health and
contributing to disease development is increasingly recognized. However, a
comprehensive understanding of this microbiome and its influencing factors
remains elusive. Utilizing whole metagenomic and whole-genome
sequencing, along with detailed metadata, we characterized the urinary
microbiome and its influencing factors in a cohort of 1579 Chinese individuals.
Our findings unveil the distinctiveness of the urinary microbiome from other
body sites, delineating five unique urotypes. We identified 137 host factors
significantly influencing the urinary microbiome, collectively explaining 13.16%
of the variance in microbial composition. Notably, gender-related factors,
including sex hormones, emerged as key determinants in defining urotype
groups and microbial composition, with the urinary microbiome exhibiting
strong predictive ability for gender (AUC=0.815). Furthermore, we discovered
43 genome-wide significant associations between host genetic loci and
specific urinary bacteria, particularly *Acinetobacter,* linked to 8 host loci ($p <$
$5x10^{-8}$). These associations were also modulated by gender and sex hormone
levels. In summary, our study provides novel insights into the impact of host

genetics and other factors on the urinary microbiome, shedding light on its implications for host health and disease.

Introduction

Mounting evidence has implicated the presence of a microbiome community within the urinary tract, commonly referred to as urinary microbiome or urobiome[1,2]. This discovery has been substantiated through high-throughput 16S rRNA/metagenomic sequencing, as well as enhanced urine culture techniques across multiple studies[3-6]. In recent years, the urinary microbiota has garnered considerable attention owning to its potential implications for urinary tract health and diseases, such as urine incontinence[7,8], urinary tract infection(UTI)[9,10], chronic kidney disease[11], and bladder cancer[12].

Understanding the urinary microbiome and elucidating the factors that influence its composition and diversity are essential for developing effective diagnostic and therapeutic interventions. Previous studies have identified certain host factors, such as sex and age, that may influence composition of the urinary microbiota[13,14]. A recent study in mainly postmenopausal women has suggested a role for host genetics in microbial variations among individuals[15]. However, these studies have predominantly utilized 16S rRNA sequencing that had relatively low resolution and allowed for the identification and quantification of bacterial taxa at the genus level. Furthermore, existing research has primarily focused on elderly or middle-aged individuals, in which the microbiome may be largely influenced by the environmental factors in the aging process. Thus, there is a gap in knowledge concerning urinary microbiota in young individuals, which our study aims to address. Moreover, there is a necessity to explore additional potential host factors, such as blood metabolites, urine factors, lifestyle, etc., as they may also impact the urinary microbiota. Current research in this area remains limited and calls for more in-depth exploration.

Here in this study, leveraging shotgun metagenomics sequencing and comprehensive metadata from 1579 Chinese individuals from the 4D-SZ cohort[16-21], which is primarily comprised of young adults around 30 years old, we conducted a thorough investigation to enhance our understanding of the urinary microbiota and its influencing factors. Initially, we characterized the urinary microbiome by analyzing its microbial composition and comparing it to microbial communities in other body sites. We then systematically investigated its influencing factors among 386 traits, encompassing sex, age, BMI, diets, lifestyles, as well as blood and urinary measurements. Moreover, by performing the so far first urinary Metagenome-genome wide association study (M-GWAS) in a subset of 687 individuals with available whole genome

88 sequencing data, we endeavor to unravel the intricate interplay between host
89 genetics and the urinary microbiome.
90
91
92
93 **Results**
94
95 **Urinary microbiome: A unique microecology distinct from other body**
96 **sites**
97
We collected urine samples of 1579 individuals, with 66.12% being females
and a mean age of 29 years. Utilizing shotgun metagenomic sequencing, we
generated an average of 19.18 ± 7.90 Gb of data per sample
(**Supplementary Table 1**; **Supplementary Fig. 1**). These samples are from
the 4D-SZ cohort[16-21], with multi-omics data collected including human whole
genome, whole metagenomes across multiple body sites, blood metabolites,
and detailed questionaries. Since we have metagenomic data from not only
the urinary tract but also the other body sites (**Fig. 1a**), including the gut
(n=1754), saliva (n=3222), and reproductive tract (n=686), it prompted us to
compare the urinary microbiome with the microbial communities inhabiting
these distinct body sites. Permutational mutivariate analysis of variance
(PERMANOVA) revealed that the urinary microbiome are significantly distinct
from the other examined microbiomes. It showed a much lower α-diversity
compared to the rich and diverse microbiome harbored in the gut and the oral,
with the highest disparity observed in the phylum level with the gut
microbiome (R^2=0.455, a higher R^2 values corresponding to a larger
difference between the compared sites, p<0.001; **Supplementary Fig. 2**) and
in the genus level with the saliva microbiome (R^2 = 0.361, p<0.001; **Fig. 1b**).
In comparison to the vaginal microbiome, which is known to characterize a
very low α-diversity, the urinary microbiome exhibited the smallest but
significant dissimilarity at both phylum (R^2=0.214) and genus level (R^2 =
0.111), with p<0.001 for both (**Supplementary Fig. 2 and Fig. 1b)**, along with
a significantly higher α-diversity (**Supplementary Fig. 3)**.
121
Further analysis was conducted on metagenomic data from 75 females to
corroborate the above findings, wherein data from all four body sites were
available. Consistent with the broader dataset, the urinary microbiome
exhibited the most pronounced differences from the gut microbiome at the
phylum level (R^2 = 0.299; **Supplementary Fig. 4**), and from the salivary
microbiome at the genus level (R^2 = 0.307; **Fig. 1c**), while displaying relatively
smaller disparities with the vaginal microbiome (R^2 = 0.11 at the phylum level
and R^2 = 0.051 at the genus level; **Supplementary Fig. 4** and
Supplementary Fig. 1c). Importantly, these variations remained statistically
significant (adjusted p < 0.001).

It is evident that these distinctions arise from variations in genus composition across different body sites (**Supplementary Table 2**). For example, *Lactobacillus* was the most abundant genus found in both urinary (an average relative abundance of 17.6%) and vaginal samples (58.9%). while its average relative abundance in saliva and gut was less than 0.13%. The second and third most prevalent genera in urine were *Variovorax* and *Acinetobacter*, with mean relative abundances of 10.6% and 9.4%, respectively; however, their average relative abundances in the other examined sites were less than 0.03%. *Prevotella* is found to be enriched in the saliva with an average relative abundance of 29.1%, while it is less abundant in the gut at 13.4%, followed by urine at 6.7% and vagina at 3.6%. The distinctive microbial compositions and species' diversities across different human body sites resulted in marked dissimilarities in the microbiota across these regions.

In summary, our findings underscore the uniqueness of urinary microbiota in comparison to other anatomical regions such as the gut, saliva, and vagina. This distinct microbial ecosystem warrants further in-depth investigation.

Characteristics of the urinary microbiome

Since the urinary microbiome represents a distinctive microecology, we further characterized its community composition in detail. First, we examined the top ten most abundant genera and species in the cohort (**Supplementary Fig. 5**). At the genus level, *Lactobacillus* had the highest relative abundance, followed by *Variovorax, Acinetobacter, Prevotella,* and *Sphingobium*. At the species level, the top five bacteria with the highest relative abundances, in descending order, were *Lactobacillus iners, Variovorax sp. PDC80, Sphingobium fluviale, Lactobacillus crispatus*, and *Gardnerella vaginalis*. Notably, we observed high Pearson correlations between the relative abundances of bacteria commonly found in the vagina, such as *Lactobacillus*, *Gardnerella*, *Prevotella*, and *Ureaplasma*, in the urinary and vaginal microbiomes (Pearson's r >0.5).

Second, using the partitioning around medoids (PAM) clustering method (Methods), the urinary microbial communities were clustered into five distinct clusters (referred to as "urotypes" analogous to the "enterotypes"). By applying Linear Discriminant Analysis (LDA) for detecting species discriminating each urotype from the rest and ranking them by LDA score (Methods), we determined the dominant species for each urotype. They are *Gardnerella vaginalis* (accounting for 23.10% of the samples)*, Sphingobium fluviale* (34.11%)*, Lactobacillus iners* (15.76%), *Variovorax sp. PDC80* (16.58%), and *Acinetobacter junii* (10.38%), respectively (**Fig. 1d**). Consistent with the clustering results, each urotype showed a significant enrichment of the respective dominant bacteria (**Fig. 1e**).

While the dominant species characterized the highest LDA score for each
urotype, there were more species identified, especially in clusters 1, 2, and 5
(Supplementary Fig. 6; **Supplementary Table 3**). For cluster 1 (*Gardnerella*
vaginalis urotype), the second-ranked species *Lactobacillus crispatus*
followed the dominant bacterium *Gardnerella vaginalis* so close in LDA score,
just like their close relative abundances with an average of 3.91% versus
3.92%. The individuals characterizing this urotype also harbored significantly
higher levels of species belonging to the genus *Prevotella*, including
Prevotella bivia (3.67%), *Prevotella timonensis* (3.43%), *Prevotella disiens*
(3.08%), *Prevotella colorans* (2.84%), and *Prevotella corporis* (2.70%). For
cluster 2 (*Sphingobium fluviale* urotype), species such as *Sphingomonas*
ursincola (3.76%), *Novosphingobium sp AAP93* (3.62%), *Erythrobacter*
donghaensis (3.59%) and *Acinetobacter johnsonli* (3.56%) were also highly
represented. For cluster 3 (*Lactobacillus iners* urotype), *Lactobacillus jensenii*
ranked second with an LDA score of 2.7 and belonged to the same genus
Lactobacillus as the dominant species. In cluster 4 (*Variovorax sp PDC80*
urotype)*, Bradyrhizobium sp CCH1 B1* also showed a high proportion with an
LDA score of 3.76. Finally, in cluster 5 (*Acinetobacter junii* urotype),
Herbaspirillum aquaticum (3.50%), *Cutibacterium acnes* (3.32%), and
Moraxella osloensis (3.31%) were also most abundant and significantly
enriched compared to the other clusters.

Gender-related factors determined urotypes and urinary microbial
communities

To understand the factors influencing urotypes, we calculated their
correlations with host factors. Notably, gender-related factors emerged as the
primary determinants of these urotypes. Menstrual cycle, serum testosterone,
serum estradiol, serum creatinine, proportion of urine, menstrual duration,
menarche age, hemoglobin, and gender were significantly associated with all
five urotypes (**Fig. 2a**, **Supplementary Table 4**). Remarkably, gender
distinctly differentiated the urotype groups. Females exhibited higher relative
abundances of *Gardnerella vaginalis* (urotype 1) and *Lactobacillus iners*
(urotype 3), aligning their urinary microbiome predominantly with these two
urotypes. In contrast, males demonstrated higher abundances of
Sphingobium fluviale (urotype 2)*, Variovorax sp. PDC80* (urotype 4), and
Acinetobacter junii (urotype 5), characterizing their urinary microbial
communities with these three respective urotypes. Additionally, the gender-
biased urotypes displayed significant correlations with gender-dependent
metadata. For instance, female-prevalent urotypes 1 and 3 were correlated
with lower levels of serum testosterone, urine specific gravity, and muscle
mass; while male-prevalent urotypes 4 and 5 were correlated with higher
basal metabolic rate. Besides, some blood amino acids and vitamins were

also related to specific urotypes. For example, homocysteine, proline, and
alpha-aminoadipic acid were positively correlated with urotype 4, while
glutamic acid, vitamin A, sarcosine, beta-alanine, and hemoglobin all showed
positive correlations with urotype 2.

In addition to individual urotypes, we further delved into host factors that
contributed to the overall composition of the urinary microbiome and identified
a total of 137 significant influencing factors (p.adjust < 0.05; **Supplementary**
Table 5). Serum testosterone, gender, urine specific gravity, muscle mass,
basal metabolic rate, serum creatine and estradiol were among the top
influencing factors associated with β-diversity of the urinary microbiome (**Fig.**
2b). The 137 significant covariates collectively explained 13.16% of the
variance of microbial composition (**Fig. 2c**), which encompassed 104 blood
factors (an explained variance of 11.03%), four urinary factors (3.87%),
gender (3.45%), ten anthropometric measurements (3.3%), six health-related
factors (1.84%), seven living habits (1.19%) and five dietary factors (0.84%).
Serum hormones had a substantial impact on the urinary microbiome, with
serum testosterone (explained R^2 = 4.13%) and estradiol (2.4%) being the
first and third most influential ones, respectively. Other metabolites like serum
creatinine (2.45%) and uric acid (1.94%) ranked second and sixth. Amino
acids in the blood, such as leucine (1.82%), isoleucine (1.69%), valine
(1.53%), and glutamic acid (1.35%), also exhibited significant explanatory
powers, as did vitamin A (1.38%) among the vitamins. The collective influence
of urinary factors ranked as the second significant influence among the seven
categories, with urine specific gravity demonstrating the highest explanatory
power (3.22%). Several anthropometric factors had explanatory powers
exceeding 1%, including muscle mass (2.62%), basal metabolic rate (2.53%),
weight (1.91%), height (1.83%), systolic pressure (1.03%), and BMI (0.76%).
Within the health-related factors, the test result of the free total prostate
specific antigen had the most significant impact on the urinary microbiome
(1.14%). In comparison, dietary factors had a comparatively lower effect, with
the highest influencing factor being fat meat favorites, which had an
explanatory power of only 0.25% (**Supplementary Table 6).**

Motivated by the insights that gender and gender-related factors are primary
influencers of the urinary microbiome, we proceeded to investigate differences
in the microbial community between genders in detail. We first observed
significantly higher α-diversity, as measured by the Shannon index, in the
urinary microbiota of females than in males ($p < 8.14 \times 10^{-12}$, **Fig. 3a**). The
microbiota composition also significantly differed between males and females
(R^2 = 0.03, $p < 1 \times 10^{-4}$, **Fig. 3b**). We next sought to identify sex-differentiated
microbes using Linear Discriminant Analysis Effect Size (LEfSe) (**Fig. 3c**).
Females were characterized by higher abundances of genera *Lactobacillus*,
Prevotella, Fannyhessea, Serratia, and *Citrobacter*. However, males exhibited

more microbiota belonging to *Acinetobacter, Bradyrhizobium, Herbaspirillum,*
Moraxella, *Cutibacterium*, and other genera. These results were in line with
the Spearman correlations between bacterial profiles and serum testosterone
(**Supplementary Fig. 7**), suggesting the important effects of sex and sex
hormones on shaping the urinary microbiota.
269
Given the striking gender-related difference in the microbial community, we
employed a random-forest-based machine learning approach to assess the
predictive capability of urinary microbiota for gender classification. The
predictive accuracy demonstrated by the urinary microbiota in determining
gender was notably high, with an area under the curve (AUC) value of
approximately 0.815 (**Fig. 3d**). Notably, the representative species of urotype
including *Lactobacillus iners*, *Acinetobacter junii*, and *Variovorax sp. PDC80,*
were among the top 10 species contributed most significantly to gender
differentiation (**Supplementary Table 7,** T-test p-values < 1.8×10^{-3}), which
echoes with the gender-dependent distribution of the urotypes. These findings
highlight the intricate relationship between microbial composition and host
biological characteristics, suggesting that the urinary microbiome's distinct
makeup is reflective of and influenced by the host's biological sex.
283
284   **Host genetics' influences on the urinary microbiota**
285
Out of a total of 1579 individuals, 687 had host whole genome sequencing
data available, which enabled us to investigate the influence of the host
genome on the urinary microbiome. To achieve this, we conducted a
metagenome-genome-wide association study (M-GWAS) on 106 urinary taxa
present in over 10% individuals (**Supplementary Table 8**). After adjusting for
covariates such as gender, age, BMI, sequencing read counts, and the top ten
host Principal Components (PCs), we identified 43 genome-wide significant
associations involving 27 host genetic loci and 10 urinary taxa ($p < 5\times10^{-8}$,
Fig. 4a, **Supplementary Table 9**).
295
Of all microbial taxa examined at the genus and species level, genus
Acinetobacter was associated with most host genetic loci (n =8; **Fig. 4b**). The
strongest association was observed for SNP rs139799338, located in the
intronic region of the *PKHD1L1* gene ($p=4.82\times10^{-12}$), negatively associated
with the relative abundance of the *Acinetobacter* genus (**Fig. 4a**). *PKHD1L1*
has been found to be significantly elevated in the urine of clear cell renal cell
carcinoma patients[22]. It was also reported to primarily express in both infant
and adult human kidneys and is associated with autosomal recessive
polycystic kidney disease[23]. The SNP rs139799338 demonstrated
associations with various metabolic traits, including the tumor marker
carbohydrate antigen 199 ($p=0.006$), glycine ($p=0.01$), vitamin A ($p=0.01$),
beta-alanine ($p=0.03$), and cobalt ($p=0.047$), as revealed by the genome-wide

association study (GWAS) analysis of host genetic variants and metabolic traits in this cohort (**Supplementary Fig. 8, Supplementary Table 10**).

The second most significant correlation identified was with rs4241281, located in the intergenic region of genes *CAVIN2* and *TMEFF2*, which was found to have a negative association with the genus *Acinetobacter* ($p=9.13\times10^{-12}$, **Fig. 4a**, **Supplementary Table 9**). This genetic locus was also correlated with the relative abundance of genus *Bosea*, ranking as the third strongest association. Additionally, it was also linked to the abundance of species *Acinetobacter johnsonii* belonging to the genus *Acinetobacter* ($p=1.75\times10^{-8}$). *CAVIN2* encodes Caveolae Associated Protein 2 in the kidney, and its removal leads to caveolae loss, while overexpression results in caveolae deformation and membrane tubulation[24]. *TMEFF2* has been identified as an epigenetic biomarker for bladder cancer[25], showing significant associations with increasing tumor grade and stage. Notably, SNP rs4241281 in the *CAVIN2 - TMEFF2* was also correlated with several blood and urinary factors, including concentrations of mercury ($p=0.007$), alpha amino adipic acid ($p=0.015$), albumin ($p=0.035$), body fat mass and rate($p=0.034\sim0.017$), and urine PH ($p=0.047$, **Supplementary Table 10**). Moreover, rs185971980 located in the intergenic region of *FAM170A* and *PRR16*, associated with *Acinetobacter johnsonii* ($p=4.46\times10^{-10}$), was correlated with levels of mean hemoglobin content, mean erythrocyte hemoglobin concentration, average RBC volume and blood iron. Rs117738684 located in the intergenic region of *TMED4* and *OGDH,* associated with *Acinetobacter johnsonii* ($p=3.44\times10^{-9}$), was correlated with levels of blood testosterone, fasting sugar, methionine, and albumin. *LINC01692*, linked to species *Novosphingobium sp. AAP93*, exhibited a positive correlation with urine pH ($\beta=0.087$, $p=9.3\times10^{-4}$). *SENP2*, related to the genus *Acinetobacter*, negatively affected urine specific gravity ($\beta=-0.087$, $p=1.8\times10^{-4}$, **Supplementary Table 10**). These results suggested host genes may influence the urinary microbiota by regulating the blood metabolites and urinary environment.

Gender and sex hormones have demonstrated significant impacts on the urinary microbiome, prompting an exploration of gender-gene interactions using the top two loci associated with specific taxa (**Fig. 4c-f**). An intriguing finding emerged where males with elevated testosterone levels exhibited increased abundance of of *Acinetobacter* compared to females, irrespective of genotype (AA or GA/GG). Interestingly, females with genotype AA exhibited a higher abundance of *Acinetobacter* compared to those with genotype GA/GG, while no discernible differences were observed in males with varying genotypes (**Fig. 4c**). Similarly, an interactive effect between rs139799338 and serum testosterone levels on *Acinetobacter* was observed, where individuals with genotype GA/GG and higher serum testosterone level exhibited higher relative abundance of *Acinetobacter* (AA: R=0.28, $p=2.7\times10^{-13}$, GA/GG:

R=0.65, *p*=0.011, **Fig. 4d**). The gender-gene interaction was also evident for another SNP, rs4241281, linked to *Acinetobacter*, where males exhibited greater average relative abundance than females, particularly with the AA allele influencing *Acinetobacter* in the presence of serum testosterone (R=0.28, *p*=3.7x10^{-13}, **Fig. 4e-f**).

Potential causal relationships between urinary microbiome and diseases

Finally, to explore the potential causal relationships between urinary microbes and diseases by leveraging microbiome-associated host genetic variations, we performed a two-sample Mendelian randomization (MR) analysis. After adjusting for 42 diseases from the BBJ cohort, we identified 7 suggestive causal relationships (p<=1.19x10^{-3}=0.05/42, **Supplementary Table 11, Supplementary Fig. 9**). The strongest MR evidence suggested a link between type 2 diabetes (T2D) and a decreased abundance of *Streptococcus* SGB38301 (β=-0.076, *p*=2.83×10^{-4}). *Paracoccus haematequi* was causally correlated with a decreased risk of T2D (β=-0.013, *p* = 5.12×10^{-4}), while the males-enriched *Variovorax sp. PDC80* linked to an increased T2D risk (β=0.017, *p* = 8.89×10^{-4}). Genus *Herbaspirillum* potentially increased the risk of atopic dermatitis (β=0.064, *p* = 4.41×10^{-4}). *Bosea robiniae* potentially increased the risk of prostate cancer (β=0.058, *p* = 4.65×10^{-4}). Keloid was associated with a lower *Bosea* (β = -0.061, *p* = 7.64×10^{-4}). These inferred relationships underscored a potential connection between urinary microbes and autoimmune diseases, warranting validation in future studies.

Discussion
This study represents the initial profiling of the urinary microbiome in comparison to well-studied human microbiomes from diverse body sites such as the gut, oral cavity, and reproductive tract. While previous studies have also compared the microbial composition across various body sites[26], they relied on databases from different populations and generally involved smaller sample sizes. In contrast, our research provides a more cohesive and comprehensive analysis by utilizing a uniform cohort.

The urinary microbiome was found to possess unique characteristics compared to other sites, evident in the composition of microbial species, as well as α and β diversity differences. The urinary microbiome exhibited significant dissimilarity from other sites, as demonstrated by PCoA analysis in both the entire sample set and the subset of 75 individuals with data from different sites. The difference in microbial communities across body sites was primarily driven by specific taxa. For instance, the gut microbiome was predominantly composed of the genera *Bacteroides* and *Prevotella*, whereas the saliva microbiome showed a gradient of abundance among *Prevotella, Neisseria*, and *Haemophilus*. The urinary microbiome was predominantly

composed of genera such as *Lactobacillus*, *Variovorax*, and *Acinetobacter*, while the vaginal microbiome exhibited more abundant *Lactobacillus* and *Gardnerella*. This highlights the significant role that specific microbial taxa play in characterizing the microbial landscape of different body sites, indicating distinct ecological niches and potentially different functional roles within the human body. Despite the differences, urine and vagina harbored several predominant bacteria in common, supporting the relatively closer relatedness between them[27] compared to other physically distal sites.

We identified five urotypes and their respective dominant bacteria. A previous study extracted microbiota from encrustations on ureteral stents in patients without urinary tract infections and conducted urotype analysis[28]. Their findings, similar to ours, identified types dominated by microbes such as *Lactobacillus*, *Gardnerella vaginalis*, *Achromobacter*, *Corynebacterium*, and *Streptococcus* (**Supplymentary Table 3**). However, they identified extra bacteria like Actinomycetales, Enterobacterales, and *Staphylococcus* which were less abundant in this cohort. These differences probably have stemmed from the distinct sources of the analyzed microbiome. Their study focused on bacteria from ureteral stents, on which nosocomial bacteria may form biofilm. They employed the 16S rRNA sequencing method, which differs from our metagenomic sequencing approach.

Although some studies indicated the urinary microbiome may be impacted by sex[29-32], they were performed in small samples and mainly based on the 16S rRNA data, resulting in limited sex-specific bacteria identification. In this study, we discovered extensive sex differences in the urinary microbiota which could well predict sex (AUC=0.815). We also observed a strong link between the urinary microbiome and sex hormone levels as well as some other sex-specific factors. These findings not only enrich our understanding of the gender-specific susceptibilities to certain urinary tract conditions or diseases but also open avenues for utilizing microbial profiles in gender-specific medical research and potential diagnostic applications.

Our study also marks the first detailed exploration of the impact of blood metabolites on the urinary microbiome. Serum hormones, particularly serum testosterone and serum estradiol, emerged as significant influencers of the urinary microbiome. Other metabolites, such as serum creatinine and serum uric acid, along with blood amino acids like leucine, isoleucine, valine, glutamic acid, and vitamin A, were also found to play crucial roles, highlighting the intricate relationship between various serum components and the composition of the urinary microbiome. Serum creatinine has been regarded as a biomarker of renal function for almost a century[33] and is widely utilized in clinical diagnostics. In our study, serum creatinine showed significant explanatory power for the β-diversity of urinary microbiota and different

urotypes, suggesting that serum creatinine levels might influence kidney
function through its effects on the composition of urinary microbiota. The
potential causal relationship warrants further investigation. This link
emphasizes the intricate connections between metabolic biomarkers and
microbiome composition, highlighting the importance of a holistic approach to
understanding renal health and disease.
446
Limited studies investigated the effect of host genome on the urinary
microbiome, except a previous study that reported considerable genetic
contribution and identified heritable *Escherichia* (A = 0.165) and *Lactobacillus*
iners (A = 0.177) by analyzing midstream urine samples from 1,600 older
females in the TwinsUK cohort[15]. This study represents a pioneering effort in
conducting a genome-wide association analysis of the urinary microbiome
along with the host genome. We have successfully identified 27 significant
genomic loci associated with 10 urinary microbes, demonstrating the influence
of genetic factors on the composition of the urinary microbiome. Genus
Acinetobacter and its species *Acinetobacter johnsonii* showed the most link to
host genetic loci. In addition to the urinary microbiome, *Acinetobacter*
johnsonii exhibited a genetic attribute to the skin microbiome[34]. In this GWAS,
we also observed an association between genus Acinetobacter and the
MARCO gene (β=-0.183, *p*=3.67x10^{-6}), which encoded macrophage receptor
with collagenous structure and correlated with stress urinary incontinence
(replication *p*= 0.003) in a GWAS of European women[35]. Moreover, we delved
deeper into exploring the impact of interactions between significant loci
genotypes, gender, and serum testosterone levels on specific urinary
microbes. The insights gained from these analyses offer valuable information
on the complex interplay between host genetics, hormonal levels, and
microbial abundance.
468
469
470 **Methods**
471
Study subjects
All the participants involved in this research belonged to the '4D-SZ' cohort,
which encompassed the collection of blood, gut, saliva, vaginal, and urine
samples, along with comprehensive metadata, for a multi-omics investigation
as previously reported[16-21]. In this study, 1579 urine samples from the cohort
were collected for whole metagenomic sequencing (**Supplementary Table 1**).
The detailed urine sampling, sequencing, and quality control design were
described below. For comparison, we also included metagenomic samples
from the gut (n=1754), saliva (n=3222), and reproductive tract (n=686), all
from the 4D-SZ cohort. Feces were collected with an MGIEasy kit, and stool
DNA was extracted in accordance with the MetaHIT protocol as described
previously[36]. For the salivary sample, we employed a 2× concentration of a

stabilizing reagent kit and collected 2 mL of saliva. The DNA extraction from
saliva samples was conducted using the MagPure Stool DNA KF Kit B (no.
MD5115-02B). For reproductive tract samples, the extraction of genomic DNA
from was performed in accordance with the established protocol[37](Hao
L,2020). The DNA concentrations from blood, stool, oral, and reproductive
tract samples were estimated by Qubit (Invitrogen), and the library preparation
and sequencing were conducted in accordance with the procedures described
in previous studies[16-21].
All recruitment and study procedures involving human subjects were
approved by the Institutional Review Boards (IRB) at BGI-Shenzhen. All
participants provided written informed consent at enrolment.
495
Urine metagenomic sampling, sequencing, and quality control
Midstream voided urine was self-collected using a 40-ml urine cup by 1579
individuals from the 4D-SZ cohort. From the urine cup, 5ml urine was flowed
into a sterile tube containing 5ml BGI stabilizing reagent[38] for the preservation
of metagenome at room temperature and then stored at -80°C within the day
before subsequent metagenomic analysis.
502
503 The samples were thawed and transferred into a 15mL centrifuge tube. Then
504 1mL TE buffer (1M Tris-HCL & 0.1M EDTA) for crystal dissolution[39] and
505 200uL 50mg/mL lysozyme were added before centrifuge at 8000rpm for
506 10min. Then DNA extraction was proceeded with MagPure Stool DNA KF Kit
507 B (MD5115, Magen)[40].
508
DNA extraction and library preparation were performed along with blank
negative controls and mock community controls. Specifically,
ZymoBIOMICS® Microbial Community Standard (D6300) and
ZymoBIOMICS® Microbial Community DNA Standard (D6305) were 1x10^4
times diluted using saline to simulate the low microbial load in urine samples
and divided into aliquots (1ml per tube) for storage at -80°C before using as
extraction and library preparation controls respectively. During extraction,
blank negative control and extraction control (D6300 diluted) were added for
each 96-well plate and processed with the same procedure as for the urine
samples. Similarly, separate blank negative control and library preparation
control (D6305 diluted) were also added for each 96-well plate during library
preparation and processed with the same procedure as for the urine samples.
Notably, none of the negative controls for both extraction and library
preparation passed the library preparation stage for PCR concentration lower
than the minimum criteria of 1ng/uL. In total, 33 extraction mock controls and
43 library mock controls generated metagenomic sequencing data.
525
526 **Filtering and Removal of Possible Contaminants**
527 As low biomass microbiome tends to be more easily influenced by

contaminants and cross-talk, multiple contamination control steps were further applied to the sequencing data.

Step 1. For all the samples in the entire cohort, species were considered as possible contaminants and collectively removed if a species meets one of the following criteria:

A. Relative abundance significantly correlated with PCR concentration (Spearman test, p-value <0.05)

B. Decontam score <0.3 calculated by 'decontam' package[41] based on relative abundance and PCR concentration or occurrence in <4 samples

C. Detected in the 76 mock controls but other than the 10 expected species comprising the mock community with $> 1 \times 10^{-4}$ mean relative abundance

D. Reads count less than 1000

Step 2. For all the samples in each specific 96-well plate, remaining species other than the 10 expected species detected in the respective mock control with $>1 \times 10^{-4}$ relative abundance were considered as contaminants specific to that 96-well plate and were removed.

Step 3. Filtering out non-bacterial microbes and retaining only bacteria taxa and renormalization

Constructing the urinary metagenomic taxonomic profile

Taxonomy assignment and functional prediction were performed using MetaPhlAn4[42] version 4.0.6 with default settings. MetaPhlAn 4, an advanced tool for microbial analysis, utilizes around 5.1 million unique clade-specific marker genes derived from approximately 1.01 million microbial genomes (236,600 reference genomes and 771,500 metagenomic assembled genomes), covering 26,970 species-level genome bins (21,978 kSGBs and 4,992 uSGBs). Compared to its predecessor, MetaPhlAn 3, MetaPhlAn 4 identifies a broader and more detailed range of urinary microbiota. Utilizing this tool, we obtained a raw microbial taxonomic dataset composed of a total of 1,063 taxa (11 phyla, 32 classes, 60 orders, 115 families, 248 genera, and 597 species).

Calculating α-, β- diversity, PCOA and urotypes

The microbial diversity metrics, including α-diversity (Shannon and Simpson indices) and β-diversity (Bray–Curtis dissimilarities), were calculated based on species-level abundance data using the 'diversity' and 'vegdist' functions within the R package 'vegan', respectively. Subsequently, principal coordinates analysis (PCoA) was conducted on the computed beta-diversity dissimilarities utilizing the 'capscale' function from the 'vegan' package.

For sample clustering based on relative species abundances, the JSD
distance metric and the Partitioning Around Medoids (PAM) clustering
algorithm were employed. The optimal number of clusters was determined
using the Calinski-Harabasz (CH) Index[43] (**Supplementary Fig. 10**).
Following this, the Linear Discriminant Analysis Effect Size (LEfSe)[44], a
nonparametric cross-sample test utilizing Linear Discriminant Analysis (LDA)
for test statistic computation, supported by traditional univariate tests for
feature selection, was utilized. The microbial taxa were ranked based on their
LDA values in descending order, and the predominant bacteria within each
urotype were identified as the species with the highest LDA value.
581
Host factors explained the variance of the urinary microbiome
We investigated the correlations between urinary microbiome compositions
and 420 host variables encompassing anthropometric measurements, blood
and urine parameters, among others. To identify variables significantly linked
to β-diversity, we employed Bray-Curtis distance-based redundancy analysis
(dbRDA) and determined the variance explained by these factors using the
'capscale' function in the vegan package. Significance of each response
variable was verified through analysis of variance (ANOVA) for the dbRDA
using the 'anova.cca()' function in the vegan package. Only variables showing
significant associations (Benjamini-Hochberg FDR < 0.05) with β-diversity in
univariable models were included in the multivariable analysis. These 137
significant factors were categorized into distinct groups including urinary
parameters, anthropometric measures, health-related indicators, lifestyle
habits, and dietary elements. To assess the combined explanatory power of
these variable groups on urinary microbiome compositions, we performed a
variation partitioning analysis using the vegan package. The 'adj.r.squared'
value was calculated using the RsquareAdj option to quantify the proportion of
variance explained by the different categories.
600
High-depth whole-genome sequencing for the blood samples
Among the 1,579 individuals, 687 with blood samples were sequenced to a
mean of 30× for the whole genome. For blood samples, buffy coat was
isolated and DNA was extracted using HiPure Blood DNA Mini Kit (Magen,
catalog no. D3111) according to the manufacturer's protocol. The library
construction and sequencing methods are consistent with those described in
previous articles[16].
For genomic analysis, whole-genome reads were aligned to the
GRCh38/hg38 human genome using BWA[45] (v0.7.15), with low-quality reads
filtered out. Alignments were processed into BAM files using Samtools[46]
(v0.1.18) and PCR duplicates were marked with Picardtools (v1.62). The
Genome Analysis Toolkit (GATK[47], v3.8) facilitated SNP and INDEL
identification and recalibration, referencing dbSNP (v150). Variant discovery
and processing were performed with GATK's HaplotypeCaller and subsequent

tools. The final dataset, comprising 7.2 million SNPs/INDELs(MAF ≥ 1%), were remained for M-GWAS analyses.

For variant inclusion, we set stringent thresholds: (i) a minimum average depth of over 8x; (ii) Hardy-Weinberg equilibrium (HWE) P-value >10−5; and (iii) a genotype calling rate >98%. Regarding sample criteria, we required: (i) mean sequencing depth >20 ×; (ii) a variant call rate >98%; (iii) absence of population stratification, confirmed through principal components analysis (PCA) conducted using PLINK[48](version 1.07); and (iv) the exclusion of related individuals, determined by calculating pairwise identity by descent (IBD, with a Pi-hat threshold of 0.1875) in PLINK. Following these rigorous quality control measures, 687 individuals were left for subsequent analysis.

Predictive analysis of gender using urinary microbiota

We employed the Random Forest algorithm to explore the potential of using the average relative abundance of urinary microbiota for gender prediction. This approach was chosen due to its efficacy in handling complex biological data and its robustness in feature selection, making it particularly suitable for high-dimensional datasets like microbiome compositions. We first divided our dataset into 70% training and 30% testing sets, ensuring a balanced representation of male and female samples. The Random Forest model was then trained on the training set with gender as the response variable and the average relative abundance of different urinary microbes as predictor variables. We used 500 trees in our model to ensure adequate learning and generalization capability. The model's performance was evaluated using the Area Under the Receiver Operating Characteristic (ROC) Curve (AUC), which provides a comprehensive measure of model accuracy across different classification thresholds. Furthermore, we conducted t-tests to identify the top 10 differentially abundant microbes between genders.

Association analysis for microbial taxa

Considering the power of GWAS tests, we refined our approach to focus on microbial taxa exhibiting occurrence rates exceeding 10% (present in at least 157 individuals) and an average relative abundance above 1×10^{-4}. After filtering, the represented genera of these microbial taxa covered 99.73% of the whole community in the cohort. Subsequently, a total of 106 microbial genera and species were included for association analyses.

We examined the relationships between host genetics and the urine microbiome using two different statistical models: a linear model based on the relative abundance and a logistic model based on the presence/absence (P/A) of microbial features. For those taxa present in over 50% individuals, we used a linear model and transformed the relative abundance data using a natural logarithm. Subsequently, we calculated the residuals by employing the 'lm' function in R: (log10(Microbe abundance) _ age + sex + BMI +

sequencing read counts + top ten PCs). These residuals were then extracted
using the residuals() function from the R stats package and applied in a
univariate linear model for association analysis with genotypes. For the
microbial taxa observed in more than 10% but less than 90% of individuals,
we converted the data into presence/absence patterns to avoid issues with
zero inflation. By dichotomizing these data, we could treat the bacterial
abundance as a binary trait for logistic regression analysis. This analysis was
adjusted for the same set of covariates as mentioned earlier. This dual
approach allowed us to comprehensively assess the influence of host
genetics on different aspects of the urine microbiome.

We searched the oral microbiome-related SNPs in the summary statistics data
from this cohort as previously reported to examine their associations with host
traits.

Additionally, we conducted GWAS association analyses similar to the steps
described above for α, β diversity, and urinary clusters, however, no genome-
wide significant signals were observed (**Supplementary Tables 12 & 13**). We
also performed gene-environment interaction analyses on the top two
significant loci, chr8:109442613 and chr2:191850055. These analyses,
focusing on interactions with gender and serum testosterone levels, were
executed using the --interaction parameter in PLINK.

Two-sample MR analysis
To explore the causal relationships between the urinary microbiome and
various diseases, we accessed summary statistics data of 42 diseases from
Biobank Japan[49,50]. The two-sample bidirectional MR analysis was performed
by applying the GSMR method[51]. Genetic variants with $P < 1 \times 10^{-5}$ and LD r^2
<0.1 were also selected as instrumental variables for phenotypes in the Japan
Biobank study.

Data availability
The metagenomic sequencing data after removing host reads have been
deposited to the CNGB Nucleotide Sequence Archive (CNSA:
https://db.cngb.org/cnsa; accession number CNP0003862). The summary
statistics that support the findings of this study including the associations
between host genetics and urinary microbiome are available from
https://db.cngb.org/search/project/CNP0005626/. According to the Human
Genetic Resources Administration of China regulation and the institutional
review board of BGI-Shenzhen related to protecting individual privacy,
sequencing data are controlled-access and are available via application on
request.

Acknowledgments

We sincerely thank the support provided by China National GeneBank. We thank all the volunteers for their time and for self-collecting the oral samples using our kit. This work was supported by the National Natural Science Foundation of China (No. 32200548).

Author contributions

X.L. and T.Z. conceived and organized this study. J.W. initiated the overall health project. X.X., X.J., L.X., Y.H., and Y.Z. contributed to organization of the cohort, the sample collection and questionnaire collection. H.L led the DNA extraction and sequencing. L.Z. and Z.Z. processed the metagenome data. X.L and X.T. processed the whole genome data. L.Z. and X.L performed the metagenome-genome-wide association analyses. L.Z., Z.Z., and X.L wrote the raw manuscript. All authors contributed to data and texts in this manuscript.

Declaration of interests

The authors declare no competing interest.

References

Wolfe, A. J. & Brubaker, L. Urobiome updates: advances in urinary microbiome research. *Nat Rev Urol* **16**, 73-74 (2019). https://doi.org:10.1038/s41585-018-0127-5

Whiteside, S. A., Razvi, H., Dave, S., Reid, G. & Burton, J. P. The microbiome of the urinary tract—a role beyond infection. *Nature Reviews Urology* **12**, 81-90 (2015). https://doi.org:10.1038/nrurol.2014.361

Pearce, M. M. *et al.* The female urinary microbiome: a comparison of women with and without urgency urinary incontinence. *mBio* **5**, e01283-01214 (2014). https://doi.org:10.1128/mBio.01283-14

Ishihara, T. *et al.* Usefulness of next-generation DNA sequencing for the diagnosis of urinary tract infection. *Drug Discoveries & Therapeutics* **14**, 42-49 (2020). https://doi.org:10.5582/ddt.2020.01000

Price, T. K. *et al.* The Clinical Urine Culture: Enhanced Techniques Improve Detection of Clinically Relevant Microorganisms. *J Clin Microbiol* **54**, 1216-1222 (2016). https://doi.org:10.1128/jcm.00044-16

Thomas-White, K. *et al.* Culturing of female bladder bacteria reveals an interconnected urogenital microbiota. *Nature Communications* **9**, 1557 (2018). https://doi.org:10.1038/s41467-018-03968-5

Komesu, Y. M. *et al.* The urinary microbiome in women with mixed urinary incontinence compared to similarly aged controls. *International Urogynecology Journal* **29**, 1785-1795 (2018). https://doi.org:10.1007/s00192-018-3683-6

Govender, Y., Gabriel, I., Minassian, V. & Fichorova, R. The Current Evidence on the Association Between the Urinary Microbiome and Urinary Incontinence in Women. *Frontiers in Cellular and Infection Microbiology* **9** (2019).

https://doi.org:10.3389/fcimb.2019.00133

9   Moustafa, A. *et al.* Microbial metagenome of urinary tract infection. *Scientific Reports*
8 (2018). https://doi.org:10.1038/s41598-018-22660-8

10   Vaughan, M. H. *et al.* The Urinary Microbiome in Postmenopausal Women with
Recurrent Urinary Tract Infections. *Journal of Urology* **206**, 1222-1231 (2021).
https://doi.org:doi:10.1097/JU.0000000000001940

11   Kramer, H. *et al.* Diversity of the midstream urine microbiome in adults with chronic
kidney disease. *International Urology and Nephrology* **50**, 1123-1130 (2018).
https://doi.org:10.1007/s11255-018-1860-7

12   Wu, P. *et al.* Profiling the Urinary Microbiota in Male Patients With Bladder Cancer in
China. *Front Cell Infect Microbiol* **8**, 167 (2018).
https://doi.org:10.3389/fcimb.2018.00167

13   Qin, J. *et al.* Characterization of the Genitourinary Microbiome of 1,165 Middle-Aged
and Elderly Healthy Individuals. *Front Microbiol* **12**, 673969 (2021).
https://doi.org:10.3389/fmicb.2021.673969

14   Ammitzbøll, N. *et al.* Pre- and postmenopausal women have different core urinary
microbiota. *Scientific Reports* **11**, 2212 (2021). https://doi.org:10.1038/s41598-021-
81790-8

15   Adebayo, A. S. *et al.* The Urinary Tract Microbiome in Older Women Exhibits Host
Genetic and Environmental Influences. *Cell Host Microbe* **28**, 298-305.e293 (2020).
https://doi.org:10.1016/j.chom.2020.06.022

16   Liu, X. *et al.* A genome-wide association study for gut metagenome in Chinese adults
illuminates complex diseases. *Cell Discovery* **7**, 9 (2021).
https://doi.org:10.1038/s41421-020-00239-w

17   Liu, X. *et al.* Metagenome-genome-wide association studies reveal human genetic
impact on the oral microbiome. *Cell Discovery* **7**, 117 (2021).
https://doi.org:10.1038/s41421-021-00356-0

18   Liu, X. *et al.* Mendelian randomization analyses support causal relationships between
blood metabolites and the gut microbiome. *Nature genetics* **54**, 52-61 (2022).
https://doi.org:10.1038/s41588-021-00968-y

19   Jie, Z. *et al.* A transomic cohort as a reference point for promoting a healthy human
gut microbiome. *Medicine in Microecology* **8**, 100039 (2021).
https://doi.org:https://doi.org/10.1016/j.medmic.2021.100039

20   Zhu, J. *et al.* Over 50,000 Metagenomically Assembled Draft Genomes for the Human
Oral Microbiome Reveal New Taxa. *Genomics, Proteomics & Bioinformatics* (2021).
https://doi.org:https://doi.org/10.1016/j.gpb.2021.05.001

21   Chen, C. *et al.* Cervicovaginal microbiome dynamics after taking oral probiotics.
Journal of Genetics and Genomics **48**, 716-726 (2021).
https://doi.org:https://doi.org/10.1016/j.jgg.2021.03.019

22   Yang, Y. *et al.* Excavation of diagnostic biomarkers and construction of prognostic
model for clear cell renal cell carcinoma based on urine proteomics. *Frontiers in*
Oncology **13** (2023). https://doi.org:10.3389/fonc.2023.1170567

23   Onuchic, L. F. *et al.* PKHD1, the Polycystic Kidney and Hepatic Disease 1 Gene,
Encodes a Novel Large Protein Containing Multiple Immunoglobulin-Like Plexin-

Transcription–Factor Domains and Parallel Beta-Helix 1 Repeats. *The American*
Journal of Human Genetics **70**, 1305-1317 (2002).
https://doi.org:https://doi.org/10.1086/340448

24   Hansen, C. G., Bright, N. A., Howard, G. & Nichols, B. J. SDPR induces membrane
curvature and functions in the formation of caveolae. *Nat Cell Biol* **11**, 807-814
(2009). https://doi.org:10.1038/ncb1887

25   Costa, V. L. *et al.* Three epigenetic biomarkers, GDF15, TMEFF2, and VIM,
accurately predict bladder cancer from DNA-based analyses of urine samples. *Clin*
Cancer Res **16**, 5842-5851 (2010). https://doi.org:10.1158/1078-0432.Ccr-10-1312

26   Adebayo, A. S. *et al.* The Urinary Tract Microbiome in Older Women Exhibits Host
Genetic and Environmental Influences. *Cell Host & Microbe* **28**, 298-305.e293 (2020).
https://doi.org:https://doi.org/10.1016/j.chom.2020.06.022

27   Komesu, Y. M. *et al.* Defining the relationship between vaginal and urinary
microbiomes. *American Journal of Obstetrics and Gynecology* **222**, 154.e151-
154.e110 (2020). https://doi.org:https://doi.org/10.1016/j.ajog.2019.08.011

28   Buhmann, M. T. *et al.* Encrustations on ureteral stents from patients without urinary
tract infection reveal distinct urotypes and a low bacterial load. *Microbiome* **7**, 60
(2019). https://doi.org:10.1186/s40168-019-0674-x

29   Siddiqui, H., Nederbragt, A. J., Lagesen, K., Jeansson, S. L. & Jakobsen, K. S.
Assessing diversity of the female urine microbiota by high throughput sequencing of
16S rDNA amplicons. *BMC Microbiol.* **11**, 244 (2011). https://doi.org:10.1186/1471-
2180-11-244

30   Fouts, D. E. *et al.* Integrated next-generation sequencing of 16S rDNA and
metaproteomics differentiate the healthy urine microbiome from asymptomatic
bacteriuria in neuropathic bladder associated with spinal cord injury. *J Transl Med* **10**,
174 (2012). https://doi.org:10.1186/1479-5876-10-174

31   Lewis, D. A. *et al.* The human urinary microbiome; bacterial DNA in voided urine of
asymptomatic adults. *Front Cell Infect Microbiol* **3**, 41 (2013).
https://doi.org:10.3389/fcimb.2013.00041

32   Gottschick, C. *et al.* The urinary microbiota of men and women and its changes in
women during bacterial vaginosis and antibiotic treatment. *Microbiome* **5**, 99 (2017).
https://doi.org:10.1186/s40168-017-0305-3

33   Bjornsson, T. D. Use of Serum Creatinine Concentrations to Determine Renal
Function1. *Clinical Pharmacokinetics* **4**, 200-222 (1979).
https://doi.org:10.2165/00003088-197904030-00003

34   Moitinho-Silva, L. *et al.* Host genetic factors related to innate immunity, environmental
sensing and cellular functions are associated with human skin microbiota. *Nat*
Commun **13**, 6204 (2022). https://doi.org:10.1038/s41467-022-33906-5

35   Cartwright, R. *et al.* Genome-Wide Association Study Identifies Two Novel Loci
Associated with Female Stress and Urgency Urinary Incontinence. *Journal of Urology*
206, 679-687 (2021). https://doi.org:doi:10.1097/JU.0000000000001822

36   Qin, J. J. *et al.* A metagenome-wide association study of gut microbiota in type 2
diabetes. *Nature* **490**, 55-60 (2012). https://doi.org:10.1038/nature11450

37   Lilan, H. & Lilan, H. DNA extraction for human microbe samples. *protocols.io* (2020).

38   Han, M. *et al.* A novel affordable reagent for room temperature storage and transport
of fecal samples for metagenomic analyses. *Microbiome* **6**, 43 (2018).
https://doi.org:10.1186/s40168-018-0429-0
39   Ackerman, A. L. *et al.* Optimization of DNA extraction from human urinary samples for
mycobiome community profiling. *PLoS One* **14**, 26 (2019).
https://doi.org:10.1371/journal.pone.0210306
40   Yang, F. M. *et al.* Assessment of fecal DNA extraction protocols for metagenomic
studies. *GigaScience* **9**, 12 (2020). https://doi.org:10.1093/gigascience/giaa071
41   Davis, N. M., Proctor, D. M., Holmes, S. P., Relman, D. A. & Callahan, B. J. Simple
statistical identification and removal of contaminant sequences in marker-gene and
metagenomics data. *Microbiome* **6**, 226 (2018). https://doi.org:10.1186/s40168-018-
0605-2
42   Blanco-Miguez, A. *et al.* Extending and improving metagenomic taxonomic profiling
with uncharacterized species using MetaPhlAn 4. *Nat. Biotechnol.* **41**, 1633-+ (2023).
https://doi.org:10.1038/s41587-023-01688-w
43   Caliński, T. & Harabasz, J. A dendrite method for cluster analysis. *Communications in*
Statistics **3**, 1-27 (1974). https://doi.org:10.1080/03610927408827101
44   Segata, N. *et al.* Metagenomic biomarker discovery and explanation. *Genome*
Biology **12**, R60 (2011). https://doi.org:10.1186/gb-2011-12-6-r60
45   Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler
transform. *Bioinformatics* **25**, 1754-1760 (2009).
https://doi.org:10.1093/bioinformatics/btp324
46   Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**,
2078-2079 (2009). https://doi.org:10.1093/bioinformatics/btp352
47   McKenna, A. *et al.* The Genome Analysis Toolkit: A MapReduce framework for
analyzing next-generation DNA sequencing data. *Genome Res.* **20**, 1297-1303
(2010). https://doi.org:10.1101/gr.107524.110
48   Purcell, S. *et al.* PLINK: A tool set for whole-genome association and population-
based linkage analyses. *Am. J. Hum. Genet.* **81**, 559-575 (2007).
https://doi.org:10.1086/519795
49   Ishigaki, K. *et al.* Large-scale genome-wide association study in a Japanese
population identifies novel susceptibility loci across different diseases. *Nature Genet.*
52, 669-+ (2020). https://doi.org:10.1038/s41588-020-0640-3
50   Kanai, M. *et al.* Genetic analysis of quantitative traits in the Japanese population links
cell types to complex human diseases. *Nature Genet.* **50**, 390-+ (2018).
https://doi.org:10.1038/s41588-018-0047-6
51   Zhu, Z. *et al.* Causal associations between risk factors and common diseases inferred
from GWAS summary data. *Nature Communications* **9**, 224 (2018).
https://doi.org:10.1038/s41467-017-02317-2
874
875
876
877

Figure captions

Figure 1. Comparative analysis of microbial communities across human body sites and characterizing the urinary microbiome.
 a) An illustrative diagram presenting the total and shared sample counts of microbial communities from four body sites: saliva, gut, urine, and the reproductive tract. **b**) A scatterplot illustrating the genus-level PCoA distribution of the total microbial samples from saliva, gut, urine, and reproductive tract, along with PERMANOVA pairwise difference values. **c**) A scatterplot showing the genus-level PCoA distribution of 75 shared samples among saliva, gut, urine, and reproductive tract, with PERMANOVA pairwise difference values. **d**) A scatterplot reflecting the species-level PCoA distribution of urinary microbes, the urotypes classification, and the dominant bacteria present in each urotype. **e**) Boxplots displaying the average relative abundance of the five dominant bacteria among individuals with five different urotypes, along with t-test differential analysis.

Figure 2. Impact of host phenotypes on urinary microbiota.
This scatterplot presents the distribution of Pearson correlation estimate values between host phenotypes and five urotypes, highlighting the top 20 significant associations for each urotype, with colors corresponding to the urine classification results in Figure 1d. Each phenotype is ranked from high to low according to the average of Pearson correlation estimate value for each urotype. **b**) A bar chart showing the top 40 significant host phenotypes sorted by their explained variance (R^2 values) in the β-diversity of urinary microbiota. **c**) A categorized bar chart displaying the explained variance of significant host phenotypic factors for the urinary microbiota β-diversity, including: 137 total host factors, 104 blood factors, 4 urinary factors, gender, 10 anthropometric factors, 6 health factors, 7 lifestyle factors, and 5 diet-related factors.

Figure 3. The urinary microbiome differed between genders and effectively distinguished gender.
a) Box plot of α-diversity (Shannon index) of the urinary microbiome between males and females. **b**) β-diversity of the urinary microbiome using principal coordinate analysis, as colored by males (cyan) and females (red). The first and second axes, PCoA1 and PCoA2, explaining the highest variance, are shown. Dashed ellipses represent the 95% confidence level of permutational multivariate analysis of variance (PERMANOVA) test. Explained variance (R^2) and p-value were also been showed. **c**) The sex differential urinary microbiota, as sorted by LDA scores in the LEfSe analysis**. d**) A ROC curve demonstrating the predictive power of the urinary microbiota for gender using a random forest model. The analysis comprises 520 male samples and 1044 female samples.

Figure 4. Influences of host genetics and their interactions with gender/sex hormones on urinary microbiota.

a) A Manhattan plot showing the genetic variants associated with urine microbial taxa (n=106), with genome-wide significance denoted by horizontal black lines (p = 5 × 10^{-8}) and gene names annotated for 27 significant loci. **b**) A bar chart displaying the urinary microbes associated with the significant 27 loci and the number of associations. **c**) Boxplots illustrating the average relative abundance of genus *Acinetobacter* among individuals with different genotypes (AA vs GA/GG) at the top locus rs139799338, as well as stratified by male and female samples. **d**) A scatterplot depicting the association between serum testosterone and the genus *Acinetobacter* relative abundance among individuals with different genotypes (AA vs GA/GG) at the top locus rs139799338. Spearman correlation's R and p values were also listed. **e**) Boxplots showing the average relative abundance of genus *Acinetobacter* among individuals with different genotypes (AA vs GA/GG) at the top locus rs4241281, as well as stratified by male and female samples. **f**) A scatterplot illustrating the association between serum testosterone and the genus *Acinetobacter* relative abundance among individuals with different genotypes (AA vs GA/GG) at the top locus rs4241281. Spearman correlation's R and p values were also listed.

a

position	total	overlap
saliva	3222	75
gut	1754	75
vagina	686	75
urine	1579	75

b

gut · saliva · urine · vagina

pairs	R2	p.value	p.adjusted
gut vs urine	0.2788688	0.001	0.001
saliva vs urine	0.3617572	0.001	0.001
vagina vs urine	0.1112906	0.001	0.001

c

gut · saliva · urine · vagina

pairs	R2	p.value	p.adjusted
gut vs urine	0.27824331	0.001	0.001
saliva vs urine	0.30737010	0.001	0.001
vagina vs urine	0.05148467	0.001	0.001

d

e