The dynamic climate adaptation of indicator warbler species complex in the

Pacific Rainforest

Silu Wang
Department of Biological Sciences
University at Buffalo
211 Mary Talbert Way, Buffalo, NY, 14260

Data management plan:

All the data involved in this study will be uploaded to Dryad.

Abstract

Understanding the functional responses of ecosystems to climate change is critical to predict and conserve global biodiversity. The evolution of indicator species provides an efficient avenue towards the functional biome response to climate change. Leveraging decades of citizen science data coupled with high-resolution spatiotemporal climate data, I dissected the climate change response of hybridizing *Setophaga* warblers, indicator species breeding in the mysterious canopy of the Pacific Rainforest. Breeding habitat temperature and precipitation powerfully predicted the breeding occupancy of the northern (*S. townsendi*) and the southern species (*S. occidentalis*). Both species showed positive climatic responses in the recent decade when the recent breeding occupancy was greater than the expected contingency. This implies the potential climate adaptation or life history plasticity in the breeding warbler populations. However, the *S. occidentalis* showed a compromised climate response in 2000-10, when the predicted breeding occupancy was significantly lower than the expected contingency. The compromised response might be due to rampant decline of July

precipitation in 2000-10, compared to previous decades. Although the July precipitation

continued to decline in 2010-20, *S. occidentalis* showed signs of recovery, after

generations of climate adaptation. I further evaluated their breeding niche competition,

reflected by the overlap of breeding occupancy between species. I found that the

competition potential was the lowest in 2000-10 when both species were at the trough

of breeding occupancy, which recovered in 2010-20. This study shed light on eco-evo

feedback in the indicating species complex and illuminated the functional response of

the rainforest ecosystem to climate change.

40   **Keywords:**

41   Climate change, Climate adaptation, plasticity, wood warbler, hybridization, evolution,

42   temperate rainforest, *Setophaga,* niche competition

Introduction

In the catastrophic climatic change (Kemp et al. 2022) and biodiversity loss (Brook et al. 2003; Dirzo and Raven 2003), efficient monitoring of the potential and trend in organismal response is crucial for conservation and restoration practice (Mawdsley et al. 2009; Dawson et al. 2011; Watson et al. 2012). The accruing citizen science data of organism occurrence (Johnston et al. 2023) and high-resolution spatiotemporal climate data collectively empower us to uncover organismal response to climate change.

The indicator species of vulnerable ecosystems are crucial stepping-stones for understanding the functional response of the ecosystems to climate change (Pearman et al. 2011; Siddig et al. 2016; Bal et al. 2018; Terrigeol et al. 2022). Here I leverage the indicating power of the wood warblers that breed in the canopy of the temperate rainforest of western North America as an avenue toward understanding the rainforest's functional response to climate change.

The sister species, *Setophaga occidentalis* (SOCC) and *S. townsendi* (STOW) are indicator species of the old-growth rainforest (Carpenter et al. 2014) in western North America (Matsuoka et al. 1997; Huang 2013; Bielski et al. 2024). The temperate rainforest of Western North America has been experiencing unprecedented ecological vulnerability related to climate change (Coops and Waring 2011; Hicke et al. 2013; Scheller et al. 2018; Buotte et al. 2019; Goss et al. 2020). The breeding warblers hold nesting territories in the canopy of coniferous trees, such as Douglas Fir, Western Hemlocks, and Western Red Cedars (Barlow 1899), and their populations reflect the

functional ecology of the temperate rainforest in Western North America (Bielski et al.

2024).

SOCC and STOW breed in divergent climatic conditions (Wang et al. 2021) yet

hybridize and compete extensively at breeding overlaps (Rohwer and Wood 1998;

Pearson 2000; Pearson and Manuwal 2000; Wang et al. 2019, 2020). The local

mitonuclear ancestries of SOCC and STOW have been shown to covary with climatic

variation in their breeding habitats (Wang et al. 2021). Whether SOCC and STOW are

adaptively responding to the rapidly changing climatic conditions over the past decades

remains an open question. Further, whether species-specific climatic response affects

their breeding niche competition remains unknown, thus hindering the understanding of

eco-evo feedback in rainforest climate change response.

I investigated the breeding warbler responses to climate change by comparing

their expected breeding occupancy based on historical contingency and present

breeding occupancy. If the warblers are positively responding to climate change, the

present breeding occupancy should be greater than the expected breeding occupancy

based on historical contingency. Evaluating the wood warbler's response to breeding

habitat climate change would shed light on the potential of climate adaptation of the

temperate rainforest ecosystem. Further, I evaluated the effect of climate change on

species boundaries as SOCC and STOW hybridize and compete upon contact (Rohwer

and Wood 1998; Pearson and Manuwal 2000; Wang et al. 2019, 2020; de Zwaan et al.

2022). The geographic overlap in the predicted breeding occupancy of SOCC and

STOW reflects the potential niche competition between the hybridizing species. A

greater overlap suggests a stronger potential for competition in their breeding niches.

Collectively, this time-series modeling synthesis sheds light on organismal, species

boundaries, and ecosystem responses to climate change.

Fig. 1 A schematic illustration of the tests of Breeding Range Shift and
Organismal Response to Climate Change. Organismal Response to Climate
Change: Step 1: calculate *expected breeding contingency* (*Exp*) with the historical
model (e.g. $m_{[1990\text{-}2000]}$) and the breeding climate of the current period (e.g. 2000-2010).
Step 2: Calculate *predicted breeding occupancy* (*Pred*) with the recent model (e.g.,
$m_{[2000\text{-}2010]}$). **Step 3**: compare *Pred* and *Exp.* A greater *Pred* than *Exp* reflects a positive
response to climate change (e.g. climate adaptation). **Breeding Range Shift:** compare
Pred within species across decades (e.g., 1990-2000 vs 2000-2010).
99

Methods
Breeding Site Climate

I downloaded the climate data from three decades: 1990-2000, 2000-2010, and

2010-2020 for western North America from WorldClim (Fick and Hijmans 2017; Harris et

al. 2020). To represent the total climatic conditions at the breeding sites, I incorporated

all the monthly climate data from CRU-TS-4.06 by using WorldClim 2.1 for bias

correction with a spatial resolution of 2.5 minutes (~21 km^2 at the equator). At this

temporal and spatial resolution, the climate variables available were: (1) average

minimum temperature (°C), (2) average maximum temperature (°C), and (3) total

precipitation (mm).

Warbler breeding occurrence

I downloaded all the historical occurrences for *Setophaga occidentalis* and *S.*

townsendi from GBIF.org (07 January 2024, GBIF.Org User 2024a,b). To focus on

breeding individuals, I extracted occurrences for June each year between 1990 to 2020.

To alleviate any identification error from the inexperienced observers, I excluded data

from observers with less than 3 submission records.

Breeding occupancy model

To train the breeding occupancy model for each decade (i.e. $m_{[1990-2000]}$, $m_{[2000-2010]}$, $m_{[2010-2020]}$), I used the breeding season climate data to predict species

presence/absence for each decade. The presence data was taken from the filtered

breeding occurrence data from GBIF (see above). For absence data, I randomly

generated pseudo-absence points (of equal sample size to the presence data) within

the square of -155° to -105° W and 30° to 65° N. The collection presence/absence data

was randomly divided into 5 folds so that ⅘ of it was used to train the occupancy model

and the remainder ⅕ of the data was saved as testing data. With the training data, I

fitted the binomial generalized linear model in which the presence/absence is predicted

by the interactions of all three climate variables. With the testing data for each model, I

calculated the corresponding threshold probability of breeding presence with

pa_evaluate function in the *predicted* package in R (Fielding and Bell 1997; Liu et al.

2011). I evaluated the model prediction accuracy as the following: (true positive + true

negative)/total testing size.

Breeding warbler response to climate change

To evaluate whether the wood warblers are responding to climate change, I

calculated the *expected breeding continency* of the future decade (e.g. 2000-2010 or

2010-2020) with a historical occupancy model ($m_{[1990-2000]}$) and climate input from the

respective future decade (**Fig. 1**). At the same time, I calculate the *predicted breeding*

occupancy with the species distribution models fitted with the climatic variation and

species occurrence of each future decade. I compared the *expected breeding*

contingency (*Exp*) and *predicted breeding occupancy* (*Pred*) within 2000-10 or 2010-20.

A smaller value in the former than in the latter could suggest a positive response to

climate change (e.g. signature of climate adaptation).

To test whether there was a significant difference between *expected breeding*

contingency (*Exp*) and *predicted breeding occupancy* (*Pred*), I generated 1000

bootstrap samples each with 10,000 randomly sampled localities. For each bootstrap

sample, I calculated *occupancy probability* as the number of localities with *breeding*

presence (see above) over the total number of localities. The same set of localities

applied for *Exp* and *Pred* extraction. With all the bootstrap samples, I ran a linear mixed

effect model with the response variable being breeding occupancy probability, fixed

effect being "Exp vs pred", and random effect being bootstrap sample IDs. I compared

the full model to the reduced model without the fixed effect to evaluate whether *Exp* and

Pred were significantly different.

To further understand any potential knock-on effect on the breeding niche

competition between SOCC and STOW, I compared the overlap of breeding occupancy

across decades. All the analysis was conducted in R (R Core Team 2023). **Results**

For all three decades, the climate variables significantly predicted the breeding

distribution of SOCC ($R^2 \approx 0.41$, accuracy ≈ 0.8, **Table 1**) and STOW ($R^2 \approx 0.8$,

accuracy ≈ 0.96, **Table 1**).

Table 1 A summary of the breeding occupancy models across decades. Climate
variables significantly explain breeding occupancy in STOW and SOCC in each decade
$m_{[1990-2000]}$, $m_{[2000-2010]}$, $m_{[2010-2020]}$.

STOW	Coef$_{1990-2000}$	p-value$_{1990-2000}$	Stats	Coef$_{2000-10}$	p-value$_{2000-10}$	Stats	Coef$_{2010-20}$	p-value$_{2010-20}$	Stats
Tmin	0.9742	$< 10^{-15}$	$R^2 = 0.42$	1.2020	$< 10^{-15}$	$R^2 = 0.47$	0.9935	$< 10^{-15}$	$R^2 = 0.42$
Tmax	-0.6752	$< 10^{-15}$		-0.8246	$< 10^{-15}$		-0.6907	$< 10^{-15}$	
Precipitation	-0.0656	$< 10^{-15}$	Accuracy	-0.0650	$< 10^{-15}$	Accuracy	-0.0734	$< 10^{-15}$	Accuracy
Tmin*Tmax	-0.0212	$< 10^{-15}$	= 0.83	-0.0228	$< 10^{-9}$	= 0.84	-0.0193	$< 10^{-15}$	= 0.82
Tmin*Prec	-0.0007	0.026		-0.0015	0.0500		0.0000	0.902	
Tmax*Prec	0.0079	$< 10^{-15}$		0.0075	$< 10^{-15}$		0.0087	$< 10^{-15}$	
Tmin*Tmax*Prec	-0.0005	$< 10^{-15}$		-0.0005	$< 10^{-10}$		-0.0006	$< 10^{-15}$	
SOCC	**Coef$_{1990-2000}$**	**p-value$_{1990-2000}$**	**Stats**	**Coef$_{2000-10}$**	**p-value$_{2000-10}$**	**Stats**	**Coef$_{2010-20}$**	**p-value$_{2010-20}$**	**Stats**
Tmin	2.4570	$< 10^{-15}$	$R^2 = 0.82$	1.1610	$< 10^{-15}$	$R^2 = 0.84$	2.6206	$< 10^{-15}$	$R^2 = 0.81$
Tmax	-0.6430	$< 10^{-15}$		-1.3090	$< 10^{-15}$		-0.6761	$< 10^{-15}$	
Precipitation	-0.1322	$< 10^{-15}$	Accuracy	-0.2959	$< 10^{-15}$	Accuracy	-0.1336	$< 10^{-15}$	Accuracy
Tmin*Tmax	-0.0991	$< 10^{-15}$	= 0.97	0.0093	$< 10^{-5}$	= 0.96	-0.0955	$< 10^{-15}$	= 0.97
Tmin*Prec	-0.0234	$< 10^{-15}$		-0.0087	$< 10^{-13}$		-0.0230	$< 10^{-15}$	
Tmax*Prec	0.0165	$< 10^{-15}$		0.0298	$< 10^{-15}$		0.0167	$< 10^{-15}$	
Tmin*Tmax*Prec	0.0005	$< 10^{-5}$		-0.0011	$< 10^{-10}$		0.0004	0.0003	

161
162
163

Table 2 The change in predicted breeding occupancy (*Pred*) of SOCC, STOW, and their breeding overlap in the past decades. Linear mixed effect model of the predicted presence among decades. The prediction was made with the occupancy model of each decade: $m_{[1990-2000]}$, $m_{[2000-2010]}$, $m_{[2010-2020]}$.

Species	Periods	Difference	95%CI	*p*-value	Mixed Effect Model
SOCC	(1990-2000)-(2000-2010)	**0.00619**	(0.00611, 0.0063)	$p < 10^{-15}$	
	(1990-2000)-(2010-2020)	**0.00318**	(0.00312, 0.00324)	$p < 10^{-15}$	F = 174291
	(2000-2010)-(2010-2020)	**-0.00301**	(-0.00306, -0.00295)	$p < 10^{-15}$	$p < 10^{-15}$
STOW	(1990-2000)-(2000-2010)	**0.01085**	(0.01073, 0.01098)	$p < 10^{-15}$	
	(1990-2000)-(2010-2020)	**-0.00944**	(-0.00956, -0.00932)	$p < 10^{-15}$	F = 496044
	(2000-2010)-(2010-2020)	**-0.02029**	(-0.02042, -0.02016)	$p < 10^{-15}$	$p < 10^{-15}$
Overlap	(1990-2000)-(2000-2010)	**0.00082**	(0.00811, 0.00823)	$p < 10^{-15}$	
	(1990-2000)-(2010-2020)	**-0.00180**	(-0.00187, -0.00179)	$p < 10^{-15}$	F = 71545
	(2000-2010)-(2010-2020)	**-0.01000**	(-0.01006, -0.00994)	$p < 10^{-15}$	$p < 10^{-15}$

I observed the lowest predicted breeding occupancy in 2000-10 for both SOCC and STOW (**Table 2**). The predicted breeding overlap between SOCC and STOW was also the lowest in 2000-10 (**Table 2**, **Fig. 5**). I detected the signature of breeding climate adaptation in 2010-20 for both species (**Fig. 2, 3; Table 3**) and STOW-only in 2000-10 (**Fig. 2; Table 3**). Specifically, in 2000-10, *Pred* was significantly greater than *Exp* in STOW (**Fig. 2 B, E, G**), but the reverse was true for SOCC (Table 3, **Fig. 3 B, E, G**). There was greater *Pred* than *Exp* in 2010-2020 for both STOW (**Table 3**; **Fig. 2 C, F, H**) and SOCC (**Table 3**; **Fig. 3 C, F, H**).

Table 3 The difference in the predicted probability between the expected contingency (*Exp*) and predicted occupancy (*Pred*) of STOW or SOCC in 2000-10 and 2010-20. Significant and negative mean differences between *Exp* and *Pred* indicate population potential of climate adaptation during the decade reflects a positive response to climate change. *Exp* calculated by the historical occupancy model ($m_{[1990-2000]}$) and the climate variation in the respective decade (2000-10 or 2010-20). *Pred* is inferred by the occupancy model ($m_{[2000-10]}$ or $m_{[2010-20]}$) in the respective decade.

Species	Probability	Mean	95%CI	T_{9999}	*p*-value
STOW	p2000-10 (exp - pred)	-0.0345	(-0.0349, -0.0341)	-164.71	< 2.2e-16
	p2010-20 (exp - pred)	-0.0038	(-0.0042, -0.0034)	-18.47	< 2.2e-16
SOCC	p2000-10 (exp - pred)	0.0104	(0.0102, 0.0106)	95.74	< 2.2e-16
	p2010-20 (exp - pred)	-0.0065	(-0.0067, -0.0063)	-52.97	< 2.2e-16

Fig. 2 The greater predicted breeding occupancy (*Pred*) than expected continency (*Exp*) indicates a positive response to climate change in *Setophaga townsendi* (STOW). **B**, **E**, **G**: In 2000-10, the *Pred* (**E**) is greater than *Exp* (**B**). **C**, **F**, **H**: A similar pattern was observed in 2010-20, where the predicted breeding occupancy (**F**) is less than the expected contingency (**C**).

Fig. 3 The heterogeneous response to climate change in *Setophaga occidentalis* (SOCC), as reflected by the difference between predicted breeding occupancy (*Pred*) and expected continency (*Exp*). B, E, G: In 2000-10, the *Pred* (**E**) is less than *Exp* (**B**). **C, F, H**: an opposite pattern was observed in 2010-20, where the *Pred* (**F**) was greater than *Exp* (**C**), which indicates climate adaptation in this decade.

204
205 **Fig. 4 Change in mean July precipitation over decades. A**, July mean precipitation
206 (mm) over each decades from 1960-70 to 2010-20. **B**, violin plot showing the change of
207 precipitation with each line being a location. There was a significant reduction of
208 precipitation after 2000 (repeated measure ANOVA, post-hoc paired T-test with
209 Bonferroni-corrected $p < 0.05$).

215
216 **Fig. 5 The change in the potential of breeding niche competition between SOCC**
217 **and STOW reflected by the overlap of predicted breeding occupancy of over three**
218 **decades.** The lowest predicted overlap occurred in 2000-10, when July precipitation
219 drastically declined.

Discussion

Understanding the rainforest ecosystem response to climate change is central to

global biodiversity conservation, as rainforests harbor the majority of terrestrial

biodiversity (McMahon et al. 2011). An efficient and powerful predictive framework for

dissecting rainforest indicator species' response to climate change is essential. The rich

citizen science data for the *Setophaga* warblers that breed in the temperate rainforest of

North America empowered us to unravel the functional response of the rainforest to

climate change. I observed a positive response to climate change in both species, as

the recent breeding occupancy was greater than the expectation based on historical

contingency (**Fig. 2, 3**). This pattern indicates breeding plasticity or adaptation to the

changing climatic conditions in the breeding habitat. The decadal trend in the indicator

species of the temperate rainforest could indicate the functional adaptability of the

rainforest ecosystem to climate change.

The climate variables significantly explained the breeding distribution of both

species (**Table 1**) with a strong effect on the breeding distribution of southern species

(SOCC, $R^2 \sim 0.8$; **Fig. 3**) and a moderate effect on the northern species (STOW, $R^2 \sim$

0.4; **Fig. 2**). A caveat of our prediction framework is that it is based on limited climatic

variables -- temperature and precipitation. Additional climate factors could elevate the

predicting power. Despite that, the variation explained by two key climate factors was

remarkable, especially for SOCC breeding occupancy.

More latent factors could contribute to STOW breeding occupancy, which spans

over heterogeneous climatic conditions. The coastal STOW populations harbor

mitonuclear ancestry from SOCC (Krosby and Rohwer 2009; Wang et al. 2021). The

divergent SOCC and inland STOW mitonuclear ancestries have been associated with

lipid metabolism in divergent climate adaptation (Wang et al. 2021). The predicted

STOW breeding occupancy in the SOCC breeding range in southern Oregon and

California (**Fig. 2**) could be attributed to the fact that coastal STOW contains

mitonuclear variants of SOCC. The elevated genetic diversity from ancient hybridization

in coastal British Columbia and Alaska (Wang et al. 2021) could predispose STOW with

greater adaptive potential. Future studies could further dissect the heterogeneity in the

breeding distribution of STOW concerning metabolic genotypes.

The excessive breeding occupancy than expected based on historical climatic

reaction contingency could reflect climate adaptation. If so, the genetic variants suitable

for the recent climatic conditions could have increased in frequency in the recent

decade resulting in change in the niche adaptability. Alternatively, the observed positive

response could be explained by breeding plasticity within the breeding niche potential.

The historical climate-occupancy contingency may only represent the historical realized

niche within a greater niche potential. However, the niche potential could also evolve

and track climate change. Future studies could differentiate the possibilities by

investigating any change in genetic frequency related to climate adaptation.

It is alarming that SOCC showed negative response to climate change in 2000-

10 when the breeding occupancy was lower than expected based on niche contingency

(**Fig. 3 G**). This pattern is consistent with their recent negative population trend of -

0.961 (95% CI: -1.675 to -0.285) (Sauer et al. 2020). Since annual precipitation is highly

predictively of SOCC breeding occupancy (Table 1), the plummeting July precipitation

since 2000-10 (**Fig. 4**) could be especially harmful. Despite that the July precipitation

did not recover in 2010-20, SOCC showed signs of recovery, where the breeding

occupancy surpassed expected breeding contingency. Further monitoring is need to

determine whether this reversal could halt SOCC population decline.

The potential of breeding habitat competition between the hybridizing STOW and

SOCC could be reflected by the overlap of their predicted breeding occupancy. The two

warbler species are ecologically similar during breeding season as specialists in the

canopy of coniferous forests (Morrison 1982). In the hybrid zones, interspecific

competition does occur besides hybridization (Pearson 2000; Pearson and Manuwal

2000; de Zwaan et al. 2022). The predicted breeding overlap covers much of the

breeding range of SOCC indicating that the absence of STOW breeding occupancy in

California and Southern Oregon could be due to breeding niche competition with SOCC.

By leveraging citizen science data from three decades, I conducted a modeling

synthesis for understanding the breeding wood warbler response to climate change

which illuminates the functional reaction of the temperate rainforest of North America.

Overall, the indicator bird species demonstrated positive responses to climate change,

despite the challenges represented by the rampant decline of July precipitation since

2000. This result reflects the climate adaptation of the hermit warblers, rendering hope

for the temperate rainforest ecosystem. However, more ecological factors in different

trophic levels need to be further evaluated to fully understand the ecosystem response

to climate change. Future investigation of the physiological, behavioral, and

phenological mechanisms underlying the organismal response to climate change could

advance our understanding of climate adaptation in rainforest biomes, where most

terrestrial biodiversity resides.

289
290

References
Bal, P., A. I. Tulloch, P. F. Addison, E. McDonald-Madden, and J. R. Rhodes. 2018.

Selecting indicator species for biodiversity management. Frontiers in Ecol &

Environ 16:589–598.

Barlow, C. 1899. Nesting of the Hermit Warbler in the Sierra Nevada Mountains,

California. The Auk 16:156–161.

Bielski, L., C. Cansler, K. McGinn, M. Z. Peery, and C. Wood. 2024. Can the Hermit

Warbler (Setophaga occidentalis) serve as an old-forest indicator species in the

Sierra Nevada? JFO 95:art4.

Brook, B. W., N. S. Sodhi, and P. K. L. Ng. 2003. Catastrophic extinctions follow

deforestation in Singapore. Nature 424:420–423.

Buotte, P. C., S. Levis, B. E. Law, T. W. Hudiburg, D. E. Rupp, and J. J. Kent. 2019.

Near-future forest vulnerability to drought and fire varies across the western

United States. Global Change Biology 25:290–303.

Carpenter, D. N., J. G. Bockheim, and P. F. Reich. 2014. Soils of temperate rainforests

of the North American Pacific Coast. Geoderma 230–231:250–264.

Coops, N. C., and R. H. Waring. 2011. Estimating the vulnerability of fifteen tree species

under changing climate in Northwest North America. Ecological Modelling

222:2119–2129.

Dawson, T. P., S. T. Jackson, J. I. House, I. C. Prentice, and G. M. Mace. 2011. Beyond

Predictions: Biodiversity Conservation in a Changing Climate. Science 332:53–

58.

de Zwaan, D. R., J. Mackenzie, E. Mikkelsen, C. Wood, and S. Wang. 2022. Pleiotropic

opposing dominance within a color gene block contributes to a nascent species

boundary via its influence on hybrid male territorial behavior. PNAS Nexus

1:pgac074.

Dirzo, R., and P. H. Raven. 2003. Global State of Biodiversity and Loss. Annu. Rev.

Environ. Resour. 28:137–167.

Fick, S. E., and R. J. Hijmans. 2017. WorldClim 2: new 1-km spatial resolution climate

surfaces for global land areas. Intl Journal of Climatology 37:4302–4315.

Fielding, A. H., and J. F. Bell. 1997. A review of methods for the assessment of

prediction errors in conservation presence/absence models. Envir. Conserv.

24:38–49.

GBIF.Org User. 2024a. Occurrence Download. [object Object].

GBIF.Org User. 2024b. Occurrence Download. [object Object].

Goss, M., D. L. Swain, J. T. Abatzoglou, A. Sarhadi, C. A. Kolden, A. P. Williams, and

N. S. Diffenbaugh. 2020. Climate change is increasing the likelihood of extreme

autumn wildfire conditions across California. Environ. Res. Lett. 15:094016.

Harris, I., T. J. Osborn, P. Jones, and D. Lister. 2020. Version 4 of the CRU TS monthly

high-resolution gridded multivariate climate dataset. Sci Data 7:109.

Hicke, J. A., A. J. H. Meddens, C. D. Allen, and C. A. Kolden. 2013. Carbon stocks of

trees killed by bark beetles and wildfire in the western United States. Environ.

Res. Lett. 8:035032.

Huang, A. 2013. Generalists and Specialists Warblers and Their Conservation Statuses

in North America. Master's Thesis. University of British Columbia.

Johnston, A., E. Matechou, and E. B. Dennis. 2023. Outstanding challenges and future

directions for biodiversity monitoring using citizen science data. Methods Ecol

Evol 14:103–116.

Kemp, L., C. Xu, J. Depledge, K. L. Ebi, G. Gibbins, T. A. Kohler, J. Rockström, M.

Scheffer, H. J. Schellnhuber, W. Steffen, and T. M. Lenton. 2022. Climate

Endgame: Exploring catastrophic climate change scenarios. Proc. Natl. Acad.

Sci. U.S.A. 119:e2108146119.

Krosby, M., and S. Rohwer. 2009. A 2000 km genetic wake yields evidence for northern

glacial refugia and hybrid zone movement in a pair of songbirds. Proc. R. Soc. B.

276:615–621.

Liu, C., M. White, and G. Newell. 2011. Measuring and comparing the accuracy of

species distribution models with presence-absence data. Ecography 34:232–243.

Matsuoka, S. M., C. M. Handel, and D. D. Roby. 1997. Nesting Ecology of Townsend's

Warblers in Relation to Habitat Characteristics in a Mature Boreal Forest. The

Condor 99:271–281.

Mawdsley, J. R., R. O'Malley, and D. S. Ojima. 2009. A Review of Climate-Change

Adaptation Strategies for Wildlife Management and Biodiversity Conservation.

Conservation Biology 23:1080–1089.

McMahon, S. M., S. P. Harrison, W. S. Armbruster, P. J. Bartlein, C. M. Beale, M. E.

Edwards, J. Kattge, G. Midgley, X. Morin, and I. C. Prentice. 2011. Improving

assessment and modelling of climate change impacts on global terrestrial

biodiversity. Trends in Ecology & Evolution 26:249–259.

Morrison, M. 1982. The Structure of Western Warbler Assemblages: Ecomorphological

Analysis of the Black-Throated Gray and Hermit Warblers. The Auk 99:503–513.

Pearman, P. B., A. Guisan, and N. E. Zimmermann. 2011. Impacts of climate change on

Swiss biodiversity: An indicator taxa approach. Biological Conservation 144:866–

875.

Pearson, S. F. 2000. Asymmetries in male aggression across an avian hybrid zone.

Behavioral Ecology 11:93–101.

Pearson, S. F., and D. A. Manuwal. 2000. Influence of Niche Overlap and Territoriality

on Hybridization Between Hermit Warblers and Townsend's Warblers. The Auk

117:175–183.

R Core Team. 2023. R: A language and environment for statistical omputing. R

Foundation for Statistical   Computing, Vienna, Austria.

Rohwer, S., and C. Wood. 1998. Three Hybrid Zones between Hermit and Townsend's

Warblers in Washington and Oregon. The Auk 115:284–310.

Sauer, J. R., W. A. Link, and J. E. Hines. 2020. The North American breeding bird

survey, analysis results 1966–2019. U.S. Geological Survey, Laurel, Maryland,

USA.

Scheller, R. M., A. M. Kretchun, E. L. Loudermilk, M. D. Hurteau, P. J. Weisberg, and C.

Skinner. 2018. Interactions Among Fuel Management, Species Composition,

Bark Beetles, and Climate Change and the Potential Effects on Forests of the

Lake Tahoe Basin. Ecosystems 21:643–656.

Siddig, A. A. H., A. M. Ellison, A. Ochs, C. Villar-Leeman, and M. K. Lau. 2016. How do

ecologists select and use indicator species to monitor ecological change?

Insights from 14 years of publication in Ecological Indicators. Ecological

Indicators 60:223–230.

Terrigeol, A., S. Ewane Ebouele, M. Darveau, C. Hébert, L.-P. Rivest, and D. Fortin.

2022. On the efficiency of indicator species for broad-scale monitoring of bird

diversity across climate conditions. Ecological Indicators 137:108773.

Wang, S., M. J. Ore, E. K. Mikkelsen, J. Lee-Yaw, D. P. L. Toews, S. Rohwer, and D.

Irwin. 2021. Signatures of mitonuclear coevolution in a warbler species complex.

Nat Commun 12:4279.

Wang, S., S. Rohwer, D. R. De Zwaan, D. P. L. Toews, I. J. Lovette, J. Mackenzie, and

D. Irwin. 2020. Selection on a small genomic region underpins differentiation in

multiple color traits between two warbler species. Evolution Letters 4:502–515.

Wang, S., S. Rohwer, K. Delmore, and D. E. Irwin. 2019. Cross-decades stability of an

avian hybrid zone. J of Evolutionary Biology 32:1242–1251.

Watson, J. E. M., M. Rao, K. Ai-Li, and X. Yan. 2012. Climate Change Adaptation

Planning for Biodiversity Conservation: A Review. Advances in Climate Change

Research 3:1–11.

397