History of recurrent short and long-term coral bleaching events in Indian coral reefs: a

systematic review of contrasting bleaching patterns, lessons learned, and future directions

Thinesh Thangadurai[1,2*], Kalyan De[3*], Sobana Murugesan[1], Sivagurunathan P[4], Riana Peter[4], Ramasamy

Pasiyappazham[5], Joseph Selvin[1], Polpass Arul Jose[6], Anthony Bellantuono[7]

[1]Department of Microbiology, Pondicherry University, Puducherry, India.

[2]Current affiliation - Centre for Ecological Sciences, Indian Institute of Science, Bangalore, India.

[3] Biological Oceanography Division, CSIR-National Institute of Oceanography, Goa, India.

[4]Department of Microbiology, Annamalai University, Chidambaram, Tamil Nadu, India.

[5]Department of Physiology, Saveetha Universty, Chennai, Tamil Nadu, India.

[6]Madurai Kamaraj University, India. Current: Department of Biological Science, Harvard University, USA.

[7]Department of Biological Sciences, Florida International University, Miami, Florida. USA.

[*]Corresponding Authors: thina.sathesh@gmail.com. Co – Correspondence - kalyandeaqua@gmail.com

[*] Equally contributed

Abstract

Climate change has intensified coral bleaching events, leading to widespread coral mortality and

threatening the coral reef survival worldwide. However, species susceptibility patterns to bleaching differ

spatially and temporally, making it crucial to understand bleaching impact and species susceptibility

patterns across latitudinal ranges, particularly from understudied locations. We critically reviewed 23 years

of bleaching episodes on Indian coral reefs (four major reefs and other patch reefs) reported from 1998 to

2020 to understand the geographical footprint of bleaching patterns, species susceptibility differences, and

their impact. We found that all four major Indian reef systems (Gulf of Kachchh, Gulf of Mannar,

Lakshadweep, and the Andaman Islands) have experienced three major bleaching episodes (1998, 2010,

and 2016) and multiple short-term bleaching events. Short-term bleaching events (DHWs> four weeks)

created differences in species susceptibility, but the disparity among species diminished to long-term

bleaching events (DHWs <4 weeks) in all reef sites. *Acropora*, *Porites*, *Pocillopora*, *Montipora,* and

Turninaria in the Gulf of Mannar, *Porites,* and *Favia* in the Gulf of Kachchh, and *Acropora* in the Andaman

and Lakshadweep were the most vulnerable genera during all the bleaching events. In terms of recovery,

Porites and other massive forms recovered better than branching forms. Furthermore, species susceptibility

pattern analysis (winners vs. losers) to 2016 long-term bleaching events found conflicting results from

studies (one-time surveys) in the same area. This result indicates that a one-time survey (survey timing)

influences results as bleaching events lengthen, necessitating repeated surveys to determine winners and

losers. Although inconsistent data from different study sites makes it difficult to create predictive models,

data collected over the past 23 years provides critical insight into coral community susceptibility patterns

and bleaching impacts, underscoring the need for systematic, uniform surveys in the future.

Keywords: Coral bleaching, Indian Ocean, Contrasting bleaching pattern, Gulf of Mannar, Andaman, Gulf

of Kachchh.

Introduction

Increasing coral bleaching frequency threatens the persistence of coral reef ecosystems, which provide livelihoods, food security, and coastal protection to over 280 million people worldwide (Moberg and Folke, 1999, Hughes et al., 2018). Coral bleaching is a multifaceted environmental problem that can arise due to various factors, such as increased seawater temperatures, extremely low or cold temperatures, salinity depletion, pollution, ocean acidification, and excessive exposure to sunlight (Hughes et al., 2018). However, the primary cause of coral bleaching is identified as increased episodes of marine heatwaves due to ongoing climate change. These climate-induced coral bleaching events occur when reef-building corals expose to a few degrees above the average summer temperature, causing a break down in the symbiotic relationship between coral and their endosymbiotic algal partners called *Symbiodinacea* (LaJeunesse et al., 2018, Hughes et al., 2018). This symbiotic relationship between corals and their intracellular algal symbiosis is the engine of the reef that forms the basis of its highly diverse ecosystem structure. Bleached autotrophic corals are physiologically deprived of regular food supply in the absence of their algal partner (Davy et al., 2012), and prolonged bleaching events led to extensive coral die-off (Glynn et al., 2001; González-Pech et al., 2021; Hughes et al., 2017b). Unfortunately, coral bleaching episodes have become more frequent and severe as the number of warmest years has increased, with nine of the ten warmest years occurring since 2005 (Bellwood et al., 2019). These frequent bleaching events reduced recovery intervals from bleaching episodes (Hughes et al., 2018). Coral bleaching events have recently shifted from average heat stress exposures of 4 to 12 °C-weeks to exceeding 24-35 °C-weeks in some reef locations (Hoegh-Guldberg 2011, Claar et al., 2018; Boyle et al., 2017; Brainard et al., 2018). The 2014-2017 global coral-bleaching event caused widespread mortality of corals and other reef-associated organisms over thousands of square kilometers (Eakin et al., 2019). Furthermore, bleaching events diminish coral recovery potentials such as reproduction, larval succession, and growth rates, which result in reduced species diversity, richness, and dynamics (Albright and Langdon, 2011; Albright and Mason, 2013; Anthony et al., 2008; Fordyce et al., 2019; Prada et al., 2017). Climate models have predicted future bleaching events with greater frequency and severity (more than 90% by 2030 and nearly all reefs by 2050) (Cai, 2014). To achieve better

prediction, climate models require historical data on sea surface temperature, light intensity, salinity,

nutrient level, species response, and other variables to identify patterns and relationships for coral bleaching

(Eakin et al., 2018, Liu et al 2018). Therefore, geographical footprints of bleaching frequency, intensity,

and their impact pattern at a broad geographical scale, particularly from the understudied locations, are

crucial to predict future coral community structure against the increasing thermal stress.

Accurate bleaching survey is important to capture differential species susceptibility patterns to

bleaching (winners and losers)

Although global warming has triggered unprecedented mass coral mortality across the ocean; the extent of

bleaching at different reef region varies due to differences in temperature magnitude, coral species

composition, past bleaching history, hydrodynamics, and several other local environmental factors

(Anthony et al., 2008; Hughes et al., 2018, 2003; Swain et al., 2016). In many instances, some corals tolerate

or recover better from bleaching than others in every geographical region including India (Kumaraguru,

2003; Manikandan et al., 2014; Pisapia and Burn, 2016; Thinesh et al., 2019a, De et al. 2022, 2023). In

India, studies are at a nascent state to identify the factors (symbiont types) responsible for differential

bleaching patterns (for example Thinesh et al., 2019; Mote et al., 2021). Generally, massive taxa such as

Porites spp. and some *Favidae* are more resistant to bleaching (Baird and Marshall, 2002). Massive coral

taxa can live in the bleaching state for longer and recover better than their branching counterparts (Grottoli

et al., 2014; Loya et al., 2001; Thinesh et al., 2019a). Historically the fast-growing branching taxa

(*Acropora* and *Pocillopora)* are thermally sensitive and susceptible to heat stress-driven bleaching (Putnam

et al., 2012). In contrast, slow-growing massive corals such as *Porites* spp. and some *Favidae* are considered

better tolerant to bleaching (Baird and Marshall, 2002, Darling et al., 2012). During the bleaching events,

massive coral taxa can live in the bleaching state for longer and recover better than their branching

counterparts, but branching corals can only stay bleached for a short time before they die or recover

(Grottoli et al., 2014; Loya et al., 2001; Thinesh et al., 2019a). This differential species-specific response

is influenced by a combination of factors, including microbiomes, types of symbionts, coral life-history

traits, tissue thickness, investment in energy storage, colony size, growth rate, protein content in the mucus,

coral species composition, past bleaching history, hydrodynamics (Berkelmans and van Oppen, 2006; Loya

et al., 2001; McClanahan et al., 2004, Anthony et al., 2008; Swain et al., 2016). These factors significantly

influence their susceptibility to bleach mortality and recovery in individual reefs, determining their future

community structure (Camp et al., 2018; Hughes et al., 2003). Therefore, data on coral community

composition, taxon-specific susceptibility, and resilience patterns to bleaching events are essential for every

reef location (Cornwell et al., 2021; Edmunds, 1994; Thinesh et al., 2019a, Dixon et al., 2022). Coral

bleaching survey is a widely used approach to record this pattern in the field. Therefore, surveying efforts

must be uniform to use the data for local and global scale comparison. During short-term bleaching

events (DHWs > 4), one-time surveys could document the complete bleaching impact as bleaching

windows narrowed. In cases of long-term bleaching events, repeated surveys are necessary to document the

impact of heat stress accurately, especially to determine the winners and losers (Heron et al., 2016, Danielle

et al., 2018). If repeated surveys are not feasible, a study must include DHWs for their locations, the

bleaching proportion of each coral genera, which allow us to validate and compare bleaching effects across

studies and regions (Claar et al., 2018).

In India, the first mass coral bleaching episode was reported in 1998. Since then, many surveys have

documented the bleaching episodes, impact, and species susceptibility pattern from multiple bleaching

events across different reef sites. As major reefs are separated geographically, most available bleaching

data across this 23-year timespan are patchy, limited to one-time surveys, or restricted to a single bleaching

episode from a specific location. The scattered data across reef sites hinder the understanding of species

susceptibility patterns and bleaching impact. In this study, we examined the available published research

on the thermal history, bleaching intensity, mortality, and species susceptibility pattern from 1998 to date;

(i) to create a comprehensive database of bleaching episodes in Indian reefs; (ii) to assess bleaching

percentage in respective to DHWs; iii) to identify susceptible and resistant coral genera during long-term

(DHWs > 4) and short-term bleaching events DHWs < 4 ; (iv) to highlight gaps in monitoring efforts; and

iv) to provide uniformed future survey recommendations.

Methods

Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) was adopted for the current review, using different scholarly databases, namely Scopus™, Web of Science™, and Google Scholar™, as these databases are robust and cover ecology, environmental studies, and oceanography. PRISMA is a published standard to conduct a systematic literature review, also suitable for the environmental management field because it clearly defines the research questions for a systematic review (Moher et al., 2009; Sierra-Correa and Cantera Kintz, 2015). We also carried out manual searches on several other journal databases such as Science Direct, Taylor Francis, Springer, and Sage, considering that they contain peer-reviewed journal articles related to the field of coral reefs. Primary literature (concerning Indian coral reefs and coral bleaching) published during the timeline, 1998 to 2021 was extracted using the following keywords: (coral reef OR tropical reef OR coral OR Scleractinia OR hard coral OR stony corals OR octocorals OR soft corals OR Coral bleaching OR Coral mortality Or climate change OR global warming OR temperature rise OR SST anomaly OR el-Nino OR la-Nina OR Indian reef OR Gulf of Kachchh OR Malvan Marine Sanctuary OR Grande Island OR Angriya bank Or Netrani Island OR Lakshadweep OR Palk Bay OR Gulf of Mannar OR Andaman Or Nicobar Or Bay of Bengal Or Arabian Sea. Only peer-reviewed articles were considered for the present review as the primary sources that offer empirical data to maintain both consistency and quality. A careful cross-referencing was also conducted to include any articles missed during the literature search. It should be noted that the review only focused on articles published in English. Other than that, only studies conducted in the Indian territorial water were selected because they are in line with the review's objective. Our search retrieved 210 articles from the databases at the first stage of the systematic review process. A thorough screening was conducted to remove duplicate reports. Based on the degree of relevance, individual papers were then eliminated by 'title' or 'abstract' alone, or by accessing the entire article. A total of 36 published research papers were reviewed as part of this effort. We avoided reports published in the newspapers as it has no data on species

164   susceptibility, percentage of bleaching, and mortality except the evidence of bleaching. Fig. 1 represents

165   the flow diagram for the PRISMA analysis.

167   **Fig. 1.** Flow diagram of the PIRSMA analysis (adapted from Moher et al., 2009)

169 **Fig. 2.** Bleaching area alert annual maximum a) 1998; b) 2010; c) 2016 (Data and Plot: NOAA coral reef

170 watch V3.1). d) Geographical distribution of the major Indian coral reefs with reported coral bleaching

171 episodes

Results

Coral reef distribution in India and the bleaching reports:

India has four major coral reef regions and a few patchy reefs. Major coral reefs are located in the i) Gulf

of Kachchh, ii) Gulf of Mannar biodiversity reserve, iii) Lakshadweep Islands, and iv) Andaman and

Nicobar Islands separated geographically. In addition, there are three patch reefs 1) Palk Bay, 2) Malvan

Marine Sanctuary, 3) Grande Islands, and a few patches at different locations as follows Ratnagiri, Redi,

Netrani, Quilon in the Kerala coast to Enayem in Tamilnadu, Parangipettai (Porto Novo), south of

Cuddalore, and Pondicherry (De et al., 2017). The estimated coral reef area in India is around 2,375 km^2

(DOD and SAC 1997). These coral reefs harbor a total of 585 species belonging to 108 genera and 23

families (De et al., 2020). Studies have documented a steady rise in SST anomaly and thermal stress in

Indian reefs (Arora et al., 2019, 2021b, De et al., 2022). The spatial distribution of the warmest SST, SST

anomaly, and DHWs data for each reef environment indicates that corals experienced thermal stress and

likely mass bleaching (DHWs $\geq$ 4) (Arora et al., 2019).

A history of bleaching events and their mortality:

Indian reefs experienced three long-term (major) and ten short-term (Minor) bleaching episodes within the

last 23 years. All three major bleaching events (1998, 2010, and 2016) affected all four major reefs, which

caused mortality. Ten short-term bleaching events are scattered among major reefs and patchy reefs out of

13 recorded bleaching events from seven reef sites. The Gulf of Mannar experienced many bleaching events

compared to other reef sites (Fig 3). The bleaching percentage and mortality report on bleaching events is

given in Table 1.

197

198 **Fig. 3.** (A) Several recorded bleaching events and their percentage from the major and patchy reefs. (B)

199 Observed bleaching-induced mortality events in seven Indian reefs (we used the highest mortality

200 percentage report for representation. Details are given in the Table 1.)

201 **Table 1:** Occurrence of different coral bleaching episodes in major coral reefs in Indian territorial

202 water

Coral reef regions	Year	Bleaching intensity	Bleaching impacted coral genera	Reference
Gulf of Kachchh	July 2016	3.9% coral bleaching	*Coscinaraea, Favia, Goniastrea, Goniopora, Leptastrea, Porites,* and *Turbinaria*	Arora et al., 2019
	September 2014	Not mentioned	*Favia, Favites,* and *Porites*	Adhavan et al. 2014
	May and June, 2010	60-70% coral bleaching	*Montipora explanata, Siderastrea savignyana, Coscinaraea spp., Goniopora spp., Porite spp.,Favia spp., Goniastrea spp., Hydnophora exesa, Symphyllia spp., Turbinaria peltata*	Joshi et al., 2014
	March 1998	30% coral mortality	Not mentioned	Wafar, 1999
	April-May 1998	11% coral bleaching	Not mentioned	Arthur, 2000

Malvan Marine Sanctuary	October 2014	14% coral bleaching	*Porites, Favites, Turbinaria, Pseudosiderastrea, Cyphastrea, Plesiastrea*	De et al., 2015
	December 2015	70.93% coral bleaching	*Pavona, Coscinaraea, Goniastrea, Favites, Favia, Cyphastrea, Leptastrea, Montastrea, Turbinaria, Goniopora, Porites.*	Raj et al., 2018
	May 2015, May 2016, May 2017, October 2018, April 2019	54.2% coral bleaching May 2015, 46.76% in May 2016, 20.22% in May 2017, 5.07% October 2018, 8.37% April 2019	*Porites lichen, Porites compressa, Favites melicerum, Turbinaria mesenterina, Pseudosiderastrea tayami, Cyphastrea serailia, Plesiastrea versipora, Goniopora spp, Siderastrea savignyana*	De et al., 2021
Grande Islands	October 2014, April 2015, November 2015, April 2016	6% and 4 % coral bleaching at two sites in October 2014; 7 % and 5.34 % in April 2015; 51 % and 50.15 % during November 2015, 7 % and 8% in April 2016	*Porites, Cyphastrea, Montastrea, Pseuodosiderastrea, Favia, Turbinaria and Leptastrea*	Hussain and Ingole, 2020
Gulf of Mannar	May 2020	38% to 57% coral bleaching	*Porite lutea, Porites solida, Goniastrea retiformis, Dipsastraea favus, D. speciose, Pocillopora damicornis, Montipora digitate, Favites halicora, F. complanata, Pavona sp, Favites halicora, F. complanata, Pavona sp.*	Sadhukhan et al. 2021

	May, 2019	-	*Acropora, Montipora, Porites, Pocillopora, Goniastrea, Platygyra, Dipsastraea, Turbinaria*	Ramesh et al., 2020
	2016	1.05±0.65% coral bleaching	*Sinularia, Xenia, and Subergorgia*	Boopathi et al., 2021
	April–May 2016	46.04% coral bleaching	*Acropora, Montipora, Symphyllia, Porites, Favia, Favites, Goniostrea, Leptastrea, Platygyra, Turbinaria, Pavona, Hydnophora, Pocillopora*	Krishnan et al., 2018
	March-June, 2016	23.92% coral bleaching	*Acropora, Montipora,* and *Pocillopora*	Edward et al., 2018
	February-May 2016	6% at Kurusadai island, 27 % at Manoli 60% at Pamban	*Porites* (34.9%), and *Acropora* (19.1%)	Ranith and Kripa, 2019
	April 2010	72% coral bleaching with 24.3% mortality	*Porites solida, P. lutea, Favia spp., Acropora cytherea, A. intermedia, A. Formosa, A. nobilis, Montipora digitata*	Manikandan et al., 2014
	2010	10% coral bleaching and <50% mortality	*Pocillopora damicornis, Acropora formosa, A. intermedia, A. nobilis, A. cytherea, Montipora digitata, M.foliosa, Favia sp. and Echinopora sp*	Edward et al., 2012
	April-June 2010	~57% coral bleaching at the Kurusadai Island during	*Porites sp.,* and *Acropora sp*	Jeevamani et al., 2013
	1998	60% coral mortality	Not mentioned	Wafar, 1999
	July 1998	89% coral bleaching and 23% mortality	Not mentioned	Arthur, 2000
Palk Bay	April to May 2019	85% coral bleaching	*Massive corals*	Sadhukhn et al. 2021
	April–May 2016	71.48% coral bleaching	*Porites, Favia, Leptastrea, Montastrea, Oulophyllia, Platygyra*	Krishnan et al., 2018

	May 2016	53.80% coral bleaching	*Acropora, Porites, Favia, Platygyra, Goniostrea, Symphyllia*	Thinesh et al., 2019b
	June 2014	53.6% coral bleaching	*Favites spp., Favites, Leptastrea* and *Porites spp*	Manikandan et al., 2016
	April 2010	75% coral bleaching with 32.2% mortality	*Porites solida, P. lutea, Favia spp., Acropora cytherea, A. intermedia, A. Formosa, A. nobilis, Montipora digitata*	Manikandan et al., 2014
	May-June 2010	24-39.6% coral bleaching	*Porites, Goniopora,* and *Favia*	Ravindran et al., 2012
	April-June 2002	50-60% coral bleaching followed by 2.7-4.5% mortality at Palk Bay	Bleaching occurred in *Acroporidae, Agariciidae, Faviidae, Poritidae, Oculinidae, Merulinidae, Mussidae, Pocilloporidae, and Dendrophyllidae* *Mortality occurred in Acropora formosa, A. digitifera, A. hyacinthus, Montipora digitata, M. foliosa, Pocilloproa verrucosa,* and *P. damicornis*	Kumaraguru, 2003
Lakshadweep Islands	2016	31% mortality	Not mentioned	Yadav et al. 2018
	2016	>60% coral bleaching	*Acropora* sp., and *Porites* sp.	Vineetha et al. 2018
	2010	65% coral bleaching	*Acropora* sp., and *Porites* sp. were severely impacted. Live coral cover decline from 36% to 26%	Idrees Babu and Suresh Kumar, 2016
	May-June 2010	73% coral bleaching	Bleaching-related mortality of sea anemones (87%) and giant clams (83%) at Agatti reported	Vinoth et al., 2012
	March 1998	43–87% mortality	Not mentioned	Wafar, 1999
	June 1998	82% bleaching and 26% mortality	Not mentioned	Arthur, 2000
Andaman Islands	2016	Not mentioned	*Acropora, Fungia*	Mohanty et al. 2017 a and b

July 2010	74 % coral bleaching at Havelock Island, 77% at Port Blair Bay	*Acropora cerealis, A. humilis, Montipora sp., Favia pallida, Diploastrea sp., Goniopora sp. Fungia concinna, Gardineroseries sp., Porites sp., Favites abdita and Lobophyllia robusta*	Marimuthu et al., 2013
April- May 2010	69.49 % coral bleaching at Havelock Island, 67.28% at South Button Island, 56.45 % at Nicolson Island, 43.39% at Red Skin Island, 41.65% at North Bay, 36.54 % at Chidiyatapu	*Acropora spp., Echinopora lamellosa, Porites solida, Diplostrea heliopora, Sinularis sp*	Krishnan et al., 2011
April, 2010	57%-90% corl bleaching at the Aves Island, Karlo Island, Sound Island, Rail Island and Interview Island	*Porites* sp, *Goniastrea* sp., *Acropora* sp, *Pocillopra* sp, *Fungia* sp., *Pachyseris* sp, *Pectinia* sp.	Sadhukhan and Raghunathan, 2012
April-July 2010	Overall 80±10.5%; Max 90.43% at Mayabunder and Minimum 69.83% at Rani Jhansi Marine National Park	Not mentioned	Mondal et al., 2014
April-May 2005	25% coral bleaching	Mortality of *Acropora* (43%), *Montipora* (22%) and *Porites* (14%)	Mohanty et al., 2013,

			Krishnan et al. (2011)
			Sarkar and Ghosh (2013)
April 2002	20-40%	Not mentioned	Krishnan et al. 2011
March 1998	up to 80% coral mortality	Not mentioned	Wafar, 1999, Ravindran et al. (1999)

204 Experienced degree heating weeks and reported bleaching percentage from four major reefs

Indian coral reefs were exposed to different degree heating weeks (DHWs; $°$C-weeks)- an indices for

assessing thermal stress based on satellite- derived SST data, and their bleaching percentage differ among

sites for all three major bleaching events. For 1998, 90% of reefs experienced bleaching in the Gulf of

Mannar, with 23% mortality to DHWs 6.7. In Andaman, 80% of reefs experienced bleaching to DHWs

4.9. In Lakshadweep, 82% of reefs experienced bleaching with 26% mortality to DHWs 6.17, while in the

Gulf of Kachchh, 11 to 30% of reefs experienced bleaching to DHWs 2.16. For 2010, In Andaman and

Nicobar, 36.5% to 77% of reefs experienced bleaching to DHWs 11.7. Whereas, 10% to 72% of reefs

experienced bleaching, and 24% mortality to DHWs 5.6 in the Gulf of Mannar. In the Gulf of Kachchh,

60-70% of reefs experienced bleaching to 11 DHWs. In 2016, In Andaman & Nicobar reef experienced

DHWs 7.21 to 9.5, but this site did not report the bleaching. In the Gulf of Mannar, 23% to 60% of reefs

experienced bleaching to DHWs 7.6 (41 days), while in the Gulf of Kachchh, 3.9% of reefs experienced

bleaching to DHWs 5.57. Above mentioned information is graphed in Fig 4, and the details of thermal

stress events on Indian reefs are given in the Table 2 and 3. Daily trends of mean SST and SST anomaly

for the period 2010 to 2021 in major Indian coral region (Data source: NOAA CRW v 3.1) were represented

in the Supplementary Fig S1-S5.

Fig. 4. Degree heating weeks and recorded bleaching percentage to three major bleaching events from

major Indian coral reefs

Table 2: Warmest month's sea surface temperature (SST), warmest month, SST anomaly, duration of

thermal stress and degree heating weeks for mass coral bleaching years in the Indian coral reef regions

based on Arora et al. 2019

Year	Coral bleaching indices	Andaman	Nicobar	Lakshadweep	Gulf of Mannar	Gulf of Kachchh
1998	Warmest month SST (°C)	30.66	30.69	30.71	30.84	29.58
	Warmest month SST anomaly (°C)	0.76	0.87	0.81	0.61	0.23
	Duration of thermal stress (days)	61	67	68	67	37
	Total no. of continuous days (%)	39.87	43.79	44.44	43.79	24.18
	Degree heating week (°C)	4.91	6.43	6.17	6.77	2.16
2010	Warmest month SST (°C)	31.43	31.26	30.77	30.87	30.44
	Warmest month SST anomaly (°C)	1.53	1.44	0.87	0.64	1.09
	Duration of thermal stress (days)	87	85	73	48	91
	Total no. of continuous days (%)	56.86	55.55	47.71	31.37	59.48
	Degree heating week (°C)	11.72	12.24	6.8	5.06	10.98
2016	Warmest month SST (°C)	31.03	31.2	30.86	31.21	29.91
t	Warmest month SST anomaly (°C)	1.13	1.38	0.96	0.98	0.56
	Duration of thermal stress (days)	53	67	91	63	70
	Total no. of continuous days (%)	34.42	43.51	59.09	40.91	45.45

Degree heating week (°C)	7.21	9.51	8.71	7.69	5.57

Table 3: Warmest months, thermal thresholds, and bleaching thresholds of Indian coral reefs regions by

using NOAA OISST data from 1982 to 2018 based on Arora et al. 2019; and De et al. 2021

Coral reef regions	Warmest month	Warmest quarter	Thermal threshold (°C)	Bleaching threshold (°C)
Gulf of Kachchh	June	March-July	29.35 ± 0.45	29.85
Malvan	May	March-May	29.39° C + 0.49	29.89
Lakshadweep	May	March-May	29.90 ± 0.49	30.4
Gulf of Mannar	April	March-May	30.23 ± 0.39	30.73
Andaman	May	April-June	29.90 ± 0.58	30.4
Nicobar	May	March-May	29.83 ± 0.63	30.33

Species susceptibility responses - winners and losers

To rank the species' susceptibility to bleaching events, we have selected two major bleaching events (2010 and 2016) and one short-term bleaching event from three major reef regions (Gulf of Mannar, Gulf of Kachchh, and Andaman reef). These reefs had species response details to 2010 and 2016 major bleaching episodes and one short-term bleaching episode, as the 1998 bleaching event does not have species response details. Four coral genera in the Gulf of Mannar, *Acropora, Pocillopora, Montipora,* and *Porites* were bleached to short-term bleaching events (lowest DHWs - 2). In addition, *Echinopora,* and *Favia* were bleached in 2010 (DHWs, 5.6). In 2016, When DHWs increased to 7.69, susceptibility differences diminished among species. In the Gulf of Kachchh, *Porites,* and *Favia* were bleached to short-term 2010 bleaching. In addition, *Goniopora, Goniastraea, Leptastrea,* and *Gascinaraea* were bleached in 2016 (DHWs- 5.57, WM 29.9). The disparity disappeared among coral genera to 2010 long-term bleaching episode (DHWs-10.98, WM-30.44). In Andaman reefs, *Acropora, Montipora,* and *Porites* were bleached to all three bleaching events. *Fungia* was bleached during 2016 in addition to short-term events. Again, the disparity in susceptibility disappeared when DHWs extended to above ten weeks (DHWs 11-12). Bleached

244　coral genera details against DHWs from different reef site details are given in Fig. 5.

Fig. 5. Bleaching response observed in coral genera to one short-term and two long-term (Major) bleaching

events from left to right 1) Gulf of Mannar (GM), 2) Gulf of Kachchh (GoK), and 3) Andaman and Nicobar

Islands (A&N).

Contrasting conclusions on species susceptibility to the 2016 long-term bleaching event in the Gulf of

Mannar and Palk Bay reef

In the Gulf of Mannar, three studies documented the impact and species susceptibility pattern during the

2016 (DHW= 7.6, highest temperature 31.2°C) long-term bleaching event. All these three surveys are

independent one-time surveys. These three studies created excellent data, but their conclusions differ on

the bleaching percentage and species susceptibility ranking. Krishnan et al. (2018) found that 46% of reefs

experiencing bleaching and 68% of bleached corals were massive forms from the Tuticorin group of

islands. They found that 22 out of the 26 massive forms were bleached, but the *Acropora* corymbose

(ACC), digitate (ACD), and encrusting coral (CE) forms had not bleached in any of their study sites. Based

on the findings, the study concluded that branching corals might have adapted to bleaching as they have

experienced repeated bleaching events in the Gulf of Mannar.

In contrast, Edward et al. (2018) found the opposite pattern. They reported that 23.92% of reefs experienced

a 2016 bleaching episode (21% of reefs in the Tuticorin reef). They found higher susceptibility and

mortality in branching corals than in the massive forms (mortality and bleaching percentage not provided)

(Edward et al., 2018). This study concluded that fast-growing branching corals (fast-growing coral forms,

Acropora, *Montipora*, and *Pocillopora* are more susceptible to bleaching than the boulders (*Porites, Favia,*

and *Favites*). Another study reported more susceptibility in massive forms than in branching forms (Ranith

and Kripa, 2019). In the case of Palk Bay, two studies contrast one another. Krishnan et al. (2018) reported

that 71% of reefs experienced bleaching, and the highest susceptibility was in *Porites*, *Favia, Leptastrea,*

Montastrea, Oulophyllia, and *Platygyra*, wherein the highest vulnerability was observed in *Acropora* than

the massive life forms by different study (Thinesh et al., 2019b).

Fig. 6. Differential coral bleaching patterns observed from Indian coral reefs. A) Partially bleached *Porites* near to healthy *Favites b)* Bleached *Acropora*, *Favites*, partially bleached *platygyra* and bleached dead *Porites* partially covered by algae c) Bleached dead *Acropora* and bleached *Fungia*, *Goniastrea* and *Porites* d) Bleached dead Acropora along with bleached other coral genera.

Discussion

Global data shows that the degree of bleaching differs at different geographical locations. Similarly, based on the literature survey, we found variation in bleaching percentage and species response among Indian reefs. The observed bleaching intensity variation between study locations is likely due to the coral composition type, type of symbionts, depth, time of exposure, and various other factors (Hoogenboom et al., 2017). The Gulf of Kachchh is least susceptible to major bleaching events among all major reefs. For instance, in 2016, DHWs in the Gulf of Kachchh reached 5.57 weeks; only 3% of reefs experienced bleaching, while in the case of the Gulf of Mannar during 2010 (DHWs- 5.6, 2010), it was 10 to 70%, and in Andaman, it was ~ 80 (DHWs-4.91, 1998). This differential susceptibility between locations is likely due to differences in the taxonomic composition. (Marshall and Baird, 2000). The highest proportion of massive forms and the absence of sensitive coral genera (*Acropora, Pocillopora*) in the Gulf of Kachchh likely favors the better resilient to bleaching as massive forms are stress-tolerant (Darling et al., 2012, Yohesh et al, 2014). The physiological mechanism that leads to higher resistance in massive corals is unknown to Indian reefs. However, attempted studies in other locations found the highest metabolic rate (Gates and Edmunds, 1999), tissue thickness (Loya et al., 2001), mass transfer rates (Nakamura and Van Woesik, 2001), light absorbing capacities (Fabricius, 2006), heterotrophy (Grottoli et al., 2006), and symbiont association (Baker et al., 2004) favors a role in resistance.

Most bleaching studies have found that branching coral genera such as *Acropora* and *Pocillopora* are more susceptible to bleaching than slow-growing massive corals (Baker et al., 2004; Loya et al., 2001; Marshall and Baird, 2000; McClanahan et al., 2004), except few studies reported contrasting patterns (Guest et al.,

2012; Krishnan et al., 2018). This observation has led to the widespread classification of fast-growing

branching species as "losers" in the face of global warming (Loya et al., 2001), at least in the short term

(Van Woesik et al., 2011). Likewise, our data shows the highest susceptibility in *Acropora* at every location

except in the Gulf of Kachchh, where the *Acropora* population diminished for unknown reasons. In contrast,

Porite's susceptibility to bleaching is also invariably recorded in short and long-term events across the site.

We infer that *Porites* and massive colony types exhibit earlier partial bleaching (pale coloration). However,

they remain in the bleaching state for the entire bleaching event and recover, unlike their sensitive

counterparts who underwent mortality. Since existing data undifferentiated partial/complete bleaching, the

Porites fall under susceptible genera. This pattern suggests that if the survey happens in the first week, one

could expect partial bleaching in the *Porites* colonies even before the susceptible branching forms start to

pale. In the Gulf of Mannar, *Montipora,* and *Pocillopora* are the other three genera equally susceptible to

short-term events. Other coral genera that have only bleached during long-term bleaching events are given

in the graph. Similar to bleaching susceptibility, coral taxa differ in their recovery pattern (Baird &

Marshall, 2002). A study from the Great Barrier Reef (GBR) recorded the highest recovery in massive

corals than the branching corals. For instance, a tagged colony study found that 88% of *Acropora*

hyacinthus, and 32% of *A. millepora* colonies died compared to 13% of *Platygyra daedalea* and 0% of

Porites lobata (Baird & Marshall, 2002). Although few studies looked at the recovery pattern in coral

genera immediately following bleaching, the study found the highest recovery in the massive type than the

branching corals (Manikandan et al., 2016b; Thinesh et al., 2019b). In support of that, a large-scale survey

in the Gulf of Mannar after the mass bleaching event documented an increased massive coral cover in the

coral community composition as branching forms were presumed to have died as a result of bleaching. (Raj

et al., 2021). In India, studies are beginning to understand factors leading to the highest resistance among

specific coral taxa and growth forms (Thinesh et al., 2019). We assume this susceptibility pattern in coral

genera may be associated with differences in the type of symbiont hosted (Berkelmans and van Oppen,

2006), pre-bleaching history, metabolic rates (Gates and Edmunds, 1999), tissue thickness (Loya et al.,

2001), and heterotrophic feeding capacity (Grottoli et al., 2006).

We found a susceptibility pattern among coral genera to short-term bleaching events. However, the

disparity diminished to long-term bleaching events similar to the study, which reported no difference in

species susceptibility to prolonged bleaching events in the GBR (Hughes et al., 2018). In addition, the

scientific community hypothesized that recurrent bleaching episodes favor coral to adapt or acclimatize to

future bleaching episodes. Studies have indicated that previous recurrent bleaching exposure protects

corals from subsequent episodes (Carilli et al., 2012; Guest et al., 2016; Palumbi et al., 2014; Trapon et al.,

2011). In contrast, one large-scale study from the GBR found no evidence of acclimation or adaptation in

the corals exposed to earlier bleaching events (Hughes et al., 2018). Our analysis of available literature

shows no evidence of protective effects from previous bleaching history in Indian coral reefs. For instance,

the Gulf of Mannar reefs experienced eight bleaching episodes, including three long-term and five short-

term. However, 23 to 60% of reefs experienced bleaching in 2016 (DHWs-7.69), and the same coral genera

show the highest susceptibility (Raj et al.2021).

Our analysis of species susceptibility reports for 2016 long-term bleaching events from the Gulf of Mannar

and Palk Bay revealed contrasting conclusions on species susceptibility ranking. Then, we examined why

the study found a contrasting pattern of results even though they surveyed the exact locations. This pattern

may be attributed to the colonies they encountered being different from one another, which might have

experienced different thermal histories. However, we assume that the survey timing is most likely to

influence the pattern difference between the studies similar to the bleaching surveys undertaken during the

1997/1998 El Nino (surveyed 10, 14, 20, 28, and 40 weeks after bleaching began; Marshall and Baird,

2000) as well as the 2015/2016 El Nino from Kiritimati Island (Claar and Baum, 2019). In GBR Marine

Park, a survey during the 6th week found the highest susceptibility to the *Acropora hyacinthus* population,

while the massive reef experienced mild bleaching. More *Acropora* died between six and ten weeks, but

others recovered completely while more massive corals bleached. When temperatures returned to normal

during the 14th week, *Acropora* either died or attained complete recovery, yet, many massive colonies

remained pale and started recovery. If we rely on a one-time survey without baseline data, one might have

erroneously concluded species susceptibility, percentage of bleaching, and mortality in a scenario like this.

By surveying bleaching at multiple time points during this event, the authors accurately captured the

dynamics of bleaching and mortality and provided a foundation for future work.

Similarly, on Kavaratti Island, during the 2015 and 2016 bleaching events, the 11th-week survey found

many tolerant corals mostly or entirely bleached, while thermally sensitive corals appeared fully pigmented.

However, when they returned after ten months, the temperature became normal. They found a reverse

pattern: susceptible corals died while resistant corals recovered in many locations. These observed

bleaching patterns in two studies from sample locations are likely due to each study's survey time

difference. This contrasting conclusion underscores the importance of repetitive surveys with baseline data

before taking up conservation management practices. Based on available information from the Gulf of

Mannar reef, one can expect the following diagrammatic bleaching scenario in the field.

Fig. 7. This diagram represents how a one-time survey during different DHWs could influence our result

in determining corals' susceptibility to long-term bleaching events. Probable scenarios 1) If the survey

happened during DHWs 1-4, the study concludes that branching corals are more tolerant than massive

corals as massive corals start to pale (partial bleaching to 10-20%) while branching corals show no

bleaching/partially bleached. 2) In DHWs at peak, there is no difference in species susceptibility but most

likely missed dead branching corals 3) DHWs 6-11 conclude the highest bleaching in massive corals as

massive corals are being recovered. In contrast, bleached branching corals died, which divers failed to

notice/record. Picture: a-b: Partially bleached Massive corals vs. unbleached branching corals; c-d:

Bleached corals were irrespective of coral genera; e-f: Bleached dead *Acropora* covered with secondary

algae.

Conclusion

The frequency and severity of coral-bleaching episodes are increasing due to ongoing climate change,

which alters species composition and reef function that determines future coral reefs. Hence, information

on bleaching episodes, intensity, and species susceptibility is essential from broad geological location to

frame conservation strategy. 23 years of concerted bleaching survey data from Indian coral reefs, i)

provides insights on DHWs and their influence in bleaching intensity and species susceptibility response

for major Indian reefs; ii) informed the highest susceptibility in branching corals and recovery in massive

corals across locations; iv) highlighted the contrasting conclusion to long-term bleaching events and

underscored the importance of repeated surveys during long-term bleaching events before concluding

species susceptibility. This meta-analysis also underlines the limitations that hinder acquiring the most

from our extensive analysis: i) one-time surveys with no information about DHWs, ii) no symbiont diversity

data, ii) partial/incomplete bleaching information, iii) the presence and absence of data on sensitive coral

genera rather than a proportion iv) lack of follow up survey to document mortality and bleaching recovery,

v) contrasting in conclusions on species susceptibility to long-term bleaching events as a result of one-time

survey. Despite these limitations, present study helps to improve the knowledge gap on bleaching episodes

and reveal the coral community responses to short-term and long-term bleaching events. This analysis also

identifies that coral bleaching studies in Indian reefs are mostly fragmented individual efforts, and based

on a one-time survey. We recommend inter-institutional collaborations, adhering to a uniform survey

procedure and including all details in the future survey to obtain the most accurate field data on bleaching episodes using a broad, data calibration for accuracy and quality control, and a transdisciplinary approach that will ultimately serve the common goal to improve coral health and conservation. This accurate data will not only allow us to compare with future events from Indian coral reefs and reefs worldwide, this will enable us to select potential species for restoration, which will ultimately determine restoration success and coral reef conservation.

Acknowledgment

TT thanks the Current CSIR Pool Scientist Scheme, India. Rufford & Idea Wild small grants for support. KD is thankful to CSIR-NIO for facilities and support. Authors are thankful to Prof. Kartik Shanker, and Dr. Titus Immanuel, Centre for Ecological sciences, Indian Institute of Science for his valuable comments to improve this manuscript.

Author contributions

Thinesh Thangadurai: Conceptualization, Methodology, Writing - original draft, Writing – review & editing. **Kalyan De**: Methodology, Data curation, Writing - original draft, Writing – review & editing. **Anthony Bellantuono**: Writing – review & editing. **Sobana**, **Riana Peter, Sivagurunathan:** Data curation. **Polpass Arul Jose, Joseph Selvin**: review & editing.

References

Adhavan, D., Kamboj, R.D., Marimuthu, N., Bhalodi, M.M., 2014. Seasonal variation and climate change influence coral bleaching in Pirotan Island, Gulf of Kachchh Marine National Park, Gujarat. Curr. Sci. 107, 1780–1781.

Albright, R., Langdon, C., 2011. Ocean acidification impacts multiple early life history processes of the Caribbean coral Porites astreoides. Glob. Chang. Biol. 17, 2478–2487. https://doi.org/10.1111/j.1365-2486.2011.02404.x

Albright, R., Mason, B., 2013. Projected Near-Future Levels of Temperature and pCO2 Reduce Coral
    Fertilization Success. PLoS One 8. https://doi.org/10.1371/journal.pone.0056468

Anthony, S.L., Page, C.A., Bourne, D.G., Willis, B.L., 2008. Newly characterized distinct phases of the
coral disease 'atramentous necrosis ' on the Great Barrier Reef Diseases of Aquatic Organisms, 81 (3).
pp. 255-259. https://doi.org/10.3354/dao01962

Arora, M., De, K., Ray Chaudhury, N., Nanajkar, M., Chauhan, Prakash Pateriya, B., 2021a. Climate
    change and recurrent coral bleaching over Gulf of Kachchh and Malvan Region, India., in: Climate
    Issues in Asia and Africa - Examining Climate, Its Flux, the Consequences, and Society's
    Responses. IntechOpen. https://doi.org/10.5772/intechopen.96806

Arora, M., Gujrati, A., Chaudhury, N.R., Chauhan, P., Patel, R.C., 2021b. Assessment of coral reef
    thermal stress over India based on remotely sensed sea surface temperature. Geocarto Int. 36, 740–
    757. https://doi.org/10.1080/10106049.2019.1624983

Arora, M., Ray Chaudhury, N., Gujrati, A., Patel, R.C., 2019. Bleaching stress on Indian coral reef
    regions during mass coral bleaching years using NOAA OISST data. Curr. Sci. 117, 242–250.

Arthur, R., 2000. Coral bleaching and mortality in three Indian reef regions during an El Niño southern
    oscillation event. Curr. Sci. 79, 1723–1729.

Baird, A.H., Marshall, P.A., 2002. Mortality Growth and Reproduction Corals GBR. Mar. Ecol. Prog.
    Ser. 237, 133–141.

Baker, A.C., Starger, C.J., McClanahan, T.R., Glynn, P.W., 2004. Corals' adaptive response to climate
    change: Shifting to new algal symbionts may safeguard devastated reefs from extinction. Nature
    430, 741. https://doi.org/10.1038/430741a

Bellwood, D.R., Streit, R.P., Brandl, S.J., Tebbett, S.B., 2019. The meaning of the term 'function' in
    ecology: A coral reef perspective. Funct. Ecol. https://doi.org/10.1111/1365-2435.13265

Berkelmans, R., van Oppen, M.J.., 2006. The role of zooxanthellae in the thermal tolerance of corals: a
    "nugget of hope" for coral reefs in an era of climate change. Proc. R. Soc. B Biol. Sci. 273, 2305–
    2312. https://doi.org/10.1098/rspb.2006.3567

Boopathi, M., Kumar, P.D., Edward, J.K.P., 2021. The impacts of climate change and post bleaching
    effects on the octocoral communities of Gulf of Mannar, India. J. Earth Syst. Sci. 130, 1–11.

https://doi.org/10.1007/S12040-021-01701-3

Brainard, R.E., Oliver, T., McPhaden, M.J., Cohen, A., Venegas, R., Heenan, A., Vargas-Angel, B.,
Rotjan, R., Mangubhai, S., Flint, E., Hunter, S.A., 2018. 5. Ecological impacts of the 2015/16 El
Niño in the central equatorial Pacific. Bull. Am. Meteorol. Soc. 99, S21–S26.
https://doi.org/10.1175/BAMS-D-17-0128.1

Brown, B. Coral bleaching: causes and consequences. *Coral Reefs* **16,** S129–S138 (1997).

Bruno, J.F., Selig, E.R., Casey, K.S., Page, C.A., Willis, B.L., Harvell, C.D., Melendy, A.M., 2007.
Thermal Stress and Coral Cover as Drivers of Coral Disease Outbreaks *PLoS Biology (2007) 5(6)*
1220-1227. https://doi.org/10.1371/journal.pbio.0050124

Camp, E.F., Schoepf, V., Mumby, P.J., Hardtke, L.A., Rodolfo-Metalpa, R., Smith, D.J., Suggett, D.J.,
2018. The future of coral reefs subject to rapid climate change: Lessons from natural extreme
environments. Front. Mar. Sci. 5, 1–21. https://doi.org/10.3389/fmars.2018.00004

Carilli, J., Donner, S.D., Hartmann, A.C., 2012. Historical temperature variability affects coral response
to heat stress. PLoS One 7, 1–9. https://doi.org/10.1371/journal.pone.0034418

Cai, W., Borlace, S., Lengaigne, M., Van Rensch, P., Collins, M., Vecchi, G., Timmermann, A., Santoso,
A., Mcphaden, M.J., Wu, L., England, M.H., Wang, G., Guilyardi, E., Jin, F.F., 2014. Increasing
frequency of extreme El Niño events due to greenhouse warming. Nat. Clim. Chang. 4, 111–116.
https://doi.org/10.1038/nclimate2100

Chandra Mohanty, P., 2017. Coral bleaching along andaman coast Due to thermal Stress during summer
months of 2016: a geospatial Assessment. Am. J. Environ. Prot. 6, 1.
https://doi.org/10.11648/j.ajep.20170601.11

Claar, D.C., Baum, J.K., 2019. Timing matters: survey timing during extended heat stress can influence
perceptions of coral susceptibility to bleaching. Coral Reefs 38, 559–565.
https://doi.org/10.1007/s00338-018-01756-7

Cornwell, B., Armstrong, K., Walker, N., Lippert, M., Nestor, V., Golbuu, Y., Palumbi, S.R., 2021.
Widespread variation in heat tolerance and symbiont load are associated with growth tradeoffs in the
coral acropora hyacinthus in palau. Elife 10, 1–3. https://doi.org/10.7554/eLife.64790

Darling, E.S., Alvarez-Filip, L., Oliver, T.A., Mcclanahan, T.R., Côté, I.M., 2012. Evaluating life-history

strategies of reef corals from species traits. Ecol. Lett. 15, 1378–1386.

https://doi.org/10.1111/j.1461-0248.2012.01861.x

Davy S. K., Allemand D., Weis V. M. 2012. Cell biology of cnidarian-dinoflagellate symbiosis.

Microbiol. Mol. Biol. Rev. 76, 229–261.

De, K., Sautya, S., Mote, S., Tsering, L., Patil, V., Nagesh, R., Ingole, B., 2015. Is climate change

triggering coral bleaching in tropical reef? Curr. Sci. 109, 1379–1880.

De, K., Venkataraman, K., Ingole, B., 2017. Current status and scope of coral reef research in India: a

bio-ecological perspective. Indian J. Geo. Mar. Sci. 46, 647–662.

De, K., Venkataraman, K., Ingole, B., 2020. The hard corals (Scleractinia) of India: a revised checklist.

Indian J. Geo. Mar. Sci. 49, 1651–1660.

De, K., Nanajkar, M., Arora, M., Nithyanandan, M., Mote, S., Ingole, B., 2022. Application of remotely

sensed sea surface temperature for assessment of recurrent coral bleaching (2014–2019) impact on a

marginal coral ecosystem. Geocarto Int. 37 (15), 4483-4508.

https://doi.org/10.1080/10106049.2021.1886345

De, K., Nanajkar, M., Mote, S., Ingole, B. (2023) Reef on the edge: Resilience failure of marginal coral

habitat by combined impact of recurrent coral bleaching, coral disease, and local stressors. Environ.

Sci. Pollut. Res. 30, 7288–7302. https://doi.org/10.1007/s11356-022-22651-3

Dixon, A.M., Forster, P.M., Heron, S.F., Stoner, A.M.K., Beger, M., 2022. Future loss of local-scale

thermal refugia in coral reef ecosystems. PLOS Clim. 1, e0000004.

https://doi.org/10.1371/journal.pclm.0000004

Edmunds, P.J., 1994. Evidence that reef-wide patterns of coral bleaching may be the result of the

distribution of bleaching-susceptible clones, Marine Biology. Marine Biology.

Edward, J.K.P., Mathews, G., Diraviya Raj, K., Laju, R.L., Selva Bharath, M., Arasamuthu, A., Dinesh

Kumar, P., Bilgi, D.S., Malleshappa, H., 2018. Coral mortality in the Gulf of Mannar, southeastern

India, due to bleaching caused by elevated sea temperature in 2016. Curr. Sci. 114, 1967–1972.

https://doi.org/10.18520/cs/v114/i09/1967-1972

Edward, J.K.P., Mathews, G., Raj, K.D., Thinesh, T., Patterson, J., Tamelander, J., Wilhelmsson, D.,

2012. Coral reefs of Gulf of Mannar, India-signs of resilience. Proc. 12th Int. Coral Reef Symp.

Cairns, Aust. 9-13 July 2012 9–13.

Fabricius, K.E., 2006. Effects of irradiance, flow, and colony pigmentation on the temperature
495   microenvironment around corals: Implications for coral bleaching? Limnol. Oceanogr. 51, 30–37.
496   https://doi.org/10.4319/lo.2006.51.1.0030

497   Fordyce, A.J., Ainsworth, T.D., Heron, S.F., Leggat, W., 2019. Marine heatwave hotspots in coral reef
498   environments: Physical drivers, ecophysiological outcomes and impact upon structural complexity.
499   Front. Mar. Sci. https://doi.org/10.3389/fmars.2019.00498

500   Gardner, T.A., Côté, I.M., Gill, J.A., Grant, A., Watkinson, A.R., 2003. Long-term region-wide declines
501   in Caribbean corals. Science, 301, 958–960. https://doi.org/10.1126/science.1086050

502   Gates, R.D., Edmunds, P., 1999. The physiological mechanisms of acclimatization in tropical reef corals.
503   Am. Zool. 43, 30–43. https://doi.org/10.1093/icb/39.1.30

504   Glynn, P.W., Maté, J.L., Baker, A.C., Calderón, M.O., 2001. Coral bleaching and mortality in Panama
505   and Ecuador during the 1997-1998 El Niño-Southern Oscillation event: Spatial/temporal patterns
506   and comparisons with the 1982-1983 event. Bull. Mar. Sci. 69, 79–109.

507   González-Pech, R.A., Stephens, T.G., Chen, Y., Mohamed, A.R., Cheng, Y., Shah, S., Dougan, K.E.,
508   Fortuin, M.D.A., Lagorce, R., Burt, D.W., Bhattacharya, D., Ragan, M.A., Chan, C.X., 2021.
509   Comparison of 15 dinoflagellate genomes reveals extensive sequence and structural divergence in
510   family Symbiodiniaceae and genus Symbiodinium. BMC Biol. 19. https://doi.org/10.1186/s12915-
511   021-00994-6

512   Grottoli, A.G., Rodrigues, L.J., Palardy, J.E., 2006. Heterotrophic plasticity and resilience in bleached
513   corals. Nature 440, 1186–1189. https://doi.org/10.1038/nature04565

514   Grottoli, A.G., Warner, M.E., Levas, S.J., Aschaffenburg, M.D., Schoepf, V., Mcginley, M., Baumann, J.,
515   Matsui, Y., 2014. The cumulative impact of annual coral bleaching can turn some coral species
516   winners into losers. Glob. Chang. Biol. 20, 3823–3833. https://doi.org/10.1111/gcb.12658

517   Guest, J.R., Baird, A.H., Maynard, J.A., Muttaqin, E., Edwards, A.J., Campbell, S.J., Yewdall, K.,
518   Affendi, Y.A., Chou, L.M., 2012. Contrasting patterns of coral bleaching susceptibility in 2010
519   suggest an adaptive response to thermal stress. PLoS One 7, 1–8.
520   https://doi.org/10.1371/journal.pone.0033353

Guest, J.R., Low, J., Tun, K., Wilson, B., Ng, C., Raingeard, D., Ulstrup, K.E., Tanzil, J.T.I., Todd, P.A.,

Toh, T.C., McDougald, D., Chou, L.M., Steinberg, P.D., 2016. Coral community response to

bleaching on a highly disturbed reef. Sci. Rep. 6, 1–10. https://doi.org/10.1038/srep20717

Hoogenboom, M.O., Frank, G.E., Chase, T.J., Jurriaans, S., Álvarez-Noriega, M., Peterson, K., Critchell,

K., Berry, K.L.E., Nicolet, K.J., Ramsby, B., Paley, A.S., 2017. Environmental drivers of variation

in bleaching severity of Acropora species during an extreme thermal anomaly. Front. Mar. Sci. 4, 1–

16. https://doi.org/10.3389/fmars.2017.00376

Hughes, T.P., Baird, A.H., Bellwood, D.R., Card, M., Connolly, S.R., Folke, C., Grosberg, R., Jackson,

J.B.C., Kleypas, J., Lough, J.M., Marshall, P., Nystro, M., 2003. Climate Change , Human Impacts ,

and the Resilience of Coral Reefs. Science. 2003 Aug 15;301(5635):929-33. doi:

10.1126/science.1085046. PMID: 12920289.

Hughes, T.P., Barnes, M.L., Bellwood, D.R., Cinner, J.E., Cumming, G.S., Jackson, J.B.C., Kleypas, J.,

Van De Leemput, I.A., Lough, J.M., Morrison, T.H., Palumbi, S.R., Van Nes, E.H., Scheffer, M.,

2017a. Coral reefs in the Anthropocene. Nature 546, 82–90. https://doi.org/10.1038/nature22901

Hughes, T.P., Kerry, J.T., Álvarez-Noriega, M., Álvarez-Romero, J.G., Anderson, K.D., Baird, A.H.,

Babcock, R.C., Beger, M., Bellwood, D.R., Berkelmans, R., Bridge, T.C., Butler, I.R., Byrne, M.,

Cantin, N.E., Comeau, S., Connolly, S.R., Cumming, G.S., Dalton, S.J., Diaz-Pulido, G., Eakin,

C.M., Figueira, W.F., Gilmour, J.P., Harrison, H.B., Heron, S.F., Hoey, A.S., Hobbs, J.P.A.,

Hoogenboom, M.O., Kennedy, E. V., Kuo, C.Y., Lough, J.M., Lowe, R.J., Liu, G., McCulloch,

M.T., Malcolm, H.A., McWilliam, M.J., Pandolfi, J.M., Pears, R.J., Pratchett, M.S., Schoepf, V.,

Simpson, T., Skirving, W.J., Sommer, B., Torda, G., Wachenfeld, D.R., Willis, B.L., Wilson, S.K.,

2017b. Global warming and recurrent mass bleaching of corals. Nature 543, 373–377.

https://doi.org/10.1038/nature21707

Hughes, T.P., Kerry, J.T., Baird, A.H., Connolly, S.R., Dietzel, A., Eakin, C.M., Heron, S.F., Hoey, A.S.,

Hoogenboom, M.O., Liu, G., McWilliam, M.J., Pears, R.J., Pratchett, M.S., Skirving, W.J., Stella,

J.S., Torda, G., 2018. Global warming transforms coral reef assemblages. Nature 556, 492–496.

https://doi.org/10.1038/s41586-018-0041-2

Hussain, A., Ingole, B., 2020. Massive coral bleaching in the patchy reef of Grande Island, along the

eastern Arabian Sea during the 2015/16 global bleaching event. Reg. Stud. Mar. Sci. 39, 101410.

https://doi.org/10.1016/j.rsma.2020.101410

Kenyon, J.C., Brainard, R.E., Hoeke, R.K., Parrish, F.A., Wilkinson, C.B., 2006. Towed-diver surveys, a
method for mesoscale spatial assessment of Benthic Reef habitat: A case study at Midway Atoll in
the Hawaiian Archipelago. Coast. Manag. 34, 339–349.
https://doi.org/10.1080/08920750600686711

Idrees Babu, K.K., Suresh Kumar, S., 2016. Status and changing trends of coral reefs in Lakshadweep
archipelago after 1998 mass bleaching event - long term monitoring survey. Int. J. Appl. pure Sci.
Agric. 163–175.

Jeevamani, J.J.J., Kamalakannan, B., Nagendran, N.A., Chandrasekaran, S., 2013. Climate change
induced coral bleaching and algal phase shift in reefs of the gulf of Mannar, India, in:
Environmental Science and Engineering (Subseries: Environmental Science). Springer Berlin
Heidelberg, pp. 87–94. https://doi.org/10.1007/978-3-642-36143-2_6

Joshi, D., Munjpara, S., Banerji, U., Parasharya, D., 2014. Coral bleaching observations in the Gulf of
Kachchh, India – a climate induced stress on the Scleractinians. J. Aquat. Biol. Fish. 2, 106–113.

Krishnan, P., Dam Roy, S., George, G., Srivastava, R.C., Anand, A., Murugesan, S., Kaliyamoorthy, M.,
Vikas, N., Soundararajan, R., 2011. Elevated sea surface temperatureduring May 2010 induces
massbleaching of corals in the Andaman. Current Science, 100 (1). pp. 111-117.

Krishnan, P., Purvaja, R., Sreeraj, C.R., Raghuraman, R., Robin, R.S., Abhilash, K.R., Mahendra, R.S.,
Anand, A., Gopi, M., Mohanty, P.C., Venkataraman, K., Ramesh, R., 2018. Differential bleaching
patterns in corals of Palk Bay and the Gulf of Mannar. Curr. Sci. 114, 679–685.
https://doi.org/10.18520/cs/v114/i03/679-685

Kumaraguru, A. K., Jayakumar, K., C.M.R., 2003. Coral bleaching 2002 in the Palk Bay, southeast coast
of India. Curr. Sci. 85 (12): 1787–1793.

LaJeunesse, T.C., Parkinson, J.E., Gabrielson, P.W., Jeong, H.J., Reimer, J.D., Voolstra, C.R., Santos,
S.R., 2018. Systematic Revision of Symbiodiniaceae Highlights the Antiquity and Diversity of
Coral Endosymbionts. Curr. Biol. 28, 2570-2580.e6. https://doi.org/10.1016/j.cub.2018.07.008

Loya, Y., Sakai, K., Nakano, Y., Woesik, R. Van, 2001. Coral bleaching: the winners and the losers.
Ecol. Lett. 4, 122–131. https://doi.org/10.1046/j.1461-0248.2001.00203.x

Manikandan, B., Ravindran, J., Vidya, P.J., Mani Murali, R., 2016a. Bleaching and recovery patterns of

corals in Palk Bay, India: An indication of bleaching resilient reef. Reg. Stud. Mar. Sci. 8, 151–156.

https://doi.org/10.1016/j.rsma.2016.07.005

Manikandan, S., Ganesapandian, S., Singh, M., Anand, M., Kumaraguru, A.K., 2014. Impact of climate

change induced coral bleaching in the northern part of Gulf of Mannar and Palk bay, southeast coast

of India. J. Biol. Sci. 14, 276–283. https://doi.org/10.3923/jbs.2014.276.283

Marimuthu, N., Jerald Wilson, J., Vinithkumar, N. V., Kirubagaran, R., 2013. Coral reef recovery status

in south Andaman Islands after the bleaching event 2010. J. Ocean Univ. China 12, 91–96.

https://doi.org/10.1007/s11802-013-2014-2

Marshall, P.A., Baird, A.H., 2000. Bleaching of corals on the Great Barrier Reef: differential

susceptibilities among taxa. Coral Reefs 19, 155–163. https://doi.org/10.1007/s003380000086

McClanahan, T.R., Baird, A.H., Marshall, P.A., Toscano, M.A., 2004. Comparing bleaching and

mortality responses of hard corals between southern Kenya and the Great Barrier Reef, Australia.

Mar. Pollut. Bull. 48, 327–335. https://doi.org/10.1016/j.marpolbul.2003.08.024

Moberg, F., Folke, C., 1999. Ecological goods and services of coral reef ecosystems. Ecol. Econ. 29,

215–233. https://doi.org/10.1016/S0921-8009(99)00009-9

Mohanty, P.C., Mahendra, R.S., Bisoyi, H., Kumar Tummula, S., Grinson, G., Nayak, S., Kumar Sahu,

B., 2013. Assessment of the coral bleaching during 2005 to decipher the thermal stress in the coral

environs of the Andaman Islands using remote sensing. Eur. J. Remote Sens. 46, 417–430.

https://doi.org/10.5721/EuJRS20134624

Mohanty, P.C., Venkateshwaran, P. Mahendra, R.S., Shiva Kumar, H., Kumar, T.S., Vinithkumar, N. V.,

Kirubagaran, R., Ramesh, S., Ramesh, R., Sathianarayanan, D., Prakash, V.D., Ramadasss, G.A.,

Shenoi, S.S.C., 2017. Coral bleaching along Andaman coast due to thermal stress during summer

months of 2016: A geospatial assessment. Am. J. Environ. Prot. 6, 1.

https://doi.org/10.11648/j.ajep.20170601.11

Moher, D., Liberati, A., Tetzlaff, J., Altman, D.G., The PRISMA Group., 2009. Preferred reporting items

for systematic reviews and MetaAnalyses: the PRISMA state- ment. PLoS Med. 6 (7) e1000097

https://doi.org/10.1371/journal.pmed1000097

Mondal, T., Raghunathan, C., Venkataraman, K., 2014. Coral bleaching in Andaman Sea-an indicator for

climate change in Andaman and Nicobar Islands. Indian J. Geo-Marine Sci. 43, 1945–1948.

Mote, S., Gupta, V., De, K., Hussain, A., More, K., Nanajkar, M., & Ingole, B. (2021). Differential
Symbiodiniaceae association with coral and coral-eroding sponge in a bleaching Impacted marginal
coral reef environment. *Frontiers in Marine Science*, **8**, 442.

Nakamura, T., Van Woesik, R., 2001. Water-flow rates and passive diffusion partially explain differential
survival of corals during the 1998 bleaching event. Mar. Ecol. Prog. Ser. 212, 301–304.
https://doi.org/10.3354/meps212301

Palumbi, S.R., Barshis, D.J., Bay, R. a, 2014. To Future Climate Change Science.23;344(6186):895-8.
doi: 10.1126/science.1251336.

Pisapia, C., Burn, D., 2016. Changes in the population and community structure of corals during recent
disturbances) on Maldivian coral reefs. https://doi.org/10.1038/s41598-019-44809-9

Prada, F., Caroselli, E., Mengoli, S., Brizi, L., Fantazzini, P., Capaccioni, B., Pasquini, L., Fabricius,
K.E., Dubinsky, Z., Falini, G., Goffredo, S., 2017. Ocean warming and acidification synergistically
increase coral mortality. Sci. Rep. 7, 1–10. https://doi.org/10.1038/srep40842

Raj, K.D., Mathews, G., Bharath, M.S., Sawant, R.D., Bhave Vishal, Apte, D., Vasudevan, N., Edward,
J.K.P., 2018. Climate change-induced coral bleaching in Malvan Marine Sanctuary, Maharashtra,
India. Curr. Sci. 114, 384–387. https://doi.org/10.18520/cs/v114/i02/384-387\

Raj, K.D., Aeby, G.S., Mathews, G., Williams, G.J., Caldwell, J.M., Laju, R.L., Bharath, M.S., Kumar,
P.D., Arasamuthu, A., Asir, N.G.G., Wedding, L.M., Davies, A.J., Moritsch, M.M., Edward, J.K.P.,
2021. Coral reef resilience differs among islands within the Gulf of Mannar, southeast India,
following successive coral bleaching events. Coral Reefs 40, 1029–1044.
https://doi.org/10.1007/s00338-021-02102-0

Ramesh, C.H., Koushik, S., Shunmugaraj, T., Murthy, M.V.R., 2020. Exhibition of bleaching resistance
via adaptive bleaching pattern by coral reefs of the Gulf of Mannar during massive bleaching event
2019. J. Mar. Sci. 2, 1–5. https://doi.org/https://doi.org/10.30564/jms.v2i2.1666

Ranith, R., Kripa, V., 2019. Post-bleaching benthic status on reef sites of Mandapam group of islands and
Palk Bay, Indian Ocean following the 2016 bleaching events. Reg. Stud. Mar. Sci. 27, 100525.
https://doi.org/10.1016/j.rsma.2019.100525

Ravindran, J., Kannapiran, E., Manikandan, B., Mani Mural, R., Joseph, A., 2012. Bleaching and
secondary threats on the corals of Palk Bay: A survey and Proactive conservation needs. Indian J.
Mar. Sci. 41, 883–890.

Sadhukhan, K., Raghunathan, C., 2012. Study on coral bleaching (2010) in Middle Andaman, Andaman
and Nicobar Islands. Rec. Zool. Surv. India 112, 27–34.

Saroj, J., Gautam, R.K., Joshi, A., Tehseen, P., 2016. Review of Coral Reefs of India: Distribution,
Status, Research and Management. Int. J. Sci. Environ. Technol. 5, 3088–3098.

Sierra-Correa, P.C., Cantera Kintz, J.R., 2015. Ecosystem-based adaptation for improving coastal
planning for sea-level rise: A systematic review for mangrove coasts. Mar. Policy 51, 385–393.
https://doi.org/10.1016/j.marpol.2014.09.013

Stuart-Smith, R.D., Brown, C.J., Ceccarelli, D.M., Edgar, G.J., 2018. Ecosystem restructuring along the
Great Barrier Reef following mass coral bleaching. Nature 560, 92–96.
https://doi.org/10.1038/s41586-018-0359-9

Swain, D.L., Horton, D.E., Singh, D., Diffenbaugh, N.S., 2016. Trends in atmospheric patterns conducive
to seasonal precipitation and temperature extremes in California. Sci. Adv. 2.
https://doi.org/10.1126/sciadv.1501344

Thinesh, T., Meenatchi, R., Jose, P.A., Kiran, G.S., Selvin, J., 2019b. Differential bleaching and recovery
pattern of southeast Indian coral reef to 2016 global mass bleaching event: Occurrence of stress-
tolerant symbiont Durusdinium (Clade D) in corals of Palk Bay. Mar. Pollut. Bull. 145, 287–294.
https://doi.org/10.1016/j.marpolbul.2019.05.033

Trapon, M.L., Pratchett, M.S., Penin, L., 2011. Comparative Effects of Different Disturbances in Coral
Reef Habitats in Moorea, French Polynesia. J. Mar. Biol. 2011, 1–11.
https://doi.org/10.1155/2011/807625

Van Woesik, R., Sakai, K., Ganase, A., Loya, Y., 2011. Revisiting the winners and the losers a decade
after coral bleaching. Mar. Ecol. Prog. Ser. 434, 67–76. https://doi.org/10.3354/meps09203

Vinoth, R., Gopi, M., Kumar, T.T.A., Thangaradjou, T., Balasubramanian, T., 2012. Coral reef bleaching
at Agatti Island of Lakshadweep atolls, India. J. Ocean Univ. China 11, 105–110.
https://doi.org/10.1007/s11802-012-1822-0

Wafar, M.V.M., 1999. Status report India, in: Coral Reef Decradation in the Indian Ocean, Status Reports
and Project Presentation. CORDIO, Stockholm, Sweden, pp. 24–25.

Yadav, S., Alcoverro, T., Arthur, R., 2018. Coral reefs respond to repeated ENSO events with increasing
resistance but reduced recovery capacities in the Lakshadweep archipelago. Coral Reefs 37, 1245–
1257. https://doi.org/10.1007/s00338-018-1735-5

668