Cryo-EM structure of the bicarbonate receptor GPR30

Shota Kaneda[1], Airi Jo-Watanabe[2‡], Hiroaki Akasaka[1], Hidetaka S. Oshima[1], Takehiko

Yokomizo[2], Wataru Shihoya[1‡], Osamu Nureki[1‡].

Affiliations:

[1]Department of Biological Sciences, Graduate School of Science, The University of

Tokyo, Bunkyo, Tokyo 113-0033, Japan.

[2]Department of Biochemistry, Juntendo University Graduate School of Medicine, Tokyo

113-8421, Japan.

‡ To whom correspondence should be addressed. E-mail: awatanabe-tky@umin.ac.jp

(A.J.-W), wtrshh9@gmail.com (W.S.), and nureki@bs.s.u-tokyo.ac.jp (O.N.)

ORCID list
Airi Jo-Watanabe 0000-0002-5219-1553
Hiroaki Akasaka 0000-0003-2118-0912
Hidetaka S. Oshima 0009-0000-0341-707X
Takehiko Yokomizo 0000-0003-0899-9915
Wataru Shihoya 0000-0003-4813-5740
Osamu Nureki 0000-0003-1813-7008
21

22 **Abstract**

G-protein-coupled receptor 30 (GPR30) is a bicarbonate receptor that plays a vital role in
cellular responses to extracellular pH and ion homeostasis. Despite its significance, the
mechanisms by which GPR30 interacts with bicarbonate ions remain elusive. There is no
consensus on a drug that targets GPR30, and the difficulty in the pharmacological analysis
has limited biological and drug discovery researches on GPR30. Here, we present the
cryo-electron microscopy structure of human GPR30 in the presence of bicarbonate ions
at 3.2 Å resolution. Our structure reveals unique extracellular pockets and critical residues
for bicarbonate binding and activation. Functional assays demonstrate that mutations in
these residues impair bicarbonate-induced GPR30 activation, underscoring their
importance in receptor function. This study also provides insights into the G-protein
coupling, highlighting the structural divergence between GPR30 and other GPCRs. Our
findings not only advance the understanding of the role of GPR30 in pH homeostasis but
also pave the way for the development of high-affinity drugs targeting GPR30 for
therapeutic interventions in diseases associated with acid-base imbalance.

38 **Introduction**

The extracellular environment regulates cellular responses, as exemplified by pH
and ion homeostasis. The acid-base balance, largely based on the bicarbonate buffer
system *in vivo*, serves as a vital mechanism that maintains the optimal pH for cellular
function.[1] Changes in the extracellular pH and ion homeostasis are monitored by
membrane channels and receptors, such as G protein-coupled receptors (GPCRs).
However, it remains unknown whether and how the acid–base balance-related ions,
protons and bicarbonate ions, bind to the receptors and cause conformational changes that
lead to intracellular signal transduction.
Previous studies established four acid–base balance-related GPCRs[2–4] that
respond to an increase in proton concentration. None of the structures of these proton-
sensing GPCRs in a proton-bound form has been determined, although the cryo-EM

structures of apo-GPR134 (G2A) and its oxidized fatty acid- and N-acyl amino acid-
bound forms have been reported.[5] While these proton-sensing GPCRs are likely to be
involved in pH homeostasis, particularly in the acidic tumor microenvironment, at
inflamed sites, and in ischemia-reperfusion injury[6], it remains unknown how protons
cooperate with these receptors working as acid–base balance-related GPCRs.

We recently reported that the physiological concentration of bicarbonate ions, a
counterpart of protons in the bicarbonate buffer system, activates G protein-coupled
receptor 30 (GPR30), which leads to G_q-coupled calcium responses[7]. Our study also
demonstrated that GPR30 in brain mural cells regulates blood flow and ischemia–
reperfusion injury. GPR30 was identified as a G protein-coupled estrogen receptor that
mediates the rapid non-genomic action of estradiol (E2)[8]. However, despite numerous
reports on the pleiotropic functions of GPR30 *in vivo*[9,10], there is controversy surrounding
the responses of GPR30 to E2 *in vitro*[11], *ex vivo*[12], and *in vivo*[13]. The broad expression of
Gpr30[14,15], including blood vessels[16,17], stomach, and lung, has also raised the possibility
of its non-estrogenic functions. We demonstrated that three amino acids, E115, Q138,
and H307, are essential for the bicarbonate-induced activation of GPR30, according to
the public homology model (https://gpcrdb.org/)[18]. However, we did not examine whether
bicarbonate ions interact with GPR30 and cause conformational changes. Moreover, to
date, there is no consensus on a high-affinity drug that targets GPR30, and the difficulty
in the pharmacological analysis has limited biological and drug discovery research on
GPR30. To elucidate the bicarbonate–GPR30 interaction and to clarify the first structural
identification of an acid-base balance-related GPCR, we report the cryo-electron
microscopy (cryo-EM) structure of human GPR30 in the presence of bicarbonate ions.

Results

Overall structure

For the structural study, we used the full-length human GPR30 sequence. To
efficiently purify the stable GPCR-G-protein complex, the receptor and mini-G_{sqi} were
incorporated in a 'Fusion-G system' by combining two complex stabilization
techniques[19,20] (Figure 1—figure supplement 1A, B). The modified receptor and G-

protein were co-expressed in HEK293 cells and purified by Flag affinity chromatography

in the presence of 200 mM $NaHCO_3$. After an incubation with nanobody 35 (Nb35) and

single chain scFv16, which binds to mini-G_{sqi}, the complex was purified by size exclusion

chromatography (Figure 1—figure supplement 1C, D). The structure of the purified

complex was determined by single-particle cryo-EM analysis, with an overall resolution

of 3.21 Å (Table 1, Figure 1—figure supplement 2, and "Methods"). As the extracellular

portion of the receptor was poorly resolved, we performed receptor focused refinement,

yielding a density map with a nominal resolution of 3.30 Å, which was combined with

the overall refined map. The resulting composite map allowed us to precisely build the

atomic model of the components, including the receptor (residues 51 to 196, 207 to 288,

and 296 to 340, G-proteins, and scFv16 (Figure 1A, B). Nb35 was not visible in the cryo-

EM map, probably because of the effect of the mini-G_s modification, although the

interactive residues with Nb35 were not changed.

The receptor consists of the canonical 7 transmembrane helices (TM) connected by

three intracellular loops (ICL1–3) and three extracellular loops (ECL1–3), and the

amphipathic helix 8 at the C-terminus (H8) (Figure 2A). Most of the TMs are kinked as

in typical GPCRs[21], but TM1 is also kinked at P71[1.44] (superscripts indicate Ballesteros-

Weinstein numbers[22]), whereas TM1 adopts a straight α-helix in most class A GPCRs.

ICL1 and ECL1 contain short α helices (Figure 2B, C). The short ICL3 was completely

visible in the cryo-EM map (Figure 2C and Figure 1—figure supplement 3), while

residues 197 to 206 in ECL2 and 289 to 295 in ECL3 were disordered (Figure 2B). ECL2

is attached to ECL1 by the disulfide bond between C130[ECL1] and C207[ECL2] (Figure 2B),

which is highly conserved in class A GPCRs[21]. The cryo-EM structure did not

superimpose well on the AlphaFold-predicted structure (Q99527 in the AlphaFold

[19,20]database) with a root mean square deviation (R.M.S.D.) of Cα atoms of 2.33 Å (Figure

2—figure supplement 1A). The conserved $D^{3.49}R^{3.50}Y^{3.51}$ motif[21] in the predicted

structure represents an inactive state (Figure 2—figure supplement 1B). Moreover, ECLs

2 and 3 are rich in cysteine residues, which form incorrect disulfide bonds in the predicted

structure (Figure 2—figure supplement 1C). These comparisons support the usefulness

of experimental structural determination.

Extracellular pockets

The interaction network between ECL1–3 covers the extracellular side of the receptor: ECL2 is anchored to ECL1 via the disulfide bond and ECL3 extends between ECL1 and ECL2, with Q296$^{7.25}$ and F298$^{7.27}$ located within the receptor cavity (Figure 3A). R299^{ECL3} and the backbone carbonyl oxygen of S297^{ECL3} interact with D210^{ECL2}, and N296$^{7.25}$ forms polar interactions with E115$^{2.60}$, Q121$^{2.66}$, and N310$^{7.39}$. These interactions among the ECLs create four extracellular pockets (Pockets A–D) (Figure 3B). While Pockets A–C are positively-charged , Pocket D is negatively, and the size of each pocket is varied. Pocket A consists of ECL1, TM1, and TM7 (Figure 3C), while pocket B is formed by TM6, TM7, and ECL1 (Figure 3D). These two pockets are separated by R122^{ECL1} and H307$^{7.36}$. Pocket A is hydrophobic and appears unsuitable for bicarbonate binding (Figure 3C). Pocket B has hydrophilic residues (Figure 3D), but given the invisible region of ECL3 (residues 289 to 295), we cannot rule out the possibility that the pocket is buried. Pocket C consists of ECL2, ECL3, TM5, and TM6, and is positively charged due to the presence of arginine and histidine residues (Figure 3B, E). However, pocket C is relatively superficial compared to the other pockets. No density corresponding to bicarbonate is observed in these pockets (A–C), indicating that they are not ligand binding sites. Nonetheless, they may serve as potential binding sites for allosteric modulators or agonist drugs targeting GPR30.

Pocket D consists of ECL1–3 and TM2–7, and is the largest among the four pockets. Although the surface of pocket D is partitioned by the C130$^{3.25}$-C207$^{45.52}$ disulfide bond and F208^{ECL2}, it is connected inside the receptor, deep enough to reach W272$^{6.48}$. Pocket D contains numerous hydrophilic residues such as aspartate, glutamate, asparagine, and glutamine, which favor bicarbonate binding, suggesting that pocket D is a good candidate for the bicarbonate binding site.

Insights into bicarbonate binding

To predict the bicarbonate binding site, we performed an exhaustive mutant analysis of hydrophilic residues in pockets A–D, which was not done in the previous study[7]. We confirmed the similar expression levels of the HA-tagged hGPR30 mutants

by western blotting (Figure 4—figure supplement 1A, B). To evaluate the bicarbonate-

induced activation of GPR30, we used the TGFα shedding assay[23]. The HA-tagged

GPR30 wild-type showed a bicarbonate-dependent response, while the H307$^{7.36}$A mutant

completely abolished the response, consistent with previous findings [7](Figure 4A, B).

These results indicate that the present experimental procedure is useful to evaluate the

bicarbonate-induced activation of GPR30. Furthermore, the C207^{ECL2}A mutation that

disrupts the disulfide bond between ECL2 and ECL1, and the P71$^{1.44}$ mutant that disrupts

the characteristic kink in TM1, also abolished the bicarbonate response. These findings

underscore the critical role of the structural integrity of the receptor, and provide

compelling evidence that the bicarbonate response is mediated through GPR30.

Among the hydrophilic residue mutants, the alanine mutants of D125^{ECL1},

S134$^{3.29}$, D210^{ECL2}, and Q215$^{5.39}$ reduced the bicarbonate response (Figure 4A, B).

Combining these data with previous studies, we mapped the residues essential for the

bicarbonate response on the current structure (P71$^{1.44}$, E115$^{2.60}$, D125^{ECL1}. S134$^{3.29}$,

Q138$^{3.33}$, C207^{ECL2}, D210^{ECL2}, Q215$^{5.39}$, and H307$^{7.36}$) (Figure 4C). Except for the

structure-contributing residues (C207^{ECL2}and P71$^{1.44}$) and H307$^{7.36}$, the essential residues

face pocket D, further supporting its role as the bicarbonate binding site. H307$^{7.36}$

separates pockets A and B (Figure 3C), and the nearby mutants Q296$^{7.25}$ and N310$^{7.39}$ had

no effect on the bicarbonate response, suggesting the indirect importance of stabilizing

the active conformation of the receptor, rather than directly binding to bicarbonate.

Given the low affinity of GPR30 for bicarbonate[7] and the challenges posed by

carboxylic acid decarboxylation in cryo-EM analysis, visualizing bicarbonate at this

resolution could be difficult. Nevertheless, we observed a weak density in pocket D,

allowing us to assign bicarbonate (Figure 4D). The bicarbonate is surrounded by Q138$^{3.33}$,

Q215$^{5.39}$, and E218$^{5.42}$ (Figure 4E). Q138$^{3.33}$ and Q215$^{5.39}$ are residues with reduced

bicarbonate responses due to mutations (Figure 4C). Most bicarbonate binding modes in

other structures require recognition by asparagine and glutamine residues, in agreement

with our model. Moreover, Q138$^{3.33}$, Q215$^{5.39}$, and E218$^{5.42}$ are evolutionarily conserved

in fish to human GPR30s（Figure 4—figure supplement 2). Below the bicarbonate

binding site, M141$^{3.36}$ forms a hydrophobic interaction with W272$^{6.48}$ (Figure 4D), a

toggle switch motif essential for receptor activation[21,24]. These observations allowed us

to speculate that bicarbonate binding induces conformational changes in TM3 and TM5,

facilitating receptor activation.

G-protein coupling

The C-terminal helix of $G\alpha_q$ ($\alpha5$-helix) is deeply enters the intracellular cavity of

GPR30, resulting in the formation of an active signaling complex. On the C-terminal side

of the $\alpha5$-helix, the backbone carbonyl of Y243[G.H5.23] (superscript indicates the common

$G\alpha$ numbering [CGN] system[25]) hydrogen bonds with R155[3.50] in the conserved

D[3.49]R[3.50]Y[3.51] motif[21] (Figure 5A). In addition, N244[G.H5.24] forms hydrogen bonds with

the peptide backbones of Y324[7.53] and F331[8.50]. ICL3 faces the middle part of the $\alpha5$-

helix, with three characteristic arginine residues (R248[6.24], R251[6.27], and R254[6.30])

surrounding D233[G.H5.13] (Figure 5B). R248[6.24] and R254[6.27] form hydrogen bonds with

D233[G.H.13] and Q237[G.H5.17], respectively. Moreover, R251[6.30] extends toward the G-

protein and hydrogen bonds with the peptide backbones of G207[G.h4s6.11] and H209[G.h4s6.13].

At ICL2, M163[34.51] fits into a hydrophobic pocket composed of the $\alpha5$-helix, the αN-$\beta2$

loop, and the $\beta2$-$\beta3$ loop of the $G\alpha$ subunit, as in other GPCR-G_q complexes[26–30] (Figure

5C–H). Above it, L159[3.54] and A162[34.50] form extensive hydrophobic interactions with

L235[G.H5.15], L236[G.H5.16], and L240[G.H5.20] in the $\alpha5$-helix. These tight interactions with the

$\alpha5$-helix would enable the coupling of GPR30 with G_q.

The position of the $\alpha5$-helix aligns with those in other GPCR-G_q complexes, but

the position of the αN-helix is rotated away from the receptor (Figure 5I). The

superimposition of the $G\alpha$ subunits indicates that the rotation is not due to the movement

of the entire G protein, but rather a 33° bend in the αN-helix (Figure 5J). This is attributed

to variations in the interactions of the αN-helix with its receptors. The αN-helix interacts

with the intracellular ends of ICL2 and TM4 in receptor-specific modes. In most G_q-

coupled structures, ICL2 adopts an α-helix and R[34.55] interacts with R38[G.hns1.03] at the end

of the αN-helix. A typical example is the head-to-toe interaction between R185[34.55] and

R32[G.hns1.03] in the serotonin 5-HT$_{2A}$ receptor[29] (Figure 5E). However, in GPR30, ICL2

does not adopt an α-helix and R[34.55] is replaced by L167[34.55], which displaces R32[G.hns1.03]

downward through their interaction. Moreover, T170[34.58] and R169[34.57] form direct

hydrogen bonds with R31[G.hns1.02] and Y243[G.H5.23], respectively, which are not observed in

other GPCR-G_q complexes. We hypothesize that these interactions may underpin the

distinct αN rotation: A similar interaction between R38[G.hns1.03] and L136[34.54] is also

observed in histamine H1 receptor (H_1R)[26] (Figure 5D), in which translational rather than

rotational motion of αN is observed (Figure 5J).

Discussion

Our study has elucidated the distinctive structure of GPR30, characterized by the

kink of TM1 and the presence of multiple extracellular pockets formed by ECLs.

Structural analysis and unbiased mutagenesis experiments on hydrophilic residues

highlighted the largest pocket, designated as pocket D, as pivotal for bicarbonate

recognition. Through putative bicarbonate modeling, we have proposed the binding mode

of bicarbonate ions and the receptor activation mechanism. Overall, the elucidated

structure of the extracellular domain of GPR30 and the insights into bicarbonate binding

could facilitate the development of high-affinity drugs targeting GPR30.

It should be noted that our current model cannot fully explain all the mutagenesis

results, such as those involving E115[2.60], D125[ECL1]. and D210[ECL2] [7]. Although negatively

charged residues generally repel bicarbonate, some structures have indicated their

involvement in bicarbonate recognition[31–34]. Thus, it is conceivable that these residues are

transiently involved in bicarbonate binding, or two bicarbonate molecules might be

accommodated within pocket D of GPR30. Future studies should focus on higher-

resolution structures and approaches such as using bicarbonate derivatives to further

evaluate bicarbonate binding.

GPR30 exhibits low homology to other GPCRs, with even closely related

receptors sharing less than 30% sequence identity. A BLAST analysis revealed that

GPR30 shares the highest sequence identity of 28% with the type 2 angiotensin II receptor

(AT_2R) (Figure 6—figure supplement 1). Consequently, we compared the current

structure of GPR30 with the ligand-bound AT_2R[35]. The superimposition of GPR30 and

AT_2R, with a Cα R.M.S.D. of 2.17 Å, suggests significant structural divergence between

the two (Figure 6A). Notably, there is no structural commonality between GPR30 and AT2R in terms of the orientations of their respective TMs. Specifically, the extracellular half of TM1 in GPR30 extends outward, and is kinked at P71$^{1.44}$ in the center of TM1 (Figure 6B). Near P71$^{1.44}$, Y65$^{1.38}$ and F70$^{1.43}$ are oriented towards the interior of the receptor, forming hydrophobic interactions with L113$^{2.58}$ and L331$^{7.40}$ to stabilize the kink. In the examination of the loops, ECL2 of AT$_2$R features a short β-sheet commonly found in peptide-activated class A GPCRs[36–38], whereas that of GPR30 adopts an elongated conformation that covers the extracellular pocket essential for the bicarbonate response (Figure 4A, B). These comparisons underscore the unique structural characteristics of GPR30.

In most GPCRs, TM1 forms a straight α-helix and is positioned away from the orthosteric pocket formed by TM2–7. However, TM1 undergoes an inward movement due to allosteric conformational changes during agonist binding. The cannabinoid receptor CB$_1$ exemplifies this, with TM1 kinked at G127$^{1.43}$ and folding inward upon ligand binding [39,40](Figure 6C). Similarly, the β$_2$ adrenergic receptor (β$_2$AR)[41,42] and adenosine A2A receptor (A$_{2A}$R)[43,44] exhibit inward movements of TM1, with TM1 bending at the glycine residues rather than kinking (Figure 6D, E). By contrast, TM1 [24,45]moves 4 Å outward in the endothelin ET$_B$ receptor (Figure 6F). In GPR30, TM1 is kinked at P71$^{1.44}$ (Figure 6B), indicating that this kink occurs regardless of the receptor activation state. Indeed, the P71$^{1.44}$A mutation, which eliminates the kink in TM1, completely abolished the bicarbonate response (Figure 4A, B). Moreover, P71$^{1.44}$ is evolutionarily preserved (Figure 4—figure supplement 2). These findings suggest the critical role of the TM1 kink in both the putative conformational changes associated with ligand binding and the functional integrity of GPR30.

Table. 1   Cryo-EM data collection, refinement and validation statistics

Data collection	GPR30-G$_q$ complex

Microscope	Titan Krios (Thermo Fisher Scientific)
Voltage (keV)	300
Electron exposure (e$^-$/Å^2)	50.660
Detector	Gatan K3 Summit camera (Gatan)
Magnification	×105,000
Defocus range (μm)	-0.8–1.6
Pixel size (Å/pix)	0.83
Number of movies	9,824
Symmetry	C1
Picked particles	11,176,478
Final particles	148,285
Map resolution (Å)	3.21
FSC threshold	0.143
Model refinement	
Atoms	
R.m.s. deviations from ideal	
Bond lengths (Å)	0.004
Bond angles (°)	0.633
Validation	
Clashscore	9.17
Rotamers Outliers(%)	0.34
Ramachandran plot	
Favored (%)	93.77
Allowed (%)	6.23
Outlier (%)	0

Figures

Figure 1. Overall structure of the GPR30-miniG$_{sqi}$β$_1$γ$_2$-scFv16 complex.

A Unsharpened cryo-EM density map of the GPR30-miniG$_{sqi}$β$_1$γ$_2$-scFv16-Nb35 complex, with the components individually colored. **B** The refined structure of the complex is shown as a ribbon representation.

Figure 2. Receptor structure.

A–C Overall structure of the receptor, viewed from the membrane plane (**A**), intracellular side (**B**), and extracellular side (**C**). **D** Extracellular view of the ribbon model overlaid the cryo-EM map.

Figure 3. Architecture of the extracellular pocket.

A Interactions between the ECLs. Black dashed lines indicate hydrogen bonds. **B** Molecular surface of the extracellular side. **C–F** Residues facing pocket A (**C**), pocket B (**D**), pocket C (**E**), and pocket D (**F**).

Figure 4. Mutational analysis.

A, B TGFα shedding assay using HEK293 cells transfected with HA-tagged hGPR30. The mutants D111A; C207A, P71A, H307A; and D125A, S134A, D210A, Q215A are highlighted in red, blue, and purple, respectively. Statistical analysis: [$] $p = 0.0001$, [#] $p < 0.0001$ compared to mock cells using two-tailed unpaired t-test with Bonferroni's correction after two-way ANOVA. ns indicates no significant difference. Data are presented as mean values (**A**) and mean values ± SEM (**B**). **C** Residues that were subjected to mutant analysis are mapped to the structure. Only residues with reduced bicarbonate responses are colored.in magenta. **D** Density corresponding to bicarbonate. **E** Putative binding mode of bicarbonate. Black dashed lines indicate hydrogen bonds.

Figure 5. G-protein coupling.

A Hydrogen-bonding interactions between the C-terminal α5-helix residues and the

receptor. **B** Electrostatic and hydrogen-binding interactions between ICL3 and the α5-
helix. **C**–**H** Interface between ICL2 and G_q, with residues involved in hydrophobic
interactions represented by CPK models. G-protein-bound GPCRs used in the
comparison are as follows: H_1R-G_q (PDB 7DFL, gray), B_2R-G_q (PDB 7F6I, yellow-
green), MRGPRX$_2$-G_q (PDB 7S8L, green), 5-HT$_{2A}$-G_q (PDB 6WHA, gray), and
GPR103-G_q (PDB 8ZH8, red). **I** Comparison of the angles and positions of α5h and αN
relative to the receptor. **J** Superimposition of the Gα subunits.

Figure 6. Structural comparison with related GPCRS.

A Structural comparison of GPR30 and AT$_2$R (PDB 5UNF). **B** Interactions around P70
in TM1 of GPR30. **C**–**F** Conformational changes of TM1 upon agonist binding in CB1
(**C**), β$_2$AR (**D**), A$_{2A}$R (**E**), and ET$_B$ (**F**). The agonist-bound states are colored with the
respective colors, while the inactive states are colored gray. The PDB codes used in this
figure are CB1-active (PDB 5XRA), CB$_1$-inactive (PDB 5TGZ), β$_2$AR-active (PDB
3SN6), β$_2$AR-inactive (PDB 2RH1), A$_{2A}$R-active(PDB 6GDG), A$_{2A}$R-inactive (PDB
3EML), ET$_B$-active (PDB 8IY5), and ET$_B$-inactive (PDB 5X93).

Figure 1.—figure supplement 1. Sample purification.
A Schematic representation of the fusion-G system. **B** Fluorescence-detection size-
exclusion chromatography (FSEC) analysis of complex formation by GPR30. Solubilized
cells expressing only GPR30 are orange, and those co-expressing GPR30 and G-protein
are blue. **C** Size-exclusion chromatography of the GPR30-G-protein complex on a
Superose 6 Increase column. The fraction shaded in blue was collected. **D** SDS-PAGE
gel of samples after size-exclusion chromatography, stained with Coomassie Brilliant
Blue.
307

308   **Figure 1.—figure supplement 2.** Cryo-EM analysis.
309   Flow chart of the cryo-EM data processing for the GPR30-G_q complex, including particle
310   projection selection, classification, and 3D density map reconstruction. The 3D density
311   map was refined with a mask on the receptor. Details are provided in the Methods section.
312

313 **Figure 3.—figure supplement 1.** Structural comparison with AlphaFold-predicted
314 structure.
315 **A–C** Superimposition of the cryo-EM (orange) and AF2 (khaki) structures. (**A**) Overall
316 view of the receptor, (**B**) focused on the intracellular side, and (**C**) focused on the
317 extracellular side.
318
319 **Figure 4.—figure supplement 1.** Cell surface expression of mutant receptors.
320 **A** Whole cell expression of HA-tagged mutants, analyzed by western blotting. **B** Cell
321 surface expression of HA-tagged mutants, using cell surface biotinylation and avidin
322 immunoprecipitation, analyzed by western blotting.
323
324 **Figure 4.—figure supplement 2.** Conservation of the GPR30 homologs.
325 Amino acid alignment of representative GPR30 homologs.
326
327 **Figure 6.—figure supplement 1.** Sequence alignment of GPR30 and AT_2R.
328

Data Availability

330   The cryo-EM density map and atomic coordinates for the LPA_1-G_i complex have
331 been deposited in the Electron Microscopy Data Bank and the PDB, under the accession
332 codes: EMD-XXXXX [https://www.ebi.ac.uk/emdb/entry/EMD-XXXXX], and PDB
333 XXXX [http://doi.org/10.2210/pdbXXXX/pdb]. Source data are provided with this paper.
334 All other data are available from the corresponding authors upon reasonable request.
335

Acknowledgements

We thank K. Ogomori and C. Harada for technical assistance. This work was
supported by grants from the JSPS KAKENHI, grant numbers 21H05037 (O.N.),
22K19371 and 22H02751 (W.S.), 23K06393 (A.J-W.); the Japan Agency for Medical
Research and Development (AMED), Innovation and Clinical Research Center Project,
PRIME, grant number JP20gm6210026 (A.J.-W.); the JST FOREST Program (Grant

number JPMJFR220S to A.J.-W.); the Takeda Science Foundation (W.S.); the Lotte

Foundation (W.S.); AMED, grant number JP233fa627001 (O.N.); and the Platform

Project for Supporting Drug Discovery and Life Science Research (Basis for Supporting

Innovative Drug Discovery and Life Science Research (BINDS)) from AMED, grant

numbers JP23ma121002 (support number 3272, O.N.) and JP23ama121012 (supporting

number 4869, T.Y.).

Author contributions

S.K. performed all the experiments involved in the structural determination,

assisted by H.A. and H.S.O. W.S. designed the project and initially tried the expression

of GPR30. A.J.-W. and T.Y. performed and oversaw the mutagenesis study. The

manuscript was mainly prepared by S.K. and W.S., with assistance from O.N.

Competing interests

O.N. is a co-founder and scientific advisor for Curreio. All other authors declare

no competing interests.

Methods

Expression and purification of scFv16 and Nb35

The gene encoding scFv16 was synthesized (GeneArt) and subcloned into a

modified pFastBac vector[46], with the resulting construct encoding the GP67 secretion

signal sequence at the N terminus, and a His$_8$ tag followed by a TEV cleavage site at the

C terminus. The His$_8$-tagged scFv16 was expressed and secreted by Sf9 insect cells, as

previously reported[44]. The Sf9 cells were removed by centrifugation at 5,000g for 10 min,

and the secreta-containing supernatant was combined with 5 mM CaCl$_2$, 1 mM NiCl$_2$,

20 mM HEPES (pH 8.0), and 150 mM NaCl. The supernatant was mixed with Ni

Superflow resin (GE Healthcare Life Sciences) and stirred for 1 h at 4 °C. The collected

resin was washed with buffer containing 20 mM HEPES (pH 8.0), 500 mM NaCl and

20 mM imidazole, and further washed with 10 column volumes of buffer containing

20 mM HEPES (pH 8.0), 500 mM NaCl and 20 mM imidazole. Next, the protein was

eluted with 20 mM Tris (pH 8.0), 500 mM NaCl and 400 mM imidazole. The eluted

fraction was concentrated and loaded onto a Superdex200 10/300 Increase size-exclusion

column, equilibrated in buffer containing 20 mM Tris (pH 8.0) and 150 mM NaCl. Peak

fractions were pooled, concentrated to 5 mg/ml using a centrifugal filter device (Millipore

10 kDa MW cutoff), and frozen in liquid nitrogen.

Nb35 was prepared as previously reported[47,48]. In brief, Nb35 was expressed in

the periplasm of *E. coli*. The harvested cells were disrupted by sonication. Nb35 was

purified by nickel affinity chromatography, followed by gel-filtration chromatography,

and frozen in liquid nitrogen.

Constructs for expression of GPR30 and G_q

The human GPR30 gene (UniProtKB, Q99527) was subcloned into a modified

pFastBac vector[49], with an N-terminal haemagglutinin signal peptide followed by the

Flag-tag epitope (DYKDDDDK) and the LgBiT fused to its C-terminus followed by a 3

C protease site and EGFP-His$_8$ tag. A 15 amino sequence of GGSGGGGSGGSSSGG was

inserted into both the N-terminal and C-terminal sides of LgBiT. Rat $G\beta_1$ and bovine $G\gamma_2$

were subcloned into the pFastBac Dual vector. In detail, rat $G\beta_1$ was cloned with a C-

terminal HiBiT connected with a 15 amino sequence of GGSGGGGSGGSSSGG.

Moreover, mini-G_{sqi} was subcloned into the C-terminus of the bovine $G\gamma_2$ with a nine

amino sequence GSAGSAGSA linker. The resulting pFastBac dual vector can express

the mini-G_{sqi} trimer.

Expression and purification of the human GPR30 – G_q `complex`

The recombinant baculovirus was prepared using the Bac-to-Bac baculovirus

expression system (Thermo Fisher Scientific). For expression, 2 L of Sf9 cells at a density

of 3×10^6 cells/mL were co-infected with baculovirus encoding GPR30 and miniGsqi

trimer at the ratio of 1:1 and the cells were incubated at 30°C. After 48 hours, the collected

cells were resuspended and dounce-homogenized in 20 mM Tris-HCl, pH 8.0(4°C),
200 mM NaCl, 10% Glycerol, 200 mM NaHCO₃, 25 mU/ml apyrase. The crude
membrane fraction was collected by ultracentrifugation at 180,000g for 1 h and
solubilized in buffer, containing 20 mM Tris-HCl, pH 8.0, 150 mM NaCl, 1% n-dodecyl-
beta-D-maltopyranoside (DDM) (Calbiochem), 0.2 % cholesteryl hemisuccinate (CHS)
(Merck), 10% glycerol, 200 mM NaHCO₃, and 25 mU/ml Apyrase, for 1 h at 4 °C. The
supernatant was separated from the insoluble material by ultracentrifugation at 180,000g
for 20 min and incubated with 5 mL of Anti-DYKDDDDK M2 resin (Sigma) for 1 h at
4°C. The resin was washed with 20 column volumes of buffer containing 20 mM Tris-
HCl, pH 8.0 500 mM NaCl, 10% Glycerol, 0.05% Lauryl Maltose Neopentyl Glycol
(LMNG) (Antrace), 0.005% CHS and 200 mM NaHCO₃. The complex was eluted in
buffer containing 20 mM Tris-HCl, pH 8.0, 150 mM NaCl, 10% Glycerol, 0.01% LMNG,
0.001% CHS, 200 mM NaHCO₃ and 0.2 mg/mL Flag peptide. The eluate was incubated
with the Nb35 and scFv16 at 4 °C. The complex was concentrated and purified by size
exclusion chromatography on a Superose 6 increase (GE) column in 20 mM Tris-HCl,
pH8.0, 150 mM NaCl, 0.01% LMNG, 0.001% CHS and 200 mM NaHCO₃. Peak fractions
were concentrated to 4.72 mg/ml for electron microscopy studies.
415

Sample vitrification and cryo-EM data acquisition

The purified complex was applied onto a freshly glow-discharged
Quantifoil UltraAu grid (R1.2/1.3, 300 mesh), and plunge-frozen in liquid ethane by
using a Vitrobot Mark IV. Data collections were performed on a 300kV Titan Krios G3i
microscope (Thermo Fisher Scientific) equipped with a BioQuantum K3 imaging filter
and a K3 direct electron detector (Gatan).
A total of 9824 movies were acquired with a calibrated pixel size of 0.83 Å pix^{-1}
and with a defocus range of -0.8 to -1.6 μm, using EPU. Each movie was acquired for
2.6 s and split into 48 frames, resulting in an accumulated exposure of about
50.660 electrons per Å².
426

Image processing

All acquired dose-fractionated movies were imported into CryoSPARC v4.4[50] and

subjected to beam-induced motion correction. The contrast transfer function (CTF)

parameters were estimated and a total of 10,148,422 particles were extracted. The

particles were subjected to 2D classifications, Ab-initio reconstruction and several rounds

of hetero refinement and Non-uniform refinement. Next, the particles were subjected to

3D classification with a mask on the receptor. Then the particle sets were subjected to

Reference Based Motion Correction. Motion-corrected 148,285 particles were subjected

to Non-uniform refinement, yielding a map with a global nominal resolution of 3.21 Å,

with the gold standard Fourier Shell Correlation (FSC=0.143) criteria[51]. Moreover, the

3D model was refined with a mask on the receptor. As a result, the receptor has a higher

resolution with a nominal resolution of 3.30 Å. The overall and receptor focused maps

were combined by phenix[52]. The processing strategy is described in Supplementary

Figure 2.

Model building and refinement

The density map was sharpened by phenix.auto_sharpen[53] and the quality of the

density map was sufficient to build a model manually in COOT[54,55]. The model building

was facilitated by the Alphafold-predicted structure. We manually fitted GPR30, the

G_q heterotrimer and scFv16 into the map. We then manually readjusted the model into

using COOT and refined it using phenix.real_space_refine[52,56] (v.1.19) with the

secondary-structure restraints using phenix secondary_structure_restraints.

Vector construction and transfection

Human *Gpr30* cDNA was obtained from human hepatoblastoma-derived HepG2 cells.

The coding sequences of human *Gpr30* were inserted into the multi-cloning site of the

plasmid vector pCXN2, which was generated in our laboratory via modification of

pCAGGS, between the KpnI and EcoRI sites. The C-terminal HA-tagged *Gpr30* was

amplified using a reverse primer containing the HA sequenceOne amino acid mutation

of human GPR30 was generated as follows: the targeted amino acid was changed to

alanine (GCC) using a two-step PCR method with the QuikChange® Primer Design

Program by Agilent (https://www.agilent.com/store/primerDesignProgram.jsp), and the

coding sequence with each mutation was inserted into the multi-cloning site of pCXN2

between the KpnI and EcoRI sites. The mutations generated were D111A, E121A, R122A,

D125A, S134A, D210A, Q215A, N276A, Q296A, C207A, P71A, and H307A.

Cell line sources and transfection

463   HEK293A cells (female origin; Thermo Fisher Scientific) were maintained at 37 °C and

464   5% CO_2. These vectors were transfected using the lipofection method (Lipofectamine™

465   2000 Transfection Reagent, 11668019, Invitrogen).

TGFα shedding assay

The TGFα shedding assay was performed according to a previously published protocol[23].

AP-TGFα expression vector was provided by Dr. Inoue and Dr. Aoki, Tohoku University.

HEK293 cells were seeded in 12-well plates at a density of 1×10^5 cells/well and cultured

for 24 h. At 70% confluency, a mixture of plasmid vectors containing GPCR (see '***Vector***

construction and transfection' section) and AP-TGFα was transfected into the cells using

Lipofectamine 2000 transfection reagent. After another 24 h of incubation, the cells were

detached with 0.05% trypsin/EDTA (32777-44, Nacalai Tesque), suspended in Hanks'

balanced salt solution, and seeded in 96-well plates. The cells were stimulated for 1 h

with $1–44 \times 10^{-3}$ M (final concentration) of $NaHCO_3$ at 37 °C, under 0.03% CO_2.

Conditioned media (CM) was transferred to another plate, and 80 µl of alkaline

phosphatase (AP) solution (40 mM Tris-HCl, pH 9.5, 40 mM NaCl, 10 mM $MgCl_2$, 10

mM p-nitrophenylphosphate disodium salt hexahydrate) was added to both plates, which

were then incubated at 37 °C. The optical density at 405 nm (OD405) was measured using

a microplate reader (iMark, Bio-Rad) at 5 min, 30 min, 1 h, and 2 h, depending on the

reaction rate. The percentage of shed AP-TGF was calculated using the following

equations:

`AP activity = ∆OD405 (1 − 0 h)`

% CM (conditioned media) = AP_{CM} / (AP_{CM} + AP_{Cell})

[15]**Analysis of cell surface expression of GPR30**

The cell surface expression of wild-type and mutant hGPR30-HA was analysed via cell

surface protein isolation using a Cell Surface Protein Isolation Kit (#89881, Thermo

Scientific™) followed by western blotting. Briefly, HEK293 cells transiently expressing

wild-type or mutant hGPR30-HA were biotinylated for 30 min at 4 °C. Then, the cells

were harvested, and biotinylated proteins were isolated with avidin binding.

Western blotting

Cell lysates were separated using sodium dodecyl sulfate-polyacrylamide gel

electrophoresis under reducing conditions and transferred to polyvinylidene difluoride

membranes (Immobilon P, IPVH00010, Millipore). Primary antibodies used were anti-

HA High Affinity, 1:1000 dilution, 11867423001, Roche; Na-K-ATPase, 1:1000 dilution,

#3010, Cell Signaling Technology; β-Actin (AC-15), 1:1000 dilution, sc-69879, Santa

Cruz Biotechnology. Secondary antibodies used were anti-rabbit IgG, HRP-linked

Antibody, 1:5000 dilution, #7074, Cell Signaling Technology; anti-mouse IgG, HRP-

linked Antibody, 1:5000 dilution, #7076, Cell Signaling Technology; anti-rat IgG, HRP-

Linked Whole Ab Goat, 1:5000 dilution, NA935, Cytiva. The membranes were probed at

4 °C overnight with the primary antibodies. The membranes were subsequently incubated

with the corresponding secondary antibodies. The signals were detected with ECL Prime

(RPN2236, Cytiva) or ImmunoStar LD (296-69901, FUJIFILM Wako Pure Chemical

Corporation) using a chemiluminescence imaging system (CFusion FX7, Vilber).

1. Brinkman, J. E. & Sharma, S. Physiology, Metabolic Alkalosis. in *StatPearls*

(StatPearls Publishing, Treasure Island (FL), 2024).

2. Ludwig, M.-G. *et al.* Proton-sensing G-protein-coupled receptors. *Nature* **425**, 93–98

(2003).

3. Murakami, N., Yokomizo, T., Okuno, T. & Shimizu, T. G2A is a proton-sensing G-
protein-coupled receptor antagonized by lysophosphatidylcholine. *J Biol Chem* **279**,
42484–42491 (2004).

4. Wang, J.-Q. *et al.* TDAG8 is a proton-sensing and psychosine-sensitive G-protein-
coupled receptor. *J Biol Chem* **279**, 45626–45633 (2004).

5. Wang, J.-L. *et al.* Functional screening and rational design of compounds targeting
GPR132 to treat diabetes. *Nat Metab* **5**, 1726–1746 (2023).

6. Imenez Silva, P. H., Camara, N. O. & Wagner, C. A. Role of proton-activated G
protein-coupled receptors in pathophysiology. *Am J Physiol Cell Physiol* **323**, C400–
C414 (2022).

7. Jo-Watanabe, A. *et al.* Bicarbonate signalling via G protein-coupled receptor
regulates ischaemia-reperfusion injury. *Nat Commun* **15**, 1530 (2024).

8. Revankar, C. M., Cimino, D. F., Sklar, L. A., Arterburn, J. B. & Prossnitz, E. R. A
transmembrane intracellular estrogen receptor mediates rapid cell signaling. *Science*
307, 1625–1630 (2005).

9. Ford, J. *et al.* GPR30 deficiency causes increased bone mass, mineralization, and
growth plate proliferative activity in male mice. *J Bone Miner Res* **26**, 298–307
(2011).

10. Sharma, G. *et al.* GPER deficiency in male mice results in insulin resistance,
dyslipidemia, and a proinflammatory state. *Endocrinology* **154**, 4136–4145 (2013).

11. Tutzauer, J. *et al.* Ligand-Independent G Protein-Coupled Estrogen Receptor/G
Protein-Coupled Receptor 30 Activity: Lack of Receptor-Dependent Effects of G-1
and 17β-Estradiol. *Mol Pharmacol* **100**, 271–282 (2021).

12. Lamprecht, M. R. & Morrison, B. GPR30 activation is neither necessary nor
sufficient for acute neuroprotection by 17β-estradiol after an ischemic injury in
organotypic hippocampal slice cultures. *Brain Res* **1563**, 131–137 (2014).

13. Otto, C. *et al.* GPR30 does not mediate estrogenic responses in reproductive organs
in mice. *Biol Reprod* **80**, 34–41 (2009).

14. Uhlén, M. *et al.* Proteomics. Tissue-based map of the human proteome. *Science* **347**,
1260419 (2015).

15. The Human Protein Atlas. https://www.proteinatlas.org/.

16. Vanlandewijck, M. *et al.* A molecular atlas of cell types and zonation in the brain
vasculature. *Nature* **554**, 475–480 (2018).

17. Winkler, E. A. *et al.* A single-cell atlas of the normal and malformed human brain
vasculature. *Science* **375**, eabi7377 (2022).

18. Kooistra, A. J. *et al.* GPCRdb in 2021: integrating GPCR sequence, structure and
function. *Nucleic Acids Res* **49**, D335–D343 (2021).

19. Sano, F. K., Akasaka, H., Shihoya, W. & Nureki, O. Cryo-EM structure of the
endothelin-1-ETB-Gi complex. *Elife* **12**, e85821 (2023).

20. Shihoya, W., Iwama, A., Sano, F. K. & Nureki, O. Cryo-EM advances in GPCR
structure determination. *J Biochem* mvae029 (2024) doi:10.1093/jb/mvae029.

21. Venkatakrishnan, A. J. *et al.* Molecular signatures of G-protein-coupled receptors.
Nature **494**, 185–194 (2013).

22. Ballesteros, J. A. & Weinstein, H. [19] Integrated methods for the construction of
three-dimensional models and computational probing of structure-function relations
in G protein-coupled receptors. in *Methods in Neurosciences* (ed. Sealfon, S. C.) vol.

25 366–428 (Academic Press, 1995).

23. Inoue, A. *et al.* TGFα shedding assay: an accurate and versatile method for detecting

GPCR activation. *Nat Methods* **9**, 1021–1029 (2012).

24. Shihoya, W., Sano, F. K. & Nureki, O. Structural insights into endothelin receptor

signalling. *J Biochem* **174**, 317–325 (2023).

25. Flock, T. *et al.* Universal allosteric mechanism for Gα activation by GPCRs. *Nature*

524, 173–179 (2015).

26. Xia, R. *et al.* Cryo-EM structure of the human histamine H1 receptor/Gq complex.

Nat Commun **12**, 2086 (2021).

27. Shen, J. *et al.* Cryo-EM structures of human bradykinin receptor-Gq proteins

complexes. *Nat Commun* **13**, 714 (2022).

28. Cao, C. *et al.* Structure, function and pharmacology of human itch GPCRs. *Nature*

600, 170–175 (2021).

29. Kim, K. *et al.* Structure of a Hallucinogen-Activated Gq-Coupled 5-HT2A Serotonin

Receptor. *Cell* **182**, 1574-1588.e19 (2020).

30. Iwama, A. *et al.* Structure and dynamics of the RF-amide QRFP receptor GPR103.

2023.12.06.570340 Preprint at https://doi.org/10.1101/2023.12.06.570340 (2023).

31. Capper, M. J. *et al.* Substrate binding and inhibition of the anion exchanger 1

transporter. *Nat Struct Mol Biol* **30**, 1495–1504 (2023).

32. Fang, S. *et al.* Molecular mechanism underlying transport and allosteric inhibition of

bicarbonate transporter SbtA. *Proc Natl Acad Sci U S A* **118**, e2101632118 (2021).

33. Koropatkin, N. M., Koppenaal, D. W., Pakrasi, H. B. & Smith, T. J. The structure of

a cyanobacterial bicarbonate transport protein, CmpA. *J Biol Chem* **282**, 2606–2614

(2007).

34. Wang, W. *et al.* Cryo-EM structure of the sodium-driven chloride/bicarbonate

exchanger NDCBE. *Nat Commun* **12**, 5690 (2021).

35. Zhang, H. *et al.* Structural basis for selectivity and diversity in angiotensin II

receptors. *Nature* **544**, 327–332 (2017).

36. Hong, C. *et al.* Structures of active-state orexin receptor 2 rationalize peptide and

small-molecule agonist recognition and receptor activation. *Nat Commun* **12**, 815

(2021).

37. Mobbs, J. I. *et al.* Structures of the human cholecystokinin 1 (CCK1) receptor bound

to Gs and Gq mimetic proteins provide insight into mechanisms of G protein

selectivity. *PLoS Biol* **19**, e3001295 (2021).

38. Tang, T. *et al.* Receptor-specific recognition of NPY peptides revealed by structures

of NPY receptors. *Sci Adv* **8**, eabm1232 (2022).

39. Hua, T. *et al.* Crystal structures of agonist-bound human cannabinoid receptor CB1.

Nature **547**, 468–471 (2017).

40. Hua, T. *et al.* Crystal Structure of the Human Cannabinoid Receptor CB1. *Cell* **167**,

750-762.e14 (2016).

41. Cherezov, V. *et al.* High-resolution crystal structure of an engineered human beta2-

adrenergic G protein-coupled receptor. *Science* **318**, 1258–1265 (2007).

42. Rasmussen, S. G. F. *et al.* Crystal structure of the β2 adrenergic receptor-Gs protein

complex. *Nature* **477**, 549–555 (2011).

43. García-Nafría, J., Lee, Y., Bai, X., Carpenter, B. & Tate, C. G. Cryo-EM structure of

the adenosine A2A receptor coupled to an engineered heterotrimeric G protein. *Elife*

7, e35946 (2018).

44. Jaakola, V.-P. *et al.* The 2.6 angstrom crystal structure of a human A2A adenosine

receptor bound to an antagonist. *Science* **322**, 1211–1217 (2008).

45. Shihoya, W. *et al.* Activation mechanism of endothelin ETB receptor by endothelin-

1. *Nature* **537**, 363–368 (2016).

46. Okamoto, H. H. *et al.* Cryo-EM structure of the human MT1-Gi signaling complex.

Nat Struct Mol Biol **28**, 694–701 (2021).

47. Nagiri, C. *et al.* Cryo-EM structure of the β3-adrenergic receptor reveals the

molecular basis of subtype selectivity. *Mol Cell* **81**, 3205-3215.e5 (2021).

48. Nureki, I. *et al.* Cryo-EM structures of the β3 adrenergic receptor bound to solabegron

and isoproterenol. *Biochem Biophys Res Commun* **611**, 158–164 (2022).

49. Sano, F. K., Akasaka, H., Shihoya, W. & Nureki, O. Cryo-EM structure of the

endothelin-1-ETB-Gi complex. *eLife* **12**, e85821 (2023).

50. Punjani, A., Rubinstein, J. L., Fleet, D. J. & Brubaker, M. A. cryoSPARC: algorithms

for rapid unsupervised cryo-EM structure determination. *Nat Methods* **14**, 290–296

(2017).

51. Rosenthal, P. B. & Henderson, R. Optimal determination of particle orientation,

absolute hand, and contrast loss in single-particle electron cryomicroscopy. *J Mol*

Biol **333**, 721–745 (2003).

52. Adams, P. D. *et al.* PHENIX: a comprehensive Python-based system for

macromolecular structure solution. *Acta Crystallogr D Biol Crystallogr* **66**, 213–221

(2010).

53. Terwilliger, T. C., Sobolev, O. V., Afonine, P. V. & Adams, P. D. Automated map

sharpening by maximization of detail and connectivity. *Acta Crystallogr D Struct*

Biol **74**, 545–559 (2018).

54. Emsley, P. & Cowtan, K. Coot: model-building tools for molecular graphics. *Acta*

Crystallogr D Biol Crystallogr **60**, 2126–2132 (2004).

55. Emsley, P., Lohkamp, B., Scott, W. G. & Cowtan, K. Features and development of

Coot. *Acta Crystallogr D Biol Crystallogr* **66**, 486–501 (2010).

56. Afonine, P. V. *et al.* Real-space refinement in PHENIX for cryo-EM and

crystallography. *Acta Crystallogr D Struct Biol* **74**, 531–544 (2018).

Figure 1. Overall structure of the GPR30-miniG$_{sqi}$β$_1$γ$_2$-scFv16 complex.
A Unsharpened cryo-EM density map of the GPR30-miniG$_{sqi}$β$_1$γ$_2$-scFv16-Nb35 complex, with the components individually colored. **B** The refined structure of the complex is shown as a ribbon representation.

Figure 2. Receptor structure.
A–C Overall structure of the receptor, viewed from the membrane plane (**A**),
extracellular side (**B**), and intracellular side (**C**).**D** Extracellular view of the ribbon model
overlaid the cryo-EM map.

Figure 3. Architecture of the extracellular pocket.
A Interactions between the ECLs. Black dashed lines indicate hydrogen bonds. **B** Molecular surface of the extracellular side. **C–F** Residues facing pocket A (**C**), pocket B (**D**), pocket C (**E**), and pocket D (**F**).

Figure 4. Mutational analysis.
A, B TGFα shedding assay using HEK293 cells transfected with HA-tagged hGPR30. The mutants D111A; C207A, P71A, H307A; and D125A S134A, D210A, Q215A are highlighted in red, blue, and purple, respectively. Statistical analysis:$^{\$}$ p = 0.0001, $^{\#}$ p < 0.0001 compared to mock cells using two-tailed unpaired t-test with Bonferroni's correction after two-way ANOVA. ns indicates no significant difference. Data are presented as mean values (c) and mean values ± SEM (d).
C Residues that were subjected to mutant analysis are mapped to the structure. Only residues with reduced bicarbonate responses are colored.in magenta. **D** Density corresponding to bicarbonate. **E** Putative binding mode of bicarbonate. Black dashed lines indicate hydrogen bonds.

Figure 5. G-protein coupling.

A Hydrogen-bonding interactions between the C-terminal α5-helix residues and the receptor. **B** Electrostatic and hydrogen-binding interactions between ICL3 and the α5-helix. **C–H** Interface between ICL2 and Gq, with residues involved in hydrophobic interactions represented by CPK models. G-protein-bound GPCRs used in the comparison are as follows: H1R-G$_q$ (PDB 7DFL, gray), 5-HT2A-G$_q$ (PDB 6WHA, purple), MRGPRX2-G$_q$ (PDB 7S8L, green), B2R-G$_q$ (PDB 7F6I, yellow-green), and GPR103-Gq (PDB 8ZH8, red). I Comparison of the angles and positions of α5h and αN relative to the receptor. J Superimposition of the Gα subunits.

Figure 6. Structural comparison with related GPCRs

A Structural comparison of GPR30 and AT2R (PDB 5UNF). **B** Interactions around P70 in TM1 of GPR30. **C–F** Conformational changes of TM1 upon agonist binding in CB1 (**C**), β2AR (**D**), A2AR (**E**), and ETB (**F**). The agonist-bound states are colored with the respective colors, while the inactive states are colored gray.

The PDB codes used in this figure are CB1-active (PDB 5XRA), CB1-inactive (PDB 5TGZ), β2AR-active (PDB 3SN6), β2AR-inactive (PDB 2RH1), A2AR-active(PDB 6GDG), A2AR-inactive (PDB 3EML), ETB-active (PDB 8IY5), and ETB-inactive (PDB 5X93)..