Cryo-EM structure of the bicarbonate receptor GPR30

Shota Kaneda[1], Airi Jo-Watanabe[2‡], Hiroaki Akasaka[1], Hidetaka S. Oshima[1], Takehiko Yokomizo[4], Wataru Shihoya[3‡], Osamu Nureki[1‡].

Affiliations:

[1]Department of Biological Sciences, Graduate School of Science, The University of Tokyo, Bunkyo, Tokyo 113-0033, Japan.

[2]Department of Ion Signaling and Response, The Sakaguchi Laboratory, Keio University School of Medicine, Shinanomachi, Tokyo 160-0016, Japan.

[3]Department of Signal Exploration, The Sakaguchi Laboratory, Keio University School of Medicine, Shinanomachi, Tokyo 160-0016, Japan.

[4]Department of Biochemistry, Juntendo University Graduate School of Medicine, Tokyo 113-8421, Japan.

‡To whom correspondence should be addressed. E-mail: awatanabe-tky@umin.ac.jp (A.J.-W.), wtrshh9@gmail.com (W.S.), and nureki@bs.s.u-tokyo.ac.jp (O.N.)

ORCID list
Airi Jo-Watanabe 0000-0003-0899-9915
Hiroaki Akasaka 0000-0003-2118-0912
Hidetaka S. Oshima 0009-0000-0341-707X
Takehiko Yokomizo 0000-0002-5219-1553
Wataru Shihoya 0000-0003-4813-5740
Osamu Nureki 0000-0003-1813-7008

Abstract

G-protein-coupled receptor 30 (GPR30) is a bicarbonate receptor that plays a vital role in cellular responses to extracellular pH and ion homeostasis. Despite its significance, the mechanisms by which GPR30 interacts with bicarbonate ions remain elusive. There is no consensus on a drug that targets GPR30, and difficulties in pharmacological analyses have limited biological and drug discovery research on GPR30. Here, we present the cryo-electron microscopy structure of human GPR30 in the presence of bicarbonate ions, at 3.15 Å resolution. Our structure reveals unique extracellular pockets and critical residues for bicarbonate binding and activation. Functional assays demonstrate that mutations in these residues impair bicarbonate-induced GPR30 activation, underscoring their importance in receptor function. This study also provides insights into G-protein coupling, highlighting the structural divergence between GPR30 and other GPCRs. Our findings not only advance the understanding of the role of GPR30 in pH homeostasis but also pave the way for the development of high-affinity drugs targeting GPR30 for therapeutic interventions in diseases associated with acid-base imbalance.

Introduction

The extracellular environment regulates cellular responses, as exemplified by pH and ion homeostasis. The acid-base balance, largely based on the bicarbonate buffer system *in vivo*, serves as a vital mechanism that maintains the optimal pH for cellular function.[1] Changes in the extracellular pH and ion homeostasis are monitored by membrane channels and receptors, such as G protein-coupled receptors (GPCRs). However, it remains unknown whether and how the acid–base balance-related ions, protons and bicarbonate ions, bind to the receptors and cause the conformational changes that lead to intracellular signal transduction. These proton-sensing GPCRs are thought to be involved in pH homeostasis, particularly in the acidic tumor microenvironment, at inflamed sites, and during ischemia-reperfusion injury[2]. In recent years, the determination of several proton receptor structures has begun to elucidate the molecular mechanisms underlying proton sensing[3–7].

We recently reported that a physiological concentration of bicarbonate ions, the

counterpart of protons in the bicarbonate buffer system, activates G protein-coupled

receptor 30 (GPR30), which leads to G_q-coupled calcium responses[8]. Our study also

demonstrated that GPR30 in brain mural cells regulates blood flow and ischemia–

reperfusion injury. GPR30 was identified as a G protein-coupled estrogen receptor that

mediates the rapid non-genomic action of estradiol (E2)[9]. However, despite numerous

reports on the pleiotropic functions of GPR30 *in vivo*[10,11], controversy remains

regarding the responses of GPR30 to E2 *in vitro*[12], *ex vivo*[13], and *in vivo*[14]. The broad

expression of GPR30[15,16], including blood vessels[17,18], stomach, and lung, has also

raised the possibility of its non-estrogenic functions. We recently demonstrated that

three amino acids, E115$^{2.61}$, Q138$^{3.33}$, and H307$^{7.36}$, are essential for the bicarbonate-

induced activation of GPR30, according to the public homology model

(https://gpcrdb.org/)[19]. However, we did not examine whether bicarbonate ions interact

with GPR30 and cause conformational changes. Moreover, to date, there is no

consensus on a high-affinity drug that targets GPR30, and difficulties in

pharmacological analyses have limited biological and drug discovery research on

GPR30. To elucidate the bicarbonate–GPR30 interaction and to establish the first

structural identification of an acid-base balance-related GPCR, we report the cryo-

electron microscopy (cryo-EM) structure of human GPR30 in the presence of

bicarbonate ions.

Results

Overall structure

For the structural study, we used the full-length human GPR30 sequence. To

efficiently purify the stable GPCR-G-protein complex, the receptor and mini-G_{sqi} were

incorporated in a 'Fusion-G system' by combining two complex stabilization

techniques[20,21] (Figure 1—figure supplement 1A, B). The modified receptor and G-

protein were co-expressed in Sf9 cells and purified by Flag affinity chromatography in

the presence of 200 mM $NaHCO_3$. After an incubation with nanobody 35 (Nb35) and

single chain scFv16, which binds to mini-G_{sqi}, the complex was purified by size

exclusion chromatography (Figure 1—figure supplement 1C, D). The structure of the
purified complex was determined by single-particle cryo-EM analysis, with an overall
resolution of 3.15 Å (Table 1, Figure 1—figure supplement 2, and "Methods"). As the
extracellular portion of the receptor was poorly resolved, we performed receptor
focused refinement, yielding a density map with a nominal resolution of 3.18 Å, which
was combined with the refined map focused on G-protein. The resulting composite map
allowed us to precisely build the atomic model of the components, including the
receptor (residues 51 to 196, 207 to 288, and 299 to 340, G-proteins, and scFv16)
(Figure 1A, B). Nb35 was not visible in the cryo-EM map, probably because of the
effect of the mini-G_s modification, although the interactive residues with Nb35 were not
changed.

The receptor consists of the canonical 7 transmembrane helices (TM) connected
by three intracellular loops (ICL1–3) and three extracellular loops (ECL1–3), and the
amphipathic helix 8 at the C-terminus (H8) (Figure 2A). Most of the TMs are kinked, as
in typical GPCRs[22]. It should be noted that TM1 is also kinked at P71[1.44] (superscripts
indicate Ballesteros-Weinstein numbers[23]), whereas TM1 adopts a straight α-helix in
most class A GPCRs. ICL1 and ECL1 contain short α helices (Figure 2B, C). The short
ICL3 was completely visible in the cryo-EM map (Figure 2C and Figure 1—figure
supplement 2), while residues 197 to 206 in ECL2 and 289 to 298 in ECL3 were
disordered (Figure 2B). ECL2 is attached to TM3 by the disulfide bond between
C130[3.25] and C207[ECL2] (Figure 2B, Figure 2—figure supplement 1A-F), which is highly
conserved in class A GPCRs[22,24]. The cryo-EM structure did not superimpose well on
the AlphaFold-predicted structure (Q99527 in the AlphaFold database)[25], with a root
mean square deviation (R.M.S.D.) of Cα atoms of 3.10 Å (Figure 2—figure supplement
2A). The conserved D[3.49]R[3.50]Y[3.51] motif in the predicted structure represents an inactive
state (Figure 2—figure supplement 1B)[22]. Moreover, ECLs 2 and 3 are rich in cysteine
residues, which form incorrect disulfide bonds in the predicted structure (Figure 2—
figure supplement 2C). These comparisons support the usefulness of experimental
structural determination.

Extracellular pockets

The interaction network between ECL1–3 covers the extracellular side of the receptor. As described above, ECL2 is anchored to TM3 via the disulfide bond, and ECL3 extends between ECL1 and ECL2, with R299^{ECL3} located within the receptor cavity (Figure 3A, B). R299^{ECL3} forms an electrostatic interaction with D210^{ECL2}. These interactions among the ECLs create three extracellular pockets (pockets A–C) (Figure 3A). Pocket B consists of ECL1, TM1, and TM7 (Figure 3C), while pocket C is formed by TM6, TM7, and ECL1 (Figure 3D). These two pockets are small, superficial, and hydrophobic and appear unsuitable for bicarbonate binding. Pocket A consists of ECL1–3 and TM2–7 and is the largest among the three pockets. It is connected to the inside of the receptor, and deep enough to reach W272$^{6.48}$ (Figure 3B). Pocket A contains numerous hydrophilic residues favorable for bicarbonate binding (Figure 3E), suggesting that pocket A is a good candidate for the bicarbonate binding site.

However, no obvious density corresponding to bicarbonate is observed in pocket A, along with pockets B and C. In cryo-EM, negative charges are generally difficult to visualize, and the local resolution of the extracellular region in this structure is moderate (Figure 1—figure supplement 2). Thus, even if bicarbonate ions are present, it is possible that they cannot be visualized.

Our previous studies have shown that the mutations of E115$^{2.61}$, Q138$^{3.33}$, and H307$^{7.36}$ abolish the bicarbonate-dependent calcium response of GPR30, suggesting their contribution to bicarbonate binding. To predict the bicarbonate binding site, we performed an exhaustive mutant analysis of the hydrophilic residues in pocket A, which was not done in the previous study[8]. We performed calcium assays using cell lines stably expressing N-terminally HA-tagged GPR30 and its mutants (Figure 3F, G), whose surface expression levels were analyzed by fluorescence-activated cell sorting (FACS) using an HA-antibody (Figure 3—figure supplement 1-3). Moreover, we performed TGFα shedding assays[26] using HEK cells transiently expressing the receptors (Figure 3—figure supplement 4A-F). The HA-tagged GPR30 wild-type showed a bicarbonate-dependent response. By contrast, the C207^{ECL2}A mutation that disrupts the disulfide bond between ECL2 and TM3, and the P71$^{1.44}$ mutation that disrupts the characteristic kink in TM1, abolished the bicarbonate-dependent response. The

expression levels of these mutants on the cell surface are significantly reduced (Figure

3—figure supplement 1, 4F), indicating that their reduced calcium responses are due to

indirect effects such as structural instability, rather than the direct binding of

bicarbonate ions. These findings underscore the critical role of the structural integrity of

the receptor and provide compelling evidence that the bicarbonate response is mediated

through GPR30.

Among the hydrophilic residue mutants, the $D125^{ECL1}A$ mutant abolished the

bicarbonate-induced responses (Figure 3F, G, and Figure 3—figure supplement 4A, B).

$Q138^{3.33}A$ and $D210^{ECL2}A$ reduced the responses and showed significant reductions in

their cell surface expression levels (Figure 3F, G, and Figure 3—figure supplement 1,

4A, B, F). However, caution is required in interpreting the results, since the cell surface

expression level of the $Q215^{5.39}A$ mutant was reduced to the same extent as $Q138^{3.33}A$

and $D210^{ECL2}A$, but its bicarbonate-induced calcium response was as high as that of WT

(Figure 3—figure supplement 1).

Combining these data with previous studies, we mapped the residues that cause

decreased bicarbonate responses due to mutations on the current structure ($P71^{1.44}$,

$E115^{2.60}$, $D125^{ECL1}$, $Q138^{3.33}$, $C207^{ECL2}$, $D210^{ECL2}$, and $H307^{7.36}$) (Figure 3E). Except

for the structure-contributing residues ($C207^{ECL2}$ and $P71^{1.44}$), the essential residues face

pocket A, further supporting its role as the bicarbonate binding site. The $D210^{ECL2}A$

mutant retained a moderate bicarbonate-induced calcium response (Figure 3F, G) and

would contribute to the structural stability of the pocket through its interaction with

$R299^{ECL3}$ (Figure 3A). The $D125^{ECL1}A$ mutant has lost its activity but is located on the

surface (Figure 3E-G). Thus, $D125^{ECL1}$ is unlikely to be a bicarbonate binding site, and

the mutational effect could be explained due to the decreased surface expression (Figure

3—figure supplement 1, 4F). Given that most bicarbonate binding modes in other

structures require recognition by positively charged residues, bicarbonate may bind to

$H307^{7.36}$, while $E115^{2.61}$, which is located near $H307^{7.36}$, may play an essential role by

coordinating cations for bicarbonate binding, as observed in other structures. These

residues are evolutionarily conserved from fish to human, further supporting their

importance (Figure 3—figure supplement 5).

 ## G-protein coupling

The C-terminal helix of $G\alpha_q$ (α5-helix) deeply enters the intracellular cavity of
GPR30, resulting in the formation of an active signaling complex. On the C-terminal
side of the α5-helix, the backbone carbonyl of Y243$^{G.H5.23}$ (superscript indicates the
common $G\alpha$ numbering [CGN] system[27]) hydrogen bonds with R155$^{3.50}$ in the
conserved D$^{3.49}$R$^{3.50}$Y$^{3.51}$ motif[22] (Figure 4A). In addition, N244$^{G.H5.24}$ forms a hydrogen
bond with the peptide backbone of Y324$^{7.53}$. ICL3 faces the middle part of the α5-helix,
with three characteristic arginine residues (R248$^{6.24}$, R251$^{6.27}$, and R254$^{6.30}$) surrounding
D233$^{G.H5.13}$ (Figure 4B). R248$^{6.24}$ and R254$^{6.30}$ form hydrogen bonds with D233$^{G.H.13}$
and Q237$^{G.H5.17}$, respectively. At ICL2, M163$^{34.51}$ fits into a hydrophobic pocket
composed of the α5-helix, the αN-β2 loop, and the β2-β3 loop of the $G\alpha$ subunit, as in
other GPCR-G_q complexes[28–32] (Figure 4C–H). Above it, L159$^{3.54}$ and A162$^{34.50}$ form
extensive hydrophobic interactions with L235$^{G.H5.15}$, L236$^{G.H5.16}$, and L240$^{G.H5.20}$ in the
α5-helix. These tight interactions with the α5-helix would enable the coupling of
GPR30 with G_q.

The position of the α5-helix aligns with those in other GPCR-G_q complexes, but
the position of the αN-helix is rotated away from the receptor (Figure 4I). The
superimposition of the $G\alpha$ subunits indicates that the rotation is not due to the
movement of the entire G protein, but rather a 33° bend in the αN-helix (Figure 4J).
This is attributed to variations in the interactions of the αN-helix with its receptors. The
αN-helix interacts with the intracellular ends of ICL2 and TM4 in receptor-specific
modes. In most G_q-coupled structures, ICL2 forms an α-helix and R$^{34.55}$ interacts with
R38$^{G.hns1.03}$ at the end of the αN-helix. A typical example is the head-to-toe interaction
between R185$^{34.55}$ and R32$^{G.hns1.03}$ in the serotonin 5-HT$_{2A}$ receptor[31] (Figure 4E).
However, in GPR30, ICL2 does not adopt an α-helix and R$^{34.55}$ is replaced by L167$^{34.55}$,
which displaces R32$^{G.hns1.03}$ downward through their interaction. Moreover, T170$^{34.58}$
and R169$^{34.57}$ form direct hydrogen bonds with R31$^{G.hns1.02}$ and Y243$^{G.H5.23}$, respectively,
which are not observed in other GPCR-G_q complexes. We hypothesize that these
interactions may underpin the distinct αN rotation: A similar interaction between
R38$^{G.hns1.03}$ and L136$^{34.54}$ is also observed in histamine H1 receptor (H$_1$R)[28] (Figure 4D),

in which translational rather than rotational motion of αN is observed (Figure 4J).

Discussion

Our study has elucidated the distinctive structure of GPR30, which is unique
among the class A GPCRs. The GPR30 structure is characterized by the kink of TM1
and the presence of the large extracellular pocket covered by ECLs. This pocket is rich
in hydrophilic residues and could be a focus for the development of high-affinity drugs
targeting GPR30. Our comprehensive mutagenesis study of the binding pocket
complemented our previous study and mapped the essential residues for the
bicarbonate-induced calcium response. Mutations in the S-S bond between ECL2 and
TM3 and the bend in TM1 support their importance in the stability and the shape of the
GPR30 structure and pocket. Furthermore, we were able to identify residues in pocket A
that are important for the bicarbonate response, although there is room for debate
regarding whether they directly contribute to binding or their mutations caused
significant reductions in membrane expression. Considering the binding sites of
negatively charged bicarbonate in other protein structures, it is reasonable to regard the
region around $H307^{7.36}$ as a bicarbonate binding site candidate. To clarify the binding
mode of bicarbonate, a higher resolution structural analysis and complementary
functional assessments are necessary.

GPR30 exhibits low homology to other GPCRs, with even closely related
receptors sharing less than 30% sequence identity. A BLAST analysis revealed that
GPR30 shares the highest sequence identity of 28% with the type 2 angiotensin II
receptor (AT_2R) (Figure 5—figure supplement 1). Consequently, we compared the
current structure of GPR30 with the ligand-bound AT_2R[33]. The superimposition of
GPR30 and AT_2R, with a Cα R.M.S.D. of 1.73 □, suggests significant structural
divergence between the two (Figure 5A). Notably, there is no structural commonality
between GPR30 and AT2R in terms of the orientations of their respective TMs.
Specifically, the extracellular half of TM1 in GPR30 extends outward, and is kinked at
$P71^{1.44}$ in the center of TM1 (Figure 5B). Near $P71^{1.44}$, $Y65^{1.38}$ and $F70^{1.43}$ are oriented

toward the interior of the receptor, forming hydrophobic interactions with L113$^{2.58}$ and

L331$^{7.40}$ to stabilize the kink. In the examination of the loops, ECL2 of AT$_2$R features a

short β-sheet commonly found in peptide-activated class A GPCRs[34–36], whereas that of

GPR30 adopts an elongated conformation that covers the extracellular pocket essential

for the bicarbonate response (Figure 5A, B). These comparisons underscore the unique

structural characteristics of GPR30.

In most GPCRs, TM1 adopts a straight α-helix and is positioned away from the

orthosteric pocket formed by TM2–7. However, TM1 undergoes an inward movement

due to allosteric conformational changes during agonist binding. The cannabinoid

receptor CB$_1$ and adenosine A2A receptor (A$_{2A}$R) exemplify this phenomenon, with

TM1 kinked at G127$^{1.43}$ and G23$^{1.49}$ folding inward upon ligand binding[37–40] (Figure 5C,

D). By contrast, TM1 moves 4 Å outward in the endothelin ET$_B$ receptor[41,42] (Figure

5E). In GPR30, TM1 is kinked at P71$^{1.44}$ (Figure 5B), indicating that this kink exists

regardless of the receptor activation state. Indeed, the P71$^{1.44}$A mutation, which

eliminates the kink in TM1, completely abolished the bicarbonate response (Figure 3F,

G). Moreover, P71$^{1.44}$ is evolutionarily conserved (Figure 3—figure supplement 5).

These findings suggest the critical role of the TM1 kink in both the putative

conformational changes associated with ligand binding and the functional integrity of

GPR30.

During the revision of this manuscript, the structures of apo-GPR30-G$_q$ (PDB

8XOG) and the exogenous ligand Lys05-bound GPR30-G$_q$ (PDB 8XOF) were

reported[43]. We compared our structure of GPR30 in the presence of bicarbonate with

these structures. In the extracellular region, the position of TM5 in GPR30 in the

presence of bicarbonate is similar to that in apo-GPR30. In contrast, the position of

TM6 is shifted outward relative to that of apo-GPR30, resembling the conformation

observed in Lys05-bound GPR30 (Figure 6A, B). Additionally, the position of ECL1 is

also shifted outward compared to that of apo-GPR30 (Figure 6B). In the GPR30

structure in the presence of bicarbonate, ECL2 was modeled, suggesting differences in

structural flexibility. These findings indicate that the structure of GPR30 in the presence

of bicarbonate is different from both the apo structure and the Lys05-bound structure,

demonstrating that the structure and the flexibility of the extracellular domain of GPR30
change depending on the type of ligand. Furthermore, focusing on the interaction with
G_q, the αN helix of G_q is not rotated in the structure bound to Lys05, in contrast to the
characteristic bending of the αN helix in our structure (Figure 6C, D). Although it is
necessary to consider variations in experimental conditions, such as salt concentration,
the differences in the G_q binding modes suggest that the downstream signals may
change in a ligand-dependent manner.

In our work, we used relatively high (mM) concentrations of bicarbonate to
activate GPR30. In parallel with its low potency, the physiological concentrations of
bicarbonate are 22-26 mM in the extracellular fluid, including interstitial fluid and
blood, and 10-12 mM in the cytoplasm. Therefore, GPR30 is activated by the
physiological concentrations of bicarbonate in tissues. The bicarbonate concentrations
are altered in various physiological and pathological conditions – metabolic acidosis in
chronic kidney disease causes a drop to 2-3 mM, and metabolic alkalosis induced by
severe emesis increases bicarbonate concentrations to over 30 mM. Thus, our work
clearly shows that GPR30 is activated by physiological concentrations of bicarbonate,
present both intracellularly and extracellularly, and that GPR30 can be deactivated or
reactivated under various pathophysiological conditions. Thus, we have identified
GPR30 as a bicarbonate receptor.

Table. 1   Cryo-EM data collection,
refinement and validation statistics

Data collection	GPR30-G_q complex
Microscope	Titan Krios (Thermo Fisher Scientific)
Voltage (keV)	300
Electron exposure (e^-/Å^2)	50.660
Detector	Gatan K3 Summit camera (Gatan)
Magnification	×105,000
Defocus range (μm)	-0.8–1.6
Pixel size (Å/pix)	1.16

Number of movies	9,824
Symmetry	C1
Picked particles	10,148,422
Final particles	522,404
Map resolution (Å)	3.15
FSC threshold	0.143
Model refinement	
Atoms	
R.m.s. deviations from ideal	
Bond lengths (Å)	0.003
Bond angles (°)	0.528
Validation	
Clashscore	9.56
Rotamers Outliers(%)	0.23
Ramachandran plot	
Favored (%)	96.32
Allowed (%)	3.68
Outlier (%)	0

Figures

Figure 1. Overall structure of the GPR30-miniG$_{sqi}$β$_1$γ$_2$-scFv16 complex.

A Unsharpened cryo-EM density map of the GPR30-miniG$_{sqi}$β$_1$γ$_2$-scFv16-Nb35 complex, with the components individually colored. **B** The refined structure of the complex is shown as a ribbon representation.

Figure 2. Receptor structure.

A–C Overall structure of the receptor, viewed from the membrane plane (**A**), intracellular side (**B**), and extracellular side (**C**).

Figure 3. Architecture of the extracellular pocket.

A Molecular surface of the extracellular side. **B** Cross section of the pocket. **C–E** Residues facing pocket B (**C**), pocket C (**D**), and pocket A (**E**). In panel (**E**), only residues with reduced bicarbonate responses are highlighted. **F, G** Calcium assay using stable HEK293 cell lines expressing the N-terminal HA-tagged GPR30 wild-type (WT) and mutants. The mutants D111$^{2.56}$A, N276$^{6.52}$A, and Q296^{ECL3}A; E121^{ECL1}A, R122^{ECL1}A, S134$^{3.29}$A, Q215$^{5.39}$A, and E218$^{5.42}$A; D210^{ECL2}A and Q138$^{3.33}$A; and D125^{ECL1}A, C207^{ECL2}A, and P71$^{1.44}$A are highlighted in red, blue, green, and purple, respectively. The cells were stimulated by the indicated concentrations of NaHCO$_3$ at the timepoint of t = 20 sec. The Y axis indicates the difference between the maximum and minimum fluorescent values during 15 to 60 sec. Statistical analysis: each sample was compared with WT cells using Dunnett's T3 multiple comparisons test after Brown-Forsythe and Welch ANOVA tests. **** $p < 0.0001$ compared to WT cells, no significant difference is not shown. Data are presented as mean values ± SEM (**F**) and each replicate with the mean ± SEM (**G**).

Figure 4. G-protein coupling.

A Hydrogen-bonding interactions between the C-terminal α5-helix residues and the receptor. **B** Electrostatic and hydrogen-binding interactions between ICL3 and the α5-helix. **C**–**H** Interface between ICL2 and G_q, with residues involved in hydrophobic interactions represented by CPK models. G-protein-bound GPCRs used in the comparison are as follows: H_1R-G_q (PDB 7DFL, gray), B_2R-G_q (PDB 7F6I, yellow-green), MRGPRX$_2$-G_q (PDB 7S8L, green), 5-HT$_{2A}$-G_q (PDB 6WHA, gray), and GPR103-G_q (PDB 8ZH8, red). **I** Comparison of the angles and positions of α5h and αN relative to the receptor. **J** Superimposition of the Gα subunits.

Figure 5. Structural comparison with related GPCRs.

A Structural comparison of GPR30 and AT$_2$R (PDB 5UNF). **B** Interactions around P71$^{1.44}$ in TM1 of GPR30. **C**–**F** Conformational changes of TM1 upon agonist binding in CB1 (**C**), A$_{2A}$R (**D**), and ET$_B$ (**E**). The agonist-bound states are colored with the respective colors, while the inactive states are colored gray. The PDB codes used in this figure are CB1-active (PDB 5XRA), CB$_1$-inactive (PDB 5TGZ), A$_{2A}$R-active (PDB 6GDG), A$_{2A}$R-inactive (PDB 3EML), ET$_B$-active (PDB 8IY5), and ET$_B$-inactive (PDB 5X93).

Figure 6. Structural comparison with GPR30 bound to Lys05.

A, B Superimposition of bicarbonate-GPR30, Lys05-binding GPR30 (PDB 8XOF) (**A**), and apo-GPR30 (PDB 8XOG) (**B**) structures, viewed from the extracellular side. **C** Comparison of the angles and positions of α5h and αN relative to the receptor. **D** Superimposition of the Gα subunits.

Figure 1.—figure supplement 1. Sample purification.
A Schematic representation of the fusion-G system. **B** Fluorescence-detection size-exclusion chromatography (FSEC) analysis of complex formation by GPR30. The trace from solubilized cells expressing only GPR30 is orange, and that from cells co-

expressing GPR30 and G-protein is blue. **C** Size-exclusion chromatography of the

GPR30-G-protein complex on a Superose 6 Increase column. The fraction shaded in

blue was collected. **D** SDS-PAGE gel of samples after size-exclusion chromatography,

stained with Coomassie Brilliant Blue.

Figure 1.—figure supplement 2. Cryo-EM analysis.

Flow chart of the cryo-EM data processing for the GPR30-G_q complex, including

particle projection selection, classification, and 3D density map reconstruction. The 3D

density map was refined with a mask on the receptor. Details are provided in the

Methods section.

Figure 2.—figure supplement 1. ECL2 density.

A-C top view of density map around disulfide bond between ECL2 and TM3 at counter

levels 8.5 (**A**), 7.25 (**B**), and 6 (**C**). **D-F** side view of density map around disulfide bond

between ECL2 and TM3 at counter levels 8.5 (**D**), 7.25 (**E**), and 6 (**F**).

Figure 2.—figure supplement 2. Structural comparison with AlphaFold-predicted

structure.

A–C Superimposition of the cryo-EM (orange) and AF3 (khaki) structures. (**A**) Overall

view of the receptor, (**B**) focused on the intracellular side, and (**C**) focused on the

extracellular side.

Figure 3.—figure supplement 1. Cell surface expression of stable wild-type and

GPR30 mutants.

Flow cytometric-based assay to analyze WT and mutant GPR30 cell surface expression.

Cell surface expression, defined as the HA-positive subset (red) compared with isotype

control staining (blue), is indicated on each histogram.

Figure 3.—Figure Supplement 2. Flow cytometric gating criteria to analyze GPR30

cell surface expression.

A, B Gating used in the analysis. **C, D** Cell surface expression is defined as the HA-

positive subset (red) compared with isotype control staining (blue). **(C)** Mock cells, **(D)**

WT GPR30-expressing cells.

Figure 3.—Figure Supplement 3. Flow cytometric-based assay to analyze the

whole-cell expression of WT GPR30 and its mutants.

A Gating used in the analysis. **B** Whole-cell expression is defined as the HA-positive

subset (red) compared with isotype control staining (blue).

Figure 3.—Figure Supplement 4. Cell surface expression and bicarbonate-induced

activation of HEK293 cells transiently expressing GPR30 mutants.

A, B TGFα shedding assay using HEK293 cells transfected with HA-tagged hGPR30.

The mutants $D111^{2.56}A$; $D125^{ECL1}A$, $C207^{ECL2}A$, $P71^{1.44}A$, and $H307^{7.36}A$; and

$S134^{3.29}A$, $D210^{ECL2}A$, and $Q215^{5.39}A$ are highlighted in red, blue, and purple,

respectively. **C** Whole-cell expression of HA-tagged mutants, analyzed by western

blotting with 20 μg of whole-cell lysate per lane. **D** Cell surface expression of HA-

tagged mutants, using cell surface biotinylation and avidin immunoprecipitation,

analyzed by western blotting with 1.5 μg of cell surface protein per lane. **E, F**

Quantitative analysis of HA expression in **C** and **D**.

Statistical analysis: $^{\$}p = 0.01$, $^{\S}p < 0.001$ compared to HA-hGPR30 cells stimulated

with vehicle, $^{\#}p < 0.0001$ compared to HA-hGPR30 cells stimulated with 11 mM

$NaHCO_3$, using two-tailed unpaired t-test with Bonferroni's correction after two-way

ANOVA. Data are presented as mean values (**A**) and mean values ± SEM (**B**).

Figure 3.—figure supplement 5. Conservation of the GPR30 homologs.

Amino acid alignment of representative GPR30 homologs.

Figure 5.—figure supplement 1. Sequence alignment of GPR30 and AT$_2$R.

Data Availability

The cryo-EM density map and atomic coordinates for the GPR30-G$_q$ complex
have been deposited in the Electron Microscopy Data Bank and the PDB, under the
accession codes: EMD-66632 [https://www.ebi.ac.uk/emdb/entry/EMD-66632]
(GPR30-G$_q$), EMD-66629 [https://www.ebi.ac.uk/emdb/entry/EMD-66629] (GPR30-G$_q$,
consensus map), EMD-66630 [https://www.ebi.ac.uk/emdb/entry/EMD-66630]
(GPR30-G$_q$, G-protein focused), EMD-66631 [https://www.ebi.ac.uk/emdb/entry/EMD-
66631] (GPR30-G$_q$, receptor focused), and PDB 9X74
[http://doi.org/10.2210/pdb9X74/pdb]. Source data are provided with this paper. All
other data are available from the corresponding authors upon reasonable request.

Acknowledgements

We thank K. Ogomori and C. Harada for technical assistance. This work was
supported by grants from the JSPS KAKENHI, grant numbers 21H05037 (O.N.),
25K02397 (W.S.), 24KJ0906 (H.A.), 23K06393, 24KK0146, and 25H01336 (T.Y); the
Japan Agency for Medical Research and Development (AMED), grant numbers
JP233fa627001 (O.N.), 25ak0101252h0001 (W.S.), and JP20gm6210026 (A.J.-W.); the
JST FOREST Program (Grant number JPMJFR220S to A.J.-W.); the Takeda Science
Foundation (W.S. and A.J.-W.); the Mochida Memorial Foundation for Medical and
Pharmaceutical Research (W.S.); Dojindo Laboratories' Foundation for Life Science
(W.S.); the Nakatani Foundation (W.S.); the Mitsubishi Foundation (W.S.); the Daiichi
Sankyo Foundation of Life Science (W.S.); the Takahashi Industrial and Economic
Research Foundation (W.S.); G-7 Scholarship Foundation (W.S.); and the Platform
Project for Supporting Drug Discovery and Life Science Research (Basis for Supporting
Innovative Drug Discovery and Life Science Research (BINDS)) from AMED, grant

numbers JP23ma121002 and JP23ama121012.

Author contributions

430   S.K. performed all the experiments involved in the structural determination,
431   assisted by H.A. and H.S.O. W.S. designed the project and initially tried the expression
432   of GPR30. A.J.-W. and T.Y. performed and oversaw the mutagenesis study. The
433   manuscript was mainly prepared by S.K., H.A., and W.S., with assistance from O.N.

Competing interests

436   The authors declare no competing interests.

Methods

Expression and purification of scFv16 and Nb35

The gene encoding scFv16 was synthesized (GeneArt) and subcloned into a
modified pFastBac vector[44], with the resulting construct encoding the GP67 secretion
signal sequence at the N terminus, and a His_8 tag followed by a TEV cleavage site at the
C terminus. The His_8-tagged scFv16 was expressed and secreted by Sf9 insect cells, as
previously reported[45]. The Sf9 cells were removed by centrifugation at 5,000g for
10 min, and the secreta-containing supernatant was combined with 5 mM $CaCl_2$, 1
mM $NiCl_2$, 20 mM HEPES (pH 8.0), and 150 mM NaCl. The supernatant was
mixed with Ni Superflow resin (GE Healthcare Life Sciences) and stirred for 1 h at
4 ℃. The collected resin was washed with buffer containing 20 mM HEPES
(pH 8.0), 500 mM NaCl and 20 mM imidazole, and further washed with
10 column volumes of buffer containing 20 mM HEPES (pH 8.0), 500 mM NaCl
and 20 mM imidazole. Next, the protein was eluted with 20 mM Tris (pH 8.0),
500 mM NaCl and 400 mM imidazole. The eluted fraction was concentrated and
loaded onto a Superdex200 10/300 Increase size-exclusion column, equilibrated in

buffer containing 20 mM Tris (pH 8.0) and 150 mM NaCl. Peak fractions were
pooled, concentrated to 5 mg/ml using a centrifugal filter device (Millipore
10 kDa MW cutoff), and frozen in liquid nitrogen.

Nb35 was prepared as previously reported[46,47]. In brief, Nb35 was expressed in
the periplasm of *E. coli*. The harvested cells were disrupted by sonication. Nb35 was
purified by nickel affinity chromatography, followed by gel-filtration chromatography,
and frozen in liquid nitrogen.

461 Constructs for expression of GPR30 and G_q

The human GPR30 gene (UniProtKB, Q99527) was subcloned into a modified
pFastBac vector[20], with an N-terminal haemagglutinin signal peptide followed by the
Flag-tag epitope (DYKDDDDK) and the LgBiT fused to its C-terminus followed by a 3
C protease site and EGFP-His$_8$ tag. A 15 amino sequence of GGSGGGGSGGGSSSGG
was inserted into both the N-terminal and C-terminal sides of LgBiT. Rat $G\beta_1$ and
bovine $G\gamma_2$ were subcloned into the pFastBac Dual vector. In detail, rat $G\beta_1$ was cloned
with a C-terminal HiBiT connected with a 15 amino sequence of
GGSGGGGSGGGSSSGG. Moreover, mini-G_{sqi} was subcloned into the C-terminus of the
bovine $G\gamma_2$ with a nine amino sequence GSAGSAGSA linker. The resulting pFastBac
dual vector can express the mini-G_{sqi} trimer.

473 Expression and purification of the human GPR30 – G_q complex

The recombinant baculovirus was prepared using the Bac-to-Bac baculovirus
expression system (Thermo Fisher Scientific). For expression, 2 L of Sf9 cells at a
density of 3×10^6 cells/mL were co-infected with baculovirus encoding GPR30 and
mini-G_{sqi} trimer at the ratio of 1:1 and the cells were incubated at 30 ☐. After 48 hours,
the collected cells were resuspended and dounce-homogenized in 20 mM Tris-HCl,
pH 8.0 (4 ☐), 200 mM NaCl, 10% Glycerol, 200 mM NaHCO$_3$, 25 mU/ml apyrase.
The crude membrane fraction was collected by ultracentrifugation at 180,000g for 1 h
and solubilized in buffer, containing 20 mM Tris-HCl, pH 8.0, 150 mM NaCl, 1%
n-dodecyl-beta-D-maltopyranoside (DDM) (Calbiochem), 0.2 % cholesteryl

hemisuccinate (CHS) (Merck), 10% glycerol, 200 mM $NaHCO_3$, and 25 mU/ml

Apyrase, for 1 h at 4□□. The supernatant was separated from the insoluble material by

ultracentrifugation at 180,000g for 20□min and incubated with 5 mL of M2 anti-flag

affinity resin (Sigma) for 1 h at 4 □. The resin was washed with 20 column volumes of

buffer containing 20□mM Tris-HCl, pH□8.0 500□mM NaCl, 10% Glycerol, 0.05%

Lauryl Maltose Neopentyl Glycol (LMNG) (Antrace), 0.005% CHS and 200 mM

$NaHCO_3$. The complex was eluted in buffer containing 20□mM Tris-HCl, pH□8.0,

150□mM NaCl, 10% Glycerol, 0.01% LMNG, 0.001% CHS, 200 mM $NaHCO_3$ and 0.2

mg/mL Flag peptide. The eluate was incubated with the Nb35 and scFv16 at 4 □. The

complex was concentrated and purified by size exclusion chromatography on a

Superose 6 increase (GE) column in 20□mM Tris-HCl, pH8.0, 150□mM NaCl, 0.01%

LMNG, 0.001% CHS and 200 mM $NaHCO_3$. Peak fractions were concentrated to 4.72

mg/ml for electron microscopy studies.

Sample vitrification and cryo-EM data acquisition

The purified complex was applied onto a freshly glow-discharged Quantifoil Au

grid (R1.2/1.3, 300 mesh), and plunge-frozen in liquid ethane by using a Vitrobot Mark

IV. Data collections were performed on a 300kV Titan Krios G3i microscope (Thermo

Fisher Scientific) equipped with a BioQuantum K3 imaging filter and a K3 direct

electron detector (Gatan).

A total of 9824 movies were acquired with a calibrated pixel size of

0.83□Å□pix^{-1} and with a defocus range of -0.8 to -1.6□μm, using EPU. Each movie

was acquired for 2.6□s and split into 48 frames, resulting in an accumulated exposure

of about 50.660□electrons per $Å^2$.

Image processing

All acquired dose-fractionated movies were imported into CryoSPARC v4.4[48]

and subjected to beam-induced motion correction. The contrast transfer function (CTF)

parameters were estimated and a total of 10,148,422 particles were extracted. The

particles were subjected to 2D classifications, Ab-initio reconstruction and several

rounds of hetero refinement and Non-uniform refinement. Next, the particles were

subjected to 3D classification with a mask on the receptor. Then the particle sets were

subjected to Reference Based Motion Correction. Motion-corrected 522,404 particles

were subjected to Non-uniform refinement, yielding a map with a global nominal

resolution of 3.15 Å, with the gold standard Fourier Shell Correlation (FSC=0.143)

criteria[49]. Moreover, the 3D model was refined with a mask on the receptor. As a result,

the receptor has a higher resolution with a nominal resolution of 3.18 Å. The overall and

receptor focused maps were combined by phenix[50]. The processing strategy is described

in Supplementary Figure 2.

Model building and refinement

The density map was sharpened by phenix.auto_sharpen[51] and the quality of the

density map was sufficient to build a model manually in COOT[52,53]. The model building

was facilitated by the Alphafold-predicted structure. We manually fitted GPR30, the

G_q heterotrimer and scFv16 into the map. We then manually readjusted the model into

using COOT and refined it using phenix.real_space_refine[50,54] (v.1.19) with the

secondary-structure restraints using phenix secondary_structure_restraints.

Vector construction and transfection

Human *Gpr30* cDNA was obtained from human hepatoblastoma-derived HepG2

cells. The coding sequences of human *Gpr30* were inserted into the multi-cloning site of

the plasmid vector pCXN2, which was generated in our laboratory via modification of

pCAGGS, between the KpnI and EcoRI sites. The N- and C-terminal HA-tagged *Gpr30*

was amplified using a reverse primer containing the HA sequenceOne amino acid

mutation of human GPR30 was generated as follows: the targeted amino acid was

changed to alanine (GCC) using a two-step PCR method with the QuikChange® Primer

Design Program by Agilent (https://www.agilent.com/store/primerDesignProgram.jsp),

and the coding sequence with each mutation was inserted into the multi-cloning site of

pCXN2 between the KpnI and EcoRI sites. The mutations generated were D111A,

E121A, R122A, D125A, S134A, Q138A, D210A, Q215A, E218A, N276A, Q296A,

C207A, P71A, and H307A.

Cell line sources and transfection

HEK293A cells (female origin; Thermo Fisher Scientific) were maintained at

37 □ and 5% CO_2. These vectors were transfected using the lipofection method

(Lipofectamine™ 2000 Transfection Reagent, 11668019, Invitrogen). Stable cell lines

were established through antibiotic drug selection. The HEK293A cells were transfected

with the N-terminal HA-tagged wild-type and mutant *Gpr30* and maintained in the

culture medium containing 1 mg/ml of G418. After 2-3 weeks of antibiotic drug

selection, stable expression of exogenous GPR30 was confirmed by three independent

flow cytometric analyses.

TGFα shedding assay

The TGFα shedding assay was performed according to a previously published

protocol[26]. The AP-TGFα expression vector was provided by Dr. Inoue and Dr. Aoki,

Tohoku University. HEK293 cells were seeded in 12-well plates at a density of 1×10^5

cells/well and cultured for 24 h. At 70% confluency, a mixture of plasmid vectors

containing GPCR (see '***Vector construction and transfection***' section) and AP-TGFα

was transfected into the cells using Lipofectamine 2000 transfection reagent. After

another 24 h of incubation, the cells were detached with 0.05% trypsin/EDTA (32777-

44, Nacalai Tesque), suspended in Hanks' balanced salt solution, and seeded in 96-well

plates. The cells were stimulated for 1 h with $1–44\times10^{-3}$ M (final concentration) of

$NaHCO_3$ at 37 □, under 0.03% CO_2. Conditioned media (CM) was transferred to

another plate, and 80 µl of alkaline phosphatase (AP) solution (40 mM Tris-HCl, pH 9.5,

40 mM NaCl, 10 mM $MgCl_2$, 10 mM p-nitrophenylphosphate disodium salt

hexahydrate) was added to both plates, which were then incubated at 37 □. The optical

density at 405 nm (OD405) was measured using a microplate reader (iMark, Bio-Rad)

at 5 min, 30 min, 1 h, and 2 h, depending on the reaction rate. The percentage of shed

AP-TGF was calculated using the following equations:

`AP activity = ΔOD405 (1 − 0 h)`

% CM (conditioned media) = AP_{CM} / (AP_{CM} + AP_{Cell})

Calcium assay

Stable HEK293 cells (2×10^4 cells/well) were seeded into a black wall and clear

bottom 96-well plate 24 h before the assay. Cells at 90–100% confluency were

incubated with HEPES buffer ($1 \times$ HBSS, 2.5 mM probenecid, 25 mM HEPES, pH 7.4)

containing 10 µM Fluo-8 AM (21080, AAT-Bio) for 60 min at 37□ and 5% CO_2. The

cells were washed twice with HEPES buffer. The cells were stimulated with indicated

concentrations of $NaHCO_3$ and ATP at 37 □ and 0.03% CO_2. The fluorescence intensity

was analysed using FlexStation 3 (Molecular Devices, Ex/Em = 490/525 nm).

Flow cytometry

HEK293A cells stably expressing the N-terminal HA-tagged wild-type and

mutant GPR30 were incubated with the calcium-free Minimum Essential Medium (S-

MEM, #11380037, Gibco™) for 30 min at 37 □ and 5.0% CO_2 before the assay. The

cells were blocked with PBS containing 2% goat serum for 10 min at 4□, incubated

with the biotin-conjugated anti-HA antibody (2.5 µg/ml, #12158167001, Roche) for 60

min at 4□, and then stained with Alexa Fluor 488-conjugated anti-rat IgG (5 µg/ml, #A-

11006, Thermo Fisher Scientific) for 40 min at 4□. LSRFortessa (BD Biosciences) was

used for analysis. For intracellular staining, the cells were fixed and stained using

eBioscience™ Fixable Viability Dye eFluor 780 (#65-0865-14, Thermo Fisher

Scientific) and eBioscience™ Fox3/Transcription Factor Staining Buffer Set (#00-5523-

00, Thermo Fisher Scientific).

Analysis of cell surface expression of GPR30

597   The cell surface expression of wild-type and mutant hGPR30-HA was analysed

598   via cell surface protein isolation using a Cell Surface Protein Isolation Kit (#89881,

599   Thermo Scientific™) followed by western blotting. Briefly, HEK293 cells transiently

600   expressing wild-type or mutant hGPR30-HA were biotinylated for 30 min at 4 □. Then,

601   the cells were harvested, and biotinylated proteins were isolated with avidin binding.

Western blotting

Cell lysates were separated using sodium dodecyl sulfate-polyacrylamide gel

electrophoresis under reducing conditions and transferred to polyvinylidene difluoride

membranes (Immobilon P, IPVH00010, Millipore). Primary antibodies used were anti-

HA High Affinity, 1:1000 dilution, 11867423001, Roche; Na-K-ATPase, 1:1000

dilution, #3010, Cell Signaling Technology; β-Actin (AC-15), 1:1000 dilution, sc-69879,

Santa Cruz Biotechnology. Secondary antibodies used were anti-rabbit IgG, HRP-linked

Antibody, 1:5000 dilution, #7074, Cell Signaling Technology; anti-mouse IgG, HRP-

linked Antibody, 1:5000 dilution, #7076, Cell Signaling Technology; anti-rat IgG, HRP-

Linked Whole Ab Goat, 1:5000 dilution, NA935, Cytiva. The membranes were probed

at 4 □ overnight with the primary antibodies. The membranes were subsequently

incubated with the corresponding secondary antibodies. The signals were detected with

ECL Prime (RPN2236, Cytiva) or ImmunoStar LD (296-69901, FUJIFILM Wako Pure

Chemical Corporation) using a chemiluminescence imaging system (CFusion FX7,

Vilber).

References

1.   Brinkman, J. E. & Sharma, S. Physiology, metabolic alkalosis. in *StatPearls*

(StatPearls Publishing, Treasure Island (FL), 2025).

2.   Imenez Silva, P. H., Camara, N. O. & Wagner, C. A. Role of proton-activated G

protein-coupled receptors in pathophysiology. *Am. J. Physiol. Cell Physiol.* **323**,

C400–C414 (2022).

3.   Guo, L. *et al.* Structural basis and biased signaling of proton sensation by GPCRs

mediated by extracellular histidine rearrangement. *Mol. Cell* **85**, 1658-1673.e7

(2025).

4.   Yue, X. *et al.* Structural basis of stepwise proton sensing-mediated GPCR

activation. *Cell Res.* **35**, 423–436 (2025).

5.   Howard, M. K. *et al.* Molecular basis of proton sensing by G protein-coupled

receptors. *Cell* **188**, 671-687.e20 (2025).

6.   Ma, Y. *et al.* Cryo-EM structure of an activated GPR4-Gs signaling complex. *Nat.*

Commun. **16**, 605 (2025).

7.   Wen, X. *et al.* Evolutionary study and structural basis of proton sensing by Mus

GPR4 and Xenopus GPR4. *Cell* **0**, (2025).

8.   Jo-Watanabe, A. *et al.* Bicarbonate signalling via G protein-coupled receptor

regulates ischaemia-reperfusion injury. *Nat. Commun.* **15**, 1530 (2024).

9.   Revankar, C. M., Cimino, D. F., Sklar, L. A., Arterburn, J. B. & Prossnitz, E. R. A

transmembrane intracellular estrogen receptor mediates rapid cell signaling.

Science **307**, 1625–1630 (2005).

10.   Ford, J. *et al.* GPR30 deficiency causes increased bone mass, mineralization, and

growth plate proliferative activity in male mice. *J. Bone Miner. Res.* **26**, 298–307

(2011).

11.   Sharma, G. *et al.* GPER deficiency in male mice results in insulin resistance,

dyslipidemia, and a proinflammatory state. *Endocrinology* **154**, 4136–4145 (2013).

12.   Tutzauer, J. *et al.* Ligand-independent G protein-coupled estrogen receptor/G

protein-coupled receptor 30 activity: Lack of receptor-dependent effects of G-1 and

17β-estradiol. *Mol. Pharmacol.* **100**, 271–282 (2021).

13.   Lamprecht, M. R. & Morrison, B., 3rd. GPR30 activation is neither necessary nor

sufficient for acute neuroprotection by 17β-estradiol after an ischemic injury in

organotypic hippocampal slice cultures. *Brain Res.* **1563**, 131–137 (2014).

14. Otto, C. *et al.* GPR30 does not mediate estrogenic responses in reproductive organs

in mice. *Biol. Reprod.* **80**, 34–41 (2009).

15. Uhlén, M. *et al.* Proteomics. Tissue-based map of the human proteome. *Science*

347, 1260419 (2015).

16. The Human Protein Atlas. https://www.proteinatlas.org/.

17. Vanlandewijck, M. *et al.* A molecular atlas of cell types and zonation in the brain

vasculature. *Nature* **554**, 475–480 (2018).

18. Winkler, E. A. *et al.* A single-cell atlas of the normal and malformed human brain

vasculature. *Science* **375**, eabi7377 (2022).

19. Kooistra, A. J. *et al.* GPCRdb in 2021: integrating GPCR sequence, structure and

function. *Nucleic Acids Res.* **49**, D335–D343 (2021).

20. Sano, F. K., Akasaka, H., Shihoya, W. & Nureki, O. Cryo-EM structure of the

endothelin-1-ETB-Gi complex. *Elife* **12**, (2023).

21. Shihoya, W., Iwama, A., Sano, F. K. & Nureki, O. Cryo-EM advances in GPCR

structure determination. *J. Biochem.* (2024) doi:10.1093/jb/mvae029.

22. Venkatakrishnan, A. J. *et al.* Molecular signatures of G-protein-coupled receptors.

Nature **494**, 185–194 (2013).

23. Ballesteros, J. A. & Weinstein, H. [19] Integrated methods for the construction of

three-dimensional models and computational probing of structure-function

relations in G protein-coupled receptors. in *Methods in Neurosciences* vol. 25 366–

428 (Elsevier, 1995).

24. Nicoli, A., Dunkel, A., Giorgino, T., de Graaf, C. & Di Pizio, A. Classification

model for the second extracellular loop of class A GPCRs. *J. Chem. Inf. Model.* **62**,

511–522 (2022).

25. Database, A. P. S. AlphaFold Protein Structure Database.

https://alphafold.ebi.ac.uk/.

26. Inoue, A. *et al.* TGFα shedding assay: an accurate and versatile method for

detecting GPCR activation. *Nat. Methods* **9**, 1021–1029 (2012).

27. Flock, T. *et al.* Universal allosteric mechanism for Gα activation by GPCRs.

Nature **524**, 173–179 (2015).

28. Xia, R. *et al.* Cryo-EM structure of the human histamine H1 receptor/Gq complex.

Nat. Commun. **12**, 2086 (2021).

29. Shen, J. *et al.* Cryo-EM structures of human bradykinin receptor-Gq proteins

complexes. *Nat. Commun.* **13**, 714 (2022).

30. Cao, C. *et al.* Structure, function and pharmacology of human itch GPCRs. *Nature*

600, 170–175 (2021).

31. Kim, K. *et al.* Structure of a hallucinogen-activated gq-coupled 5-HT2A serotonin

receptor. *Cell* **182**, 1574-1588.e19 (2020).

32. Iwama, A. *et al.* Structure and dynamics of the pyroglutamylated RF-amide peptide

QRFP receptor GPR103. *Nat. Commun.* **15**, 4769 (2024).

33. Zhang, H. *et al.* Structural basis for selectivity and diversity in angiotensin II

receptors. *Nature* **544**, 327–332 (2017).

34. Hong, C. *et al.* Structures of active-state orexin receptor 2 rationalize peptide and

small-molecule agonist recognition and receptor activation. *Nat. Commun.* **12**, 815

(2021).

35. Mobbs, J. I. *et al.* Structures of the human cholecystokinin 1 (CCK1) receptor

bound to Gs and Gq mimetic proteins provide insight into mechanisms of G

protein selectivity. *PLoS Biol.* **19**, e3001295 (2021).

36. Tang, T. *et al.* Receptor-specific recognition of NPY peptides revealed by

structures of NPY receptors. *Sci. Adv.* **8**, eabm1232 (2022).

37. Hua, T. *et al.* Crystal structure of the human cannabinoid receptor CB1. *Cell* **167**,

750-762.e14 (2016).

38. Hua, T. *et al.* Crystal structures of agonist-bound human cannabinoid receptor CB1.

Nature **547**, 468–471 (2017).

39. García-Nafría, J., Lee, Y., Bai, X., Carpenter, B. & Tate, C. G. Cryo-EM structure

of the adenosine A2A receptor coupled to an engineered heterotrimeric G protein.

Elife **7**, (2018).

40. Jaakola, V.-P. *et al.* The 2.6 angstrom crystal structure of a human A2A adenosine

receptor bound to an antagonist. *Science* **322**, 1211–1217 (2008).

41. Shihoya, W. *et al.* Activation mechanism of endothelin ETB receptor by

endothelin-1. *Nature* **537**, 363–368 (2016).

42. Shihoya, W., Sano, F. K. & Nureki, O. Structural insights into endothelin receptor

signalling. *J. Biochem.* **174**, 317–325 (2023).

43. Liu, H. *et al.* Structural and functional evidence that GPR30 is not a direct estrogen

receptor. *Cell Res.* **34**, 530–533 (2024).

44. Okamoto, H. H. *et al.* Cryo-EM structure of the human MT1-Gi signaling complex.

Nat. Struct. Mol. Biol. **28**, 694–701 (2021).

45. Akasaka, H., Sano, F. K., Shihoya, W. & Nureki, O. Structural mechanisms of

potent lysophosphatidic acid receptor 1 activation by nonlipid basic agonists.

Commun. Biol. **7**, 1444 (2024).

46. Nagiri, C. *et al.* Cryo-EM structure of the β3-adrenergic receptor reveals the

molecular basis of subtype selectivity. *Mol. Cell* **81**, 3205-3215.e5 (2021).

47. Nureki, I. *et al.* Cryo-EM structures of the β3 adrenergic receptor bound to

solabegron and isoproterenol. *Biochem. Biophys. Res. Commun.* **611**, 158–164

(2022).

48. Punjani, A., Rubinstein, J. L., Fleet, D. J. & Brubaker, M. A. cryoSPARC:

algorithms for rapid unsupervised cryo-EM structure determination. *Nat. Methods*

14, 290–296 (2017).

49. Rosenthal, P. B. & Henderson, R. Optimal determination of particle orientation,

absolute hand, and contrast loss in single-particle electron cryomicroscopy. *J. Mol.*

Biol. **333**, 721–745 (2003).

50. Adams, P. D. *et al.* PHENIX: a comprehensive Python-based system for

macromolecular structure solution. *Acta Crystallogr. D Biol. Crystallogr.* **66**, 213–

221 (2010).

51. Terwilliger, T. C., Sobolev, O. V., Afonine, P. V. & Adams, P. D. Automated map

sharpening by maximization of detail and connectivity. *Acta Crystallogr. D Struct.*

Biol. **74**, 545–559 (2018).

52. Emsley, P., Lohkamp, B., Scott, W. G. & Cowtan, K. Features and development of

coot. *Acta Crystallogr. D Biol. Crystallogr.* **66**, 486–501 (2010).

53. Emsley, P. & Cowtan, K. Coot: model-building tools for molecular graphics. *Acta*

Crystallogr. D Biol. Crystallogr. **60**, 2126–2132 (2004).

54. Afonine, P. V. *et al.* Real-space refinement in PHENIX for cryo-EM and

crystallography. *Acta Crystallogr. D Struct. Biol.* **74**, 531–544 (2018).

Figure 1. Overall structure of the GPR30-miniG$_{sqi}$β$_1$γ$_2$-scFv16 complex.
A Unsharpened cryo-EM density map of the GPR30-miniG$_{sqi}$β$_1$γ$_2$-scFv16-Nb35 complex, with the components individually colored. **B** The refined structure of the complex is shown as a ribbon representation.

Figure 2. Receptor structure.
A–C Overall structure of the receptor, viewed from the membrane plane (**A**), extracellular side (**B**), and intracellular side (**C**).

Figure 3. Architecture of the extracellular pocket.
A Molecular surface of the extracellular side. **B** Cross section of the pocket. **C–E** Residues facing pocket B (**C**), pocket C (**D**), and pocket A (**E**). In panel (**E**), only residues with reduced bicarbonate responses are highlighted. **F**, **G** Calcium assay using stable HEK293 cell lines expressing the N-terminal HA-tagged GPR30 wild-type (WT) and mutants. The mutants D111$^{2.56}$A, N276$^{6.52}$A, and Q296^{ECL3}A; E121^{ECL1}A, R122^{ECL1}A, S134$^{3.29}$A, Q215$^{5.39}$A, and E218$^{5.42}$A; D210^{ECL2}A and Q138$^{3.33}$A; and D125^{ECL1}A, C207^{ECL2}A, and P71$^{1.44}$A are highlighted in red, blue, green, and purple, respectively. The cells were stimulated by the indicated concentrations of NaHCO$_3$ at the timepoint of t = 20 sec. The Y axis indicates the difference between the maximum and minimum fluorescent values during 15 to 60 sec. Statistical analysis: each sample was compared with WT cells using Dunnett's T3 multiple comparisons test after Brown-Forsythe and Welch ANOVA tests. **** $p < 0.0001$ compared to WT cells, no significant difference is not shown. Data are presented as mean values ± SEM (**F**) and each replicate with the mean ± SEM (**G**).

Figure 4. G-protein coupling.

A Hydrogen-bonding interactions between the C-terminal α5-helix residues and the receptor. **B** Electrostatic and hydrogen-binding interactions between ICL3 and the α5-helix. **C–H** Interface between ICL2 and Gq, with residues involved in hydrophobic interactions represented by CPK models. G-protein-bound GPCRs used in the comparison are as follows: H1R-Gq (PDB 7DFL, gray), 5-HT2A-Gq (PDB 6WHA, purple), MRGPRX2-Gq (PDB 7S8L, green), B2R-Gq (PDB 7F6I, yellow-green), and GPR103-Gq (PDB 8ZH8, red). **I** Comparison of the angles and positions of α5h and αN relative to the receptor. **J** Superimposition of the Gα subunits.

Figure 5. Structural comparison with related GPCRs

A Structural comparison of GPR30 and AT_2R (PDB 5UNF). **B** Interactions around $P71^{1.44}$ in TM1 of GPR30. **C–E** Conformational changes of TM1 upon agonist binding in CB_1 (**C**), $A_{2A}R$ (**D**), and ET_B (**E**). The agonist-bound states are colored with the respective colors, while the inactive states are colored gray. The PDB codes used in this figure are CB_1-active (PDB 5XRA), CB_1-inactive (PDB 5TGZ), $A_{2A}R$-active(PDB 6GDG), $A_{2A}R$-inactive (PDB 3EML), ET_B-active (PDB 8IY5), and ET_B-inactive (PDB 5X93).

Figure 6. structural comparison with GPR30 bound to Lys05.
A, B superimposition of bicarbonate-GPR30, Lys05-binding GPR30 (PDB 8XOF), and apo-GPR30 (PDB 8XOG) structures, viewed from the extracellular side.
C comparison of the angles and positions of α5h and αN relative to the receptor. **D** superimposition of the Gα subunits.