Transcriptional Signatures of Domestication Revealed by Meta-Analysis of Pig, Chicken, Wild Boar, and Red Junglefowl Gene Expression Data

Motoki Uno [1] and Hidemasa Bono [1,2]

1   Graduate School of Integrated Sciences for Life, Hiroshima University, 3-10-23 Kagamiyama, Higashi-Hiroshima, Hiroshima 739-0046, Japan; m230361@hiroshima-u.ac.jp

2   Genome Editing Innovation Center, Hiroshima University, 3-10-23 Kagamiyama, Higashi-Hiroshima, Hiroshima 739-0046, Japan; bonohu@hiroshima-u.ac.jp

*   Correspondence: bonohu@hiroshima-u.ac.jp; Tel.: +81-82-424-4013

Simple Summary

Domesticated animals, such as pigs and chickens, have undergone significant genetic changes compared to their wild ancestors. We collected gene expression data from pigs, chickens, wild boars, and red junglefowl, and performed a comprehensive analysis. We identified differentially expressed genes by comparing the gene expression patterns of domesticated animals and their wild counterparts. These genes are involved in various biological processes, including immune response, metabolism, and stress response. Notably, domesticated animals exhibited higher expression of genes related to viral resistance and bone weakness, whereas their wild ancestors showed higher expression of genes associated with stress response and energy metabolism. Identifying these differentially expressed genes provides valuable insights into the genetic changes that occurred during the domestication process. Furthermore, these findings have highlighted potential candidate genes that could be targeted in breeding programs to improve the health and productivity of domesticated animals, ultimately contributing to the development of sustainable livestock production and addressing the growing global food demand.

Abstract

Domesticated animals have undergone significant changes in their behavior, morphology, and physiological functions during domestication. To identify the changes in gene expression associated with domestication, we collected RNA-seq data of pigs, chickens, wild boars, and red junglefowl from public databases and performed a meta-analysis. Gene expression was quantified and the expression ratio between domesticated animals and their wild ancestors (DW-ratio) was calculated. Genes were classified as "upregulated," "downregulated," or "unchanged" based on their DW-ratio, and DW-score was calculated for each gene. Gene set enrichment analysis revealed that genes upregulated in pigs were related to defense from viral infection, whereas those upregulated in chickens were associated with aminoglycan and carbohydrate derivative catabolic processes. Genes commonly upregulated in pigs and chickens are involved in the immune response, olfactory learning, epigenetic regulation, cell division, and extracellular matrix. In contrast, genes upregulated in wild boar and red junglefowl are related to stress response, cell proliferation, cardiovascular function, neural regulation, and energy metabolism. These findings provide valuable insights into the genetic basis of the domestication process and highlight potential candidate genes for breeding applications.

Keywords: Sus scrofa domesticus, Sus scrofa, Gallus gallus, domesticated animals, breeding, gene set enrichment analysis

1. Introduction

With the rapid growth of the global population, food shortage has become a serious concern, and it is estimated that the global food demand in 2050 will be approximately 1.5 times that of 2010 [1]. Pigs and chickens are crucial protein sources for humans [2] and their breeding and improvement are essential to address this challenge. Pigs provide meat, whereas chickens supply both meat and eggs, and both are used in pharmaceutical manufacturing [3].

Pigs were domesticated from wild boars [4], while chickens are presumed to have been domesticated from red junglefowl [5]. Through domestication, these animals have been brought under human control and have been

selectively bred for specific traits useful to humans, such as faster growth rate, improved meat quality, and increased 48

litter size [6,7]. This process has significantly altered the animals' behavior, morphology, and physiological functions. 49

However, excessive selective breeding has led to health problems and reduced their reproductive capacity [8]. Further 50

improvement of these animals is necessary to address the food crisis and solve these problems. 51

Quantitative trait locus (QTL) analysis has been used to identify trait-associated loci, such as the NR6A1 locus in 52

pigs, which is associated with the vertebral number [9]. However, QTL analysis has limitations in directly identifying 53

genes related to traits, capturing changes in gene expression, and detecting genes without sequence variation. In con- 54

trast, the relationship between changes in gene expression and traits has become apparent, as exemplified by the myo- 55

statin gene, where decreased expression leads to increased muscle mass [10]. 56

Transcriptomic analysis is a powerful method for identifying domestication-associated changes in gene expres- 57

sion. However, small sample sizes and biases in sampled tissues and breeds make it difficult to identify genes that 58

universally contribute to domestication. Meta-analyses overcome these challenges by integrating data from different 59

breeds, tissues, and growth stages, thereby enabling the elucidation of insights underlying domestication that individ- 60

ual studies cannot reveal [11]. 61

Here, we aimed to identify gene expression changes associated with domestication by collecting gene expression 62

data for pigs, chickens, wild boars, and red junglefowl from public databases and performing a meta-analysis. To the 63

best of our knowledge, this is the first meta-analysis to demonstrate the differences in gene expression between domes- 64

ticated animals and their wild ancestors, specifically pigs and wild boars, and chickens and red junglefowl. Furthermore, 65

this meta-analysis is the first to integrate multiple studies and identify genes that are commonly differentially expressed 66

across species, both in domesticated animals and their wild ancestors. The meta-analysis revealed that genes related to 67

viral infection defense and leucine uptake were differentially expressed between pigs and wild boars, while genes in- 68

volved in aminoglycan catabolic processes and the response to corticotropin-releasing hormone were differentially 69

expressed between chickens and red junglefowl. Furthermore, we identified genes commonly upregulated in domesti- 70

cated animals, including those involved in the immune response, olfactory learning, and epigenetic regulation. In con- 71

trast, genes associated with stress response, cell proliferation, cardiovascular function, and energy metabolism were 72

found to be commonly upregulated in the wild ancestors. The insights gained from comparing domesticated animals 73

with their wild ancestors will provide a genetic basis for domestication, which can be combined with technologies such 74

as genome editing to enable more efficient and sustainable livestock breeding. This meta-analysis is expected to capture 75

changes in gene expression that may have been overlooked in individual studies. The findings of this study are expected 76

to contribute to addressing food problems and developing sustainable livestock breeding strategies. 77

2. Materials and Methods 78

2.1. Curation of Gene Expression Data from Public Databases 80

RNA-seq data were obtained from public databases, the National Center for Biotechnology Information Gene 81

Expression Omnibus (NCBI GEO) [12] and ArrayExpress from European Bioinformatics Institute BioStudies (EBI Bio- 82

Studies) [13]. For NCBI GEO, a search was conducted using "Sus scrofa" and "Gallus gallus." To further refine the 83

search results, the filter "Expression profiling by high throughput sequencing" was applied. In EBI BioStudies' Ar- 84

rayExpress, a search was performed with the following parameters: Organism set to either "Sus scrofa" or "Gallus 85

gallus," Technology set to "sequencing assay," and Assay by Molecule set to "rna assay" to narrow down the results. 86

Based on search results, manual curation was performed to collect pairs of domesticated animals and their wild ances- 87

tors from the same project. 88

2.2. Gene Expression Quantification 90

FASTQ format files were obtained from the NCBI GEO using the fsterq-dump (version 3.8.0) command from the 91

SRA toolkit [14], based on the respective accession numbers. Quality control of the raw reads was performed using Trim 92

Galore (version 0.6.10) [15]. Transcript quantification was performed using Salmon (version 1.10.2) [16], with Sscrofa11.1 93

being used as the reference cDNA for pig and wild boar samples, and bGalgal1.mat.broiler.GRCg7b being used as the reference cDNA for chicken and red junglefowl samples. The resulting quantified RNA-seq data were expressed as transcripts per million (TPM). Transcript-level TPM values were summarized at the gene level using the tximport (version 1.28.0) [17] package in R.

2.3. Calculation of DW-Ratio

Gene expression data were normalized to the DW-ratio. Each letter of DW respectively stands for Domesticated (D) and Wild (W). The DW-ratio was calculated using the following formula:

$$DW - ratio = \frac{DTPM + 1}{WTPM + 1}$$

When calculating the DW-ratio, 1 was added to the TPM values to avoid division by zero and to handle genes with zero expression values.

2.4. Classification of Differential Expressed Genes (DEGs)

To evaluate the genes that exhibited changes in expression between domesticated animals and their wild ancestors, all genes were classified into three groups. When the DW-ratio was above a specified threshold, the gene was classified as "upregulated." In contrast, when the DW-ratio was below a specified threshold, the gene was classified as "downregulated." Genes that did not fall into either category were classified as "unchanged." For the upregulated category, thresholds of 1.5-, 2-, 3-, 4-, and 5-fold were tested and a 1.5-fold threshold was adopted. For the downregulated category, thresholds of 1/1.5-, 1/2-, 1/3-, 1/4-, and 1/5- were tested, and a 1/1.5-fold threshold was adopted.

2.5. Calculation of DW-Score

To integrate and evaluate the DEGs across different experiments, the DW-score was calculated. The DW-score for each gene was determined by subtracting the number of domesticated-wild pairs in which the gene was classified as "downregulated" from the number of pairs in which it was classified as "upregulated." The DW-ratio and DW-score were computed using code that had been utilized in a previous study [18].

2.6. Gene Set Enrichment Analysis

Based on the obtained DW-score, the top- and bottom-ranked genes were selected, and gene set enrichment anal-

ysis was performed using ShinyGO 0.80 [19]. For the pig and wild boar analysis, the species was set to "Pig-Duroc genes

Sscrofa11.1", and the "GO Biological Process" was selected as the Pathway database. All other parameters were main-

tained at their default settings. For the chicken and red junglefowl comparison, no enriched terms were obtained when

using the "Chicken maternal Broiler genes bGalGal1.mat.broiler.GRCg7b" annotation. Therefore, the gene IDs were

converted to human gene IDs, and the enrichment analysis was performed using "Human genes GRCh38.p13".

2.7. Commonly Upregulated Genes in Domesticated Animals and Wild Ancestors

To enable cross-species analysis, the gene IDs from each species were converted into human gene IDs (GRCh38).

Based on the DW-score, ranked lists of the top- and bottom-ranked genes were generated. The overlap of commonly

upregulated genes between domesticated animals and wild ancestors was investigated and visualized using the Venn

diagram tool [20].

3. Results

3.1. Overview of this study

In this study, we performed a meta-analysis to identify DEGs between domesticated animals and their wild an-

cestors. The research workflow is illustrated in Fig 1.

First, we manually curated and obtained gene expression data from public databases for pigs, chickens, wild boars,

and red jungle fowl. The acquired RNA-seq datasets were processed for gene expression quantification and calculation

of the expression ratio between domesticated animals and their wild ancestors (DW-ratio).

Next, we classified genes as "upregulated," "downregulated," or "unchanged" based on their DW-ratio and cal- 142

culated a DW-score for each gene. Gene set enrichment analysis was conducted using the top-ranked genes based on 143

their DW-score for each combination of domesticated animals and wild ancestors. 144

Finally, to identify genes that are commonly differentially expressed across different species, we converted gene 145

IDs to human gene IDs and examined the overlap of DEGs between domesticated animals and their wild ancestors. 146

Figure 1. The workflow of this study. Gene expression data of domesticated animals and their wild ancestors were obtained from public databases. The datasets were processed to quantify gene expression and calculate the expression ratio between domesticated animals and wild ancestors (DW-ratio). Genes were classified based on their DW-ratio, and a DW-score was calculated for each gene. Gene set enrichment analysis was performed using top-ranked genes for each domesticated-wild pair. Gene IDs were converted to human gene IDs to identify commonly differentially expressed genes across species.

DEG, differentially expressed gene; DW, domestic-wild.

3.2. Curation and breakdown of RNA-seq data obtained from public databases

In this study, we obtained 177 RNA-seq datasets from nine projects in the NCBI GEO [12] and EBI BioStudies [13] databases. To enable the integration and analysis of data from different research projects, microarray data were excluded and only RNA-seq data were obtained. Additionally, all pairs formed were within single projects to minimize the differences in gene expression between projects. We acquired 46 pairs of pigs and their wild ancestors, wild boars, and 59 pairs of chickens and their presumed wild ancestors, red junglefowl. In the pig-wild boar pairs, muscle tissue was the most common tissue examined, being found in 18 pairs, followed by the spleen in 12 pairs, and the brain in 11 pairs. In pairs of chickens and red junglefowl, the brain was the most common, with 19 pairs, followed by the testes with 10 pairs, and the eyes with 8 pairs. The metadata of the curated dataset, including accession IDs, breeds, sampled tissues, ages, and sex of the collected RNA-seq data, are presented in Supplementary Tables 1 and 2.

Figure 2. Tissue distribution of the curated RNA-seq data pairs. A total of 46 pig-wild boar pairs and 59 chicken-red junglefowl pairs of RNA-seq data were obtained. The left pie chart shows the proportion of pairs for each tissue type in the pig-wild boar pairs, with muscle tissue having the highest proportion. The right pie chart represents the tissue distribution in the chicken-red junglefowl pairs, with brain tissue having the highest proportion.

3.3. Characterization of DEGs and Enrichment Analysis of Pigs and Boars

After gene expression quantification using Salmon [16] (Supplementary Tables 3 and 4), the expression ratio between domesticated animals and their wild ancestors (DW-ratio) was calculated. A 1.5-fold threshold was chosen for both upregulated and downregulated genes to capture a comprehensive set of DEGs, while minimizing the omission of potentially relevant candidates. For each sample, genes with a DW-ratio ≥1.5 were classified as "upregulated," those with a DW-ratio less than or equal to 1/1.5 were classified as "downregulated," and the remaining genes were classified as "unchanged." For each gene a DW-score was calculated by subtracting the number of samples in which the gene was classified as "downregulated" from the number of samples in which it was classified as "upregulated." By ranking the genes based on their DW-score, a list of DEGs between domesticated animals and their wild ancestors was obtained. The rankings of DEGs at different thresholds are shown in Supplementary Tables 5 and 6. To investigate whether there were any functional biases in the list of DEGs, enrichment analysis was performed using ShinyGO 0.80 [19].

Figure 3. Scatter plot of the DW-score and gene set enrichment analysis of upregulated in pigs and wild boars. (A) The scatter plot shows the DW-score of each gene, with red dots representing genes upregulated in pigs (positive DW-score values) and blue dots representing genes upregulated in wild boars (negative DW-score values). (B) Gene set enrichment analysis for genes upregulated in pigs. The analysis was performed using the top 218 genes based on their DW-scores. (C) Gene set enrichment analysis for genes upregulated in wild boars. The analysis was performed the top 239 genes based on their DW-scores.

For the pig-wild boar pairs, enrichment analysis was conducted using DEGs (218 and 239 genes upregulated in pigs and wild boars, respectively; Supplementary Tables 7 and 8). The threshold for the number of genes used in enrichment analysis was determined based on the DW-scores that showed biases in Gene Ontology (GO) term function, corresponding to approximately 1% of the total number of genes in the reference genome. The enrichment analysis of the gene list upregulated in pigs revealed a significant enrichment of GO terms related to viral infection defense, such as "Negative regulation of viral genome replication" and "Positive regulation of interferon-beta production." The enrichment analysis results for genes upregulated in wild boars showed significant enrichment of GO terms related to leucine uptake, such as "Leucine import across plasma membrane" and "L-leucine import across plasma membrane" (Supplementary Tables 9 and 10).

3.4. Characterization of DEGs and enrichment analysis in chickens and red junglefowl

An enrichment analysis was performed for chickens and their wild ancestor, red junglefowl, based on the obtained DW-score. In this case, no functional enrichment was observed; therefore, the gene IDs were converted to human gene IDs, and an enrichment analysis was performed using 223 and 172 chicken and red junglefowl genes, respectively (Supplementary Tables 11 and 12). The enrichment analysis results for genes upregulated in chickens revealed two terms: "Aminoglycan catabolic process" and "Carbohydrate derivative catabolic process." The GO term "Response to corticotropin-releasing hormone" was the top-ranked term for genes upregulated in red junglefowl (Supplementary Tables 13 and 14).

Figure 4. Scatter plot of the DW-score and gene set enrichment analysis of upregulated genes in chickens and red junglefowl. (A) The scatter plot shows the DW-score of each gene, with red dots representing genes upregulated in chickens (positive DW-score values) and blue dots representing genes upregulated in red junglefowl (negative DW-score values). (B) Gene set enrichment analysis for genes upregulated in chickens. The analysis was performed using the top 223 genes based on their DW-scores. (C) Gene set enrichment analysis for genes upregulated in red junglefowl. The analysis was performed with the top 172 genes based on their DW-scores.

3.5. Common DEGs in domesticated animals and their wild ancestors

To investigate whether genes were commonly differentially expressed across different domesticated species, the gene IDs of each species were converted into human gene IDs (Supplementary Tables 6 and 15). The overlap of DEGs was examined between domesticated animals (pigs and chickens) and their wild ancestors (wild boars and red junglefowl). When examining the overlap between the top 240 upregulated genes in pigs and the top 223 upregulated genes in chickens (Supplementary Tables 16 and 17), we found that 10 genes were upregulated in both pigs and chickens. These genes were EZH2, DPP7, PRTFDC1, CEP20, CMPK2, CX3CR1, USP18, NID1, CXCL13, and PHF11 (Table 1). When examining the overlap between the top 206 and 172 genes upregulated in wild boars and red junglefowl, respectively (Supplementary Tables 18 and 19), we found that eight genes were upregulated in both wild boars and red junglefowl: TAGLN, HSPA8, POPDC2, MAFF, HSPB8, SCN1B, NT5C1A, and BAG3 (Table 2).

Table 1. Upregulated genes in common between domesticated animals (pig and chicken)

Gene ID	Gene Name	Description
ENSG00000106462	*EZH2*	Enhancer of zeste 2 polycomb repressive complex 2 subunit
ENSG00000176978	*DPP7*	Dipeptidyl peptidase 7
ENSG00000184979	*USP18*	Ubiquitin specific peptidase 18
ENSG00000099256	*PRTFDC1*	Phosphoribosyl transferase domain containing 1
ENSG00000133393	*CEP20*	Centrosomal protein 20
ENSG00000116962	*NID1*	Nidogen 1
ENSG00000156234	*CXCL13*	C-X-C motif chemokine ligand 13
ENSG00000134326	*CMPK2*	Cytidine/uridine monophosphate kinase 2
ENSG00000168329	*CX3CR1*	C-X3-C motif chemokine receptor 1
ENSG00000136147	*PHF11*	PHD finger protein 11

Table 2. Upregulated genes in common between wild ancestors (wild boar and red junglefowl)

Gene ID	Gene Name	Description
ENSG00000149591	*TAGLN*	Transgelin
ENSG00000109971	*HSPA8*	Heat shock protein family A (Hsp70) member 8
ENSG00000121577	*POPDC2*	Popeye domain containing 2
ENSG00000185022	*MAFF*	MAF bZIP transcription factor F
ENSG00000152137	*HSPB8*	Heat shock protein family B (small) member 8
ENSG00000105711	*SCN1B*	Sodium voltage-gated channel beta subunit 1
ENSG00000116981	*NT5C1A*	5'-nucleotidase, cytosolic IA
ENSG00000151929	*BAG3*	BAG cochaperone 3

Figure 5. Upregulated genes in common among domesticated animals (pig and chicken) and their wild ancestors (wild boar and red junglefowl). (A) Venn diagram showing the overlap of the top upregulated genes in pigs (240 genes) and chickens (223 genes). Ten genes were found to be upregulated in both domesticated animals. (B) Venn diagram showing the overlap of the top upregulated genes in wild boars (206 genes) and red junglefowl (RJF) (172 genes). Eight genes were found to be upregulated in both wild ancestors.

4. Discussion

In this study, the gene expression data of domesticated animals and their wild ancestors were manually curated

and obtained from public databases [12,13]. Ultimately, 46 pigs-wild boars pairs and 59 chicken-red junglefowl pairs

were obtained. Using these data, a meta-analysis was performed to identify DEGs between domesticated animals and

their wild ancestors. To characterize DEGs, enrichment analysis was conducted for each combination of domesticated

animals and their wild ancestors. Furthermore, to identify genes that were commonly differentially expressed across

different species, all gene IDs were converted to human gene IDs. Subsequently, the DEGs shared between domesticated

animals, pigs, and chickens, and between their wild ancestors, wild boars and red junglefowl, respectively, were inves-

tigated. The list of DEGs included several genes that have been reported to be involved in the traits of domesticated

animals. This suggests that the gene expression dataset and meta-analysis method used in this study accurately reflect

the scientific knowledge regarding gene expression differences between domesticated animals and their wild ancestors.

The list of genes upregulated in pigs included several genes previously reported to be associated with pig traits,

such as muscle development and fat accumulation, including IGF2BP3, ENSSSCG00000035293 (IGF2), NCAPG, FGF11,

and FABP4. IGF2 is a well-known QTL associated with muscle growth and back fat thickness in pigs [21,22], and its

expression level is correlated with skeletal muscle mass [23]. Additionally, IGF2BP3 is involved in regulating IGF2 ex-

pression [24], suggesting that it may play a role in controlling skeletal muscle growth in pigs [25]. NCAPG, which is

essential for maintaining chromosome structure [26], is associated with traits such as body size and muscle development

in various domesticated animals [27–34]. FDF11 has been suggested to regulate adipocyte differentiation through the

expression of PPARγ, a key transcription factor controlling adipocyte differentiation [35]. FABP4, which binds to free

fatty acids with high affinity and functions as a lipid transport protein [36], is upregulated in muscle-specific PPARγ-

overexpressing pigs and is thought to be involved in intramuscular fat deposition [37]. Moreover, among the genes

upregulated in pigs, genes involved in viral or bacterial infections, such as IFI44L, IL33, and RSAD2, were ranked at the 259

top of the list. Furthermore, when enrichment analysis was performed using 218 genes upregulated in pigs, GO terms 260

related to viral infection defense, such as "Negative regulation of viral genome replication" and "Positive regulation of 261

interferon-beta production," were ranked among the top terms. Notably, the expression of sensor proteins that bind to 262

viral nucleic acids, such as RIGI, IFIH1 (encoding MDA5), DHX58 (encoding LGP2), RIF2AK2 (encoding PKR), OAS1, 263

and OAS2 [38–40], was found to be elevated. These findings suggest that, during their domestication process, pigs with 264

higher resistance to viruses and pathogenic bacteria were selected. 265

The enrichment analysis using 239 genes upregulated in wild boars revealed that GO terms related to leucine 266

uptake, such as "Leucine import across plasma membrane" and "L-leucine import across plasma membrane," were the 267

most significantly enriched. Like other animals, leucine is also an essential amino acid for pigs [41] and is known to play 268

a crucial role in muscle synthesis. Moreover, leucine has been suggested to promote muscle growth by enhancing 269

mTORC1 activation and its downstream effectors [42]. Furthermore, leucine supplementation suppresses obesity in 270

obese mice fed a high-fat diet [43]. In addition, leucine supplementation may improve leptin sensitivity and suppress 271

appetite [44]. Genes upregulated in wild boars can also be considered downregulated in pigs. Therefore, active leucine 272

uptake in wild boars, which is impaired in pigs, may contribute to fat accumulation and appetite in pigs. 273

When enrichment analysis was performed on chicken and red junglefowl pairs using chicken GO annotations, no 274

enriched functions were observed. Therefore, when chicken gene IDs were converted to human gene IDs, and enrich- 275

ment analysis was performed using human GO annotations, enriched functions were obtained for both chicken and red 276

junglefowl. This may indicate that current genetic research on chickens is insufficient. Enrichment analysis using a list 277

of 223 upregulated genes that were upregulated in chickens yielded two terms: "Aminoglycan catabolic proc." and 278

"Carbohydrate derivative catabolic process." Selection for rapid growth rates resulted in the skeletal development of 279

broilers not keeping pace with their body weight gain, leading to fractures and locomotor disorders [8]. Aminoglycans, 280

a more general term for glycosaminoglycans, promote bone and joint healing [45]. Research is also being conducted to 281

promote the structural development of bones and joints by incorporating glycosaminoglycans into feed [46]. The en- 282

richment of the "Aminoglycan catabolic proc." term in the enrichment analysis using the gene set upregulated in chick- 283

ens may reflect weakness of chicken bones and locomotor disorders. 284

Enrichment analysis using 172 upregulated genes in red junglefowl revealed a strong enrichment for the GO term 285

"Response to corticotropin-releasing hormone," with three genes – NR4A1, NR4A2, and NR4A3 – assigned to this GO 286

term. The NR4A family, also known as the orphan nuclear receptors, consists of NR4A1 (Nur77), NR4A2 (Nurr1), and 287

NR4A3 (Nor-1), which are thought to be involved in diverse functions in various tissues, including cell proliferation, 288

apoptosis, and fatty acid metabolism [47,48]. All three genes belonging to the NR4A family are known to be rapidly and 289

transiently induced following corticotropin-releasing hormone treatment [49] and activate the hypothalamic-pituitary- 290

adrenal (HPA) axis, suggesting that NR4A members may be involved in CRH-mediated stress responses via the HPA 291

axis [50]. The results of the enrichment analysis using the list of upregulated genes in red junglefowl may reflect an 292

active stress response, which could be attributed to the high level of vigilance exhibited by this species. 293

We examined the overlap of the DEGs among the domesticated animals; 10 genes (DPP7, CX3CR1, USP18, 294

CXCL13, PHF11, CMPK2, PRTFDC1, CEP20, EZH2, and NID1) were upregulated in pigs (240 genes) and chickens (223 295

genes). In contrast, eight genes (MAFF, HSPA8, HSPB8, BAG3, TAGLN, POPDC2, SCN1B, and NT5C1A) were com- 296

monly upregulated in wild boars (206 genes) and red junglefowl (172 genes). The upregulated genes in the domesticated 297

animals were involved in various biological processes, including immune response (DPP7 [51], CX3CR1 [52], USP18 298

[53], CXCL13 [54], PHF11 [55], CMPK2 [56]), olfactory learning (PRTFDC1 [57]), epigenetic regulation (EZH2 [58]), cell 299

division (CEP20 [59]), and extracellular matrix (NID1 [60]). These changes in gene expression may contribute to the 300

adaptation of the immune system, changes in olfaction, regulation of growth and development, and morphological 301

changes associated with domestication. In contrast, the upregulated genes in the wild counterparts were related to stress 302

response (MAFF [61], HSPA8 [62], HSPB8 [63], BAG3 [63]), cell proliferation and differentiation (TAGLN [64]), cardio-vascular function (POPDC2 [65]), neural regulation (SCN1B [66]), and energy metabolism (NT5C1A [67]). These changes in gene expression suggest that their wild counterparts maintained the physiological functions necessary for adaptation to the natural environment. Furthermore, these DEGs were upregulated in domesticated animals and their wild ancestors, suggesting a strong association with the domestication process. This indicates that these genes may serve as universal candidates for breeding applications.

This study has several limitations that should be taken into consideration. First, the selection of DEGs was not based on statistical methods, necessitating caution in interpreting the results. Future studies should employ rigorous statistical approaches to identify DEGs with higher confidence. Second, potential biases may have been introduced due to undisclosed metadata. The lack of complete and accurate metadata for some datasets could have influenced the interpretation of the results and the comparison of gene expression patterns across different studies. Efforts should be made to ensure the transparency and completeness of metadata in public databases to facilitate more reliable meta-analyses. Third, the reference transcriptome used for the wild ancestors (wild boar and red junglefowl) were based on their domesticated counterparts (pig and chicken) due to the lack of well-annotated data. This limitation may have affected the accuracy of gene expression quantification and the identification of species-specific genes. Future research will require high-quality reference data for the wild ancestors to enable accurate comparisons between domesticated animals and their wild ancestors. Despite these limitations, this meta-analysis provides a valuable starting point for understanding the gene expression changes associated with domestication and highlights the need for further research to address these challenges and validate the findings.

To our knowledge, this study is the first meta-analysis to demonstrate differences in gene expression between these domesticated animals and their wild ancestors. This is the first meta-analysis to identify genes that are commonly differentially expressed across different species in both domesticated animals and their wild ancestors. The insights

gained from this meta-analysis provide a valuable foundation for understanding the genetic basis of the domestication   325

process and highlighting potential candidate genes for breeding applications.   326

5. Conclusions   328

In conclusion, this meta-analysis identified DEGs associated with the domestication process. These genes are in-   329

volved in traits such as muscle development, fat accumulation, viral resistance, and bone weakness in domesticated   330

animals. Genes commonly upregulated in domesticated animals were related to the immune response, olfactory learn-   331

ing, epigenetic regulation, cell division, and the extracellular matrix, whereas those upregulated in wild ancestors were   332

associated with the stress response, cell proliferation, cardiovascular function, neural regulation, and energy metabo-   333

lism.   334

The findings of this study provide valuable insights into the genetic basis of domestication and highlight potential   335

candidate genes for breeding applications. The DEGs identified in this study may contribute to the adaptation of do-   336

mesticated animals to their environment and the development of desired traits. Furthermore, the genes commonly up-   337

regulated across different domesticated species suggest a strong association with the domestication process and their   338

potential as universal candidate genes for breeding. In future research, candidate genes for breeding should be identi-   339

fied using a wider range of species, and the possibility of applying these candidate genes to efficient breeding programs   340

should be explored using techniques such as genome editing.   341

Supplementary Materials: The Supplementary Material for this article can be found online   343
(https://doi.org/10.6084/m9.figshare.25793967).   344

Author Contributions: Conceptualization, M.U. and H.B.; methodology, M.U. and H.B.; software, M.U. and H.B.; vali-   346
dation, M.U. and H.B.; formal analysis, M.U. and H.B.; investigation, M.U.; resources, M.U. and H.B.; data curation,   347
M.U.; writing—original draft preparation, M.U.; writing—review and editing, M.U. and H.B.; visualization, M.U.; su-   348
pervision, H.B.; project administration, H.B.; funding acquisition, H.B. All authors have read and agreed to the pub-   349
lished version of the manuscript.   350

Funding: This study was supported by the Center of Innovation for Bio-Digital Transformation (BioDX), an open innovation platform for industry-academia co-creation (COI-NEXT), and the Japan Science and Technology Agency (JST, COI-NEXT; grant number JPMJPF2010).

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: The data presented in this study are publicly available at FigShare (https://doi.org/10.6084/m9.figshare.25793967).

Conflicts of Interest: The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript; or in the decision to publish the results.

References

1. van Dijk, M.; Morley, T.; Rau, M.L.; Saghai, Y. A Meta-Analysis of Projected Global Food Demand and Population at Risk of Hunger for the Period 2010–2050. *Nat. Food* **2021**, *2*, 494–501, doi:10.1038/s43016-021-00322-9.
2. OECD *OECD-FAO Agricultural Outlook 2023-2032*; Organisation for Economic Co-operation and Development: Paris, 2023;
3. Ulmer, J.B.; Valley, U.; Rappuoli, R. Vaccine Manufacturing: Challenges and Solutions. *Nat. Biotechnol.* **2006**, *24*, 1377–1383, doi:10.1038/nbt1261.
4. Giuffra, E.; Kijas, J.M.H.; Amarger, V.; Carlborg, Ö.; Jeon, J.-T.; Andersson, L. The Origin of the Domestic Pig: Independent Domestication and Subsequent Introgression. *Genetics* **2000**, *154*, 1785–1791, doi:10.1093/genetics/154.4.1785.
5. Lawal, R.A.; Martin, S.H.; Vanmechelen, K.; Vereijken, A.; Silva, P.; Al-Atiyat, R.M.; Aljumaah, R.S.; Mwacharo, J.M.; Wu, D.-D.; Zhang, Y.-P.; et al. The Wild Species Genome Ancestry of Domestic Chickens. *BMC Biol.* **2020**, *18*, 13, doi:10.1186/s12915-020-0738-1.
6. Quiniou, N.; Dagorn, J.; Gaudré, D. Variation of Piglets' Birth Weight and Consequences on Subsequent Performance. *Livest. Prod. Sci.* **2002**, *78*, 63–70, doi:10.1016/S0301-6226(02)00181-1.
7. Zuidhof, M.J.; Schneider, B.L.; Carney, V.L.; Korver, D.R.; Robinson, F.E. Growth, Efficiency, and Yield of Commercial Broilers from 1957, 1978, and 20051. *Poult. Sci.* **2014**, *93*, 2970–2982, doi:10.3382/ps.2014-04291.
8. Olkowski, A.A.; Laarveld, B.; Wojnarowicz, C.; Chirino-Trejo, M.; Chapman, D.; Wysokinski, T.W.; Quaroni, L. Biochemical and Physiological Weaknesses Associated with the Pathogenesis of Femoral Bone Degeneration in Broiler Chickens. *Avian Pathol.* **2011**, *40*, 639–650, doi:10.1080/03079457.2011.626017.
9. Mikawa, S.; Morozumi, T.; Shimanuki, S.-I.; Hayashi, T.; Uenishi, H.; Domukai, M.; Okumura, N.; Awata, T. Fine Mapping of a Swine Quantitative Trait Locus for Number of Vertebrae and Analysis of an Orphan Nuclear Receptor, Germ Cell Nuclear Factor (NR6A1). *Genome Res.* **2007**, *17*, 586–593, doi:10.1101/gr.6085507.
10. McPherron, A.C.; Lawler, A.M.; Lee, S.-J. Regulation of Skeletal Muscle Mass in Mice by a New TGF-p Superfamily Member. *Nature* **1997**, *387*, 83–90, doi:10.1038/387083a0.

11. Bono, H.; Hirota, K. Meta-Analysis of Hypoxic Transcriptomes from Public Databases. *Biomedicines* **2020**, *8*, 10, doi:10.3390/biomedicines8010010.

12. Barrett, T.; Wilhite, S.E.; Ledoux, P.; Evangelista, C.; Kim, I.F.; Tomashevsky, M.; Marshall, K.A.; Phillippy, K.H.; Sherman, P.M.; Holko, M.; et al. NCBI GEO: Archive for Functional Genomics Data Sets—Update. *Nucleic Acids Res.* **2013**, *41*, D991–D995, doi:10.1093/nar/gks1193.

13. Athar, A.; Füllgrabe, A.; George, N.; Iqbal, H.; Huerta, L.; Ali, A.; Snow, C.; Fonseca, N.A.; Petryszak, R.; Papatheodorou, I.; et al. ArrayExpress Update – from Bulk to Single-Cell Expression Data. *Nucleic Acids Res.* **2019**, *47*, D711–D715, doi:10.1093/nar/gky964.

14. Download SRA Sequences from Entrez Search Results. Available online: https://www.ncbi.nlm.nih.gov/sra/docs/sradownload/ (accessed on 1 June 2023).

15. Krueger, F. Babraham Bioinformatics-Trim Galore! Available online: https://www.bioinformatics.babraham.ac.uk/projects/trim_galore/ (accessed on 1 July 2023).

16. Patro, R.; Duggal, G.; Love, M.I.; Irizarry, R.A.; Kingsford, C. Salmon Provides Fast and Bias-Aware Quantification of Transcript Expression. *Nat. Methods* **2017**, *14*, 417–419, doi:10.1038/nmeth.4197.

17. Tximport Available online: http://bioconductor.org/packages/tximport/ (accessed on 1 September 2023).

18. Ono, Y.; Bono, H. Multi-Omic Meta-Analysis of Transcriptomes and the Bibliome Uncovers Novel Hypoxia-Inducible Genes. *Biomedicines* **2021**, *9*, 582, doi:10.3390/biomedicines9050582.

19. Ge, S.X.; Jung, D.; Yao, R. ShinyGO: A Graphical Gene-Set Enrichment Tool for Animals and Plants. *Bioinformatics* **2020**, *36*, 2628–2629, doi:10.1093/bioinformatics/btz931.

20. Draw Venn Diagram Available online: https://bioinformatics.psb.ugent.be/webtools/Venn/ (accessed on 1 January 2024).

21. Jeon, J.-T.; Carlborg, Ö.; Törnsten, A.; Giuffra, E.; Amarger, V.; Chardon, P.; Andersson-Eklund, L.; Andersson, K.; Hansson, I.; Lundström, K.; et al. A Paternally Expressed QTL Affecting Skeletal and Cardiac Muscle Mass in Pigs Maps to the IGF2 Locus. *Nat. Genet.* **1999**, *21*, 157–158, doi:10.1038/5938.

22. Nezer, C.; Moreau, L.; Brouwers, B.; Coppieters, W.; Detilleux, J.; Hanset, R.; Karim, L.; Kvasz, A.; Leroy, P.; Georges, M. An Imprinted QTL with Major Effect on Muscle Mass and Fat Deposition Maps to the IGF2 Locus in Pigs. *Nat. Genet.* **1999**, *21*, 155–156, doi:10.1038/5935.

23. Van Laere, A.-S.; Nguyen, M.; Braunschweig, M.; Nezer, C.; Collette, C.; Moreau, L.; Archibald, A.L.; Haley, C.S.; Buys, N.; Tally, M.; et al. A Regulatory Mutation in IGF2 Causes a Major QTL Effect on Muscle Growth in the Pig. *Nature* **2003**, *425*, 832–836, doi:10.1038/nature02064.

24. Yin, H.; He, H.; Shen, X.; Zhao, J.; Cao, X.; Han, S.; Cui, C.; Chen, Y.; Wei, Y.; Xia, L.; et al. miR-9-5p Inhibits Skeletal Muscle Satellite Cell Proliferation and Differentiation by Targeting IGF2BP3 through the IGF2-PI3K/Akt Signaling Pathway. *Int. J. Mol. Sci.* **2020**, *21*, 1655, doi:10.3390/ijms21051655.

25. Yang, Y.; Fan, X.; Yan, J.; Chen, M.; Zhu, M.; Tang, Y.; Liu, S.; Tang, Z. A Comprehensive Epigenome Atlas Reveals DNA Methylation Regulating Skeletal Muscle Development. *Nucleic Acids Res.* **2021**, *49*, 1313–1329, doi:10.1093/nar/gkaa1203.

26. Kimura, K.; Cuvier, O.; Hirano, T. Chromosome Condensation by a Human Condensin Complex in *Xenopus* Egg Extracts*. *J. Biol. Chem.* **2001**, *276*, 5417–5420, doi:10.1074/jbc.C000873200.

27. Tetens, J.; Widmann, P.; Kühn, C.; Thaller, G. A Genome-Wide Association Study Indicates LCORL/NCAPG as a Candidate Locus for Withers Height in German Warmblood Horses. *Anim. Genet.* **2013**, *44*, 467–471, doi:10.1111/age.12031.

28. Signer-Hasler, H.; Flury, C.; Haase, B.; Burger, D.; Simianer, H.; Leeb, T.; Rieder, S. A Genome-Wide Association Study Reveals Loci Influencing Height and Other Conformation Traits in Horses. *PLOS ONE* **2012**, *7*, e37282, doi:10.1371/journal.pone.0037282.

29. Lindholm-Perry, A.K.; Sexten, A.K.; Kuehn, L.A.; Smith, T.P.; King, D.A.; Shackelford, S.D.; Wheeler, T.L.; Ferrell, C.L.; Jenkins, T.G.; Snelling, W.M.; et al. Association, Effects and Validation of Polymorphisms within the NCAPG - LCORL Locus Located on BTA6 with Feed Intake, Gain, Meat and Carcass Traits in Beef Cattle. *BMC Genet.* **2011**, *12*, 103, doi:10.1186/1471-2156-12-103.

30. Hoshiba, H.; Setoguchi, K.; Watanabe, T.; Kinoshita, A.; Mizoshita, K.; Sugimoto, Y.; Takasuga, A. Comparison of the Effects Explained by Variations in the Bovine PLAG1 and NCAPG Genes on Daily Body Weight Gain, Linear Skeletal Measurements and Carcass Traits in Japanese Black Steers from a Progeny Testing Program. *Anim. Sci. J.* **2013**, *84*, 529–534, doi:10.1111/asj.12033.

31. Zhang, W.; Li, J.; Guo, Y.; Zhang, L.; Xu, L.; Gao, X.; Zhu, B.; Gao, H.; Ni, H.; Chen, Y. Multi-Strategy Genome-Wide Association Studies Identify the DCAF16-NCAPG Region as a Susceptibility Locus for Average Daily Gain in Cattle. *Sci. Rep.* **2016**, *6*, 38073, doi:10.1038/srep38073.

32. Liu, Y.; Duan, X.; Chen, S.; He, H.; Liu, X. *NCAPG* Is Differentially Expressed during Longissimus Muscle Development and Is Associated with Growth Traits in Chinese Qinchuan Beef Cattle. *Genet. Mol. Biol.* **2015**, *38*, 450–456, doi:10.1590/S1415-475738420140287.

33. Setoguchi, K.; Watanabe, T.; Weikard, R.; Albrecht, E.; Kühn, C.; Kinoshita, A.; Sugimoto, Y.; Takasuga, A. The SNP c.1326T>G in the Non-SMC Condensin I Complex, Subunit G (NCAPG) Gene Encoding a p.Ile442Met Variant Is Associated with an Increase in Body Frame Size at Puberty in Cattle. *Anim. Genet.* **2011**, *42*, 650–655, doi:10.1111/j.1365-2052.2011.02196.x.

34. Al-Mamun, H.A.; Kwan, P.; Clark, S.A.; Ferdosi, M.H.; Tellam, R.; Gondro, C. Genome-Wide Association Study of Body Weight in Australian Merino Sheep Reveals an Orthologous Region on OAR6 to Human and Bovine Genomic Regions Affecting Height and Weight. *Genet. Sel. Evol.* **2015**, *47*, 66, doi:10.1186/s12711-015-0142-4.

35. Lee, K.W.; Jeong, J.-Y.; An, Y.J.; Lee, J.-H.; Yim, H.-S. FGF11 Influences 3T3-L1 Preadipocyte Differentiation by Modulating the Expression of PPARγ Regulators. *FEBS Open Bio* **2019**, *9*, 769–780, doi:10.1002/2211-5463.12619.

36. Tontonoz, P.; Hu, E.; Graves, R.A.; Budavari, A.I.; Spiegelman, B.M. mPPAR Gamma 2: Tissue-Specific Regulator of an Adipocyte Enhancer. *Genes Dev.* **1994**, *8*, 1224–1234, doi:10.1101/gad.8.10.1224.

37. Gu, H.; Zhou, Y.; Yang, J.; Li, J.; Peng, Y.; Zhang, X.; Miao, Y.; Jiang, W.; Bu, G.; Hou, L.; et al. Targeted Overexpression of PPARγ in Skeletal Muscle by Random Insertion and CRISPR/Cas9 Transgenic Pig Cloning Enhances Oxidative Fiber Formation and Intramuscular Fat Deposition. *FASEB J.* **2021**, *35*, e21308, doi:10.1096/fj.202001812RR.

38. Zhu, Z.; Zhang, M.; Yuan, L.; Xu, Y.; Zhou, H.; Lian, Z.; Liu, P.; Li, X. LGP2 Promotes Type I Interferon Production To Inhibit PRRSV Infection via Enhancing MDA5-Mediated Signaling. *J. Virol. 97*, e01843-22, doi:10.1128/jvi.01843-22.

39. Dar, A.C.; Dever, T.E.; Sicheri, F. Higher-Order Substrate Recognition of eIF2α by the RNA-Dependent Protein Kinase PKR. *Cell* **2005**, *122*, 887–900, doi:10.1016/j.cell.2005.06.044.

40. Minks, M.A.; West, D.K.; Benvin, S.; Baglioni, C. Structural Requirements of Double-Stranded RNA for the Activation of 2',5'-Oligo(A) Polymerase and Protein Kinase of Interferon-Treated HeLa Cells. *J. Biol. Chem.* **1979**, *254*, 10180–10183, doi:10.1016/S0021-9258(19)86690-5.

41. Rezaei, R.; Wang, W.; Wu, Z.; Dai, Z.; Wang, J.; Wu, G. Biochemical and Physiological Bases for Utilization of Dietary Amino Acids by Young Pigs. *J. Anim. Sci. Biotechnol.* **2013**, *4*, 7, doi:10.1186/2049-1891-4-7.

42. Suryawan, A.; Jeyapalan, A.S.; Orellana, R.A.; Wilson, F.A.; Nguyen, H.V.; Davis, T.A. Leucine Stimulates Protein Synthesis in Skeletal Muscle of Neonatal Pigs by Enhancing mTORC1 Activation. *Am. J. Physiol. - Endocrinol. Metab.* **2008**, *295*, E868–E875, doi:10.1152/ajpendo.90314.2008.

43. Li, H.; Xu, M.; Lee, J.; He, C.; Xie, Z. Leucine Supplementation Increases SIRT1 Expression and Prevents Mitochondrial Dysfunction and Metabolic Disorders in High-Fat Diet-Induced Obese Mice. *Am. J. Physiol. - Endocrinol. Metab.* **2012**, *303*, E1234–E1244, doi:10.1152/ajpendo.00198.2012.

44. Yuan, X.-W.; Han, S.-F.; Zhang, J.-W.; Xu, J.-Y.; Qin, L.-Q. Leucine Supplementation Improves Leptin Sensitivity in High-Fat Diet Fed Rats. *Food Nutr. Res.* **2015**, *59*, doi:10.3402/fnr.v59.27373.

45. Kamarul, T.; Abdul Rahim, S.; Tumin, M.; Selvaratnam, L.; Ahmad, T.S. A Preliminary Study of the Effects of Glucosamine Sulphate and Chondroitin Sulphate on Surgically Treated and Untreated Focal Cartilage Damage. *Eur. Cells Mater. J.* **2011**, *21*, 259–271.

46. Martins, J.M.S.; dos Santos Neto, L.D.; Sgavioli, S.; Araújo, I.C.S.; Reis, A.A.S.; Santos, R.S.; de Araújo, E.G.; Leandro, N.S.M.; Café, M.B. Effect of Glycosaminoglycans on the Structure and Composition of Articular Cartilage and Bone of Broilers. *Poult. Sci.* **2023**, *102*, 102916, doi:10.1016/j.psj.2023.102916.

47. Pearen, M.A.; Muscat, G.E.O. Minireview: Nuclear Hormone Receptor 4A Signaling: Implications for Metabolic Disease. *Mol. Endocrinol.* **2010**, *24*, 1891–1903, doi:10.1210/me.2010-0015.

48. Herring, J.A.; Elison, W.S.; Tessem, J.S. Function of Nr4a Orphan Nuclear Receptors in Proliferation, Apoptosis and Fuel Utilization Across Tissues. *Cells* **2019**, *8*, 1373, doi:10.3390/cells8111373.

49. Maira, M.; Martens, C.; Philips, A.; Drouin, J. Heterodimerization between Members of the Nur Subfamily of Orphan Nuclear Receptors as a Novel Mechanism for Gene Activation. *Mol. Cell. Biol.* **1999**, *19*, 7549–7557.

50. Li, R.; Chen, S.; Gu, X.; An, S.; Wang, Z. Role of the Nuclear Receptor Subfamily 4a in Mast Cells in the Development of Irritable Bowel Syndrome. *Comput. Struct. Biotechnol. J.* **2022**, *20*, 1198–1207, doi:10.1016/j.csbj.2022.02.017.

51. Danilova, O.; Li, B.; Szardenings, A.K.; Huber, B.T.; Rosenblum, J.S. Synthesis and Activity of a Potent, Specific Azabicyclo[3.3.0]-Octane-Based DPP II Inhibitor. *Bioorg. Med. Chem. Lett.* **2007**, *17*, 507–510, doi:10.1016/j.bmcl.2006.10.012.

52. Imai, T.; Hieshima, K.; Haskell, C.; Baba, M.; Nagira, M.; Nishimura, M.; Kakizaki, M.; Takagi, S.; Nomiyama, H.; Schall, T.J.; et al. Identification and Molecular Characterization of Fractalkine Receptor CX3CR1, Which Mediates Both Leukocyte Migration and Adhesion. *Cell* **1997**, *91*, 521–530, doi:10.1016/S0092-8674(00)80438-9.

53. Ritchie, K.J.; Hahn, C.S.; Kim, K.I.; Yan, M.; Rosario, D.; Li, L.; de la Torre, J.C.; Zhang, D.-E. Role of ISG15 Protease UBP43 (USP18) in Innate Immunity to Viral Infection. *Nat. Med.* **2004**, *10*, 1374–1378, doi:10.1038/nm1133.

54. Gunn, M.D.; Ngo, V.N.; Ansel, K.M.; Ekland, E.H.; Cyster, J.G.; Williams, L.T. A B-Cell-Homing Chemokine Made in Lymphoid Follicles Activates Burkitt's Lymphoma Receptor-1. *Nature* **1998**, *391*, 799–803, doi:10.1038/35876.

55. Clarke, E.; Rahman, N.; Page, N.; Rolph, M.S.; Stewart, G.J.; Jones, G.J. Functional Characterization of the Atopy-Associated Gene *PHF11*. *J. Allergy Clin. Immunol.* **2008**, *121*, 1148-1154.e3, doi:10.1016/j.jaci.2008.02.028.

56. El-Diwany, R.; Soliman, M.; Sugawara, S.; Breitwieser, F.; Skaist, A.; Coggiano, C.; Sangal, N.; Chattergoon, M.; Bailey, J.R.; Siliciano, R.F.; et al. CMPK2 and BCL-G Are Associated with Type 1 Interferon–Induced HIV Restriction in Humans. *Sci. Adv.* **2018**, *4*, eaat0843, doi:10.1126/sciadv.aat0843.

57. Mukilan, M.; Rajathei, D.M.; Jeyaraj, E.; Kayalvizhi, N.; Rajan, K.E. MiR-132 Regulated Olfactory Bulb Proteins Linked to Olfactory Learning in Greater Short-Nosed Fruit Bat *Cynopterus Sphinx*. *Gene* **2018**, *671*, 10–20, doi:10.1016/j.gene.2018.05.107.

58. Viré, E.; Brenner, C.; Deplus, R.; Blanchon, L.; Fraga, M.; Didelot, C.; Morey, L.; Van Eynde, A.; Bernard, D.; Vanderwinden, J.-M.; et al. The Polycomb Group Protein EZH2 Directly Controls DNA Methylation. *Nature* **2006**, *439*, 871–874, doi:10.1038/nature04431.

59. Shen, M.; Cai, Y.; Yang, Y.; Yan, X.; Liu, X.; Zhou, T. Centrosomal Protein FOR20 Is Essential for S-Phase Progression by Recruiting Plk1 to Centrosomes. *Cell Res.* **2013**, *23*, 1284–1295, doi:10.1038/cr.2013.127.

60. Fox, J.W.; Mayer, U.; Nischt, R.; Aumailley, M.; Reinhardt, D.; Wiedemann, H.; Mann, K.; Timpl, R.; Krieg, T.; Engel, J. Recombinant Nidogen Consists of Three Globular Domains and Mediates Binding of Laminin to Collagen Type IV. *EMBO J.* **1991**, *10*, 3137–3146, doi:10.1002/j.1460-2075.1991.tb04875.x.

61. Katsuoka, F.; Motohashi, H.; Ishii, T.; Aburatani, H.; Engel, J.D.; Yamamoto, M. Genetic Evidence That Small Maf Proteins Are Essential for the Activation of Antioxidant Response Element-Dependent Genes. *Mol. Cell. Biol.* **2005**, *25*, 8044–8051, doi:10.1128/MCB.25.18.8044-8051.2005.

62. Arndt, V.; Dick, N.; Tawo, R.; Dreiseidler, M.; Wenzel, D.; Hesse, M.; Fürst, D.O.; Saftig, P.; Saint, R.; Fleischmann, B.K.; et al. Chaperone-Assisted Selective Autophagy Is Essential for Muscle Maintenance. *Curr. Biol.* **2010**, *20*, 143–148, doi:10.1016/j.cub.2009.11.022.

63. Carra, S.; Seguin, S.J.; Lambert, H.; Landry, J. HspB8 Chaperone Activity toward Poly(Q)-Containing Proteins Depends on Its Association with Bag3, a Stimulator of Macroautophagy *. *J. Biol. Chem.* **2008**, *283*, 1437–1444, doi:10.1074/jbc.M706304200.

64. Elsafadi, M.; Manikandan, M.; Dawud, R.A.; Alajez, N.M.; Hamam, R.; Alfayez, M.; Kassem, M.; Aldahmash, A.; Mahmood, A. Transgelin Is a TGFβ-Inducible Gene That Regulates Osteoblastic and Adipogenic Differentiation of Human Skeletal Stem Cells through Actin Cytoskeleston Organization. *Cell Death Dis.* **2016**, *7*, e2321–e2321, doi:10.1038/cddis.2016.196.

65. Kirchmaier, B.C.; Poon, K.L.; Schwerte, T.; Huisken, J.; Winkler, C.; Jungblut, B.; Stainier, D.Y.; Brand, T. The Popeye Domain Containing 2 (*Popdc2*) Gene in Zebrafish Is Required for Heart and Skeletal Muscle Development. *Dev. Biol.* **2012**, *363*, 438–450, doi:10.1016/j.ydbio.2012.01.015.

66. Chen, C.; Westenbroek, R.E.; Xu, X.; Edwards, C.A.; Sorenson, D.R.; Chen, Y.; McEwen, D.P.; O'Malley, H.A.; Bharucha, V.; Meadows, L.S.; et al. Mice Lacking Sodium Channel B1 Subunits Display Defects in Neuronal Excitability, Sodium Channel Expression, and Nodal Architecture. *J. Neurosci.* **2004**, *24*, 4030–4042, doi:10.1523/JNEUROSCI.4139-03.2004.

67. Kulkarni, S.S.; Karlsson, H.K.R.; Szekeres, F.; Chibalin, A.V.; Krook, A.; Zierath, J.R. Suppression of 5′-Nucleotidase Enzymes Promotes AMP-Activated Protein Kinase (AMPK) Phosphorylation and Metabolism in Human and Mouse Skeletal Muscle*. *J. Biol. Chem.* **2011**, *286*, 34567–34574, doi:10.1074/jbc.M111.268292.