Biochemical and neurophysiological effects of deficiency of the mitochondrial import protein TIMM50

Eyal Paz[1,2, ¶], Sahil Jain[1, ¶], Irit Gottfried[1], Orna Staretz-Chacham[3], Muhammad Mahajnah[4,5], Pritha Bagchi[6,7,8], Nicholas T. Seyfried[6,7], Uri Ashery[1,2,*], Abdussalam Azem[1,2,*]

[1]School of Neurobiology, Biochemistry and Biophysics, George S. Wise Faculty of Life Sciences, Tel Aviv University, Tel Aviv, Israel

[2]Sagol School of Neuroscience, Tel Aviv University, Tel Aviv, Israel

[3]Metabolic Disease Unit, Soroka Medical Center, Faculty of Health Sciences, Ben-Gurion University of the Negev, Beer Sheba, Israel

[4]The Ruth and Bruce Rappaport Faculty of Medicine, Technion-Israel Institute of Technology, Haifa, Israel

[5]Child Neurology and Development Center, Hillel Yaffe Medical Center, Hadera, Israel

[6]Center for Neurodegenerative Diseases, Emory University, Atlanta, GA, USA

[7]Department of Biochemistry, Emory University, Atlanta, GA, USA

[8]Emory Integrated Proteomics Core, Emory University, Atlanta, GA, USA

*Corresponding Authors

E-mail: azema@tauex.tau.ac.il (AA)

E-mail: uria@tauex.tau.ac.il (UA)

[¶]- These authors contributed equally to this work.

Abstract

TIMM50, an essential TIM23 complex subunit, is suggested to facilitate the import of ~60% of the mitochondrial proteome. In this study, we characterized a *TIMM50* disease causing mutation in human fibroblasts, and noted significant decreases in TIM23 core protein levels (TIMM50, TIMM17A/B, and TIMM23). Strikingly, TIMM50 deficiency had no impact on the steady state levels of most of its substrates, challenging the currently accepted import dogma of the essential general import role of TIM23 and suggesting that fully functioning TIM23 complex is not essential for maintaining the steady state level of the majority of mitochondrial proteins. As TIMM50 mutations have been linked to severe neurological phenotypes, we aimed to characterize TIMM50 defects in manipulated mammalian neurons. TIMM50 knockdown in mouse neurons had a minor effect on the steady state level of most of the mitochondrial proteome, supporting the results observed in patient fibroblasts. Amongst the few affected TIM23 substrates, a decrease in the steady state level of components of the intricate oxidative phosphorylation and mitochondrial ribosome complexes was evident. This led to declined respiration rates in fibroblasts and neurons, reduced cellular ATP levels and defective mitochondrial trafficking in neuronal processes, possibly contributing to the developmental defects observed in patients with TIMM50 disease. Finally, increased electrical activity was observed in TIMM50 deficient mice neuronal cells, which correlated with reduced levels of KCNJ10 and KCNA2 plasma membrane potassium channels, likely underlying the patients' epileptic phenotype.

Keywords: Mitochondria / mitochondrial protein import / potassium channels / TIM23 complex / TIMM50

Introduction

The mitochondrion is a vital organelle found in nearly all eukaryotic cells, where it is involved in numerous important cellular functions and metabolic pathways, including supplying cellular energy, assembling iron-sulphur clusters, and regulating the cell cycle, cell growth and differentiation, programmed cell death, and synaptic transmission (1–3). In humans, these functions are executed by ~1,500 different mitochondrial proteins, of which only 13 are encoded by the mitochondrial genome (4). The remaining mitochondrial proteins are nuclear-encoded and thus imported into the mitochondria. The translocated proteins are sorted to their specific mitochondrial compartment via several intricate protein translocation pathways, including the presequence pathway, which is used for import of nearly ~60% of the mitochondrial proteins (5,6).

The TIM23 complex mediates the import of some intermembrane space (IMS) proteins, many mitochondrial inner membrane (MIM) proteins, and all mitochondrial matrix proteins (7). The TIM23 complex in yeast comprises three essential subunits, Tim23, Tim17 and Tim50 (TIMM23, TIMM17A/B and TIMM50 in mammals). Association of Tim21 (TIMM21 in mammals) and Mgr2 (ROMO1 in mammals) promotes the lateral translocation of proteins into the MIM, while association of the presequence translocase-associated motor (PAM) complex with the TIM23 core promotes the import of matrix proteins (8–10). Recent structural analysis showed that Tim17 forms the protein translocation path, whereas the associated Tim23 protein likely plays a structural role, serving as a platform that mediates the association of other complex subunits (11,12).

Tim50, first discovered in yeast some two decades ago (13,14), is thought to be the first TIM23 complex component to interact with presequences of precursor proteins as they emerge from the Tom40 channel, thus playing a pivotal role in presequence-containing protein sorting (15). It was further suggested that normal Tim50 functionality is required for maintaining the mitochondrial membrane potential (16). Additionally, TIMM50, the mammalian homologue of Tim50, was shown to be involved in steroidogenesis and play a preventive role in pathological cardiac hypertrophy and several types of cancer (17–21).

75 Recently, TIMM50 has generated immense interest in human health research, as mutations in the
76 encoding gene have been linked in geographically and ethnically varied populations to the
77 development of a severe disease characterized by mitochondrial epileptic encephalopathy,
78 developmental delay, optic atrophy, cardiomyopathy, and 3-methylglutaconic aciduria. To date,
79 seven different mutations have been identified in children from ten unrelated families (22–27).
80
Although TIMM50 mutants are mostly associated with neurological disorders, functional
characterization of TIMM50 has yet to be reported in neuronal cells. Additionally, despite being
involved in the import of nearly 60% of the mitochondrial proteins, the impact of TIMM50
deficiency on the entire mitochondrial proteome has yet to be characterized. In this study, using a
proteomics approach, we show the impact of TIMM50 deficiency on the mitochondrial and
cellular proteome and characterize for the first time the neurological role of TIMM50 by
knocking it down in mouse neurons. Surprisingly, functional and physiological analysis points to
the possibility that TIMM50 and a fully functional TIM23 complex are not essential for
maintaining steady-state levels of most presequence-containing proteins, yet does affect some
critical mitochondrial properties and neuronal activity.
91

Results

TIMM50 disease-causing mutation reduces the levels of TIM23 complex components in patient fibroblasts

A previous study reported the clinical features of a rare genetic disease in several patients from two unrelated Arab families residing in Israel and Palestine (23). Patients suffering from the disease exhibited various neurological and metabolic disorders, including epilepsy, developmental delay, optic atrophy, cardiomyopathy, and 3-methylglutaconic aciduria. The disease was linked to a homozygous missense mutation resulting in T252M replacement in the *TIMM50* gene. To unravel the molecular basis of the disease, we examined the effect of the mutation on the proteome of primary fibroblasts collected from two affected family members (P1 and P2), as well as from a healthy relative (HC).

We first examined the effect of the mutation on TIMM50 levels by immunoblotting and found that the mutation leads to a significant decrease in TIMM50 levels in patient fibroblasts (Fig 1A and B). We next examined the effect of the *TIMM50* mutation on other components of the TIM23 complex. Notably, our analysis revealed a significant reduction in the levels of the core TIM23 complex subunits, namely, TIMM23, TIMM17A and TIMM17B (Fig 1A and B). These results are in agreement with previous reports showing that two other *TIMM50* mutations (resulting in R217Q+G372S and S112*+G190A replacements, both compound heterozygous) also led to major decreases in TIMM50, TIMM23 and TIMM17A/B levels (24,25).

In contrast to the decreased levels of TIM23 complex core components seen in the patient fibroblasts, the levels of subunits belonging to the PAM complex were not affected, despite their expected import dependency on the TIM23 complex. TIMM21 also showed minimal to no change in amount. Additionally, as subunits of the TOM complex that serves as the general import pore in the outer membrane were shown to interact with TIM23 complex subunits, including TIMM50 (28), we also considered the possible effect of reduced TIMM50 levels on the expression of TOM complex subunits. No changes in the levels of TOM complex subunits addressed in patient fibroblasts were noted (Fig 1A and B).

122   Finally, as TIM23 is known to be the sole gateway into the mitochondrial matrix (7), we
123   examined the effect of *TIMM50* mutation on the steady state levels of the matrix proteins
124   aconitase 2 and mtHsp60. Surprisingly, the steady state levels of both proteins were unchanged
125   in patient fibroblasts (Fig 1A and B), despite the significant loss of TIMM50, TIMM23, and
126   TIMM17A/B.

Generating a neuronal model system to study the neurophysiological effects of TIMM50 deficiency

As *TIMM50* mutations lead to severe neurodevelopmental symptoms and to a significant
reduction in steady state TIMM50 levels (Fig 1 and (24,25)) we wanted to study the effects of
TIMM50 deficiency in neuronal cells by knocking down TIMM50 in mouse primary cortical
neurons. For this purpose, we designed three shRNA sequences (termed Sh1, Sh2 and Sh3;
vector schemes are presented in Fig 2A) and cloned them into a lentiviral vector that also allows
for EGFP expression under control of the hSyn promotor (29). These vectors allow TIMM50
knockdown (KD) while specifically labeling neurons with EGFP, allowing for efficient
visualization in single cell experiments. We compared TIMM50 expression relative to three
controls, namely, an untreated control (i.e., cultures that were not transduced), a pLL3.7 control
(i.e., cultures that were transduced but did not transcribe the shRNA sequence, yet expressed
EGFP), and a shRNA system activation control (i.e., cultures that were transduced with a
scrambled shRNA sequence). All the three targeting shRNA sequences had an impact on
TIMM50 levels, in comparison to the controls. Sh2 had the most significant and consistent
effect, reducing TIMM50 levels by ~80% (Fig 2B and C). Therefore, Sh2 was chosen as the KD
vector for all subsequent experiments on neurons.

We next examined the effects of TIMM50 KD in neurons, addressing the same components as
were tested in fibroblasts. In complete agreement with the fibroblast results, the levels of
TIMM23 and TIMM17A/B were significantly decreased, while the amounts of TIMM21, PAM
subunits, TOM subunits and the matrix substrates tested were not affected (Fig 2B and C).
Hence, similarly to what was detected in fibroblasts, steady state levels of many mitochondrial
proteins were not affected in TIMM50 KD neurons.

TIMM50 deficiency does not affect the steady state levels of a majority of its substrates

It was expected that the significant decrease in the levels of TIM23 core components seen upon TIMM50 deficiency would decrease the levels of substrates processed by this translocation complex. The observation that steady state levels of two mitochondrial matrix substrates, mtHsp60 and aconitase 2, were not affected by the significant loss of TIM23 core components in both research systems (Fig 1A and B and Fig 2B and C), motivated us to examine the impact of TIMM50 KD on the total mitochondrial and cellular proteomes. For this purpose, we performed untargeted mass spectrometry analysis of fibroblasts from both patients, as compared to the healthy control, and of mice primary neurons transduced with either Sh2 or the Scr control.

In the case of fibroblasts, 127 MIM and 190 matrix proteins were detected using mass spectrometry (Supplemental material S1 and S2 Datasets). Surprisingly, we noticed that the levels of 83 (~65%) MIM proteins and 135 (~71%) matrix proteins were not affected in either patient, as compared to the HC (Fig 3A and B, left and middle panels and supplemental material S1 and S2 Datasets). Among the MIM proteins that were not affected by the *TIMM50* mutation, we identified multiple proteins involved in calcium homeostasis (such as MICU2, SLC25A3 and LETM1), heme synthesis (such as PPOX and CPOX), and cardiolipin synthesis (HADHA). Among matrix proteins that were not affected by *TIMM50* mutation, we identified multiple proteins involved in Fe-S cluster biosynthesis (such as NFS1, GLRX5 and ISCU), detoxification (such as PRDX5, SOD2, ABHD10 and GSTK1), fatty acid oxidation (such as DECR1, ECHS1 and ETFA), and amino acid metabolism (PYCR1, ALDH18A1 and HIBCH). Also, the majority of TCA cycle proteins, such as ACO2, DLST, IDH3B and OGDH, found in the matrix, were not affected in patient fibroblasts. Unexpectedly, a few matrix proteins (ALDH2, GRSF-1, AK4, LACTB2 and OAT) showed increased steady state levels in both patients (Fig 3A and B, left and middle panels and supplemental material S1 and S2 Datasets). Of these proteins, ALDH2 and GRSF-1 showed 15 and 6-fold increases, respectively, as was also confirmed by immunoblot (S1A Fig).

Oxidative phosphorylation subunits and mitochondrial ribosomal proteins comprise the majority of down-regulated proteins in TIMM50-deficient fibroblasts

A total of 69 oxidative phosphorylation (OXPHOS) and 27 mitochondrial ribosomal proteins (MRP) were detected in fibroblasts by mass spectrometry. Remarkably, of the 18 MIM proteins found in decreased amounts in both patients, 17 belong to the OXPHOS family, while of the 7 matrix proteins found in decreased amounts in both patients, 4 proteins belong to the MRP family (Fig 4A and B, left and middle panels). The OXPHOS complexes most affected were CI, CII and CIV. In the case of CI, one membrane core subunit (MT-ND1), two hydrophilic core subunits (NDUFS1 and NDUFV2), and four super-numerary subunits (NDUFA2, NDUFA10, NDUFB3, and NDUFS5) exhibited a significant negative fold-change in both patients (supplemental material S1 and S2 Datasets). The levels of two CII subunits, namely, SDHA and SDHB, and eight CIV subunits, including the two catalytic subunits MT-CO1 and MT-CO2, were also significantly decreased in both patients. The MT-subunits are encoded and translated by mitochondria, and as such, the decrease in their levels cannot be directly related to TIM23 deficiency. Still, the decreased levels of MRP subunits could be responsible for the decrease in mitochondrially translated OXPHOS subunits in patient fibroblasts.

The impact of TIMM50 deficiency on the mitochondrial proteome in neurons

In TIMM50 KD neurons, 170 MIM and 215 matrix proteins were detected by mass spectrometry (supplemental material S3 Dataset). We observed a lesser decrease in the levels of multiple OXPHOS and MRP proteins in TIMM50 KD neurons, as compared to the levels seen in patient fibroblasts (Fig 4A and B, right panels). Yet, the trends seen for these proteins were similar to those observed in patient fibroblasts, with most of the OXPHOS and MRP proteins being detected at lower levels in the TIMM50 KD neuronal cells than in the corresponding control. The difference in the extent of fold-change in neurons, as compared to patient fibroblasts, could be due to the short duration of KD in the neuronal experiment, as compared to that with the patient fibroblasts, that constantly carry the deficiency.

Additionally, to determine whether the changes observed in the levels of OXPHOS and MRP proteins were an indirect effect resulting from general mitochondrial DNA loss, we performed qPCR assessment of the mitochondrial DNA content. Our results showed no effect on mitochondrial DNA levels in either model system (Fig 4C and D), thereby confirming that the observed effects on the OXPHOS and MRP protein levels were the direct and specific result of TIMM50 deficiency, and not indirectly due to mitochondrial DNA loss.

Overall, we conclude that translocation of the majority of MIM and matrix proteins was not affected in both patient fibroblasts and TIMM50 KD mice neurons, even after significant disruption of TIMM50 and the TIM23 core subunits. However, TIMM50 deficiency severely affected two major mitochondria complex systems, namely, the OXPHOS and MRP protein machineries.

TIMM50 deficiency affects ATP production

The observed decrease in steady state levels of OXPHOS subunits led us to examine oxygen consumption in TIMM50-deficient cells. For this purpose, we performed the Seahorse XF cell Mito Stress test (Fig 5A). Comparing oxygen consumption rates in the HC, P1 and P2 fibroblasts, and in the Sh2- and Scr control-transduced neuronal cells revealed significant impairment of both basal and maximal respiration rates and of OXPHOS-dependent energy production rate in the non-control cells (Fig 5C and D). Overall, these results suggest that TIMM50 deficiency severely affects the OXPHOS and MRP protein machineries and leads to OXPHOS-dependent ATP deficiency in both systems.

Moreover, as cells are able to meet their energetic requirements via glycolysis when the OXPHOS apparatus malfunctions (30), we measured the glycolytic capabilities of TIMM50-deficient cells by performing a Seahorse XF glycolysis stress test (Fig 5B). In both systems, the basal glycolysis level remained stable, in comparison to controls, suggesting that the cells did not make the metabolic switch so as to increasingly rely on the non-mitochondrial energy production pathway that is glycolysis (Fig 5E and F, left panel). Moreover, both glycolytic capacity and glycolytic reserves were significantly reduced, indicative of an impaired ability of both patient

241 fibroblasts and neurons to switch their energetic emphasis to glycolysis when needed (Fig 5E and

242 F, middle and right panels).

TIMM50 KD impairs mitochondrial trafficking in neuronal cells

The transport of mitochondria within neuronal processes is crucial for cell survival (31–33).

Therefore, we investigated the effect of TIMM50 deficiency on mitochondrial trafficking in

neuronal cells. To track individual mitochondria in neuronal processes, we used a transfection

method, instead of transduction, that results in low expression efficiency in neurons (34). This

allowed us to visualize individual processes and track single mitochondria with minimal

background noise. To visualize neuronal cells and their mitochondria, we co-transfected our

cultures with a dsRed-mito-encoding plasmid, together with the KD or control plasmids. We

then performed live cell imaging of individual neuronal processes and tracked the movement of

individual mitochondria in these structures (for examples, see Fig 6A and S1 Movie). The live

imaging sets were converted into kymographs and calibrated in time and space, which allowed

extraction of different trafficking parameters, such as distance of movement, speed, and

percentage of moving mitochondria (Fig 6B).

Our results showed a two-fold decrease in the percentage of mobile mitochondria in TIMM50-

deficient neuronal cells, as compared to control cells (Fig 6C). Moreover, mobile mitochondria

in TIMM50-deficient neuronal cells tend to cover less distance and travel at a lower average

travelling speed (Fig 6D and E). This indicates that TIMM50 deficiency causes neuronal cell

mitochondria to be more static, which could consequently lead to further energy deprivation in

regions where mitochondria are needed but cannot be shipped.

Mitochondrial movement along neuronal processes is coordinated by a motor/adaptor complex.

The motors kinesin and dynein use ATP to move organelles along microtubules. Mitochondria

are assembled onto these motors via a mitochondrial outer membrane protein called Miro, a

cytosolic adaptor called Milton (also known as TRAK1/2), and a few cytosolic accessory

proteins (35). Although TIM23 and TIMM50 are not directly involved in the biogenesis of any

of these proteins, we, nonetheless, examined their expression levels following TIMM50 KD. Our

proteomics data revealed no major changes in the levels of proteins involved in mitochondrial

trafficking (Fig 6F), suggesting that the observed effect on neuronal cell mitochondrial
trafficking to be most likely indirect and resulting from the ATP deficiency. To examine this
possibility, we measured cellular ATP levels in Scr control-transduced and TIMM50 KD
neuronal cells. As expected from the impaired ATP production (Fig. 5), we found that TIMM50
KD led to a significant reduction of about 25% in cellular ATP levels (Fig 6G).

TIMM50 KD leads to excess neuronal activity and increased action
potential frequency

All TIMM50 mutant patients studied thus far displayed severe neurological pathologies, which
include epilepsy, developmental delay, and loss of movement abilities. Such abnormalities can
be attributed to alternations in basic neuronal function. To assess whether such functions are
altered upon TIMM50 KD in neuronal cells, we measured intrinsic neuronal excitability, as well
as spontaneous neurotransmitter release, using the whole cell patch clamp technique.

Initially, we measured spontaneous excitatory activity of the cells in the presence of tetrodotoxin
(TTX) (representative traces are presented in S2A Fig). We quantified the average miniature
excitatory post-synaptic current (mEPSC) amplitude, area and frequency in each of the measured
neuronal cells and found no significant differences between any of these measures in KD cells,
as compared to controls (S2B Fig). Moreover, the relative frequency distribution of the
amplitude measurements showed an even distribution pattern (S2C Fig). Examination of the
cumulative distribution function confirmed that no significant differences in mEPSC amplitude
distribution exists in the neuronal cells (S2D Fig).

We subsequently measured the minimal current required to induce an action potential by slowly
increasing the stimulation current in a stepwise manner (Fig 7A). This allowed us to estimate the
rheobase of the neuronal TIMM50 KD and control cells. We found that there were no significant
differences in the current needed to trigger an action potential between the TIMM50 KD and Scr
control-transduced cells (Fig 7D). We also assessed characteristics of the first observed action
potential in each measurement. Both the half-width and rate of fall of the first action potential
were similar in TIMM50 KD and Scr control-transduced neurons (Fig 7B, E and F). However,

TIMM50 KD cells displayed shorter action potential latency (Fig 7G) and a significant increase

in the maximum number of action potentials as compared to the Scr control-transduced cells (Fig

7C and H). Overall, these results suggest that TIMM50 KD causes the cells to fire more action

potentials without decreasing the firing threshold, probably due to a faster recovery time between

successive action potentials.

An increase in the firing frequencies of action potentials can be explained by a reduced presence

of voltage-dependent potassium channels, which are known to control spike frequency (36).

Indeed, our proteomics results confirmed a decrease of about 2.5-fold in the levels of the

KCNA2 and KCNJ10 potassium channels in the TIMM50-deficient neuronal cells, supporting

the increase in their action potential frequency (Fig 7I). KCNA2 levels were also tested via

immunoblot and confirmed to be dramatically decreased (Fig S1B). To further test how KCNA2

reduction impacts cellular electrical activity, we used α-dendrotoxin (α-DTX), a known KCNA2

channel blocker (37,38), to mimic a reduction in KCNA2. The number of action potentials fired

was measured before and after application of 100 nM α-DTX to Scr control-transduced and

TIMM50 KD neuronal cells (Fig S2E). As expected, the difference in the number of action

potentials after and before α-DTX treatment increased significantly more in Scr control-

transduced cells than in TIMM50 KD cells (Fig S2F), given how TIMM50 KD cells initially

contain less KCNA2 channels than do the corresponding controls. These data indicate that a

reduction in KCNA2 contributes to the observed increase in firing rate in TIMM50 KD neurons.

Discussion

The TIMM50 protein is a pivotal member of the TIM23 complex that is suggested to participate in the import of nearly 60% of the mitochondrial proteome (5,6). Human *TIMM50* mutations lead to neurological effects, including mitochondrial epileptic encephalopathy, intellectual disability, seizure disorders like infantile spasms, and severe hypotonia accompanied by 3-methylglutaconic aciduria (22–27). Despite being involved in the import of the majority of the mitochondrial proteome, no study thus far characterized the effects of TIMM50 deficiency on the entire mitochondrial proteome. Additionally, despite the fact that human TIMM50 mutations lead mainly to neurological symptoms, no study has yet addressed the effects of TIMM50 deficiency in brain cells. Therefore, in this study, we utilized two research models – TIMM50 mutant patient fibroblasts and TIMM50 KD primary mouse neuronal cultures, to study the impact of TIMM50 deficiency on the mitochondrial proteome and its impact on neurophysiology.

Interestingly, in both model systems, TIMM50 deficiency reduced the levels of TIM23 core subunits, yet did not alter steady state levels of a majority of TIM23 substrates. This is surprising, as TIM23 is thought to be indispensable for the translocation of presequence-containing mitochondrial proteins (14,15,39–41). Even more surprising was that the amounts of some TIM23 substrates related to intricate metabolic and maintenance activities (e.g., ALDH2, GRSF-1, OAT, etc.) were increased. Moreover, of those TIM23 substrates that decreased in amount, a majority belong to the OXPHOS and MRP machineries. These observations can be explained by several plausible mechanisms: it is possible that unlike what occurs in yeast, fully functional mammalian TIMM50 and TIM23 complex are mainly essential for maintaining the steady state levels of intricate complexes/assemblies. Another explanation for this scenario is that the normal quantities of unaffected matrix proteins might be low, and hence, even ~10-20% of functional TIMM50 protein might be sufficient to maintain their steady state levels. Alternatively, the presequence of such proteins might contain mitochondrial targeting signals (MTSs) that receive priority over other presequence-containing precursor proteins, thus enabling their translocation even in the presence of very few functional TIM23 complexes. However, further experiments examining these possibilities are needed for understanding the compromised TIM23-mediated protein import.

As stated earlier, loss of TIMM50 led to a significant reduction in the steady state levels of other TIM23 core subunits (i.e., TIMM23 and TIMM17A/B). Notably, TIMM23 and TIMM17A/B are thought to be imported by the TIM22 complex (42). However, we observed that the steady state levels of TIM22 complex subunits had not been affected by TIMM50 deficiency (supplemental material S1-3 Datasets). This indicates a hitherto unknown relation between steady state levels of TIMM50 and other TIM23 core subunits, which might affect the complex assembly process. Various structural and biochemical studies have attempted to elucidate the intricate structure of the TIM23 complex and the complicated interactions between its different subunits (11,12,43), however, little is presently known of the dynamic assembly processes of the complex. The fact that TIMM50 KD leads to a major reduction in the levels of TIMM23 and TIMM17A/B, despite a lack of direct dependency of the import of these proteins on TIMM50, and does not affect the levels of other TIM23 complex subunits, could provide a basis for future studies examining the assembly process of import complexes.

Our proteomics data and Seahorse XF analysis, paired with cellular ATP measurements, indicated that lower ATP levels were present in TIMM50-deficient cells. Specifically in the case of neurons, such ATP deficiency could be responsible for the negative impact seen on mitochondrial trafficking in neuronal cell processes, which could contribute to the various neurodegenerative phenotypes linked to the TIMM50 disease. Additionally, the detected increase in action potential firing rate in TIMM50-deficient neurons can explain the presence of epileptic seizures, a hallmark of all TIMM50 patients studied thus far. A plausible explanation for the increased action potential firing rate could be the 2.5-fold reduction in the levels of the KCNA2 ($K_v1.2$) and KCNJ10 ($K_{ir}4.1$) voltage dependent potassium channels, as revealed by our proteomics and immunoblot analysis. The KCNA2 ($K_v1.2$) channel is a slowly inactivating channel that regulates neuronal excitability and firing rate (44). In peripheral sensory neurons, $K_v1.2$ helps to determine spike frequency, while in neurons of the medial trapezoid body, the Kv1.1, Kv1.2, and Kv1.6 subunits are important regulators of repetitive spiking (38,45). While the inactivation of several of the potassium channels like $K_v1.2$, $K_v1.1$ and Kcnj10 ($K_{ir}4.1$) are important for neuronal excitability, action potential width, and firing properties, in general, mutations in the genes encoding these channels that alter their inactivation are known to lead to

temporal lobe epilepsy (44,46,47). In addition, blocking the activation of the $K_v1.2$ by α-DTX

had no effect on the resting membrane potential and only small effects on the amplitude and

duration of the action potential (38), similar to what we observed in TIMM50 KD neurons, that

showed a reduction in $K_v1.2$ levels. Furthermore, blocking $K_v1.2$ activation by α-DTX increased

the frequency of action potentials in visceral sensory neurons (38), which, again, agrees with our

observation that α-DTX application increased the firing rate in Scr control-transduced neurons,

while hardly affecting the firing rate of TIMM50 KD neurons as they express lower levels of

$K_v1.2$. Hence, similar changes in the neuronal firing rate, as observed in our TIMM50 KD

neurons, might occur in TIMM50 patients, and could lead to the epileptic phenotype seen in

these patients. However, more studies are needed to verify this hypothesis.

In summary, our results challenge the main dogma that TIMM50 is essential for maintaining the

mitochondrial matrix and inner membrane proteome, as steady state level of most mitochondrial

matrix and inner membrane proteins did not change in either patient fibroblasts or mouse

neurons following a significant decrease in TIMM50 levels. Nevertheless, reductions in

TIMM50 levels led to a decrease of many OXPHOS and MRP complex subunits, which

indicates that TIMM50 might be essential only for maintaining the steady state levels and

assembly of intricate complex proteins. The consequently reduced cellular ATP levels and the

detected mitochondrial abnormalities in neurons provide a plausible link between the TIMM50

mutation and the observed developmental defects in the patients. Moreover, the increased

electrical activity resulting from decreased steady state levels of KCNA2 and KCNJ10 potassium

channels plausibly link the TIMM50 mutation to the epileptic phenotype of patients, thus,

providing a new direction for therapeutic efforts.

Materials and methods

Generation of primary human fibroblasts

Four mm punch biopsy samples were obtained from two TIMM50 patients (Patient 1 (P1) and Patient 2 (P2) carrying the mutation c.755C>T; p.Thr252Met and a normal family member that served as a healthy control (HC). Primary fibroblast cells were generated using standard procedures (48). In brief, each biopsy sample was cut into 12-15 pieces and 2-3 pieces were placed in six-well plate wells containing complete Dulbecco's modified Eagle's medium (DMEM; DMEM, 20% fetal bovine serum (FBS), 1% sodium pyruvate, 1% penicillin-streptomycin) and previously coated with 0.1% gelatin. The fibroblasts were grown for 2-3 weeks and passaged into 10 cm plates. Cells obtained from the first three passages were frozen and stored for further use. Following the generation of the primary fibroblast cells, genomic DNA purification and sequencing were performed to verify the presence of the mutation. Sequencing primers are found in S1 Table.

Generation of TIMM50 knockdown (KD) mice primary cortical neuronal cultures

Mouse primary cortical neurons were harvested from P_0/P_1 pups and cultured using a previously described procedure (49). Plates were pre-coated with Matrigel (Corning, 354234) (diluted 1:1000 in Hank's balanced salt solution (Satorius, 02-018-1A) with 10 mM HEPES, pH 7.4 (Fisher bioreagents, BP310-500)). For TIMM50 knockdown, three targeting shRNA sequences (Sh1, Sh2 and Sh3) and a scrambled (Scr) control sequence were designed. All shRNA sequences were cloned into the third-generation lentiviral vector pLL3.7 for expression under control of the U6 promotor. The same plasmid also encoded EGFP under control of the hSyn promotor, which allowed us to visualize and differentiate neuronal cells from other cells in the culture. To produce lentiviral particles, HEK293T/17 (ATCC, CRL-11268) cells were co-transfected with each of the designed shRNA vectors, together with the lentiviral helper constructs pMDLg-pRRE, pRSV-REV and CMV-VSVG, via calcium phosphate transfection. To generate TIMM50 KD neurons, the neuronal cultures were transduced on 4 days *in vitro* (DIV) with the generated lentiviruses and grown until 18 DIV. Immunoblotting with anti-TIMM50

438 antibodies were used to assess KD efficiency. shRNA oligo sequences and pLL3.7 sequencing
439 primers are found in S1 Table.

Immunoblotting

Fibroblasts were grown to ~90% confluency on 10 cm cell culture plates, harvested using
trypsin, washed twice with PBS, and then lysed using 50 μl of solubilization buffer (50 mM Na-
HEPES, pH 7.4, 150 mM NaCl, 1.5 mM $MgCl_2$, 10% glycerol, 1% Triton X-100, 1 mM EDTA,
1 mM EGTA supplemented with 400 μM of PMSF and 1:1000 dilution of protease inhibitor
cocktail (GenDEPOT, P3200-020). Neurons (~$1x10^6$ cells) were cultured in the wells of a six-
well plate, transduced on 4 DIV, grown until 18 DIV, and lysed by adding 50 μl of solubilization
buffer to each plate well, following by scraping with a cell scraper. The protein concentration of
both lyzed cultures was measured using Bradford reagent (BioRad, 500-0006) and appropriate
protein amounts (20-100 μg) were loaded onto homemade polyacrylamide gels (12/14/16%) and
separated using sodium dodecyl sulfate-polyacrylamide gel electrophoresis. The separated
proteins were then transferred to a PVDF membrane (Millipore, IPVH00010) and
immunodetection was carried out using antibodies against the target proteins. The list of
antibodies used can be found in S2 Table. For fibroblasts, actin or GAPDH served as a loading
control. For neurons, tubulin was used as loading control. ImageJ was used for densitometry
analysis of protein expression levels. At least three biological repeats were performed for each
immunoblotted protein (S1 and S2 Raw images).

Quantitative protein assessment and analysis

Label-free quantitative mass spectrometry was performed based on a published procedure (50).
Spectra were searched against the Uniprot/Swiss-Prot mouse database (17,041 target sequences)
for neuronal cells or the Uniprot/Swiss-Prot human database (20,379 target sequences) for
fibroblast cells using the Andromeda search engine integrated into MaxQuant. Methionine
oxidation (+15.9949 Da), asparagine and glutamine deamidation (+0.9840 Da), and protein N-
terminal acetylation (+42.0106 Da) were variable modifications (up to 5 allowed per peptide),
while cysteine was assigned a fixed carbamidomethyl modification (+57.0215 Da). Trypsin-
cleaved peptides with up to two missed cleavages were considered in the database search. A
precursor mass tolerance of ±20 ppm was applied prior to mass accuracy calibration and ±4.5

ppm after internal MaxQuant calibration. Other search settings included a maximum peptide

mass of 6,000 Da, a minimum peptide length of 6 residues, 0.05 Da tolerance for Orbitrap and

0.6 Da tolerance for ion trap MS/MS scans. The false discovery rates for peptide spectral

matches, proteins, and site decoy fractions were all set to 1 percent. Quantification settings were

as follows: Re-quantification in a second peak finding attempt after protein identification;

matched MS1 peaks between runs; a 0.7 min retention time match window after an alignment

function was found with a 20-minute RT search space. Protein quantitation was performed using

summed peptide intensities provided by MaxQuant. The quantitation method only considered

razor plus unique peptides for protein level quantitation.

Data were obtained from three biological repeats, each involving three technical repeats (i.e.,

nine samples in total) for every cell type (neurons: Sh2 and Scr control; fibroblasts: P1, P2 and

HC). Statistical analysis was performed using Perseus software. Protein levels were considered

to be increased or decreased in TIMM50-deficient cells if they were significantly different (p-

value < 0.05) and had a fold-change of at least 1.414, relative to what was measured in control

cells. Protein classification was performed manually by comparing the obtained data with the

MitoCarta3.0 human and mouse databases (51).

Seahorse XF mito-stress and glycolysis stress tests

Fibroblasts were plated in Seahorse XF 96-well plates (Agilent, 103775-100) at 20-30%

confluency, with experiments being carried out at ~90% confluency. For neurons, ~1×10^5 cells

were plated in each well of Seahorse XF 96-well plates, transduced with the Scr control or Sh2

constructs on 4 DIV and the experiment was carried out on 18 DIV. One-two hours prior to the

experiment, the fibroblasts or neuronal cultures were washed and the medium was replaced with

Seahorse XF DMEM, pH 7.4 (Agilent, 103575-100).

For the mito-stress test, the medium was supplemented with 1 mM sodium pyruvate (Sigma,

S8636-100ML), 10 mM glucose (Merck, 1.08337.1000) and 2 mM glutamine (Biological

Industries, 03-020-1B). The plates were loaded with oligomycin (Sigma-Aldrich, O4876),

carbonyl cyanide-p-trifluoromethoxyphenylhydrazone (FCCP, Sigma-Aldrich, C2920), and

rotenone (Sigma-Aldrich, R8875) together with antimycin A (Sigma-Aldrich, A8674), at final

well concentrations of 1, 2, 0.5 and 0.5 µM, respectively. For the glycolysis stress test, the

medium was only supplemented with glutamine. The plates were loaded with glucose,

oligomycin and 2-deoxy-D-glucose (2-DG, Sigma-Aldrich, D8375-1G)) at final well

concentrations of 10 mM, 1 µM and 50 mM, respectively. Plates were then loaded into the

Seahorse XFe96 Extracellular Flux Analyzer and the experiments were carried using the

manufacturer's protocol. To normalize the oxygen consumption rate (OCR) or extracellular

acidification rate (ECAR) values, fibroblasts were dyed with SynaptoGreen (Biotium, 70022)

immediately at the end of each experiment and fluorescence levels were measured using a plate

reader (BioTek Synergy HTX). In the case of neurons, the cells were dyed with DRAQ5

(BioLegend, 424101) immediately at the end of each experiment and imaged on an Incucyte SX5

live cell imaging and analysis system. Dyed nuclei were counted using the ImageJ particle

analysis function. Three to six biological repeats, each involving three to six technical repeats,

were performed for each experimental and control group.

Mitochondrial DNA content

Total DNA was isolated form near-confluent fibroblasts or 18 DIV transduced neuronal cultures

using a GeneElute Mammalian Genomic DNA Miniprep kit (Sigma, G1N350-1KT).

Quantitative real-time PCR (StepOnePlus Real-Time PCR System) was then performed on each

sample in the presence of SYBR green (PCR Biosystems, PB20.16-05). Expression levels were

determined using the comparative cycle threshold ($2^{-\Delta\Delta Ct}$) method, with the hypoxanthine

guanine phosphoribosyl transferase (HPRT)-encoding gene serving as housekeeping gene.

Primer sequences used are listed in S1 Table.

Mitochondrial trafficking

Neuronal cultures were plated in a similar manner as described for membrane potential

measurements. Neuronal cultures (4 DIV) were co-transfected with the TIMM50 KD or control

plasmids, as well as a mito-dsRed-expressing plasmid (52), using 0.6 µg of each plasmid and 0.6

µl of Lipofectamine 2000 (Invitrogen, 11668-027). A transfection rate of 2-5% was seen the next

day. This sparse transfection allowed for visualization of single neurons and their mitochondria,

given the dramatically reduced background that allowed for tracking of mitochondrial movement

in individual neurites. Following transfection, 10 DIV cultures were live-imaged with an iMIC

inverted microscope equipped with a Polychrome V system (TILL photonics) and an ANDOR

iXon DU 888D EMCCD camera (Andor, Belfast, Northern Ireland). Cells were imaged using a

60x oil immersion objective (Olympus), under temperature and CO_2 control. Fields of view

containing neurites stretching for at least 50 μm and not more than 100 μm from the cell body

were imaged for 5 minutes, with a 3 second interval between each image. Individual neurites in

each image set were selected using the segmented line tool in ImageJ. The images were then

calibrated in time and space and turned into kymographs using the KymoToolBox ImageJ plugin

(53). Each individual mitochondrion present in the neurite was then manually tracked on the

kymograph using the segmented line tool to extract various parameters. Mitochondria were

defined as static if they moved at a speed lower than 0.02 μm/sec.

Determination of cellular ATP levels

Neuronal cells were plated into a 96-well cell culture plate at a density of $5x10^4$ cells per well.

Cells were transduced with the Sh2 or Scr control constructs on 4 DIV and cellular ATP

measurements were performed on 18 DIV using a Luminescent ATP Detection Assay Kit

(Abcam, ab113849). To block luminescence signal contamination from adjacent wells, the lysis

step of the assay was performed in the clear cell culture plate and the lysates were transferred to

a white, flat bottom 96-well plate. Luminescence signals were read using the GloMax Navigator

System.

Intrinsic neuronal excitability and spontaneous activity

Neurons were plated at a density of $1.5x10^5$ cells/well of a 12-well plate and transduced on 4

DIV, with experiments being carried out on 16-20 DIV. Conventional whole cell recordings

were performed with borosilicate thin wall glass capillaries (World Precision Instruments,

TW150-3) with an input resistance of 4-5 MΩ. Series resistance ranged from 8-25 MΩ. An EPC-

9 patch clamp amplifier was used in conjunction with PatchMaster software (HEKA Electronik,

Lambrecht, Germany). The external solution consisted of 140 mM NaCl, 3 mM KCl, 2 mM

$CaCl_2$, 1 mM $MgCl_2$, 10 mM HEPES, supplemented with 2 mg/ml glucose, pH 7.4, osmolarity

adjusted to 305 mOsm. The internal solution consisted of 110 mM K gluconate, 10 mM KCl, 2

mM MgCl$_2$, 10 mM HEPES, 10 mM glucose, 10 mM Na creatine phosphate, and 10 mM EGTA,

pH 7.4. Osmolarity adjusted to 285 mOsm.

To measure the rheobase and assess repetitive and maximal firing, long (250 ms) depolarization

square current steps of varying intensity (from -100 to 600 pA, at 2 pA intervals) were applied.

Signals were filtered at 2 kHz and sampled at 5 kHz. For α-dendrotoxin (α-DTX) measurements,

long (250 ms) depolarizing square current steps of varying intensity (From 0-500 pA, at 100 pA

intervals) were applied to the cells, before and after application of an external solution containing

100 nM α-DTX (Alomone Labs, D-350). Clear traces in the range of 100-300 pA were chosen in

the before-α-DTX measurements and compared to the respective traces in the after-α-DTX

measurements. To measure spontaneous activity, an external solution containing 1 μM

tetrodotoxin (TTX; Alomone Labs, T550_1mg) was perfused onto the cells, followed by a high

current injection to assure that no action potentials were being generated. The mode was then

switched to voltage clamp, the holding voltage was adjusted to -60 mV and mEPSCs were

recorded for 2 minutes (at repeating 10 second intervals). The data were analyzed with Igor Pro

software (Wavemetrics, Lake Oswego, OR). The TaroTools procedure set (for Igor Pro) was

used to analyze spontaneous activity measurements.

Acknowledgements

The authors thank Professor Bernard Attali, Dr. Celeste Weiss Katz, Dr. Natalia Borovok and Dr. Amit Kessel for their valuable input and guidance. This work was supported by the Emory University Emory Integrated Proteomics Core Facility (RRID:SCR_023530). A.A. is incumbent of The Louise and Nahum Barag Chair in Molecular Genetics of Cancer. U.A. is the incumbent of the Michael Gluck Chair in Neuropharmacology and A.L.S Research.

Funding

Work in the Azem lab was supported by grants #1389/18 and #1057/22 from the Israel Science Foundation, and an Emory University and Tel Aviv University Collaborative Research Grant. This research was also supported by the Ministry of Innovation, Science & Technology, Israel (1001576154), Israel Science Foundation (ISF grant: 2141/20), BrightFocus grant (A2022029S), NIH grant 1R21AG074846-01A1, and the Michael J. Fox Foundation (MJFF-022407) (to U.A).

Conflict of interests

The authors declare that they have no conflict of interest.

Author contributions

Eyal Paz: Writing-original draft; data curation; formal analysis; investigation. **Sahil Jain:** Writing-original draft; data curation; formal analysis; investigation. **Irit Gottfried:** resources; writing-review and editing; validation; **Orna Staretz-Chacham**: Resources. **Muhammad Mahajnah**: Resources. **Pritha Bagchi:** Investigation; writing-review and editing. **Nicholas T. Seyfried:** Supervision; methodology. **Uri Ashery:** Conceptualization; supervision; funding acquisition; validation; methodology; writing-review and editing. **Abdussalam Azem:** Conceptualization; supervision; funding acquisition; validation; methodology; writing-review and editing.

References

1. Reichert AS, Neupert W. Mitochondriomics or what makes us breathe. Trends Genet [Internet]. 2004;20(11):555–62. Available from: https://www.sciencedirect.com/science/article/pii/S0168952504002495

2. Stehling O, Wilbrecht C, Lill R. Mitochondrial iron–sulfur protein biogenesis and human disease. Biochimie [Internet]. 2014 May;100:61–77. Available from: https://www.sciencedirect.com/science/article/pii/S0300908414000285

3. Pfanner N, Warscheid B, Wiedemann N. Mitochondrial proteins: from biogenesis to functional networks. Nat Rev Mol Cell Biol. 2019;20(5):267–84.

4. Filograna R, Mennuni M, Alsina D, Larsson NG. Mitochondrial DNA copy number in human disease: the more the better? FEBS Lett. 2021;595(8):976–1002.

5. Schmidt O, Pfanner N, Meisinger C. Mitochondrial protein import: From proteomics to functional mechanisms. Nat Rev Mol Cell Biol. 2010;11(9):655–67.

6. Pfanner N. Mitochondrial Machineries for Import and Assembly of Proteins. Biochim Biophys Acta - Bioenerg. 2016 Aug;1857:e4.

7. Mokranjac D, Neupert W. The many faces of the mitochondrial TIM23 complex. Biochim Biophys Acta - Bioenerg [Internet]. 2010;1797(6–7):1045–54. Available from: http://dx.doi.org/10.1016/j.bbabio.2010.01.026

8. Mokranjac D, Popov-[Cbreve]eleketiÄ D, Hell K, Neupert W. Role of Tim21 in Mitochondrial Translocation Contact Sites. J Biol Chem [Internet]. 2005 Jun;280(25):23437–40. Available from: http://www.jbc.org/content/280/25/23437.abstract

9. Ieva R, Schrempp SG, Opalinski L, Wollweber F, Hoss P, Heisswolf AK, et al. Mgr2 functions as lateral gatekeeper for preprotein sorting in the mitochondrial inner membrane. Mol Cell. 2014 Dec;56(5):641–52.

10. Richter F, Dennerlein S, Nikolov M, Jans DC, Naumenko N, Aich A, et al. ROMO1 is a constituent of the human presequence translocase required for YME1L protease import. J Cell Biol. 2019;218(2):598–614.

11. Sim SI, Chen Y, Lynch DL, Gumbart JC, Park E. Structural basis of mitochondrial protein import by the TIM23 complex. Nature. 2023;621(September).

12. Fielden LF, Busch JD, Merkt SG, Ganesan I, Steiert C, Hasselblatt HB, et al. Central role of Tim17 in mitochondrial presequence protein translocation. Nature. 2023;621(September).

13. Geissler A, Chacinska A, Truscott KN, Wiedemann N, Brandner K, Sickmann A, et al. The Mitochondrial Presequence Translocase. Cell [Internet]. 2002;111(4):507–18. Available from: http://www.sciencedirect.com/science/article/pii/S0092867402010735

14. Hell K, Nargang FE, Neupert W, Kozany C, Prokisch H, Paschen SA, et al. Tim50, a novel component of the TIM23 preprotein translocase of mitochondria. EMBO J [Internet]. 2003 Feb;22(4):816–25. Available from: http://dx.doi.org/10.1093/emboj/cdg090

15. Yamamoto H, Esaki M, Kanamori T, Tamura Y, Nishikawa S, Endo T. Tim50 Is a Subunit of the TIM23 Complex that Links Protein Translocation across the Outer and Inner Mitochondrial Membranes. Cell [Internet]. 2002;111(4):519–28. Available from: http://www.sciencedirect.com/science/article/pii/S009286740201053X

16. Meinecke M, Wagner R, Kovermann P, Guiard B, Mick DU, Hutu DP, et al. Tim50 maintains the permeability barrier of the mitochondrial inner membrane. Science. 2006 Jun;312(5779):1523–6.

17. Bose HS, Gebrail F, Marshall B, Perry EW, Whittal RM. Inner Mitochondrial Translocase Tim50 Is Central in Adrenal and Testicular Steroid Synthesis. Mol Cell Biol. 2019;39(1).

18. Tang K, Zhao Y, Li H, Zhu M, Li W, Liu W, et al. Translocase of Inner Membrane 50 Functions as a Novel Protective Regulator of Pathological Cardiac Hypertrophy. J Am Heart Assoc. 2017;

19.   Sankala H, Vaughan C, Wang J, Deb S, Graves PR. Upregulation of the mitochondrial transport protein, Tim50, by
mutant p53 contributes to cell growth and chemoresistance. Arch Biochem Biophys [Internet]. 2011;512(1):52–60.
Available from: http://dx.doi.org/10.1016/j.abb.2011.05.005

20.   Gao SP, Sun HF, Jiang HL, Li LD, Hu X, Xu XE, et al. Loss of TIM50 suppresses proliferation and induces apoptosis in
breast cancer. Tumour Biol. 2016 Jan;37(1):1279–87.

21.   Zhang X, Han S, Zhou H, Cai L, Li J, Liu N, et al. TIMM50 promotes tumor progression via ERK signaling and predicts
poor prognosis of non-small cell lung cancer patients. Mol Carcinog. 2019;58(5):767–76.

22.   F. S. A distinct type of 3-methylglutaconic aciduria due to a novel mutation in the Translocase of inner mitochondrial
membrane 50 (TIMM50) gene. European Journal of Neurology. 2015.

23.   Shahrour MA, Staretz-Chacham O, Dayan D, Stephen J, Weech A, Damsh N, et al. Mitochondrial epileptic
encephalopathy, 3-methylglutaconic aciduria and variable complex V deficiency associated with TIMM50 mutations.
Clin Genet. 2017 May;91(5):690–6.

24.   Tort F, Ugarteburu O, Texidó L, Gea-Sorlí S, García-Villoria J, Ferrer-Cortès X, et al. Mutations in TIMM50 cause
severe mitochondrial dysfunction by targeting key aspects of mitochondrial physiology. Hum Mutat. 2019;

25.   Reyes A, Melchionda L, Burlina A, Robinson AJ, Ghezzi D, Zeviani M. Mutations in TIMM50 compromise cell
survival in OxPhos☐dependent metabolic conditions . EMBO Mol Med. 2018;

26.   Mir A, Hadab S, Sammak M, Alhazmi R, Housawi Y, Bashir S. Complete resolution of epileptic spasms with vigabatrin
in a patient with 3-methylglutaconic aciduria caused by TIMM50 gene mutation. Clinical Genetics. 2020.

27.   Moudi M, Vahidi Mehrjardi MY, Hozhabri H, Metanat Z, Kalantar SM, Taheri M, et al. Novel variants underlying
autosomal recessive neurodevelopmental disorders with intellectual disability in Iranian consanguineous families. J Clin
Lab Anal. 2022;36(2):1–10.

28.   Shiota T, Mabuchi H, Tanaka-Yamano S, Yamano K, Endo T. In vivo protein-interaction mapping of a mitochondrial
translocator protein Tom22 at work. Proc Natl Acad Sci U S A [Internet]. 2011 Sep;108(37):15179–83. Available from:
https://www.jstor.org/stable/41352066

29.   Ben-Simon Y, Rodenas-Ruano A, Alviña K, Lam AD, Stuenkel EL, Castillo PE, et al. A Combined Optogenetic-
Knockdown Strategy Reveals a Major Role of Tomosyn in Mossy Fiber Synaptic Plasticity. Cell Rep. 2015;12(3):396–
404.

30.   Bhattacharya S. Metabolic Reprogramming in Nerve Cells. Am J Biomed Sci Res. 2023;18(2):116–30.

31.   Detmer SA, Chan DC. Functions and dysfunctions of mitochondrial dynamics. Nature Reviews Molecular Cell Biology.
2007.

32.   Johri A, Beal MF. Mitochondrial dysfunction in neurodegenerative diseases. Journal of Pharmacology and Experimental
Therapeutics. 2012.

33.   Wang W, Zhao F, Ma X, Perry G, Zhu X. Mitochondria dysfunction in the pathogenesis of Alzheimer's disease: Recent
advances. Molecular Neurodegeneration. 2020.

34.   Course MM, Hsieh CH, Tsai PI, Codding-Bui JA, Shaltouki A, Wang X. Live imaging mitochondrial transport in
neurons. In: Neuromethods. 2017.

35.   Sheng ZH. Mitochondrial trafficking and anchoring in neurons: New insight and implications. J Cell Biol.
2014;204(7):1087–98.

36.   D'adamo MC, Liantonio A, Rolland JF, Pessia M, Imbrici P. Kv1.1 channelopathies: Pathophysiological mechanisms
and therapeutic approaches. Int J Mol Sci. 2020;21(8):1–21.

37.   Harvey AL. Twenty years of dendrotoxins. Toxicon. 2001;39(1):15–26.

38.   Glazebrook PA, Ramirez AN, Schild JH, Shieh C, Doan T, Wible BA, et al. Potassium channels Kv1.1, Kv1.2 and

685   Kv1.6 influence excitability of Rat Visceral Sensory Neurons. 2002;467–82.

686   39.   Vögtle FN, Wortelkamp S, Zahedi RP, Becker D, Leidhold C, Gevaert K, et al. Global Analysis of the Mitochondrial N-

687   Proteome Identifies a Processing Peptidase Critical for Protein Stability. Cell. 2009;139(2):428–39.

688   40.   Pfanner Nikolaus, Geissler Andreas. Versatility of the mitochondrial protein import machinery. Nat Rev Mol Cell Biol

689   [Internet]. 2001;2(May):339–49. Available from: https://www.nature.com/articles/35073006

690   41.   Chacinska A, Lind M, Frazier AE, Dudek J, Meisinger C, Geissler A, et al. Mitochondrial presequence translocase:

691   Switching between TOM tethering and motor recruitment involves Tim21 and Tim17. Cell [Internet]. 2005;120(6):817–

692   29. Available from: http://dx.doi.org/10.1016/j.cell.2005.01.011

693   42.   Káldi K, Bauer MF, Sirrenberg C, Neupert W, Brunner M. Biogenesis of Tim23 and Tim17, integral components of the

694   TIM machinery for matrix-targeted preproteins. EMBO J. 1998;17(6):1569–76.

695   43.   Gevorkyan-Airapetov L, Zohary K, Popov-ÄŒeleketiÄ D, Mapa K, Hell K, Neupert W, et al. Interaction of Tim23 with

696   Tim50 Is Essential for Protein Translocation by the Mitochondrial TIM23 Complex. J Biol Chem [Internet]. 2009

697   Feb;284(8):4865–72. Available from: http://www.jbc.org/content/284/8/4865.abstract

698   44.   Shen W, Hernandez-Lopez S, Tkatch T, Held JE, Surmeier DJ. Kv1.2-Containing K+ Channels Regulate Subthreshold

699   Excitability of Striatal Medium Spiny Neurons. J Neurophysiol. 2004;91(3):1337–49.

700   45.   Dodson PD, Barker MC, Forsythe ID. Two heteromeric Kv1 potassium channels differentially regulate action potential

701   firing. J Neurosci. 2002;22(16):6953–61.

702   46.   Kaczmarek LK. Policing the Ball: A New Potassium Channel Subunit Determines Inactivation Rate. Neuron. 2006 Mar

703   2;49(5):642–4.

704   47.   Schulte U, Thumfart JO, Klöcker N, Sailer CA, Bildl W, Biniossek M, et al. The epilepsy-linked Lgi1 protein assembles

705   into presynaptic Kv1 channels and inhibits inactivation by Kvβ1. Neuron. 2006;49(5):697–706.

706   48.   Vangipuram M, Ting D, Kim S, Diaz R, Schüle B. Skin punch biopsy explant culture for derivation of primary human

707   fibroblasts. J Vis Exp. 2013;(77):9–11.

708   49.   Lavi A, Sheinin A, Shapira R, Zelmanoff D, Ashery U. DOC2B and Munc13-1 differentially regulate neuronal network

709   activity. Cereb Cortex. 2014;

710   50.   Seyfried NT, Dammer EB, Swarup V, Nandakumar D, Duc M, Yin L, et al. A Multi-network Approach Identifies

711   Protein-Specific Co-expression in Asymptomatic and Symptomatic Alzheimer's Disease. 2018;4(1):60–72. Available

712   from: https://www.synapse.org/#!Synapse:syn2580853

713   51.   Rath S, Sharma R, Gupta R, Ast T, Chan C, Durham TJ, et al. MitoCarta3.0: An updated mitochondrial proteome now

714   with sub-organelle localization and pathway annotations. Nucleic Acids Res. 2021;49(D1):D1541–7.

715   52.   Lindenboim L, Ferrando-May E, Borner C, Stein R. Non-canonical function of Bax in stress-induced nuclear protein

716   redistribution. Cell Mol Life Sci. 2013;70(16):3013–27.

717   53.   Zala D, Hinckelmann MV, Yu H, Lyra Da Cunha MM, Liot G, Cordelières FP, et al. Vesicular glycolysis provides on-

718   board energy for fast axonal transport. Cell. 2013;152(3):479–91.

Figure Legends

Fig 1. Differential effects of *TIMM50* mutation on the expression of TIM23, PAM and TOM subunits and matrix-destined proteins.

(A) Healthy control- and patients-derived primary fibroblasts were lysed and analyzed by immunoblot with the indicated antibodies. Actin was used as loading control. Full results and original blots are found in supplemental material S1 Raw images. (B) Band density analysis of the blots presented in A (and in supplemental material S1 Raw images), showing significant decrease in levels of TIMM50, TIMM23 and TIMM17A/B, but not in TIMM21, PAM subunits, TOM subunits and matrix TIM23 complex substrates. Band signals were normalized to the loading control and compared to the level measured in the healthy control sample, taken as 100%. Data are shown as means ± SEM, n = 6 biological repeats for TIMM50 antibody, n = 3-5 biological repeats for all other antibodies, *p-value < 0.05, **p-value < 0.01, ***p-value < 0.001, ****p-value < 0.0001, Ordinary one-way ANOVA.

Fig 2. TIMM50 knockdown in mouse primary cortical neurons serves as a model system to study TIMM50 deficiency in mammalian neurons.

(A) Schematic depictions of the different constructs used in this study. "pLL3.7 control" is a control for EGFP expression. "Scr control" is a control for the shRNA system activation with a scrambled Sh3 sequence and Sh1-Sh3 are three plasmids expressing different TIMM50 targeting shRNA sequences. (B) Neuronal cultures were transduced to express the indicated constructs, lysed and analyzed by immunoblot with the indicated antibodies. Tubulin was used as loading control. "Untreated" control are cells that were not transduced. Full results and original blots are found in supplemental material S2 Raw images. (C) Band density analysis of the blots presented in B (and in supplemental material S2 Raw images) showing significant decrease in levels of TIMM50, TIMM23 and TIMM17A/B, but not in TIMM21, PAM subunits, TOM subunits and matrix TIM23 complex substrates. Band signals were normalized to the loading control and compared to the Scr control, taken as 100%. Data are shown as means ± SEM, n = 14 biological repeats for TIMM50 antibody, n = 3-4 biological repeats for all other antibodies, **p-value < 0.01, ****p-value < 0.0001, unpaired Student's t-test.

Fig 3. TIMM50 deficiency does not affect the majority of MIM and mitochondrial matrix proteome.

Untargeted mass spectrometry analysis of fibroblasts from both patients (P1, P2), as compared to the healthy control, and of mice primary neurons transduced with either Sh2 or the Scr control. In the volcano plots displayed in (A, B), the y-axis cut off of >1.301 corresponds to –log (0.05) or p-value = 0.05, while the x-axis cut off of <-0.5 and >0.5 corresponds to a ±1.414-fold change. Each dot in the graph represents a protein. Proteins depicted on the right side of the x-axis cut-off and above the y-axis cut off were considered to be increased in amount, while proteins depicted on the left side of the x-axis cut-off and above the y-axis cut off were considered to be decreased in amount. All relevant proteins that were increased/decreased in amount are identified by their gene name, next to the dot. Statistical analysis was performed using Student's t-test and a p-value <0.05 was considered statistically significant. For fibroblasts, n = 9 per group (three biological repeats in triplicate), P1 and P2 results were compared to HC results. For neurons, n = 9 per group (three biological repeats in triplicates), Sh2 results were compared to Scr control results. Full list of differentially expressed proteins in fibroblasts and neurons is found in supplemental material S1-3 Datasets. (A) The steady state levels of a majority of MIM proteins detected in patient fibroblasts and TIMM50 KD neuronal cells were not affected. MIM proteins are colored red, while other detected mitochondrial proteins are colored grey. (B) The steady state levels of a majority of matrix proteins detected in patient fibroblasts and TIMM50 KD neuronal cells were not affected. Matrix proteins are colored red, while other detected mitochondrial proteins are colored grey.

Fig 4. TIMM50 deficiency leads to decreased protein levels of the OXPHOS and MRP machineries.

Untargeted mass spectrometry analysis of fibroblasts from both patients (P1, P2), as compared to the healthy control, and of mice primary neurons transduced with either Sh2 or the Scr control. In the volcano plots displayed in (A, B), the y-axis cut off of >1.301 corresponds to –log (0.05) or p-value = 0.05, while the x-axis cut off of <-0.5 and >0.5 corresponds to a ±1.414-fold change. Each dot in the graph represents a protein. Proteins depicted on the right side of the x-axis cut-off and above the y-axis cut off were considered to be increased in amount, while proteins depicted on the left side of the x-axis cut-off and above the y-axis cut off were

considered to be decreased in amount. All relevant proteins that were increased/decreased in amount are identified by their gene name, next to the dot. Statistical analysis was performed using Student's t-test and a p-value <0.05 was considered statistically significant. For fibroblasts, n = 9 per group (three biological repeats in triplicate), P1 and P2 results were compared to HC results. For neurons, n = 9 per group (three biological repeats in triplicates), Sh2 results were compared to Scr control results. Full list of differentially expressed proteins in fibroblasts and neurons is found in supplemental material S1-3 Datasets. (A) Decreased steady state levels of multiple OXPHOS subunits was observed in patient fibroblasts. A similar trend was observed in TIMM50 KD neuronal cells. OXPHOS proteins are colored red, while other detected mitochondrial proteins are colored grey. (B) Decreased steady state levels of multiple MRP subunits was observed in patient fibroblasts and TIMM50 KD neuronal cells. MRP proteins are colored red, while other detected mitochondrial proteins are colored grey. (C, D) Mitochondrial DNA content in fibroblasts (C) and neurons (D) was not affected in TIMM50 deficient cells compared to control cells. Mitochondrial DNA content was estimated by measuring the ratio between mitochondrial and nuclear DNA. Dloop1 expression was measured by qPCR relative to HPRT; TERT served as a control of a nuclear-encoded gene. Ethidium bromide (EtBr; 100 ng/ml) was used as positive control. Data are shown as means ± SEM, n = 6 (samples from three biological replicates, each performed twice), *p-value < 0.05, ***p-value < 0.001, Kruskal-Wallis test.

Fig 5. TIMM50 deficiency negatively impacts OXPHOS machinery and glycolysis functions.

(A) A Seahorse XF cell mito-stress assay was used to measure mitochondrial oxygen consumption rates at basal levels and in response to the indicated effectors in fibroblasts (upper panel) and neuronal cells (lower panel). (B) A Seahorse XF cell glycolysis stress assay was used to measure glycolysis at basal levels and in response to the indicated effectors in fibroblasts (upper panel) and neuronal cells (lower panel). (C) Basal respiration, maximal respiration and ATP-linked respiration were reduced in patient fibroblast cells compared to HC cells. Data are shown as means ± SEM, *p-value < 0.05, ***p-value < 0.001, ****p-value < 0.0001, Ordinary one-way ANOVA. (D) Basal respiration, maximal respiration and ATP-linked respiration were reduced in TIMM50 KD neuronal cells compared to Scr control-transduced neuronal cells. Data

are shown as means ± SEM, ***p-value < 0.001, ****p-value < 0.0001, unpaired Student's t-

test. (E) Basal glycolysis remained similar, while glycolytic capacity and glycolytic reserves

were reduced in patient fibroblast cells compared to HC cells. Data are shown as means ± SEM,

*p-value < 0.05, ***p-value < 0.001, ****p-value < 0.0001, Kruskal-Wallis test. (F) Basal

glycolysis remained similar, while glycolytic capacity and glycolytic reserves were reduced in

TIMM50 KD neuronal cells compared to Scr control-transduced neuronal cells. Data are shown

as means ± SEM, ***p-value < 0.001, ****p-value < 0.0001, unpaired Student's t-test. For all

the experiments shown in (A-F), n = 4-6 biological repeats of 3-6 technical repeats each.

819 Fig 6. TIMM50 deficiency leads to defective mitochondrial trafficking along neuronal

820 processes.

(A) Representative images of neuronal processes in neuronal cultures that were co-transfected

with a dsRed-mito plasmid and either a Scr control plasmid (left panel) or a TIMM50 KD

plasmid (right panel). (B) Kymographs of the same processes in A, showing the displacement of

mitochondria over time. Y axis length is 5 minutes, X axis length is about 100 μm. (C) Lower

percentage of moving mitochondria was observed in TIMM50 KD neuronal processes compared

to controls. Each dot in the graph represents the percentage of moving mitochondria out of the

total observed mitochondria in a single neurite. Data are shown as means ± SEM, n = 13-31

neurites per condition, analyzed from three biological repeats, ****p-value < 0.0001, Ordinary

one-way ANOVA. (D) Decreased mitochondrial cumulative travelling distance was observed in

TIMM50 KD neuronal cells compared to controls. Each dot in the graph represents a single

moving mitochondrion. n = 84-130 mitochondria per condition, analyzed from three biological

repeats, *p-value < 0.05, Kruskal-Wallis test. (E) Slower mitochondrial movement was observed

in TIMM50 KD neuronal cells compared to controls. Each dot in the graph represents a single

moving mitochondrion. n = 84-130 mitochondria per condition, analyzed from three biological

repeats, ****p-value < 0.0001, Kruskal-Wallis test. (F) No change in mitochondrial trafficking

proteins was observed TIMM50 KD neuronal cells. The y-axis cut off of >1.301 corresponds to

–log (0.05) or p-value = 0.05, while the x-axis cut off of <-0.5 and >0.5 corresponds to a ±1.414-

fold change. Each dot in the graph represents a protein. Proteins depicted on the right side of the

x-axis cut-off and above the y-axis cut off were considered to be increased in amount, while

proteins depicted on the left side of the x-axis cut-off and above the y-axis cut off were

considered to be decreased in amount. Statistical analysis was performed using Student's t-test
and a p-value <0.05 was considered statistically significant. n = 9 per group (three biological
repeats in triplicate). Full list of differentially expressed proteins in neurons is found in S3
Dataset. (G) Lower cellular ATP levels were observed in TIMM50 KD neuronal cells compared
to Scr control-transduced cells. Data are shown as mean ± SEM, n = 27 quantified wells (each
containing $5x10^4$ cells) per condition, from three biological repeats with nine technical repeats
each, ***p-value < 0.001, unpaired Student's t-test.

Fig 7. TIMM50 deficiency leads to a significant decrease in the levels of KCNA2 and
KCNJ10 potassium channels and an increased electrical activity.
(A) Example of voltage traces in response to increased depolarization of TIMM50 KD neurons.
The bottom panel shows the protocol applied, while the top panel shows the typical response
pattern that was measured. (B) Representative traces of a single action potential, received at
rheobase level, from a TIMM50 KD neuron, as compared to Scr control-transduced neuron. (C)
Representative traces of the maximal stimulus for each condition, showing the maximal amount
of action potentials measured for each group. (D) No change in rheobase was observed in
TIMM50 KD neuronal cells, as compared to Scr control-transduced cells. Rheobase was
measured as the first current step that caused firing of an action potential. Data are shown as
means ± SEM, n = 28 / 37 cells (for Scr control-/-Sh2-transduced cells, respectively) from three
biological repeats, Mann-Whitney test. (E) No change in the action potential half-width was
observed in TIMM50 KD neuronal cells, as compared to Scr control-transduced cells. Half-width
was measured as the time between the rising and falling phases of the action potential, at the
half-point between the tip of the peak and the bottom of the voltage rising curve. Data are shown
as means ± SEM, n = 28 / 37 cells (for Scr control-/ Sh2-transduced cells, respectively) from
three biological repeats, Mann-Whitney test. (F) No change in the action potential rate of fall
was observed in TIMM50 KD neuronal cells, as compared to Scr control-transduced cells. Rate
of fall was measured as the time between the action potential peak and the baseline following the
action potential, divided by the change in voltage between the same two points ($\Delta X/\Delta Y$). Data
are shown as means ± SEM, n = 28 / 37 cells (for Scr control / Sh2 respectively) from three
biological repeats, Mann-Whitney test. (G) Decreased action potential latency was observed in
TIMM50 KD neuronal cells compared to Scr control-transduced cells. Latency was measured as

the time difference between the beginning of the pulse to the peak of the action potential. Each

dot in the graph represents the average latency of the first five consecutive action potentials

appearing after rheobase level. Data are shown as means ± SEM, n = 40 / 50 cells (for Scr

control-/Sh2-transduced cells, respectively) from four biological repeats, *p-value < 0.05,

unpaired Student's t-test. (H) An increase in the maximal number of action potentials fired in a

single stimulus was observed in TIMM50 KD neuronal cells, as compared to Scr control-

transduced cells. Data are shown as means ± SEM, n = 28 / 37 cells (for Scr control-/Sh2-

transduced cells, respectively) from three biological repeats, ****p-value < 0.0001, unpaired

Student's t-test. (I) Amongst the detected ion channel proteins, a specific decrease in KCNA2

and KCNJ10 potassium channels was observed in TIMM50 KD neuronal cells. The y-axis cut

off of >1.301 corresponds to –log (0.05) or p-value = 0.05, while the x-axis cut off of <-0.5 and

>0.5 corresponds to a ±1.414-fold change. Each dot in the graph represents a protein. Proteins

depicted on the right side of the x-axis cut-off and above the y-axis cut off were considered to be

increased in amount, while proteins depicted on the left side of the x-axis cut-off and above the

y-axis cut off were considered to be decreased in amount. Statistical analysis was performed

using Student's t-test and a p-value <0.05 was considered statistically significant. n = 9 per group

(three biological repeats in triplicate). Full list of differentially expressed proteins in neurons is

found in S3 Dataset.

Supporting information

S1 Fig. Immunoblot confirmation of mass spectrometry-based proteomics findings.

(A) Verification of the measured increase in the levels of ALDH2 and GRSF1 in fibroblasts.

Three biological repeats were performed for each antibody. Full results and original blots are

found in S1 Raw images. (B) Verification of the measured decrease in the levels of the KCNA2

potassium channel in neuronal cultures. Three biological repeats were performed. Full results

and original blots are found in S2 Raw images.

S2 Fig. Spontaneous excitatory activity recordings and the effect of α-DTX on TIMM50 KD

neuronal cells.

(A) Representative traces of mEPSCs measurements taken in the presence of 1 μM TTX. (B) No change in the average mEPSC amplitude, frequency and area under the peak was observed in TIMM50 KD neuronal cells compared to Scr control-transduced cells. Data are shown as means ± SEM, n = 23 / 20 cells (for Scr control-/Sh2-transduced cells, respectively) from three biological repeats, Mann-Whitney test. (C) Histogram of the relative frequency distribution of the mEPSC amplitude measurements. (D) Cumulative distribution function of the mEPSC amplitude measurements. (E) Measurements of the number of action potentials fired before and after treatment with 100 nM α-DTX for Scr control- and Sh2-transduced neuronal cells. (F) A higher difference in the number of action potentials fired after and before α-DTX application was observed in Scr control-transduced cells compared to TIMM50 KD neuronal cells. n = 21 / 20 cells (for Scr control-/Sh2-transduced cells, respectively) from three biological repeats, ***p-value < 0.001, unpaired Student's t-test.

S1 Table. Oligonucleotides used in this study.

S2 Table. Reagents and tools used in this study.

S1 Raw images. Original blots – Fibroblasts.

S2 Raw images. Original blots – Neurons.

S1 Dataset. HC vs P1 all proteins fold change.

S2 Dataset. HC vs P2 all proteins fold change.

S3 Dataset. Scr vs Sh2 all proteins fold change.

S1 Movie. Representative mitochondrial trafficking movies. Scr control neurons and TIMM50 KD neurons co-transfected with the corresponding control/KD plasmid and mito-DsRed plamid. The cells were imaged for 5 minutes with a 3 second interval between each image. The videos displayed are sped up by ~15x.