The ontogeny of immune tolerance: a model of early-life secretory IgA - gut microbiome interactions

Burcu Tepekule[1*], Ai Ing Lim[2], C. Jessica E. Metcalf[1*]

[1] Dept of Ecology and Evolutionary Biology, Princeton University, Princeton, 08544, USA.
[2] Dept of Molecular Biology, Princeton University, Princeton, 08544, USA.

*Corresponding authors. Emails: burcutepekule@gmail.com, cjmetcalf@princeton.edu

Abstract: To achieve immune and microbial homeostasis during adulthood, the developing immune system must learn to identify which microbes to tolerate and which to defend against. How such 'immune education' unfolds remains a major knowledge gap. We address this gap by synthesizing existing literature to develop a mechanistic mathematical model representing the interplay between gut ecology and adaptive immunity in early life. Our results indicate that the inflammatory tone of the microenvironment is the mediator of information flow from pre- to post-weaning periods. We evaluate the power of postnatal fecal samples for predicting immunological trajectories, and explore breastfeeding scenarios when maternal immunological conditions affect breastmilk composition. Our work establishes a quantitative basis for 'immune education', yielding insights into questions of applied relevance.

Introduction

The global burden of immune-mediated diseases is rapidly growing (*1*). Epidemiological data indicate that early life exposures are key determinants of immune-mediated diseases later in life (*2*), such as allergies (*3*), asthma (*4*), type 1 diabetes (*5*), and inflammatory bowel disease (IBD) (*6*). This impact of early life is primarily attributed to interactions between the microbiome and the immune system during a critical developmental window, which enables hosts to establish tolerance to commensal bacteria (*7*), ensuring the maintenance of immune and microbial homeostasis into adulthood, while appropriately defending against pathogens (*8*, *9*). When this crosstalk is perturbed, pathological imprinting may develop, characterized by excessive immune reactivity and increased susceptibility to inflammatory diseases in adulthood (*10*). Both inherent microbial factors and maternal cues determine how microbe-immune interactions unfold during early life: symbiotic commensals provide metabolic products that establish regulatory pathways facilitating a balanced immune response, whereas pathogens stimulate the immune system to develop defense mechanisms. Breastmilk delivers microbes and nutrients, but also antibodies that dictate the timing and nature of bacterial antigen presentation to the infant's developing immune system, establishing a transgenerational cycle of immune priming (*11*, *12*). The collective influence of these processes on immune ontogeny and maturation is encapsulated by the term 'immune education' (*13*).

The information available to tackle the establishment of 'immune education' is now considerable (*14*). To date, thousands of papers have been published, ranging from observational data from human populations to experimental perturbations in animal models. However, experimental methods are inevitably reductionist, focusing on a limited set of mechanisms, while larger scale descriptive analysis rooted in observational data may illuminate patterns, but ultimately yield verbal descriptions lacking mathematical characterization. The time is ripe to integrate these layers

of evidence into a systems biology framework that formally accounts for the multiple potential interacting components.

To this end, we introduce a mathematical framework describing the reciprocal imprinting of the human gut microbiome and the antigen-specific endogenous mucosal Secretory Immunoglobulin A (SIgA) response during the first two years of life. While immune outcomes are undoubtedly multifaceted and influenced by numerous factors, many lines of evidence indicate the importance of the SIgA response, as its reactivity to microbial antigens plays a crucial role in mediating gut immune responses and their dysregulation (*15, 16*). The SIgA response also integrates both the B-cell and T-cell arms of immune ontogeny, and manifests the dual functionality of the gut mucosal immune system by selectively neutralizing pathogens while tolerating commensal bacteria beneficial for immune homeostasis (*17, 18*). Furthermore, SIgA represents the transgenerational aspect of immune priming: maternal SIgA modulates antigen presentation while simultaneously regulating the gut community composition (*11, 19*).

Our mechanistic modeling strategy (Fig. 1) blends flexibility with tractability in reflecting the effects of maternal factors, feeding practices, consumer resource dynamics within microbial communities, and multifunctionality of SIgA (Fig. 1B); and introduces a quasi-stochastic mathematical model of germinal center reactions embedded in a combination of ordinary and delay differential equations (Fig. 1C). We parameterize our framework using data from infant fecal samples (*20–23*), formally mapping microbial abundance measurements into expectations for the dynamics of the lumen and the germinal centers of the gut, contexts traditionally obscured by the impracticality of direct sampling (*24*) (Fig. 1A). The mechanistic model structure entails a high dimensional parameter space, but qualitative results are consistent across parameter combinations (see materials and methods).

Fig 1. Modeling framework. A) Key assumptions and mechanisms enabling predictions for the gut lumen and germinal centers (GCs). 2Y: 2 years; mSIgA and eSIgA: maternal and endogenous secretory Immunoglobulin A. **B)** Model dynamics. Breastfeeding (BF) and complementary feeding (CF) introduces human milk oligosaccharides (HMOs), plant-derived polysaccharides (PDPs), mSIgA, and commensal bacteria to the gut lumen as SIgA-bacteria complexes, or bacteria alone. These bacterial populations modulate the inflammatory tone of the microenvironment of the gut lumen (σ), influencing epithelial cell-mediated communication to the Peyer's Patches (PPs). Composition of antigens sampled by Microfold (M) cells drives the balance between antigen-specific regulatory versus helper T cells. This balance and σ shape the selection threshold δ_i for taxon i, which determines the BCR affinity ranges that lead to different B cell fates: apoptosis, continued circulation, or terminal differentiation into IgA-secreting plasma cells. Alongside with the HMOs, PDPs, and the oxygen concentration (O_2) based on commensals' metabolism, IgA secreted by the plasma cells regulates the microbiome. VB: Vaginal Birth. **C)** GC reactions (steps 1-7, materials and methods). Naïve B cells with a wide BCR affinity distribution migrate to the PPs, and their affinity distribution is combined with that of circulating B-cells. B-cells activate upon encountering bacterial antigens. Depending on δ_i, cells undergo positive selection, receiving adequate T cell help to continue circulating or differentiate to IgA-secreting plasma cells, or become apoptotic. Circulating cells undergo somatic hypermutation (SHM), increasing the standard deviation of their BCR affinity distribution, continuing the cycle.

Overall, our model brings mathematical formalism to the concept of 'immune education', opening the way to investigating persistent questions in early life immunology, including the power of fecal sampling data in predicting immunological outcomes, and the determinants of transmission of immune information from early infancy to adulthood. Finally, it establishes hypotheses regarding breastfeeding practices guided by maternal immunological profiles.

Results

Model outcomes unveil the dynamics in gut lumen, successfully predicting out-of-sample data

Capturing immune 'education' requires encompassing exogenous inputs into the gut lumen alongside endogenous dynamics in the gut lumen and Peyer's patches (PPs, Fig. 1A). Exogenous inputs quantified based on different feeding practices of infants (*25–27*) (Eqns. S1.1.18-S1.1.21), include caloric intake from human milk oligosaccharides (HMOs) and plant-derived polysaccharides (PDPs), and maternal secretory Immunoglobulin A (mSIgA) concentrations. HMOs and PDPs differentially modulate growth rates of different bacterial taxa based on their metabolism (*28–32*) (Eqn. S1.1.4). Timing of endogenous immune system activation is dictated by the maturation of Microfold (M) cells, which serve as the primary route for antigen transport to PPs (Fig. 1B), the key site for the development of the gut adaptive immune response (*33*). M cells appear shortly before weaning, primarily due to decreasing concentrations of maternal steroids in breastmilk (*34*), for which we use mSIgA as a proxy, given a lack of quantitative data. M cell maturation translates into antigenic sampling — and thus the start of immune response maturation (Fig. 2A).

To fit our model to fecal microbiome data (*20–23*) and the established timeline of immune ontogeny (*14, 34*), we leverage two observations: i) the infant develops an SIgA profile resembling their nursing mother's (*21*), and ii) the infant's immune response and microbial community structure stabilize after two years (*33*) (Fig. 1A). Specifically, the matured offspring secretes endogenous antibodies with quantitatively similar affinities to bacterial groups as their nursing mother; and relative abundances of bacterial taxa converge to their equilibrium values in host's fecal samples from day 720. We estimate a single set of affinity values shared between the mother and the matured offspring, reflecting maternal antibodies in the gut lumen before activation and after the stabilization of the host's immune response. For this step of inference, we define a composite optimization criterion that integrates the relative and absolute abundances of key taxonomic groups observed in fecal samples of infants and their IgA indexes (indicating enrichment in the SIgA coated (SIgA+) fractions). Optimization provides initial estimates which then guide inference for the interim phase of endogenous immune system maturation. A second composite optimization criterion is applied to match the average SIgA coating ratio (*36*) in the gut lumen and the matured endogenous affinity values to the maternal ones.

To capture immunological and competitive interactions across ontogeny, we selected four bacterial taxonomic groups to reflect broad categories of host/bacterial interactions: *Enterobacteriaceae* encompasses the *Escherichia-Shigella* genus with potentially pathogenic (*37, 38*) bacteria encountered pre-weaning when the endogenous system is more vulnerable to enteric infections. *Bifidobacteriaceae* represents early colonizers with anti-inflammatory properties that regulate the microenvironment and provide colonization resistance. *Bacteroidaceae* and *Clostridiales* are post-

weaning bacteria with potentially anti-inflammatory properties that contribute to a balanced gut environment when properly regulated by SIgA (Table S2). Taxa are differentiated by two parameters: *i)* their invasiveness (*35*), defined as their potential to penetrate colonic epithelial cells, and *ii)* their immunostimulatory potential, defined as their capacity to modulate the inflammatory tone of the microenvironment (*40*, *41*) via their metabolic products and epithelial conditioning (*42*). While bacterial niche (such as proximity to the epithelium (*43*)) could serve as an additional parameter for differentiating taxa in relation to SIgA responses, it has not been included in the model due to lack of existing quantitative data.

There is substantial evidence that the transfer of maternal antibodies to the infant gut through breastmilk plays a crucial role in shaping the neonatal gut microbiome and the immune system's development (*19*, *44*, *45*). We characterize SIgA's functionality into two mutually exclusive categories: masking(M), which represents SIgA's non-neutralizing functionality, promoting the intestinal residency of the symbiotic commensals by limiting their interactions with the immune system (*46*, *47*); and neutralizing (N), contributing to the expulsion of bacteria from the gut (*43*). The masking (M) or neutralizing (N) rates per one unit of SIgA concentration are both modeled as a graduated response to affinity levels (*48*) (Eqn. S1.1.1-S1.1.2). This dual function of SIgA creates three subpopulations in the gut lumen — uncoated (SIgA-), masked (SIgA+ (M)), and neutralized (SIgA+ (N)) — of which only the uncoated and masked are metabolically active. The neutralized subpopulation neither interacts with other subpopulations nor stimulates the immune system. Both masked and neutralized populations are coated with SIgA, meaning that techniques like IgA-Seq or BugFACS would detect SIgA binding in both cases. The SIgA coating ratio, which measures the proportion of bacteria coated with SIgA, would reflect contributions from both masked and neutralized bacteria. The functional dichotomy between high-affinity neutralizing and low-affinity masking SIgA is not absolute, as SIgA binding can influence bacterial growth or colonization in species-specific ways, involving mechanisms such as enchained growth, biofilm formation, or altered gene expression (*17*). Given the scale of data available, our model does not attempt to capture these nuances (see supplementary material, section *Model Limitations*), but instead focuses on a fundamental, bacteria-agnostic mechanism that aims to capture the core features of SIgA-microbiome interactions. We based our model on the observation that SIgA targeting pathogens or pathobionts generally exhibits higher affinity compared to SIgA targeting symbiotic commensals, which often neutralizes rather than promotes their presence (*49*). Rather than relying on species-specific rules, our framework explores how affinity and functionality shift dynamically under different physiological conditions such as homeostasis and inflammation.

Our model assumes a common set of mechanisms for endogenous SIgA maturation across all commensals, allowing rates of antibody masking and neutralization to emerge from maturing affinity levels. Our quasi-stochastic formulation of germinal center reactions assumes a temporally dynamic distribution of B-cell receptor (BCR) binding variability that dictates B cell fates; mean and standard deviation of binding variability are adjusted at each proliferation, somatic hypermutation, and selection (P-SHM-S) cycle (Fig. 1C). Naïve B cells migrate to the PPs, are activated depending on the antigen uptake by M cells (Eqns. S1.1.5-S1.1.7) and start participating in GC reactions (*50*). Each P-SHM-S cycle is informed by the microenvironmental stimulation in the PPs, and the selection pressure imposed by an implicit model of antigen-specific follicular helper and regulatory T cells (Materials and Methods, *Modeling the role of Germinal Centres*, and Eqn. S1.2.8), eventually determining the size and average affinity of the plasma cell pool.

Microenvironmental stimulation is modeled as a composite variable (Eqn. S1.1.8) reflecting the cumulative effects of cytokines and microbial products; akin to the inflammatory tone of the local environment (*51*). We infer mSIgA affinities (Fig. 2B) assuming the nursing mother with a balanced microbiome-immune response transfers masking antibodies for the symbiotic commensals (*Bifidobacteriaceae*, *Bacteroidaceae* and *Clostridiales*); and neutralizing antibodies for the pathogenic taxon *Enterobacteriaceae* (*48*, *52–56*). Model fit estimates illustrate the trajectory of microenvironmental stimulation (Fig. 2C), average affinity of the endogenous antibodies converging toward maternal values (Fig. 2D), along with the normalized concentrations of antibody levels (Fig. 2E) proportional to the maturation of plasma cells being imprinted as the host ages.

Bacterial colonization is also affected by the oxygen concentration (O_2) in the gut lumen (Eqn. S1.1.4). O_2 inhibits the growth of obligate anaerobic bacteria and is consumed by facultative anaerobes (Eqn. S1.2.3) (*57*). From an initially normalized level of 1, O_2 degrades at a rate inferred during the model fitting process (Fig. 2C).

To parametrize our model, we use relative bacterial abundance and IgA-Seq data derived from infants' fecal samples (*20–23*) as a proxy for the ecological and immunological dynamics within the gut lumen (*58*); however, equating fecal samples with the gut lumen overlooks SIgA's role in microbial turnover (*46*, *47*). We reconstruct the relative abundances per fecal content by calculating the weighted sum of neutralized, coated, and uncoated subpopulations in the gut lumen using distinct observation rates (Eqns. S1.1.14-S1.1.17) which translates the microbial abundances in the gut lumen to abundances detected in fecal samples (Fig. 1A), allowing us to align the model output with the 16S rRNA sequencing analysis provided in *Tsukuda et al* (*20*). We apply the same reconstruction using SIgA−, SIgA+(C), and SIgA+(N) subpopulations to compute the IgA indexes for each taxon, allowing us to include the IgA-seq profiles of *Enterobacteriaceae* and *Bifidobacteriaceae* provided in *Planer et al.* (*21*). Our estimates led to an average coating ratio of 37.2%, with negative IgA indexes for *Bifidobacteriaceae*, *Bacteroidaceae*, *Clostridiales*, and a positive IgA index for *Enterobacteriaceae* (Fig. 2F).

Relative abundance estimates for periods of inactivity (neonatal, early postnatal) or stability (after 2 years) of the endogenous immune system demonstrate the reliability of our model fitting procedure (Fig. 2G). Furthermore, out-of-sample data corresponding to the period of maturation— not used for parameter inference (Fig. 2G, gray area)—was also accurately predicted by the model. This alignment provides indirect validation of the underlying assumptions regarding the immune response's impact on microbial community dynamics, lending credibility to the model estimates of the gut lumen (Fig. 2H) for which direct data is unavailable (Fig. 1A).

Fig 2. Illustration of Core Mechanisms and Model Fit. (**A**) Model inputs including normalized maternal secretory immunoglobulin A (mSIgA) concentration, human milk oligosaccharide (HMOs) and plant-derived polysaccharides (PDPs) caloric inputs, and timing of the endogenous immune system activation over time. EBM: exclusive breastfeeding; MF: mixed feeding; ECF: exclusive complementary feeding. (**B**) Rate and type of SIgA function relative to the level of antibody affinity, where mSIgA values for key taxa are highlighted. (**C**) Normalized oxygen concentration and microenvironmental stimulation (normalized by its peak value, reflecting the inflammatory tone of the environment) over time. (**D**) Temporal progression of average

endogenous SIgA (eSIgA) affinities (solid lines) gradually approaching the mSIgA affinities (dashed lines), (**E**) and the normalized antibody concentrations in the gut lumen for each key taxon. (**F**) Fractions of coated and neutralized bacteria and IgA indexes for each key taxon at DOL 735. (**G**) Relative abundances in fecal samples aligned with observational data, where the gray area highlights the out-of-sample data during parameter inference. (**H**) Absolute abundances in the gut lumen for key taxa distinguished by coating status. Model fit estimates for all parameters with their respective description, unit, and prior distribution are provided in Table S3. DOL: Day of life; E: *Enterobacteriaceae*; B: *Bifidobacteriaceae*; BC: *Bacteroidaceae*; C: *Clostridiales*.

Relative abundance estimates aligned with data points are visualized in Fig. S1, and additional assumptions implicit to the model structure are provided in Table S3. A shiny app is available, to allow exploration of model parameters and emergent phenomena.

IgA-bound Enterobacteriaceae abundance can predict immune phenotypes but only at specific ontogenetic phases

Our ability to develop effective intervention strategies for preventing pathological imprinting hinges on our capacity to accurately predict the trajectory of 'immune education' as the infant gut matures. While there are studies correlating the time course of microbiome composition and IgA-binding patterns with eventual disease susceptibility (*59–61*), the predictive potential of such analyses at different stages of ontogeny remains underexplored. To evaluate the potential of using taxonomic and IgA-seq profiles of fecal samples to predict the trajectory of infant's immune maturation, we systematically explored variation in ecological and immunological outcomes by simulating our model for a total duration of 720 days across a comprehensive range of exclusive breastfeeding (EBF) and mixed feeding (MF) durations. Across 91.1% of all feeding scenarios, the host's gut mucosal immune response matured to tolerate commensals by simultaneously developing masking SIgA against all taxa (Fig. 3A), although the underlying rate of masking varied (Fig. 3B). The infant immune response diverges to a hyperreactive profile—developing neutralizing antibodies against symbiotic commensals rather than masking them—across 3.6%, 9.5%, and 8.4%, of all possible feeding scenarios for *Bifidobacteriaceae*, *Bacteroidaceae*, *Clostridiales* respectively (Fig. 3B), and 8.9% of all possible feeding scenarios for at least one of the commensals (Fig. 3A). These results suggest robust convergence towards a tolerogenic profile across a wide spectrum of feeding patterns, in line with previously reported stereotypical convergence of the systemic immune response (*2*).

Using this generated data, we investigated how well the matured immunological status after 2 years can be predicted using the total and SIgA-bound *Enterobacteriaceae* abundance from fecal samples collected at various stages of ontogeny. We fitted three generalized linear models to this binary outcome (tolerant or hyperreactive) including as covariates i) total, ii) SIgA-bound and iii) total and SIgA-bound *Enterobacteriaceae* abundances per day. The third model significantly improved prediction over both the first (deviance reduction 68501, $p < 2.2e{-}16$) and second model (deviance reduction 58251, $p < 2.2e{-}16$), confirming that combining total and SIgA-bound abundances yields a significant marginal gain in predicting tolerance versus hyperreactivity. Total *Enterobacteriaceae* abundances had the lowest predictive capacity across time (Fig. 3C, 2E), which can be explained by considering the two compensatory mechanisms regulating it: *i)* ecological competition preventing pathogen overgrowth and *ii)* early activation of the endogenous

immune system (earliest at DOL 30) when maternal antibodies are not sufficiently neutralizing. In contrast, SIgA-bound *Enterobacteriaceae* abundance alone during the first 15 days of life provides valuable information regarding the immunological trajectory (Fig. 3D, 3E), as this quantity reflects the efficiency of maternal antibodies (*60*).

Higher levels of SIgA-bound *Enterobacteriaceae* abundance flip from reflecting tolerant to reflecting hyperreactive immunological states over the time course of ontogeny (Fig. 3D). Lower SIgA-bound *Enterobacteriaceae* observed for hyperreactive phenotypes during early phases of ontogeny are indicative of insufficient *Enterobacteriaceae* neutralization by maternal antibodies. As the host matures, this pattern alters, with the hyperreactive phenotype exhibiting an increase in SIgA-bound *Enterobacteriaceae* abundance relative to the tolerant state, echoing the enrichment in SIgA-coated *Enterobacteriaceae* in adults with IBD-like phenotypes (*62*, *63*). Interestingly, samples collected during the transition from exclusively maternal to exclusively endogenous SIgA presence in the gut lumen provided the least information regarding the immunological trajectory of the host, demonstrating non-monotonic temporal patterns in the predictive value of 16S rRNA and IgA-seq analyses.

Fig 3. Immunological outcomes and *Enterobacteriaceae* abundances for the first 2 years of life for different combinations of exclusive breastfeeding (EBF) and mixed feeding (MF) durations. (A) Immune functionality across a spectrum of EBF and MF durations, where 'tolerant' corresponds to developing non-neutralizing but masking antibodies against all symbiotic commensals; and 'hyperreactive' otherwise. EBF and MF durations vary from 0 to 180 and to 300 days respectively **(B)** Distribution of masking (M) and neutralizing (N) rates of endogenous SIgA at DOL 720. Distribution of **(C)** total and **(D)** SIgA-bound *Enterobacteriaceae* abundances over the first two years of life segregated by tolerant and hyperreactive states. The dataset is balanced for each time point to reflect the typical sample size of recent cohort studies (*64*). A Wilcoxon rank-sum test is used to calculate the effect size (rank-biserial correlation from -1 to +1; values closer to ±1 indicating stronger associations) and the significance (*** for $p < 0.001$, ** for $p < 0.01$, * for $p < 0.05$, . for $p < 0.1$, and ns for $p \geq 0.1$), displayed at the top of each boxplot. **(E)** Mean predictive power of *Enterobacteriaceae* abundance segregated by data types. Predictive power of 0 indicates no better prediction of a tolerogenic/hyperreactive phenotype than random chance; 1 indicates perfect accuracy. Shaded areas represent the interquartile range covering the 25th to 75th percentiles. DOL: Day of life.

Ecology versus the microenvironment: what carries the information?

An enduring mystery in mucosal immunology is how postnatal influences on the immune system persist into adulthood, even though many core cell types are not present when these initial influences occur (*11*). One way to test whether this persistent influence is also reflected in our model is to investigate the impact of EBF duration on affinity maturation: antigenic sampling is delayed if M cells remain immature as a result of persistently high levels of mSIgA, and this postpones the activation of B and T cells until the end of this period. Thus, we expect the influence of the EBF period – if any – to rapidly fade unless other components of the system have enduring effects, since mSIgA has a strict half-life and the delivery of HMOs ceases instantaneously with the cessation of breastfeeding (Fig. S2A).

We first investigate the differential impacts of EBF and MF durations in determining endogenous SIgA (eSIgA) affinity across a comprehensive range of EBF and MF durations. Given the presence of pre-weaning commensals (*Bifidobacteriaceae*, *Enterobacteriaceae*) in the gut lumen during both EBF and MF, eSIgA affinity maturation towards these taxa will be influenced by the duration of both feeding practices. Conversely, a significant role for EBF duration in the affinity maturation against post-weaning commensals (*Bacteroidaceae*, *Clostridiales*) would align with the persistent postnatal influences described above. As hypothesized, immune responses against pre-weaning and post-weaning commensals are impacted by EBF and MF durations differently shown by a predictor importance analysis (Fig 4A), where we compare the relative importance of EBF and MF durations in explaining the variance in eSIgA affinities. Recapitulating the presented dilemma, the influence of EBF duration remains prominent in explaining the variance in eSIgA affinities against post-weaning commensals, despite MF duration being 1.5 times more influential. This influence is more prominent when EBF is followed by Exclusive Complementary Feeding (ECF) without any intervening MF period, which makes it challenging for the host to develop tolerance against post-weaning commensals (Fig. S2B).

Although mSIgA itself may not persist, its initial modifications may continue to shape both the ecology and the inflammatory tone of the microenvironment; similarly, HMOs may have shaped the ecology by selecting for bacteria based on their metabolic capacity, leading to a sustained temporal influence. Therefore, we compare the impact of pre-weaning commensal abundance and the microenvironmental stimulation on eSIgA affinities for post-weaning commensals to identify which components of our system carry information from the EBF period onward. We employed the Random Forests method to handle correlation between microbial abundances and microenvironmental stimulation. Intriguingly, our findings reveal a significantly greater influence of the microenvironmental stimulation over taxonomic composition (Fig. 4B). These findings suggest that the conditions and signals within the gut microenvironment—such as epithelial permeability and the tolerogenic bias of follicular T cells which are directly modulated by the microenvironmental stimulation in our model—play a pivotal role in shaping the infant's immune response relative to the presence or abundance of specific bacterial taxa. Moreover, these results indicate that breastfeeding's influence extends beyond immediate transmission of HMOs, antibodies, and microbes, conditioning the gut microenvironment in ways that produce effects outlasting the transient presence of such factors.

Fig 4. Impact of different feeding durations and environmental variables in determining the matured eSIgA values for key taxa. (**A**) Relative importance analysis in determining the eSIgA values for key taxa, comparing the impact of EBF and MF durations. (**B**) Relative importance analysis in determining the eSIgA values for post-weaning commensals, comparing the impact of the abundances of pre-weaning commensals and microenvironmental stimulation. Due to the similarity of results for *Bacteroidaceae* and *Clostridiales*, average values are provided in (**A**) and (**B**). Directionality of the importance values are determined using the directionality of the correlation between the outcome variable and the predictors.

Maternal protection against pathogens depends on mother's immunological profile

Despite the well-established health benefits of breastfeeding, it presents a complex decision-making process for mothers experiencing immune dysregulation (*65*). While epigenetic regulation

of various immune cells, including dendritic cells, stromal cells, T cells, alveolar macrophages, and epithelial cells, has been implicated in this persistence (*12*, *59*), other mechanisms may also play a role. By meticulously eliminating other possibilities such as genetic and epigenetic factors, microbiota variation, and differences in milk-derived metabolites, recent work by Ramanan *et al.* demonstrated that the primary maternal factor allowing for transgenerational immune priming in mice is vertical transmission of maternal IgA (*11*). This observation opens up the possibility of transmission of inflammatory phenotypes via maternal IgA, as well as tolerogenic ones. Indeed, evidence suggests that Inflammatory Bowel Disease (IBD) is significantly more prevalent in the offspring of IBD mothers, who tend to produce breastmilk with higher inflammatory potential and lower antibody levels (*66–68*); and inadequate levels of maternal SIgA to food allergens have been linked to infantile allergic diseases in breastfed infants, showing a stronger correlation with symptoms than parental atopic history (*69*). However, breastmilk's influence goes beyond immune priming and antibody transfer, also enhancing colonization resistance against pathogens through nutritional and microbial support. Therefore, it is not immediately apparent under what circumstances the benefits of breastfeeding outweigh its risk of inflammatory priming.

To address this question, we compare five different feeding scenarios with varying maternal immunological profiles : *i)* control group recapitulating the scenario in Fig. 2, *ii)* excessive targeting of commensal bacteria by maternal antibodies (high affinity mSIgA against *Bifidobacteriaceae*, *Bacteroidaceae*, and *Clostridiales*), mimicking an IBD-like phenotype (*70*), *iii)* breastmilk devoid of mIgA but supplying HMOs and *Bifidobacteriaceae*, indicative of a maternal SIgA deficiency, *iv)* exclusive complementary feeding (ECF), with *Bifidobacteriaceae* introduced at a lower inoculum size and delayed introduction of post-weaning taxa (*Bacteroidaceae* and *Clostridiales*), mimicking a scenario with no breastfeeding, and *v)* ECF with *Bacteroidaceae* and *Clostridiales* transfer from the start, mimicking additional probiotic supplementation. In all scenarios, *Enterobacteriaceae* are introduced at the same inoculum size to assess the balance between pathogenic and symbiotic commensals (*Bifidobacteriaceae*, *Bacteroidaceae*, and *Clostridiales*) during the first 30 days of life, a critical period when the infant is particularly vulnerable to pathogen overgrowth.

In all scenarios apart from the control, inflammatory phenotypes were transgenerationally transmitted: excessive commensal targeting by mSIgA led to hyperreactive endogenous SIgA maturation in the infant, pointing to the importance of SIgA masking in immune imprinting (Fig S4). However, scenarios differed in degree of pathogen control, demonstrating two opposite trends (Fig. 5). ECF with probiotic supplementation (Fig. 5A, 5C; dashed light blue line) and SIgA-deficient breastfeeding (Fig. 5A, 5C; dashed orange line) initially showed an increase in the pathogenic-to-symbiotic commensal ratio followed by a decrease that brought them to a similar level by day 10, echoing their similarity in leading to NEC susceptibility (*60*). SIgA-deficient breastfeeding had an initial advantage suggesting that HMOs and *Bifidobacteriaceae* can partially compensate for the lack of mSIgA by directly modulating the gut microbiome composition and function. In contrast, ECF without probiotic supplementation (Fig. 5A, 5C; dashed yellow line) demonstrated a ten fold increase in the pathogenic-to-symbiotic commensal ratio, emphasizing the importance of a diverse microbiome in providing defense against pathogens. Interestingly, the worst outcomes were observed with the IBD-like phenotype (Fig. 5 dotted red line), even though it provides neutralizing antibodies against the pathogenic commensal *Enterobacteriaceae*. The transfer of hyperreactive mSIgA against symbiotic commensals not only led to inflammatory

imprinting but allowed for excessive *Enterobacteriaceae* overgrowth by neutralizing *Bifidobacteriaceae*, leading to significantly higher relative abundances of SIgA-bound *Enterobacteriaceae* in the fecal samples compared to the control group (Fig. 5B).

Fig 5. Comparison of various breastfeeding scenarios and their impact on *Enterobacteriaceae* growth. Temporal progression of (**A**) Pathogenic-to-symbiotic commensal ratio in the gut lumen, (**B**) relative SIgA-bound (SIgA+) *Enterobacteriaceae* abundance in fecal samples, and (**C**) relative SIgA-free (SIgA-) *Enterobacteriaceae* abundance in fecal samples of the infant for the first 30 days of life (DOL) for four different breastfeeding scenarios : Control, breastmilk (BM) with hyperreactive mSIgA, mSIgA deficient BM, and only exclusive complementary feeding (ECF).

Discussion

'Immune education' has proved a powerful concept in understanding the long shadow of early life on health outcomes. Yet, to date, the concept has remained largely qualitative, informally encompassing a wide range of individual components. This diversity and complexity of interactions make integrating these effects by experimental methods alone intractable, quite apart from the question of the degree to which animal models translate to humans. We take the first steps towards formalizing the concept of immune education in the context of gut mucosal SIgA - microbiome interactions, leveraging the power of mathematical frameworks to robustly integrate known feedback loops connecting the ecology and immunology of the gut, and inferring core parameters using data on early life immunity and microbial community dynamics.

Our analysis adds a predictive dimension to the interpretation of gut microbial composition data. Traditional studies, employing 16S rRNA and IgA-seq techniques on fecal samples, have primarily focused on identifying associations between early-life microbiome compositions and later immune-mediated health outcomes. We add mechanistic understanding as to why and how the IgA coating patterns vary over the course of ontogeny by disentangling the influences of maternal and endogenous antibodies. We show that IgA-seq analysis of fecal samples collected at early and late stages of ontogeny are informative in predicting immunological outcomes (albeit with opposing patterns of SIgA-bound Enterobacteriaceae abundances) but are not informative during the transitional period. These insights call for a discerning approach when using fecal samples from specific postnatal periods to predict disease outcomes.

Mixed feeding duration emerges as a critical determinant of the infant's immune trajectory: the sustained presence of maternal antibodies after weaning enables the introduction of novel antigens to the immune system in a non-inflammatory manner, promoting the development of tolerance. Exclusive breastfeeding still plays a role in shaping the immune response against microbes introduced after its cessation, by its remnant effects on microenvironmental stimulation, or inflammatory tone within the gut lumen and PPs. This emergent phenomenon resolves the perennial question of how information from the early postnatal phase propagates to further stages of immune development: our model indicates that the information is not specifically carried by either symbiotic or pathogenic taxa, but, rather, results from the cumulative capacity of their SIgA-coated and uncoated subpopulations to modulate the inflammatory tone of the microenvironment.

Beyond these fundamental insights, our analysis systematically explores the applied question of how to approach breastfeeding when maternal immunological conditions such as allergies, IBD, or SIgA deficiency affect breastmilk composition. Both lack of SIgA in breastmilk, or excessive targeting of symbiotic commensals by SIgA can lead to inflammatory imprinting in the infant, indicating that maternal SIgA is the main mechanism driving the transgenerational transmission of immunological phenotypes. Although transfer of *Bifidobacteriaceae* and HMOs cannot entirely compensate for maternal SIgA's role in tolerogenic imprinting, it can partially compensate for the absence of SIgA in pathogen control. However, these benefits do not negate the risk of inflammatory imprinting when maternal SIgA is hyperreactive, disrupting colonization resistance by excessive targeting of symbiotic commensals.

This finding, combined with the influence of microenvironmental stimulation on endogenous SIgA affinity maturation suggests the possibility of alternative therapeutic approaches for promoting tolerogenic imprinting when maternal immunity is hyperreactive or insufficient. Our results suggest that interventions could focus on strategies to modulate gut inflammation beyond probiotic supplementation. Further strategies might include targeting Toll-like receptors such as TLR4, which has been proposed to tackle preterm birth and fetal inflammatory injury (*71*). Preliminary results from our model indicate the potential effectiveness of this intervention (Fig. S4), suggesting that tolerance could be induced by reducing the inflammatory potential of Gram-negative commensals, similar to the effects observed with TLR4 antagonists. However, a complete exploration of such a scenario would necessitate expanding our model to explicitly incorporate cellular immunity.

Our blend of mechanistic and semi-mechanistic modeling formally merging multiple data sources to recapitulate out of sample immunological and ecological trajectories represents a first step towards an integrated model of 'immune education'. Given the immense complexity and dynamism of early life, our model is necessarily a simplification, tailored to the range of data currently available, and excluding factors such as innate immune mechanisms, changes in gut permeability, and strain-specific shifts in the gut microbiome. While qualitative conclusions are likely to be robust to these features, quantitative predictions may change (see supplementary material, section *Model Limitations*). Novel experimental and measurement approaches are likely to open the way to evaluating the role of such processes, as well as providing a path to evaluating our underlying mechanistic assumptions (see supplementary material, section *Experimental Model Validation*). Our quantitative analysis yields both fundamental insights into pressing questions in the field and has therapeutic implications relevant in the context of a growing global burden of immune-mediated diseases.

References and Notes:

1. J.-F. Bach, The effect of infections on susceptibility to autoimmune and allergic diseases. *N. Engl. J. Med.* **347**, 911–920 (2002).

2. A. Olin, E. Henckel, Y. Chen, T. Lakshmikanth, C. Pou, J. Mikes, A. Gustafsson, A. K. Bernhardsson, C. Zhang, K. Bohlin, P. Brodin, Stereotypic Immune System Development in Newborn Children. *Cell* **174**, 1277-1292.e14 (2018).

3. H. Steininger, J. Moltzau-Anderson, S. V. Lynch, Contributions of the early-life microbiome to childhood atopy and asthma development. *Semin. Immunol.* **69**, 101795 (2023).

4. I. Laforest-Lapointe, M.-C. Arrieta, Patterns of Early-Life Gut Microbial Colonization during Human Immune Development: An Ecological Perspective. *Front. Immunol.* **8** (2017).

5. L. Hall, Faculty Opinions recommendation of Temporal development of the gut microbiome in early childhood from the TEDDY study. *Fac. Opin. – Post-Publ. Peer Rev. Biomed. Lit.*, doi: 10.3410/f.734278907.793574624 (2020).

6. L. Zhang, M. Agrawal, S. C. Ng, T. Jess, Early-life exposures and the microbiome: implications for IBD prevention. *Gut*, gutjnl-2023-330002 (2023).

7. K. A. Knoop, J. K. Gustafsson, K. G. McDonald, D. H. Kulkarni, P. E. Coughlin, S. McCrate, D. Kim, C.-S. Hsieh, S. P. Hogan, C. O. Elson, P. I. Tarr, R. D. Newberry, Microbial antigen encounter during a preweaning interval is critical for tolerance to gut bacteria. *Sci. Immunol.* **2** (2017).

8. Y. Belkaid, T. W. Hand, Role of the Microbiota in Immunity and Inflammation. *Cell* **157**, 121–141 (2014).

9. T. Gensollen, S. S. Iyer, D. L. Kasper, R. S. Blumberg, How colonization by microbiota in early life shapes the immune system. *Science* **352**, 539–544 (2016).

10. Z. Al Nabhani, S. Dulauroy, R. Marques, C. Cousu, S. Al Bounny, F. Déjardin, T. Sparwasser, M. Bérard, N. Cerf-Bensussan, G. Eberl, A Weaning Reaction to Microbiota Is Required for Resistance to Immunopathologies in the Adult. *Immunity* **50**, 1276-1288.e5 (2019).

11. D. Ramanan, E. Sefik, S. Galván-Peña, M. Wu, L. Yang, Z. Yang, A. Kostic, T. V. Golovkina, D. L. Kasper, D. Mathis, C. Benoist, An Immunologic Mode of Multigenerational Transmission Governs a Gut Treg Setpoint. *Cell* **181**, 1276-1290.e13 (2020).

12. K. A. Fernandes, A. I. Lim, Maternal-driven immune education in offspring. *Immunol. Rev.*, doi: 10.1111/imr.13315 (2024).

13. Z. Al Nabhani, G. Eberl, Imprinting of the immune system by the microbiota early in life. *Mucosal Immunol.* **13**, 183–189 (2020).

14. N. Torow, T. W. Hand, M. W. Hornef, Programmed and environmental determinants driving neonatal mucosal immune development. *Immunity* **56**, 485–499 (2023).

15. M. Wei, R. Shinkura, Y. Doi, M. Maruya, S. Fagarasan, T. Honjo, Mice carrying a knock-in mutation of Aicda resulting in a defect in somatic hypermutation have impaired gut homeostasis and compromised mucosal defense. *Nat. Immunol.* **12**, 264–270 (2011).

16. E. Michaud, L. Waeckel, R. Gayet, R. Goguyer-Deschaumes, B. Chanut, F. Jospin, K. Bathany, M. Monnoye, C. Genet, A. Prier, C. Tokarski, P. Gérard, X. Roblin, N. Rochereau, S. Paul, Alteration of microbiota antibody-mediated immune selection contributes to dysbiosis in inflammatory bowel diseases. *EMBO Mol. Med.* **14** (2022).

17. O. Pabst, E. Slack, IgA and the intestinal microbiota: the importance of being specific. *Mucosal Immunol.* **13**, 12–21 (2020).

18. O. Pabst, V. Cerovic, M. Hornef, Secretory IgA in the Coordination of Establishment and Maintenance of the Microbiota. *Trends Immunol.* **37**, 287–296 (2016).

19. E. W. Rogier, A. L. Frantz, M. E. Bruno, L. Wedlund, D. A. Cohen, A. J. Stromberg, C. S. Kaetzel, Lessons from mother: Long-term impact of antibodies in breast milk on the gut microbiota and intestinal immune system of breastfed offspring. *Gut Microbes* **5**, 663–668 (2014).

20. N. Tsukuda, K. Yahagi, T. Hara, Y. Watanabe, H. Matsumoto, H. Mori, K. Higashi, H. Tsuji, S. Matsumoto, K. Kurokawa, T. Matsuki, Key bacterial taxa and metabolic pathways affecting gut short-chain fatty acid profiles in early life. *ISME J.* **15**, 2574–2590 (2021).

21. J. D. Planer, Y. Peng, A. L. Kau, L. V. Blanton, I. M. Ndao, P. I. Tarr, B. B. Warner, J. I. Gordon, Development of the gut microbiota and mucosal IgA responses in twins and gnotobiotic mice. *Nature* **534**, 263–266 (2016).

22. K. Pan, The Effects of Feeding Type on the Gut Microbiota of Neonates and Early Infants. *J. Infect. Dis. Case Rep.*, 1–5 (2022).

23. C. Palmer, E. M. Bik, D. B. DiGiulio, D. A. Relman, P. O. Brown, Development of the human infant intestinal microbiota. *PLoS Biol.* **5**, e177 (2007).

24. J. M. Rodríguez, K. Murphy, C. Stanton, R. P. Ross, O. I. Kober, N. Juge, E. Avershina, K. Rudi, A. Narbad, M. C. Jenmalm, J. R. Marchesi, M. C. Collado, The composition of the gut microbiota throughout life, with an emphasis on early life. *Microb. Ecol. Health Dis.* **26** (2015).

25. K. Wambach, J. Riordan, *Breastfeeding and Human Lactation* (Jones & Bartlett Learning, 2016).

26. S. S. N. P. for W. (U.S.) Infants, and Children, C. S. F. P. (U.S.), *Infant Nutrition and Feeding: A Guide for Use in the WIC and CSF Programs* (United States Department of Agriculture, 2009).

27. R. Mannel, P. J. Martens, M. Walker, *Core Curriculum for Lactation Consultant Practice* (Jones & Bartlett Publishers, 2012).

28. M. J. Pichler, C. Yamada, B. Shuoker, C. Alvarez-Silva, A. Gotoh, M. L. Leth, E. Schoof, T. Katoh, M. Sakanaka, T. Katayama, C. Jin, N. G. Karlsson, M. Arumugam, S. Fushinobu, M. Abou Hachem, Butyrate producing colonic Clostridiales metabolise human milk oligosaccharides and cross feed on mucin via conserved pathways. *Nat. Commun.* **11** (2020).

29. J. Wang, C. Chen, Z. Yu, Y. He, Q. Yong, D. S. Newburg, Relative fermentation of oligosaccharides from human milk and plants by gut microbes. *Eur. Food Res. Technol.* **243**, 133–146 (2016).

30. D. A. Sela, Bifidobacterial utilization of human milk oligosaccharides. *Int. J. Food Microbiol.* **149**, 58–64 (2011).

31. H. J. Flint, K. P. Scott, S. H. Duncan, P. Louis, E. Forano, Microbial degradation of complex carbohydrates in the gut. *Gut Microbes* **3**, 289–306 (2012).

32. A. Marcobal, M. Barboza, J. W. Froehlich, D. E. Block, J. B. German, C. B. Lebrilla, D. A. Mills, Consumption of Human Milk Oligosaccharides by Gut-Related Microbes. *J. Agric. Food Chem.* **58**, 5334–5340 (2010).

33. N. Torow, R. Li, T. C. A. Hitch, C. Mingels, S. Al Bounny, N. van Best, E.-L. Stange, B. Simons, T. Maié, L. Rüttger, N. M. K. P. Gubbi, D. A. Abbott, A. Benabid, M. Gadermayr, S. Runge, N. Treichel, D. Merhof, S. P. Rosshart, N. Jehmlich, T. W. Hand, M. von Bergen, F. Heymann, O. Pabst, T. Clavel, F. Tacke, H. Lelouard, I. G. Costa, M. W. Hornef, M cell maturation and cDC activation determine the onset of adaptive immune priming in the

neonatal Peyer's patch. *Immunity* **56**, 1220-1238.e7 (2023).

34. C. Wagner, N. Torow, M. W. Hornef, H. Lelouard, Spatial and temporal key steps in early-life intestinal immune system development and education. *FEBS J.* **289**, 4731–4757 (2021).

35. M.-L. Wernroth, S. Peura, A. M. Hedman, S. Hetty, S. Vicenzi, B. Kennedy, K. Fall, B. Svennblad, E. Andolf, G. Pershagen, others, Development of gut microbiota during the first 2 years of life. *Sci. Rep.* **12**, 9080 (2022).

36. L. A. van der Waaij, F. G. Kroese, A. Visser, G. F. Nelis, B. D. Westerveld, P. L. Jansen, J. O. Hunter, Immunoglobulin coating of faecal bacteria in inflammatory bowel disease. *Eur. J. Gastroenterol. Hepatol.* **16**, 669–674 (2004).

37. M. Pop, A. W. Walker, J. Paulson, B. Lindsay, M. Antonio, M. A. Hossain, J. Oundo, B. Tamboura, V. Mai, I. Astrovskaya, Diarrhea in young children from low-income countries leads to large-scale alterations in intestinal microbiota composition. *Genome Biol.* **15**, 1–12 (2014).

38. D. Ley, J.-L. Desseyn, V. Gouyer, S. Plet, S. Tims, I. Renes, M. Mischke, F. Gottrand, Early life nutrition influences susceptibility to chronic inflammatory colitis in later life. *Sci. Rep.* **9**, 18111 (2019).

39. T. Ohkusa, T. Yoshida, N. Sato, S. Watanabe, H. Tajiri, I. Okayasu, Commensal bacteria can enter colonic epithelial cells and induce proinflammatory cytokine secretion: a possible pathogenic mechanism of ulcerative colitis. *J. Med. Microbiol.* **58**, 535–545 (2009).

40. O. Lakhdari, J. Tap, F. Béguet-Crespel, K. Le Roux, T. de Wouters, A. Cultrone, M. Nepelska, F. Lefèvre, J. Doré, H. M. Blottière, Identification of NF-κB Modulation Capabilities within Human Intestinal Commensal Bacteria. *J. Biomed. Biotechnol.* **2011**, 1–9 (2011).

41. M. Schirmer, S. P. Smeekens, H. Vlamakis, M. Jaeger, M. Oosting, E. A. Franzosa, R. ter Horst, T. Jansen, L. Jacobs, M. J. Bonder, A. Kurilshikov, J. Fu, L. A. B. Joosten, A. Zhernakova, C. Huttenhower, C. Wijmenga, M. G. Netea, R. J. Xavier, Linking the Human Gut Microbiome to Inflammatory Cytokine Production Capacity. *Cell* **167**, 1897 (2016).

42. L. W. Peterson, D. Artis, Intestinal epithelial cells: regulators of barrier function and immune homeostasis. *Nat. Rev. Immunol.* **14**, 141–153 (2014).

43. N. W. Palm, M. R. de Zoete, T. W. Cullen, N. A. Barry, J. Stefanowski, L. Hao, P. H. Degnan, J. Hu, I. Peter, W. Zhang, E. Ruggiero, J. H. Cho, A. L. Goodman, R. A. Flavell, Immunoglobulin A Coating Identifies Colitogenic Bacteria in Inflammatory Bowel Disease. *Cell* **158**, 1000–1010 (2014).

44. K. Donald, C. Petersen, S. E. Turvey, B. B. Finlay, M. B. Azad, Secretory IgA: Linking microbes, maternal health, and infant health through human milk. *Cell Host Microbe* **30**, 650–659 (2022).

45. P. Brandtzaeg, Mucosal immunity: integration between mother and the breast-fed infant. *Vaccine* **21**, 3382–3388 (2003).

46. G. P. Donaldson, M. S. Ladinsky, K. B. Yu, J. G. Sanders, B. B. Yoo, W.-C. Chou, M. E. Conner, A. M. Earl, R. Knight, P. J. Bjorkman, S. K. Mazmanian, Gut microbiota utilize immunoglobulin A for mucosal colonization. *Science* **360**, 795–800 (2018).

47. M. L. Van Tassell, M. J. Miller, Lactobacillus Adhesion to Mucus. *Nutrients* **3**, 613–636 (2011).

48. A. Cerutti, Location, location, location: B-cell differentiation in the gut lamina propria. *Mucosal Immunol.* **1**, 8–10 (2008).

49. J. J. Bunker, A. Bendelac, IgA responses to microbiota. *Immunity* **49**, 211–224 (2018).

50. N. Y. Lycke, M. Bemark, The role of Peyer's patches in synchronizing gut IgA responses. *Front. Immunol.* **3** (2012).
51. D. Artis, Epithelial-cell recognition of commensal bacteria and maintenance of immune homeostasis in the gut. *Nat. Rev. Immunol.* **8**, 411–420 (2008).
52. M. Bemark, D. Angeletti, Know your enemy or find your friend?—Induction of IgA at mucosal surfaces. *Immunol. Rev.* **303**, 83–102 (2021).
53. A. Cerutti, M. Rescigno, The biology of intestinal immunoglobulin A responses. *Immunity* **28**, 740–750 (2008).
54. Y. Cong, T. Feng, K. Fujihashi, T. R. Schoeb, C. O. Elson, A dominant, coordinated T regulatory cell-IgA response to the intestinal microbiota. *Proc. Natl. Acad. Sci.* **106**, 19256–19261 (2009).
55. K. W. Ng, A. Hobbs, C. Wichmann, G. D. Victora, G. P. Donaldson, "B cell responses to the gut microbiota" in *Advances in Immunology* (Elsevier, 2022; http://dx.doi.org/10.1016/bs.ai.2022.08.003), pp. 95–131.
56. B. Pietrzak, K. Tomela, A. Olejnik-Schmidt, A. Mackiewicz, M. Schmidt, Secretory IgA in intestinal mucosal secretions as an adaptive barrier against microbial cells. *Int. J. Mol. Sci.* **21**, 9254 (2020).
57. V. Senn, D. Bassler, R. Choudhury, F. Scholkmann, F. Righini-Grunder, R. N. Vuille-dit-Bille, T. Restin, Microbial Colonization From the Fetus to Early Childhood—A Comprehensive Review. *Front. Cell. Infect. Microbiol.* **10** (2020).
58. Q. Tang, G. Jin, G. Wang, T. Liu, X. Liu, B. Wang, H. Cao, Current Sampling Methods for Gut Microbiota: A Call for More Precise Devices. *Front. Cell. Infect. Microbiol.* **10** (2020).
59. A. Sarkar, J. Y. Yoo, S. Valeria Ozorio Dutra, K. H. Morgan, M. Groer, The Association between Early-Life Gut Microbiota and Long-Term Health and Diseases. *J. Clin. Med.* **10**, 459 (2021).
60. K. P. Gopalakrishna, B. R. Macadangdang, M. B. Rogers, J. T. Tometich, B. A. Firek, R. Baker, J. Ji, A. H. P. Burr, C. Ma, M. Good, M. J. Morowitz, T. W. Hand, Maternal IgA protects against the development of necrotizing enterocolitis in preterm infants. *Nat. Med.* **25**, 1110–1115 (2019).
61. A. D. Kostic, D. Gevers, H. Siljander, T. Vatanen, T. Hyötyläinen, A.-M. Hämäläinen, A. Peet, V. Tillmann, P. Pöhö, I. Mattila, H. Lähdesmäki, E. A. Franzosa, O. Vaarala, M. de Goffau, H. Harmsen, J. Ilonen, S. M. Virtanen, C. B. Clish, M. Orešič, C. Huttenhower, M. Knip, R. J. Xavier, The Dynamics of the Human Infant Gut Microbiome in Development and in Progression toward Type 1 Diabetes. *Cell Host Microbe* **17**, 260–273 (2015).
62. M. Viladomiu, C. Kivolowitz, A. Abdulhamid, B. Dogan, D. Victorio, J. G. Castellanos, V. Woo, F. Teng, N. L. Tran, A. Sczesnak, C. Chai, M. Kim, G. E. Diehl, N. J. Ajami, J. F. Petrosino, X. K. Zhou, S. Schwartzman, L. A. Mandl, M. Abramowitz, V. Jacob, B. Bosworth, A. Steinlauf, E. J. Scherl, H.-J. J. Wu, K. W. Simpson, R. S. Longman, IgA-coated E. coli enriched in Crohn's disease spondyloarthritis promote T H 17-dependent inflammation. *Sci. Transl. Med.* **9** (2017).
63. H. L. DuPont, Z.-D. Jiang, A. S. Alexander, A. W. DuPont, E. L. Brown, Intestinal IgA-Coated Bacteria in Healthy- and Altered-Microbiomes (Dysbiosis) and Predictive Value in Successful Fecal Microbiota Transplantation. *Microorganisms* **11**, 93 (2022).
64. C. J. Stewart, N. J. Ajami, J. L. O'Brien, D. S. Hutchinson, D. P. Smith, M. C. Wong, M. C. Ross, R. E. Lloyd, H. Doddapaneni, G. A. Metcalf, D. Muzny, R. A. Gibbs, T. Vatanen, C. Huttenhower, R. J. Xavier, M. Rewers, W. Hagopian, J. Toppari, A.-G. Ziegler, J.-X. She, B.

Akolkar, A. Lernmark, H. Hyoty, K. Vehik, J. P. Krischer, J. F. Petrosino, Temporal development of the gut microbiome in early childhood from the TEDDY study. *Nature* **562**, 583–588 (2018).

65. E. G. Barrett, Maternal influence in the transmission of asthma susceptibility. *Pulm. Pharmacol. Ther.* **21**, 474–484 (2008).

66. I. Dotan, A. Alper, D. Rachmilewitz, E. Israeli, S. Odes, I. Chermesh, T. Naftali, G. Fraser, A. B.-G. Shitrit, V. Peles, S. Reif, Maternal inflammatory bowel disease has short and long-term effects on the health of their offspring: A multicenter study in Israel. *J. Crohns Colitis* **7**, 542–550 (2013).

67. X. Meng, G. Dunsmore, P. Koleva, Y. Elloumi, R. Y. Wu, R. T. Sutton, L. Ambrosio, N. Hotte, V. Nguyen, K. L. Madsen, L. A. Dieleman, H. Chen, V. Huang, S. Elahi, The Profile of Human Milk Metabolome, Cytokines, and Antibodies in Inflammatory Bowel Diseases Versus Healthy Mothers, and Potential Impact on the Newborn. *J. Crohns Colitis* **13**, 431–441 (2018).

68. J. Torres, J. Hu, A. Seki, C. Eisele, N. Nair, R. Huang, L. Tarassishin, B. Jharap, J. Cote-Daigneault, Q. Mao, I. Mogno, G. J. Britton, M. Uzzan, C.-L. Chen, A. Kornbluth, J. George, P. Legnani, E. Maser, H. Loudon, J. Stone, M. Dubinsky, J. J. Faith, J. C. Clemente, S. Mehandru, J.-F. Colombel, I. Peter, Infants born to mothers with IBD present with altered gut microbiome that transfers abnormalities of the adaptive immune system to germ-free mice. *Gut* **69**, 42–51 (2019).

69. S. MacHtinger, R. Moss, Cow's milk allergy in breast-fed infants: The role of allergen and maternal secretory IgA antibody. *J. Allergy Clin. Immunol.* **77**, 341–347 (1986).

70. E. V. Russler-Germain, S. Rengarajan, C.-S. Hsieh, Antigen-specific regulatory T-cell responses to intestinal microbiota. *Mucosal Immunol.* **10**, 1375–1386 (2017).

71. S. A. Robertson, M. R. Hutchinson, K. C. Rice, P. Chin, L. M. Moldenhauer, M. J. Stark, D. M. Olson, J. A. Keelan, Targeting Toll-like receptor-4 to tackle preterm birth and fetal inflammatory injury. *Clin. Transl. Immunol.* **9** (2020).

72. N. Torow, M. W. Hornef, The Neonatal Window of Opportunity: Setting the Stage for Life-Long Host-Microbial Interaction and Immune Homeostasis. *J. Immunol.* **198**, 557–563 (2017).

73. M. Bemark, M. J. Pitcher, C. Dionisi, J. Spencer, Gut-associated lymphoid tissue: a microbiota-driven hub of B cell immunity. *Trends Immunol.* **45**, 211–223 (2024).

74. H. Sugahara, S. Okai, T. Odamaki, C. B. Wong, K. Kato, E. Mitsuyama, J.-Z. Xiao, R. Shinkura, Decreased Taxon-Specific IgA Response in Relation to the Changes of Gut Microbiota Composition in the Elderly. *Front. Microbiol.* **8** (2017).

75. R. M. May, Stability and Complexity in Model Ecosystems. doi: 10.1515/9780691206912 (2001).

76. V. Bucci, B. Tzen, N. Li, M. Simmons, T. Tanoue, E. Bogart, L. Deng, V. Yeliseyev, M. L. Delaney, Q. Liu, B. Olle, R. R. Stein, K. Honda, L. Bry, G. K. Gerber, MDSINE: Microbial Dynamical Systems INference Engine for microbiome time-series analyses. *Genome Biol.* **17** (2016).

77. R. R. Stein, V. Bucci, N. C. Toussaint, C. G. Buffie, G. Rätsch, E. G. Pamer, C. Sander, J. B. Xavier, Ecological modeling from time-series inference: insight into dynamics and stability of intestinal microbiota. *PLoS Comput. Biol.* **9**, e1003388 (2013).

78. S. Marino, N. T. Baxter, G. B. Huffnagle, J. F. Petrosino, P. D. Schloss, Mathematical modeling of primary succession of murine intestinal microbiota. *Proc. Natl. Acad. Sci.* **111**,

439–444 (2014).

79. M. Molari, K. Eyer, J. Baudry, S. Cocco, R. Monasson, Quantitative modeling of the effect of antigen dosage on B-cell affinity distributions in maturating germinal centers. *Elife* **9**, e55678 (2020).

80. M. J. Thomas, U. Klein, J. Lygeros, M. Rodríguez Martínez, A Probabilistic Model of the Germinal Center Reaction. *Front. Immunol.* **10** (2019).

81. L. Buchauer, H. Wardemann, Calculating germinal centre reactions. *Curr. Opin. Syst. Biol.* **18**, 1–8 (2019).

82. S. Wang, J. Mata-Fink, B. Kriegsman, M. Hanson, D. J. Irvine, H. N. Eisen, D. R. Burton, K. D. Wittrup, M. Kardar, A. K. Chakraborty, Manipulating the selection forces during affinity maturation to generate cross-reactive HIV antibodies. *Cell* **160**, 785–797 (2015).

83. A. Koshkin, U. Herbach, M. R. Martínez, O. Gandrillon, F. Crauste, Stochastic modeling of a gene regulatory network driving B cell development in germinal centers. *Plos One* **19**, e0301022 (2024).

84. A. Pélissier, Y. Akrout, K. Jahn, J. Kuipers, U. Klein, N. Beerenwinkel, M. Rodríguez Martínez, Computational model reveals a stochastic mechanism behind germinal center clonal bursts. *Cells* **9**, 1448 (2020).

85. J. Mikulic, S. Longet, L. Favre, J. Benyacoub, B. Corthesy, Secretory IgA in complex with Lactobacillus rhamnosus potentiates mucosal dendritic cell-mediated Treg cell differentiation via TLR regulatory proteins, RALDH2 and secretion of IL-10 and TGF-β. *Cell. Mol. Immunol.* **14**, 546–556 (2016).

86. C. Atyeo, G. Alter, The multifaceted roles of breast milk antibodies. *Cell* **184**, 1486–1499 (2021).

87. J. C. Onyiah, S. P. Colgan, Cytokine responses and epithelial function in the intestinal mucosa. *Cell. Mol. Life Sci.* **73**, 4203–4212 (2016).

88. J. R. Turner, Intestinal mucosal barrier function in health and disease. *Nat. Rev. Immunol.* **9**, 799–809 (2009).

89. A. Dillon, D. D. Lo, M Cells: Intelligent Engineering of Mucosal Immune Surveillance. *Front. Immunol.* **10** (2019).

90. O. Pabst, New concepts in the generation and functions of IgA. *Nat. Rev. Immunol.* **12**, 821–832 (2012).

91. F. Fransen, E. Zagato, E. Mazzini, B. Fosso, C. Manzari, S. El Aidy, A. Chiavelli, A. M. D'Erchia, M. K. Sethi, O. Pabst, M. Marzano, S. Moretti, L. Romani, G. Penna, G. Pesole, M. Rescigno, BALB/c and C57BL/6 Mice Differ in Polyreactive IgA Abundance, which Impacts the Generation of Antigen-Specific IgA and Microbiota Diversity. *Immunity* **43**, 527–540 (2015).

92. Y. Liu, J. Rhoads, Communication between B-Cells and Microbiota for the Maintenance of Intestinal Homeostasis. *Antibodies* **2**, 535–553 (2013).

93. M. Ding, B. Yang, R. P. Ross, C. Stanton, J. Zhao, H. Zhang, W. Chen, Crosstalk between sIgA-Coated Bacteria in Infant Gut and Early-Life Health. *Trends Microbiol.* **29**, 725–735 (2021).

94. M. A. Berkowska, J.-N. Schickel, C. Grosserichter-Wagener, D. de Ridder, Y. S. Ng, J. J. M. van Dongen, E. Meffre, M. C. van Zelm, Circulating Human CD27−IgA+ Memory B Cells Recognize Bacteria with Polyreactive Igs. *J. Immunol.* **195**, 1417–1426 (2015).

95. K. Iyer, L. Erkert, C. Becker, Know your neighbors: microbial recognition at the intestinal barrier and its implications for gut homeostasis and inflammatory bowel disease. *Front. Cell*

Dev. Biol. **11** (2023).

96. M. Mahapatro, L. Erkert, C. Becker, Cytokine-Mediated Crosstalk between Immune Cells and Epithelial Cells in the Gut. *Cells* **10**, 111 (2021).

97. Y. Goto, H. Kiyono, Epithelial barrier: an interface for the cross-communication between gut flora and immune system. *Immunol. Rev.* **245**, 147–163 (2011).

98. C. Martin-Gallausiaux, L. Marinelli, H. M. Blottière, P. Larraufie, N. Lapaque, SCFA: mechanisms and functional importance in the gut. *Proc. Nutr. Soc.* **80**, 37–49 (2020).

99. Y. Li, H. S. Faden, L. Zhu, The Response of the Gut Microbiota to Dietary Changes in the First Two Years of Life. *Front. Pharmacol.* **11** (2020).

100. M. F. Laursen, Gut Microbiota Development: Influence of Diet from Infancy to Toddlerhood. *Ann. Nutr. Metab.* **77**, 21–34 (2021).

101. O. Appert, A. R. Garcia, R. Frei, C. Roduit, F. Constancias, V. Neuzil-Bunesova, R. Ferstl, J. Zhang, C. Akdis, R. Lauener, C. Lacroix, C. Schwab, Initial butyrate producers during infant gut microbiota development are endospore formers. *Environ. Microbiol.* **22**, 3909–3921 (2020).

102. L. V. Hooper, D. R. Littman, A. J. Macpherson, Interactions Between the Microbiota and the Immune System. *Science* **336**, 1268–1273 (2012).

103. J. Zhu, W. E. Paul, Peripheral CD4+ T-cell differentiation regulated by networks of cytokines and transcription factors. *Immunol. Rev.* **238**, 247–262 (2010).

104. X. Ma, S. Nakayamada, J. Wang, Multi-Source Pathways of T Follicular Helper Cell Differentiation. *Front. Immunol.* **12** (2021).

105. A. Reboldi, J. G. Cyster, Peyer's patches: organizing B-cell responses at the intestinal frontier. *Immunol. Rev.* **271**, 230–245 (2016).

106. Y. Kumamoto, T. Hirai, P. W. Wong, D. H. Kaplan, A. Iwasaki, CD301b+ dendritic cells suppress T follicular helper cells and antibody responses to protein antigens. *eLife* **5** (2016).

107. H.-J. Ko, S.-Y. Chang, Regulation of Intestinal Immune System by Dendritic Cells. *Immune Netw.* **15**, 1 (2015).

108. L. Favre, F. Spertini, B. Corthésy, Secretory IgA Possesses Intrinsic Modulatory Properties Stimulating Mucosal and Systemic Immune Responses. *J. Immunol.* **175**, 2793–2800 (2005).

109. H. Morbach, E. M. Eichhorn, J. G. Liese, H. J. Girschick, Reference values for B cell subpopulations from infancy to adulthood. *Clin. Exp. Immunol.* **162**, 271–279 (2010).

110. K. S. Cashman, S. A. Jenks, M. C. Woodruff, D. Tomar, C. M. Tipton, C. D. Scharer, F. Eun-Hyung Lee, J. M. Boss, I. Sanz, Understanding and measuring human B-cell tolerance and its breakdown in autoimmune disease. *Immunol. Rev.* **292**, 76–89 (2019).

111. C. L. Davis, R. Wahid, F. R. Toapanta, J. K. Simon, M. B. Sztein, D. Levy, Applying Mathematical Tools to Accelerate Vaccine Development: Modeling Shigella Immune Dynamics. *PLoS ONE* **8**, e59465 (2013).

112. J. Tellier, S. L. Nutt, The secret to longevity, plasma cell style. *Nat. Immunol.* **23**, 1507–1508 (2022).

113. C.-A. Siegrist, R. Aspinall, B-cell responses to vaccination at the extremes of age. *Nat. Rev. Immunol.* **9**, 185–194 (2009).

114. E. C. Semmes, J.-L. Chen, R. Goswami, T. D. Burt, S. R. Permar, G. G. Fouda, Understanding Early-Life Adaptive Immunity to Guide Interventions for Pediatric Health. *Front. Immunol.* **11** (2021).

115. C. E. Gustafson, D. Higbee, A. R. Yeckes, C. C. Wilson, E. F. De Zoeten, P. Jedlicka, E.

N. Janoff, Limited expression of APRIL and its receptors prior to intestinal IgA plasma cell development during human infancy. *Mucosal Immunol.* **7**, 467–477 (2014).

116. S. Trend, T. Strunk, M. L. Lloyd, C. H. Kok, J. Metcalfe, D. T. Geddes, C. T. Lai, P. Richmond, D. A. Doherty, K. Simmer, A. Currie, Levels of innate immune factors in preterm and term mothers' breast milk during the 1st month postpartum. *Br. J. Nutr.* **115**, 1178–1193 (2016).

117. E. Goonatilleke, J. Huang, G. Xu, L. Wu, J. T. Smilowitz, J. B. German, C. B. Lebrilla, Human Milk Proteins and Their Glycosylation Exhibit Quantitative Dynamic Variations during Lactation. *J. Nutr.* **149**, 1317–1325 (2019).

118. J. Lis-Kuberka, M. Berghausen-Mazur, M. Orczyk-Pawiłowicz, Lactoferrin and Immunoglobulin Concentrations in Milk of Gestational Diabetic Mothers. *Nutrients* **13**, 818 (2021).

119. Stan Development Team. (2020). RStan: the R interface to Stan. http://mc-stan.org/.

120. C. M. Bishop, *Pattern Recognition and Machine Learning* (Springer, 2016).

121. Q. S. Damaceno, J. P. Souza, J. R. Nicoli, R. L. Paula, G. B. Assis, H. C. Figueiredo, V. Azevedo, F. S. Martins, Evaluation of Potential Probiotics Isolated from Human Milk and Colostrum. *Probiotics Antimicrob. Proteins* **9**, 371–379 (2017).

122. L. W. Chia, M. Mank, B. Blijenberg, R. S. Bongers, K. van Limpt, H. Wopereis, S. Tims, B. Stahl, C. Belzer, J. Knol, Cross-feeding between Bifidobacterium infantis and Anaerostipes caccae on lactose and human milk oligosaccharides. *Benef. Microbes* **12**, 69–83 (2021).

123. R. Vidanaarachchi, M. Shaw, S.-L. Tang, S. Halgamuge, IMPARO: inferring microbial interactions through parameter optimisation. *BMC Mol. Cell Biol.* **21** (2020).

124. Y. Huang, R. U. Sheth, S. Zhao, L. A. Cohen, K. Dabaghi, T. Moody, Y. Sun, D. Ricaurte, M. Richardson, F. Velez-Cortes, T. Blazejewski, A. Kaufman, C. Ronda, H. H. Wang, High-throughput microbial culturomics using automation and machine learning. *Nat. Biotechnol.* **41**, 1424–1433 (2023).

125. A. S. Weiss, A. G. Burrichter, A. C. Durai Raj, A. von Strempel, C. Meng, K. Kleigrewe, P. C. Münch, L. Rössler, C. Huber, W. Eisenreich, L. M. Jochum, S. Göing, K. Jung, A. Sanchez, B. Stecher, "Exploring the interaction network of a synthetic gut bacterial community" (Cold Spring Harbor Laboratory, 2021); https://doi.org/10.1101/2021.02.25.432904.

126. P. Fernandez-Julia, D. M. Commane, D. van Sinderen, J. Munoz-Munoz, Cross-feeding interactions between human gut commensals belonging to the Bacteroides and Bifidobacterium genera when grown on dietary glycans. *Microbiome Res. Rep.*, doi: 10.20517/mrr.2021.05 (2022).

127. E. V. Khokhlova, V. V. Smeianov, B. A. Efimov, L. I. Kafarskaia, S. I. Pavlova, A. N. Shkoporov, Anti-inflammatory properties of intestinal Bifidobacterium strains isolated from healthy infants. *Microbiol. Immunol.* **56**, 27–39 (2012).

128. H.-J. Shon, Y.-M. Kim, K. S. Kim, J.-O. Choi, S.-H. Cho, S. An, S.-H. Park, Y.-J. Cho, J.-H. Park, S.-U. Seo, J.-Y. Cho, W.-U. Kim, D. Kim, Protective role of colitis in inflammatory arthritis via propionate-producing Bacteroides in the gut. *Front. Immunol.* **14** (2023).

129. H. Sokol, B. Pigneur, L. Watterlot, O. Lakhdari, L. G. Bermúdez-Humarán, J.-J. Gratadoux, S. Blugeon, C. Bridonneau, J.-P. Furet, G. Corthier, C. Grangette, N. Vasquez, P. Pochart, G. Trugnan, G. Thomas, H. M. Blottière, J. Doré, P. Marteau, P. Seksik, P.

Langella, Faecalibacterium prausnitzii   is an anti-inflammatory commensal bacterium identified by gut microbiota analysis of Crohn disease patients. *Proc. Natl. Acad. Sci.* **105**, 16731–16736 (2008).

130.   G. D. Victora, M. C. Nussenzweig, Germinal Centers. *Annu. Rev. Immunol.* **30**, 429–457 (2012).

131.   C. D. C. Allen, T. Okada, J. G. Cyster, Germinal-Center Organization and Cellular Dynamics. *Immunity* **27**, 190–202 (2007).

132.   P. de Sousa-Pereira, J. M. Woof, IgA: Structure, Function, and Developability. *Antibodies* **8**, 57 (2019).

133.   R. B. Silverstein, I. U. Mysorekar, Group therapy on in utero colonization: seeking common truths and a way forward. *Microbiome* **9** (2021).

134.   L. Xiao, F. Zhao, Microbial transmission, colonisation and succession: from pregnancy to infancy. *Gut* **72**, 772–786 (2023).

135.   W. W. Agace, K. D. McCoy, Regionalized Development and Maintenance of the Intestinal Adaptive Immune Landscape. *Immunity* **46**, 532–548 (2017).

136.   K. A. Knoop, R. D. Newberry, Isolated Lymphoid Follicles are Dynamic Reservoirs for the Induction of Intestinal IgA. *Front. Immunol.* **3** (2012).

137.   F. Barone, P. Patel, J. Sanderson, J. Spencer, Gut-associated lymphoid tissue contains the molecular machinery to support T-cell-dependent and T-cell-independent class switch recombination. *Mucosal Immunol.* **2**, 495–503 (2009).

138.   U. M. Mörbe, P. B. Jørgensen, T. M. Fenton, N. von Burg, L. B. Riis, J. Spencer, W. W. Agace, Human gut-associated lymphoid tissues (GALT); diversity, structure, and function. *Mucosal Immunol.* **14**, 793–802 (2021).

139.   L. M. Sollid, R. Iversen, Tango of B cells with T cells in the making of secretory antibodies to gut bacteria. *Nat. Rev. Gastroenterol. Hepatol.* **20**, 120–128 (2022).

140.   D. Gibbons, J. Spencer, Mouse and human intestinal immunity: same ballpark, different players; different rules, same score. *Mucosal Immunol.* **4**, 148–157 (2011).

141.   C. R. Nowosad, L. Mesin, T. B. R. Castro, C. Wichmann, G. P. Donaldson, T. Araki, A. Schiepers, A. A. K. Lockhart, A. M. Bilate, D. Mucida, G. D. Victora, Tunable dynamics of B cell selection in gut germinal centres. *Nature* **588**, 321–326 (2020).

142.   S. Vergani, K. G. Muleta, C. Da Silva, A. Doyle, T. A. Kristiansen, S. Sodini, N. Krausse, G. Montano, K. Kotarsky, J. Nakawesi, H. Åkerstrand, S. Vanhee, S. L. Gupta, D. Bryder, W. W. Agace, K. Lahl, J. Yuan, A self-sustaining layer of early-life-origin B cells drives steady-state IgA responses in the adult gut. *Immunity* **55**, 1829-1842.e6 (2022).

143.   Q. Zhao, C. O. Elson, Adaptive immune education by gut microbiota antigens. *Immunology* **154**, 28–37 (2018).

144.   M. A. Jackson, C. Pearson, N. E. Ilott, K. E. Huus, A. N. Hegazy, J. Webber, B. B. Finlay, A. J. Macpherson, F. Powrie, L. H. Lam, Accurate identification and quantification of commensal microbiota bound by host immunoglobulins. *Microbiome* **9**, 1–22 (2021).

145.   M. Liang, Z. Liwen, S. Jianguo, D. Juan, D. Fei, Z. Yin, W. Changping, C. Jianping, Fecal microbiota transplantation controls progression of experimental autoimmune hepatitis in mice by modulating the TFR/TFH immune imbalance and intestinal microbiota composition. *Front. Immunol.* **12**, 728723 (2021).

146.   J. Spencer, L. M. Sollid, The human intestinal B-cell response. *Mucosal Immunol.* **9**, 1113–1124 (2016).

147.   E. Gullberg, Å. V. Keita, Y. S. Sa'ad, M. Andersson, K. D. Caldwell, J. D. Söderholm, P.

Artursson, Identification of cell adhesion molecules in the human follicle-associated epithelium that improve nanoparticle uptake into the Peyer's patches. *J. Pharmacol. Exp. Ther.* **319**, 632–639 (2006).

148. K. Hase, K. Kawano, T. Nochi, G. S. Pontes, S. Fukuda, M. Ebisawa, K. Kadokura, T. Tobe, Y. Fujimura, S. Kawano, Uptake through glycoprotein 2 of FimH+ bacteria by M cells initiates mucosal immune response. *Nature* **462**, 226–230 (2009).

149. T. Rollenske, S. Burkhalter, L. Muerner, S. von Gunten, J. Lukasiewicz, H. Wardemann, A. J. Macpherson, Parallelism of intestinal secretory IgA shapes functional microbial fitness. *Nature* **598**, 657–661 (2021).

150. A. Goenka, T. R. Kollmann, Development of immunity in early life. *J. Infect.* **71**, S112–S120 (2015).

151. T. Vatanen, A. D. Kostic, E. d'Hennezel, H. Siljander, E. A. Franzosa, M. Yassour, R. Kolde, H. Vlamakis, T. D. Arthur, A.-M. Hämäläinen, Variation in microbiome LPS immunogenicity contributes to autoimmunity in humans. *Cell* **165**, 842–853 (2016).

152. H. Chardin, K. Mercier, C. Frydman, N. Vollmer, Surface plasmon resonance imaging: a method to measure the affinity of the antibodies in allergy diagnosis. *J. Immunol. Methods* **405**, 23–28 (2014).

153. R. B. Schasfoort, J. van Weperen, M. van Amsterdam, J. Parisot, J. Hendriks, M. Koerselman, M. Karperien, A. Mentink, M. Bennink, H. Krabbe, Presence and strength of binding of IgM, IgG and IgA antibodies against SARS-CoV-2 during CoViD-19 infection. *Biosens. Bioelectron.* **183**, 113165 (2021).

154. T. Noy-Porat, R. Alcalay, A. Mechaly, E. Peretz, E. Makdasi, R. Rosenfeld, O. Mazor, Characterization of antibody-antigen interactions using biolayer interferometry. *STAR Protoc.* **2**, 100836 (2021).

155. Y. Huang, Z. Chen, H. Wang, X. Ba, P. Shen, W. Lin, Y. Wang, K. Qin, Y. Huang, S. Tu, Follicular regulatory T cells: a novel target for immunotherapy? *Clin. Transl. Immunol.* **9**, e1106 (2020).

156. J. B. Wing, W. Ise, T. Kurosaki, S. Sakaguchi, Regulatory T cells control antigen-specific expansion of Tfh cell number and humoral immune responses via the coreceptor CTLA-4. *Immunity* **41**, 1013–1025 (2014).

157. D. Zotos, J. M. Coquet, Y. Zhang, A. Light, K. D'Costa, A. Kallies, L. M. Corcoran, D. I. Godfrey, K.-M. Toellner, M. J. Smyth, IL-21 regulates germinal center B cell differentiation and proliferation through a B cell–intrinsic mechanism. *J. Exp. Med.* **207**, 365–378 (2010).

158. M. E. M. El Shikh, R. El Sayed, N. A. R. Aly, E. Prediletto, R. Hands, L. Fossati-Jimack, M. Bombardieri, M. J. Lewis, C. Pitzalis, Follicular dendritic cell differentiation is associated with distinct synovial pathotype signatures in rheumatoid arthritis. *Front. Med.* **9**, 1013660 (2022).

159. K. Haniuda, T. Nojima, D. Kitamura, In vitro-induced germinal center b cell culture system. *Germinal Cent. Methods Protoc.*, 125–133 (2017).

160. X. Huang, W. Yang, S. Yao, A. J. Bilotta, Y. Lu, Z. Zhou, P. Kumar, S. M. Dann, Y. Cong, IL-21 promotes intestinal memory IgA responses. *J. Immunol.* **205**, 1944–1952 (2020).

161. X.-F. Lei, Y. Ohkawara, M. R. Stämpfli, C. Mastruzzo, R. A. Marr, D. Snider, Z. Xing, M. Jordana, Disruption of antigen-induced inflammatory responses in CD40 ligand knockout mice. *J. Clin. Invest.* **101**, 1342–1353 (1998).

162. J. Xu, T. M. Foy, J. D. Laman, E. A. Elliott, J. J. Dunn, T. J. Waldschmidt, J. Elsemore,

R. J. Noelle, R. A. Flavell, Mice deficient for the CD40 ligand. *Immunity* **1**, 423–431 (1994).

163. A. Purwada, S. B. Shah, W. Béguelin, A. August, A. M. Melnick, A. Singh, Ex vivo synthetic immune tissues with T cell signals for differentiating antigen-specific, high affinity germinal center B cells. *Biomaterials* **198**, 27–36 (2019).

164. A. Purwada, M. K. Jaiswal, H. Ahn, T. Nojima, D. Kitamura, A. K. Gaharwar, L. Cerchietti, A. Singh, Ex vivo engineered immune organoids for controlled germinal center reactions. *Biomaterials* **63**, 24–34 (2015).

165. C. Corinaldesi, A. B. Holmes, Q. Shen, E. Grunstein, L. Pasqualucci, R. Dalla-Favera, K. Basso, Tracking immunoglobulin repertoire and transcriptomic changes in germinal center B cells by single-cell analysis. *Front. Immunol.* **12**, 818758 (2022).

166. A. Biram, Z. Shulman, Evaluation of B cell proliferation in vivo by EdU incorporation assay. *Bio-Protoc.* **10**, e3602–e3602 (2020).

Acknowledgments:
We are grateful to Prof. Dr. Emma Slack and Prof. Dr. M.D. Mathias Hornef for their valuable time and insightful discussions, which significantly improved our work.

Funding:
SNSF postdoc.mobility grant P500PB_206889 (BT)
Branco Weiss Fellowship—Society in Science administered by the ETH Zurich and Rutgers Cancer Institute of New Jersey New Investigator Award (AIL)
N/A (CJEM)

Authors contributions:
Conceptualization: BT, AIL, CJEM
Data curation: BT
Methodology: BT, CJEM
Formal Analysis: BT, CJEM
Visualization: BT
Software: BT
Supervision: AIL, CJEM
Writing – original draft: BT, CJEM
Writing – review & editing: BT, AIL, CJEM

Competing interests: The authors declare that they have no competing interests.

Data and materials availability: All data and code used in this work is available at https://github.com/burcutepekule/ONIMTOL under the GNU General Public License v3.0.

Supplementary Materials
Materials and Methods
Supplementary Text
Figs. S1 to S13
Tables S1 to S5
References (72–166)

A Functionality of the Endogenous Immune System After 2 Years (DOL 720)

B Distribution of Masking (M) and Neutralizing (N) Rate of eSIgA After 2 Years (DOL 720)

C Relative Enterobacteriaceae Abundance in Fecal Samples

D Relative IgA−bound Enterobacteriaceae Abundance in Fecal Samples

E Predictive Power of Enterobacteriaceae Abundance over Time

A Pathogenic/Symbiotic commensals, lumen

B SIgA+ Enterobacteriaceae, fecal samples

C SIgA− Enterobacteriaceae, fecal samples

Feeding
— Control
····· Hyperreactive SIgA BM
– – SIgA Deficient BM
–·– Only ECF
— Only ECF + Probiotics

A Average eSIgA Affinity

B eSIgA Concentration

C IgA-secreting plasma cell count

Bacteroidaceae
Bifidobacteriaceae
Clostridiales
Enterobacteriaceae

A Average eSIgA Affinity

B Cumulative Enterobacteriaceae Load in the gut lumen

C Cumulative Enterobacteriaceae Load in PPs