The novel role of Kallistatin in linking metabolic syndromes and cognitive memory deterioration by inducing amyloid-β plaques accumulation and tau protein hyperphosphorylation

Weiwei Qi[1,2#], Yanlan Long[3#], Ziming Li[4#], Zhen Zhao[1], Jinhui Shi[1], Wanting Xie[1], Laijian Wang[4], Yandan Tan[1], Ti Zhou[1], Minting Liang[1], Ping Jiang[5*], Bin Jiang[4*], Xia Yang[1,6*], Guoquan Gao[1,7,8,9*]

[1] Department of Biochemistry and molecular biology, Zhongshan School of Medicine, Sun Yat-sen University, Guangzhou, China

[2] Advanced Medical Technology Center, The First Affiliated Hospital, Zhongshan School of Medicine, Sun Yat-sen University

[3] Guangdong Key Laboratory of Nanomedicine, CAS-HK Joint Lab of Biomaterials, Shenzhen Institute of Advanced Technology (SIAT), Chinese Academy of Sciences (CAS), Shenzhen, China

[4] Guangdong Province Key Laboratory of Brain Function and Disease, School of Medicine, Sun Yat-sen University, Shenzhen, China

[5] Department of Clinical Medical Laboratory, Guangzhou First People Hospital, School of Medicine, South China University of Technology, Guangzhou, China

[6] China Key Laboratory of Tropical Disease Control (Sun Yat-sen University), Ministry of Education, Guangzhou, China

[7] Guangdong Engineering & Technology Research Center for Gene Manipulation and Biomacromolecular Products (Sun Yat-sen University), Guangzhou, China

[8] Guangdong Province Key Laboratory of Brain Function and Disease,

Zhongshan School of Medicine, Sun Yat-sen University, Guangzhou, China

[9] Guangdong Provincial Key Laboratory of Diabetology& Guangzhou Municipal

Key Laboratory of Mechanistic and Translational Obesity Research, Medical Center

for Comprehensive Weight Control, The Third Affiliated Hospital of Sun Yat-sen

University Guangzhou, Guangdong, China

#These authors contributed equally to this study.

*Corresponding author: Gao GQ: gaogq@mail.sysu.edu.cn; Yang X:

yangxia@mail.sysu.edu.cn; Jiang B: jiangb3@mail.sysu.edu.cn; Ping J:

jiangp45@mail2.sysu.edu.cn

Highlights

● Kallistatin-transgenic (KAL-TG) mice defined its cognitive memory impairment

phenotype accompanied by increased Aβ deposition and tau phosphorylation.

● Kallistatin could directly bind to the Notch1 receptor and thereby upregulate

BACE1 expression by inhibiting PPARγ signaling.

● Fenofibrate could alleviate cognitive memory impairment and down-regulate the

serum Kallistatin level.

Abstract

Accumulation of amyloid β (Aβ) peptides and hyperphosphorylated tau proteins

in the hippocampus triggers cognitive memory decline in Alzheimer's disease (AD).

The incidence and mortality of sporadic AD were tightly associated with diabetes and

hyperlipidemia, while the exact linked molecular is uncertain. Here, we reported that

serum Kallistatin concentrations were meaningfully higher in AD patients, with a

higher concentration of fasting blood glucose and triglyceride. In addition, the

constructed Kallistatin-transgenic (KAL-TG) mice defined its cognitive memory

impairment phenotype and lower LTP in hippocampal CA1 neurons accompanied by

increased Aβ deposition and tau phosphorylation. Mechanistically, Kallistatin could

directly bind to the Notch1 receptor and thereby upregulate BACE1 expression by

inhibiting PPARγ signaling, resulting in Aβ cleavage and production. Besides,

Kallistatin could promote the phosphorylation of tau by activating GSK-3β.

Fenofibrate, a hypolipidemic drug, could alleviate cognitive memory impairment by

down-regulating Aβ and tau phosphorylation of KAL-TG mice. Collectively, our data

clarified   a   novel   mechanism   for   Aβ   accumulation   and   tau   protein

hyperphosphorylation regulation by Kallistatin, which might play a crucial role in

linking metabolic syndromes and cognitive memory deterioration, and suggested that

fenofibrate might have the potential for treating metabolism-related AD.

Key words: Alzheimer's disease, Kallistatin, Aβ42, BACE1, Diabetes

Introduction

Alzheimer's disease (AD), the most prevalent irreversible neurodegenerative disorder associated with dementia in elderly individuals, is marked by a gradual decline in cognitive memory. Pathologically, AD is identified by the presence of extracellular amyloid-β (Aβ) plaques and intracellular neurofibrillary tangles (NFTs) [1, 2, 3]. The Aβ cascade and tau protein hyperphosphorylation are the two primary hypotheses concerning AD. According to the Aβ cascade hypothesis, the excessive production of Aβ disrupts normal cellular functions, leading to synaptic dysfunction, neurodegeneration, tau hyperphosphorylation, and neuroinflammation, which ultimately result in memory impairment in individuals with AD and dementia [4, 5]. Aβ peptides are derived from the sequential cleavage of amyloid precursor protein (APP) by β-secretase (β-site APP cleaving enzyme 1, BACE1) and γ-secretase, thus making this cleavage process significant in AD pathology [6, 7]. BACE1 is considered a highly promising therapeutic target. Several potent BACE1 inhibitors have progressed to advanced stages in clinical trials, emphasizing the role of BACE1 in Aβ production [8, 9, 10]. Tau, a microtubule-associated protein, naturally occurs in axons and regulates microtubule dynamics and axonal transport [11]. In AD, tau undergoes a multistep transformation from a natively unfolded monomer to large aggregated forms, such as NFTs, another defining feature of AD [12, 13]. Glycogen synthase kinase-3 (GSK3) is a key kinase involved in the initial steps of tau phosphorylation, with Wnt signaling

being crucial in activating GSK-3β and GSK-3β-mediated tau phosphorylation [14]. The

physiological mechanisms underlying their interaction remain poorly understood.

There is a close relationship between metabolic disorders and cognitive

impairment across the AD spectrum [15, 16]. Nearly 95% of AD patients are categorized

as sporadic patients, whose increasing incidence and mortality are strongly associated

with type 2 diabetes mellitus (T2DM), obesity, and hyperlipidemia [17, 18, 19]. About 37%

of comorbidities between AD and diabetes have been reported in the Alzheimer's

Association Report [20, 21]. As a result of the strong association and shared mechanism

between AD and T2DM, AD has been termed "type 3 diabetes" by some researchers

[22, 23, 24, 25]. Several studies have demonstrated that diabetes confers a 1.6-fold increased

risk of developing dementia [26, 27]. Similarly, central obesity and high body mass index

(BMI) during middle age are associated with an about 3.5-fold increased risk of

dementia later in life [28]. Therefore, controlling blood glucose and lipids is expected to

be a strategy for preventing or moderating cognitive decline during aging.

Nevertheless, the exact link and key associated regulators between metabolic

abnormalities and AD are still unclear.

Kallistatin is a serine proteinase inhibitor previously identified as a tissue

kallikrein-binding protein [29]. It is produced predominantly in the liver and is widely

expressed in body tissues, where it has antiangiogenic, antifibrotic, antioxidative

stress, and antitumor effects [30, 31]. Furthermore, Kallistatin was found to be increased

in patients with obesity, prediabetes, and diabetes [32, 33, 34]. The concentration of

Kallistatin was positively correlated with the triglyceride glucose index [35], which was

proven to be an independent risk factor for dementia [36]. In addition, our previous

study revealed that the concentration of serum Kallistatin in T2DM patients was

significantly increased and further revealed that Kallistatin suppressed wound healing

in T2DM patients by promoting local inflammation, which suggested that Kallistatin

plays a critical role in the progression of T2DM [37]. Furthermore, our recent research

revealed that Kallistatin can cause memory and cognitive dysfunction by disrupting

the glutamate-glutamine cycle[38].

To explore the relationships among T2DM, AD and Kallistatin, we constructed

Kallistatin transgenic (KAL-TG) mice to explore whether Kallistatin could cause

cognitive impairment through the upregulation of Aβ production. Taken together, our

results suggest that a novel regulatory mechanism of Aβ production and tau protein

hyperphosphorylation by Kallistatin is involved in the progression of metabolic

abnormality-related AD.

Results

Kallistatin increases in AD patients and AD model mice

To explore the relevance of AD in T2DM, a GAD disease enrichment analysis

was initially conducted on differentially expressed genes in neurons of T2DM patients

and normal controls, revealing a close relationship between AD and T2DM

(GSE161355) (Fig. S1A). Additionally, PFAM analysis using the DAVID database

identified enrichment of the Serpin family protein domain (Fig. S1B)

(https://david.ncifcrf.gov/). Our previous studies revealed that Kallistatin (serpin

family a member 4) was elevated in the serum of T2DM patients and was associated

with an adverse prognosis of diabetes complications [44]. We collected 11 serum

samples from dementia patients at Sun Yat-sen Memorial Hospital and reported that

the concentration of Kallistatin was greater than that in normal controls[38]. In this study,

56 AD patients and 61 healthy controls were enrolled from four hospitals in

Guangdong Province to further investigate the potential relevance of Kallistatin and

AD. The clinical and biochemical characteristics of the participants are provided in

Tables S1 and S2. In addition, the serum Kallistatin (12.78 ±2.80 μg/mL) content in

patients with AD was greater than that in normal controls (9.78 ±1.93 μg/mL) (Fig.

1A). Similarly, fasting blood glucose (FBG) and triglyceride (TG) levels were greater

in AD patients than in healthy controls (Fig. 1B). We further grouped all the AD

patients according to diabetes status and found that the Kallistatin (13.79 ±3.05

μg/mL) and TG contents were further elevated in AD patients with diabetes (Fig. 1C-

D). Similarly, Kallistatin expression was increased in the hippocampus of the AD

model mouse SAMP8 compared with that in the control mouse SAMR1 (Fig. S1C-D).

Taken together, these results indicate that the Kallistatin concentration is increased in

metabolic abnormality-related AD patients.

Kallistatin impairs cognitive memory in mice

The above experiments demonstrated that Kallistatin was increased in AD

patients and AD model mice. We subsequently generated KAL-TG mice and assessed

their behavioral performance through the Morris water maze (MWM) and Y maze

tests. Notably, the latency to escape the platform was longer, and the number of

platform crossings, percentage of time spent, and spontaneous alternation were

significantly lower in KAL-TG mice than in age-matched WT mice (Fig. 1E-I).

Furthermore, long-term potentiation (LTP) was measured using whole-cell voltage-

clamp recordings of CA1 neurons in acute hippocampal slices from KAL-TG and WT

mice to assess changes in hippocampal synapses. The LTP in KAL-TG mice was

significantly reduced compared to that in WT mice (Fig. 1J). These results showed

that Kallistatin could impair cognitive memory in mice.

Kallistatin promotes Aβ deposition and tau phosphorylation

We evaluated Aβ deposition and tau phosphorylation in these experimental

mouse hippocampal tissues via immunohistochemistry (IHC) and western blotting.

Predictably, the plaque density and tau phosphorylation in KAL-TG mice were much

greater than those in age-matched WT mice (Fig. 2A-C, 3A-D). Consistent with these

results, ELISA detection of the Aβ42 content in hippocampal tissue revealed that Aβ

production was extraordinarily increased in KAL-TG mice compared with WT mice

(Fig. 2D). These results suggested that Kallistatin promoted Aβ deposition and tau

phosphorylation.

Kallistatin positively regulates Aβ generation by promoting β-secretase rather

than γ-secretase

Western blot and ELISA analyses revealed that the Aβ levels in primary

hippocampal neurons (immunofluorescence identified with the neural marker MAP2,

Fig. S2G) infected with the Kallistatin adenovirus were greater than those in the

control groups (Fig. 2E-G), as were those in the HT22 cells (Fig. S2A-C). Amyloid-

beta precursor protein (APP) undergoes proteolytic processing to generate peptide

fragments [45]. β-Secretase (BACE1) and γ-secretase, which are composed of presenilin

1 (PS1), nicastrin, and Pen-2, are crucial enzymes for Aβ generation [46, 47]. We

determined the levels of APP, BACE1, and PS1 in hippocampal tissue. Compared

with those in WT mice, BACE1 protein and mRNA levels were greater in KAL-TG

mice (Fig. 4A-C, S2D), whereas no significant difference in APP, PS1, or α-secretase

(ADAM9, ADAM10, or ADAM17) expression was detected (Fig. 4A, S2E).

Consistent with the above results, the activity of BACE1 increased (Fig. 4D), whereas

PS1 activity did not change (Fig. S2F). Similarly, the expression and activity of

BACE1 were found to be increased in primary mouse neurons and HT22 cells

transfected with Kallistatin adenovirus (Fig. 5A-C, S3A-C), while PS1 expression and

activity remained unchanged (Fig. 5A, 5D, S3A). Additionally, the effect of

Kallistatin was attenuated by the BACE1 inhibitor verubecestat or siBACE1 03,

which was the most effective (Fig. 5E-F, S3D). These results indicate that Kallistatin

can promote Aβ generation through the upregulation of BACE1 expression.

Kallistatin suppresses PPARγ activation to promote BACE1 expression

The transcription factors SP1, YY1, and PPAR reportedly regulate BACE1

expression at the transcriptional level. Among them, PPARγ can downregulate

BACE1 expression [48, 49, 50]. PPARγ decreased in the hippocampal tissue of KAL-TG

mice, as detected by western blot and immunohistochemical analysis (Fig. 5G-J).

However, no significant differences in YY1 or SP1 expression were detected (Fig.

5G). In addition, Kallistatin downregulated the expression of PPARγ in primary

hippocampal neurons and HT22 cells, thus increasing BACE1 and Aβ expression (Fig.

5K, S3E). Treatment with rosiglitazone, a PPARγ agonist, reversed the decrease in

PPARγ caused by Kallistatin (Fig. 5K-L). Predictably, rosiglitazone inhibited the

ability of Kallistatin to promote BACE1 and Aβ (Fig. 5K and Fig. S3E).

Kallistatin promotes Aβ production *via* direct binding to the Notch1 receptor and

activating the Notch1 pathway

Our results indicated that Notch1 was highly expressed in the hippocampal

tissues of KAL-TG mice (Fig. 6A-D). Furthermore, Notch1 expression was

upregulated in primary hippocampal neurons and HT22 cells infected with adenovirus

expressing Kallistatin *in vitro* (Fig. S4A-B). Additionally, Kallistatin could directly

bind to the Notch1 receptor and activate the Notch1 pathway (Fig. 6E-F and Fig. S4C-

D). Treatment with siNotch1 03, which was the most effective (Fig. S4E), to knock

down Notch1 inhibited the effect of Kallistatin on the activation of the Notch1

signaling pathway, leading to the downregulation of HES1, upregulation of PPARγ,

and downregulation of BACE1 and Aβ (Fig. 6G). HES1, an essential downstream

effector of the Notch1 signaling pathway, has been reported to suppress the expression

of *PPARG* (the gene name of PPARγ) in neurons [51, 52]. Similarly, HES1 shRNA 1, the

most effective (Fig. S4F), upregulated PPARγ and decreased the production of

BACE1 and Aβ when neurons were infected with adenovirus to overexpress

Kallistatin (Fig. 6H). These results suggest that Kallistatin promotes Aβ production

via direct binding to the Notch1 receptor and activating the Notch1 pathway.

Kallistatin promotes the phosphorylation of tau by activating the Wnt signaling

pathway.

Glycogen synthase kinase3-β (GSK-3β) is a crucial element in the

phosphorylation of tau [53]. When Wnt signaling is activated, the LRP6 PPPSPxS motif

can directly interact with GSK-3β and phosphorylate it [54]. Consequently, when Wnt

signaling is inhibited, GSK-3β becomes activated and dephosphorylated, allowing

nonphosphorylated GSK-3β to add phosphate groups to serine/threonine residues of

tau [14]. Kallistatin has already been reported as a competitive inhibitor of the canonical

Wnt signaling pathway [55]. Consistent with previous reports, our results demonstrated

that GSK-3β was activated in the hippocampus of KAL-TG mice (Fig. 7A-B).

Moreover, an increase in tau phosphorylation was observed with the activation of

GSK-3β induced by Kallistatin overexpression (Fig. 7C-D, Fig. S5A), which was

reversed by LiCl, an inhibitor of GSK-3β (Fig. 7E-F, Fig. S5B). These findings

confirmed that Kallistatin promoted tau phosphorylation by activating the Wnt

signaling pathway.

Fenofibrate alleviates memory and cognitive impairment in KAL-TG mice

Hyperlipidemia and hyperlipidemia account for the development of AD [56, 57].

Here, a hypolipidemic drug (fenofibrate) and a hypoglycemic drug (rosiglitazone)

were used to treat KAL-TG mice (Fig. 8A). Compared with that of the KAL-TG

group, the behavioral performance of the treated group was improved, as measured by

the MWM and Y-maze tests. The latency to reach the escape platform on the fifth

training day was significantly decreased (Fig. 8B), and the number of platform

crossings (Fig. 8C), percentage of time spent (Fig. 8D), and spontaneous alternation

(Fig. 8F) were significantly greater in the fenofibrate-treated group than in the

rosiglitazone-treated group. Similarly, the path trace heatmap indicated that the mice

in the fenofibrate-treated group stayed in the target quadrant longer (Fig. 8E). In

addition, decreased serum Kallistatin levels, Aβ and BACE1 levels, phosphorylation

of tau, and activation of GSK3β were detected in the fenofibrate-treated KAL-TG

group (Fig. 8G-K). However, there was no significant difference between the

rosiglitazone-treated group and the KAL-TG group (Fig. 8G-K).

Mechanism summary

In patients with metabolic abnormality-related AD, the concentration of

Kallistatin is elevated, which could increase Aβ deposition through the

Notch1/HES1/PPARγ/BACE1 pathway and induce tau hyperphosphorylation by

activating GSK-3β. Consequently, elevated Kallistatin impaired cognitive memory by

inducing Aβ deposition and tau hyperphosphorylation (Fig. S5C).

Discussion

This study demonstrated that Kallistatin is a novel regulator of amyloid-β plaque accumulation, tau protein hyperphosphorylation, and metabolic abnormality-related cognitive memory impairment. It was shown that Kallistatin levels were increased in the serum of patients with AD and diabetes-related AD, as well as in the hippocampus of AD model mice. Additionally, the KAL-TG mice exhibited cognitive memory impairment and lower LTP in hippocampal CA1 neurons, along with increased Aβ deposition and tau phosphorylation. Mechanistically, Kallistatin transcriptionally upregulates BACE1 expression by suppressing the transcriptional repressor PPARγ, leading to Aβ cleavage and production. Most importantly, our studies revealed that Kallistatin could bind directly to the Notch1 receptor and activate the Notch1/HES1 pathway, causing a decrease in PPARγ, overproduction of BACE1, and increased Aβ42 generation. Moreover, Kallistatin can induce tau phosphorylation by activating GSK-3β, which results from the inhibition of LRP6. Finally, prolonged stimulation with high concentrations of Kallistatin impaired cognitive memory in mice. Finally, the hypolipidemic drug fenofibrate decreased Aβ expression, tau phosphorylation, and the serum Kallistatin level in KAL-TG mice, alleviating memory and cognitive impairment. For the first time, these observations establish an association between high Kallistatin levels and metabolic abnormality-related AD and provide a new drug candidate (fenofibrate) for AD patients with metabolic syndromes.

Increasing evidence suggests that diabetes mellitus and AD are closely linked.

About 80% of AD patients are insulin resistant or have T2DM [58]; additionally, T2DM

patients have a higher risk of up to 73% dementia than healthy controls do [26]. In line

with these observations, the process of cognitive decline in T2DM patients appears to

begin in the prediabetic phase of insulin resistance [59, 60]. A GAD disease enrichment

analysis of differentially expressed genes in the neurons of T2DM patients and normal

controls revealed a close relationship between AD and T2DM. Additionally, PFAM

analysis identified an enrichment of the Serpin family protein domain (Fig. S1A, B).

We and other researchers reported that the level of Kallistatin (which belongs to the

serpin family) was increased in T2DM patients [32, 37]. Although the relationship

between Kallistatin and AD has not been reported to date, we speculate that

Kallistatin might be a critical molecule that could establish a relationship between

Alzheimer's disease and diabetes. Our data indicated that AD patients exhibit

metabolic disorders and elevated Kallistatin levels (Fig. 1A-D, S1C-D). Additionally,

KAL-TG mice showed impaired memory, cognitive function, and synaptic plasticity

(Fig. 1E-J). These results suggest that Kallistatin represents a novel connection

between AD and diabetes.

A hallmark of AD is the aggregation of Aβ into amyloid plaques and tau

phosphorylation in patients' brains. Aβ, a small peptide with a high propensity to form

aggregates, is widely believed to be central and initial to the pathogenesis of this

disease [61]. Correspondingly, it was discovered that Kallistatin could lead to Aβ

overproduction through the Notch1/HES1/PPARγ/BACE1 signaling pathway. GSK-

3β, a vital kinase that regulates the process of tau phosphorylation [53], can be activated

by inhibiting the Wnt signaling pathway. Kallistatin has been identified as a

competitive inhibitor of LRP6, the Wnt receptor [55]. Consistent with previous reports,

Kallistatin increased tau phosphorylation by activating GSK-3β. It was demonstrated

for the first time that Kallistatin promoted AD by increasing Aβ production and tau

phosphorylation in the central nervous system.

Previous studies have shown that Notch signaling is closely related to AD. For

example, a NOTCH mutation was reported to cause AD-like pathology [62]. In addition,

the level of Notch1 was found to be increased in AD patients [63]. In addition, the

Notch1/HES1 signaling pathway has been reported to suppress the expression of

PPARγ [64]. Our previous studies demonstrated that Kallistatin can activate Notch1

signaling [37]. Consequently, we detected Notch1 in our animal and cell models. Indeed,

Notch1 was upregulated by Kallistatin (Fig. 6, S4), as was Aβ deposition. Notch

signaling is initiated by receptor□ligand interactions at the cell surface. In mammals,

there are five ligands encoded by JAG1, JAG2, DLL1, DLL3, and DLL4 [65]. Here, our

results revealed that Kallistatin could activate Notch1 by binding directly to it,

indicating that Kallistatin is a new ligand of the Notch1 receptor.

The treatment of AD has always been a prominent and challenging issue in

neurology. Multiple strategies have been proposed to reduce the pathogenicity of Aβ

and tau. Unfortunately, several Aβ-targeted therapies tested in phase III clinical trials

have failed to slow cognitive decline, although they can effectively reduce the Aβ load

[66, 67, 68, 69, 70, 71]. BACE1 inhibitors have not only failed to improve the cognitive

function of AD patients but have also resulted in clinical deterioration and liver

function impairment [72, 73, 74]. Two possible reasons for the failure of clinical trials with

BACE1 inhibitors are: first, the reduction in BACE1 activity could lead to the

accumulation of full-length APP [75]; second, the size of the BACE1 active site is

relatively large, and the use of a small molecule may not be sufficient to occupy the

active site [76]. Consequently, synaptic damage caused by BACE1 inhibitors or their

insufficient effect may lead to the failure of clinical trials. Therapeutic strategies

targeting tau include tau aggregation blockers (TRx0014, TRx0237), antibody vaccine

therapy (e.g., RO7105705, BIIB092), the inhibition of tau phosphorylation (Anavex2-

73), and the use of microtubule stabilizers (Anavex2-73) [77]. Some of these drugs have

been partially discontinued, while others are still undergoing clinical testing and have

shown protective benefits. Nonetheless, several obstacles remain to the

commercialization of tau treatments when they reach maturity.

Because of the failure of clinical trials, some researchers have proposed

alternative options for AD therapeutics to address modifiable risk factors for the

development of AD, such as type 2 diabetes [78, 79, 80]. Previous studies revealed that

Kallistatin is a multifunctional protein strongly associated with diabetes and that a

Kallistatin neutralizing antibody improves diabetic wound healing [44]. This study

demonstrated that Kallistatin induced memory-related cognitive dysfunction by

promoting Aβ deposition and tau phosphorylation. Thus, it is speculated that

increased Kallistatin could be a promising candidate for T2DM-related AD therapy.

PPARγ is a ligand-activated transcription factor and a master modulator of glucose

and lipid metabolism, organelle differentiation, and inflammation [81, 82]. Growing

evidence revealed that PPARγ agonists (rosiglitazone) could rescue memory

impairment of AD model mice [83, 84, 85]. In clinical trials, it is controversial whether

rosiglitazone has a protective effect on memory cognitive function [86, 87, 88]. In this

study, although Aβ expression had a downward trend, the memory and cognition of

KAL-TG mice were unchanged after treatment with rosiglitazone for a month (Fig.8).

This might be caused by insufficient treatment time and the unchanged Kallistatin

level.

Fenofibrate is a fibric acid derivative for clinically lowering blood lipids, mainly

triglycerides [89]. Studies showed that fenofibrate could prevent memory disturbances,

maintain hippocampal neurogenesis, and protect against Parkinson's disease (PD) [90, 91].

Specifically, fenofibrate has a neuroprotective effect on memory impairment induced

by Aβ through targeting α- and β- secretase [92]. Recently, our study proved that the

fenofibrate could repair the disrupted glutamine-glutamate cycle by upregulating

glutamine synthetase, while there is currently no fenofibrate treatment of AD in

clinical trials[38]. In this study, we proved fenofibrate was beneficial for memory and

cognitive impairment of KAL-TG mice. In addition, Aβ, BACE1, phosphorylated tau,

and serum Kallistatin level of KAL-TG mice could be downregulated after the

treatment of fenofibrate. All of these suggested that fenofibrate might be helpful for

metabolic abnormalities-related AD patients. Therefore, fenofibrate administration in

patients with metabolic syndrome play an early role in preventing and treating AD.

In summary, we affirmed that Kallistatin concentrations were increased in

diabetes-related AD patients. In addition, our study demonstrated for the first time that

Kallistatin positively regulated Aβ42 through Notch1/HES1/PPARγ/BACE1, and

increased phosphorylated tau through inhibition of the Wnt signaling pathway.

Kallistatin might play a crucial role in linking diabetes and cognitive memory

deterioration. Moreover, fenofibrate could decrease the serum Kallistatin level,

BACE1, Aβ, and phosphorylated tau of KAL-TG mice, leading to alleviating memory

and cognitive impairment. These findings might provide new insight into AD and

possibly other neurodegenerative disorders.

Materials and methods

Ethics approval and consent to participate

All patients involved in this study gave their informed consent. This study

obtained the institutional review board approval of Medical Ethics of Zhongshan

Medical College No. 072 in 2021 and Animal Experiment Ethics of Zhongshan

Medical College, Approval No.: SYSU-IACUC-2019-B051.

Human Subjects

The study was approved by the experimental ethics committee of Guangdong

Academy of Medical Sciences and Sun Yat-sen University and carried out in strict

accordance with the ethical principles, and each participant was provided written

informed consent before collecting samples. We certify that the study was performed

in accordance with the 1964 declaration of HELSINKI and later amendments. We

collected 61 normal human samples, 56 AD patient samples, of whom 36 normal

human samples were from the Zhongshan City People's Hospital; 14 normal human

samples and 22 AD patient samples were from Zhongshan Third People's Hospital;

11 normal human samples and 14 AD patient samples were from Sun Yat-sen

Memorial Hospital. AD patient was clinically diagnosed according to ICD-10

(International Classification of Diseases) and NINCDS-ADRDA (the National

Institute of Neurological and Communicative Disorders and Stroke and the

Alzheimer's Disease and Related Disorders Association) criteria, and 20 AD patient

samples were clinically diagnosed according to MMSE (Mini-Mental State

Examination), were collected from Guangdong Provincial People's Hospital. All

subjects' Clinical characteristics were presented in Supplementary Tables (Table S1-2).

Experimental Animals and Protocols

All animal experiment procedures were carried out in an environment without

specific pathogens (Specific pathogen free, SPF) with the approval of the Animal

Care and Use Committee of Sun Yat-sen University (approval ID: SYXK 2015-0107).

The wild type mice (WT, C57BL/6) were purchased from the Animal Center of

Guangdong Province (Production license No.: SCXK 2013-0002, Guangzhou, China).

The SAMR1 and SAMP8 mice (7 months old) were purchased from Tianjin

University of Traditional Chinese Medicine (Tianjin, China). Kallistatin transgenic

mice (KAL-TG) were C57BL/6 strain provided by Dr. Jianxing Ma (University of

Oklahoma Health Sciences Center) [39]. The KAL-TG mice genotype was identified by

PCR technology (forward primer: 5'-AGGGAAG-ATTGTGGATTTGG-3', reverse

primer: 5'-ATGAAGATACCAGTGATGCTC-3'). KAL-TG mice aged 6 months were

randomly divided into three groups: control group (KAL-TG), fenofibrate-treated

group (KAL-TG-Feno, 0.3 g/kg/d), and rosiglitazone-treated group (KAL-TG-RSG,

0.005 g/kg/d). Fenofibrate (Sigma-Aldrich, cat. no. F6020) and rosiglitazone

(Selleck, cat. no. S2556) were administered to mice by oral gavage. In three groups,

the serum Kallistatin were examined in the 0 week and 4 week after drug treatment

from the blood taken from mouse orbit. In addition, the Morris water maze and Y-

maze test were performed one week after the second blood collection.

Morris water maze (MWM)

The KAL-TG and WT mice were employed for the Morris water maze test

including the behavioral test, latency experiment (for 6 days), and the probe test (the

7th day). In addition, the MWM was performed as described previously [40]. Mice were

brought into the testing room and handled for 1 day before the training experiment. In

the 6-day training experiment, each mouse was trained with four daily trials. The mice

facing the wall were placed into the maze, exploring the maze from different

directions (east, south, west, and north). This trial was completed as soon as the

mouse found the platform, or 90 s elapsed. If the mice could discover and climb the

submerged platform within 90 s, the system would automatically record the latency

time and path immediately, and then the mouse was guided to and placed on the

submerged platform for extra 20 s. On day 7, the platform was removed, and a probe

test was performed to examine the strength and integrity of the animal spatial memory

24 h after the last testing trial. During the probe test, the mice were gently brought

into the water from the fixed monitoring point, and the mice were allowed to swim for

90 s without the platform. Finally, all of the measured behavioral parameters were

analyzed using SMART software.

Y-maze test

A Y maze test was performed to assess the mice's spatial memory. The Y maze

was separated by 120°, consisting of three identical arms (30 cm long, 7 cm wide, and

15 cm high) made of blue PVC. The mice were placed first in one of the arms, and

over the next 10 minutes, the sequence and number of their entry into the three arms

were monitored. An alternation is defined when a mouse visits three straight arms

(namely, ABC, BCA, or CAB, but not ABA, BAB, or CAC). Spontaneous alternation

(%) $=$ [(number of alternations)/(total number of arms-2)]$\times$ 100.

Electrophysiology

Hippocampal slices (300-400 μm) from KAL-TG and WT mice were cut as

described [41]. Coronal slices from hippocampus (400 μm thick) were prepared from

different age groups KAL-TG mice and their WT littermates using a tissue slicer

(Vibratome 3000; Vibratome) in ice-cold dissection buffer containing the following

(in mM): 212.7 sucrose, 3 KCl, 1.25 NaH_2PO_4, 3 $MgCl_2$, 1 $CaCl_2$, 26 $NaHCO_3$, and

10 dextrose, bubbled with 95% O_2/5% CO_2. The slices were immediately transferred

to ACSF at 35 °C for 30 min before recordings. The recipe of ACSF was similar to the

dissection buffer, except that sucrose was replaced with 124 mM NaCl, and the

concentrations of $MgCl_2$ and $CaCl_2$ were changed to 1 mM and 2 mM, respectively.

All recordings were performed at 28-30 °C. Pyramidal cells in CA1 areas were

identified visually under infrared differential interference contrast optics based on

their pyramidal somata and prominent apical dendrites. Whole-cell was performed

using an Integrated Patch-Clamp Amplifier (Sutter Instrument, Novato, CA, USA)

controlled by Igor 7 software (WaveMetrics, Portland, OR, USA) filtered at 5 kHz

and sampled at 20 kHz. Igor 7 software was also used for acquisition and analysis.

Only cells with series resistance <20 MΩ and input resistance >100 MΩ were studied.

Cells were excluded if input resistance changed >15% or series resistance

changed >10% over the experiment. A concentric bipolar stimulating electrode with a

tip diameter of 125 μm (FHC) was placed in the stratum radiatum. The recording and

stimulating electrode distances were kept at 50-100 μm. Patch pipettes (2-4MΩ) were

filled with the internal solution consisting of the following (in mM): 120 Cs-

methylsulfonate, 10 Na-phosphocreatine, 10 HEPES, 4 ATP, 5 lidocaine N-ethyl

bromide (QX-314), 0.5 GTP; the pH of the solution was 7.2–7.3, and the osmolarity

was 270-285 mOsm.

To induce LTP, a pairing protocol was applied. In brief, conditioning stimulation

consisted of 360 pulses at 2 Hz paired with continuous postsynaptic depolarization

(180 s) to 0 mV. 50 μM picrotoxin was added to the recording bath to suppress

excessive polysynaptic activity, and the concentration of Ca^{2+} and Mg^{2+} was elevated

to 4 mM to reduce the recruitment of polysynaptic responses. A test pulse was

delivered at 0.067 Hz to monitor baseline amplitude for 10 min before and 30 min

following paired stimulation. To calculate LTP, the EPSC amplitude was normalized

to the mean baseline amplitude during 10 min baseline. Potentiation was defined as

the mean normalized EPSC amplitude 25–40 min after paired stimulation.

ELISA

To quantify serum Kallistatin, the collected samples were centrifuged at 4 □ for

10min at 5000 rpm. It was detected using the KBP ELISA kit (#DY1669, R&D

systems, MN, USA) as per the instructions of the manufacturer. The levels of Aβ42 in

brain tissue produced from mouse primary neuron cells and HT22 cells were

measured with a mouse Aβ42 Elisa Kit (27721, IBL, Germany). To measure Aβ42 in

brain tissue, 0.05 g of mouse brain tissues were weighed and homogenized using 2ml

PBS with a protease inhibitor (cocktail, IKM1020, Solarbio). After centrifugalization

at 4 □ for 30 min at 12000 g, the extracts' supernatants were analyzed using the

ELISA method after total protein quantification. To quantify levels of Aβ42 produced

from primary neuron cells, the cell supernatants were ultrafiltrated with an

Ultrafiltration tube (4-kD Millipore), centrifugalization, and testing. Cell homogenate

was prepared in 1ml PBS with cocktail and quantified using the BCA method before

being measured by ELISA.

Immunohistochemistry

Tissue slices were prepared as described before [42]. The sections were incubated

with Aβ (ab201060, Abcam, Cambridge, UK), BACE1 (#5606S, Cell Signaling

Technology, Boston, USA), PPARγ (#2435, Cell Signaling Technology, Boston, USA),

Notch1 (#3608, Cell Signaling Technology, Boston, USA), p-tau S202 (ab108387,

Abcam, Cambridge, UK), p-tau T231(ab151559, Abcam, Cambridge, UK), p-tau

S396 (ab109390, Abcam, Cambridge, UK), tau (ab75714, Abcam, Cambridge, UK)

antibodies overnight at 4°C and then incubated with Alexa Fluor 488-donkey anti-

rabbit IgG (H□+□L) (1:200, Life Technologies, Gaithersburg, MD, USA, #A21208)

for 1h, then incubated with a biotin-conjugated secondary antibody for 30□min,

followed by incubation with DAB for 10□s and hematoxylin staining for 30□s. The

IHC signals were analyzed using ImageJ.

Cell culture experiments

495   HT22 cells were purchased from the Cell Bank of the Chinese Academy of

496   Sciences (Shanghai, China). HT22 cells were cultured and grown to confluence in

497   DMEM supplemented with 10% FBS (Gibco BRL), 100 U/mL penicillin, and 100

498   U/mL streptomycins (Gibco BRL).

Primary culture of hippocampal neurons

Primary neurons were obtained from the hippocampus of C57/BL6J mice at age

1-3 days. Before culturing, the newborn pup was euthanized and dipped into 70%

ethanol for 3 min. First, the infant pup hippocampus was isolated with eye tweezers

observed under the stereomicroscope, and excess soft tissue was removed. Second,

hippocampal tissue in PBS buffer was cut up with scissors gently and blown with a

1ml pipette until it was not visible. Next, the cell suspension was transferred to a 15ml

centrifuge tube and centrifuged at 1000 rpm for 5 min at room temperature. Cell

precipitation was suspended and cultured with 2-3mL primary neural stem cell (NSC)

suspension (Thermo Fisher Scientific, 21103049) in a 37 °C, 5% CO_2 cell incubator

for 3 days, changing half medium every 2 days. After 7 days, the cell suspension was

transferred to a 15ml centrifuge tube, centrifuged, and recultured with neurobasal,

10%FBS, 1:50 B27(Thermo Fisher Scientific, A3582801), and 1:100 bFGF (Thermo

Fisher Scientific, #RP-8626). One day later, the medium was changed to neurobasal

(2%FBS, 1:50 B27, and 1:100 bFGF), culturing for 21 more days. The

immunofluorescence technique was used with the neuron-specific marker (MAP2,

#4542, Cell Signaling Technology, Boston, USA) to determine the purity of neurons.

siRNA, shRNA, and adenovirus transfection

Notch1 siRNA and control siRNA were purchased from RiboBio (Guangzhou,

China). HES1 shRNA and control shRNA were purchased from Qingke (Guangzhou,

China). Green fluorescent protein-adenovirus (Ad-GFP) and Kallistatin-adenovirus

(Ad-KAL) were provided by Dr. Jianxing Ma (University of Oklahoma Health

Sciences Center). According to the manufacturer's instructions, the transfections were

performed at approximately 60% confluency using Lipofectamine®3000 transfection

reagent (Invitrogen) or RNAiMAX. After 24□h, interference confirmation was

conducted using real-time quantitative PCR (RT-qPCR) and Western blot.

RNA isolation and quantitative RT-PCR

Total RNA extraction, reverse transcription of cDNA, and real-time quantitative

PCR were performed as described previously [43]. BACE1 forward:

GGAGCCCTTCTTTGACTCCC; BACE1 reverse: CAATGATCATGCTCCCTCCCA;

ADAM9 forward: GGAAGGCTCCCTACTCTCTGA; ADAM9 reverse: CAATTCC-

AAAACTGGCATTCTCC; ADAM10 forward: ATGGTGTTGCCGACAGTGTTA;

ADAM10 reverse: GTTTGGCACGCTGGTGTTTTT; ADAM17 forward: GGAT-

CTACAGTCTGCGACACA; ADAM17 reverse: TGAAAAGCGTTCGGTACTTGAT;

β-actin forward: GCACTCTTCCAGCTTCCTT; β-actin reverse:

GTTGGCGTACAG-GTCTTTGC.

Western blot

Western blot was performed as described previously [40, 43]. Equal amounts of

protein were subjected to western blot analysis. Blots were probed with antibodies

against Kallistatin (ab187656, Abcam, Cambridge, UK), Aβ (ab201060, Abcam,

Cambridge, UK), Presenilin-1 (ab76083, Abcam, Cambridge, UK), BACE1 (#5606S,

Cell Signaling Technology, Boston, USA), APP (#2452S, Cell Signaling Technology,

Boston, USA), MAP2 (#4542, Cell Signaling Technology, Boston, USA), PPARγ

（#2435, Cell Signaling Technology, Boston, USA), SP1 (#9389, Cell Signaling

Technology, Boston, USA), YY1 (#46395, Cell Signaling Technology, Boston,

USA）, Notch1 (#3608, Cell Signaling Technology, Boston, USA), Hes1 (#11988,

Cell Signaling Technology, Boston, USA), p-tau S202 (ab108387, Abcam,

Cambridge, UK), p-tau T231(ab151559, Abcam, Cambridge, UK), p-tau S396

(ab109390, Abcam, Cambridge, UK), tau (ab75714, Abcam, Cambridge, UK),

GSK3β(#70109S, Cell Signaling Technology, Boston, USA), p-GSK3β Ser9 (#9323,

Cell Signaling Technology, Boston, USA), β-actin (A5441-2ml, Sigma, CA, USA),

Caveolin-1(SZ02-01, Huabio, China), GAPDH (200306-7E4, Zen-bio, China), anti-

Mouse (#PI200, Vector Laboratories, Burlingame, CA, USA), anti-Rabbit (#PI1000,

Vector Laboratories, Burlingame, CA, USA). The signal intensity was quantified

using ImageJ (NIH).

Statistical Analysis

The results are expressed as mean ± SD. Student's *t*-test was applied for

comparisons of parametric data between two groups, and one-way ANOVA followed

by LSD *t*-test was used to compare differences between more than two different

groups (GraphPad Prism software). A *P* value less than 0.05 was considered statistical

significance.

List of abbreviations

Aβ	amyloid β
p-tau	hyperphosphorylated tau
AD	Alzheimer's disease
T2DM	type 2 diabetes mellitus

FBG	fasting blood glucose
TG	triglyceride
KAL-TG	Kallistatin-transgenic
WT	wild type mice
APP	amyloid precursor protein
BACE1	β-site APP cleaving enzyme 1
BMI	body mass index
ICD-10	The International Statistical Classification of Diseases and Related Health Problems 10th Revision
NINCDS-ADRDA	the National Institute of Neurological and Communicative Disorders and Stroke and the Alzheimer's Disease and Related Disorders Association
MMSE	mini-Mental State Examination
KAL-TG-RSG	rosiglitazone-treated group
KAL-TG-Feno	fenofibrate-treated group
MWM	morris water maze
RT-qPCR	real-time quantitative PCR

Declarations

Ethics approval and consent to participate

All patients involved in this study gave their informed consent. This study

obtained the institutional review board approval of Medical Ethics of Zhongshan

Medical College No. 072 in 2021 and Animal Experiment Ethics of Zhongshan

Medical College, Approval No.: SYSU-IACUC-2019-B051.

Consent for publication

Not applicable.

Availability of data and materials

All the data supporting the conclusions of the current study are presented in the

figures and they are available from the corresponding authors upon reasonable request.

There are no restrictions on data availability. Source data are provided with this paper.

Competing interests

The authors declare that they have no competing interests.

Acknowledgements

We are thankful to Zhongshan City People's Hospital, Zhongshan Third People's

Hospital, Guangdong Provincial People's Hospital, and Sun Yat-sen Memorial

Hospital for kindly providing serum samples for the analyses of this manuscript. We

thank Professor Boxing Li for his valuable advice on our study.

Funding

This study was supported by The National Natural Science Foundation of China

(Grants 82070888, 82070882, 82100917, 82273116, 82203661, 81901557, 81870869,

and 32071101); Guangdong Natural Science Fund (Grant 2022A1515012423,

2021A1515010434, 2023A1515010316, and 2023A1515010214); Key Sci-Tech

Research Project of Guangzhou Municipality, China (Grants 202201010820,

202102020955); Key Project of Nature Science Foundation of Guangdong Province,

China (Grant 2019B1515120077); National Key R&D Program of China (Grant

2018YFA0800403); Guangdong Special Support Program for Young Top Scientist

(Grant 201629046); China Postdoctoral Science Foundation (Grant 2021M703679,

2022M713594, BX20220360); Health Commission of Guangdong Province (Grants

A2022161) ; Guangzhou Municipal Science and Technology Bureau (Grants

SL2022A04J01575); Guangdong Key Project in "Development of new tools for

diagnosis and treatment of Autism" (2018B030335001), Research and Development

Plan of Key Areas of Guangzhou Science and Technology Bureau (2020070030001),

Open Research Funds of State Key Laboratory of Ophthalmology (2020KF03);

Guangzhou Key Laboratory for Metabolic Diseases(202102100004); The Science and

Technology Planning Project of Guangdong Province 2023B1212060018.

References

1. Grundke-Iqbal I, Iqbal K, Tung YC, Quinlan M, Wisniewski HM, Binder LI. Abnormal
phosphorylation of the microtubule-associated protein tau (tau) in Alzheimer cytoskeletal pathology.
Proceedings of the National Academy of Sciences of the United States of America 1986, **83**(13)**:**
4913-4917.

2. Haass C, Selkoe DJ. Soluble protein oligomers in neurodegeneration: lessons from the Alzheimer's
amyloid beta-peptide. *Nature Reviews Molecular Cell Biology* 2007, **8**(2)**:** 101-112.

3. Holtzman DM, Morris JC, Goate AM. Alzheimer's Disease: The Challenge of the Second Century.
Science Translational Medicine 2011, **3**(77).

4. Leng F, Edison P. Neuroinflammation and microglial activation in Alzheimer disease: where do we
go from here? *Nature Reviews Neurology* 2021, **17**(3)**:** 157-172.

5. De Strooper B, Karran E. The Cellular Phase of Alzheimer's Disease. *Cell* 2016, **164**(4)**:** 603-615.

6. Tomita T. Molecular mechanism of intramembrane proteolysis by gamma-secretase. *Journal of*
Biochemistry 2014, **156**(4)**:** 195-201.

7. LaFerla FM, Green KN, Oddo S. Intracellular amyloid-beta in Alzheimer's disease. *Nature Reviews*
Neuroscience 2007, **8**(7)**:** 499-509.

8. Das B, Yan R. A Close Look at BACE1 Inhibitors for Alzheimer's Disease Treatment. *Cns Drugs*
2019, **33**(3)**:** 251-263.

9. Crunkhorn, Sarah. Alzheimer disease: BACE1 inhibitor reduces β-amyloid production in humans.
Nature Reviews Drug Discovery 2016, **16**(1)**:** 18-18.

10. Yan R, Vassar R. Targeting the β secretase BACE1 for Alzheimer's disease therapy. *Lancet*
Neurology 2014, **13**(3)**:** 319-329.

11. Wang Y, Mandelkow E. Tau in physiology and pathology. *Nature reviews Neuroscience* 2016, **17**(1)**:**
5-21.

12. Hausrat TJ, Janiesch PC, Breiden P, Lutz D, Hoffmeister-Ullerich S, Hermans-Borgmeyer I, *et al.*
Disruption of tubulin-alpha4a polyglutamylation prevents aggregation of hyper-phosphorylated tau
and microglia activation in mice. *Nature communications* 2022, **13**(1)**:** 4192.

13. Braak E, Braak H, Mandelkow EM. A sequence of cytoskeleton changes related to the formation of
neurofibrillary tangles and neuropil threads. *Acta neuropathologica* 1994, **87**(6)**:** 554-567.

14. Boonen RA, van Tijn P, Zivkovic D. Wnt signaling in Alzheimer's disease: up or down, that is the
question. *Ageing Res Rev* 2009, **8**(2)**:** 71-82.

15. Biessels GJ, Despa F. Cognitive decline and dementia in diabetes mellitus: mechanisms and clinical
implications. *Nature Reviews Endocrinology* 2018, **14**(10)**:** 591-604.

16. Li W, Huang E, Gao S. Type 1 Diabetes Mellitus and Cognitive Impairments: A Systematic Review.
Journal of Alzheimers Disease 2017, **57**(1)**:** 29-36.

17. Cardoso SM, Correia SC, Carvalho C, Moreira PI. Mitochondria in Alzheimer's Disease and
Diabetes-Associated Neurodegeneration: License to Heal! In: Singh H, Sheu SS (eds).
Pharmacology of Mitochondria, vol. 240, 2017, pp 281-308.

18. Schipper HM. Apolipoprotein E: Implications for AD neurobiology, epidemiology and risk
assessment. *Neurobiology of Aging* 2011, **32**(5)**:** 778-790.

19. Folch J, Patraca I, Martinez N, Pedros I, Petrov D, Ettcheto M*, et al.* The role of leptin in the
sporadic form of Alzheimer's disease. Interactions with the adipokines amylin, ghrelin and the
pituitary hormone prolactin. *Life Sciences* 2015, **140:** 19-28.

20. Association As. 2016 Alzheimer's disease facts and figures. *Alzheimer's & Dementia* 2016, **12**(4)**:** 459-509.

21. Baglietto-Vargas D, Shi J, Yaeger DM, Ager R, LaFerla FM. Diabetes and Alzheimer's disease crosstalk. *Neuroscience & Biobehavioral Reviews* 2016, **64:** 272-287.

22. Kandimalla R, Thirumala V, Reddy PH. Is Alzheimer's disease a Type 3 Diabetes? A critical appraisal. *Biochimica Et Biophysica Acta-Molecular Basis of Disease* 2017, **1863**(5)**:** 1078-1089.

23. Zhang Y, Huang N-q, Yan F, Jin H, Zhou S-y, Shi J-s, *et al.* Diabetes mellitus and Alzheimer's disease: GSK-3 beta as a potential link. *Behavioural Brain Research* 2018, **339:** 57-65.

24. Steen E, Terry BM, Rivera EJ, Cannon JL, Neely TR, Tavares R, *et al.* Impaired insulin and insulin-like growth factor expression and signaling mechanisms in Alzheimer's disease - is this type 3 diabetes? *Journal of Alzheimers Disease* 2005, **7**(1)**:** 63-80.

25. Candasamy M, Mohamed Elhassan SA, Kumar Bhattamisra S, Hua WY, Sern LM, Binti Busthamin NA, *et al.* Type 3 diabetes (Alzheimer's disease): new insight for promising therapeutic avenues. *Panminerva Medica* 2020, **62**(3)**:** 155-163.

26. Koekkoek PS, Kappelle LJ, van den Berg E, Rutten GE, Biessels GJ. Cognitive function in patients with diabetes mellitus: guidance for daily care. *The Lancet Neurology* 2015, **14**(3)**:** 329-340.

27. Hildreth KL, Van Pelt RE, Schwartz RS. Obesity, insulin resistance, and Alzheimer's disease. *Obesity (Silver Spring, Md)* 2012, **20**(8)**:** 1549.

28. Anjum I, Fayyaz M, Wajid A, Sohail W, Ali A. Does obesity increase the risk of dementia: a literature review. *Cureus* 2018, **10**(5).

29. Ma C, Luo C, Yin H, Zhang Y, Xiong W, Zhang T, *et al.* Kallistatin inhibits lymphangiogenesis and lymphatic metastasis of gastric cancer by downregulating VEGF-C expression and secretion. *Gastric Cancer* 2018, **21**(4)**:** 617-631.

30. Yang Y, He X, Cheng R, Chen Q, Shan C, Chen L, *et al.* Diabetes-induced upregulation of Kallistatin levels exacerbates diabetic nephropathy via RAS activation. *Faseb Journal* 2020, **34**(6)**:** 8428-8441.

31. Ma C, Yin H, Zhong J, Zhang Y, Luo C, Che D, *et al.* Kallistatin exerts anti-lymphangiogenic effects by inhibiting lymphatic endothelial cell proliferation, migration and tube formation. *International Journal of Oncology* 2017, **50**(6)**:** 2000-2010.

32.El-Asrar MA, Andrawes NG, Ismail EA, Salem SMH. Kallistatin as a marker of microvascular
complications in children and adolescents with type 1 diabetes mellitus: Relation to carotid intima
media thickness. *Vascular Medicine* 2015, **20**(6)**:** 509-517.

33.Gateva A, Assyov Y, Velikova T, Kamenov Z. Increased Kallistatin levels in patients with obesity
and prediabetes compared to normal glucose tolerance. *Endocrine Research* 2017, **42**(2)**:** 163-168.

34.Campbell DJ, Kladis A, Zhang Y, Jenkins AJ, Prior DL, Yii M*, et al.* Increased tissue kallikrein
levels in type 2 diabetes. *Diabetologia* 2010, **53**(4)**:** 779-785.

35.Nowicki GJ, Ślusarska B, Polak M, Naylor K, Kocki T. Relationship between Serum Kallistatin and
Afamin and Anthropometric Factors Associated with Obesity and of Being Overweight in Patients
after Myocardial Infarction and without Myocardial Infarction. *Journal of clinical medicine* 2021,
10(24).

36.Hong S, Han K, Park CY. The insulin resistance by triglyceride glucose index and risk for dementia:
population-based study. *Alzheimer's research & therapy* 2021, **13**(1)**:** 9.

37.Feng J, Dong C, Long YL, Mai LF, Ren M, Li LY*, et al.* Elevated Kallikrein-binding protein in
diabetes impairs wound healing through inducing macrophage M1 polarization. *Cell
Communication and Signaling* 2019, **17**.

38.Long Y, Zhao Z, Xie W, Shi J, Yang F, Zhu D*, et al.* Kallistatin leads to cognition impairment via
downregulating glutamine synthetase. *Pharmacol Res* 2024, **202:** 107145.

39.McBride JD, Jenkins AJ, Liu XC, Zhang B, Lee K, Berry WL*, et al.* Elevated Circulation Levels of
an Antiangiogenic SERPIN in Patients with Diabetic Microvascular Complications Impair Wound
Healing through Suppression of Wnt Signaling. *Journal of Investigative Dermatology* 2014, **134**(6)**:**
1725-1734.

40.Huang M, Qi WW, Fang SH, Jiang P, Yang C, Mo YS*, et al.* Pigment Epithelium-Derived Factor
Plays a Role in Alzheimer's Disease by Negatively Regulating A beta 42. *Neurotherapeutics* 2018,
15(3)**:** 728-741.

41.Guo D, Peng Y, Wang L, Sun X, Wang X, Liang C*, et al.* Autism-like social deficit generated by
Dock4 deficiency is rescued by restoration of Rac1 activity and NMDA receptor function. *Mol
Psychiatry* 2021, **26**(5)**:** 1505-1519.

42.Li C, Huang Z, Zhu L, Yu X, Gao T, Feng J*, et al.* The contrary intracellular and extracellular
functions of PEDF in HCC development. *Cell Death & Disease* 2019, **10**.

43.Jiang P, Huang M, Qi WW, Wang FH, Yang TY, Gao TX*, et al.* FUBP1 promotes neuroblastoma

proliferation via enhancing glycolysis-a new possible marker of malignancy for neuroblastoma. *Journal of Experimental & Clinical Cancer Research* 2019, **38**(1).

44. Feng J, Dong C, Long Y, Mai L, Ren M, Li L*, et al.* Elevated Kallikrein-binding protein in diabetes impairs wound healing through inducing macrophage M1 polarization. *Cell Communication and Signaling* 2019, **17**.

45. De Strooper B, Annaert W. Proteolytic processing and cell biological functions of the amyloid precursor protein. *Journal of Cell Science* 2000, **113**(11)**:** 1857-1870.

46. LaFerla FM, Oddo S. Alzheimer's disease: A beta, tau and synaptic dysfunction. *Trends in Molecular Medicine* 2005, **11**(4)**:** 170-176.

47. Scheuner D, Eckman C, Jensen M, Song X, Citron M, Suzuki N*, et al.* Secreted amyloid beta-protein similar to that in the senile plaques of Alzheimer's disease is increased in vivo by the presenilin 1 and 2 and APP mutations linked to familial Alzheimer's disease. *Nature Medicine* 1996, **2**(8)**:** 864-870.

48. Christensen MA, Zhou WH, Qing H, Lehman A, Philipsen S, Song WH. Transcriptional regulation of BACE1, the beta-amyloid precursor protein beta-secretase, by Sp1. *Molecular and Cellular Biology* 2004, **24**(2)**:** 865-874.

49. Lin N, Chen L-m, Pan X-d, Zhu Y-g, Zhang J, Shi Y-q*, et al.* Tripchlorolide Attenuates beta-amyloid Generation via Suppressing PPAR gamma-Regulated BACE1 Activity in N2a/APP695 Cells. *Molecular Neurobiology* 2016, **53**(9)**:** 6397-6406.

50. Nowak K, Lange-Dohna C, Zeitschel U, Gunther A, Luscher B, Robitzki A*, et al.* The transcription factor Yin Yang 1 is an activator of BACE1 expression. *Journal of Neurochemistry* 2006, **96**(6)**:** 1696-1707.

51. Maniati E, Bossard M, Cook N, Candido JB, Emami-Shahri N, Nedospasov SA*, et al.* Crosstalk between the canonical NF-κB and Notch signaling pathways inhibits Pparγ expression and promotes pancreatic cancer progression in mice. *The Journal of clinical investigation* 2011, **121**(12)**:** 4685-4699.

52. Herzig S, Hedrick S, Morantte I, Koo SH, Galimi F, Montminy M. CREB controls hepatic lipid metabolism through nuclear hormone receptor PPAR-gamma. *Nature* 2003, **426**(6963)**:** 190-193.

53. Kanno T, Tsuchiya A, Tanaka A, Nishizaki T. Combination of PKCε Activation and PTP1B Inhibition Effectively Suppresses Aβ-Induced GSK-3β Activation and Tau Phosphorylation. *Mol Neurobiol* 2016, **53**(7)**:** 4787-4797.

54.Piao S, Lee SH, Kim H, Yum S, Stamos JL, Xu Y, *et al.* Direct inhibition of GSK3beta by the
phosphorylated cytoplasmic domain of LRP6 in Wnt/beta-catenin signaling. *PLoS One* 2008, **3**(12)**:**
e4046.

55.Liu X, Zhang B, McBride JD, Zhou K, Lee K, Zhou Y, *et al.* Antiangiogenic and
antineuroinflammatory effects of Kallistatin through interactions with the canonical Wnt pathway.
Diabetes 2013, **62**(12)**:** 4228-4238.

56.Kubis-Kubiak A, Wiatrak B, Piwowar A. Hyper-glycemia and insulinemia induce morphological
changes and modulate secretion of S100B, S100A8, amyloid β 1-40 and amyloid β 1-42, in a model
of human dopaminergic neurons. *Biomedicine & pharmacotherapy = Biomedecine &*
pharmacotherapie 2022, **156:** 113869.

57.Rivas-Domínguez A, Mohamed-Mohamed H, Jimenez-Palomares M, García-Morales V, Martinez-
Lopez L, Orta ML, *et al.* Metabolic Disturbance of High-Saturated Fatty Acid Diet in Cognitive
Preservation. *International journal of molecular sciences* 2023, **24**(9).

58.Suzanne M. Type 3 diabetes is sporadic Alzheimer′ s disease: mini-review. *European*
Neuropsychopharmacology 2014, **24**(12)**:** 1954-1960.

59.Biessels GJ, Reagan LP. Hippocampal insulin resistance and cognitive dysfunction. *Nature reviews*
Neuroscience 2015, **16**(11)**:** 660-671.

60.Xu W, Caracciolo B, Wang HX, Winblad B, Bäckman L, Qiu C, *et al.* Accelerated progression from
mild cognitive impairment to dementia in people with diabetes. *Diabetes* 2010, **59**(11)**:** 2928-2935.

61.Selkoe DJ. Toward a comprehensive theory for Alzheimer's disease - Hypothesis: Alzheimer's
disease is caused by the cerebral accumulation and cytotoxicity of amyloid beta-protein. In:
Khachaturian ZS, Mesulam MM (eds). *Alzheimers Disease: A Compendium of Current Theories*,
vol. 924, 2000, pp 17-25.

62.Thijs V, Robberecht W, De Vos R, Sciot R. Coexistence of CADASIL and Alzheimer's disease.
Journal of Neurology Neurosurgery and Psychiatry 2003, **74**(6)**:** 790-792.

63.Brai E, Raio NA, Alberi L. Notch1 hallmarks fibrillary depositions in sporadic Alzheimer's disease.
Acta Neuropathologica Communications 2016, **4**.

64.Liu B, Wang D, Xiong T, Liu Y, Jing X, Du J, *et al.* Inhibition of Notch Signaling Promotes the
Differentiation of Epicardial Progenitor Cells into Adipocytes. *Stem Cells Int* 2021, **2021:** 8859071.

65.D'Souza B, Meloty-Kapella L, Weinmaster G. CANONICAL AND NON-CANONICAL NOTCH
LIGANDS. In: Kopan R (ed). *Notch Signaling*, vol. 92, 2010, pp 73-129.

66. Salloway S, Sperling R, Fox NC, Blennow K, Klunk W, Raskind M, *et al.* Two Phase 3 Trials of
Bapineuzumab in Mild-to-Moderate Alzheimer's Disease. *New England Journal of Medicine* 2014,
370(4)**:** 322-333.

67. Ostrowitzki S, Lasser RA, Dorflinger E, Scheltens P, Barkhof F, Nikolcheva T, *et al.* A phase III
randomized trial of gantenerumab in prodromal Alzheimer's disease. *Alzheimers Research &*
Therapy 2017, **9**.

68. Honig LS, Vellas B, Woodward M, Boada M, Bullock R, Borrie M, *et al.* Trial of Solanezumab for
Mild Dementia Due to Alzheimer's Disease. *New England Journal of Medicine* 2018, **378**(4)**:** 321-
330.

69. Henley D, Raghavan N, Sperling R, Aisen P, Raman R, Romano G. Preliminary Results of a Trial of
Atabecestat in Preclinical Alzheimer's Disease. *New England Journal of Medicine* 2019, **380**(15)**:**
1483-1485.

70. Egan MF, Kost J, Voss T, Mukai Y, Aisen PS, Cummings JL, *et al.* Randomized Trial of
Verubecestat for Prodromal Alzheimer's Disease. *New England Journal of Medicine* 2019, **380**(15)**:**
1408-1420.

71. Panza F, Lozupone M, Logroscino G, Imbimbo BP. A critical appraisal of amyloid-beta targeting
therapies for Alzheimer disease. *Nature Reviews Neurology* 2019, **15**(2)**:** 73-88.

72. Wessels AM, Lines C, Stern RA, Kost J, Voss T, Mozley LH, *et al.* Cognitive outcomes in trials of
two BACE inhibitors in Alzheimer's disease. *Alzheimer's & dementia : the journal of the*
Alzheimer's Association 2020, **16**(11)**:** 1483-1492.

73. Novak G, Streffer JR, Timmers M, Henley D, Brashear HR, Bogert J, *et al.* Long-term safety and
tolerability of atabecestat (JNJ-54861911), an oral BACE1 inhibitor, in early Alzheimer's disease
spectrum patients: a randomized, double-blind, placebo-controlled study and a two-period extension
study. *Alzheimers Res Ther* 2020, **12**(1)**:** 58.

74. Timmers M, Streffer JR, Russu A, Tominaga Y, Shimizu H, Shiraishi A, *et al.* Pharmacodynamics of
atabecestat (JNJ-54861911), an oral BACE1 inhibitor in patients with early Alzheimer's disease:
randomized, double-blind, placebo-controlled study. *Alzheimers Res Ther* 2018, **10**(1)**:** 85.

75. Nigam SM, Xu S, Ackermann F, Gregory JA, Lundkvist J, Lendahl U, *et al.* Endogenous APP
accumulates in synapses after BACE1 inhibition. *Neuroscience research* 2016, **109:** 9-15.

76. Citron M. Emerging Alzheimer's disease therapies: inhibition of beta-secretase. *Neurobiol Aging*
2002, **23**(6)**:** 1017-1022.

77.Long JM, Holtzman DM. Alzheimer Disease: An Update on Pathobiology and Treatment Strategies.
Cell 2019, **179**(2)**:** 312-339.

78.Livingston G, Sommerlad A, Orgeta V, Costafreda SG, Huntley J, Ames D*, et al.* Dementia
prevention, intervention, and care. *Lancet* 2017, **390**(10113)**:** 2673-2734.

79.Butterfield DA, Di Domenico F, Barone E. Elevated risk of type 2 diabetes for development of
Alzheimer disease: A key role for oxidative stress in brain. *Biochimica Et Biophysica Acta-*
Molecular Basis of Disease 2014, **1842**(9)**:** 1693-1706.

80.Arnold SE, Arvanitakis Z, Macauley-Rambach SL, Koenig AM, Wang H-Y, Ahima RS*, et al.* Brain
insulin resistance in type 2 diabetes and Alzheimer disease: concepts and conundrums. *Nature*
Reviews Neurology 2018, **14**(3)**:** 168-181.

81.Zhao QY, Wu XH, Yan S, Xie XF, Fan YH, Zhang JQ*, et al.* The antidepressant-like effects of
pioglitazone in a chronic mild stress mouse model are associated with PPAR gamma-mediated
alteration of microglial activation phenotypes. *Journal of neuroinflammation* 2016, **13**.

82.Guo M, Li C, Lei Y, Xu S, Zhao D, Lu XY. Role of the adipose PPAR gamma-adiponectin axis in
susceptibility to stress and depression/anxiety-related behaviors. *Molecular Psychiatry* 2017, **22**(7)**:**
1056-1068.

83.Toledo EM, Inestrosa NC. Activation of Wnt signaling by lithium and rosiglitazone reduced spatial
memory impairment and neurodegeneration in brains of an APPswe/PSEN1DeltaE9 mouse model
of Alzheimer's disease. *Mol Psychiatry* 2010, **15**(3)**:** 272-285, 228.

84.Escribano L, Simón AM, Gimeno E, Cuadrado-Tejedor M, López de Maturana R, García-Osta A*, et*
al. Rosiglitazone rescues memory impairment in Alzheimer's transgenic mice: mechanisms
involving a reduced amyloid and tau pathology. *Neuropsychopharmacology : official publication of*
the American College of Neuropsychopharmacology 2010, **35**(7)**:** 1593-1604.

85.O'Reilly JA, Lynch M. Rosiglitazone improves spatial memory and decreases insoluble Aβ(1-42) in
APP/PS1 mice. *Journal of neuroimmune pharmacology : the official journal of the Society on*
NeuroImmune Pharmacology 2012, **7**(1)**:** 140-144.

86.Watson GS, Cholerton BA, Reger MA, Baker LD, Plymate SR, Asthana S*, et al.* Preserved
cognition in patients with early Alzheimer disease and amnestic mild cognitive impairment during
treatment with rosiglitazone: a preliminary study. *The American journal of geriatric psychiatry :*
official journal of the American Association for Geriatric Psychiatry 2005, **13**(11)**:** 950-958.

87.Tzimopoulou S, Cunningham VJ, Nichols TE, Searle G, Bird NP, Mistry P*, et al.* A multi-center

randomized proof-of-concept clinical trial applying [¹□F]FDG-PET for evaluation of metabolic
therapy with rosiglitazone XR in mild to moderate Alzheimer's disease. *Journal of Alzheimer's*
disease : JAD 2010, **22**(4)**:** 1241-1256.

88.Harrington C, Sawchak S, Chiang C, Davies J, Donovan C, Saunders AM*, et al.* Rosiglitazone does
not improve cognition or global function when used as adjunctive therapy to AChE inhibitors in
mild-to-moderate Alzheimer's disease: two phase 3 studies. *Current Alzheimer research* 2011, **8**(5)**:**
592-606.

89.Keating GM, Croom KF. Fenofibrate: a review of its use in primary dyslipidaemia, the metabolic
syndrome and type 2 diabetes mellitus. *Drugs* 2007, **67**(1)**:** 121-153.

90.Barbiero JK, Santiago R, Tonin FS, Boschen S, da Silva LM, Werner MF*, et al.* PPAR-α agonist
fenofibrate protects against the damaging effects of MPTP in a rat model of Parkinson's disease.
Prog Neuropsychopharmacol Biol Psychiatry 2014, **53:** 35-44.

91.Ouk T, Gautier S, Pétrault M, Montaigne D, Maréchal X, Masse I*, et al.* Effects of the PPAR-α
agonist fenofibrate on acute and short-term consequences of brain ischemia. *Journal of Cerebral*
Blood Flow & Metabolism 2014, **34**(3)**:** 542-551.

92.Assaf N, El-Shamarka ME, Salem NA, Khadrawy YA, El Sayed NS. Neuroprotective effect of
PPAR alpha and gamma agonists in a mouse model of amyloidogenesis through modulation of the
Wnt/beta catenin pathway via targeting alpha- and beta-secretases. *Prog Neuropsychopharmacol*
Biol Psychiatry 2020, **97:** 109793.

Authors' contributions

G. Gao, X Yang, B Jiang, and P Jiang were involved in the concept and design of

the study. W. Qi, Y. Long, Z. Li, Z. Zhao J. Shi, D Zhu, Z Zhao, W. Xie, L. Wang, T

Zhou, Mingting Liang were responsible for conducting the experiments. W. Qi, Y.

Long and T Zhou drafted the manuscript and G. Gao revised the manuscript. P Jiang,

Y. Long and Z. Li were responsible for data analysis. All authors contributed to the

interpretation of data and provided revisions to the manuscript. G. Gao will act as

Figure legend

Fig.1 Increased Kallistatin was presented in AD patients and could impair cognitive memory in mice. (A-B) Serum Kallistatin(A), fasting blood glucose (FBG), triglyceride (TG), and total cholesterol (TC) (B) of AD patients and their corresponding normal control subjects. (C-D) Serum Kallistatin(C), TG, and TC(D) of AD patients with DM and their corresponding normal control subjects (Student's t-test). (E-J) The behavioral performance of KAL-TG mice was assessed through the Morris water maze test, Y-maze test, and electrophysiology. (E)The escape latency time of different months of KAL-TG mice (3M, 6M, 9M, 12M) and corresponding WT mice were presented during 1-6 day (two-way ANOVA). (F-H) Cognitive functions were evaluated by spatial probe test at day 7 (Student's t-test), the representative each group mice traces were shown (F), then analyzing each group mice crossing platform times (G) and time percent in the targeted area (H), n=4 to 9 per group. (I) Spontaneous alternation of Y-maze test. (J) LTP was measured by whole-cell voltage-clamp recordings of CA1 neurons in acute hippocampal slices of KAL-TG (3M, 6M, 12M) and WT mice (Student's t-test, n=6-12 cells from 3 mice per group). Error bars represent the standard deviation (SD); one asterisk, $p < 0.05$, two asterisks, $p < 0.01$; four asterisks, $p < 0.0001$.

Fig.2 Kallistatin promoted Aβ generation. (A-B) Immunohistochemistry staining of Aβ(A) was carried out in KAL-TG and WT mice hippocampal tissue. *Scale bar*,

100μm. The statistical analysis of Aβ plaques (B) in hippocampal tissue of KAL-TG

and WT mice, n=4-5 per group. (C) Protein levels of Aβ were tested by western blot

analysis in hippocampal tissue, n=3 per group, then statistically analyzed the above

results. (D) Hippocampal tissue Aβ42 contents were performed by ELISA in KAL-TG

and WT groups, n=3 per group. (E) Primary mouse neurons were isolated, then

infected with adenovirus to overexpress Kallistatin for 72h. Aβ42 concentration of

primary hippocampal neurons supernate and cell lysate were quantified by ELISA,

n=3 per group. (F-G) Western blot analysis of Aβ protein level in primary

hippocampal neurons infected with overexpressing Kallistatin adenovirus and control

groups, then statistical analysis of Aβ protein levels, n=3 per group. Error bars

represent the standard deviation (SD); one asterisk, $p < 0.05$; two asterisks, $p < 0.01$;

three asterisks, $p < 0.001$; Student's t-test.

Fig.3 Kallistatin promoted tau phosphorylation. (A-B) Immunohistochemistry

staining of phosphorylated tau (p-tau S396, p-tau T231, p-tau S202) and tau(A) was

carried out in KAL-TG and WT mice hippocampal tissue. Scale bar, 100μm. The

statistical analysis of phosphorylated tau(B) in hippocampal tissue of KAL-TG and

WT mice, n=3 per group. (C-D) Protein levels of phosphorylated tau (p-tau S396, p-

tau T231, p-tau S202) and tau were tested by western blot analysis in hippocampal

tissue, then statistically analyzed the above results. Error bars represent the standard

deviation (SD); one asterisk, $p < 0.05$; two asterisks, $p < 0.01$; three asterisks, $p <$

0.001; Student's t-test.

Fig.4 Kallistatin transgenic mice exhibited increased BACE1 expression and

activity in the hippocampus. (A) Western blot analysis of relevant proteins, such as

APP, PS1, and BACE1 during Aβ generation in hippocampal tissue of each time point

(6M, 9M, 12M) KAL-TG mice and corresponding WT control groups, n=3 per group,

then statistical analysis of APP, PS1 and BACE1 protein levels. (B)

Immunohistochemistry staining of BACE1 was carried out in KAL-TG and WT mice

hippocampal tissue at each time point (6M, 9M, 12M). n=3 to 5 per group. *Scale bar*,

100 μm. (C) Statistical analysis of BACE1 immunohistochemistry staining, n=3 to 4

per group. (D) ELISA measured the β-secretase activity of each group's hippocampal

tissue, n=3 per group. Error bars represent the standard deviation (SD); one asterisk, *p*

< 0.05, two asterisks, $p < 0.01$; Student's t-test.

1002

Fig.5 *In vitro*, Kallistatin promoted BACE1 expression to augment Aβ by

suppressing PPARγ activation. (A) The relevant protein levels in primary mouse

neurons infected with overexpressing Kallistatin adenovirus during Aβ generation

were determined by western blot analysis. (B) Statistical analysis of BACE1

expression in primary neurons. (C-D) β-secretase(C) and γ- secretase(D) activity of

primary hippocampal neurons infected with overexpressing Kallistatin adenovirus and

control adenovirus was measured by ELISA. (E) Primary hippocampal neurons were

treated with BACE1 inhibitor verubecestat (50nM), then infected with adenovirus to

overexpress Kallistatin. Western blot analysis of Aβ, BACE1, and Kallistatin protein

levels, β-actin served as a loading control. (F) HT22 cells were infected with BACE1

siRNA, then infected with adenovirus to overexpress Kallistatin. Western blot

analysis of Aβ and BACE1 protein levels, β-actin served as a loading control. (G) The

relevant proteins involved in BACE1 transcriptional expressions, such as PPARγ,

YY1, and SP1 were measured by western blot analysis in hippocampal tissue. β-actin

served as a loading control. (H) Statistical analysis of PPARγ in hippocampal tissue

each group. (I) The representative diagrams of PPARγ expression in hippocampal

tissue were presented in the above graphs. *Scale bar*, 100μm. (J) Statistical analysis of

PPARγ immunohistochemistry staining in hippocampal tissue of each group, n=3 to 4

per group. (K)Primary hippocampal neurons were treated with PPARγ agonist

rosiglitazone (10nM) for 12h, then infected with adenovirus to overexpress Kallistatin

for 72h. Western blot analysis of Aβ and BACE1 protein levels. β-actin served as a

loading control. (L) Statistical analysis of PPARγ protein levels in each group. Error

bars represent the standard deviation (SD), one asterisk, $p < 0.05$, two asterisks, $p <$

0.01; three asterisks, $p < 0.001$; ns means no significant difference; Student's *t*-test.

1027

Fig.6 Kallistatin directly bonded to the Notch1 receptor, which activated the

Notch1 pathway to promote Aβ production. (A) Notch1 expression was measured

by western blot analysis in hippocampal tissue. β-actin served as a loading control. (B)

Statistical analysis of Notch1 in hippocampal tissue of each group. (C) The

representative diagrams of Nocth1 expression in hippocampal tissue were presented in

the above graphs. Scale bar, 100μm. (D) Statistical analysis of Notch1

immunohistochemistry staining in hippocampal tissue of each group. (E-F) Primary

hippocampal neurons were infected with overexpressing Kallistatin adenovirus for

72h, then Co-IP analysis (E) and membrane extraction experiment (F) was performed

to verify whether Kallistatin can bind to the Notch1 receptor. β-actin served as a

loading control. (G-H) HT22 cells were treated with siRNA (Notch1) and shRNA

(HES1) to knock down Notch1 and HES1 for 12h, then infected with adenovirus to

overexpress Kallistatin for 24h. Western blot analysis was used to detect the Notch1

signaling pathway. Error bars represent the standard deviation (SD), one asterisk, $p <$

0.05; Student's t-test.

1043

Fig.7 Kallistatin promoted phosphorylation of tau by suppressing Wnt signaling

pathway. (A-B) GSK-3β and p-GSK-3β expression was measured by western blot

analysis in hippocampal tissue, then statistically analyzed the above results. (C-D)

Primary hippocampal neurons were treated with overexpressing Kallistatin adenovirus

for 72h, then western blot analysis was used to detect the content of GSK-3β, p-GSK-

3β, tau, p-tau (Ser9, T231, S396), and statistically analyzed the above results. (E-F)

Primary hippocampal neurons were treated with overexpressing Kallistatin adenovirus

for 48h, then treated with LiCl (10mM) for 24h, western blot analysis was used to

detect the content of GSK-3β, p-GSK-3β, p-tau (Ser9, T231, S396), and statistical

analysis of the above results. Error bars represent the standard deviation (SD), two

asterisk, p < 0.01; three asterisk, p < 0.001; Student's t-test.

1055

Fig.8 Fenofibrate could alleviate memory and cognitive impairment of KAL-TG

mice. (A) Illustration of experimental protocols. Fenofibrate (0.3 g/kg/d × 5week, i.g.)

or rosiglitazone (5mg/kg/d × 5week, i.g.) were given to KAL-TG mice. The serum for

Kallistatin measuring was collected at week 0 and week 4. And Morris water maze

and Y-maze test were performed at week 5. (B-E) Behavioral performance was

assessed through the Morris water maze test and the Y-maze test. (B) The escape

latency time was presented during 1-5 day. (C-E) Cognitive functions were evaluated

by spatial probe test at day 6, then analyzing each group of mice crossing platform

times(C), time percent in the targeted area (D), and the path traces heatmap (E), n=5

to 6 per group. (F) Spontaneous alternation of Y-maze test. (G) Kallistatin decreased

ratio was calculated by dividing the serum Kallistatin concentration of KAL-TG mice

before Fenofibrate/ rosiglitazone treatment by the serum Kallistatin concentration of

KAL-TG mice after a-month treatment, and serum Kallistatin concentration was

measured by ELISA. (H-I) Protein levels of Aβ and BACE1 were tested by western

blot analysis in hippocampal tissue, then statistically analyzing the above results. (J-K)

Protein levels of p-tau(231), tau, p-GSK-3β(Ser9) and GSK-3β were tested by western

blot analysis in hippocampal tissue , then statistically analyzing the above results.

Error bars represent the standard deviation (SD); one asterisk, $p < 0.05$; Student's t-

test.

1075

1076 Table.S1 Clinical characteristic of AD patients

	NC	AD	*p value* *(Dementia vs NC)*
N	61	56	NA
Age	49-82	52-85	****
Average age	64±8.53	73.02±9.43	NA
Male	29	25	NS
Female	32	31	
GLU (mmol/L)	5.07±0.65	6.73±2.81	****
TC (mmol/L)	4.63±0.58	4.50±1.12	NS
TG (mmol/L)	1.09±0.50	1.41±0.65	**

1077 N is an abbreviation for number; GLU is an abbreviation for glucose; TC is an

1078 abbreviation for total cholesterol; TG is an abbreviation for triglyceride; NA indicates

1079 not available; NS indicates no significance. two asterisk, $p < 0.01$, four asterisk, $p <$

1080 0.0001.

1081

1082 Table.S2 Clinical characteristic of AD patients with DM

	NC	AD + DM	*p value* *(Dementia+DM vs NC)*
N	61	26	NA
Age	49-82	52-92	****
Average age	64±8.53	76.77±8.69	NA
Male	29	15	NS
Female	32	11	
GLU (mmol/L)	5.07±0.65	8.12±3.05	****

TC (mmol/L)	4.63±0.58	4.29±1.04	NS
TG (mmol/L)	1.08±0.50	1.69±0.68	****

1083 N is an abbreviation for number; GLU is an abbreviation for glucose; TC is an

1084 abbreviation for total cholesterol; TG is an abbreviation for triglyceride; NA indicates

1085 not available; NS indicates no significance. four asterisk, $p < 0.0001$.

1086

Figure legend

Fig.S1 (A-B) GAD disease enrichment analysis (A) and PFAM analysis (B) result.

Differentially expressed genes in neurons were obtained from GSE161355, and GAD

disease enrichment was analyzed on David database. (C-D) Western blot analysis of

Kallistatin expression in aging model SAMP8 and corresponding control SAMR1

mice hippocampal tissue samples, then statistically analyzing the above results. β-

Actin served as a loading control. Error bars represent the standard deviation (SD);

one asterisks, $p < 0.05$.

1095

Fig.S2 (A) HT22 cells were infected with adenovirus to overexpress Kallistatin for

48h. Aβ42 concentration of supernate and cell lysate was quantified by ELISA. (B-C)

Western blot analysis of Aβ protein level in HT22 cells infected with overexpressing

Kallistatin adenovirus and control groups for 48h, then statistical analysis of

Kallistatin protein levels. (D) BACE1 mRNA expression in hippocampal tissue. (E) α-

secretases (ADAM9, ADAM10, ADAM17) mRNA expression in hippocampal tissue.

(F) γ-secretase activity of each group's hippocampal tissue was measured by ELISA.

(G) Primary hippocampal neurons were identified with MAP2. Error bars represent

the standard deviation (SD), one asterisk, $p < 0.05$, two asterisks, $p < 0.01$.

1105

Fig.S3 (A) The relevant protein levels in HT22 cells infected with overexpressing

Kallistatin adenovirus during Aβ generation were determined by western blot analysis.

(B) Statistical analysis of BACE1 expression in HT22 cells. (C) β-secretase activity of

HT22 cells infected with overexpressing Kallistatin adenovirus and control

adenovirus was measured by ELISA. (D) Primary hippocampal neurons were infected

with BACE1 siRNA for 72h. Western blot analysis of BACE1 protein levels. (E)

HT22 cells were treated with PPARγ agonist rosiglitazone (10nM) for 12h, then

infected with adenovirus to overexpress Kallistatin for 48h. Western blot analysis of

Aβ and BACE1 protein levels. β-actin served as a loading control. Error bars

represent the standard deviation (SD), two asterisks, $p < 0.01$.

1116

Fig.S4 (A-B) Western blot analysis of Notch1 protein level in primary hippocampal

neurons and HT22 cells infected with overexpressing Kallistatin adenovirus and

control groups. (C) HT22 cells were infected with overexpressing Kallistatin

adenovirus for 48h, and Co-IP analysis was conducted to verify whether Kallistatin

can bind to the Notch1 receptor. β-actin served as a loading control. (D) Primary

hippocampal neurons were treated with Kallistatin protein for 72h, then IP analysis.

(E) Primary hippocampal neurons were infected with Notch1 siRNA for 72h. Western

blot analysis of BACE1 protein levels. (F) HT22 cells were infected with HES1

shRNA for 48h. Western blot analysis of HES1 protein levels.

1126

Fig.S5 (A) Western blot analysis of GSK-3β, p-GSK-3β, tau, p-tau(Ser9, T231, S396)

in HT22 cells infected with overexpressing Kallistatin adenovirus and control groups.

(B) Western blot analysis of GSK-3β, p-GSK-3β, p-tau(Ser9, T231, S396) in HT22

cells infected with overexpressing Kallistatin adenovirus and control groups for 24h,

then treated with LiCl(10 mM) for 24h. (C) Simplified model depicting the pathway

of Aβ regulated by Kallistatin.

1133

Fig.1

Fig.2

Fig.3

Fig.4

Fig.5

Fig.6

Fig.7

Fig.8

Fig.S1

A

GAD Disease Enrichment

NC neuron vs T2DM neuron

B

PFAM Enrichment

NC neuron vs T2DM neuron

C

D

primary hippocampal neurons

Fig.S3

Fig.S4

Fig.S5

A

HT22

	CON	GFP	KAL	
p-tau S202				52KD
p-tau T231				52KD
p-tau S396				52KD
tau				52KD
p-GSK3β				65KD
GSK3β				65KD
KAL				50KD
β-actin				42KD

B

HT22

Ad-GFP	+	—	—	
Ad-KAL	—	+	+	
LiCl(10mM)	—	—	+	
p-GSK3β(Ser9)				65KD
GSK3β				65KD
p-tau S202				52KD
p-tau T231				52KD
p-tau S396				52KD
KAL				50KD
β-actin				42KD

C