The novel role of Kallistatin in linking metabolic syndromes and cognitive memory deterioration by inducing amyloid-β plaques accumulation and tau protein hyperphosphorylation

Weiwei Qi[1,2#], Yanlan Long[3#], Ziming Li[4#], Zhen Zhao[1], Jinhui Shi[1], Wanting Xie[1], Laijian Wang[4], Yandan Tan[1], Ti Zhou[1], Ping Jiang[5*], Bin Jiang[4*], Xia Yang[1,6*], Guoquan Gao[1,7,8,9*]

[1] Department of Biochemistry and molecular biology, Zhongshan School of Medicine, Sun Yat-sen University, Guangzhou, China

[2] Advanced Medical Technology Center, The First Affiliated Hospital, Zhongshan School of Medicine, Sun Yat-sen University

[3] Guangdong Key Laboratory of Nanomedicine, CAS-HK Joint Lab of Biomaterials, Shenzhen Institute of Advanced Technology (SIAT), Chinese Academy of Sciences (CAS), Shenzhen, China

[4] Guangdong Province Key Laboratory of Brain Function and Disease, School of Medicine, Sun Yat-sen University, Shenzhen, China

[5] Department of Clinical Medical Laboratory, Guangzhou First People Hospital, School of Medicine, South China University of Technology, Guangzhou, China

[6] China Key Laboratory of Tropical Disease Control (Sun Yat-sen University), Ministry of Education, Guangzhou, China

[7] Guangdong Engineering & Technology Research Center for Gene Manipulation and Biomacromolecular Products (Sun Yat-sen University), Guangzhou, China

[8] Guangdong Province Key Laboratory of Brain Function and Disease,

Zhongshan School of Medicine, Sun Yat-sen University, Guangzhou, China

[9] Guangdong Provincial Key Laboratory of Diabetology& Guangzhou Municipal

Key Laboratory of Mechanistic and Translational Obesity Research, Medical Center

for Comprehensive Weight Control, The Third Affiliated Hospital of Sun Yat-sen

University Guangzhou, Guangdong, China

[#]These authors contributed equally to this study.

*Corresponding   author: Gao   GQ: gaogq@mail.sysu.edu.cn;   Yang   X:

yangxia@mail.sysu.edu.cn;   Jiang   B: jiangb3@mail.sysu.edu.cn;   Ping   J:

jiangp45@mail2.sysu.edu.cn

Highlights

● Kallistatin-transgenic(KAL-TG) mice defined its cognitive memory impairment

phenotype accompanied by increased Aβ deposition and tau phosphorylation.

● Kallistatin could directly bind to the Notch1 receptor and thereby upregulate

BACE1 expression by inhibiting PPARγ signaling.

● Fenofibrate could alleviate cognitive memory impairment and down-regulate the

serum Kallistatin level.

Abstract

Accumulation of amyloid β (Aβ) peptides and hyperphosphorylated tau proteins

in the hippocampus triggers cognitive memory decline in Alzheimer's disease (AD).

The incidence and mortality of sporadic AD were tightly associated with diabetes and

hyperlipidemia, while the exact linked molecular is uncertain. Here, we reported that

serum Kallistatin concentrations were meaningfully higher in AD patients, with a

higher concentration of fasting blood glucose and triglyceride. In addition, the

constructed Kallistatin-transgenic (KAL-TG) mice defined its cognitive memory

impairment phenotype and lower LTP in hippocampal CA1 neurons accompanied by

increased Aβ deposition and tau phosphorylation. Mechanistically, Kallistatin could

directly bind to the Notch1 receptor and thereby upregulate BACE1 expression by

inhibiting PPARγ signaling, resulting in Aβ cleavage and production. Besides,

Kallistatin could promote the phosphorylation of tau by activating GSK-3β.

Fenofibrate, a hypolipidemic drug, could alleviate cognitive memory impairment by

down-regulating Aβ and tau phosphorylation of KAL-TG mice. Collectively, our data

clarified a novel mechanism for Aβ accumulation and tau protein

hyperphosphorylation regulation by Kallistatin, which might play a crucial role in

linking metabolic syndromes and cognitive memory deterioration, and suggested that

fenofibrate might have the potential for treating metabolism-related AD.

Key words: Alzheimer's disease, Kallistatin, Aβ42, BACE1, Diabetes

Introduction

Alzheimer's disease (AD), the most common irreversible neurodegenerative disease involving dementia in the elderly, is characterized by progressive deterioration of cognitive memory. Pathologically, the characteristic hallmarks of AD are extracellular amyloid-β (Aβ) plaques and intracellular neurofibrillary tangles (NFTs) [1, 2, 3]. Aβ cascade and the hyperphosphorylation of tau protein are two main hypotheses related to AD. The Aβ cascade hypothesis dominates that overproduction of Aβ changes the normal cellular state, resulting in synaptic dysfunction, neurodegeneration, tau hyperphosphorylation, and neuroinflammation, eventually causing memory loss in AD and dementia patients [4, 5]. Aβ peptides are produced from the sequential cleavage of amyloid precursor protein (APP) by β-secretase (β-site APP cleaving enzyme 1, BACE1) and γ-secretase, therefore this cleavage step contributes heavily to AD pathology [6, 7]. BACE1 is one of the most potential therapeutic targets. Some potent BACE1 inhibitors have advanced to late stages in clinical trials, highlighting the significance of BACE1 in the Aβ generation [8, 9, 10]. Tau is a protein found in microtubules. It exists physiologically in axons and regulates microtubule dynamics and axonal transport [11]. Tau goes through a multi-step process in AD, transitioning from a natively unfolded monomer to massive aggregated aggregates like NFTs, another hallmark of AD [12, 13]. One of the central kinases responsible for initiating tau phosphorylation is glycogen synthase kinase-3 (GSK3) and Wnt signaling play a vital role in activating GSK-3β and GSK-3β-mediated tau

phosphorylation [14]. The physiological mechanisms governing their relationship are

still poorly understood.

There has been a close relationship between metabolic disorders and cognitive

impairment across the AD spectrum [15, 16]. Accounting for nearly 95% of AD patients

were categorized as sporadic ones, whose increasing incidence and mortality were

tightly associated with type 2 diabetes mellitus (T2DM), obesity, and hyperlipidemia

[17, 18, 19]. There was about 37 percent comorbidity between AD and diabetes reported

by the Alzheimer's Association Report [20, 21]. As a result of the strong association and

share mechanism between AD and T2DM, AD was termed "Type 3 Diabetes" by

some researchers [22, 23, 24, 25]. Several studies demonstrate that diabetes confers a 1.6-

fold increased risk of developing dementia [26, 27]. Similarly, central obesity and high

body mass index (BMI) during middle age have about 3.5 times increased risk of

dementia later in life [28]. Therefore, control of blood glucose and lipids is expected to

be a strategy for preventing or moderating cognitive decline in aging. Nevertheless,

the exact link and key associated regulator between metabolic abnormalities and AD

is still unclear.

Kallistatin was a serine proteinase inhibitor, previously identified as a tissue

kallikrein-binding protein [29]. It was predominantly produced in the liver and widely

expressed in the body tissues and conferred anti-angiogenesis, anti-fibrosis, anti-

oxidative stress, and anti-tumor growth [30, 31]. Furthermore, Kallistatin was found

increased in patients with obesity, prediabetes, and diabetes [32, 33, 34]. The concentration

of Kallistatin was positively correlated with the triglyceride glucose index [35], which

was proven to be an independent risk factor for dementia [36]. In addition, our previous

study revealed that the concentration of serum Kallistatin in T2DM patients was

significantly increased and further clarified that Kallistatin suspended wound healing

in T2DM patients by promoting local inflammation, which suggested that Kallistatin

played a critical role in the progression of T2DM [37]. Furthermore, our recent research

has found that kallistatin can cause memory and cognitive dysfunction by disrupting

the glutamate-glutamine cycle[38].

According to the relationship between T2DM, AD and Kallistatin, we

constructed Kallistatin transgenic (KAL-TG) mice to explore whether it could cause

cognitive impairment through up-regulation of Aβ production. Taken together, our

results suggest that a novel regulatory mechanism of Aβ production and tau protein

hyperphosphorylation by Kallistatin is involved in the progression of metabolic

abnormalities-related AD.

Results

Kallistatin was increased in AD patients and AD model mice.

To explore the relevance of AD in T2DM (Fig.S1A), we first performed a GAD

disease enrichment analysis of differentially expressed genes in neurons of T2DM and

normal controls, and found AD was closely related to T2DM (GSE161355). We

further identified enrichment of the Serpin family protein domain using PFAM

analysis on the David database (Fig.S1B) (https://david.ncifcrf.gov/). Our previous studies found that Kallistatin (serpin family a member 4) was elevated in the serum of T2DM patients and caused an adverse prognosis of diabetes complications [44]. We have collected 11 serum samples of dementia patients from Sun Yat-sen Memorial Hospital and found that the concentration of Kallistatin was higher compared to the normal control[38]. Here, we enrolled 56 AD patients and 61 healthy controls from four hospitals in Guangdong province to further explore the potential relevance of Kallistatin and AD. The Clinical and biochemical characteristics are presented in Table S1, S2. In addition, we measured the serum Kallistatin (12.78±2.80 μg/ml) content in patients with AD was higher than in normal controls (9.78±1.93 μg/ml) (Fig.1A). Likewise, the fasting blood glucose (FBG) and triglyceride (TG) were more elevated in AD patients than in healthy human (Fig. 1B). We further grouped all the AD patients according to whether they had diabetes, then found Kallistatin (13.79±3.05 μg/ml) and TG content were further elevated in AD patients with diabetes (Fig.1C-D). Similarly, Kallistatin expression was increased in AD model mouse SAMP8 compared with its control mouse SAMR1 in hippocampus tissue (Fig.S1C-D). Taken together, these results indicated that Kallistatin concentration was increased in metabolic abnormalities-related AD patients.

Kallistatin could impair cognitive memory in mice.

The above experiments proved that Kallistatin increased in AD patients and AD model mice. Then, we constructed KAL-TG mice and assessed their behavioral

performance through the Morris water maze (MWM) and Y-maze tests. Notably, the

latency to escape platform was increased, and crossing platform times, time in

percent, and spontaneous alternation were significantly decreased in KAL-TG mice

compared with age-matched WT mice (Fig.1E-I). Furthermore, long-term potentiation

(LTP) was measured by whole-cell voltage-clamp recordings of CA1 neurons in acute

hippocampal slices of KAL-TG and WT mice to determine the changes in

hippocampal synapses. LTP of KAL-TG mice was significantly lower than WT mice

(Fig.1J). These results showed Kallistatin could impair cognitive memory in mice.

Kallistatin promoted Aβ deposition and tau phosphorylation.

We evaluated Aβ deposition and tau phosphorylation in these experimental mice

hippocampus tissues by immunohistochemistry staining and western blot. Predictably,

the plaque density and tau phosphorylation of KAL-TG mice were much higher than

age-matched WT mice (Fig.2A-C, 3A-D). Consistent with this result, ELISA

detection of Aβ42 contents in hippocampus tissue showed that Aβ production was

extraordinarily increased in KAL-TG mice compared to WT mice (Fig.2D). These

results suggested that Kallistatin promoted Aβ deposition and tau phosphorylation.

Kallistatin positively regulated Aβ generation by promoting β-secretase rather

than γ-secretase.

Western blot and ELISA analysis of Aβ levels in primary hippocampal neurons

(immunofluorescence identified with neural marker MAP2, Fig.S2G) infected with

overexpressing Kallistatin adenovirus were higher than in control groups (Fig.2E-G),

as well as HT22 cells (Fig.S2A-C). Amyloid-beta precursor protein (APP) undergoes

proteolytic processing to generate peptide fragments [45]. β-secretase (BACE1) and γ-

secretase, composed of presenilin 1 (PS1), nicastrin, and Pen-2, were crucial enzymes

for Aβ generation [46, 47]. We determined the levels of APP, BACE1, and PS1 in

hippocampus tissue. BACE1 protein and mRNA accumulate to higher levels in KAL-

TG mice compared with WT mice (Fig.4A-C, S2D, whereas no significant difference

appeared in APP, PS1, and α-secretases (ADAM9, ADAM10, and ADAM17)

expression (Fig.4A, S2E). Consistent with the above results, the activity of BACE1

was increased (Fig.4D), while PS1 activity was not changed (Fig.S2F). Similarly, the

expression and activity of BACE1 were further determined to increase in primary

mouse neurons, and HT22 cells transfected with Kallistatin adenovirus (Fig.5A-C,

S3A-C), and PS1 expression and activity were still not changed (Fig.5A, 5D, S3A).

Furthermore, the effect of Kallistatin was attenuated by the BACE1 inhibitor

(verubecestat) or siBACE1 03, the most effective one (Fig.5E-F, S3D). Therefore,

these results indicated that Kallistatin could promote Aβ generation through

upregulating BACE1 expression.

Kallistatin suppressed PPARγ activation to promote BACE1 expression.

Transcription factors SP1, YY1, and PPAR were reported to regulate BACE1

expression at the transcription level. Among them, PPARγ could down-regulated

BACE1 expression [48, 49, 50]. As elaborated, PPARγ decreased in the hippocampal tissue

of KAL-TG mice detected by western blot and immunohistochemical analysis

(Fig.5G-J). However, no statistical differences appeared in the YY1 and SP1

expressions (Fig.5G). In addition, Kallistatin downregulated the expression of PPARγ

on primary hippocampal neurons and HT22 cells, thus increasing BACE1 and Aβ

expression (Fig.5K, S3E). Treatment with rosiglitazone, the PPARγ agonist, could

reverse the decrease of PPARγ caused by Kallistatin (Fig.5K-L). Predictably,

rosiglitazone could inhibit the promoting effect of Kallistatin on BACE1 and Aβ

(Fig.5K and Fig.S3E).

Kallistatin promoted Aβ production *via* directly binding to the Notch1 receptor

and activating the Notch1 pathway.

Our results indicated that Notch1 was highly expressed in the hippocampal

tissues of KAL-TG mice (Fig.6A-D). Moreover, primary hippocampal neurons and

HT22 cells infected with overexpressing Kallistatin adenovirus had upregulated

Notch1 expression *in vitro* (Fig.S4A-B). In addition, Kallistatin could directly bind to

the Notch1 receptor and activate the Notch1 pathway (Fig.6E-F and Fig.S4C-D).

Treatment with the siNotch1 03, which was the most effective (Fig.S4E), to knock

down Notch1 inhibited the effect of Kallistatin on the activation of the Notch1

signaling pathway, resulting in the downregulation of HES1, upregulation of PPARγ,

and downregulation of BACE1 and Aβ (Fig.6G). HES1, an essential downstream

effector of Notch1 signaling pathway, was reported to suppress the expression

of *PPARG* (gene name of PPARγ) in neuron cells [51, 52]. Similarly, the HES1 shRNA

01, the most effective (Fig.S4F), could upregulate PPARγ and decrease the production

of BACE1 and Aβ when the neuron cells are infected with adenovirus to overexpress

Kallistatin (Fig.6H). The above results suggested that Kallistatin promotes Aβ

production *via* directly binding to the Notch1 receptor and activating the Notch1

pathway.

Kallistatin promoted the phosphorylation of tau by activating the Wnt signaling

pathway.

Glycogen synthase kinase 3-β (GSK-3β) is a crucial element in the

phosphorylation of tau [53]. When the Wnt signaling was activated, LRP6 PPPSPxS

motif could directly interact with GSK3-3β and phosphorylate it [54]. So when the Wnt

signaling was inhibited, GSK3β was activated and dephosphorylated, and non-

phosphorylated GSK3β was able to add phosphate groups to serine/threonine residues

of tau [14]. Kallistatin has already been reported as a competitive inhibitor of the

canonical Wnt signaling pathway [55]. Consistent with previous reports, our results

showed that GSK-3β was activated in the hippocampus of KAL-TG mice (Fig.7A-B).

Furthermore, visible increase in tau phosphorylation with the activation of GSK-3β

stimulated by Kallistatin overexpression (Fig.7C-D), which could be reversed by

LiCl, the inhibitor of GSK-3β (Fig. 7E-F). These results proved that Kallistatin could

promote the phosphorylation of tau by activating the Wnt signaling pathway.

Fenofibrate could alleviate memory and cognitive impairment of KAL-TG mice.

Hyperlipidemia and hyperlipidemia were account for the development of AD [56,

57]. Here, a hypolipidemic drug (fenofibrate) and a hypoglycemic drug (rosiglitazone)

were used to treat the KAL-TG mice (Fig.8A). After a-month treatment with

fenofibrate, the behavioral performance measured by MWM and Y-maze test of

treated group was improved compared to KAL-TG group. The latency to the escape

platform on the fifth training day was significantly decreased (Fig.8B), and crossing

platform times (Fig.8C), time in percent (Fig.8D), and spontaneous alternation

(Fig.8F) were significantly increased in the fenofibrate-treated group, which

performed better than rosiglitazone-treated group. Similarly, the path traces heatmap

indicated mice stayed in the target quadrant longer in the fenofibrate-treated group

(Fiure.8E). Besides, the decreased serum Kallistatin level, Aβ, BACE1,

phosphorylation of tau, and the activation of GSK3β was detected in the fenofibrate-

treated KAL-TG group (Fig.8G-I). However, no significant difference showed in the

rosiglitazone-treated group compared with the KAL-TG mice (Fig.8G-I).

Mechanism summary

In metabolic abnormalities-related AD patients, the concentration of Kallistatin is

elevated, which could increase Aβ deposition through Notch1/HES1/PPARγ/BACE1

pathway, and induce tau hyperphosphorylation by activating GSK-3β. Finally,

elevated Kallistatin caused impaired cognitive memory via inducing Aβ deposition

and tau hyperphosphorylation (Fig.8J).

Discussion

This study demonstrated that Kallistatin is a novel regulator of amyloid-β plaque

accumulation, tau protein hyperphosphorylation, and metabolic abnormalities-related

cognitive memory impairment. We proved that Kallistatin was increased in the serum

of AD and diabetes-related AD patients and the hippocampus of AD model mice. In

addition, the constructed KAL-TG mice defined its cognitive memory impairment

phenotype and lower LTP in hippocampal CA1 neurons accompanied by increased Aβ

deposition and tau phosphorylation. Mechanistically, Kallistatin transcriptionally

upregulates BACE1 expression by suppressing the transcriptional repressor PPARγ,

resulting in Aβ cleavage and production. Most importantly, our studies proved

Kallistatin could bind to the Notch1 receptor directly and activate the Notch1/HES1

pathway, causing the decrease of PPARγ, overproduction of BACE1, and elevated

Aβ42 generation. Moreover, Kallistatin could induce tau phosphorylation by

activating GSK-3β, which result from the inhibition of LRP6. Finally, prolonged

stimulation with high concentrations of Kallistatin could impair cognitive memory in

mice. Lastly, the hypolipidemic drug-fenofibrate decreasing of Aβ expression,

phosphorylation of tau, and the serum Kallistatin level of KAL-TG mice, causing

alleviating memory and cognitive impairment. For the first time, these observations

established an association between high Kallistatin levels and metabolic

abnormalities-related AD, and provided a new drug candidate (fenofibrate) for AD

patients with metabolic syndromes.

Growing shreds of evidence suggest that Diabetes mellitus and AD are closely

linked. Approximately 80% of AD subjects are insulin resistant or T2DM [58];

additionally, the T2DM patients had a higher risk of up to 73% of dementia than the

healthy control population [26]. In line with these observations, the process of cognitive

decline in T2DM patients appears to begin in the prediabetic phase of insulin

resistance [59, 60]. We performed a GAD disease enrichment analysis of differentially

expressed genes in neurons of T2DM and normal controls, which showed AD is

closely related to T2DM, besides identified enrichment of the Serpin family protein

domain using PFAM analysis (Fig.S1A, B). We and other researchers found that

Kallistatin (which belongs to the Serpin family) was increased in T2DM [32, 37].

Although the interrelationship between Kallistatin and AD has not been reported to

date, we have enough reason to speculate Kallistatin might be one of the critical

molecules which could establish the relationship between Alzheimer's disease and

Diabetes. Our data suggested that AD patients indeed have metabolic disorders and

elevated Kalllistatin (Fig.1A-D, S1C-D). In addition, memory cognitive function and

synaptic plasticity of KAL-TG mice were impaired (Fig.1E-J). These results

suggested that Kallistatin was a new link between AD and diabetes.

A hallmark of AD is the aggregation of Aβ into amyloid plaques and tau

phosphorylation in patients' brains. Aβ, a small peptide with a high propensity to form

aggregates, is widely believed to be central and initial to the pathogenesis of the

disease [61]. Correspondingly, we discovered Kallistatin could lead to Aβ

overproduction through Notch1/HES1/PPARγ/BACE1 signaling pathway. GSK-3β, a

vital kinase regulating the process of tau phosphorylation [53], could be activated by

inhibiting the Wnt signaling pathway. Kallistatin was already promoted as a

competitive inhibitor of LRP6, the Wnt receptor [55]. Consistent with previous reports,

Kallistatin could increase tau phosphorylation by activating GSK-3β. We

demonstrated for the first time that Kallistatin could promote AD by increasing Aβ

production and tau phosphorylation in the central nervous system.

Previous studies have shown that Notch signaling is closely related to AD. For

example, NOTCH mutation was reported to cause AD-like pathology [62]. And Notch1

was found to increase in AD patients [63]. Besides, the Notch1/HES1 signaling pathway

was reported to suppress the expression of PPARγ [64]. Our previous studies proved that

Kallistatin could activate Notch1 signaling [37]. Consequently, we detect Notch1 in our

animal and cell models. Indeed, Notch1 was upregulated by Kallistatin (Fig.6, S4), as

well as Aβ deposition. Notch signaling is initiated by receptor-ligand interaction at the

cell surface. In mammals, there are five ligands encoded by JAG1, JAG2, DLL1,

DLL3, and DLL4 [65]. Here, our results found that Kallistatin could activate Notch1 by

binding to it directly, which means Kallistatin was a new ligand of the Notch1

receptor.

Treatment of AD has always been a hot and challenging problem in neurology.

Multiple strategies have been proposed to reduce the pathogenicity of Aβ and tau.

Unfortunately, several Aβ-targeted therapies tested in the phase III clinical trials have

failed as they have not been able to slow down cognitive decline, although they could

effectively reduce the Aβ load [66, 67, 68, 69, 70, 71]. BACE1 inhibitors were failed to

improve the cognitive function of AD patients but also resulted in clinical

deterioration and impairment of liver function [72, 73, 74]. Here are two possible reasons

for the failure of of clinical trials with the BACE1 inhibitor. First, the reduction of

BACE1 activity could lead to an accumulation of full-length APP [75]. Second, the size

of the BACE1 active site is relatively large; the use of a small molecule may not be

sufficient to occupy the active site [76]. Therefore, synaptic damage caused by BACE1

inhibitors or the insufficient effect of BACE1 inhibitors may lead to the failure of

clinical trials. Therapeutic strategies targeting tau include tau aggregation blockers

(TRx0014, TRx0237), antibody vaccine therapy (e.g., RO7105705, BIIB092),

inhibition of tau phosphorylation (Anavex2-73), and microtubule stabilizers

(Anavex2-73) [77]. Some of the aforementioned drugs have been partially stopped,

while others are still undergoing a clinical test and shown protective benefits. There

are still several obstacles to the commercialization of tau treatments when they reach

maturity.

Because of the failures of clinical trials, some researchers propose an alternative

option for AD therapeutics to address the modifiable risk factors for developing AD,

such as type 2 diabetes [78, 79, 80]. Our previous study proved that Kallistatin is a

multifunctional protein strongly associated with diabetes, and Kallistatin neutralizing

antibody improves diabetic wound healing [44]. This study demonstrated that Kallistatin

induced memory cognitive dysfunction by promoting Aβ deposition and tau

phosphorylation. Thus, we speculate the increased Kallistatin is a decent candidate for

T2DM-related AD therapy. PPARγ is a ligand-activated transcription factor and a

master modulator of glucose and lipid metabolism, organelle differentiation, and

inflammation [81, 82]. Growing evidence revealed that PPARγ agonists (rosiglitazone)

could rescue memory impairment of AD model mice [83, 84, 85]. In clinical trials, it is

controversial whether rosiglitazone has a protective effect on memory cognitive

function [86, 87, 88]. In this study, although Aβ expression had a downward trend, the

memory and cognition of KAL-TG mice were unchanged after treatment with

rosiglitazone for a month (Fig.8). This might be caused by insufficient treatment time

and the unchanged Kallistatin level.

Fenofibrate is a fibric acid derivative for clinically lowering blood lipids, mainly

triglycerides [89]. Studies showed that fenofibrate could prevent memory disturbances,

maintain hippocampal neurogenesis, and protect against Parkinson's disease (PD) [90,

91]. Specifically, fenofibrate has a neuroprotective effect on memory impairment

induced by Aβ through targeting α- and β- secretase [92]. Recently, our study proved

that the fenofibrate could repair the disrupted glutamine-glutamate cycle by

upregulating glutamine synthetase, while there is currently no fenofibrate treatment of

AD in clinical trials[38]. In this study, we proved fenofibrate was beneficial for memory

and cognitive impairment of KAL-TG mice. In addition, Aβ, BACE1, phosphorylated

tau and serum Kallistatin level of KAL-TG mice could be downregulated after the

treatment of fenofibrate. All of these suggested that fenofibrate might be helpful for

metabolic abnormalities-related AD patients. Therefore, fenofibrate administration in

patients with metabolic syndrome play an early role in preventing and treating AD.

In summary, we affirmed that Kallistatin concentrations were increased in

diabetes-related AD patients. In addition, our study demonstrated for the first time that

Kallistatin positively regulated Aβ42 through Notch1/HES1/PPARγ/BACE1, and

increased phosphorylated tau through inhibition of the Wnt signaling pathway.

Kallistatin might play a crucial role in linking diabetes and cognitive memory

deterioration. Moreover, fenofibrate could decrease the serum Kallistatin level,

BACE1, Aβ, and phosphorylated tau of KAL-TG mice, leading to alleviating memory

and cognitive impairment. These findings might provide new insight into AD and

possibly other neurodegenerative disorders.

Materials and methods

Ethics approval and consent to participate

All patients involved in this study gave their informed consent. This study

obtained the institutional review board approval of Medical Ethics of Zhongshan

Medical College No. 072 in 2021 and Animal Experiment Ethics of Zhongshan

Medical College, Approval No.: SYSU-IACUC-2019-B051.

Human Subjects

The study was approved by the experimental ethics committee of Guangdong

Academy of Medical Sciences and Sun Yat-sen University and carried out in strict

accordance with the ethical principles, and each participant was provided written informed consent before collecting samples. We certify that the study was performed in accordance with the 1964 declaration of HELSINKI and later amendments. We collected 61 normal human samples, 56 AD patient samples, of whom 36 normal human samples were from the Zhongshan City People's Hospital; 14 normal human samples and 22 AD patient samples were from Zhongshan Third People's Hospital；11 normal human samples and 14 AD patient samples were from Sun Yat-sen Memorial Hospital. AD patient was clinically diagnosed according to ICD-10 (International Classification of Diseases) and NINCDS-ADRDA (the National Institute of Neurological and Communicative Disorders and Stroke and the Alzheimer's Disease and Related Disorders Association) criteria, and 20 AD patient samples were clinically diagnosed according to MMSE (Mini-Mental State Examination), were collected from Guangdong Provincial People's Hospital. All subjects' Clinical characteristics were presented in Supplementary Tables (Table S1-2).

Experimental Animals and Protocols

All animal experiment procedures were carried out in an environment without specific pathogens (Specific pathogen free, SPF) with the approval of the Animal Care and Use Committee of Sun Yat-sen University (approval ID: SYXK 2015-0107). The wild type mice (WT, C57BL/6) were purchased from the Animal Center of Guangdong Province (Production license No.: SCXK 2013-0002, Guangzhou, China).

The SAMR1 and SAMP8 mice (7 months old) were purchased from Tianjin

University of Traditional Chinese Medicine (Tianjin, China). Kallistatin transgenic

mice (KAL-TG) were C57BL/6 strain provided by Dr. Jianxing Ma (University of

Oklahoma Health Sciences Center) [39]. The KAL-TG mice genotype was identified by

PCR technology (forward primer: 5'-AGGGAAG-ATTGTGGATTTGG-3', reverse

primer: 5'-ATGAAGATACCAGTGATGCTC-3'). KAL-TG mice aged 6 months were

randomly divided into three groups: control group (KAL-TG), fenofibrate-treated

group (KAL-TG-Feno, 0.3 g/kg/d), and rosiglitazone-treated group (KAL-TG-RSG,

0.005 g/kg/d). Fenofibrate (Sigma-Aldrich, cat. no. F6020) and rosiglitazone

(Selleck, cat. no. S2556) were administered to mice by oral gavage. In three groups,

the serum Kallistatin were examined in the 0 week and 4 week after drug treatment

from the blood taken from mouse orbit. In addition, the Morris water maze and Y-

maze test were performed one week after the second blood collection.

Morris water maze (MWM)

The KAL-TG and WT mice were employed for the Morris water maze test

including the behavioral test, latency experiment (for 6 days), and the probe test (the

7th day). In addition, the MWM was performed as described previously [40]. Mice were

brought into the testing room and handled for 1 day before the training experiment. In

the 6-day training experiment, each mouse was trained with four daily trials. The mice

facing the wall were placed into the maze, exploring the maze from different

directions (east, south, west, and north). This trial was completed as soon as the

mouse found the platform, or 90 s elapsed. If the mice could discover and climb the

submerged platform within 90 s, the system would automatically record the latency

time and path immediately, and then the mouse was guided to and placed on the

submerged platform for extra 20 s. On day 7, the platform was removed, and a probe

test was performed to examine the strength and integrity of the animal spatial memory

24 h after the last testing trial. During the probe test, the mice were gently brought

into the water from the fixed monitoring point, and the mice were allowed to swim for

90 s without the platform. Finally, all of the measured behavioral parameters were

analyzed using SMART software.

Y-maze test

A Y maze test was performed to assess the mice's spatial memory. The Y maze

was separated by 120°, consisting of three identical arms (30 cm long, 7 cm wide, and

15 cm high) made of blue PVC. The mice were placed first in one of the arms, and

over the next 10 minutes, the sequence and number of their entry into the three arms

were monitored. An alternation is defined when a mouse visits three straight arms

(namely, ABC, BCA, or CAB, but not ABA, BAB, or CAC). Spontaneous alternation

(%) = [(number of alternations)/(total number of arms−2)]× 100.

Electrophysiology

Hippocampal slices (300-400 μm) from KAL-TG and WT mice were cut as

described [41]. Coronal slices from hippocampus (400 μm thick) were prepared from

different age groups KAL-TG mice and their WT littermates using a tissue slicer

(Vibratome 3000; Vibratome) in ice-cold dissection buffer containing the following

(in mM): 212.7 sucrose, 3 KCl, 1.25 NaH_2PO_4, 3 $MgCl_2$, 1 $CaCl_2$, 26 $NaHCO_3$, and

10 dextrose, bubbled with 95% O_2/5% CO_2. The slices were immediately transferred

to ACSF at 35 °C for 30 min before recordings. The recipe of ACSF was similar to the

dissection buffer, except that sucrose was replaced with 124 mM NaCl, and the

concentrations of $MgCl_2$ and $CaCl_2$ were changed to 1 mM and 2 mM, respectively.

All recordings were performed at 28-30 °C. Pyramidal cells in CA1 areas were

identified visually under infrared differential interference contrast optics based on

their pyramidal somata and prominent apical dendrites. Whole-cell was performed

using an Integrated Patch-Clamp Amplifier (Sutter Instrument, Novato, CA, USA)

controlled by Igor 7 software (WaveMetrics, Portland, OR, USA) filtered at 5 kHz and

sampled at 20 kHz. Igor 7 software was also used for acquisition and analysis. Only

cells with series resistance <20 MΩ and input resistance >100 MΩ were studied. Cells

were excluded if input resistance changed >15% or series resistance changed >10%

over the experiment. A concentric bipolar stimulating electrode with a tip diameter of

125 μm (FHC) was placed in the stratum radiatum. The recording and stimulating

electrode distances were kept at 50-100 μm. Patch pipettes (2-4MΩ) were filled with

the internal solution consisting of the following (in mM): 120 Cs-methylsulfonate, 10

Na-phosphocreatine, 10 HEPES, 4 ATP, 5 lidocaine N-ethyl bromide (QX-314), 0.5

GTP; the pH of the solution was 7.2–7.3, and the osmolarity was 270-285 mOsm.

To induce LTP, a pairing protocol was applied. In brief, conditioning stimulation

consisted of 360 pulses at 2 Hz paired with continuous postsynaptic depolarization

(180 s) to 0 mV. 50 μM picrotoxin was added to the recording bath to suppress

excessive polysynaptic activity, and the concentration of Ca^{2+} and Mg^{2+} was elevated

to 4 mM to reduce the recruitment of polysynaptic responses. A test pulse was

delivered at 0.067 Hz to monitor baseline amplitude for 10 min before and 30 min

following paired stimulation. To calculate LTP, the EPSC amplitude was normalized

to the mean baseline amplitude during 10 min baseline. Potentiation was defined as

the mean normalized EPSC amplitude 25–40 min after paired stimulation.

ELISA

To quantify serum Kallistatin, the collected samples were centrifuged at 4 °C for

10min at 5000 rpm. It was detected using the KBP ELISA kit (#DY1669, R&D

systems, MN, USA) as per the instructions of the manufacturer. The levels of Aβ42 in

brain tissue produced from mouse primary neuron cells and HT22 cells were

measured with a mouse Aβ42 Elisa Kit (27721, IBL, Germany). To measure Aβ42 in

brain tissue, 0.05 g of mouse brain tissues were weighed and homogenized using 2ml

PBS with a protease inhibitor (cocktail). After centrifugalization at 4 °C for 30 min at

12000 g, the extracts' supernatants were analyzed using the ELISA method after total

protein quantification. To quantify levels of Aβ42 produced from primary neuron

cells, the cell supernatants were ultrafiltrated with an Ultrafiltration tube (4-kD

Millipore), centrifugalization, and testing. Cell homogenate was prepared in 1ml PBS

with a protease inhibitor (cocktail) and quantified using the BCA method before being

measured by ELISA.

Immunohistochemistry

Tissue slices were prepared as described before [42]. The sections were incubated

with Aβ (ab201060, Abcam, Cambridge, UK), BACE1 (#5606S, Cell Signaling

Technology, Boston, USA), PPARγ (#2435, Cell Signaling Technology, Boston,

USA), Notch1 (#3608, Cell Signaling Technology, Boston, USA), p-tau S202

(ab108387, Abcam, Cambridge, UK), p-tau T231(ab151559, Abcam, Cambridge,

UK), p-tau S396 (ab109390, Abcam, Cambridge, UK), tau (ab75714, Abcam,

Cambridge, UK) antibodies overnight at 4°C and then incubated with Alexa Fluor

488-donkey anti-rabbit IgG (H + L) (1:200, Life Technologies, Gaithersburg, MD,

USA, #A21208) for 1h, then incubated with a biotin-conjugated secondary antibody

for 30 min, followed by incubation with DAB for 10 s and hematoxylin staining for

30 s. The IHC signals were analyzed using ImageJ.

Cell culture experiments

495 HT22 cells were purchased from the Cell Bank of the Chinese Academy of

496 Sciences (Shanghai, China). HT22 cells were cultured and grown to confluence in

497 DMEM supplemented with 10% FBS (Gibco BRL), 100 U/mL penicillin, and 100

498 U/mL streptomycins (Gibco BRL).

Primary culture of hippocampal neurons

Primary neurons were obtained from the hippocampus of C57/BL6J mice at age

1-3 days. Before culturing, the newborn pup was euthanized and dipped into 70%

ethanol for 3 min. First, the infant pup hippocampus was isolated with eye tweezers

observed under the stereomicroscope, and excess soft tissue was removed. Second,

hippocampal tissue in PBS buffer was cut up with scissors gently and blown with a

1ml pipette until it was not visible. Next, the cell suspension was transferred to a 15ml

centrifuge tube and centrifuged at 1000 rpm for 5 min at room temperature. Cell

precipitation was suspended and cultured with 2-3mL primary neural stem cell (NSC)

suspension (Thermo Fisher Scientific, 21103049) in a 37 °C, 5% CO_2 cell incubator

for 3 days, changing half medium every 2 days. After 7 days, the cell suspension was

transferred to a 15ml centrifuge tube, centrifuged, and recultured with neurobasal,

10%FBS, 1:50 B27(Thermo Fisher Scientific, A3582801), and 1:100 bFGF (Thermo

Fisher Scientific, #RP-8626). One day later, the medium was changed to neurobasal

(2%FBS, 1:50 B27, and 1:100 bFGF), culturing for 21 more days. The

immunofluorescence technique was used with the neuron-specific marker (MAP2,

#4542, Cell Signaling Technology, Boston, USA) to determine the purity of neurons.

siRNA, shRNA, and adenovirus transfection

Notch1 siRNA and control siRNA were purchased from RiboBio (Guangzhou,

China). HES1 shRNA and control shRNA were purchased from Qingke (Guangzhou,

China). Green fluorescent protein-adenovirus (Ad-GFP) and Kallistatin-adenovirus

(Ad-KAL) were provided by Dr. Jianxing Ma (University of Oklahoma Health

Sciences Center). According to the manufacturer's instructions, the transfections were

performed at approximately 60% confluency using Lipofectamine®3000 transfection

reagent (Invitrogen) or RNAiMAX. After 24 h, interference confirmation was

conducted using real-time quantitative PCR (RT-qPCR) and Western blot.

RNA isolation and quantitative RT-PCR

Total RNA extraction, reverse transcription of cDNA, and real-time quantitative

PCR were performed as described previously [43]. BACE1 forward:

GGAGCCCTTCTTTGACTCCC; BACE1 reverse:

CAATGATCATGCTCCCTCCCA; ADAM9 forward:

GGAAGGCTCCCTACTCTCTGA; ADAM9 reverse: CAATTCC-

AAAACTGGCATTCTCC; ADAM10 forward: ATGGTGTTGCCGACAGTGTTA;

ADAM10 reverse: GTTTGGCACGCTGGTGTTTTT; ADAM17 forward: GGAT-

CTACAGTCTGCGACACA; ADAM17 reverse:

TGAAAAGCGTTCGGTACTTGAT; β-actin forward:

GCACTCTTCCAGCTTCCTT; β-actin reverse: GTTGGCGTACAG-GTCTTTGC.

Western blot

Western blot was performed as described previously [40, 43]. Equal amounts of

protein were subjected to western blot analysis. Blots were probed with antibodies

against Kallistatin (ab187656, Abcam, Cambridge, UK), Aβ (ab201060, Abcam,

Cambridge, UK), Presenilin-1 (ab76083, Abcam, Cambridge, UK), BACE1 (#5606S,

Cell Signaling Technology, Boston, USA), APP (#2452S, Cell Signaling Technology,

Boston, USA), MAP2 (#4542, Cell Signaling Technology, Boston, USA), PPARγ

（#2435, Cell Signaling Technology, Boston, USA), SP1 (#9389, Cell Signaling

Technology, Boston, USA), YY1 (#46395, Cell Signaling Technology, Boston,

USA）, Notch1 (#3608, Cell Signaling Technology, Boston, USA), Hes1 (#11988,

Cell Signaling Technology, Boston, USA), p-tau S202 (ab108387, Abcam,

Cambridge, UK), p-tau T231(ab151559, Abcam, Cambridge, UK), p-tau S396

(ab109390, Abcam, Cambridge, UK), tau (ab75714, Abcam, Cambridge, UK),

GSK3β(#70109S, Cell Signaling Technology, Boston, USA), p-GSK3β Ser9 (#9323,

Cell Signaling Technology, Boston, USA), β-actin (A5441-2ml, Sigma, CA, USA),

anti-Mouse (#PI200, Vector Laboratories, Burlingame, CA, USA), anti-Rabbit

(#PI1000, Vector Laboratories, Burlingame, CA, USA). The signal intensity was

quantified using ImageJ (NIH).

Statistical Analysis

The results are expressed as mean ± SD. Student's t-test was applied for

comparisons of parametric data between two groups, and one-way ANOVA followed

by LSD t-test was used to compare differences between more than two different

groups (GraphPad Prism software). A P value less than 0.05 was considered statistical

significance.

List of abbreviations

Aβ	amyloid β
p-tau	hyperphosphorylated tau
AD	Alzheimer's disease
T2DM	type 2 diabetes mellitus

FBG	fasting blood glucose
TG	triglyceride
KAL-TG	Kallistatin-transgenic
WT	wild type mice
APP	amyloid precursor protein
BACE1	β-site APP cleaving enzyme 1
BMI	body mass index
ICD-10	The International Statistical Classification of Diseases and Related Health Problems 10th Revision
NINCDS-ADRDA	the National Institute of Neurological and Communicative Disorders and Stroke and the Alzheimer's Disease and Related Disorders Association
MMSE	mini-Mental State Examination
KAL-TG-RSG	rosiglitazone-treated group
KAL-TG-Feno	fenofibrate-treated group
MWM	morris water maze
RT-qPCR	real-time quantitative PCR

Declarations

Ethics approval and consent to participate

All patients involved in this study gave their informed consent. This study

obtained the institutional review board approval of Medical Ethics of Zhongshan

Medical College No. 072 in 2021 and Animal Experiment Ethics of Zhongshan

Medical College, Approval No.: SYSU-IACUC-2019-B051.

Consent for publication

Not applicable.

Availability of data and materials

All the data supporting the conclusions of the current study are presented in the figures and they are available from the corresponding authors upon reasonable request. There are no restrictions on data availability. Source data are provided with this paper.

Competing interests

The authors declare that they have no competing interests.

Acknowledgements

We are thankful to Zhongshan City People's Hospital, Zhongshan Third People's Hospital, Guangdong Provincial People's Hospital, and Sun Yat-sen Memorial Hospital for kindly providing serum samples for the analyses of this manuscript. We thank Professor Boxing Li for his valuable advice on our study.

Funding

This study was supported by The National Natural Science Foundation of China (Grants 82070888, 82070882, 82100917, 82273116, 82203661, 81901557, 81870869, and 32071101); Guangdong Natural Science Fund (Grant 2022A1515012423, 2021A1515010434, 2023A1515010316, and 2023A1515010214); Key Sci-Tech Research Project of Guangzhou Municipality, China (Grants 202201010820, 202102020955); Key Project of Nature Science Foundation of Guangdong Province,

China (Grant 2019B1515120077); National Key R&D Program of China (Grant

2018YFA0800403); Guangdong Special Support Program for Young Top Scientist

(Grant 201629046); China Postdoctoral Science Foundation (Grant 2021M703679,

2022M713594, BX20220360); Health Commission of Guangdong Province (Grants

A2022161) ; Guangzhou Municipal Science and Technology Bureau (Grants

SL2022A04J01575); Guangdong Key Project in "Development of new tools for

diagnosis and treatment of Autism" (2018B030335001), Research and Development

Plan of Key Areas of Guangzhou Science and Technology Bureau (2020070030001),

Open Research Funds of State Key Laboratory of Ophthalmology (2020KF03);

Guangzhou Key Laboratory for Metabolic Diseases(202102100004); The Science and

Technology Planning Project of Guangdong Province 2023B1212060018.

References

1. Grundke-Iqbal I, Iqbal K, Tung YC, Quinlan M, Wisniewski HM, Binder LI. Abnormal phosphorylation of the microtubule-associated protein tau (tau) in Alzheimer cytoskeletal pathology. *Proceedings of the National Academy of Sciences of the United States of America* 1986, **83**(13)**:** 4913-4917.

2. Haass C, Selkoe DJ. Soluble protein oligomers in neurodegeneration: lessons from the Alzheimer's amyloid beta-peptide. *Nature Reviews Molecular Cell Biology* 2007, **8**(2)**:** 101-112.

3. Holtzman DM, Morris JC, Goate AM. Alzheimer's Disease: The Challenge of the Second Century. *Science Translational Medicine* 2011, **3**(77).

4. Leng F, Edison P. Neuroinflammation and microglial activation in Alzheimer disease: where do we go from here? *Nature Reviews Neurology* 2021, **17**(3)**:** 157-172.

5. De Strooper B, Karran E. The Cellular Phase of Alzheimer's Disease. *Cell* 2016, **164**(4)**:** 603-615.

6. Tomita T. Molecular mechanism of intramembrane proteolysis by gamma-secretase. *Journal of*
Biochemistry 2014, **156**(4)**:** 195-201.

7. LaFerla FM, Green KN, Oddo S. Intracellular amyloid-beta in Alzheimer's disease. *Nature Reviews*
Neuroscience 2007, **8**(7)**:** 499-509.

8. Das B, Yan R. A Close Look at BACE1 Inhibitors for Alzheimer's Disease Treatment. *Cns Drugs*
2019, **33**(3)**:** 251-263.

9. Crunkhorn, Sarah. Alzheimer disease: BACE1 inhibitor reduces β-amyloid production in humans.
Nature Reviews Drug Discovery 2016, **16**(1)**:** 18-18.

10. Yan R, Vassar R. Targeting the β secretase BACE1 for Alzheimer's disease therapy. *Lancet*
Neurology 2014, **13**(3)**:** 319-329.

11. Wang Y, Mandelkow E. Tau in physiology and pathology. *Nature reviews Neuroscience* 2016,
17(1)**:** 5-21.

12. Hausrat TJ, Janiesch PC, Breiden P, Lutz D, Hoffmeister-Ullerich S, Hermans-Borgmeyer I*, et al.*
Disruption of tubulin-alpha4a polyglutamylation prevents aggregation of hyper-phosphorylated tau
and microglia activation in mice. *Nature communications* 2022, **13**(1)**:** 4192.

13. Braak E, Braak H, Mandelkow EM. A sequence of cytoskeleton changes related to the formation of
neurofibrillary tangles and neuropil threads. *Acta neuropathologica* 1994, **87**(6)**:** 554-567.

14. Boonen RA, van Tijn P, Zivkovic D. Wnt signaling in Alzheimer's disease: up or down, that is the
question. *Ageing Res Rev* 2009, **8**(2)**:** 71-82.

15. Biessels GJ, Despa F. Cognitive decline and dementia in diabetes mellitus: mechanisms and clinical
implications. *Nature Reviews Endocrinology* 2018, **14**(10)**:** 591-604.

16. Li W, Huang E, Gao S. Type 1 Diabetes Mellitus and Cognitive Impairments: A Systematic Review.
Journal of Alzheimers Disease 2017, **57**(1)**:** 29-36.

17. Cardoso SM, Correia SC, Carvalho C, Moreira PI. Mitochondria in Alzheimer's Disease and
Diabetes-Associated Neurodegeneration: License to Heal! In: Singh H, Sheu SS (eds).
Pharmacology of Mitochondria, vol. 240, 2017, pp 281-308.

18. Schipper HM. Apolipoprotein E: Implications for AD neurobiology, epidemiology and risk
assessment. *Neurobiology of Aging* 2011, **32**(5)**:** 778-790.

19. Folch J, Patraca I, Martinez N, Pedros I, Petrov D, Ettcheto M*, et al.* The role of leptin in the

sporadic form of Alzheimer's disease. Interactions with the adipokines amylin, ghrelin and the
pituitary hormone prolactin. *Life Sciences* 2015, **140:** 19-28.
661

662 20.Association As. 2016 Alzheimer's disease facts and figures. *Alzheimer's & Dementia* 2016, **12**(4)**:**
663   459-509.
664

665 21.Baglietto-Vargas D, Shi J, Yaeger DM, Ager R, LaFerla FM. Diabetes and Alzheimer's disease
666   crosstalk. *Neuroscience & Biobehavioral Reviews* 2016, **64:** 272-287.
667

668 22.Kandimalla R, Thirumala V, Reddy PH. Is Alzheimer's disease a Type 3 Diabetes? A critical
669   appraisal. *Biochimica Et Biophysica Acta-Molecular Basis of Disease* 2017, **1863**(5)**:** 1078-1089.
670

671 23.Zhang Y, Huang N-q, Yan F, Jin H, Zhou S-y, Shi J-s*, et al.* Diabetes mellitus and Alzheimer's
672   disease: GSK-3 beta as a potential link. *Behavioural Brain Research* 2018, **339:** 57-65.
673

674 24.Steen E, Terry BM, Rivera EJ, Cannon JL, Neely TR, Tavares R*, et al.* Impaired insulin and insulin-
675   like growth factor expression and signaling mechanisms in Alzheimer's disease - is this type 3
676   diabetes? *Journal of Alzheimers Disease* 2005, **7**(1)**:** 63-80.
677

678 25.Candasamy M, Mohamed Elhassan SA, Kumar Bhattamisra S, Hua WY, Sern LM, Binti Busthamin
679   NA*, et al.* Type 3 diabetes (Alzheimer's disease): new insight for promising therapeutic avenues.
680   *Panminerva Medica* 2020, **62**(3)**:** 155-163.
681

682 26.Koekkoek PS, Kappelle LJ, van den Berg E, Rutten GE, Biessels GJ. Cognitive function in patients
683   with diabetes mellitus: guidance for daily care. *The Lancet Neurology* 2015, **14**(3)**:** 329-340.
684

685 27.Hildreth KL, Van Pelt RE, Schwartz RS. Obesity, insulin resistance, and Alzheimer's disease.
686   *Obesity (Silver Spring, Md)* 2012, **20**(8)**:** 1549.
687

688 28.Anjum I, Fayyaz M, Wajid A, Sohail W, Ali A. Does obesity increase the risk of dementia: a
689   literature review. *Cureus* 2018, **10**(5).
690

691 29.Ma C, Luo C, Yin H, Zhang Y, Xiong W, Zhang T*, et al.* Kallistatin inhibits lymphangiogenesis and
692   lymphatic metastasis of gastric cancer by downregulating VEGF-C expression and secretion.
693   *Gastric Cancer* 2018, **21**(4)**:** 617-631.
694

695 30.Yang Y, He X, Cheng R, Chen Q, Shan C, Chen L*, et al.* Diabetes-induced upregulation of
696   kallistatin levels exacerbates diabetic nephropathy via RAS activation. *Faseb Journal* 2020, **34**(6)**:**
697   8428-8441.
698

31.Ma C, Yin H, Zhong J, Zhang Y, Luo C, Che D*, et al.* Kallistatin exerts anti-lymphangiogenic
effects by inhibiting lymphatic endothelial cell proliferation, migration and tube formation.

International Journal of Oncology 2017, **50**(6)**:** 2000-2010.

32.El-Asrar MA, Andrawes NG, Ismail EA, Salem SMH. Kallistatin as a marker of microvascular

complications in children and adolescents with type 1 diabetes mellitus: Relation to carotid intima

media thickness. *Vascular Medicine* 2015, **20**(6)**:** 509-517.

33.Gateva A, Assyov Y, Velikova T, Kamenov Z. Increased kallistatin levels in patients with obesity

and prediabetes compared to normal glucose tolerance. *Endocrine Research* 2017, **42**(2)**:** 163-168.

34.Campbell DJ, Kladis A, Zhang Y, Jenkins AJ, Prior DL, Yii M*, et al.* Increased tissue kallikrein

levels in type 2 diabetes. *Diabetologia* 2010, **53**(4)**:** 779-785.

35.Nowicki GJ, Ślusarska B, Polak M, Naylor K, Kocki T. Relationship between Serum Kallistatin and

Afamin and Anthropometric Factors Associated with Obesity and of Being Overweight in Patients

after Myocardial Infarction and without Myocardial Infarction. *Journal of clinical medicine* 2021,

10(24).

36.Hong S, Han K, Park CY. The insulin resistance by triglyceride glucose index and risk for dementia:

population-based study. *Alzheimer's research & therapy* 2021, **13**(1)**:** 9.

37.Feng J, Dong C, Long YL, Mai LF, Ren M, Li LY*, et al.* Elevated Kallikrein-binding protein in

diabetes impairs wound healing through inducing macrophage M1 polarization. *Cell*

Communication and Signaling 2019, **17**.

38.Long Y, Zhao Z, Xie W, Shi J, Yang F, Zhu D*, et al.* Kallistatin leads to cognition impairment via

downregulating glutamine synthetase. *Pharmacol Res* 2024, **202:** 107145.

39.McBride JD, Jenkins AJ, Liu XC, Zhang B, Lee K, Berry WL*, et al.* Elevated Circulation Levels of

an Antiangiogenic SERPIN in Patients with Diabetic Microvascular Complications Impair Wound

Healing through Suppression of Wnt Signaling. *Journal of Investigative Dermatology* 2014, **134**(6)**:**

1725-1734.

40.Huang M, Qi WW, Fang SH, Jiang P, Yang C, Mo YS*, et al.* Pigment Epithelium-Derived Factor

Plays a Role in Alzheimer's Disease by Negatively Regulating A beta 42. *Neurotherapeutics* 2018,

15(3)**:** 728-741.

41.Guo D, Peng Y, Wang L, Sun X, Wang X, Liang C*, et al.* Autism-like social deficit generated by

Dock4 deficiency is rescued by restoration of Rac1 activity and NMDA receptor function. *Mol*

Psychiatry 2021, **26**(5)**:** 1505-1519.

42.Li C, Huang Z, Zhu L, Yu X, Gao T, Feng J*, et al.* The contrary intracellular and extracellular

functions of PEDF in HCC development. *Cell Death & Disease* 2019, **10**.

43. Jiang P, Huang M, Qi WW, Wang FH, Yang TY, Gao TX, *et al.* FUBP1 promotes neuroblastoma proliferation via enhancing glycolysis-a new possible marker of malignancy for neuroblastoma. *Journal of Experimental & Clinical Cancer Research* 2019, **38**(1).

44. Feng J, Dong C, Long Y, Mai L, Ren M, Li L, *et al.* Elevated Kallikrein-binding protein in diabetes impairs wound healing through inducing macrophage M1 polarization. *Cell Communication and Signaling* 2019, **17**.

45. De Strooper B, Annaert W. Proteolytic processing and cell biological functions of the amyloid precursor protein. *Journal of Cell Science* 2000, **113**(11)**:** 1857-1870.

46. LaFerla FM, Oddo S. Alzheimer's disease: A beta, tau and synaptic dysfunction. *Trends in Molecular Medicine* 2005, **11**(4)**:** 170-176.

47. Scheuner D, Eckman C, Jensen M, Song X, Citron M, Suzuki N, *et al.* Secreted amyloid beta-protein similar to that in the senile plaques of Alzheimer's disease is increased in vivo by the presenilin 1 and 2 and APP mutations linked to familial Alzheimer's disease. *Nature Medicine* 1996, **2**(8)**:** 864-870.

48. Christensen MA, Zhou WH, Qing H, Lehman A, Philipsen S, Song WH. Transcriptional regulation of BACE1, the beta-amyloid precursor protein beta-secretase, by Sp1. *Molecular and Cellular Biology* 2004, **24**(2)**:** 865-874.

49. Lin N, Chen L-m, Pan X-d, Zhu Y-g, Zhang J, Shi Y-q, *et al.* Tripchlorolide Attenuates beta-amyloid Generation via Suppressing PPAR gamma-Regulated BACE1 Activity in N2a/APP695 Cells. *Molecular Neurobiology* 2016, **53**(9)**:** 6397-6406.

50. Nowak K, Lange-Dohna C, Zeitschel U, Gunther A, Luscher B, Robitzki A, *et al.* The transcription factor Yin Yang 1 is an activator of BACE1 expression. *Journal of Neurochemistry* 2006, **96**(6)**:** 1696-1707.

51. Maniati E, Bossard M, Cook N, Candido JB, Emami-Shahri N, Nedospasov SA, *et al.* Crosstalk between the canonical NF-κB and Notch signaling pathways inhibits Pparγ expression and promotes pancreatic cancer progression in mice. *The Journal of clinical investigation* 2011, **121**(12)**:** 4685-4699.

52. Herzig S, Hedrick S, Morantte I, Koo SH, Galimi F, Montminy M. CREB controls hepatic lipid metabolism through nuclear hormone receptor PPAR-gamma. *Nature* 2003, **426**(6963)**:** 190-193.

53. Kanno T, Tsuchiya A, Tanaka A, Nishizaki T. Combination of PKCε Activation and PTP1B Inhibition Effectively Suppresses Aβ-Induced GSK-3β Activation and Tau Phosphorylation. *Mol*

Neurobiol 2016, **53**(7)**:** 4787-4797.

54. Piao S, Lee SH, Kim H, Yum S, Stamos JL, Xu Y*, et al.* Direct inhibition of GSK3beta by the
phosphorylated cytoplasmic domain of LRP6 in Wnt/beta-catenin signaling. *PLoS One* 2008, **3**(12)**:**
e4046.

55. Liu X, Zhang B, McBride JD, Zhou K, Lee K, Zhou Y*, et al.* Antiangiogenic and
antineuroinflammatory effects of kallistatin through interactions with the canonical Wnt pathway.
Diabetes 2013, **62**(12)**:** 4228-4238.

56. Kubis-Kubiak A, Wiatrak B, Piwowar A. Hyper-glycemia and insulinemia induce morphological
changes and modulate secretion of S100B, S100A8, amyloid β 1-40 and amyloid β 1-42, in a model
of human dopaminergic neurons. *Biomedicine & pharmacotherapy = Biomedecine &*
pharmacotherapie 2022, **156:** 113869.

57. Rivas-Domínguez A, Mohamed-Mohamed H, Jimenez-Palomares M, García-Morales V, Martinez-
Lopez L, Orta ML*, et al.* Metabolic Disturbance of High-Saturated Fatty Acid Diet in Cognitive
Preservation. *International journal of molecular sciences* 2023, **24**(9).

58. Suzanne M. Type 3 diabetes is sporadic Alzheimer′s disease: mini-review. *European*
Neuropsychopharmacology 2014, **24**(12)**:** 1954-1960.

59. Biessels GJ, Reagan LP. Hippocampal insulin resistance and cognitive dysfunction. *Nature reviews*
Neuroscience 2015, **16**(11)**:** 660-671.

60. Xu W, Caracciolo B, Wang HX, Winblad B, Bäckman L, Qiu C*, et al.* Accelerated progression from
mild cognitive impairment to dementia in people with diabetes. *Diabetes* 2010, **59**(11)**:** 2928-2935.

61. Selkoe DJ. Toward a comprehensive theory for Alzheimer's disease - Hypothesis: Alzheimer's
disease is caused by the cerebral accumulation and cytotoxicity of amyloid beta-protein. In:
Khachaturian ZS, Mesulam MM (eds). *Alzheimers Disease: A Compendium of Current Theories*,
vol. 924, 2000, pp 17-25.

62. Thijs V, Robberecht W, De Vos R, Sciot R. Coexistence of CADASIL and Alzheimer's disease.
Journal of Neurology Neurosurgery and Psychiatry 2003, **74**(6)**:** 790-792.

63. Brai E, Raio NA, Alberi L. Notch1 hallmarks fibrillary depositions in sporadic Alzheimer's disease.
Acta Neuropathologica Communications 2016, **4**.

64. Liu B, Wang D, Xiong T, Liu Y, Jing X, Du J*, et al.* Inhibition of Notch Signaling Promotes the
Differentiation of Epicardial Progenitor Cells into Adipocytes. *Stem Cells Int* 2021, **2021:** 8859071.

65. D'Souza B, Meloty-Kapella L, Weinmaster G. CANONICAL AND NON-CANONICAL NOTCH
LIGANDS. In: Kopan R (ed). *Notch Signaling*, vol. 92, 2010, pp 73-129.
829

830 66. Salloway S, Sperling R, Fox NC, Blennow K, Klunk W, Raskind M*, et al.* Two Phase 3 Trials of
831 Bapineuzumab in Mild-to-Moderate Alzheimer's Disease. *New England Journal of Medicine* 2014,
832 **370**(4)**:** 322-333.
833

834 67. Ostrowitzki S, Lasser RA, Dorflinger E, Scheltens P, Barkhof F, Nikolcheva T*, et al.* A phase III
835 randomized trial of gantenerumab in prodromal Alzheimer's disease. *Alzheimers Research &*
836 *Therapy* 2017, **9**.
837

838 68. Honig LS, Vellas B, Woodward M, Boada M, Bullock R, Borrie M*, et al.* Trial of Solanezumab for
839 Mild Dementia Due to Alzheimer's Disease. *New England Journal of Medicine* 2018, **378**(4)**:** 321-
840 330.
841

842 69. Henley D, Raghavan N, Sperling R, Aisen P, Raman R, Romano G. Preliminary Results of a Trial of
843 Atabecestat in Preclinical Alzheimer's Disease. *New England Journal of Medicine* 2019, **380**(15)**:**
844 1483-1485.
845

846 70. Egan MF, Kost J, Voss T, Mukai Y, Aisen PS, Cummings JL*, et al.* Randomized Trial of
847 Verubecestat for Prodromal Alzheimer's Disease. *New England Journal of Medicine* 2019, **380**(15)**:**
848 1408-1420.
849

850 71. Panza F, Lozupone M, Logroscino G, Imbimbo BP. A critical appraisal of amyloid-beta targeting
851 therapies for Alzheimer disease. *Nature Reviews Neurology* 2019, **15**(2)**:** 73-88.
852

853 72. Wessels AM, Lines C, Stern RA, Kost J, Voss T, Mozley LH*, et al.* Cognitive outcomes in trials of
854 two BACE inhibitors in Alzheimer's disease. *Alzheimer's & dementia : the journal of the*
855 *Alzheimer's Association* 2020, **16**(11)**:** 1483-1492.
856

857 73. Novak G, Streffer JR, Timmers M, Henley D, Brashear HR, Bogert J*, et al.* Long-term safety and
858 tolerability of atabecestat (JNJ-54861911), an oral BACE1 inhibitor, in early Alzheimer's disease
859 spectrum patients: a randomized, double-blind, placebo-controlled study and a two-period extension
860 study. *Alzheimers Res Ther* 2020, **12**(1)**:** 58.
861

862 74. Timmers M, Streffer JR, Russu A, Tominaga Y, Shimizu H, Shiraishi A*, et al.* Pharmacodynamics of
863 atabecestat (JNJ-54861911), an oral BACE1 inhibitor in patients with early Alzheimer's disease:
864 randomized, double-blind, placebo-controlled study. *Alzheimers Res Ther* 2018, **10**(1)**:** 85.
865

866 75. Nigam SM, Xu S, Ackermann F, Gregory JA, Lundkvist J, Lendahl U*, et al.* Endogenous APP
867 accumulates in synapses after BACE1 inhibition. *Neuroscience research* 2016, **109:** 9-15.
868

76. Citron M. Emerging Alzheimer's disease therapies: inhibition of beta-secretase. *Neurobiol Aging*
2002, **23**(6): 1017-1022.

77. Long JM, Holtzman DM. Alzheimer Disease: An Update on Pathobiology and Treatment Strategies.
Cell 2019, **179**(2): 312-339.

78. Livingston G, Sommerlad A, Orgeta V, Costafreda SG, Huntley J, Ames D*, et al.* Dementia
prevention, intervention, and care. *Lancet* 2017, **390**(10113): 2673-2734.

79. Butterfield DA, Di Domenico F, Barone E. Elevated risk of type 2 diabetes for development of
Alzheimer disease: A key role for oxidative stress in brain. *Biochimica Et Biophysica Acta-*
Molecular Basis of Disease 2014, **1842**(9): 1693-1706.

80. Arnold SE, Arvanitakis Z, Macauley-Rambach SL, Koenig AM, Wang H-Y, Ahima RS*, et al.* Brain
insulin resistance in type 2 diabetes and Alzheimer disease: concepts and conundrums. *Nature*
Reviews Neurology 2018, **14**(3): 168-181.

81. Zhao QY, Wu XH, Yan S, Xie XF, Fan YH, Zhang JQ*, et al.* The antidepressant-like effects of
pioglitazone in a chronic mild stress mouse model are associated with PPAR gamma-mediated
alteration of microglial activation phenotypes. *Journal of neuroinflammation* 2016, **13**.

82. Guo M, Li C, Lei Y, Xu S, Zhao D, Lu XY. Role of the adipose PPAR gamma-adiponectin axis in
susceptibility to stress and depression/anxiety-related behaviors. *Molecular Psychiatry* 2017, **22**(7):
1056-1068.

83. Toledo EM, Inestrosa NC. Activation of Wnt signaling by lithium and rosiglitazone reduced spatial
memory impairment and neurodegeneration in brains of an APPswe/PSEN1DeltaE9 mouse model
of Alzheimer's disease. *Mol Psychiatry* 2010, **15**(3): 272-285, 228.

84. Escribano L, Simón AM, Gimeno E, Cuadrado-Tejedor M, López de Maturana R, García-Osta A*, et*
al. Rosiglitazone rescues memory impairment in Alzheimer's transgenic mice: mechanisms
involving a reduced amyloid and tau pathology. *Neuropsychopharmacology : official publication of*
the American College of Neuropsychopharmacology 2010, **35**(7): 1593-1604.

85. O'Reilly JA, Lynch M. Rosiglitazone improves spatial memory and decreases insoluble Aβ(1-42) in
APP/PS1 mice. *Journal of neuroimmune pharmacology : the official journal of the Society on*
NeuroImmune Pharmacology 2012, **7**(1): 140-144.

86. Watson GS, Cholerton BA, Reger MA, Baker LD, Plymate SR, Asthana S*, et al.* Preserved
cognition in patients with early Alzheimer disease and amnestic mild cognitive impairment during
treatment with rosiglitazone: a preliminary study. *The American journal of geriatric psychiatry :*
official journal of the American Association for Geriatric Psychiatry 2005, **13**(11): 950-958.

87. Tzimopoulou S, Cunningham VJ, Nichols TE, Searle G, Bird NP, Mistry P*, et al.* A multi-center randomized proof-of-concept clinical trial applying [¹⁸F]FDG-PET for evaluation of metabolic therapy with rosiglitazone XR in mild to moderate Alzheimer's disease. *Journal of Alzheimer's disease : JAD* 2010, **22**(4)**:** 1241-1256.

88. Harrington C, Sawchak S, Chiang C, Davies J, Donovan C, Saunders AM*, et al.* Rosiglitazone does not improve cognition or global function when used as adjunctive therapy to AChE inhibitors in mild-to-moderate Alzheimer's disease: two phase 3 studies. *Current Alzheimer research* 2011, **8**(5)**:** 592-606.

89. Keating GM, Croom KF. Fenofibrate: a review of its use in primary dyslipidaemia, the metabolic syndrome and type 2 diabetes mellitus. *Drugs* 2007, **67**(1)**:** 121-153.

90. Barbiero JK, Santiago R, Tonin FS, Boschen S, da Silva LM, Werner MF*, et al.* PPAR-α agonist fenofibrate protects against the damaging effects of MPTP in a rat model of Parkinson's disease. *Prog Neuropsychopharmacol Biol Psychiatry* 2014, **53:** 35-44.

91. Ouk T, Gautier S, Pétrault M, Montaigne D, Maréchal X, Masse I*, et al.* Effects of the PPAR-α agonist fenofibrate on acute and short-term consequences of brain ischemia. *Journal of Cerebral Blood Flow & Metabolism* 2014, **34**(3)**:** 542-551.

92. Assaf N, El-Shamarka ME, Salem NA, Khadrawy YA, El Sayed NS. Neuroprotective effect of PPAR alpha and gamma agonists in a mouse model of amyloidogenesis through modulation of the Wnt/beta catenin pathway via targeting alpha- and beta-secretases. *Prog Neuropsychopharmacol Biol Psychiatry* 2020, **97:** 109793.

Authors' contributions

G. Gao, X Yang, B Jiang, and P Jiang were involved in the concept and design of the study. W. Qi, Y. Long, Z. Li, Z. Zhao J. Shi, D Zhu, Z Zhao, W. Xie, L. Wang, T Zhou were responsible for conducting the experiments. W. Qi, Y. Long and T Zhou drafted the manuscript and G. Gao revised the manuscript. P Jiang, Y. Long and Z. Li were responsible for data analysis. All authors contributed to the interpretation of data

and provided revisions to the manuscript. G. Gao will act as guarantor for the study.

All authors read and approved the final manuscript. W. Qi, Y. Long and Z. Li

contributed equally to this study.

Figure legend

Fig.1 Increased Kallistatin was presented in AD patients and could impair

cognitive memory in mice. (A-B) Serum Kallistatin(A), fasting blood glucose

(FBG), triglyceride (TG), and total cholesterol (TC). (B) of AD patients and their

corresponding normal control subjects. (C-D) Serum Kallistatin(C), TG, and TC(D) of

AD patients with DM and their corresponding normal control subjects. The behavioral

performance of KAL-TG mice was assessed through the Morris water maze test. (E-

H), Y-maze test(I), and electrophysiology (J). (E)The escape latency time of different

months of KAL-TG mice (3M, 6M, 9M, 12M) and corresponding WT mice were

presented during 1-6 day. (F-H) Cognitive functions were evaluated by spatial probe

test at day 7, the representative each group mice traces were shown (F), then

analyzing each group mice crossing platform times (G) and time percent in the

targeted area (H), n=4 to 9 per group. (I) Spontaneous alternation of Y-maze test. (J)

LTP was measured by whole-cell voltage-clamp recordings of CA1 neurons in acute

hippocampal slices of KAL-TG (3M, 6M, 12M) and WT mice, n = 6-12 cells from 3

mice per group. Error bars represent the standard deviation (SD); one asterisk, $p <$

0.05, two asterisks, $p < 0.01$; three asterisks, $p < 0.001$; four asterisks, $p < 0.0001$.

Fig.2 Kallistatin promoted Aβ generation. (A-B) Immunohistochemistry staining of

Aβ(A) was carried out in KAL-TG and WT mice hippocampal tissue. *Scale bar*,

100μm. The statistical analysis of Aβ plaques (B) in hippocampal tissue of KAL-TG

and WT mice, n=4-5 per group. (C) Protein levels of Aβ were tested by western blot

analysis in hippocampal tissue, n=3 per group, then statistically analyzed the above

results. (D) Hippocampal tissue Aβ42 contents were performed by ELISA in KAL-TG

and WT groups, n=3 per group. (E) Primary mouse neurons were isolated, then

infected with adenovirus to overexpress Kallistatin for 72h. Aβ42 concentration of

primary hippocampal neurons supernate and cell lysate were quantified by ELISA,

n=3 per group. (F-G) Western blot analysis of Aβ protein level in primary

hippocampal neurons infected with overexpressing Kallistatin adenovirus and control

groups, then statistical analysis of Aβ protein levels, n=3 per group. Error bars

represent the standard deviation (SD); one asterisk, $p < 0.05$; two asterisks, $p < 0.01$;

three asterisks, $p < 0.001$.

Fig.3 Kallistatin promoted tau phosphorylation. (A-B) Immunohistochemistry

staining of phosphorylated tau (p-tau S396, p-tau T231, p-tau S202) and tau(A) was

carried out in KAL-TG and WT mice hippocampal tissue. Scale bar, 100μm. The

statistical analysis of phosphorylated tau(B) in hippocampal tissue of KAL-TG and

WT mice, n=3 per group. (C-D) Protein levels of phosphorylated tau (p-tau S396, p-

tau T231, p-tau S202) and tau were tested by western blot analysis in hippocampal

tissue, then statistically analyzed the above results. Error bars represent the standard

deviation (SD); one asterisk, $p < 0.05$; two asterisks, $p < 0.01$; three asterisks, $p <$

0.001.

Fig.4 Kallistatin transgenic mice exhibited increased BACE1 expression and activity in the hippocampus. (A) Western blot analysis of relevant proteins, such as APP, PS1, and BACE1 during Aβ generation in hippocampal tissue of each time point (6M, 9M, 12M) KAL-TG mice and corresponding WT control groups, n=3 per group, then statistical analysis of APP, PS1 and BACE1 protein levels. (B) Immunohistochemistry staining of BACE1 was carried out in KAL-TG and WT mice hippocampal tissue at each time point (6M, 9M, 12M). n=3 to 5 per group. *Scale bar*, 100 μm. (C) Statistical analysis of BACE1 immunohistochemistry staining, n=3 to 4 per group. (D) ELISA measured the β-secretase activity of each group's hippocampal tissue, n=3 per group. Error bars represent the standard deviation (SD); one asterisk, *p* < 0.05, two asterisks, *p* < 0.01.

Fig.5 *In vitro*, Kallistatin promoted BACE1 expression to augment Aβ by suppressing PPARγ activation. (A) The relevant protein levels in primary mouse neurons infected with overexpressing Kallistatin adenovirus during Aβ generation were determined by western blot analysis. (B) Statistical analysis of BACE1

expression in primary neurons. (C-D) β-secretase (C) and γ- secretase (D) activity of

primary hippocampal neurons infected with overexpressing Kallistatin adenovirus and

control adenovirus was measured by ELISA. (E) Primary hippocampal neurons were

treated with BACE1 inhibitor verubecestat (50nM), then infected with adenovirus to

overexpress Kallistatin. Western blot analysis of Aβ, BACE1, and Kallistatin protein

levels, β-actin served as a loading control. (F) HT22 cells were infected with BACE1

siRNA, then infected with adenovirus to overexpress Kallistatin. Western blot

analysis of Aβ and BACE1 protein levels, β-actin served as a loading control. (G) The

relevant proteins involved in BACE1 transcriptional expressions, such as PPARγ,

YY1, and SP1 were measured by western blot analysis in hippocampal tissue. β-actin

served as a loading control. (H) Statistical analysis of PPARγ in hippocampal tissue of

each group. (I) The representative diagrams of PPARγ expression in hippocampal

tissue were presented in the above graphs. *Scale bar*, 100μm. (J) Statistical analysis of

PPARγ immunohistochemistry staining in hippocampal tissue of each group, n=3 to 4

per group. (K)Primary hippocampal neurons were treated with PPARγ agonist

rosiglitazone (10nM) for 12h, then infected with adenovirus to overexpress Kallistatin

for 72h. Western blot analysis of Aβ and BACE1 protein levels. β-actin served as a

loading control. (L) Statistical analysis of PPARγ protein levels in each group. Error

bars represent the standard deviation (SD), one asterisk, $p < 0.05$, two asterisks, $p <$

0.01; three asterisks, $p < 0.001$; ns means no significant difference.

1028

Fig.6 Kallistatin directly bonded to the Notch1 receptor, which activated the

Notch1 pathway to promote Aβ production. (A) Notch1 expression was measured

by western blot analysis in hippocampal tissue. β-actin served as a loading control.

(B) Statistical analysis of Notch1 in hippocampal tissue of each group. (C) The

representative diagrams of Nocth1 expression in hippocampal tissue were presented in

the above graphs. Scale bar, 100μm. (D) Statistical analysis of Notch1

immunohistochemistry staining in hippocampal tissue of each group. (E-F) Primary

hippocampal neurons were infected with overexpressing Kallistatin adenovirus for

72h, then Co-IP analysis (E) and membrane extraction experiment (F) was performed

to verify whether Kallistatin can bind to the Notch1 receptor. β-actin served as a

loading control. (G-H) HT22 cells were treated with siRNA (Notch1) and shRNA

(HES1) to knock down Notch1 and HES1 for 12h, then infected with adenovirus to

overexpress Kallistatin for 24h. Western blot analysis was used to detect the Notch1

signaling pathway. Error bars represent the standard deviation (SD), one asterisk, $p <$

0.05.

1044

1045 **Fig.7 Kallistatin promoted phosphorylation of tau by suppressing Wnt signaling**

1046 **pathway.** (A-B) GSK-3β and p-GSK-3β expression was measured by western blot

1047 analysis in hippocampal tissue, then statistically analyzed the above results. (C-D)

1048 Primary hippocampal neurons were treated with overexpressing Kallistatin adenovirus

1049 for 72h, then western blot analysis was used to detect the content of GSK-3β, p-GSK-

3β, tau, p-tau (Ser9, T231, S396), and statistically analyzed the above results. (E-F)

Primary hippocampal neurons were treated with overexpressing Kallistatin adenovirus

for 48h, then treated with LiCl (10mM) for 24h, western blot analysis was used to

detect the content of GSK-3β, p-GSK-3β, p-tau (Ser9, T231, S396), and statistical

analysis of the above results.

Fig.8 Fenofibrate could alleviate memory and cognitive impairment of KAL-TG mice. (A) Illustration of experimental protocols. Fenofibrate (0.3 g/kg/d × 5week, i.g.) or rosiglitazone (5mg/kg/d × 5week, i.g.) were given to KAL-TG mice. The serum for Kallistatin measuring was collected at week 0 and week 4. And Morris water maze and Y-maze test were performed at week 5. (B-E) Behavioral performance was assessed through the Morris water maze test (B-E) and the Y-maze test (F). (B) The escape latency time was presented during 1-5 day. (C-E) Cognitive functions were evaluated by spatial probe test at day 6, then analyzing each group of mice crossing platform times(C), time percent in the targeted area (D), and the path traces heatmap (E), n=5 to 6 per group. (F) Spontaneous alternation of Y-maze test. (G) Kallistatin decreased ratio was calculated by dividing the serum Kallistatin concentration of KAL-TG mice before Fenofibrate/ rosiglitazone treatment by the serum Kallistatin concentration of KAL-TG mice after a-month treatment, and serum Kallistatin concentration was measured by ELISA. (H) Protein levels of Aβ, BACE1, p-tau, tau, p-GSK-3β and GSK-3β were tested by western blot analysis in hippocampal tissue.

1071   Error bars represent the standard deviation (SD); one asterisk, $p < 0.05$.

1072

1073   Table.S1 Clinical characteristic of AD patients

	NC	AD	*p value* *(Dementia vs NC)*
N	61	56	NA
Age	49-82	52-85	****
Average age	64±8.53	73.02±9.43	NA
Male	29	25	NS
Female	32	31	
GLU (mmol/L)	5.07±0.65	6.73±2.81	****
TC (mmol/L)	4.63±0.58	4.50±1.12	NS
TG (mmol/L)	1.09±0.50	1.41±0.65	**

1074   N is an abbreviation for number; GLU is an abbreviation for glucose; TC is an

1075   abbreviation for total cholesterol; TG is an abbreviation for triglyceride; NA indicates

1076   not available; NS indicates no significance. two asterisk, $p < 0.01$, four asterisk, $p <$

1077   0.0001.

1078

1079   Table.S2 Clinical characteristic of AD patients with DM

	NC	AD + DM	*p value* *(Dementia+DM vs NC)*
N	61	26	NA
Age	49-82	52-92	****
Average age	64±8.53	76.77±8.69	NA
Male	29	15	NS
Female	32	11	
GLU (mmol/L)	5.07±0.65	8.12±3.05	****
TC (mmol/L)	4.63±0.58	4.29±1.04	NS
TG (mmol/L)	1.08±0.50	1.69±0.68	****

1080   N is an abbreviation for number; GLU is an abbreviation for glucose; TC is an

1081   abbreviation for total cholesterol; TG is an abbreviation for triglyceride; NA indicates

1082   not available; NS indicates no significance. four asterisk, $p < 0.0001$.

1083

Figure legend

Fig.S1 (A-B) GAD disease enrichment analysis (A) and PFAM analysis (B) result.

Differentially expressed genes in neurons were obtained from GSE161355, and GAD

disease enrichment was analyzed on David database. (C-D) Western blot analysis of

Kallistatin expression in aging model SAMP8 and corresponding control SAMR1

mice hippocampal tissue samples, then statistically analyzing the above results. β-

Actin served as a loading control. Error bars represent the standard deviation (SD);

three asterisks, $p < 0.001$.

1092

Fig.S2 (A) HT22 cells were infected with adenovirus to overexpress Kallistatin for

48h. Aβ42 concentration of supernate and cell lysate was quantified by ELISA. (B-C)

Western blot analysis of Aβ protein level in HT22 cells infected with overexpressing

Kallistatin adenovirus and control groups for 48h, then statistical analysis of

Kallstatin protein levels. (D) BACE1 mRNA expression in hippocampal tissue. (E) α-

secretases (ADAM9, ADAM10, ADAM17) mRNA expression in hippocampal tissue.

(F) γ-secretase activity of each group's hippocampal tissue was measured by ELISA.

(G) Primary hippocampal neurons were identified with MAP2. Error bars represent

the standard deviation (SD), one asterisk, $p < 0.05$, two asterisks, $p < 0.01$.

1102

Fig.S3 (A) The relevant protein levels in HT22 cells infected with overexpressing

Kallistatin adenovirus during Aβ generation were determined by western blot

analysis. (B) Statistical analysis of BACE1 expression in HT22 cells. (C) β-secretase

activity of HT22 cells infected with overexpressing Kallistatin adenovirus and control

adenovirus was measured by ELISA. (D) Primary hippocampal neurons were infected

with BACE1 siRNA for 72h. Western blot analysis of BACE1 protein levels. (E)

HT22 cells were treated with PPARγ agonist rosiglitazone (10nM) for 12h, then

infected with adenovirus to overexpress Kallistatin for 48h. Western blot analysis of

Aβ and BACE1 protein levels. β-actin served as a loading control. Error bars

represent the standard deviation (SD), two asterisks, $p < 0.01$.

1113

Fig.S4 (A-B) Western blot analysis of Notch1 protein level in primary hippocampal

neurons and HT22 cells infected with overexpressing Kallistatin adenovirus and

control groups. (C) HT22 cells were infected with overexpressing Kallistatin

adenovirus for 48h, and Co-IP analysis was conducted to verify whether Kallistatin

can bind to the Notch1 receptor. β-actin served as a loading control. (D) Primary

hippocampal neurons were treated with Kallistatin protein for 72h, then IP analysis.

(E) Primary hippocampal neurons were infected with Notch1 siRNA for 72h. Western

blot analysis of BACE1 protein levels. (F) HT22 cells were infected with HES1

shRNA for 48h. Western blot analysis of HES1 protein levels.

1123

1124 Fig.S5 (A) Western blot analysis of GSK-3β, p-GSK-3β, tau, p-tau(Ser9, T231, S396)

1125 in HT22 cells infected with overexpressing Kallistatin adenovirus and control groups.

1126 (B) Western blot analysis of GSK-3β, p-GSK-3β, p-tau(Ser9, T231, S396) in HT22

1127 cells infected with overexpressing Kallistatin adenovirus and control groups for 24h,

1128 then treated with LiCl(10 mM) for 24h.

1129

Fig.1

Fig.2

Fig.3

Fig.4

A

| 6M | 9M | 12M |

APP
PS1
BACE1
KAL
β-actin

WT   KAL-TG   WT   KAL-TG   WT   KAL-TG

Relative expression

WT   KAL-TG

APP   PS1   BACE1

*

B

BACE1   6M   9M   12M

WT

KAL-TG

C

BACE1 mean intensity

WT   KAL-TG

6M   9M   12M

**   *   **

D

β-secretase activity(AU)

WT   KAL-TG

6M   9M   12M

**   **   **   **   *   **

Fig.5

Fig.6

Fig.7

Fig.8

Fig.S1

A

B

C **D**

Fig.S2

Fig.S3

A

B

C

D

E

Fig.S4

Fig.S5

A

HT22

	CON	GFP	KAL
p-tau S202			
p-tau T231			
p-tau S396			
tau			
p-GSK3β			
GSK3β			
KAL			
β-actin			

B

HT22

Ad-GFP	+	–	–
Ad-KAL	–	+	+
LiCl(10mM)	–	–	+

p-GSK3β(Ser9)
GSK3β
p-tau S202
p-tau T231
p-tau S396
KAL
β-actin