Rtf1 HMD domain facilitates global histone H2B monoubiquitination

and regulates morphogenesis and virulence in the meningitis-causing

pathogen *Cryptococcus neoformans*

Yixuan Jiang[1,4], Ying Liang[1,4], Fujie Zhao[1], Zhenguo Lu[1], Siyu Wang[1],

Yao Meng[1], Zhanxiang Liu[1], Jing Zhang[1]*, Youbao Zhao[1,2,3]*

1 College of Veterinary Medicine, Henan Agricultural University, Zhengzhou, Henan

450046, China

2 Key Laboratory of Quality and Safety Control of Poultry Products, Ministry of

Agriculture and Rural Affairs, Zhengzhou, Henan 450046, China

3 Henan Province Key Laboratory of Animal Food Pathogens Surveillance, Zhengzhou,

Henan 450046, China

4 These authors contribute equally.

*Corresponding to: zhangjing@henau.edu.cn and zhaoyoubao@henau.edu.cn

14
15

Abstract

Rtf1 is generally considered to be a subunit of the Paf1 complex (Paf1C), which is a

multifunctional protein complex involved in histone modification and RNA biosynthesis at

multiple stages. Rtf1 is stably associated with the Paf1C in *Saccharomyces cerevisiae*, but not in

other species including humans. Little is known about its function in human fungal pathogens.

Here, we show that Rtf1 is required for facilitating H2B monoubiquitination (H2Bub1), and

regulates fungal morphogenesis and pathogenicity in the meningitis-causing fungal pathogen

Cryptococcus neoformans. Rtf1 is not tightly associated with the Paf1C, and its histone

modification domain (HMD) is sufficient to promote H2Bub1 and the expression of genes

related to fungal mating and filamentation. Moreover, Rtf1 HMD fully restores fungal

morphogenesis and pathogenicity; however, it fails to restore defects of thermal tolerance and

melanin production in the *rtf1Δ* strain background. The present study establishes a role for

cryptococcal Rtf1 as a Paf1C-independent regulator in regulating fungal morphogenesis and

pathogenicity, and highlights the function of HMD in facilitating global H2Bub1 in *C.*

neoformans.

Keywords: Rtf1; histone modification domain; H2B ubiquitination;

Cryptococcus neoformans; yeast-to-hypha transition; virulence

1. Introduction

In eukaryotes, gene transcription is regulated by dynamic changes in chromatin. The

posttranslational modifications of core histones, including acetylation, methylation, and

ubiquitination, represent major mechanisms by which cells alter the chromatin structural

properties and regulate gene transcription [1, 2]. Among them, the monoubiquitination of a

lysine (K) residue on the C-terminal of histone H2B (H2Bub1) is a conserved modification that

occurs on H2B K120 residue in *Homo sapiens* and K123 residue in *Saccharomyces cerevisiae*

[3, 4]. H2Bub1 is enriched at regions of active transcription but plays roles in both gene

activation and repression [5-7]. In addition, H2Bub1 is required for di- and trimethylation of H3

K4 and H3 K79, subsequently modulating chromatin accessibility [8-12].

The ubiquitin conjugase (E2) Rad6 and the ubiquitin ligase (E3) Bre1 are responsible

for H2Bub1 in *S. cerevisiae* [13-15]. In addition, H2Bub1 is regulated by additional factors in

yeast and other eukaryotes, among which the conserved polymerase-associated factor 1 (Paf1)

complex (Paf1C) is the prominent one [4, 16-19] Paf1C is a multi-functional protein complex,

which impacts RNA synthesis at multiple stages [20-28]. Paf1C consists of the subunits Paf1,

Ctr9, Cdc73, Rtf1, and Leo1, and the five subunits are stably associated within the complex in

S. cerevisiae [22, 29-31]. In contrast, Rtf1 appears not to be stably associated with Paf1C in

human cells, despite the Paf1C is structurally and functionally conserved [32-34]. Interestingly,

the histone modification domain (HMD) within Rtf1 is both necessary and sufficient for

stimulating H2Bub1 in yeast [4, 18]. Expression of the Rtf1 HMD alone restores H2Bub1 levels

in *S. cerevisiae* mutants deleted for the *RTF1* gene or all five Paf1C subunits-encoding genes [3,

4, 18]. These studies show that Rtf1 is the only Paf1C subunit that is strictly required for

deposition of H2Bub1 in vivo. However, little is known about its role in human fungal

pathogens.

Cryptococcus neoformans, the top-ranked fungus in the WHO fungal pathogen priority

list, is a globally distributed opportunistic fungal pathogen that can cause life-threatening

cryptococcosis [35, 36]. The mortality rate of cryptococcosis is alarmingly high, especially in

patients with HIV infection, in whom it ranges from 41% to 61% [37, 38]. *C. neoformans* can

be classified into two serotypes: the serotype A *C. neoformans* and the serotype D *C.*

deneoformans. Both *C. neoformans* and *C. deneoformans* undergo yeast-to-hypha transition

under inducing conditions, which has been shown to be associated with fungal virulence [39,

40]. Thus, deciphering the regulatory mechanisms on fungal morphogenesis and pathogenesis in

C. neoformans is critical for comprehensive understanding of the nature of pathogen and

combating against cryptococcal infection.

In our previous study, we characterized the subunits of complex-associated with Set1

(COMPASS) and found that COMPASS-mediated H3K4 methylation (H3K4me) affects yeast-

to-hypha transition and virulence in both *C. neoformans* and *C. deneoformans* [41]. We also

preliminarily showed that H2Bub1 is required for COMPASS-mediated H3K4me by deletion of

RAD6 and *RTF1* in *C. neoformans* and *C. deneoformans* [41]. However, the Here, we set out to

characterize the roles of Rtf1 in facilitating global H2Bub1 and to gain comprehensive insights

into the epigenetic regulation on fungal morphogenesis and pathogenesis in the human fungal

pathogen *C. neoformans*.

2. Materials and methods

2.1. Strains, culture conditions, and microscopy examination

Strains used in this study are listed in the Supplemental Table S1. *C. deneoformans* and

C. neoformans strains were maintained on YPD medium unless specified otherwise.

Transformants obtained from transient CRISPR-Cas9 coupled with electroporation (TRACE)

were selected on YPD medium with 100 μg/mL of nourseothricin, 100 μg/mL of G418, or

200 μg/mL of hygromycin.

Strains for phenotypic assays were grown overnight in liquid YPD medium at 30°C

with shaking. The cells were washed with sterile water, adjusted to an optical density at 600 nm

(OD_{600}) of 3.0, and serially diluted. For filamentation tests, aliquots (3 µL) of cell suspensions

(OD_{600} = 3.0) were spotted onto V8 plates and cultured at room temperature in the dark. For

morphological examinations, all strains were examined under a stereoscope. For spotting

assays, aliquots (3 µL) of serial dilutions starting from OD_{600} = 3.0 were spotted onto agar

medium with supplements and cultured under the noted conditions.

2.2. Gene manipulation

Cryptococcal genes were deleted following the TRACE protocol [42, 43]. In brief, a

deletion construct with approximately 1 kb of homologous arms flanking a target gene and the

dominant marker was cloned through fusion PCR. This construct was mixed with PCR products

of *CAS9* and a relevant guide RNA (gRNA), and the mixture was introduced into recipient

strains by electroporation as described previously [43]. Resulting yeast colonies were screened

by two rounds of diagnostic PCR. The first round of PCR was performed to detect the

integration of the construct into the corresponding locus of the target gene. The second round of

PCR was performed to confirm knockout of the target fragment. All primers used to make gene

deletion mutants are listed in the Supplemental Table S1.

For gene complementation, the ORFs plus approximately 1.0 kb of their upstream

regions were amplified by PCR and cloned into vectors through T5 exonuclease-dependent

assembly as previously described [44]. For gene overexpression with inducible or constitutively

active promoters, the constructs were obtained by amplifying the entire ORF by PCR and

cloning the PCR products into vectors at the downstream of *CTR4*, *TEF1*, or *GPD1* promoter.

All plasmids were confirmed by restriction enzyme digestion and sequencing. The confirmed

constructs, together with PCR products of *CAS9* and gRNA targeting the Safe Haven locus [45,

46], were introduced into recipient *Cryptococcus* strains. The transformants were passaged once

per day for five days and cultured on selection plates to obtain stable transformants. Then, two

rounds of diagnostic PCR were performed to confirm the integration and orientation of

constructs into the Safe Haven locus. All primers and plasmids used for gene complementation

and overexpression are listed in the Supplemental Table S1.

2.3. Protein extraction and western blotting

Proteins were extracted from *Cryptococcus* cells according to a previously described

method [47]. Aliquots of proteins were separated on 4%-to-12% gradient SDS-PAGE gels and

then transferred to a polyvinylidene difluoride membrane for Western blot analyses. Antibodies

used in this study are listed in the Supplemental Table S1. For co-immunoprecipitation assays

coupled with mass spectrometry (CoIP/MS), whole cell extracts of experimental strains were

incubated with FLAG-trap (Sigma) according to the manufacturer' s instructions. Proteins in the

eluted samples were loaded in SDS-PAGE gel, digested, and analysed by the proteome facility

centre of Institute of Microbiology, Chinese Academy of Sciences.

120 *2.4. RNA extraction and qPCR assays*

Cryptococcus strains were cultured in liquid YPD with shaking at 220 rpm at 30 °C

overnight, or on solid V8 medium at room temperature in the dark for 24 h. The cultures were

collected, flash frozen in liquid nitrogen, and lyophilized for 24 h. Total RNA was isolated with

the PureLink RNA Mini Kit (Invitrogen), and first strand cDNA was synthesized using the

GoScript Reverse Transcription System (Promega) following the manufacturer's instructions.

The Power SYBR Green system (Invitrogen) was used for RT-PCR. All the primers used here

are listed in the Supplemental Table S1. Relative transcript levels were determined using the

ΔΔCt method as described previously. Three biological replicates were included for all tests.

Statistical significance was determined using a Student's t-test. Differences for which $p < 0.05$

were considered statistically significant.

131 *2.5. RNA-seq and data analysis*

For RNA-seq analyses, strains were cultured in YPD liquid medium at 30 °C overnight.

The cells were washed with ddH2O and spotted on V8 medium to stimulate unisexual

reproduction. The level and integrity of RNA in each sample were evaluated using a Qubit RNA

Assay Kit on a Qubit 2.0 Fluorometer (Life Technologies, CA, USA) and RNA Nano 6000

Assay Kit with the Bioanalyzer 2100 system (Agilent Technologies, CA, USA), respectively.

RNA purity was assessed using a Nano Photometer spectrophotometer (IMPLEN, CA, USA).

The transcriptome libraries were generated using the VAHTS mRNA-seq v2 Library Prep Kit

(Vazyme Biotech Co., Ltd, Nanjing, China) according to the manufacturer's instructions.

The transcriptome libraries were sequenced by Annoroad Gene Technology Co., Ltd

(Beijing, China) on an Illumina platform. For RNA-seq analysis, the quality of sequenced clean

data was analyzed using FastQC software. Subsequently, sequences from approximately 2 GB

of clean data for each sample were mapped to the genome sequence of *C. deneoformans*

XL280α using STAT. Gene expression levels were measured in transcripts per million (TPM)

by Stringtie to determine unigenes. All unigenes were subsequently aligned against the well-

annotated genome of JEC21, which served as the parent strain to generate XL280α through a

cross with B3501α. The differential expression of genes (DEGs) was assessed using DEseq2 of

the R package and defined based on the fold change criterion (log2|fold-change | > 1.0, adjusted

p value < 0.05).

2.6. Virulence trait assays

Strains for examining virulence factors were grown overnight in liquid YPD at 30 °C with shaking. The overnight cultures were washed with sterile water, adjusted to $OD_{600} = 3.0$, and serially diluted. For thermal tolerance, melanin production, and capsule formation assay on solid plates, aliquots (3 µL) of serially diluted cell suspensions were spotted onto YPD plate, L-dopamine media, and 10% fetal bovine serum (FBS) media [48], respectively. Thermal tolerance was test at 30, 37, and 39 °C; melanin production was tested at 30 °C in the dark; capsule formation was tested at 37 °C with 5% CO_2 for 48 h. The capsule was visualized by staining with India ink and observed under a microscope. Images were captured using Zeiss Axio Imager M2 microscope. At least 50 cells were quantified and processed in ImageJ software to measure capsule and body sizes. All assays were repeated at least three times.

For melanin production assay in liquid media, the overnight cultures were washed with sterile water, adjusted to $OD_{600} = 0.3$ and transferred to L-DOPA medium (containing 0.1% L-asparagine, 0.1% dextrose, 3 mg/ml KH_2PO_4, 0.25 mg/ml thiamine, 5 ng/ml biotin 0.2 mg/ml $MgSO_4 \cdot 7 H_2O$ and 1 µg/ml L-DOPA). H99 strains grown for 48h at 30°C, each strain counted and diluted at $2x10^7$ cells/mL, while XL280 cells was induced at 30°C for 96h and diluted at $7x10^7$ cells/mL. OD_{490} was measured to determine melanin production.

2.7. Murine models of cryptococcosis

Intranasal infection model Female Balb/C mice of 8–10 weeks old were purchased from the Laboratory Animal Center of Zhengzhou University, China. Cryptococcal strains were inoculated in 3 mL of liquid YPD medium with the initial $OD_{600}=0.2$ (approximately 10^6 cell/mL) and incubated for 15 hr at 30 °C with shaking. Prior to intranasal infection, cells were washed with sterile saline three times and adjusted to the final concentration of $2×10^6$ cell/mL. Once the mice were sedated with ketamine and xylazine via intraperitoneal injection, 50 µL of the cell suspension ($1×10^5$ cells per mouse) were inoculated intranasally as previously described [41, 49-51]. Mice were monitored daily for disease progression. Animals were euthanized at 10 DPI, and lungs were dissected for fungal burden quantification.

Intravenous infection model Prior to intravenous infections, cryptococcal cells were washed with sterile saline three times and adjusted to the final concentration of $2×10^6$ cell/mL. Mice were sedated with Isoflurane. 50 µL of the cell suspension ($1×10^5$ cells per mouse) were injected intravenously as previously described [41, 49-51]. After DPI 7, animals were euthanized, and the brain, lungs, kidneys, and spleens were dissected.

For fungal burden quantifications, dissected organs were homogenized in 2 mL of cold

sterile PBS. Tissue suspensions were serially diluted in PBS and plated onto YNB agar medium

and incubated at 30°C for 2 days before counting the CFUs.

2.8. DAPI staining

DAPI (4',6-diamidino-2-phenylindole) staining assays were performed as previously

described [52]. Briefly, yeast cells or hyphae were collected and fixed with 3.7% formaldehyde

and permeabilized in 1% Triton X-100. The cells were then washed three times with PBS and

incubated in 2 µg/mL DAPI before being dropped onto a glass slide for fluorescent microscopic

observation.

2.9. Data availability

All RNA-seq data are going to be available at the NCBI (SUB14425795).

3. Results

3.1. Rtf1-mediated global H2Bub1 regulates cryptococcal yeast-to-hypha

transition

PAF1C subunit Rtf1 functions at the interface between Paf1C and Rad6/Bre1, and is

required for deposition of H2Bub1 in all the eukaryotic species examined [53]. We showed that

Rtf1 is also required for H2Bub1 and subsequent COMPASS-mediated H3K4me in the *C.*

deneoformans reference strain XL280α background (Figure 1A and B) [41]. Interestingly, loss

of H2Bub1 through deleting *RTF1* blocked unisexual yeast-to-hypha transition in *C.*

deneoformans (Supplementary Figure S1A) [41]. To establish the role of Rtf1 in regulating

cryptococcal filamentation during bisexual mating, we obtained *RTF1* deletion mutant in the *C.*

deneoformans reference strain XL280**a** background through spore dissection from cross

between *rtf1*Δα and XL280**a**, and conducted bisexual cross assay under mating-inducing

condition on V8 media. The mating hyphae during unilateral mating between *rtf1*Δα and

XL280**a** were produced at a slightly reduced level compared to mating between reference

partners XL280α and XL280**a**, while filamentation was significantly reduced during bilateral

mating between *rtf1*Δα and *rtf1*Δ**a** (Figure 1C).

During bisexual mating in *C. deneoformans*, mating pheromone (MF) is produced in

cells and secreted through the transporter Ste6 [54]. Secreted pheromone induces mating

response by binding to the compatible receptor on the cell surface of opposite mating type (Mf**a**

to Ste3α or Mfα to Ste3**a**) [55, 56]. In addition, Mat2, which is a direct downstream

transcription factor of the Cpk1 MAPK pathway, regulates the transcription of genes encoding

the above-mentioned pheromone sensing proteins [57] (Figure 1D). Given the bisexual mating

hyphae reduction caused by *RTF1* deletion, we further investigated the effects of *RTF1* deletion

on genes involved in bisexual mating at transcript level via qPCR. In comparison to the mating-

suppressing condition (YPD media), the transcript levels of *MAT2*, *MFα2*, *STE6*, and *STE3α*

were all highly induced under mating-inducing condition (V8 media). However, these

inductions were significantly impaired by deletion of *RTF1* (Figure 1D). These results strongly

indicated that Rtf1 facilitates H2Bub1 and regulates the expression of genes involved in fungal

morphogenesis in *C. deneoformans*.

222 *3.2. Ectopic expression of HMD restores global H2Bub1 levels and cryptococcal*
223 *yeast-to-hypha transition*

As the key subunit of Paf1C in mediating histone H2Bub1, Rtf1 is conserved across

eukaryotes and consists of two conserved domains, a histone modification domain HMD and a

domain that contains three highly conserved positively charged residues (Plus3) (Figure 1E). It

is worth noting that Rtf1 protein and Plus3 domain in *C. neoformans* is evolutionarily close to

higher eukaryotes, such as *H. sapiens* and *Drosophila melanogaster* (Figure 1F and

Supplementary Figure S1B and C), while the HMD domain is distant from higher eukaryotes

(Figure 1G). To further dissect the roles of Rtf1 HMD and Plus3 in facilitating histone H2Bub1

in *C. deneoformans*, we constructed the truncated versions of Rtf1 that encode HMD (53-145)

or Plus3 (227-333) with a nuclear localization sequence (NLS) added in their N terminus,

respectively, driven by the constitutive promoter and tagged with FLAG (Figure 1H).

Interestingly, overexpression of HMD domain itself significantly promoted H2Bub1 to an even

higher level in the *rtf1Δ* strain, compared to that in WT strain and the strain overexpressing the

full length of *RTF1* (Figure 1I), while overexpression of the Plus3 failed to restore H2Bub1

(Figure 1I). These results demonstrated that HMD alone is sufficient to facilitate the global

H2Bub1 level in *C. deneoformans*.

Our previous studies have demonstrated that H2Bub1 is positively related to the

filamentation in *C. neoformans* [41]. Consistently, overexpressing either the full length of *RTF1*

or the HMD domain, but not Plus3, promoted the filamentation in *rtf1Δ* strain (Figure 1J). To

gain an overview of effects on gene expression patterns by the overexpression of HMD domain,

we conducted transcriptome profiling by RNA-seq under filamentation-inducing condition (on

V8 media). The results showed that the expression levels of 668 genes were significantly

changed due to the *RTF1* deletion compared to the WT on V8 media ($|log2FC| > 1$, adjusted P

value < 0.05), with 308 genes significantly upregulated and 360 genes downregulated (Figure

2A, Supplementary Data S1). It is worth noting that the downregulated genes are mainly

enriched in GO categories related to sexual reproduction, pheromone-dependent signaling, and

filamentous growth (Supplementary Figure S2). Strikingly, overexpression of HMD domain

alone in *rtf1Δ* strain successfully restored the expression of these genes to similar levels as those

in wild-type XL280 strain, while overexpression of Plus3 domain failed to do so (Figure 2A). In

particular, the expression levels of marker genes of filamentous growth (*ZNF2* and *CFL1*) [57,

58]and genes involved in sexual reproduction and pheromone-dependent signaling (*MFα*,

STE3α, and *STE6*) as shown in Figure 1D were restored by overexpressing HMD domain alone

in the rtf1Δ background (Figure 2B, Supplementary Figure S3). These findings from

transcriptome analyses were further confirmed by qPCR (Figure 2C). In addition, the

downregulated genes in *rtf1Δ*/Plus3 cells were significantly enriched in common GO categories

as the downregulated genes in *rtf1Δ* cells (Supplementary Figure S2), relative to the wild-type

XL280 strain. Together, these results strongly suggested that HMD domain is sufficient to

facilitate global H2Bub1 to promote expression of genes associated with filamentation.

3.3. HMD is sufficient to facilitate global H2Bub1 and the consequent yeast-to-

hypha transition

The full length Rtf1 or HMD domain should properly translocate into the nucleus to

facilitate histone H2Bub1. To confirm the function of Rtf1 and HMD domain in facilitating

H2Bub1, we artificially intervened their sub-cellular localizations and evaluated the effects of

non-nuclear (cell membrane) and nuclear localizations on H2Bub1, H3K43me, and

filamentation. To achieve cell membrane localization, we fused the full length Rtf1 and HMD

with a cell membrane RGS2-mNeonGreen tag [59, 60] (Figure 3A), and introduced the

constructs into the *rtf1Δ* strain, respectively. As indicated by the mNeonGreen fluorescence,

HMD domain and the originally nuclear-localized full length Rtf1 translocated to cell

membrane after fusing with the RGS2-mNeongreen tag (Figure 3B). Both nuclear-localized

Rtf1 and HMD domain restored the levels of H2Bub1, H3K4me, and filamentation (Figure 3C

and D). In contrast, the non-nuclear-localized full length Rtf1 or HMD domain failed to restore

the levels of H2Bub1, H3K4me, or filamentation in the *rtf1Δ* strain (Figure 3C and D). These

results further supported the role of Rtf1 HMD domain in facilitating H2Bub1.

Rtf1 HMD domain is conserved from various eukaryotic species, and the residue of

glutamine at position 95 (E95, Figure 3E, F and G) has been shown to be critical for the

function of Rtf1 [28]. It is noteworthy that the residue of phenylalanine at position 118 (F118)

in *C. neoformans* is as conserved as the residue in *S. cerevisiae* (Figure 3E), which is critical for

H2Bub1 in yeast, although it is not conserved in other eukaryotic species [28]. To investigate

their roles in cryptococcal Rtf1 HMD domain, we constructed site-mutated alleles of full length

Rtf1 (Rtf1[E95A] and Rtf1[F118A]) and HMD domain (HMD[E95A] and HMD[F118A]) and introduced them

into the $rtf1\Delta$ strain, respectively. Both Rtf1^{E95A} and HMD^{E95A} failed to restored H2Bub1 and

H3K4me levels in the $rtf1\Delta$ strain, while Rtf1^{F118A} and HMD^{F118A} partially restored H2Bub1 and

H3K4me levels (Figure 3H). In consistent with the histone modification outputs, the mutants

expressing Rtf1^{E95A} or HMD^{E95A} showed non-filamentous phenotypes similar as the staring

$rtf1\Delta$ strain, while mutants expressing Rtf1^{F118A} or HMD^{F118A} produced more filaments than the

starting $rtf1\Delta$ strain (Figure 3I). Together, these results demonstrated that Rtf1 HMD domain

itself is sufficient to facilitate H2Bub1 and consequent cryptococcal filamentation with E95 as a

critical conserved residue.

3.4. Roles of the global H2Bub1 level in cryptococcal virulence factor

production

To investigate the role of HMD-mediated H2Bub1 in cryptococcal virulence, we

constructed *RTF1* deletion strain and mutants overexpressing the full length Rtf1, HMD

domain, Plus3 domain, or mutated alleles of Rtf1^{E95A} and HMD^{E95A}, respectively, in the

clinically isolated serotype A *C. neoformans* H99 strain background. Consistent with what we

observed in the serotype D *C. deneoformans*, deletion of *RTF1* abolished H2Bub1 and

H3K4me, and overexpressing the full length of Rtf1 and HMD domain alone, but not the Plus3

domain, Rtf1^{E95A} or HMD^{E95A}, successfully restored H2Bub1 and H3K4me (Figure 4A). Next,

we investigated whether Rtf1 HMD domain is involved in the production of major virulence

factors in vitro and pathogenicity in murine models of cryptococcosis. As shown in Figure 4B,

the $rtf1\Delta$ mutant had severe growth defect at 39°C, and overexpression of the full length Rtf1,

but not the Plus3 domain, Rtf1^{E95A} or HMD^{E95A}, partially restored the thermal sensitivity of the

$rtf1\Delta$ mutant (Figure 4B). Interestingly, overexpression of HMD domain alone restored the

growth defect of $rtf1\Delta$ mutant at 39°C to a level that was worse than the expression of the full

length of Rtf1 (Figure 4B). Furthermore, the $rtf1\Delta$ mutant was incapable to produce melanin,

and only the full length of Rtf1 restored its melanin production, while the HMD domain alone

failed to do so (Figure 4B). These results strongly indicate that the full length of Rtf1, but not

only the levels of global H2Bub1, is required to regulate thermal tolerance and melanin

production in *C. neoformans*.

Capsule production and cell size are known factors that are tightly associated with

cryptococcal virulence. We tested the capsule production in *RTF1*-related mutants on the FBS

solid media cultured at 37 °C with 5% CO_2. The control mutant strain $nrg1\Delta$ produced no

capsule under this condition, and the $rtf1\Delta$ mutant produced slightly less capsule compared to

the wild-type H99 strain in terms of capsule thickness (Figure 4C). Interestingly, the cell size of

the $rtf1\Delta$ mutant were significantly larger than the H99 strain under this condition (Figure

4C&D), and only overexpression of the full length of Rft1 restored the cell size enlargement

318 phenotype (Figure 4D). Consistently, the ratio between diameters of capsule layer and cell body

319 in *rtf1Δ* was significantly smaller than the ratio in the H99 strain (Figure 4E). In addition,

320 overexpression of the full length Rtf1 partially restored the ratio to the level as in the H99 strain,

321 and the HMD, Plus3, Rtf1^{E95A} or HMD^{E95A} failed to do so (Figure 4E). These results

322 showed that the global distribution of H2Bub1 across the chromosome play critical

323 roles in regulating capsule production and cell size in *C. neoformans*.

324   We further investigated the effects of Rtf1 on virulence traits in *C. deneoformans*.

325 Given the poor growth of serotype D strains under higher temperature, we only determined the

326 production of melanin and capsule in *C. deneoformans* strains. Consistent with what we

327 observed in *C. neoformans* H99 strain background, the *rtf1Δ* mutant produced less melanin in

328 L-DOPA media, and overexpression of the full length Rtf1 restored the melanin production and

329 the HMD domain failed to do so (Figure 4F). In addition, the cell size and ratio between

330 diameters of capsule layer and cell body in *rtf1Δ* were significantly smaller than the those in the

331 XL280 reference strain (Figure 4G&H). Overexpression of the full length Rtf1 partially restored

332 the ratio to the level as in the XL280 strain, while overexpression the HMD, failed to do so

333 (Figure 4H). Surprisingly, overexpression of the Plus3 domain, Rtf1^{E95A} or HMD^{E95A} partially

334 restored the melanin production and the ratio between diameters of capsule layer and cell body

335 in the *rtf1Δ* mutant in *C. deneoformans* (Figure 4F&H). These results indicated that Rtf1 protein

336 may have serotype-specific regulatory mode-of-actions in *C. neoformans* and *C. deneoformans*

337 regarding to melanin and capsule production.

3.5. Roles of the HMD-mediated H2Bub1 in regulating cryptococcal pathogenicity

340   To further investigate the role of HMD-mediated H2Bub1 in the pathogenicity of *C.*

341 *neoformans*, we tested the fungal burdens and survival rates of wild-type, *rtf1Δ*, and

342 complemented strains in intranasal and intravenous murine models of cryptococcosis (Figure

343 5A). Our results showed that both intranasally and intravenously infected lungs by the *rtf1Δ*

344 mutant had significantly reduced fungal burden compared to lungs infected by wild-type, full

345 length *RTF1*-complemented, or HMD-complemented strains (Figure 5B and C). The lungs

346 infected by Plus3-, Rtf1^{E95A}- or HMD^{E95A}-complemented strain had comparable fungal burden

347 relative to the *rtf1Δ*-infected lungs (Figure 5B and C). Similar trends in effects on fungal burden

348 were observed in other intravenously infected organs, including brain, kidney, and spleen

349 (Figure 5D, E and F). In consistent with the fungal burden analysis, the pathogenicity of *rtf1Δ*

350 mutant in the intravenous model of cryptococcosis were significantly attenuated compared to

351 the wild-type strain and strains complemented with the full length of Rtf1 or the HMD domain,

352 while the Plus3, Rtf1^{E95A}, or HMD^{E95A} failed to complement the attenuated virulence of the

353 *rtf1Δ* mutant (Figure 5G). Together, our findings suggest that HMD-mediated H2Bub1 is

354 essential for the successful survival and proliferation of *C. neoformans* during infection.

4. Discussion

In this study, we investigated the role of Rtf1 in promoting H2Bub1 and consequently

regulating cryptococcus yeast-to-hypha transition and virulence. Here, we demonstrated that the

global H2Bub1 plays pleiotropic roles in the sexual reproduction, morphogenesis, melanin

production, thermal tolerance, and pathogenicity of *C. deneoformans* and *C. neoformans*.

Interestingly, the Rtf1 HMD domain alone is sufficient to facilitate global H2Bub1 and

subsequent H3K4me. The HMD domain could fully restore the deficiencies on filamentation in

vitro and pathogenicity in a murine model of cryptococcosis. Our results fit a model in which

Rtf1 facilitates the global H2Bub1 and subsequent H3K4me levels, in order to promote

expression of genes involved in morphogenesis and pathogenicity in *C. neoformans*.

Paf1C was first identified as the RNA polymerase II transcriptional regulator

functioning in transcription elongation, and is also required for Rad6/Bre1-mediated H2Bub1

and subsequent H3K4me. Whether and how these two roles interplay with each other remain

unclear. In yeast, Paf1C contains five highly conserved core subunits Paf1, Leo1, Ctr9, Cdc73,

and Rtf1, which is stably associated with the other subunits within Paf1C. However, the human

core Paf1C was shown to interact with RNA polymerase II, in the absence of human Rtf1

homolog, indicating the dispensable role of human Rtf1 in the function of Paf1C [61]. The

Paf1C is conserved and consists of five subunits in *C. neoformans* [41]. To investigate the

association of Rtf1 with Paf1C in *C. neoformans*, we conducted co-immunoprecipitation

coupled with mass spectrometry (CoIP/MS) assays. None of the other four Paf1C subunits

could be detected with either FLAG-tagged full length of Rtf1, HMD, or Plus3 as bait

(Supplementary Data S2), strongly indicating that Rtf1 is not stably associated with the Paf1C

in *C. neoformans*.

Rtf1 is critical for H2Bub1 levels, and its deletion abolishes global H2Bub1 in both

yeast and humans. It is reasoned that Rtf1 may play dual roles in regulating elongation of gene

transcription and deposition of H2Bub1. To gain a comprehensive insight into the function of

Rtf1 in eukaryotes, we investigated the role of Rtf1 in human fungal pathogens *C.*

deneoformans and *C. neoformans* that belong to Basidiomycota. Besides its conserved functions

in facilitating global H2Bub1, we also found it is required for fungal morphogenesis and

pathogenicity. We showed that HMD domain alone is sufficient to restore cryptococcal

filamentation and virulence (Figure 1, 3 and Figure 5), concomitant with the restoration of

global H2Bub1 levels (Figure 1, 3 and 4). Given that HMD domain alone lacks regions of full

length Rtf1 required for its interactions with other Paf1C subunits and transcribed regions of

genes [4, 62], our results on the HMD domain support for a model in which the function of Rtf1
in regulating H2Bub1 is uncoupled from interaction with other subunits of Paf1C, and it is
required for cryptococcal development and virulence (Figure 5H). Biochemical and biophysical
studies on the association of Rtf1 with Paf1C subunits and Rad6/Bre1 would provide further
insights into its mode of action in regulating establishment and deposition of H2Bub1.

Rtf1 contains two conserved domains Plus3 and HMD (Figure 1E). The Plus3 domain
has been shown to interact with single-stranded DNA, indicating a role for Rtf1 in the
elongation bubble during transcription elongation [62]. In addition, Plus3 may also function in
facilitating proper positioning of H2Bub1, especially in regions of actively transcribed genes
[4]. Here, we showed that the Plus3 domain alone has no effects on global H2Bub1, while the
HMD domain alone could facilitate H2Bub1 (Figure 1A). Moreover, the *rtf1Δ* mutant showed
reduced thermal tolerance with growth defects at 37°C and 39°C, compared to the wild-type
strain (Figure 4C). The full length of Rtf1 or HMD domain alone fully restored the growth
defect of *rtf1Δ* mutant at 37°C; However, both of them only partially restored the growth defect
at 39°C (Figure 4C). In addition, only the full length of Rtf1 restored melanin production in the
rtf1Δ mutant (Figure 4D), although the HMD domain alone fully restored the global H2Bub1
levels in *C. neoformans* (Figure 4A). There are two possibilities that may lead to these
observations: (1) H2Bub1 was not properly depositioned with expression of only HMD domain,
although the global level of H2Bub1 seems normal; (2) production of the virulence factors may
require functions of Rtf1 and/or Paf1C in transcription elongation, which is absent in HMD-
complemented strain [4, 18]. These results on the HMD domain in regulating virulence factors
provide insights into the function of full-length Rtf1 and interactions with other subunits of
Paf1C. A detailed comparison of H2Bub1 occupancies across chromosomes between cells
expressing the full length of Rtf1 and HMD alone would be of great interest. In addition,
overexpression of truncated version of Rtf1 in serotype A and D background *rtf1Δ* mutants give
different output in terms of melanin and capsule production. These unexpected findings indicate
that the Rtf1 protein may have different regulatory function in serotype A and D strains, in
addition to the shared function in mediating H2Bub1. Further studies are required to uncover the
roles of Paf1C and Rtf1 in facilitating proper deposition of H2Bub1 to regulate fungal
morphogenesis and pathogenicity.

Acknowledgements

We thank the Zhao lab for their continued interest and ideas. We thank the Big Data
Center and Bioinformatics Center at Department of Veterinary Medicine, Henan Agricultural
University for providing high performance computing service. This work was supported by

Disclosure statement

No potential conflict of interest was reported by the author(s).

References

1.   Taylor, B.C. and N.L. Young, *Combinations of histone post-translational
modifications.* Biochemical Journal, 2021. **478**(3): p. 511-532.
2.   Yun, M., et al., *Readers of histone modifications.* Cell Res, 2011. **21**(4): p. 564-
78.
3.   Fetian, T., et al., *Paf1 complex subunit Rtf1 stimulates H2B ubiquitylation by
interacting with the highly conserved N-terminal helix of Rad6.* Proc Natl Acad
Sci U S A, 2023. **120**(22): p. e2220041120.
4.   Piro, A.S., et al., *Small region of Rtf1 protein can substitute for complete Paf1
complex in facilitating global histone H2B ubiquitylation in yeast.* Proc Natl
Acad Sci U S A, 2012. **109**(27): p. 10837-42.
5.   Batta, K., et al., *Genome-wide function of H2B ubiquitylation in promoter and
genic regions.* Genes Dev, 2011. **25**(21): p. 2254-65.
6.   Sun, Z.W. and C.D. Allis, *Ubiquitination of histone H2B regulates H3
methylation and gene silencing in yeast.* Nature, 2002. **418**(6893): p. 104-8.
7.   Briggs, S.D., et al., *Gene silencing: trans-histone regulatory pathway in
chromatin.* Nature, 2002. **418**(6897): p. 498.
8.   Worden, E.J. and C. Wolberger, *Activation and regulation of H2B-Ubiquitin-
dependent histone methyltransferases.* Curr Opin Struct Biol, 2019. **59**: p. 98-
106.
9.   Dover, J., et al., *Methylation of histone H3 by COMPASS requires
ubiquitination of histone H2B by Rad6.* J Biol Chem, 2002. **277**(32): p. 28368-
71.
10. Kim, T. and S. Buratowski, *Dimethylation of H3K4 by Set1 recruits the Set3
histone deacetylase complex to 5' transcribed regions.* Cell, 2009. **137**(2): p.
259-72.
11. Kim, T., et al., *Set3 HDAC mediates effects of overlapping noncoding
transcription on gene induction kinetics.* Cell, 2012. **150**(6): p. 1158-69.
12. Kim, J., et al., *RAD6-Mediated transcription-coupled H2B ubiquitylation
directly stimulates H3K4 methylation in human cells.* Cell, 2009. **137**(3): p. 459-
71.
13. Hwang, W.W., et al., *A conserved RING finger protein required for histone H2B
monoubiquitination and cell size control.* Mol Cell, 2003. **11**(1): p. 261-6.
14. Robzyk, K., J. Recht, and M.A. Osley, *Rad6-dependent ubiquitination of histone
H2B in yeast.* Science, 2000. **287**(5452): p. 501-4.
15. Wood, A., et al., *Bre1, an E3 ubiquitin ligase required for recruitment and
substrate selection of Rad6 at a promoter.* Mol Cell, 2003. **11**(1): p. 267-74.
16. Krogan, N.J., et al., *The Paf1 complex is required for histone H3 methylation by
COMPASS and Dot1p: linking transcriptional elongation to histone
methylation.* Mol Cell, 2003. **11**(3): p. 721-9.

466	17.	Ng, H.H., S. Dole, and K. Struhl, *The Rtf1 component of the Paf1 transcriptional elongation complex is required for ubiquitination of histone H2B.* J Biol Chem, 2003. **278**(36): p. 33625-8.
467		
468		
469	18.	Van Oss, S.B., et al., *The Histone Modification Domain of Paf1 Complex Subunit Rtf1 Directly Stimulates H2B Ubiquitylation through an Interaction with Rad6.* Mol Cell, 2016. **64**(4): p. 815-825.
470		
471		
472	19.	Wood, A., et al., *The Paf1 complex is essential for histone monoubiquitination by the Rad6-Bre1 complex, which signals for histone methylation by COMPASS and Dot1p.* J Biol Chem, 2003. **278**(37): p. 34739-42.
473		
474		
475	20.	Francette, A.M., S.A. Tripplehorn, and K.M. Arndt, *The Paf1 Complex: A Keystone of Nuclear Regulation Operating at the Interface of Transcription and Chromatin.* J Mol Biol, 2021. **433**(14): p. 166979.
476		
477		
478	21.	Jaehning, J.A., *The Paf1 complex: platform or player in RNA polymerase II transcription?* Biochim Biophys Acta, 2010. **1799**(5-6): p. 379-88.
479		
480	22.	Squazzo, S.L., et al., *The Paf1 complex physically and functionally associates with transcription elongation factors in vivo.* Embo j, 2002. **21**(7): p. 1764-74.
481		
482	23.	Kim, J., M. Guermah, and R.G. Roeder, *The human PAF1 complex acts in chromatin transcription elongation both independently and cooperatively with SII/TFIIS.* Cell, 2010. **140**(4): p. 491-503.
483		
484		
485	24.	Chen, Y., et al., *DSIF, the Paf1 complex, and Tat-SF1 have nonredundant, cooperative roles in RNA polymerase II elongation.* Genes Dev, 2009. **23**(23): p. 2765-77.
486		
487		
488	25.	Mueller, C.L., et al., *The Paf1 complex has functions independent of actively transcribing RNA polymerase II.* Mol Cell, 2004. **14**(4): p. 447-56.
489		
490	26.	Penheiter, K.L., et al., *A posttranscriptional role for the yeast Paf1-RNA polymerase II complex is revealed by identification of primary targets.* Mol Cell, 2005. **20**(2): p. 213-23.
491		
492		
493	27.	Sheldon, K.E., D.M. Mauger, and K.M. Arndt, *A Requirement for the Saccharomyces cerevisiae Paf1 complex in snoRNA 3' end formation.* Mol Cell, 2005. **20**(2): p. 225-36.
494		
495		
496	28.	Tomson, B.N., et al., *Identification of a role for histone H2B ubiquitylation in noncoding RNA 3'-end formation through mutational analysis of Rtf1 in Saccharomyces cerevisiae.* Genetics, 2011. **188**(2): p. 273-89.
497		
498		
499	29.	Koch, C., et al., *A role for Ctr9p and Paf1p in the regulation G1 cyclin expression in yeast.* Nucleic Acids Res, 1999. **27**(10): p. 2126-34.
500		
501	30.	Mueller, C.L. and J.A. Jaehning, *Ctr9, Rtf1, and Leo1 are components of the Paf1/RNA polymerase II complex.* Mol Cell Biol, 2002. **22**(7): p. 1971-80.
502		
503	31.	Costa, P.J. and K.M. Arndt, *Synthetic lethal interactions suggest a role for the Saccharomyces cerevisiae Rtf1 protein in transcription elongation.* Genetics, 2000. **156**(2): p. 535-47.
504		
505		
506	32.	Rozenblatt-Rosen, O., et al., *The tumor suppressor Cdc73 functionally associates with CPSF and CstF 3' mRNA processing factors.* Proc Natl Acad Sci U S A, 2009. **106**(3): p. 755-60.
507		
508		
509	33.	Chu, X., et al., *Structural insights into Paf1 complex assembly and histone binding.* Nucleic Acids Res, 2013. **41**(22): p. 10619-29.
510		
511	34.	Zhu, B., et al., *The human PAF complex coordinates transcription with events downstream of RNA synthesis.* Genes Dev, 2005. **19**(14): p. 1668-73.
512		
513	35.	Zhao, Y., et al., *Cryptococcus neoformans, a global threat to human health.* Infect Dis Poverty, 2023. **12**(1): p. 20.
514		

36. May, R.C., et al., *Cryptococcus: from environmental saprophyte to global*
pathogen. Nat Rev Microbiol, 2016. **14**(2): p. 106-17.
37. Rajasingham, R., et al., *The global burden of HIV-associated cryptococcal*
infection in adults in 2020: a modelling analysis. Lancet Infect Dis, 2022.
22(12): p. 1748-1755.
38. Iyer, K.R., et al., *Treatment strategies for cryptococcal infection: challenges,*
advances and future outlook. Nature Reviews Microbiology, 2021. **19**(7): p.
454-466.
39. Zhao, Y., et al., *Life Cycle of Cryptococcus neoformans.* Annu Rev Microbiol,
2019. **73**: p. 17-42.
40. Lin, X. and J. Heitman, *The biology of the Cryptococcus neoformans species*
complex. Annu Rev Microbiol, 2006. **60**: p. 69-105.
41. Liu, R., et al., *The COMPASS Complex Regulates Fungal Development and*
Virulence through Histone Crosstalk in the Fungal Pathogen Cryptococcus
neoformans. J Fungi (Basel), 2023. **9**(6).
42. Fan, Y. and X. Lin, *Multiple Applications of a Transient CRISPR-Cas9 Coupled*
with Electroporation (TRACE) System in the Cryptococcus neoformans Species
Complex. Genetics, 2018. **208**(4): p. 1357-1372.
43. Lin, J., Y. Fan, and X. Lin, *Transformation of Cryptococcus neoformans by*
electroporation using a transient CRISPR-Cas9 expression (TRACE) system.
Fungal Genet Biol, 2020. **138**: p. 103364.
44. Xia, Y., et al., *T5 exonuclease-dependent assembly offers a low-cost method for*
efficient cloning and site-directed mutagenesis. Nucleic Acids Res, 2019. **47**(3):
p. e15.
45. Fan, Y. and X. Lin, *An intergenic "safe haven" region in Cryptococcus*
neoformans serotype D genomes. Fungal Genet Biol, 2020. **144**: p. 103464.
46. Upadhya, R., et al., *A fluorogenic C. neoformans reporter strain with a robust*
expression of m-cherry expressed from a safe haven site in the genome. Fungal
Genet Biol, 2017. **108**: p. 13-25.
47. Zhao, Y. and X. Lin, *A PAS Protein Directs Metabolic Reprogramming during*
Cryptococcal Adaptation to Hypoxia. mBio, 2021. **12**(2).
48. Vartivarian, S.E., et al., *Regulation of cryptococcal capsular polysaccharide by*
iron. J Infect Dis, 1993. **167**(1): p. 186-90.
49. Lin, J., et al., *Immunoprotection against Cryptococcosis Offered by Znf2*
Depends on Capsule and the Hyphal Morphology. mBio, 2022. **13**(1): p.
e0278521.
50. Zhao, Y., et al., *Activation of Meiotic Genes Mediates Ploidy Reduction during*
Cryptococcal Infection. Curr Biol, 2020. **30**(8): p. 1387-1396.e5.
51. Zhai, B., et al., *Congenic strains of the filamentous form of Cryptococcus*
neoformans for studies of fungal morphogenesis and virulence. Infect Immun,
2013. **81**(7): p. 2626-37.
52. Zhao, Y., S. Upadhyay, and X. Lin, *PAS Domain Protein Pas3 Interacts with*
the Chromatin Modifier Bre1 in Regulating Cryptococcal Morphogenesis.
mBio, 2018. **9**(6).
53. Francette, A.M., S.A. Tripplehorn, and K.M. Arndt, *The Paf1 Complex: A*
Keystone of Nuclear Regulation Operating at the Interface of Transcription and
Chromatin. Journal of Molecular Biology, 2021. **433**(14).
54. Hsueh, Y.P. and W.C. Shen, *A homolog of Ste6, the a-factor transporter in*
Saccharomyces cerevisiae, is required for mating but not for monokaryotic
fruiting in Cryptococcus neoformans. Eukaryot Cell, 2005. **4**(1): p. 147-55.

55.   Wang, P. and J. Heitman, *Signal transduction cascades regulating mating,*
filamentation, and virulence in Cryptococcus neoformans. Curr Opin Microbiol,
1999. **2**(4): p. 358-62.
56.   Shen, W.C., et al., *Pheromones stimulate mating and differentiation via*
paracrine and autocrine signaling in Cryptococcus neoformans. Eukaryot Cell,
2002. **1**(3): p. 366-77.
57.   Lin, X., et al., *Transcription factors Mat2 and Znf2 operate cellular circuits*
orchestrating opposite- and same-sex mating in Cryptococcus neoformans.
PLoS Genet, 2010. **6**(5): p. e1000953.
58.   Wang, L., et al., *Fungal adhesion protein guides community behaviors and*
autoinduction in a paracrine manner. Proc Natl Acad Sci U S A, 2013. **110**(28):
p. 11571-6.
59.   Chen, S.Y., et al., *Optogenetic Control Reveals Differential Promoter*
Interpretation of Transcription Factor Nuclear Translocation Dynamics. Cell
Syst, 2020. **11**(4): p. 336-353.e24.
60.   Heximer, S.P., et al., *Mechanisms governing subcellular localization and*
function of human RGS2. J Biol Chem, 2001. **276**(17): p. 14195-203.
61.   Vos, S.M., et al., *Structure of activated transcription complex Pol II-DSIF-PAF-*
SPT6. Nature, 2018. **560**(7720): p. 607-612.
62.   de Jong, R.N., et al., *Structure and DNA binding of the human Rtf1 Plus3*
domain. Structure, 2008. **16**(1): p. 149-59.

Figure 1. Rtf1 regulates cryptococcal bisexual mating by facilitating H2Bub1 via

the HMD domain. (A) Immunoblot analysis of H2Bub1 in *C. deneoformans* wild-type

XL280, *rtf1Δ*, and *rtf1Δ*/RTF1 strains. (B) Immunoblot analysis of H3K4me (including

H3K4me1, H3K4me2, and H3K4me3) in *C. deneoformans* wild-type XL280, *rtf1Δ*, and

rtf1Δ/RTF1 strains. (C) Colony morphology of cells during bisexual mating between

indicated strains. The same volume of cells with opposite mating type at $OD_{600} = 3$ were

mixed, and 3 μL of mixtures were spotted and cultured on V8 media for 2 days at room

temperature in dark. (D) Schematic diagram of pheromone-dependent signaling

pathway and transcript levels of genes involved in pheromone signaling. The mating

cells were prepared and cultured on V8 media following the same protocol as the

colony morphology assay. After 24 h, cells were collected for total RNA extraction and

qPCR. (E) Domain structure of Rtf1 homologs in indicated eukaryotes. Cn, *C.*

deneoformans, Ca, *Candida albicans*, Af, *Aspergillus fumigatus*, Fg, *Fusarium*

graminearum, Nc, *Neurospora crassa*, Sp, *Schizosaccharomyces pombe*, Sc, *S.*

cerevisiae, Dm, *Drosophila melanogaster*, Hs, *Homo sapiens*. (F and G) Neighbor-

joining tree of Rtf1 homologs and their corresponding HMD in indicated eukaryotes.

(H) Schematic diagram of overexpressing constructs of Rtf1, HMD domain, and Plus3

domain. The constitutive promoter of *TEF1* gene was used to drive gene expression. (I)

Immunoblot analysis of H2Bub1 in strains expressing the indicated proteins. (J)

Unisexual hyphal formation of indicated strains on V8 media.

Figure 2. The expression of HMD domain alone rescues the downregulation of

genes involved in pheromone signaling and filamentous growth due to the deletion

of *RTF1*. (A) Volcano plots of differentially expressed genes in *rtf1Δ*, *rtf1Δ*/HMD, and

rtf1Δ/Plus3 relative to wild-type XL280, respectively, during unisexual mating on V8

media. Genes involved in pheromone signaling and filamentous growth were indicated.

(B) Reads coverage of indicated gene loci in XL280, *rtf1Δ*, *rtf1Δ*/HMD, and

rtf1Δ/Plus3 strains. Reads coverage at *TEF1* locus served as control. (C) qPCR

quantification of transcript levels of indicated genes in XL280, *rtf1Δ*, *rtf1Δ*/HMD, and

rtf1Δ/Plus3 strains cultured on V8. The transcript level of indicated genes were relative

to its transcript level in XL280 cells cultured in YPD.

Figure 3. The HMD domain alone is sufficient to facilitate global H2Bub1 and

restore hyphal formation in *rtf1Δ* strain. (A) Schematic diagram of constructs

expressing mNG-labelled Rtf1 or HMD domain with cell membrane tag (RGS2) or

NLS. (B) Fluorescence analysis of sub-cellular localizations of Rtf1 and HMD with

RGS2 and NLS. (C) Immunoblot analysis of H2Bub1 and H3K4me in strains

expressing the indicated proteins. (D) Colony morphology of indicated strains during

unisexual mating on V8. (E) ClustalW multiple amino acid sequence alignment of the

HMD domain in the indicated eukaryotes. The E95 and F118 residues in *C.de*

neoformans were indicated with asterisk and dot, respectively. Cn, *C. deneoformans*,

Ca, *Candida albicans*, Af, *Aspergillus fumigatus*, Fg, *Fusarium graminearum*, Nc,

Neurospora crassa, Sp, *Schizosaccharomyces pombe*, Sc, *S. cerevisiae*, Dm,

Drosophila melanogaster, Hs, *Homo sapiens*. (F and G) The 3D structure of *C.*

deneoformans HMD domain predicted by SWISS-MODEL with the 3D structure of *S.*

cerevisiae HMD domain (5emx) as the template. The conserved E95 residue was

indicated in red. (H) Immunoblot analysis of H2Bub1 and H3K4me in strains

expressing the indicated proteins. (I) Colony morphology of indicated strains during

unisexual mating on V8.

Figure 4. Rtf1 and HMD domain regulate virulence traits in both *C. neoformans*

and *C. deneoformans*. (A) Immunoblot analysis of H2Bub1 and H3K4me in strains

expressing the indicated proteins in *C. neoformans* strain background. (B) Thermal

tolerance and melanin production in *C. neoformans* strain background under indicated

conditions. (C) Capsule production in the indicated *C. neoformans* strains on capsule-

inducing media. The non-capsule-producing strain *nrg1Δ* was used as control. (D) The

indicated strains in *C. neoformans* background were cultured on capsule-inducing

media. Cell body diameter of 50 cells were quantified with ImageJ. (E) The ratio between

diameters of capsule layer and cell body of the indicated strains in *C. neoformans*

background were quantified. 50 cells for each strain were analysed. (F) Melanin

production of indicated strains in *C. deneoformans* background in liquid L-DOPA

media were quantified by determining OD_{490}. (G) Capsule production in the indicated

C. deneoformans strains on capsule-inducing media. (H) The ratio between diameters of

capsule layer and cell body of the indicated strains in *C. deneoformans* background were

quantified. 50 cells for each strain were analysed.

Figure 5. Rtf1 and HMD domain regulate cryptococcal pathogenicity in murine

models of cryptococcosis. (A) Schematic diagram of the intranasal and intravenous

infection models of cryptococcosis. The inoculum and time for detecting fungal burden

in these two infection models were indicated, respectively. (B) The fungal burden of

lungs infected by indicated strains through intranasal infection. (C-F) The fungal burden

of lungs, brain, kidney, and spleen infected by indicated strains through intravenous

infection. (G) The survival curve of animals infected by indicated strains through

intravenous infection. The inoculum was the same as the fungal burden assay for

intravenous infection. (H) Schematic diagram of the working model depicting the role

of Rtf1 and its HMD domain in regulating fungal morphogenesis in *C. deneoformans*.

Supplementary Figure S1. (A) *RTF1* gene regulates fungal filamentation in *C.*

deneoformans. (B) The neighbor-joining tree of Plus3 domain in the indicated

eukaryotes. (C) ClustalW multiple amino acid sequence alignment of Plus3 domain in

the indicated eukaryotes. Cn, *C. deneoformans*, Ca, *Candida albicans*, Af, *Aspergillus*

fumigatus, Fg, *Fusarium graminearum*, Nc, *Neurospora crassa*, Sp,

Schizosaccharomyces pombe, Sc, *S. cerevisiae*, Dm, *Drosophila melanogaster*, Hs,

Homo sapiens.

Supplementary Figure S2. Overexpression of Plus3 domain alone failed to rescue

the downregulation of genes enriched in sexual reproduction, pheromone

signaling, and filamentous growth due to *RTF1* deletion. (A) Venn diagram of

significantly enriched (*p*-value < 0.05) GO terms of downregulated genes in *rtf1Δ*,

rtf1Δ/HMD, and *rtf1Δ*/Plus3 relative to the wild-type XL280 strain on V8. (B) The 19

GO terms of downregulated genes shared by *rtf1Δ* and *rtf1Δ*/Plus3 strains.

Supplementary Figure S3. HMD domain alone successfully restores the expression

of *MAT2* and *MFα1* in *rtf1Δ* strain.

Supplementary Table S1. Strains, plasmids, primers and antibodies used in this study.

Supplementary Data S1. List of differentially expressed genes in *rtf1Δ*, *rtf1Δ*/HMD,

and *rtf1Δ*/Plus3 relative to the wild-type XL280 strain on V8.

Supplementary Data S2. Potential interacting proteins with Rtf1, HMD, and Plus3

identified by CoIP/MS.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Supplementary Figure S1

Supplementary Figure S2

A

Enriched GO terms of downregulated genes

B

19 GO terms of downregulated genes shared by *rtf1*Δ and *rtf1*Δ/Plus3

GO ID	GO name	*rtf1*Δ vs XL280		*rtf1*Δ/Plus3 vs XL280	
		Fold enrichment	*p*-value	Fold enrichment	*p*-value
GO:0019953	sexual reproduction	4.47	1.00E-05	7.13	6.20E-04
GO:0044703	multi-organism reproductive process	4.47	1.00E-05	7.13	6.20E-04
GO:0071444	cellular response to pheromone	10.98	3.92E-05	25.23	1.81E-04
GO:0032005	signal transduction involved in positive regulation of conjugation with cellular fusion	14.64	5.85E-05	42.05	3.13E-05
GO:0000750	pheromone-dependent signal transduction involved in conjugation with cellular fusion	14.64	5.85E-05	42.05	3.13E-05
GO:0019236	response to pheromone	9.99	6.92E-05	22.93	2.47E-04
GO:0040007	growth	4.31	7.80E-05	9.89	2.64E-05
GO:0030447	filamentous growth	4.6	1.05E-04	11.73	9.63E-06
GO:0000749	response to pheromone triggering conjugation with cellular fusion	12.55	1.32E-04	36.04	5.43E-05
GO:0022414	reproductive process	2.92	1.61E-04	4.47	2.09E-03
GO:0000003	reproduction	2.89	1.78E-04	4.43	2.18E-03
GO:0031136	positive regulation of conjugation	10.98	2.54E-04	31.54	8.61E-05
GO:0031139	positive regulation of conjugation with cellular fusion	10.98	2.54E-04	31.54	8.61E-05
GO:2000243	positive regulation of reproductive process	8.79	7.08E-04	25.23	1.81E-04
GO:0046999	regulation of conjugation	7.99	1.07E-03	22.93	2.47E-04
GO:0031137	regulation of conjugation with cellular fusion	7.99	1.07E-03	22.93	2.47E-04
GO:0007618	mating	4.16	1.18E-03	9.09	8.95E-04
GO:2000241	regulation of reproductive process	5.49	1.66E-03	12.61	1.58E-03
GO:0030448	hyphal growth	9.41	2.85E-03	24.03	2.81E-03

Supplementary Figure S3

Strains, plasmids, primers and antibodies used in this study.

Strain name	Genotype	Background	Sources and comments
XL280α	WT	XL280α	[51]
XC27	*rtf1::NAT*	XL280α	This study
XC111	*rtf1::NAT, P$_{CTR4}$-3xFLAG-RTF1-NEO*	XL280α	This study
YJ3	*rtf1::NAT, P$_{TEF1}$-HMD-2xFLAG-HYG*	XL280α	This study
TW3	*rtf1::NAT, P$_{TEF1}$-Plus3-2x FLAG-HYG*	XL280α	This study
YJ53	*rtf1::NAT, P$_{CTR4}$-3xFLAG-RTF1-E95A-NEO*	XL280α	This study
YL27	*rtf1::NAT, P$_{TEF1}$-HMD-E95A-2xFLAG-HYG*	XL280α	This study
YJ55	*rtf1::NAT, P$_{CTR4}$-3xFLAG-RTF1-F118A-NEO*	XL280α	This study
YL30	*rtf1::NAT, P$_{TEF1}$-HMD-F118A-2xFLAG-HYG*	XL280α	This study
YJ70	*rtf1::NAT,P$_{H3}$-RGS2membrane-mNeonGreen-RTF1-NEO*	XL280α	This study
YJ73	*rtf1::NAT,P$_{H3}$-RGS2membrane-mNeonGreen-HMD-NEO*	XL280α	This study
YJ74	*rtf1::NAT,P$_{TEF1}$-RTF1-mNeonGreen-NEO*	XL280α	This study
YJ15	*rtf1::NAT, P$_{TEF1}$-NLS-RTF1-HMD-mNeonGreen-NEO*	XL280α	This study
XL280**a**	WT	XL280**a**	[51]
XC129	*rtf1::NAT*	XL280**a**	This study
H99α	WT	H99α	Lab stock
RL226	*rtf1::NAT*	H99α	Madhani's deletion set
YJ62	*rtf1::NAT, P$_{CTR4}$-3xFLAG-RTF1-NEO*	H99α	This study
YJ64	*rtf1::NAT, P$_{TEF1}$-HMD-2xFLAG-HYG*	H99α	This study
YJ67	*rtf1::NAT, P$_{TEF1}$-Plus3-2xFLAG-HYG*	H99α	This study
YJ78	*rtf1::NAT, P$_{CTR4}$-3xFLAG-RTF1-E95A-NEO*	H99α	This study
YJ82	*rtf1::NAT, P$_{TEF1}$-HMD-E95A-2xFLAG-HYG*	H99α	This study

Plasmid	Genotype	Sources
pYZ175	P$_{CTR4}$-3×FLAG-NEO	This study
pYZ41	P$_{TEF1}$-2xFLAG-HYG	This study
pYZ25	P$_{TEF1}$-mNeonGreen-NEO	This study
pYZ281	P$_{CTR4}$-3xFLAG-RTF1-NEO	This study
pYZ255	P$_{TEF1}$-HMD-core-2xFLAG-NEO	This study
pYZ278	P$_{TEF1}$-ATG-NLS-Plus3-2xFLAG-HYG	This study
pYZ463	P$_{CTR4}$-3xFLAG-RTF1-E95A-NEO	This study
pYZ455	P$_{TEF1}$-HMD-E95A-2xFLAG-HYG	This study
pYZ467	P$_{CTR4}$-3xFLAG-RTF1-F118A-NEO	This study
pYZ459	P$_{TEF1}$-HMD-F118A-2xFLAG-HYG	This study

pYZ317	P*H3*-*RGS2membrane-mNeonGreen-NEO*	This study
pYZ506	P*H3*-*RGS2membrane-mNeonGreen-RTF1-NEO*	This study
pYZ509	P*H3*-*RGS2membrane-mNeonGreen-HMD-NEO*	This study
pYZ554	P*TEF1*-*RTF1-mNeonGreen-NEO*	This study
pYZ269	P*TEF1*-*NLS-HMD-mNeonGreen-NEO*	This study

Primer name	Sequence (5' to 3')	Description
M13F	GTAAAACGACGGCCAGT	NAT/NEO/HYG cassette and TRACE constructs from plasmid
M13R	CAGGAAACAGCTATGAC	NAT/NEO/HYG cassette and TRACE constructs from plasmid
ZhaoLab0003/YZ	TTGGATGCTGGATGCTGGGT	NAT-F
ZhaoLab0004/YZ	CCGTCTTCACCTGCATCTGATT	NAT-split-R
ZhaoLab0015/YZ	TGTGGATGCTGGCGGAGGATA	NAT-split-R for positive PCR of mutant
ZhaoLab0192/YZ	AACTGAGATACCTACAGCGTGAG	gRNA-scaffold-Far-R
ZhaoLab0193/YZ	ACTCCCTGGTCCCATCCCT	CnU6-Far-F
ZhaoLab0011/YZ	CCATCGATTTGCATTAGAACTAAAAACAAAGCA	U6 promoter NF
ZhaoLab0012/YZ	CCGCTCGAGTAAAACAAAAAAGCACCGAC	gRNA scaffold NR
ZhaoLab0054/YZ	GATAGATACTGAGGAGGACAT	PGPD1_F for Cas9
ZhaoLab0055/YZ	GGGCCCCTCTTCACGTGG	TGPD1_R for Cas9
ZhaoLab0013/YZ	AGACTCCACAGCCTAAGATCAACAGTATACCCTGCCGGTG	*SH2* gRNA, paired with 0193 (U6 promoter R)
ZhaoLab0014/YZ	GATCTTAGGCTGTGGAGTCTGTTTTAGAGCTAGAAATAGCAAGTT	*SH2* gRNA, paired with 0192 (gRNA scaffold F)
ZhaoLab0105/YZ	GTTCTCTGACCCAAAACATCGTTTTAGAGCTAGAAATAGCAAGTT	*SH3* gRNA, paired with 0193 (U6 promoter R)
ZhaoLab0106/YZ	GATGTTTTGGGTCAGAGAACAACAGTATACCCTGCCGGTG	*SH3* gRNA, paired with 0192 (gRNA scaffold F)
ZhaoLab0188/YZ	CGAAGGATGGTTGTCGCTC	screening insertion into SH3 in serotype D
ZhaoLab0189/YZ	GTATCGTCTTGCTCTTCATTCC	screening insertion into SH3 in serotype D
ZhaoLab0190/YZ	GTTGTTTCAGGCCTGCGGATG	screening insertion into SH2 in serotype A
ZhaoLab0191/YZ	GACTCATTCCTATGCCGTTC	screening insertion into SH2 in serotype A
ZhaoLab0044/YZ	TCTGACGCTGCGCCTTTG	*RTF1* deletion in XL280 LF
ZhaoLab0045/YZ	CTGGCCGTCGTTTTACATGGGATGCTGATGAGATTGCT	*RTF1* deletion in XL280 LR

ZhaoLab0046/YZ	GCTTCGGGCACCACTAAC	*RTF1* deletion in XL280 NLF
ZhaoLab0047/YZ	GTCATAGCTGTTTCCTGGGGAGAAGGATGACTACGA	*RTF1* deletion in XL280 RF
ZhaoLab0048/YZ	AAAGCGAACTGTGGACGA	*RTF1* deletion in XL280 RR
ZhaoLab0049/YZ	GACAAGAAAGAAACCGA	*RTF1* deletion in XL280 NRR
ZhaoLab0050/YZ	AGCATGGTAGGCCAAAGTACGTTTTAGAGCTAGAAATAGCAAGTT	*RTF1* deletion in XL280 gRNA, paired with 0192 (gRNA scaffold F)
ZhaoLab0051/YZ	GTACTTTGGCCTACCATGCTAACAGTATACCCTGCCGGTG	*RTF1* deletion in XL280 gRNA, paired with 0193 (U6 promoter R)
ZhaoLab0275/YZ	AGAGTGCGAGGGTTAGTAGG	*RTF1* deletion in XL280 Test F
ZhaoLab0276/YZ	GACAAGCAAAGCCCGAGT	*RTF1* deletion in XL280 Test R
ZhaoLab0306/YZ	GTGGCGGTGGGCCGGCCTCTGACCTCGAGAACGAGCTTT	*RTF1* forward with FseI for FLAG tag
ZhaoLab0307/YZ	CTGCTACTGTAACCCTTAATTCAGAAATCTCCCAAATCTAGATCCAGCTG	*RTF1* reverse with PacI for FLAG tag
ZhaoLab0177/YZ	CACAGAAAACTTCAAACCCATGGATAAAGCGGAATTAATTCCCGAGCCTCCAAAAAAGAAGAGAAAGGTCTCTGAAGATGATGAACCAGC	*HMD* forward with FseI for FLAG tag
ZhaoLab0178/YZ	TGCGATCGCGGCCGGCCCGCTCACACTAGTGTGCTTG	*HMD* reverse with SmaI for FLAG tag
ZhaoLab0310/YZ	CACAGAAAACTTCAAACCCATGGATAAAGCGGAATTAATTCCCGAGCCTCCAAAAAAGAAGAGAAAGGTCGCGCCCAACAGGGCCGA	*Plus3* reverse with FseI for FLAG tag
ZhaoLab0311/YZ	TGCGATCGCGGCCGGCCCAGCCTTTATTTCATCGTGCT	*Plus3* reverse with Sma for FLAG tag
ZhaoLab1054/YJ	AGAAAACTTCAAAGGCCGGCCCATGTCTGACCTCGAGAACG	*RTF1* forward with FseI for mNeonGreen tag
ZhaoLab1055/YJ	CCCTTGGACACCATTGCGATGAAATCTCCCAAATCTAGAT	*RTF1* reverse with AsisI for mNeonGreen tag
ZhaoLab0267/YZ	AAACTTCAAAGGCCGGATGGATAAAGCGGAATTAATTCCCGAGCCTCC	*HMD* forward with FseI for mNeonGreen tag
ZhaoLab0268/YZ	TGGACACCATTGCGATCGCGCTCACACTAGTGTGCT	*HMD* reverse with AsisI for mNeonGreen tag
ZhaoLab0315/YZ	CTCGGTACCCGGGGCGGCCGCGAGCTCGGCAGATACGATATGTTG	RGS2membrane Forward with NotI for mNeonGreen tag
ZhaoLab0316/YJ	TGGACACCATGGGCCCAGAACCACTTCCGCCCGA	RGS2membrane reverse with ApaI for mNeonGreen tag
ZhaoLab1047/YJ	CGGTGGCTCTGGGCCGGCCAGGAGCATGATCCGATACAT	*RTF1* forward with FseI for RGS2-mNeonGreen tag
ZhaoLab1048/YJ	CTGCTACTGTAACCCTTAATTAAGAAAT	*RTF1* reverse with AsisI

	CTCCCAAATCTAG	for RGS2-mNeonGreen tag
ZhaoLab1049/YJ	GCTACTGTAACCCTTAATTAAGCTCACACTAGTGTGCTTG	*HMD reverse with AsisI for mNeonGreen tag*
ZhaoLab0872/YJ	GGAAATCGAGAGAGCAAACATCTTGGCGT	Primer for RTF1^{E95A} (GAA-GCA)
ZhaoLab0873/YJ	ACGCCAAGATGTTTGCTCTCTCGATTTCC	Primer for RTF1^{E95A} (GAA-GCA)
ZhaoLab0874/YJ	AGCGCTTGATGCGATGGCCAAGACTGCTCATGGT	Primer for RTF1^{F118A} (UUC-GCC)
ZhaoLab0875/YJ	ACCATGAGCAGTCTTGGCCATCGCATCAAGCGCT	Primer for RTF1^{F118A} (UUC-GCC)
ZhaoLab0823/SW	TTGGAAAGTGCGAGGGTT	*RTF1* deletion in H99 Test F
ZhaoLab0824/SW	GATTATGTCGGAGTTGAGC	*RTF1* deletion in H99 Test R
ZhaoLab0792/RL	CTCTGGTTGGCACGGTG	*real time primer for testing JEC21 DNase effect*
ZhaoLab0793/RL	CGTCGGTCAATCTTCTCG	*real time primer for testing JEC21 DNase effect*
ZhaoLab0794/RL	CGTCACCACTGAAGTCAAGT	*TEF1 real time primer*
ZhaoLab0795/RL	AGAAGCAGCCTCCATAGG	*TEF1 real time primer*
ZhaoLab0593/YJ	AATGGTGGCACGAACGATCT	*CFL1 real time primer*
ZhaoLab0594/YJ	GTTGTCGCAATCGGGTTCAG	*CFL1 real time primer*
ZhaoLab0595/YJ	GTGATGACGACAAGGAGGCTGTT	*FAD1 real time primer*
ZhaoLab0596/YJ	GAGACGCCAGGGATGTTGATGAA	*FAD1 real time primer*
ZhaoLab0599/YJ	CAGGGTTGTAAGTTCGTTCG	*FAS1 real time primer*
ZhaoLab0600/YJ	TCGCGACTCCTCGAAAT	*FAS1 real time primer*
ZhaoLab0605/YJ	GCCATCTTACCCCTACCATCTAC	*ZNF2 real time primer*
ZhaoLab0606/YJ	TGGACATAGGAACGCTGACAAT	*ZNF2 real time primer*
ZhaoLab0601/YJ	TAGCGGAGCGGACTGGAAAGA	*STE3alpha real time primer*
ZhaoLab0602/YJ	CTCGACCGAGACGGCAATCATTA	*STE3alpha real time primer*
ZhaoLab0603/YJ	GCGAATCCACCACCGAATCAATC	*STE6 real time primer*
ZhaoLab0604/YJ	CGACGACTGCAACGCACTCT	*STE6 real time primer*
ZhaoLab0607/YJ	ATCTTCACCACCTTCACTTCT	*MFalpha2 real time primer*
ZhaoLab0608/YJ	CTAGGCGATGACACAAAGG	*MFalpha2 real time primer*
ZhaoLab1252/YJ	GCTCCTCGCTACATCTCCTCA	*MAT2 real time primer F*
ZhaoLab1253/YJ	TGTTTCGGTCTACGATACCAGTT	*MAT2 real time primer R*
ZhaoLab1254/YJ	TTGTTGGAGGATTTCAGGTTGA	*PUM1 real time primer F*
ZhaoLab1255/YJ	GTCTTCAGGAGTGGCGGTTT	*PUM1 real time primer R*

Antibody	Item number	Dilution	Brand
Ubiquityl-Histone H2B-K120 Rabbit mAb	5546T	1:1000	Cell signaling
TriMethyl-Histone H3-K4 Rabbit pAb	A2357	1:2000	ABclonal
DiMethyl-Histone H3-K4 Rabbit pAb	A2356	1:2000	ABclonal
MonoMethyl-Histone H3-K4 Rabbit pAb	A2355	1:2000	ABclonal
Histone H3 Rabbit pAb	A2348	1:5000	ImmunoWay
Flag-Tag Mouse mAb	AB0008	1:2000	Abways
Goat Anti-Mouse	RS0001	1:20000	ImmunoWay
Goat Anti-Rabbit	RS0002	1:20000	ImmunoWay