Promoters constrain evolution of expression levels of essential genes in

Escherichia coli

Saburo Tsuru[1,†,*], Naoki Hatanaka[1,†] and Chikara Furusawa[1,2,3,*]

[1]Universal Biology Institute, Graduate School of Science, The University of Tokyo, 7-3-1

Hongo, Bunkyo-ku, Tokyo 113-0033, Japan

[2]Department of Physics, Graduate School of Science, The University of Tokyo, 7-3-1 Hongo,

Bunkyo-ku, Tokyo 113-0033, Japan

[3]Center for Biosystems Dynamics Research (BDR), RIKEN, 6-2-3 Furuedai, Suita, Osaka 565-

0874, Japan

[†]These authors are equally contributed.

*Corresponding authors:

Saburo Tsuru

E-mail: saburotsuru@gmail.com

Chikara Furusawa

chikara.furusawa@riken.jp

Abstract

Variability in expression levels in response to random genomic mutations varies among genes, influencing both the facilitation and constraint of phenotypic evolution in organisms. Despite its importance, both the underlying mechanisms and evolutionary origins of this variability remain largely unknown due to the mixed contributions of *cis*- and *trans*-acting elements. To address this issue, we focused on the mutational variability of *cis*-acting elements, that is, promoter regions, in *Escherichia coli*. Random mutations were introduced into the natural and synthetic promoters to generate mutant promoter libraries. By comparing the variance in promoter activity of these mutant libraries, we found no significant difference in mutational variability in promoter activity between promoter groups, suggesting the absence of a signature of natural selection for mutational robustness. In contrast, the promoters controlling essential genes exhibited a remarkable bias in mutational variability, with mutants displaying higher activities than the wild types being relatively rare compared to those with lower activities. Our evolutionary simulation on a rugged fitness landscape provided a rationale for this vulnerability. These findings suggest that past selection created non-uniform mutational variability in promoters biased toward lower activities of random mutants now constrain future evolution of downstream essential genes toward higher expression levels.

Introduction

The rate and direction of phenotypic evolution largely depend on phenotypic variability, the tendency to vary phenotypically (Lynch and Walsh 1998; Walker 2007). It is widely recognized that organisms often exhibit nonuniform phenotypic variability, with certain variants arising more frequently than others in response to different genetic and environmental perturbations, even in the absence of natural selection(Darwin 1859; Waddington 1957; Smith, et al. 1985; Noble, et al. 2019). Therefore, it is crucial to elucidate the mechanisms underlying the bias in phenotypic variability to understand why phenotypic evolution proceeds as it does(Uller, et al. 2018). Transcriptional variability of genes across different perturbations plays a significant role in phenotypic evolution because the evolution of many traits in organisms largely depends on the evolution of expression levels and their regulation(Wittkopp, et al. 2004; Zheng, et al. 2011). Interestingly, previous studies have identified differences in transcriptional variability from gene to gene in response to various perturbations(Denver, et al. 2005; Rifkin, et al. 2005; Landry, et al. 2007; Tsuru and Furusawa 2024), where genes related to cellular growth and maintenance tend to exhibit lower transcriptional variability against random genetic perturbations to the genome in yeast(Landry, et al. 2007) and bacteria(Tsuru and Furusawa 2024). These studies, supported by theoretical evidence(Kaneko 2009; Draghi and Whitlock 2012; Furusawa and Kaneko 2018), suggest that the transcriptional variability of genes is not a random occurrence devoid of biological significance but rather a product of natural selection.

What molecular mechanisms account for gene-to-gene differences in transcriptional variability? In principle, both *cis*- and *trans*-acting elements can influence transcriptional variability(Wittkopp, et al. 2004; Tirosh, et al. 2006; Landry, et al. 2007; Payne and Wagner 2015; Tsuru and Furusawa 2024). Previous studies(Denver, et al. 2005; Rifkin, et al. 2005; Landry, et al. 2007; Tsuru and Furusawa 2024) explored the transcriptional variability against different genetic perturbations using mutation accumulation (MA) experiments(Halligan and Keightley 2009), where independent mutants were generated by accumulating genomic mutations through genetic drift, and their transcriptome profiles were obtained under identical environmental conditions. Although MA experiments provide a powerful method to quantify transcriptional variability in the absence of strong natural selection, the indiscriminate accumulation of mutations on *cis*- and *trans*-acting elements(Tsuru, et al. 2015) complicates the identification of the molecular mechanism underlying transcriptional variability. Accordingly, to overcome this difficulty, it is necessary to consider the contributions of *cis*- and *trans*-acting elements separately. Although such approaches have been extensively employed in yeast(Hornung, et al. 2012; Metzger, et al. 2015; Duveau, et al. 2021), it remains largely unknown which elements practically contribute to gene-to-gene differences in transcriptional variability associated with functions of gene products.

Why have genes related to cellular growth and maintenance acquired low mutational variability? The evolutionary origin of mutational variability is debatable (de Visser, et al. 2003).

Mutational variability could result from direct natural selection for mutational

robustness(Waddington 1942), or could be a byproduct of selection for robustness against

environmental perturbations(Meiklejohn and Hartl 2002; de Visser, et al. 2003). Several

biological traits, such as metabolic flux(Ho and Zhang 2016) and cellular morphology(Ho and

Zhang 2014), show signs of adaptive evolution through direct selection for mutational

robustness, where traits highly related to cellular growth fitness are likely to acquire lower

mutational variability through direct selection for mutational robustness. On the other hand,

other studies have identified congruence in trait variability between environmental and genetic

perturbations, such as in RNA structure(Szollosi and Derenyi 2009) and gene expression

levels(Landry, et al. 2007; Tsuru and Furusawa 2024), supporting the relevance of the byproduct

of selection for environmental robustness. As an alternative scenario, mutational variability may

spontaneously emerge with trait evolution, regardless of selection for robustness(Siegal and

Bergman 2002; de Visser, et al. 2003). To date, the most appropriate scenario for low

transcriptional variability of genes related to cellular growth and maintenance remains largely

unknown.

To address these questions, we focused on the mutational variability governed by cis-

acting elements, that is, promoter regions, of genes in *Escherichia coli*. We created a mutant

library for each promoter through random mutagenesis and analyzed the changes in promoter

activity from the wild type to mutant using flow cytometry. To explore the relevant evolutionary

scenarios for the mutational variability of promoters, we compared promoters controlling essential genes for cellular growth with those controlling nonessential genes. These groups were expected to experience different natural selection pressures on their transcriptional variability in the past. In addition, random sequences were used to construct synthetic promoters that experienced no selection for mutational robustness. Interestingly, when comparing the variance in promoter activity of the mutant libraries, these three promoter groups exhibited similar mutational variability in promoter activity, indicating the absence of signatures of natural selection for mutational robustness. In contrast, the promoters for essential genes showed a remarkable vulnerability to mutations compared to the other promoter groups, where variants with higher activity than the wild type arose less frequently than variants with lower activity. Our evolutionary simulation on a rugged fitness landscape explains the higher vulnerability of the promoters for essential genes. These results suggest that past selection created a bias in mutational variability in promoters now constrains future evolution of downstream essential genes toward higher expression levels.

Results

Higher expression levels and lower transcriptional variability of essential genes in *E. coli*

First, we analyzed the transcriptional variability of essential genes in *E. coli* in response to various genomic mutations (**Fig. 1A**). Previously(Tsuru and Furusawa 2024), we obtained transcriptome profiles of six independent mutants that had accumulated genomic mutations through genetic drift via the mutation accumulation (MA) experiment(Tsuru, et al. 2015), termed the Mut dataset. Using this dataset, we quantified the mean and standard deviation of the expression levels for each of thousands of genes (**Fig. 1B**). To compensate for the nonmonotonic mean-dependency of the standard deviations, we calculated the vertical distance from the smoothed spline of the running median of the standard deviations, termed DM_{mut}. The DM_{mut} values reflect the mean transcriptional variability in response to different genetic perturbations(Tsuru and Furusawa 2024). We categorized thousands of genes into essential and nonessential genes for cell growth, following the definition of gene essentiality by Goodall *et al*(Goodall, et al. 2018). We confirmed that essential genes showed higher expression levels than nonessential genes across the different mutants (**Fig. 1C**). We also found that essential genes had lower DM_{mut} than nonessential genes (**Fig. 1D**).

Next, we explored the transcriptional variability of essential genes in response to different environmental perturbations (**Fig. 1E,F**) using a public dataset(Sastry, et al. 2019; Rychel, et al. 2021) of the transcriptome profiles of a single strain of *E. coli*, MG1655, cultured

under 76 different environmental conditions, termed the Env dataset. Previously(Tsuru and

Furusawa 2024), we quantified the mean-controlled standard deviations, termed DM_{env}, using

a method similar to DM_{mut}. The DM_{env} reflects transcriptional variability in response to different

environmental perturbations. We confirmed that the essential genes exhibited higher expression

levels (**Fig. 1G**) and lower variability (**Fig. 1H**) than the nonessential genes across different

environmental conditions. Taken together, essential genes exhibited higher expression levels

and lower transcriptional variability in response to different perturbations, which is similar to

the characteristics of essential genes in yeast(Tirosh, et al. 2006).

135   In *E. coli*, genes with many chromosomal copies or genes located at near the origin of

replication (*oriC*) tend to have higher expression levels(Ying, et al. 2014). However, essential

genes shown in **Fig. 1** are present as only single copies on the chromosome. In addition, these

essential genes are widely spread on the chromosome and exhibited higher expression levels

than neighboring nonessential genes over the chromosome (**Fig. 1I**). Therefore, gene dosage

did not seem to explain the higher expression levels of essential genes. To explore whether the

higher expression levels of essential genes are caused by chromosomal position or by higher-

order chromosomal structures, we analyzed promoter-mediated gene expression from a plasmid

based on the experimental dataset obtained by Silander *et al*(Silander, et al. 2012). Using flow

cytometry, Silander *et al* quantified promoter activity of plasmid copies of different promoter

regions in *E. coli*, where green fluorescent protein (GFP) was transcriptionally fused to plasmid

copies of promoters. Based on fluorescence intensity from GFP, we compared promoter activity

between promoters controlling essential genes (termed essential promoters) and those

controlling nonessential genes only (termed nonessential promoters) (**supplementary fig. S1A**).

We found that essential promoters tended to have higher activities than nonessential promoters

in spite of plasmid copies. This result highlighted the importance of promoter regions in

accounting for higher expression levels of essential genes independent of chromosomal

positions or structures.

Like eukaryotes, bacteria often employ postreplicative DNA methylation for the

epigenetic control of gene expression. In *E. coli*, DNA adenine methyltransferase plays a major

role in the epigenetic control and the target GATC motifs are enriched in promoter

regions(Oshima, et al. 2002). Despite that adenine methylation at GATC sites in promoters can

inhibit transcription for some genes(Casadesus and Low 2006), we found that differences in

promoter activity between essential and nonessential promoters were almost independent of the

presence or absence of GATC motifs in the promoter regions (**supplementary fig. S1B**). These

results implied that the higher expression levels of essential genes reflected higher promoter

activities of essential promoters without known epigenetic controls.

Construction of mutant libraries of *E. coli* promoters to measure mutational effects

To explore the molecular mechanisms and signs of natural selection underlying the lower

transcriptional variability of essential genes, we focused on the promoter regions of genes with

different essentialities. In particular, we explored how promoter activity varies in response to

new mutations in the promoter regions and how this variability relates to essentiality of the

downstream genes. To this end, we used 56 natural promoters from the *E. coli* Promoter

Collection(Zaslaver, et al. 2006). Importantly, to ensure an unbiased comparison of changes in

promoter activities between different essentialities and to avoid measurement of promoter

activity below the lower detection limit in flow cytometry, we avoided the use of natural

promoters with very low activity that were slightly enriched in the promoters of nonessential

genes. To achieve this, we selected natural promoters with moderate to high activity before

mutagenesis. This selection was based on the known promoter activities of the collection as

measured by Silander et al(Silander, et al. 2012). Using the RegulonDB database(Tierrafria, et

al. 2022), we retrieved information regarding known transcriptional units regulated by these

promoters. Based on the presence of essential genes in the transcription units, we categorized

natural promoters into nonessential and essential promoters, termed Nes and Ess, respectively

(**Fig. 2A**). Nonessential promoters control transcriptional units comprising only nonessential

genes, whereas essential promoters regulate those that include at least a single essential gene.

In addition to these natural promoters, we created synthetic promoters, termed Syn, as controls

that experienced no or minimal selection for mutational robustness. To this end, we placed

random sequences (~200 bp) upstream of the *gfp* gene on a low-copy-number plasmid, pUA66-

mCherry (**Fig. 2B**, **supplementary fig. S2**). The library of mixed transformants containing this

plasmid was analyzed using flow cytometry. The top 7% of cells showing higher expression

levels of GFP (red shaded area in **Fig. 2B**) were sorted, and 16 unique clones were subsequently

isolated. Importantly, these synthetic promoters experienced selection for their expression

levels only once, and no longer experienced further mutagenesis or selection for expression

level. Accordingly, they can be regarded as a control group that is almost free from selection

for mutational robustness. Consequently, in the upstream of the GFP gene on the plasmid, we

constructed three promoter groups, Nes, Ess, and Syn, comprising different sequences with low

homology (**supplementary fig. S3, supplementary note**). The green fluorescence from GFP

was used as a measure of promoter activity(Zaslaver, et al. 2006; Silander, et al. 2012). Despite

the diversity in sequence, we confirmed that the mean activities of the three promoter groups

were comparable to each other, as designed (**Fig. 2C**). We also confirmed that the obtained

promoter activities were highly consistent with those measured by Silander et al(Silander, et al.

2012) (Spearman's R=0.94, p<0.05, **supplementary fig. S4A**). Compared to the Env dataset,

the activities of the natural promoters were positively correlated with known mRNA expression

levels of the genes downstream of the chromosomal promoters in the wild-type strain

(Spearman's R=0.64, **supplementary fig. S4B**), supporting that the wild-type promoter

activities monitored by our method considerably reflect the activities of chromosomal

promoters and their impact on the expression levels of downstream genes.

For each of these promoters, we introduced mutations by error-prone PCR at a low

frequency (6.3 mutations/kb on average, **Fig. 2D**) and integrated them into pUA66-mCherry to

create a mutant library. Each mutant library had a genetic diversity of 10^5 or more genotypes.

The promoter activity of each mutant library was measured by flow cytometry based on the

green fluorescence derived from downstream *gfp*, and was compared with the green

fluorescence of the corresponding wild type (**Fig. 2E**). The introduction of mutations was

limited to the promoter regions of the plasmid, while the chromosomal copies of the natural

promoters were intact to avoid any confounding effects. The green fluorescence intensity was

$\log_{10}$-transformed after subtracting the background (**supplementary fig. S5**), and the mean and

standard deviation of the GFP distribution were calculated (**Fig. 2E, supplementary fig. S6**).

Mutational variability in promoter activity was similar between different promoter

groups

We confirmed that all mutant libraries showed larger variations than the corresponding wild-

type clones, as designed (**Fig. 3A**), supporting that the difference in variations of GFP

distributions between wild types and the mutant libraries reflected new genetic variations

introduced by error-prone PCR rather than environmental changes or cell-to-cell heterogeneity.

In addition, we also confirmed the well-known negative correlation between mean expression

levels and standard deviations in GFP distributions regardless of the introduction of mutations

or whether the promoters were natural or synthetic (**Fig. 3B,C**), which is consistent with

previous observations(Zaslaver, et al. 2006; Wolf, et al. 2015). To compare different promoters,

we compensated for the mean-dependency of the standard deviations of both wild types and

mutant libraries by subtracting the regression line obtained from wild types (**Fig. 3B,C**). We

subsequently calculated MV_{prm}, a measure of mutational variability in promoter activity, by

subtracting the compensated standard deviation of wild types from that of mutant libraries.

Unexpectedly, MV_{prm}, showed similar values across the three promoter groups (Wilcoxon test,

adjusted p-values>0.05), even though each promoter might show unique values (**Fig. 3D**).

These results indicated that mutational variability of the focal essential promoters showed little

detectable signature of natural selection for mutational robustness despite their importance in

cellular growth, even though such selection might be active. Moreover, MV_{prm} showed no

significant positive correlations with DM_{env} and DM_{mut} of the downstream genes (**Fig. 3E,F**).

The DM values were defined in **Fig. 1**. Using the RegulonDB and EcoCyc(Keseler, et al. 2021)

databases to retrieve the known direct regulatory relationships between genes, we counted the

number of direct and unique regulations that each promoter receives from transcriptional

regulators and explored the impact of the number of regulations on mutational variability. We

found that a larger number of activatory transcriptional regulators facilitated mutational

variability in promoter activity (Spearman's R=0.31, **Fig. 3G**), probably due to a larger number

of mutational target sites. The relationship between MV_{prm} and the number of inhibitory

transcriptional regulators was unclear because promoters with null regulation dominated the

results (**Fig. 3H**). Despite that the number of activations was likely to make a difference in

MV_{prm}, the lack of significant correlations between MV_{prm} and DM values suggests that the

mutational variability of promoters provides no considerable explanation for the observed gene-

to-gene differences in transcriptional variability of the downstream genes in response to

different perturbations at the genome scale (**Fig. 1**). These results suggest that the gene-to-gene

differences in transcriptional variability in response to genome-wide perturbations may reflect

a certain *trans*-acting molecular mechanism, mainly other than the *cis*-acting one, consistent

with previous observations(Tsuru and Furusawa 2024). This implication was also rational

because promoter regions are shorter than trans-acting mutational sites for a single gene.

Vulnerability in essential promotors against mutations

Next, we explored the directionality of the mutational changes in promoter activity. Consistent

with the low mutation rate in error-prone PCR, the mean expression levels of mutant libraries

were similar to those of wild types (Spearman's R=0.99) (**Fig. 4A**). Nevertheless, we found that

log_{10}-fold changes in mean expression levels from wild types to mutant libraries followed a

global tendency over expression levels, where the promoters with higher activities in wild types

tended to decrease their activity in response to mutations (Spearman's R=–056) (**Fig. 4B**). This

tendency was retained even when the mean expression levels were calculated for the cells

within the narrow gates with the fixed size over expression levels (**supplementary figs. S6,7**),

supporting that the negative correlation was not a technical biproduct of our limitation of

detecting a decrease in expression levels. This tendency may reflect bias in the mode, activation

or inhibition, of transcriptional regulation(Kinney, et al. 2010; Belliveau, et al. 2018; Ireland,

et al. 2020). We hypothesized that promoters that receive more activation than inhibition might

tend to reduce their activity in response to mutations. To test this possibility, we defined

activation bias as the number of known activations subtracted by the number of inhibitions

(**supplementary table S1**). We found that the activation bias explained the observed mutational

changes (**Fig. 4C**). Furthermore, the activation bias also explained the $\log_{10}$-fold changes after

controlling the global trend using the regression line (**Fig. 4D**), indicating that a promoter

becomes more vulnerable as it receives more activation among promoters with similar activities.

Interestingly, synthetic and nonessential promoters showed similar mutational changes

around zero, whereas essential promoters showed biased directionality in mutational changes

toward negative values (**Fig. 4E**), even though the mean activities were almost the same among

the three promoter groups (**Fig. 2C**). There were no significant differences in promoter length

and GC content between the essential and nonessential promoters (Wilcoxon test, p>0.05). In

addition, there was no significant difference in the mutation rate in error-prone PCR between

the essential and nonessential promoters (**supplementary fig. S8A, supplementary table S2**).

The mutational spectrum in error-prone PCR was quite similar between the two promoter

groups (**supplementary fig. S8B**). These results supported the notion that the observed bias

was not derived from a technical bias. This bias in essential promoters was retained for the

normalized $\log_{10}$-fold changes (**Fig. 4F**). Accordingly, essential promoters were more

vulnerable than other groups in terms of promoter activity. In other words, this biased

directionality indicated the low frequency of the variants with higher activities than the wild-

types in essential promoters. This biased variability might limit the transcriptional variability

of essential genes in response to mutations in cis-regulatory regions, independently from

transcriptional variability in response to genomic mutations (**Fig. 1**, **3F**).

Vulnerability in essential promotors against mutations may reflect past selection for

higher activity

The difference in mutational vulnerability between essential and nonessential promoters (**Fig.**

4E, **F**) could reflect the difference in past selection pressures acting on higher expression levels

between essential and nonessential genes. Based on the observation that essential genes tended

to have higher expression levels than nonessential genes (**Fig. 1**), we hypothesized that the

remarkable vulnerability of essential promoters reflects local optima in terms of promoter

activity. That is, essential promoters might have evolved toward higher activity through past

evolution and might have been stuck in local optima because of the high essentiality of

downstream genes. In contrast, nonessential promoters might have experienced relatively

relaxed selection for higher expression levels because of the low essentiality of downstream

genes. To explore the relevance of our assumption, we compared the impact of knockdown on

growth fitness of *E. coli* between essential and nonessential genes using a public experimental

dataset reported by Hawkins *et al*(Hawkins, et al. 2020). We found that knockdown of essential

genes tended to be more deleterious for growth fitness than nonessential genes (**supplementary**

fig. S9A), supporting our assumption that essential genes are under stronger selection pressure

toward higher expression levels than nonessential genes. To investigate the emergence of

differences in vulnerability depending on selection pressure, we performed a numerical

simulation of promoter evolution on a simple rugged fitness landscape (**Fig. 5**). For simplicity,

we modeled promoter genotype as 10-bit binary numbers such as 1101011011. That is, our

model promoters are comprising of biallelic sites (0 or 1), which is simplified from real

promoters comprising of quadallelic sites (A, T, G, and C). We assumed that the expression

landscape is rugged and genotypes with higher expression levels are relatively rare, consistent

with the previous experimental observations(Wolf, et al. 2015; Yona, et al. 2018). To satisfy this

assumption, the mapping form genotype to expression level, equivalent to promoter activity,

was defined by the Rough Mt. Fuji (RMF) model, as illustrated in **Fig. 5A, D**. In short, the

genotype with the highest expression level, g^*, was first chosen at random. The expression

levels of the other genotypes were decided as a decreasing linear function of the Hamming

distance from g^*. Subsequently, stochastic noise was added to generate ruggedness. Finally,

expression levels were set to range from 0 (lower bound) to 1 (upper bound) as detailed in the

Materials and Methods section. Using this expression landscape, mutational changes were

calculated for each genotype as the mean difference in the expression levels of all nearest

neighbors with a single bit difference, or one Hamming distance, from the expression level of

the focal genotype (**Fig. 5B, E**). This simple expression landscape captured the negative

correlation between expression levels and mutational changes (**Fig. 5E**), which is consistent

with the global trend observed in our experiment (**Fig. 4B**). This trend was likely due to the

prevalence of local sinks (green dots) at lower expression levels and frequent local peaks (red

dots) at higher expression levels (**Fig. 5B, E**). To clarify this point, we calculated the fraction

of the nearest neighboring genotypes with higher expression levels for each genotype, termed

uphill ratio. Local sinks tend to have higher uphill ratios, while local peaks tend to have lower

uphill ratios. We confirmed that genotypes with higher uphill ratios dominated at lower

expression levels, while genotypes with lower uphill ratios dominated at higher expression

levels (**Fig. 5F**). Importantly, even at the same expression levels, local peaks tended to have

negative mutational changes, whereas local sinks tended to show positive changes. The fitness

landscape was then modeled by adding the basal fitness, F_b, to the expression levels multiplied

by $1 - F_b$, where F_b ranged from 0 to 1 (**Fig. 5C, G**). That is, expression level is under positive

directional selection in our model, where strength of selection pressure depends on F_b. A

smaller F_b corresponds to a steeper fitness landscape, while a larger F_b results in a flatter fitness

landscape. This simple model assumes a positive linear relationship between expression level

and fitness (**Fig. 5H**), which was relevant to the experimental evidence in most cases

(**supplementary fig. S9B**). Based on the fact that knockdown of essential genes tends to be

more deleterious for cell growth than nonessential genes (**supplementary fig. S9A**), we

assigned lower and higher values of F_b for essential and nonessential promoters, respectively.

Thus, in our model, essential and nonessential promoters shared the same expression landscape

but experienced different strengths of selection pressures. Using the constructed fitness

landscape with different F_b values (0.3, 0.5, and 0.99), we performed an evolutionary

simulation based on the standard Wright-Fisher model(Otto and Day 2007) with asexual

haploid population without recombination and with reversible mutation. Initial genotypes were

randomly generated for each simulation. To mimic ongoing evolution of real organisms, the

evolutionary simulation was performed for relatively short generations. In practice, the most

abundant mutants within the evolved populations after 30 generations were isolated and

subjected to measurement of mutational changes (**Fig. 5 G,H**). Despite the short generations,

we confirmed that the evolved isolates in steeper fitness landscapes exhibited higher expression

levels (**Fig. 5I**), which is consistent with the designed selection pressure. Comparing the

mutational changes between different F_b values, we found that the mutants that evolved with

smaller F_b showed negatively larger mutational changes, even at the same expression levels

(**Fig. 5H**). These results suggest that higher activity and vulnerability in essential promoters are

linked, and both can emerge simultaneously through stronger positive selection acting on higher

expression levels of downstream essential genes.

Despite that our simple linear model basically agreed with the known relationship

between expression level and growth fitness in most cases (**supplementary fig. S9**), an

alternative model might be suitable for some genes such as *yidC* (**supplementary fig. S9B**).

These genes exhibited almost no changes in growth fitness in response to small knockdown

level, while growth fitness gradually decreases with increases in knockdown level beyond

certain knockdown levels. To deal with such biphasic cases, we constructed the linear-plateau

model (**supplementary fig. S9**). In short, the linear-plateau model has two phases: a positive

linear phase and a flat plateau phase, where the joint point of the two phases is characterized by

E_p ranging from 0 to 1 (**supplementary fig. S10D**). A larger E_p corresponds to a higher

optimal expression level. Using the linear-plateau model with different E_p values, we

performed an evolutionary simulation by following the same procedure as the linear model

(**supplementary fig. S10E, F**). We found that the evolved isolates under the larger E_p exhibited

higher expression levels (**supplementary fig. 10G**) and negatively larger mutational changes,

even at the same expression levels (**supplementary fig. 10F**). These results seemed to be

relevant by considering neutral evolution prevalent in the fitness landscapes with the smaller

E_p. These results highlighted an impact of optimal expression level on activity and vulnerability

in promoters, suggesting that higher optimal expression levels could also contribute to higher

activity and mutational vulnerability in essential promoters. Overall, our findings imply that

both higher optimal expression and stronger directional selection could contribute to mutational

vulnerability in expression levels for essential genes. Although the expression of essential

promoters tended to decrease due to mutations (**Fig. 4E**), a few essential promoters with lower

mean expression levels exhibited neutral to positive $\log_{10}$-fold changes in expression response

to mutations. A prominent example was the essential promoter controlling *tilS*, which exhibited

the lowest expression level among wild-type essential groups (**supplementary Table S2**). This

promoter showed a positive value in $\log_{10}$-fold change in response to mutations, indicating the

frequent occurrence of mutants with higher expression levels than wild-types. These essential

promoters might suggest weaker directional selection or lower optimal expression levels for the

downstream essential genes.

Consistent results were also obtained using the NK model(Kauffman and Weinberger

1989) for rugged expression landscapes (**supplementary figs. S11 and S12**, for the linear and

linear-plateau models, respectively). The NK model, similar to the RMF model, frequently

provided sinks (genotypes with higher uphill ratios) at lower expression levels and peaks

(genotypes with lower uphill ratios) at higher expression levels (**supplementary fig. S11C**).

However, compared to the RMF model (**Fig. 5F**), sinks were relatively rare at middle to high

expression levels (beyond 0.3 in expression level). Interestingly, peaks were also rare at middle

expression levels (0.3–0.6 in expression level) in the NK model. That is, genotypes with

moderate uphill ratios were relatively frequent at middle to high expression levels. The low

frequency of sinks and peaks indicates fewer genotypes with higher or lower mutational

changes. Consequently, the difference in mutational changes between different genotypes at the

same expression levels was relatively small in the NK model. Therefore, the difference in

mutational changes between different F_b (or E_p) values in the NK model was small relative to

the RMF model (**supplementary figs. S11G,12F**). Despite these minor differences, both

expression landscapes showed consistent outcomes for the linear (**Fig. 5J**,**K** and

supplementary fig. S11G,**H** for the RMF and NK model, respectively) and linear-plateau

models (**supplementary fig. S10F**,**G** and **supplementary fig. S12F**,**G** for the RMF and NK

model, respectively). These results support the broad applicability of our key assumptions

underlying rugged expression landscapes, suggesting the robustness of the outcomes derived

from our simple models.

Discussion

Transcriptional variability differs between genes. Using public transcriptome datasets for *E. coli*, we found that essential genes exhibited lower transcriptional variability across different mutants and environmental conditions than nonessential genes. To explore the underlying mechanism, we focused on the promoter regions. We quantified the changes in promoter activity changes in response to mutations by constructing mutant libraries of natural and synthetic promoters. As a result, essential promoters did not exhibit a significant difference in mutational variability compared with nonessential and synthetic promoters. Unexpectedly, we found that essential promoters showed remarkable vulnerability to mutations, where variants with higher activities than wild-type were rare and those with lower activity were abundant. These results imply that essential promoters have evolved to reach local optima in terms of promoter activity through past selections for higher expression levels of downstream essential genes. Our numerical evolutionary simulation confirmed the relevance of this scenario. Overall, our results suggest the evolutionary impact of essentiality on the directionality of the mutational variability of promoters and imply that the directional bias in mutational variability of promoter regions might constrain further evolution of expression levels of essential genes toward higher expression levels.

There are three well-known evolutionary scenarios for the emergence of mutational robustness in many biological systems(de Visser, et al. 2003), namely adaptive(Waddington

1942), congruent(Meiklejohn and Hartl 2002), and intrinsic(Siegal and Bergman 2002)

scenarios. The adaptive scenarios predict that essential traits have evolved to reduce mutational

variability and minimize the emergence of harmful mutants through direct selection for

mutational robustness. The congruent scenarios predict that mutational robustness is a

byproduct of evolution toward low variability against environmental perturbations. Contrary to

these two scenarios assuming adaptive evolution toward low variability, the intrinsic scenarios

predict that the mutational robustness of the traits emerges spontaneously and inherently along

with the emergence of traits without selection against variability. In the case of promoter activity,

an intrinsic scenario corresponds to the mutational robustness of promoters emerging

spontaneously with the emergence of promoter activity without selection for mutational

robustness, which corresponds to the mutational variability of synthetic promoters. Our results

demonstrated that both adaptive and congruent scenarios failed to explain the observed

mutational variability in natural promoters because essential promoters showed similar

mutational variability to the other promoters, and the natural promoters exhibited no significant

positive correlation with the transcriptional variability of downstream genes against different

environmental perturbations. We note that the adaptive scenario might explain the higher

expression levels, not transcriptional variability, of essential genes (**Fig. 1C, G**), which might

minimize the risk of mutational knock down for essential genes to a lethal level. In contrast, the

intrinsic scenario explained the observed mutational variability because the natural promoters

showed similar mutational variability to that of synthetic promoters. Thus, the mutational

variability of essential promoters is unlikely to be a consequence of adaptive evolution toward

robustness against genetic and environmental perturbations.

Why did essential promoters show no less mutational variability than other promoters,

despite the fact that the reduction in expression levels of essential genes is more likely to be

harmful to cell growth than nonessential genes(Cui, et al. 2018)? One possible reason is that

promoter sequences with significant mutational robustness are rare, although several possible

mechanisms underlying mutational robustness in promoter regions are hypothesized(Payne and

Wagner 2015). In other words, the expression landscape of the promoters might be rugged

everywhere as we modeled in **Fig. 5A**. In fact, synthetic promoters derived from different

random sequences achieved higher promoter activities equivalent to natural promoters, which

was consistent with previous observations(Wolf, et al. 2015; Yona, et al. 2018). A second

possible reason is that selection for higher expression levels on a rugged fitness landscape (**Fig.**

5C) prevents the essential promoters at local optima from traversing the expression landscape

to reach genotypes with higher mutational robustness. Consistently, essential promoters showed

biased mutational effects, where most mutations in essential promoters tended to reduce

promoter activity (**Fig. 4C**). Interestingly, this scenario agreed with the predictions from a

machine-learning model based on an empirical expression landscape of yeast

promoters(Vaishnav, et al. 2022). Thus, our results suggest several plausible hypotheses that

may be shared by different organisms. Ideally, future studies should explore evolutionary

scenarios for diverse promoters that were not tested in this study, and validate the hypothetical

landscape of promoters more thoroughly. The increase of the sample set of promoters would

also compensate for our small sample sets and would enhance power to detect a potential subtle

difference in mutational variability between different promoter groups.

In this study, we focused only on mutations in promoter regions to explore the

mechanism underlying the lower transcriptional variability in essential genes. As a result, we

found that mutants with higher promoter activity were relatively rare for essential promoters.

However, this result does not exclude the contribution of any other mechanism underlying the

observed transcriptional variability in response to genome-wide perturbations (**Fig. 1**). Notably,

trans-acting mechanisms, such as transcriptional regulators, may also play a role. As it is, the

previous studies reported that genes exhibiting greater transcriptional variability across diverse

environmental and genetic perturbations tend to be regulated by a larger number of unique

transcriptional regulators in *E. coli*(Tsuru and Furusawa 2024) and yeast(Landry, et al. 2007).

A higher number of regulators provides more *trans*-mutational target sites for the regulated

genes, thereby enhancing transcriptional variability in response to genomic perturbations.

Based on this evidence, it is plausible that essential genes have less transcriptional regulation.

To support this hypothesis, we used data from RegulonDB and EcoCyc to compare the number

of regulations between essential and nonessential genes (**supplementary fig. S13**). First, we

found a smaller fraction of essential genes (90 of 248 genes, 36%) lacking known

transcriptional regulators than nonessential genes (1976 of 3995, 49%) (**supplementary fig. S13A**). This finding aligns with the positive activation bias values observed (**Fig. 4F**), probably attributable to the rich activation by sigma factors aimed at enhancing expression levels. On the other hand, among the genes regulated by at least one transcriptional regulator, we observed that essential genes received fewer transcriptional regulations than nonessential genes (**supplementary fig. S13B**), supporting this hypothesis. This observation is consistent with previous findings in yeast(Bilu and Barkai 2005). Thus, our study underscores the dual contributions of both *cis*- and *trans*-acting mechanisms including their epistatic interactions in constraining the transcriptional variability of essential genes.

Materials and Methods

Bacterial strains and medium

We used *E. coli* DH5α Electro-Cells (Takara Bio, Japan, 9027) to construct the pUA66-mCherry plasmid and its derivatives. LB medium (LB Broth, Miller, BD Difco, USA, 244620) supplemented with 25 μg/mL kanamycin (Kanamycin Sulfate, Wako, Japan, 115-00342), referred to as LB/Km medium, served as the selective medium. The SOC medium, derived from SOB medium (SOB Medium, BD Difco, USA, 244310) supplemented with 22 mM Glucose (Wako, Japan, 049-31165), was used for recovery culture pos-transformation. The *E. coli* K-12 MG1655 strain served as the host for the constructed plasmids and was employed for measuring promoter activity via flow cytometry. Bacterial cells for flow cytometry were cultured in a synthetic medium, a derivative of M9 minimal medium (pH 7.2), supplemented with 47.7 mM Na_2HPO_4 (Disodium Hydrogenphosphate 12-Water, Wako, Japan, 196-02835), 22.0 mM KH_2PO_4 (Wako, Japan, 164-22635), 8.55 mM NaCl (Wako, Japan, 191-01665), 9.35 mM NH_4Cl (Wako, Japan, 014-03005), 11 mM Glucose, 1 mM $MgSO_4$ (Magnesium Sulfate Heptahydrate , Wako, Japan, 138-00415), 50 μM $CaCl_2$ (Wako, Japan, 038-24985), and 3.8 μM thiamin (Thiamin Hydrochloride, Wako, Japan, 201-00852), with supplementation of 25 μg/mL Kanamycin when required.

Selection of natural promoters

The *E. coli* Promoter Collection(Zaslaver, et al. 2006) (Horizon Discovery Ltd., UK, PEC3877)

was used to obtain the promoter regions of the natural promoters in *E. coli*. Initially, we filtered

out the promoters with low activities that were near our detection limit based on the promoter

activities of this library measured by Silander et al(Silander, et al. 2012). This filtration

excluded 12% and 17% of essential and nonessential promoters, respectively. Subsequently,

115 promoters were randomly selected from this subset library carrying the pUA66 plasmid.

The DNA sequences of these promoter regions were verified using Sanger sequencing (Azenta

Life Sciences, USA) and compared with the reference genome sequence of *E. coli* K-12

MG1655(Blattner, et al. 1997) (GenBank ID: U00096). Subsequently, 63 of the 115 strains

without mutations in their promoter sequences were identified, and plasmids were extracted

from these colonies. Among these, 56 strains were randomly selected for the subsequent

experiments (**supplementary table S2**). Each promoter was denoted by the "NP" (natural

promoter), followed by the name of the ORF present in the immediate downstream sequence in

the genome. For example, NPmetG represents the promoter regions upstream of the *metG* gene

on the chromosome.

Plasmid construction

Plasmid pUA66 was modified by replacing the promoter regions of *gfp*, flanked by the BamHI

and XhoI restriction sites, with a red fluorescence protein (mCherry) expression cassette. This

cassette comprised dual terminators RNAI and TSAL(Reynolds, et al. 1992), promoter PLtetO-

1(Lutz 1997), Shine-Dalgarno sequence SD8(Ringquist, et al. 1992) and codon-optimized

mCherry(Dunlop, et al. 2008) coding sequences. The terminators were obtained from pNS2-

σVL(Dunlop, et al. 2008)(Addgene, Plasmid #26756) and the assembled sequences of the other

elements were commercially synthesized (Fasmac Co., Ltd., Japan). The assembled cassette

was inserted between the BamHI and XhoI recognition sequences of the pUA66 plasmid in the

direction opposite to that of *gfp* (**supplementary fig. S2**), yielding plasmid pUA66-mCherry.

PCR amplification was performed using KOD One PCR Master Mix Blue (Toyobo, Japan,

KMM-201) and pUA66-mCherry as a template with primers A

(GCTTTCCAGGGATCCTCTAGATTTAAGAAGG) and B

(GAGCAGTACTCTCGAGGTGAAGACGAAAGG) to include all sequences except for the

mCherry expression cassette. The PCR product was column purified using the FastGene

Gel/PCR Extraction Kit (NIPPON Genetics, Japan, FG-91302). The purified PCR product was

double-digested with BamHI (Takara Bio, Japan, 1010A) and XhoI (Takara Bio, Japan, 1094A)

overnight at 37 °C, followed by column purification. The digested product is referred to as the

vector.

Construction of promoter library comprising of random sequences

The construction of the synthetic promoters was based on a previous study(Wolf, et al. 2015).

Single-stranded oligonucleotides (CCTTTCGTCTTCACCTCGAG-(N200)-

GGATCCTCTAGATTTAAGAAGG) were synthesized by Eurofins Genomics (Ebersberg,

Germany). The 5' and 3' ends of the oligos contained XhoI and BamHI restriction sites,

respectively, and were homologous to primers C (CCTTTCGTCTTCACCTCGA) and D

(CCTTCTTAAATCTAGAGGATCC), respectively (**supplementary fig. S2**). A 200 bases

region of random nucleotides was present between them. PCR amplification using primers C

and D was performed using a mixture of single-stranded oligos as a template to create double-

stranded DNA. The PCR products were column purified and double digested using BamHI and

XhoI for 3 h at 37 °C, followed by column purification. Thus, an insert for the following ligation

reaction was obtained. The ligation reaction was performed using T4 DNA Ligase (Takara Bio,

Japan, 2011A) overnight at 16 °C involved mixing the vector and insert at a molar ratio of 1: 3.

The resulting ligation product solution was column purified and DNA was extracted using 20

µL of sterile water. All purified ligation product solutions were added to 50 µL of DH5α

Electro-Cells and electroporated at 1800 V (Eporator, Eppendroph, Germany, 4309000027).

Transformed cells were incubated at 37 °C for 1 h in 1 mL of SOC medium, followed by

overnight incubation at 37 °C in 1 mL of LB medium supplemented with 2 µL of 25 mg/mL

kanamycin solution.

Selection of synthetic promoters using flow cytometry

Green and red fluorescence, as well as forward scatter of the cells in the DH5α-transformed

random promoter library were measured using a flow cytometer (FACSAria III, BD, USA).

Gates were set for the green and red channels to exclude cells with low GFP and high mCherry

fluorescence, respectively. Subsequently, 10,000 cells without red fluorescence but with green

fluorescence within the selected range (**Fig. 2B**) were collected by cell sorting, according to the

manufacturer's instructions. The sorted cells were spread on LB/Km agar plates and incubated

overnight at 37 °C. Bacterial colonies were randomly picked and their inserted synthetic

promoter regions were sequenced by Sanger sequencing. Consequently, 16 synthetic promoters

without improper mutations at the restriction sites were identified and named SP1–16,

(**supplementary table S1**).

Random mutagenesis using error-prone PCR

Error-prone PCR was conducted using 0.1 ng of each promoter region as a template, with 35

cycles of amplification performed using the GeneMorph II Random Mutagenesis Kit (Agilent

Technologies, USA, 200550) and primers C and D. The resulting PCR products underwent gel

purification and were double digested with BamHI and XhoI for 3 hours at 37 °C, followed by

column purification to create mutant inserts. Ligation was performed overnight at 16 °C using

T4 DNA Ligase, with the vector and mutant inserts mixed at a molar ratio of 1:3. The ligation

reaction solution was column purified, and DNA was extracted using 20 µL of sterile water.

Construction of mutant promoter library

The purified ligation product solution was added to 50 µL of DH5α Electro-Cells and

electroporated at 1800 V. Transformed cells were cultured at 37 °C for 1 h in 1 mL of SOC

medium and a portion of the culture was plated on LB/Km agar medium for overnight growth

at 37 °C. Bacterial colonies were counted to determine the transformation efficiency. If the

transformation efficiency exceeded 10^5 cells/reaction, the next step was performed; otherwise,

the process was repeated. The transformed culture was diluted with 5 mL of LB/Km medium

and grown overnight. Plasmid extraction from an overnight culture was performed using the

FastGene Plasmid Mini Kit (NIPPON Genetics, Japan, FG-90502). The obtained plasmid

solution was added to 50 µL of MG1655 electrocompetent cells and electroporated at 1800 V.

Transformed cells were cultured at 37 °C for 1 h in 1 mL of SOC medium, and a portion of the

solution was plated on LB/Km agar medium for overnight growth at 37 °C to count the number

of colonies and determine transformation efficiency. If the transformation efficiency exceeded

10^5 cells/reaction, the next step was performed; otherwise, the process was repeated. A sterile

2 mL 96 well deep well plate (Greiner Bio-One, Austria, 780271) was used to add 1 mL of the

synthetic medium to each well, followed by the addition of 10 µL of the overnight culture

containing the MG1655 transformants. The bacterial cells were grown at 37 °C with shaking at

800 rpm until confluent growth was achieved. Subsequently, another plate was prepared with 1

mL of fresh synthetic medium added to each well, and 10 µL of the confluently grown cultures

was diluted into each well. The cultures were grown for 2.5 h at 37 °C with shaking at 800 rpm

and were used as mutant libraries for flow cytometry.

Verification of PCR random mutagenesis

The number of mutations introduced into the promoters by error-prone PCR was determined

using Sanger sequencing. Three to four colonies were randomly selected from six randomly

selected mutant libraries, including essential and nonessential promoters (**supplementary table**

S2). Colony PCR was performed using KOD One PCR Master Mix Blue with primers E

(TTACTTTGCAGGGCTTCCCAA) and F (CCAGTCTTTCGACTGAGCCT) to amplify the

regions including the promoter regions (**supplementary fig. S2**). PCR products were column

purified and verified by Sanger sequencing. The average mutation rate was calculated to be 6.2

mutations/kb. Mutation rates and mutational spectrum for each promoter group were shown in

supplementary fig. S8.

Construction of wild-type promoter control group

To ensure accurate measurement of GFP fluorescence from cells with wild-type promoters,

wild-type plasmids were transformed into MG1655 cells by electroporation. Transformed

cultures were plated on LB/Km agar medium and incubated overnight, and the resulting

colonies were cultured in LB/Km medium. Cell cultures of these control groups in the synthetic

medium were prepared and used as wild types for flow cytometry.

Flow cytometry

Bacterial cells were dispensed in M9 buffer prior to analysis using a flow cytometer (BD, USA,

FACSAria III). Fluorescence intensity from GFP (GFP FI) and mCherry (mCherry FI) was

measured using a 488 nm laser with 515–545-nm emission filter, and a 561 nm laser with 563–

588-nm emission filter, respectively. The following PMT voltage settings were used: forward

scatter (FSC), 300; side scatter (SSC), 340; green fluorescence, 700; red fluorescence, 600. The

events exceeding the defined thresholds of FSC (200) and SSC (200) were recorded.

Total,000,000 events were measured per library. The MG1655 strain was used as a control for

autofluorescence of GFP FI.

Data analysis of flow cytometry

FCS files were converted to CSV format using the flowCore(Ellis, et al. 2024) package in R(R

Core Team 2023). Subsequent analyses were conducted using custom R scripts. To compare

green fluorescence among cells in similar physiological states, a narrow gate for the forward

scatter channel was set to capture 40% of the events, including the most frequent values for

MG1655 (**supplementary fig. S5A**). The top 0.5% value of the green fluorescence intensity

distribution of MG1655 within the gate was determined as the autofluorescence intensity

(**supplementary fig. S5B**). For the events of wild types and mutant libraries, the

autofluorescence intensity was subtracted from the green fluorescence intensity of the events

within the gate. Events with a negative green fluorescence intensity after subtracting the

autofluorescence intensity were excluded from the analysis. In addition, events with a red

fluorescence intensity greater than 10^3 [a.u.] were regarded as the cells carrying pUA66-

mCherry and were excluded from the analysis. Clonal cell populations exhibit huge cell-to-cell

heterogeneity in gene expression even under identical environmental conditions, resulting in

long-tailed or log-normal shaped distributions in protein numbers(Tsuru, et al. 2009; Taniguchi,

et al. 2010). This stochastic nature often results in rare cells exhibiting very highly expression

levels within cell populations, which greatly influences variance or standard deviation of protein number distributions. To ensure robust measurement of variation in fluorescent distributions, the green fluorescence intensity after subtracting the autofluorescence intensity was $\log_{10}$-transformed and used to calculate the means and standard deviations. We note that this log-transformation might limit detection of possible small differences in mutational variability in promoter activities between different promoter groups.

Classification of promoter based on essentiality of transcribed unit

Gene essentiality was based on a previous study(Goodall, et al. 2018), which identified 248 genes designated as essential across different datasets. In bacteria, clusters of genes are controlled by the same promoter sequences and are transcribed as single mRNAs, termed transcription units. Promoters controlling at least one essential gene in the transcription units were defined as essential promoters (**Fig. 2A**). The known transcription units were retrieved from the RegulonDB(Tierrafria, et al. 2022) database. Of the 56 promoters, 18 were classified as essential and 38 as nonessential as listed in **supplementary table S1**.

Estimation of transcriptional variability

Transcriptional variability of each gene in *E. coli* in response to different genetic perturbations was estimated using known transcriptome profiles obtained from six independent mutants derive from MG1655 cultured under a single environmental condition, termed the Mut dataset(Tsuru and Furusawa 2024). These mutants were subjected to mutation accumulation

(MA) experiments to accumulate genomic mutations via genetic drift(Halligan and Keightley

2009; Tsuru, et al. 2015). In addition, to quantify the transcriptional variability in response to

different environmental conditions, known transcriptome profiles of MG1655 cultured under

76 different environmental conditions(Sastry, et al. 2019; Rychel, et al. 2021; Lamoureux, et al.

2022), termed the Env dataset(Tsuru and Furusawa 2024), were utilized. The expression levels

were determined by $\log_2$-transformation and quantile normalization. The means and standard

deviations in expression levels for each gene across different genetic perturbations and

environmental conditions were quantified. To compensate for the mean-dependency of the

standard deviations (**Fig. 1B,G**), the DM values for each gene were calculated as the vertical

distance of each standard deviation from a smoothed running median of the standard deviations

for each dataset, as detailed previously(Tsuru and Furusawa 2024). The resulting DM values

were termed DM_{mut} and DM_{env} for the Mut and Env datasets, respectively. The DM values were

used as measures of transcriptional variability in response to genetic and environmental

perturbations.

Phylogenetic analysis of promoter sequence of length 73 bases

Synthetic promoters underwent a blastn(Altschul, et al. 1990) search to confirm that there was

no considerable similarity to any natural sequences (**supplementary note**). Posterior sequences

of 73 bases were aligned using the msa(Bodenhofer, et al. 2015) package in R (Clustal Omega),

where 73 bases represented the shortest length of a synthetic promoter. A phylogenetic tree was

constructed using the Neighbor-Joining method.

Phylogenetic analysis of core elements of sigma70 promoters

679   We identified sigma70 promoters from our natural promoters using RegulonDB. The core

680   elements comprising the –10/–35 elements of these promoters were concatenated and subjected

681   to phylogenetic analysis. The Levenshtein distance(Levenshtein 1966) between the focal

682   elements and the consensus elements, which consist of TTGACA and TATAAT, was calculated.

683   A phylogenetic tree was constructed as previously described.

Identification of the number of transcriptional regulators regulating the natural

promoters

686   Reported regulatory interactions between genes were compiled using all interactions with

687   experimental evidence from RegulonDB and EcoCyc(Keseler, et al. 2021) for both transcription

688   factors and sigma factors. The mode of regulation was classified as either activation or

689   inhibition, and the mode of regulation by sigma factors was classified as activation.

Numerical simulation of promoter evolution

691   The promoter genotype, g, was modeled by a bit string comprising either zeros or ones of length

692   N=10 (e.g., 1110001100). The expression level, E, was defined by the Rough Mt. Fuji (RMF)

693   model(Aita, et al. 2000; Neidhart, et al. 2014) as:

$$E(g) = -cD(g, g^*) + \eta(g),$$

where c is a scaling coefficient equal to one in this study, g^* is the genotype with the highest

expression level before adding noise, $D(g, g^*)$ is the Hamming distance between g and g^*, and

η is an independent and identically distributed random variable. A standard normal distribution

with a mean equal to zero and a standard deviation equal to one for η was used to generate a

rugged expression landscape. To construct an expression landscape, g^* was first chosen at

random from the genotype space. $E(g)$ was subsequently calculated using the above equation.

The expression landscape was then normalized using the minimal and maximal values of $E(g)$

to range from zero to one, as follows:

703
$$\hat{E}(g) = \frac{E(g) - min\big(E(g)\big)}{max\big(E(g)\big) - min\,\big(E(g)\big)} \; .$$

704 The fitness of a given genotype g was defined as:

705
$$F(g) = (1 - F_b)\hat{E}(g) + F_b \; ,$$

where $1 \geq F_b \geq 0$. F_b represents the basal fitness when the expression level is zero and determines

the strength of the selection pressure acting on the expression level. Different values of F_b (0.3,

0.5, and 0.99) were used to demonstrate the evolution under strong positive selection, moderate

positive selection, and nearly neutral evolution, respectively. It was assumed that essential

genes underwent strong positive selection on their expression levels, while nonessential genes

underwent relatively moderate selection for their expression levels, based on the empirical fact

that knock-down of essential genes was more harmful for cell growth than nonessential

genes(Li, et al. 2016). Ten independent fitness landscapes were constructed for each F_b.

Evolutionary simulation on the fitness landscapes were performed using the standard Wright-

Fisher model(Otto and Day 2007) assuming a haploid asexual population with a population size

of 10^5 and a mutation rate of 10^{-3} mutations per genotype per generation. The initial genotypes

were randomly selected for each simulation, and 600 independent simulations were examined

for each fitness landscape. For each generation, mutants with genotypes differing by one

Hamming distance from the current population were generated at the mutation rate. The

frequency of each genotype within the population was updated based on the fitness landscape.

Subsequently, a new population for the next generation was formed by sampling mutants, with

replacement, from the current population using a binomial distribution with the given

population size. The most frequent mutants in a population after 30 generations were isolated

in each simulation, resulting in 6,000 mutants for each F_b. Mutational changes in expression

levels were calculated for each isolate by averaging the expression levels of all nearest

neighboring genotypes that differed by one Hamming distance from the focal isolate. Custom

R scripts were used for the numerical simulations, modified from scripts constructed in a

previous study(Obolski, et al. 2018; Song and Zhang 2021).

Data visualization

All figures were generated in R and the following packages were used. Phylogenetic trees of

the promoters were visualized using the ape(Paradis and Schliep 2019) and ggtree(Yu, et al.

2017) packages. Other illustrative plots were generated using the ggplot2(Wickham 2016) and

ggpubr(Kassambara 2023) packages.

Data Availability

Custom R scripts used in the manuscript are available at GitHub (https://github.com/tsuruubi/Variability_PromoterAct_E_coli).

The raw fcs data are available at Zenodo (wild types for https://doi.org/10.5281/zenodo.11108555; mutant libraries for https://doi.org/10.5281/zenodo.11108786).

Acknowledgments

The authors thank Hiroyo Koike for technical assistance of Sanger sequencing. The authors also appreciate Dr. Olin Silander and Dr. Luise Wolf for fruitful technical advice on the construction of a synthetic promoter library. The pNS2-σVL plasmid was a gift from Dr. Michael Elowitz (Addgene, plasmid # 26756).

Funding

This study was supported by the Japan Society for the Promotion of Science (JSPS) KAKENHI (18H02427, 22H05403 to S.T.;17H06389 to C.F. and S.T.; 22K21344 to C.F.), and the Japan Science and Technology Agency (JST) ERATO (JPMJER1902 to S.T. and C.F.).

References

Aita T, Uchiyama H, Inaoka T, Nakajima M, Kokubo T, Husimi Y. 2000. Analysis of a local

fitness landscape with a model of the rough Mt. Fuji-type landscape: application to prolyl

endopeptidase and thermolysin. Biopolymers 54:64-79.

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search

tool. Journal of Molecular Biology 215:403-410.

Belliveau NM, Barnes SL, Ireland WT, Jones DL, Sweredoski MJ, Moradian A, Hess S,

Kinney JB, Phillips R. 2018. Systematic approach for dissecting the molecular mechanisms of

transcriptional regulation in bacteria. Proc Natl Acad Sci U S A 115:E4796-E4805.

Benjamini Y, Hochberg Y. 1995. Controlling the False Discovery Rate: A Practical and

Powerful Approach to Multiple Testing. Journal of the Royal Statistical Society: Series B

(Methodological) 57:289-300.

Bilu Y, Barkai N. 2005. The design of transcription-factor binding sites is affected by

combinatorial regulation. Genome Biol 6:R103.

Blattner FR, Plunkett G, Bloch CA, Perna NT, Burland V, Riley M, Collado-Vides J,

Glasner JD, Rode CK, Mayhew GF, et al. 1997. The complete genome sequence of

Escherichia coli K-12. Science 277:1453-1462.

Bodenhofer U, Bonatesta E, Horejs-Kainrath C, Hochreiter S. 2015. msa: an R package for

multiple sequence alignment. Bioinformatics 31:3997-3999.

Casadesus J, Low D. 2006. Epigenetic gene regulation in the bacterial world. Microbiol

Mol Biol Rev 70:830-856.

Cui L, Vigouroux A, Rousset F, Varet H, Khanna V, Bikard D. 2018. A CRISPRi screen in

E. coli reveals sequence-specific toxicity of dCas9. Nature Communications 9:1-10.

Darwin C. 1859. On the origin of species by means of natural selection, or preservation of

favoured races in the struggle for life: London : John Murray.

de Visser JA, Hermisson J, Wagner GP, Ancel Meyers L, Bagheri-Chaichian H, Blanchard

JL, Chao L, Cheverud JM, Elena SF, Fontana W, et al. 2003. Perspective: Evolution and

detection of genetic robustness. Evolution 57:1959-1972.

Denver DR, Morris K, Streelman JT, Kim SK, Lynch M, Thomas WK. 2005. The

transcriptional consequences of mutation and natural selection in *Caenorhabditis elegans*. Nat

Genet 37:544-548.

Draghi JA, Whitlock MC. 2012. Phenotypic plasticity facilitates mutational variance,

genetic variance, and evolvability along the major axis of environmental variation. Evolution

66:2891-2902.

Dunlop MJ, Cox RS, Levine JH, Murray RM, Elowitz MB. 2008. Regulatory activity

revealed by dynamic correlations in gene expression noise. Nature Genetics 40:1493-1498.

Duveau F, Vande Zande P, Metzger BP, Diaz CJ, Walker EA, Tryban S, Siddiq MA, Yang

B, Wittkopp PJ. 2021. Mutational sources of trans-regulatory variation affecting gene

expression in Saccharomyces cerevisiae. Elife 10.

Ellis B, Haaland P, Hahne F, Meur L, Nolwenn, Gopalakrishnan N, Spidlen J, Jiang M,

Finak G, Granjeaud S. 2024. flowCore: flowCore: Basic structures for flow cytometry data. R

package version 2.14.2:https://bioconductor.org/packages/flowCore/.

Furusawa C, Kaneko K. 2018. Formation of dominant mode by evolution in biological

systems. Phys Rev E 97:042410.

Goodall ECA, Robinson A, Johnston IG, Jabbari S, Turner KA, Cunningham AF, Lund PA,

Cole JA, Henderson IR. 2018. The essential genome of *Escherichia coli* K-12. mBio 9:1-18.

Halligan DL, Keightley PD. 2009. Spontaneous mutation accumulation studies in

evolutionary genetics. Annual Review of Ecology, Evolution, and Systematics 40:151-172.

Hawkins JS, Silvis MR, Koo BM, Peters JM, Osadnik H, Jost M, Hearne CC, Weissman

JS, Todor H, Gross CA. 2020. Mismatch-CRISPRi Reveals the Co-varying Expression-

Fitness Relationships of Essential Genes in *Escherichia coli* and *Bacillus subtilis*. Cell Syst

11:523-535 e529.

Ho WC, Zhang J. 2016. Adaptive Genetic Robustness of *Escherichia coli* Metabolic

Fluxes. Molecular Biology and Evolution 33:1164-1176.

Ho WC, Zhang J. 2014. The genotype-phenotype map of yeast complex traits: basic

parameters and the role of natural selection. Mol Biol Evol 31:1568-1580.

Hornung G, Oren M, Barkai N. 2012. Nucleosome organization affects the sensitivity of

gene expression to promoter mutations. Mol Cell 46:362-368.

Ireland WT, Beeler SM, Flores-Bautista E, McCarty NS, Roschinger T, Belliveau NM,

Sweredoski MJ, Moradian A, Kinney JB, Phillips R. 2020. Deciphering the regulatory

genome of *Escherichia coli*, one hundred promoters at a time. Elife 9.

Kaneko K. 2009. Relationship among phenotypic plasticity, phenotypic fluctuations,

robustness, and evolvability; Waddington's legacy revisited under the spirit of Einstein. J

Biosci 34:529-542.

Kassambara A. 2023. ggpubr: 'ggplot2' Based Publication Ready Plots. R package version

0.6.0:https://rpkgs.datanovia.com/ggpubr/.

Kauffman SA, Weinberger ED. 1989. The NK model of rugged fitness landscapes and its

application to maturation of the immune response. J Theor Biol 141:211-245.

Keseler IM, Gama-Castro S, Mackie A, Billington R, Bonavides-Martinez C, Caspi R,

Kothari A, Krummenacker M, Midford PE, Muniz-Rascado L, et al. 2021. The EcoCyc

Database in 2021. Front Microbiol 12:711077.

Kinney JB, Murugan A, Callan CG, Jr., Cox EC. 2010. Using deep sequencing to

characterize the biophysical mechanism of a transcriptional regulatory sequence. Proc Natl

Acad Sci U S A 107:9158-9163.

Lamoureux CR, Decker KT, Sastry AV, Rychel K, Gao Y, McConn JL, Zielinski DC,

Palsson BO. 2022. A multi-scale transcriptional regulatory network knowledge base for

Escherichia coli. bioRxiv:2021.2004.2008.439047.

Landry CR, Lemos B, Rifkin SA, Dickinson WJ, Hartl DL. 2007. Genetic properties

influencing the evolvability of gene expression. Science 317:118-121.

Levenshtein VI editor. Soviet physics doklady. 1966.

Li XT, Jun Y, Erickstad MJ, Brown SD, Parks A, Court DL, Jun S. 2016. tCRISPRi:

tunable and reversible, one-step control of gene expression. Sci Rep 6:39076.

Lutz R. 1997. Independent and tight regulation of transcriptional units in *Escherichia coli*

via the LacR/O, the TetR/O and AraC/I1-I2 regulatory elements. Nucleic Acids Research

25:1203-1210.

Lynch M, Walsh B. 1998. Genetics and analysis of quantitative traits: Sinauer Sunderland,

MA.

Meiklejohn CD, Hartl DL. 2002. A single mode of canalization. Trends in Ecology &

Evolution 17:468-473.

Metzger BP, Yuan DC, Gruber JD, Duveau F, Wittkopp PJ. 2015. Selection on noise

constrains variation in a eukaryotic promoter. Nature 521:344-347.

Neidhart J, Szendro IG, Krug J. 2014. Adaptation in tunably rugged fitness landscapes: the

rough Mount Fuji model. Genetics 198:699-721.

Noble DWA, Radersma R, Uller T. 2019. Plastic responses to novel environments are

biased towards phenotype dimensions with high additive genetic variation. Proc Natl Acad

Sci U S A 116:13452-13461.

Obolski U, Ram Y, Hadany L. 2018. Key issues review: evolution on rugged adaptive

landscapes. Rep Prog Phys 81:012602.

Oshima T, Wada C, Kawagoe Y, Ara T, Maeda M, Masuda Y, Hiraga S, Mori H. 2002.

Genome-wide analysis of deoxyadenosine methyltransferase-mediated control of gene

expression in Escherichia coli. Mol Microbiol 45:673-695.

Otto SP, Day T. 2007. A Biologist's Guide to Mathematical Modeling in Ecology and

Evolution: Princeton University Press.

Paradis E, Schliep K. 2019. ape 5.0: an environment for modern phylogenetics and

evolutionary analyses in R. Bioinformatics 35:526-528.

Payne JL, Wagner A. 2015. Mechanisms of mutational robustness in transcriptional

regulation. Front Genet 6:322.

R Core Team. 2023. R: A Language and Environment for Statistical Computing.

Reynolds R, Bermúdez-Cruz RM, Chamberlin MJ. 1992. Parameters affecting transcription

termination by *Escherichia coli* RNA polymerase. Journal of Molecular Biology 224:31-51.

Rifkin SA, Houle D, Kim J, White KP. 2005. A mutation accumulation assay reveals a

broad capacity for rapid evolution of gene expression. Nature 438:220-223.

Ringquist S, Shinedling S, Barrick D, Green L, Binkley J, Stormo GD, Gold L. 1992.

Translation initiation in *Escherichia coli*: sequences within the ribosome-binding site.

Molecular Microbiology 6:1219-1229.

Rychel K, Decker K, Sastry AV, Phaneuf PV, Poudel S, Palsson BO. 2021. iModulonDB: a

knowledgebase of microbial transcriptional regulation derived from machine learning.

Nucleic Acids Res 49:D112-D120.

Sastry AV, Gao Y, Szubin R, Hefner Y, Xu S, Kim D, Choudhary KS, Yang L, King ZA,

Palsson BO. 2019. The *Escherichia coli* transcriptome mostly consists of independently

regulated modules. Nat Commun 10:5536.

Siegal ML, Bergman A. 2002. Waddington's canalization revisited: developmental stability

and evolution. Proc Natl Acad Sci U S A 99:10528-10532.

Silander OK, Nikolic N, Zaslaver A, Bren A, Kikoin I, Alon U, Ackermann M. 2012. A

genome-wide analysis of promoter-mediated phenotypic noise in *Escherichia coli*. PLoS

Genet 8:e1002443.

Smith JM, Burian R, Kauffman S, Alberch P, Campbell J, Goodwin B, Lande R, Raup D,

Wolpert L. 1985. Developmental constraints and evolution: a perspective from the Mountain

Lake conference on development and evolution. The Quarterly Review of Biology 60:265-

287.

Song S, Zhang J. 2021. Unbiased inference of the fitness landscape ruggedness from

imprecise fitness estimates. Evolution 75:2658-2671.

Szollosi GJ, Derenyi I. 2009. Congruent evolution of genetic and environmental robustness

in micro-RNA. Mol Biol Evol 26:867-874.

Taniguchi Y, Choi PJ, Li GW, Chen H, Babu M, Hearn J, Emili A, Xie XS. 2010.

Quantifying *E. coli* proteome and transcriptome with single-molecule sensitivity in single

cells. Science 329:533-538.

Tierrafria VH, Rioualen C, Salgado H, Lara P, Gama-Castro S, Lally P, Gomez-Romero L,

Pena-Loredo P, Lopez-Almazo AG, Alarcon-Carranza G, et al. 2022. RegulonDB 11.0:

Comprehensive high-throughput datasets on transcriptional regulation in *Escherichia coli* K-

12. Microb Genom 8.

Tirosh I, Weinberger A, Carmi M, Barkai N. 2006. A genetic signature of interspecies

variations in gene expression. Nat Genet 38:830-834.

Tsuru S, Furusawa C. 2024. Genetic properties underlying transcriptional variability in

Escherichia coli. bioRxiv:2024.2004.2015.589659.

Tsuru S, Ichinose J, Kashiwagi A, Ying BW, Kaneko K, Yomo T. 2009. Noisy cell growth

rate leads to fluctuating protein concentration in bacteria. Phys Biol 6:036015.

Tsuru S, Ishizawa Y, Shibai A, Takahashi Y, Motooka D, Nakamura S, Yomo T. 2015.

Genomic confirmation of nutrient-dependent mutability of mutators in *Escherichia coli*.

Genes Cells 20:972-981.

Uller T, Moczek AP, Watson RA, Brakefield PM, Laland KN. 2018. Developmental Bias

and Evolution: A Regulatory Network Perspective. Genetics 209:949-966.

Vaishnav ED, de Boer CG, Molinet J, Yassour M, Fan L, Adiconis X, Thompson DA, Levin

JZ, Cubillos FA, Regev A. 2022. The evolution, evolvability and engineering of gene

regulatory DNA. Nature 603:455-463.

Waddington CH. 1942. Canalization of development and the inheritance of acquired

characters. Nature 150:563-565.

Waddington CH. 1957. The strategy of the genes: Routledge.

Walker JA. 2007. A general model of functional constraints on phenotypic evolution. Am

Nat 170:681-689.

Wickham H. 2016. ggplot2: Elegant Graphics for Data

Analysis.https://ggplot2.tidyverse.org.

Wittkopp PJ, Haerum BK, Clark AG. 2004. Evolutionary changes in cis and trans gene

regulation. Nature 430:85-88.

Wolf L, Silander OK, van Nimwegen E. 2015. Expression noise facilitates the evolution of

gene regulation. Elife 4:1-48.

Ying BW, Tsuru S, Seno S, Matsuda H, Yomo T. 2014. Gene expression scaled by distance

to the genome replication site. Mol Biosyst 10:375-379.

Yona AH, Alm EJ, Gore J. 2018. Random sequences rapidly evolve into de novo

promoters. Nat Commun 9:1530.

Yu G, Smith DK, Zhu H, Guan Y, Lam TT-Y. 2017. ggtree: an r package for visualization

and annotation of phylogenetic trees with their covariates and other associated data. Methods

in Ecology and Evolution 8:28-36.

Zaslaver A, Bren A, Ronen M, Itzkovitz S, Kikoin I, Shavit S, Liebermeister W, Surette

MG, Alon U. 2006. A comprehensive library of fluorescent transcriptional reporters for

Escherichia coli. Nature Methods 3:623-628.

Zheng W, Gianoulis TA, Karczewski KJ, Zhao H, Snyder M. 2011. Regulatory variation

within and between species. Annu Rev Genomics Hum Genet 12:327-346.

    Figures and Figure legends

Fig. 1: Transcriptional variability of essential genes in *E. coli*.

(**A–D**) Transcriptional profiles of six independent mutants of *E. coli* cultured under identical

environmental conditions, termed the Mut dataset. The mutants were created through the MA

experiment to accumulate random genomic mutations by genetic drift. (**E–I**) Transcriptional

profiles of a single strain, MG1655, cultured under 76 different environmental conditions,

termed the Env dataset. (**A**, **E**) Schematics of the datasets. (**B**, **F**) Relationship between the

means and standard deviations of mRNA expression levels for thousands of genes. The magenta

and cyan lines represent running medians and their smoothed splines, respectively. Spearman's

rank correlation coefficients (R) and p-values are shown. (**C**, **G**) Mean expression levels of

essential and nonessential genes. (**D**, **H**) Transcriptional variability, represented as either DM_{mut}

or DM_{env}, of essential and nonessential genes. The DM values were calculated as the vertical

distance of the standard deviations from the smoothed splines of the running median of standard

deviations in panels **B** and **F**. The lower and upper edges of the boxes in panels **C**, **D**, **G**, **H**

represent the first (q1) and third (q3) quartiles, respectively. The horizontal lines in the boxes

represent the medians (m). The whiskers from the boxes extend to the most extreme observed

values inside inner fences, $m\pm1.5(q3-q1)$. (**I**) Relationship between mean expression levels and

the shorter distance from *oriC* on the *E. coli* chromosome. Red and blue dots in panels **B**, **F**,

and **I** represent essential and nonessential genes, respectively.

Fig. 2: Experimental design to measure expression levels of mutant promoter library.

(**A**) Natural promoters of *E. coli* were categorized into nonessential and essential promoters,

termed Nes and Ess, respectively. Nonessential promoters control transcriptional units

comprising only nonessential genes, while essential promoters regulate transcriptional units

containing at least one essential gene. Gene essentiality was based on the requirement for cell

growth, as defined by Goodall et al(Goodall, et al. 2018). The promoter regions were obtained

from the *E. coli* Promoter Collection(Zaslaver, et al. 2006). (**B**) Synthetic promoters, termed

Syn, were derived from a random oligo DNA pool. Transformants harboring a plasmid on which

the random sequences were placed upstream of the *gfp* gene were analyzed using flow

cytometry. The 16 unique transformants, satisfying a defined green fluorescence (red shaded

area), were isolated. The obtained synthetic promoters were named SP1–16. (**C**) Natural and

synthetic promoters were placed upstream of *gfp* on a plasmid. Transformants harboring each

of these plasmids were analyzed using flow cytometry. Mean green fluorescence of each

population is shown as small dots. The large dots and error bars represent the means and

standard deviations within the groups, respectively. The number of promoters is indicated inside

the panel. The p-value corresponds to ANOVA. (**D**) Schematic showing construction of mutant

library from a given promoter region. The promoter region was subjected to random

mutagenesis by error-prone PCR. A mutant library was created through transformation to yield

more than 100,000 genetically-different mutants in a population. Unmutagenized plasmid was

also transformed to create wild type clones. (**E**) Representative examples of $\log_{10}$-transformed

green fluorescence intensity distributions of wild type and mutant library for the promoter

region of the *metG* gene in *E. coli* (NPmetG). The means (vertical lines) and standard deviations

(horizontal arrows) were calculated in each population.

Fig. 3: Mutational variability in promoter activity.

(**A**) Comparison of the standard deviations in the GFP distributions between wild types and

mutant libraries. The solid line represents y=x. (**B**, **C**) Relationship between the mean and

standard deviation of the GFP distributions for wild types (**B**) and mutant libraries (**C**). Red

lines represent a regression line for wild types and were used to compensate for the mean-

dependency of the standard deviations of both wild types and mutant libraries by subtraction.

(**D**) Mutational variability in promoter activity, term MV_{prm}, is shown for each promoter group.

The MV_{prm} was calculated by subtracting the compensated standard deviations of wild types

from those of mutant libraries. The large dots and error bars are the means and standard

deviations, respectively. Pairwise Wilcoxon test was examined, where ns represents no

significance. (**E**, **F**) The relationships between MV_{prm} and the DM values (DM_{env} for **E** and

DM_{mut} for **F**). The DM values were defined in **Fig. 1**. (**G**, **H**) Relationship between MV_{prm} and

the number of known activation (**G**) and inhibition (**H**) of natural promoters by transcriptional

regulators. The relationship between promoters and known transcriptional regulators were

based on RegulonDB(Tierrafria, et al. 2022) and EcoCyc(Keseler, et al. 2021). The grey, blue,

and red dots in each panel represent synthetic (Syn), nonessential (Nes), and essential (Ess)

promoters (defined in **Fig. 2**), respectively. Spearman's R and p-values are shown in panes **A**–

C and **E**–**H**.

Fig. 4: Biased mutational changes in activity of essential promoters.

(**A**) Comparison of the means in the GFP distributions between wild types and mutant libraries.

The solid line represents y=x. (**B**) Relationship between $\log_{10}$-fold change of mean expression

levels from wild types to mutant libraries and the expression levels of wild types. The blue line

represents a regression line used to compensate for the mean-dependency of the $\log_{10}$-fold

changes by subtraction. (**C**, **D**) Relationship between activation bias in natural promoters and

the unnormalized (**C**) and normalized (**D**) $\log_{10}$-fold changes. The activation bias was calculated

by subtracting the number of known inhibitions from those of activations by transcriptional

regulators. Spearman's R and p-values are shown in panes **A–D**. Grey, blue, and red dots in

each panel represent synthetic (Syn), nonessential (Nes), and essential (Ess) promoters (defined

in **Fig. 2**), respectively. (**E**, **F**) Unnormalized (**E**) and normalized (**F**) $\log_{10}$-fold changes for

each promoter group. The large dots and error bars are the means and standard deviations,

respectively. Asterisks represent significance levels in pairwise Wilcoxon test (ns: $p > 0.05$; *:

$p \leq 0.05$; **: $p \leq 0.01$). The p-values were adjusted using the BH method(Benjamini and

Hochberg 1995).

1012   Fig. 5: Numerical simulation of promoter evolution on a ragged fitness landscape.

(**A–C**) Schematics of landscapes of expression levels (**A**, **B**) and fitness (**C**). (**A**) Expression

landscape was constructed by the rough Mt. Fuji (RNF) model. (**B**) Mutational change of each

genotype was calculated as the mean of differential expression levels of the nearest neighboring

genotypes (arrow heads, mutant types) from the focal genotype (colored circles, wild type). The

nearest neighbors differ from the focal genotype by one Hamming distance. The color bar

represents the fraction of the genotypes with higher expression levels than the focal genotype

among the nearest neighbors, termed uphill ratio. (**C**) Fitness landscape was modeled by adding

basal fitness (F_b, red line) to the expression landscape and normalizing it. (**D**, **E**)

Representatives of expression landscape (**D**) and mutational change (**E**). Genotype was

modeled as 10-bit binary numbers. Hamming distance represents the distance from the

genotype with the global optima in expression level and fitness. (**F**) Relationship between

expression level and uphill ratio. The uphill ratio was equally divided into three classes (Low,

Mid, and High). Expression level was also equally divided into ten classes. Frequency of each

class of uphill ratio was calculated as the number of corresponding genotypes divided by the

total number of genotypes within the class. (**G**, **H**) Representatives of fitness landscape (**G**) and

relationship between fitness and expression level (**H**) (F_b= 0.3). (**I**) Mutational changes of the

evolved isolates obtained through evolutionary simulation using the standard Wright-Fisher

model. The basal fitness was varied by mimicking strong positive (F_b=0.3, left), moderate

positive (F_b=0.5, middle), and nearly neutral selection (F_b=0.99, right) acting upon higher

expression levels. Ten independent expression/fitness landscapes were generated for each

condition of basal fitness. In each fitness landscape, 600 independent evolutionary simulations

were examined where initial genotypes were randomly selected. Dots represents the most

abundant mutants within the population after 30 generations in each run. Solid lines represent

the running medians. Dashed lines are guides for the eye. **(J)** Enlarged figure of the running

medians in **(I)**. **(K)** Expression level of the evolved isolates. Asterisks represent significance

levels of the adjusted p-value (the BH method) in the Wilcoxon test (****: $p \leq 0.0001$).