Promoter constrains evolution of expression levels of essential genes in

Escherichia coli

Saburo Tsuru[1,†,*], Naoki Hatanaka[1,†] and Chikara Furusawa[1,2,3,*]

[1]Universal Biology Institute, Graduate School of Science, The University of Tokyo, 7-3-1

Hongo, Bunkyo-ku, Tokyo 113-0033, Japan

[2]Department of Physics, Graduate School of Science, The University of Tokyo, 7-3-1 Hongo,

Bunkyo-ku, Tokyo 113-0033, Japan

[3]Center for Biosystems Dynamics Research (BDR), RIKEN, 6-2-3 Furuedai, Suita, Osaka 565-

0874, Japan

[†]These authors are equally contributed.

*Corresponding authors:

Saburo Tsuru

E-mail: saburotsuru@gmail.com

Chikara Furusawa

chikara.furusawa@riken.jp

Abstract

Variability in expression levels in response to random genomic mutations varies among genes, influencing both the facilitation and constraint of phenotypic evolution in organisms. Despite its importance, both the underlying mechanisms and evolutionary origins of this variability remain largely unknown due to the mixed contributions of *cis-* and *trans-*acting elements. To address this issue, we focused on the mutational variability of *cis*-acting elements, that is, promoter regions, in *Escherichia coli*. Random mutations were introduced into the natural and synthetic promoters to generate mutant promoter libraries. By comparing the variance in promoter activity of these mutant libraries, we found no significant difference in mutational variability in promoter activity between promoter groups, suggesting the absence of a signature of natural selection for mutational robustness. In contrast, the promoters controlling essential genes exhibited a remarkable bias in mutational variability, with mutants displaying higher activities than the wild types being relatively rare compared to those with lower activities. Our evolutionary simulation on a rugged fitness landscape provided a rationale for this vulnerability. These findings suggest that bias in mutational variability in promoters may constrain the evolution of downstream essential genes toward higher expression levels.

Introduction

The rate and direction of phenotypic evolution largely depend on phenotypic variability(Lynch and Walsh 1998; Walker 2007). It is widely recognized that organisms often exhibit nonuniform phenotypic variability, with certain variants arising more frequently than others in response to different genetic and environmental perturbations, even in the absence of natural selection(Darwin 1859; Waddington 1957; Smith, et al. 1985; Noble, et al. 2019). Therefore, it is crucial to elucidate the mechanisms underlying the bias in phenotypic variability to understand why phenotypic evolution proceeds as it does(Uller, et al. 2018). Transcriptional variability of genes across different perturbations plays a significant role in phenotypic evolution because the evolution of many traits in organisms largely depends on the evolution of expression levels and their regulation(Wittkopp, et al. 2004; Zheng, et al. 2011). Interestingly, previous studies have identified differences in transcriptional variability from gene to gene in response to various perturbations(Denver, et al. 2005; Rifkin, et al. 2005; Landry, et al. 2007; Tsuru and Furusawa 2024), where genes related to cellular growth and maintenance tend to exhibit lower transcriptional variability against random genetic perturbations to the genome in yeast(Landry, et al. 2007) and bacteria(Tsuru and Furusawa 2024). These studies, supported by theoretical evidence(Kaneko 2009; Draghi and Whitlock 2012; Furusawa and Kaneko 2018), suggest that the transcriptional variability of genes is not a random occurrence devoid of biological significance but rather a product of natural selection.

What molecular mechanisms account for gene-to-gene differences in transcriptional

variability? In principle, both *cis*- and *trans*-acting elements can influence transcriptional

variability(Wittkopp, et al. 2004; Tirosh, et al. 2006; Landry, et al. 2007; Payne and Wagner

2015; Tsuru and Furusawa 2024). However, it remains largely unknown which elements

practically contribute to gene-to-gene differences and how. Previous studies(Denver, et al.

2005; Rifkin, et al. 2005; Landry, et al. 2007; Tsuru and Furusawa 2024) explored the

transcriptional variability against different genetic perturbations using mutation accumulation

(MA) experiments(Halligan and Keightley 2009), where independent mutants were generated

by accumulating genomic mutations through genetic drift, and their transcriptome profiles were

obtained under identical environmental conditions. Although MA experiments provide a

powerful method to quantify transcriptional variability in the absence of strong natural selection,

the indiscriminate accumulation of mutations on *cis*- and *trans*-acting elements(Tsuru, et al.

2015) complicates the identification of the molecular mechanism underlying transcriptional

variability. Accordingly, to overcome this difficulty, it is necessary to consider the contributions

of *cis*- and *trans*-acting elements separately.

Why have genes related to cellular growth and maintenance acquired low mutational

variability? The evolutionary origin of mutational variability is highly controversial(de Visser,

et al. 2003). Mutational variability could result from direct natural selection for mutational

robustness(Waddington 1942), or could be a byproduct of selection for robustness against

environmental perturbations(Meiklejohn and Hartl 2002; de Visser, et al. 2003). Several

biological traits, such as metabolic flux(Ho and Zhang 2016) and cellular morphology(Ho and

Zhang 2014), show signs of adaptive evolution through direct selection for mutational

robustness, where traits highly related to cellular growth fitness are likely to acquire lower

mutational variability through direct selection for mutational robustness. On the other hand,

other studies have identified congruence in trait variability between environmental and genetic

perturbations, such as in RNA structure(Szollosi and Derenyi 2009) and gene expression

levels(Landry, et al. 2007; Tsuru and Furusawa 2024), supporting the relevance of the byproduct

of selection for environmental robustness. As an alternative scenario, mutational variability may

spontaneously emerge with trait evolution, regardless of selection for robustness(Siegal and

Bergman 2002; de Visser, et al. 2003). To date, the most appropriate scenario for low

transcriptional variability of genes related to cellular growth and maintenance remains largely

unknown.

To address these questions, we focused on the mutational variability governed by cis-

acting elements, that is, promoter regions, of genes in *Escherichia coli*. We created a mutant

library for each promoter through random mutagenesis and analyzed the changes in promoter

activity from the wild type to mutant using flow cytometry. To explore the relevant evolutionary

scenarios for the mutational variability of promoters, we compared promoters controlling

essential genes for cellular growth with those controlling nonessential genes. These groups were

expected to experience different natural selection pressures on their transcriptional variability

in the past. In addition, random sequences were used to construct synthetic promoters that

experienced no selection for mutational robustness. Interestingly, when comparing the variance

in promoter activity of the mutant libraries, these three promoter groups exhibited similar

mutational variability in promoter activity, indicating the absence of signatures of natural

selection for mutational robustness. In contrast, the promoters for essential genes showed a

remarkable vulnerability to mutations compared to the other promoter groups, where variants

with higher activity than the wild type arose less frequently than variants with lower activity.

Our evolutionary simulation on a rugged fitness landscape explains the higher vulnerability of

the promoters for essential genes. These results suggest that biased mutational variability in

promoters may constrain evolution of downstream essential genes toward higher expression

levels.

Results

Higher expression levels and lower transcriptional variability of essential genes in _E. coli_

First, we analyzed the transcriptional variability of essential genes in _E. coli_ in response to various genetic perturbations (**Fig. 1A**). Previously(Tsuru and Furusawa 2024), we obtained transcriptome profiles of independent mutants that had accumulated genomic mutations through genetic drift via the mutation accumulation (MA) experiment(Tsuru, et al. 2015), termed the Mut dataset. Using this dataset, we quantified the mean and standard deviation of the expression levels for each of thousands of genes (**Fig. 1B**). To compensate for the mean-dependency of the standard deviations, we calculated the vertical distance from the smoothed spline of the running median of the standard deviations, termed DM_{mut}. The DM_{mut} values reflect the mean transcriptional variability in response to different genetic perturbations(Tsuru and Furusawa 2024). We categorized thousands of genes into essential and nonessential genes for cell growth, following the definition of gene essentiality by Goodall _et al_(Goodall, et al. 2018). We confirmed that essential genes showed higher expression levels than nonessential genes across the different mutants (**Fig. 1C**). We also found that essential genes had lower DM_{mut} than nonessential genes (**Fig. 1D**).

Next, we explored the transcriptional variability of essential genes in response to different environmental perturbations (**Fig. 1E,F**) using a public dataset(Sastry, et al. 2019; Rychel, et al. 2021) of the transcriptome profiles of a single strain of _E. coli_, MG1655, cultured

under 76 different environmental conditions, termed the Env dataset. Previously(Tsuru and

Furusawa 2024), we quantified the mean-controlled standard deviations, termed DM_{env}, using

a method similar to DM_{evo}. The DM_{env} reflects transcriptional variability in response to different

environmental perturbations. We confirmed that the essential genes exhibited higher expression

levels (**Fig. 1G**) and lower variability (**Fig. 1H**) than the nonessential genes across different

environmental conditions. Taken together, essential genes exhibited higher expression levels

and lower transcriptional variability in response to different perturbations, which is similar to

the characteristics of essential genes in yeast(Tirosh, et al. 2006).

Construction of mutant libraries of *E. coli* promoters to measure mutational effects

To explore the molecular mechanisms and signs of natural selection underlying the lower

transcriptional variability of essential genes, we focused on the promoter regions of genes with

different essentialities. In particular, we explored how promoter activity varies in response to

new mutations in the promoter regions and how this variability relates to essentiality of the

downstream genes. To this end, we used 56 natural promoters from the *E. coli* Promoter

Collection(Zaslaver, et al. 2006). Importantly, to ensure an unbiased comparison of changes in

promoter activities between different essentialities, we avoided the use of natural promoters

with very low activity that were highly enriched in the promoters of nonessential genes. To

achieve this, we selected natural promoters with moderate to high activity before mutagenesis.

This selection was based on the known promoter activities of the collection as measured by

Silander et al(Silander, et al. 2012). Using the RegulonDB database(Tierrafria, et al. 2022), we

retrieved information regarding known transcriptional units regulated by these promoters.

Based on the presence of essential genes in the transcription units, we categorized natural

promoters into nonessential and essential promoters, termed Nes and Ess, respectively (**Fig.**

2A). Nonessential promoters control transcriptional units comprising only nonessential genes,

whereas essential promoters regulate those that include at least a single essential gene. In

addition to these natural promoters, we created synthetic promoters, termed Syn, as controls

that experienced no or minimal selection for mutational robustness. To this end, we placed

random sequences (~200 bp) upstream of the *gfp* gene on a low-copy-number plasmid, pUA66-

mCherry (**Fig. 2B**, **supplementary fig. S1**). The library of mixed transformants containing this

plasmid was analyzed using flow cytometry. The top 7% of cells showing higher expression

levels of GFP (red shaded area in **Fig. 2B**) were sorted, and 16 unique clones were subsequently

isolated. Importantly, these synthetic promoters experienced selection for their expression

levels only once, and no longer experienced further mutagenesis or selection for expression

level. Accordingly, they can be regarded as a control group that is almost free from selection

for mutational robustness. Consequently, in the upstream of the GFP gene on the plasmid, we

constructed three promoter groups, Nes, Ess, and Syn, comprising different sequences with low

homology (**supplementary fig. S2, supplementary note**). The green fluorescence from GFP

was used as a measure of promoter activity(Zaslaver, et al. 2006; Silander, et al. 2012). Despite

the diversity in sequence, we confirmed that the mean activities of the three promoter groups

were comparable to each other, as designed (**Fig. 2C**). We also confirmed that the obtained

promoter activities were highly consistent with those measured by Silander et al(Silander, et al.

2012) (Spearman's R=0.94, p<0.05, **supplementary fig. S3A**). Compared to the Env dataset,

the activities of the natural promoters were positively correlated with known mRNA expression

levels of the genes downstream of the chromosomal promoters in the wild-type strain

(Spearman's R=0.64, **supplementary fig. S3B**), supporting that the wild-type promoter

activities monitored by our method considerably reflect the activities of chromosomal

promoters and their impact on the expression levels of downstream genes.

For each of these promoters, we introduced mutations by error-prone PCR at a low

frequency (6.3 mutations/kb on average, **Fig. 2D**) and integrated them into pUA66-mCherry to

create a mutant library (MT). Each mutant library had a genetic diversity of 10^5 or more

genotypes. The promoter activity of each mutant library was measured by flow cytometry based

on the green fluorescence derived from downstream *gfp*, and was compared with the green

fluorescence of the corresponding wild-type (WT) (**Fig. 2E**). The introduction of mutations was

limited to the promoter regions of the plasmid, while the chromosomal copies of the natural

promoters were intact to avoid any confounding effects. The green fluorescence intensity was

$\log_{10}$-transformed after subtracting the background (**supplementary fig. S4**), and the mean and

standard deviation, SD, of the GFP distribution were calculated (**Fig. 2E, supplementary fig.**

S5).

Mutational variability in promoter activity was similar between different promoter groups

184   We confirmed that all mutant libraries showed larger variations than the corresponding wild-type clones, as designed (**Fig. 3A**). In addition, we also confirmed the well-known negative correlation between mean expression levels and standard deviations in GFP distributions regardless of the introduction of mutations or whether the promoters were natural or synthetic (**Fig. 3B,C**), which is consistent with previous observations(Zaslaver, et al. 2006; Wolf, et al. 2015). To compare different promoters, we compensated for the mean-dependency of the standard deviations of both WT and MT by subtracting the regression line obtained from WT (**Fig. 3B,C**). We subsequently calculated MV_{prm}, a measure of mutational variability in promoter activity, by subtracting the compensated SD of WT from that of MT. Unexpectedly, MV_{prm}, showed similar values across the three promoter groups (Wilcoxon test, adjusted p-values>0.05), even though each promoter might show unique values (**Fig. 3D**). These results indicated that mutational variability of essential promoters showed no signature of natural selection for mutational robustness despite their importance in the regulation of genes essential for cellular growth. Moreover, MV_{prm} showed no significant positive correlations with DM_{env} and DM_{mut} of the downstream genes (**Fig. 3E,F**). Using the RegulonDB and EcoCyc(Keseler, et al. 2021) databases to retrieve the known regulatory relationships between genes, we counted

the number of regulations that each promoter receives and explored the impact of the number

of regulations on mutational variability. We found that a larger number of transcriptional

regulators, in particular activations (**Fig. 3G**), facilitated mutational variability in promoter

activity (**Fig. 3G,H**), probably due to a larger number of mutational target sites. Nevertheless,

the lack of significant correlations between MV_{prm} and DM values suggests that the mutational

variability of promoters provides no considerable explanation for the observed gene-to-gene

differences in transcriptional variability of the downstream genes in response to different

perturbations at the genome scale (**Fig. 1**). These results suggest that the gene-to-gene

differences in transcriptional variability in response to genome-wide perturbations may reflect

a certain *trans*-acting molecular mechanism, mainly other than the *cis*-acting one, consistent

with previous observations(Tsuru and Furusawa 2024). This implication was also rational

because promoter regions are relatively short in bacterial genomes, where the mutational target

sites within the promoter regions are small relative to the trans-acting mutational target sites for

genomic mutations.

Vulnerability in essential promotors against mutations

Next, we explored the directionality of the mutational changes in promoter activity. Consistent

with the low mutation rate in error-prone PCR, the mean expression levels of MT were similar

to those of WT (Spearman's R=0.99) (**Fig. 4A**). Nevertheless, we found that $\log_{10}$-fold changes

in mean expression levels from WT to MT followed a global tendency over expression levels,

where the promoters with higher activities in WT tended to decrease their activity in response

to mutations (Spearman's R=–056) (**Fig. 4B**). This tendency may reflect bias in the mode,

activation or inhibition, of transcriptional regulation(Kinney, et al. 2010; Belliveau, et al. 2018;

Ireland, et al. 2020). We hypothesized that promoters that receive more activation than

inhibition might tend to reduce their activity in response to mutations. To test this possibility,

we defined activation bias as the number of known activations subtracted by the number of

inhibitions (**supplementary table S1**). We found that the activation bias explained the observed

mutational changes (**Fig. 4C**). Furthermore, the activation bias also explained the $\log_{10}$-fold

changes after controlling the global trend using the regression line (**Fig. 4D**), indicating that a

promoter becomes more vulnerable as it receives more activation among promoters with similar

activities.

Interestingly, synthetic and nonessential promoters showed similar mutational changes

around zero, whereas essential promoters showed biased directionality in mutational changes

toward negative values (**Fig. 4E**), even though the mean activities were almost the same among

the three promoter groups (**Fig. 2C**). In addition, there was no significant difference in the

mutation rate in error-prone PCR between the essential and nonessential promoters

(**supplementary fig. S6, supplementary table S2**), supporting the notion that the observed

bias was not derived from a technical bias. This bias in essential promoters was retained after

compensating for the global trend (**Fig. 4F**). Accordingly, essential promoters were more

vulnerable than other groups in terms of promoter activity. In other words, this biased

directionality indicated the low frequency of the variants with higher activities than the wild-

types in essential promoters. This biased variability might limit the transcriptional variability

of essential genes in response to mutations in cis-regulatory regions, independently from

transcriptional variability in response to genomic mutations (**Fig. 1**, **3F**).

Vulnerability in essential promotors against mutations may reflect past selection for

higher activity

The difference in vulnerability between essential and nonessential promoters could reflect the

variation in past selection pressures acting on higher expression levels between essential and

nonessential genes. Based on the observation that essential genes tended to have higher

expression levels than nonessential genes (**Fig. 1**), we hypothesized that the remarkable

vulnerability of essential promoters reflects local optima in terms of promoter activity. That is,

essential promoters might have evolved toward higher activity through past evolution and might

have been stuck in local optima because of the high essentiality of downstream genes. In

contrast, nonessential promoters might have experienced relatively relaxed selection for higher

expression levels because of the low essentiality of downstream genes. To investigate the

emergence of differences in vulnerability depending on selection pressure, we performed a

numerical simulation of promoter evolution on a simple rugged fitness landscape (**Fig. 5**). For

simplicity, we modeled promoter genotype as 10-bit binary numbers such as 1101011011, and

expression level, equivalent to promoter activity, was defined by the Rough Mt. Fuji (RMF)

model, as illustrated in **Fig. 5A, D**. Expression levels were set to range from 0 to 1. Using this

expression landscape, mutational changes were calculated for each genotype as the mean

difference in the expression levels of all nearest neighbors with a single bit difference, or one

Hamming distance, from the expression level of the focal genotype (**Fig. 5B, E**). This simple

expression landscape captured the negative correlation between expression levels and

mutational changes (**Fig. 5E**), which is consistent with the global trend observed in our

experiment (**Fig. 4B**). This trend was likely due to the prevalence of local sinks (green dots) at

lower expression levels and frequent local peaks (red dots) at higher expression levels.

Importantly, even at the same expression levels, local peaks tended to have negative mutational

changes, whereas local sinks tended to show positive changes. The fitness landscape was then

modeled by adding the basal fitness, F_b, to the expression levels multiplied by $1 - F_b$, where

F_b ranged from 0 to 1 (**Fig. 5C, F**). Consequently, a smaller F_b corresponds to a steeper fitness

landscape, while a larger F_b results in a flatter fitness landscape. Based on the fact that

knockdown of essential genes is more deleterious for cell growth than nonessential genes(Li,

et al. 2016), we assigned lower and higher values of F_b for essential and nonessential promoters,

respectively. Thus, in our model, essential and nonessential promoters shared the same

expression landscape but experienced different strengths of selection pressures. Using the

constructed fitness landscape with different F_b values (0.3, 0.5, and 0.99), we performed an

evolutionary simulation based on the standard Wright-Fisher model(Otto and Day 2007). Initial

genotypes were randomly generated for each simulation. The most abundant mutants within the

evolved populations after 30 generations were isolated and subjected to measurement of

mutational changes (**Fig. 5 G,H**). We confirmed that the evolved isolates in steeper fitness

landscapes exhibited higher expression levels (**Fig. 5I**), which is consistent with the designed

selection pressure. Comparing the mutational changes between different F_b values, we found

that the mutants that evolved with smaller F_b showed negatively larger mutational changes,

even at the same expression levels (**Fig. 5H**). These results suggest that higher activity and

vulnerability in essential promoters are linked, and both can emerge simultaneously through

stronger positive selection acting on higher expression levels of downstream essential genes.

Similar results were also obtained using the NK model(Kauffman and Weinberger 1989),

suggesting the robustness of the trend (**supplementary fig. S7**).

Discussions

Transcriptional variability differs between genes. Using public transcriptome datasets for *E. coli*, we found that essential genes exhibited lower transcriptional variability across different mutants and environmental conditions than nonessential genes. To explore the underlying mechanism, we focused on the promoter regions. We quantified the changes in promoter activity changes in response to mutations by constructing mutant libraries of natural and synthetic promoters. As a result, essential promoters did not exhibit a significant difference in mutational variability compared with nonessential and synthetic promoters. Unexpectedly, we found that essential promoters showed remarkable vulnerability to mutations, where variants with higher activities than wild-type were rare and those with lower activity were abundant. These results imply that essential promoters have evolved to reach local optima in terms of promoter activity through past selections for higher expression levels of downstream essential genes. Our numerical evolutionary simulation confirmed the relevance of this scenario. Overall, our results suggest the evolutionary impact of essentiality on the directionality of the mutational variability of promoters and imply that the directional bias in mutational variability of promoter regions might constrain further evolution of expression levels of essential genes toward higher expression levels.

There are three well-known evolutionary scenarios for the emergence of mutational robustness in many biological systems(de Visser, et al. 2003), namely adaptive(Waddington

1942), congruent(Meiklejohn and Hartl 2002), and intrinsic(Siegal and Bergman 2002)

scenarios. The adaptive scenarios predict that essential traits have evolved to reduce mutational

variability and minimize the emergence of harmful mutants through direct selection for

mutational robustness. The congruent scenarios predict that mutational robustness is a

byproduct of evolution toward low variability against environmental perturbations. Contrary to

these two scenarios assuming adaptive evolution toward low variability, the intrinsic scenarios

predict that the mutational robustness of the traits emerges spontaneously and inherently along

with the emergence of traits without selection against variability. In the case of promoter activity,

an intrinsic scenario corresponds to the mutational robustness of promoters emerging

spontaneously with the emergence of promoter activity without selection for mutational

robustness, which corresponds to the mutational variability of synthetic promoters. Our results

demonstrated that both adaptive and congruent scenarios failed to explain the observed

mutational variability in natural promoters because essential promoters showed similar

mutational variability to the other promoters, and the natural promoters exhibited no significant

positive correlation with the transcriptional variability of downstream genes against different

environmental perturbations. We note that the adaptive scenario might explain the higher

expression levels, not transcriptional variability, of essential genes (**Fig. 1C, G**), which might

minimize the risk of mutational knock down for essential genes to a lethal level. In contrast, the

intrinsic scenario explained the observed mutational variability because the natural promoters

showed similar mutational variability to that of synthetic promoters. Thus, the mutational

variability of essential promoters is unlikely to be a consequence of adaptive evolution toward

robustness against genetic and environmental perturbations.

Why did essential promoters show no less mutational variability than other promoters,

despite the fact that the reduction in expression levels of essential genes is more likely to be

harmful to cell growth than nonessential genes(Cui, et al. 2018)? One possible reason is that

promoter sequences with significant mutational robustness are rare, although several possible

mechanisms underlying mutational robustness in promoter regions are hypothesized(Payne and

Wagner 2015). In other words, the expression landscape of the promoters might be rugged

everywhere as we modeled in **Fig. 5A**. In fact, synthetic promoters derived from different

random sequences achieved higher promoter activities equivalent to natural promoters, which

was consistent with previous observations(Wolf, et al. 2015; Yona, et al. 2018). A second

possible reason is that selection for higher expression levels on a rugged fitness landscape (**Fig.**

5C) prevents the essential promoters at local optima from traversing the expression landscape

to reach genotypes with higher mutational robustness. Consistently, essential promoters showed

biased mutational effects, where most mutations in essential promoters tended to reduce

promoter activity (**Fig. 4C**). Interestingly, this scenario agreed with the predictions from a

machine-learning model based on an empirical expression landscape of yeast

promoters(Vaishnav, et al. 2022). Thus, our results suggest several plausible hypotheses that

may be shared by different organisms. Ideally, future studies should explore evolutionary

scenarios for diverse promoters that were not tested in this study, and validate the hypothetical

landscape of promoters more thoroughly.

In this study, we focused only on mutations in promoter regions to explore the

mechanism underlying the lower transcriptional variability in essential genes. As a result, we

found that mutants with higher promoter activity were relatively rare for essential promoters.

However, this result does not exclude the contribution of any other mechanism underlying the

observed transcriptional variability in response to genome-wide perturbations (**Fig. 1**). Notably,

trans-acting mechanisms, such as transcriptional regulators, may also play a role. As it is, we

previously reported that genes exhibiting greater transcriptional variability across diverse

environmental and genetic perturbations tend to be regulated by a larger number of unique

transcriptional regulators in *E. coli*(Tsuru and Furusawa 2024). A higher number of regulators

provides more *trans*-mutational target sites for the regulated genes, thereby enhancing

transcriptional variability in response to genomic perturbations. Based on this evidence, it is

plausible that essential genes have less transcriptional regulation. To support this hypothesis,

we used data from RegulonDB and EcoCyc to compare the number of regulations between

essential and nonessential genes (**supplementary fig. S8**). First, we found a smaller fraction of

essential genes (90 of 248 genes, 36%) lacking known transcriptional regulators than

nonessential genes (1976 of 3995, 49%) (**supplementary fig. S8A**). This finding aligns with

the positive activation bias values observed (**Fig. 4F**), probably attributable to the rich

activation by sigma factors aimed at enhancing expression levels. On the other hand, among

the genes regulated by at least one transcriptional regulator, we observed that essential genes

received fewer transcriptional regulations than nonessential genes (**supplementary fig. S8B**),

supporting this hypothesis. This observation is consistent with previous findings in yeast(Bilu

and Barkai 2005). Thus, our study underscores the dual contributions of both *cis-* and *trans-*

acting mechanisms in constraining the transcriptional variability of essential genes.

Materials and Methods

Bacterial strains and medium

We used *E. coli* DH5α Electro-Cells (Takara Bio, Japan, 9027) to construct the pUA66-mCherry plasmid and its derivatives. LB medium (LB Broth, Miller, BD Difco, USA, 244620) supplemented with 25 μg/mL kanamycin (Kanamycin Sulfate, Wako, Japan, 115-00342), referred to as LB/Km medium, served as the selective medium. The SOC medium, derived from SOB medium (SOB Medium, BD Difco, USA, 244310) supplemented with 22 mM Glucose (Wako, Japan, 049-31165), was used for recovery culture pos-transformation. The *E. coli* K-12 MG1655 strain served as the host for the constructed plasmids and was employed for measuring promoter activity via flow cytometry. Bacterial cells for flow cytometry were cultured in a synthetic medium, a derivative of M9 minimal medium (pH 7.2), supplemented with 47.7 mM Na_2HPO_4 (Disodium Hydrogenphosphate 12-Water, Wako, Japan, 196-02835), 22.0 mM KH_2PO_4 (Wako, Japan, 164-22635), 8.55 mM NaCl (Wako, Japan, 191-01665), 9.35 mM NH_4Cl (Wako, Japan, 014-03005), 11 mM Glucose, 1 mM $MgSO_4$ (Magnesium Sulfate Heptahydrate , Wako, Japan, 138-00415), 50 μM $CaCl_2$ (Wako, Japan, 038-24985), and 3.8 μM thiamin (Thiamin Hydrochloride, Wako, Japan, 201-00852), with supplementation of 25 μg/mL Kanamycin when required.

Selection of natural promoters

The *E. coli* Promoter Collection(Zaslaver, et al. 2006) (Horizon Discovery Ltd., UK, PEC3877)

was used to obtain the promoter regions of the natural promoters in *E. coli*. Initially, we filtered

out the promoters with low activities that were near our detection limit based on the promoter

activities of this library measured by Silander et al(Silander, et al. 2012). Subsequently, 115

promoters were randomly selected from this subset library carrying the pUA66 plasmid. The

DNA sequences of these promoter regions were verified using Sanger sequencing (Azenta Life

Sciences, USA) and compared with the reference genome sequence of *E. coli* K-12

MG1655(Blattner, et al. 1997) (GenBank ID: U00096). Subsequently, 63 of the 115 strains

without mutations in their promoter sequences were identified, and plasmids were extracted

from these colonies. Among these, 56 strains were randomly selected for the subsequent

experiments (**supplementary table S1**). Each promoter was denoted by the "NP" (natural

promoter), followed by the name of the ORF present in the immediate downstream sequence in

the genome. For example, NPmetG represents the promoter regions upstream of the *metG* gene

on the chromosome.

Plasmid construction

Plasmid pUA66 was modified by replacing the promoter regions of *gfp*, flanked by the BamHI

and XhoI restriction sites, with a red fluorescence protein (mCherry) expression cassette. This

cassette comprised dual terminators RNAI and TSAL(Reynolds, et al. 1992), promoter PLtetO-

1(Lutz 1997), Shine-Dalgarno sequence SD8(Ringquist, et al. 1992) and codon-optimized

mCherry(Dunlop, et al. 2008) coding sequences. The terminators were obtained from pNS2-

σVL(Dunlop, et al. 2008)(Addgene, Plasmid #26756) and the assembled sequences of the other

elements were commercially synthesized (Fasmac Co., Ltd., Japan). The assembled cassette

was inserted between the BamHI and XhoI recognition sequences of the pUA66 plasmid in the

direction opposite to that of *gfp* (**supplementary fig. S1**), yielding plasmid pUA66-mCherry.

PCR amplification was performed using KOD One PCR Master Mix Blue (Toyobo, Japan,

KMM-201)   and   pUA66-mCherry   as   a   template   with   primers   A

(GCTTTCCAGGGATCCTCTAGATTTAAGAAGG)   and   B

(GAGCAGTACTCTCGAGGTGAAGACGAAAGG) to include all sequences except for the

mCherry expression cassette. The PCR product was column purified using the FastGene

Gel/PCR Extraction Kit (NIPPON Genetics, Japan, FG-91302). The purified PCR product was

double-digested with BamHI (Takara Bio, Japan, 1010A) and XhoI (Takara Bio, Japan, 1094A)

overnight at 37 °C, followed by column purification. The digested product is referred to as the

vector.

Construction of promoter library comprising of random sequences

The construction of the synthetic promoters was based on a previous study(Wolf, et al. 2015).

Single-stranded   oligonucleotides   (CCTTTCGTCTTCACCTCGAG-(N200)-

GGATCCTCTAGATTTAAGAAGG) were synthesized by Eurofins Genomics (Ebersberg,

Germany). The 5' and 3' ends of the oligos contained XhoI and BamHI restriction sites,

respectively, and were homologous to primers C (CCTTTCGTCTTCACCTCGA) and D

(CCTTCTTAAATCTAGAGGATCC), respectively (**supplementary fig. S1**). A 200 bases

region of random nucleotides was present between them. PCR amplification using primers C

and D was performed using a mixture of single-stranded oligos as a template to create double-

stranded DNA. The PCR products were column purified and double digested using BamHI and

XhoI for 3 h at 37 °C, followed by column purification. Thus an insert for the following ligation

reaction was obtained. The ligation reaction was performed using T4 DNA Ligase (Takara Bio,

Japan, 2011A) overnight at 16 °C involved mixing the vector and insert at a molar ratio of 1: 3.

The resulting ligation product solution was column purified and DNA was extracted using 20

μL of sterile water. All purified ligation product solutions were added to 50 μL of DH5α

Electro-Cells and electroporated at 1800 V (Eporator, Eppendroph, Germany, 4309000027).

Transformed cells were incubated at 37 °C for 1 h in 1 mL of SOC medium, followed by

overnight incubation at 37 °C in 1 mL of LB medium supplemented with 2 μL of 25 mg/mL

kanamycin solution.

Selection of synthetic promoters using flow cytometry

Green and red fluorescence, as well as forward scatter of the cells in the DH5α-transformed

random promoter library were measured using a flow cytometer (FACSAria III, BD, USA).

Gates were set for the green and red channels to exclude cells with low GFP and high mCherry

fluorescence, respectively. Subsequently, 10,000 cells without red fluorescence but with green

fluorescence within the selected range (**Fig. 2B**) were collected by cell sorting, according to the

manufacturer's instructions. The sorted cells were spread on LB/Km agar plates and incubated

overnight at 37 °C. Bacterial colonies were randomly picked and their inserted synthetic

promoter regions were sequenced by Sanger sequencing. Consequently, 16 synthetic promoters

without improper mutations at the restriction sites were identified and named SP1–16,

(**supplementary table S1**).

Random mutagenesis using error-prone PCR

Error-prone PCR was conducted using 0.1 ng of each promoter region as a template, with 35

cycles of amplification performed using the GeneMorph II Random Mutagenesis Kit (Agilent

Technologies, USA, 200550) and primers C and D. The resulting PCR products underwent gel

purification and were double digested with BamHI and XhoI for 3 hours at 37 °C, followed by

column purification to create mutant inserts. Ligation was performed overnight at 16 °C using

T4 DNA Ligase, with the vector and mutant inserts mixed at a molar ratio of 1:3. The ligation

reaction solution was column purified, and DNA was extracted using 20 µL of sterile water.

Construction of mutant promoter library

The purified ligation product solution was added to 50 µL of DH5α Electro-Cells and

electroporated at 1800 V. Transformed cells were cultured at 37 °C for 1 h in 1 mL of SOC

medium and a portion of the culture was plated on LB/Km agar medium for overnight growth

at 37 °C. Bacterial colonies were counted to determine the transformation efficiency. If the

transformation efficiency exceeded 10^5 cells/reaction, the next step was performed; otherwise,

the process was repeated. The transformed culture was diluted with 5 mL of LB/Km medium

and grown overnight. Plasmid extraction from an overnight culture was performed using the

FastGene Plasmid Mini Kit (NIPPON Genetics, Japan, FG-90502). The obtained plasmid

solution was added to 50 µL of MG1655 electrocompetent cells and electroporated at 1800 V.

Transformed cells were cultured at 37 °C for 1 h in 1 mL of SOC medium, and a portion of the

solution was plated on LB/Km agar medium for overnight growth at 37 °C to count the number

of colonies and determine transformation efficiency. If the transformation efficiency exceeded

10^5 cells/reaction, the next step was performed; otherwise, the process was repeated. A sterile

2 mL 96 well deep well plate (Greiner Bio-One, Austria, 780271) was used to add 1 mL of the

synthetic medium to each well, followed by the addition of 10 µL of the overnight culture

containing the MG1655 transformants. The bacterial cells were grown at 37 °C with shaking at

800 rpm until confluent growth was achieved. Subsequently, another plate was prepared with 1

mL of fresh synthetic medium added to each well, and 10 µL of the confluently grown cultures

was diluted into each well. The cultures were grown for 2.5 h at 37 °C with shaking at 800 rpm

and were used as MT for flow cytometry.

Verification of PCR random mutagenesis

The number of mutations introduced into the promoters by error-prone PCR was determined

using Sanger sequencing. Three to four colonies were randomly selected from six randomly

selected mutant libraries, including essential and nonessential promoters (**supplementary table**

S2). Colony PCR was performed using KOD One PCR Master Mix Blue with primers E

(TTACTTTGCAGGGCTTCCCAA) and F (CCAGTCTTTCGACTGAGCCT) to amplify the

regions including the promoter regions (**supplementary fig. S1**). PCR products were column

purified and verified by Sanger sequencing. The average mutation rate was calculated to be 6.2

mutations/kb.

Construction of WT promoter control group

To ensure accurate measurement of GFP fluorescence from cells with wild-type promoters,

wild-type plasmids were transformed into MG1655 cells by electroporation. Transformed

cultures were plated on LB/Km agar medium and incubated overnight, and the resulting

colonies were cultured in LB/Km medium. Cell cultures of these control groups in the synthetic

medium were prepared and used as WT for flow cytometry.

Flow cytometry

Bacterial cells were dispensed in M9 buffer prior to analysis using a flow cytometer (BD, USA,

FACSAria III). Fluorescence intensity from GFP (GFP FI) and mCherry (mCherry FI) was

measured using a 488 nm laser with 515–545-nm emission filter, and a 561 nm laser with 563–

588-nm emission filter, respectively. The following PMT voltage settings were used: forward

scatter (FSC), 300; side scatter (SSC), 340; green fluorescence, 700; red fluorescence, 600. The

events exceeding the defined thresholds of FSC (200) and SSC (200) were recorded.

Tota1,000,000 events were measured per library. The MG1655 strain was used as a control for

autofluorescence of GFP FI.

Data analysis of flow cytometry

FCS files were converted to CSV format using the flowCore(Ellis, et al. 2024) package in R(R

Core Team 2023). Subsequent analyses were conducted using custom R scripts. To compare

green fluorescence among cells in similar physiological states, a narrow gate for the forward

scatter channel was set to capture 40% of the events, including the most frequent values for

MG1655 (**supplementary fig. S4A**). The top 0.5% value of the green fluorescence intensity

distribution of MG1655 within the gate was determined as the autofluorescence intensity

(**supplementary fig. S4B**). For the events of WT and MT, the autofluorescence intensity was

subtracted from the green fluorescence intensity of the events within the gate. Events with a

negative green fluorescence intensity after subtracting the autofluorescence intensity were

excluded from the analysis. In addition, events with a red fluorescence intensity greater than

10^3 [a.u.] were regarded as the cells carrying pUA66-mCherry and were excluded from the

analysis. The green fluorescence intensity after subtracting the autofluorescence intensity was

$\log_{10}$-transformed and used to calculate the means and standard deviations.

Classification of promoter based on essentiality of transcribed unit

Gene essentiality was based on a previous study(Goodall, et al. 2018), which identified 248

genes designated as essential across different datasets. Promoters containing at least one

essential gene in the transcription units were defined as essential promoters. The known

transcription units were retrieved from the RegulonDB(Tierrafria, et al. 2022) database. Of the

56 promoters, 18 were classified as essential and 38 as nonessential as listed in **supplementary**

table S1.

Estimation of transcriptional variability

Transcriptional variability of each gene in *E. coli* in response to different genetic perturbations

was estimated using known transcriptome profiles obtained from six independent mutants

derive from MG1655 cultured under a single environmental condition, termed the Mut

dataset(Tsuru and Furusawa 2024). These mutants were subjected to mutation accumulation

(MA) experiments to accumulate genomic mutations via genetic drift(Halligan and Keightley

2009; Tsuru, et al. 2015). In addition, to quantify the transcriptional variability in response to

different environmental conditions, known transcriptome profiles of MG1655 cultured under

76 different environmental conditions(Sastry, et al. 2019; Rychel, et al. 2021; Lamoureux, et al.

2022), termed the Env dataset(Tsuru and Furusawa 2024), were utilized. The expression levels

were determined by $\log_2$-transformation and quantile normalization. The means and standard

deviations in expression levels for each gene across different genetic perturbations and

environmental conditions were quantified. To compensate for the mean-dependency of the

standard deviations (**Fig. 1B,G**), the DM values for each gene were calculated as the vertical

distance of each standard deviation from a smoothed running median of the standard deviations

for each dataset, as detailed previously(Tsuru and Furusawa 2024). The resulting DM values

were termed DM_{mut} and DM_{env} for the Mut and Env datasets, respectively. The DM values were used as measures of transcriptional variability in response to genetic and environmental perturbations.

Phylogenetic analysis of promoter sequence of length 73 bases

Synthetic promoters underwent a blastn(Altschul, et al. 1990) search to confirm that there was no considerable similarity to any natural sequences (**supplementary note**). Posterior sequences of 73 bases were aligned using the msa(Bodenhofer, et al. 2015) package in R (Clustal Omega), where 73 bases represented the shortest length of a synthetic promoter. A phylogenetic tree was constructed using the Neighbor-Joining method.

Phylogenetic analysis of core elements of sigma70 promoters

We identified sigma70 promoters from our natural promoters using RegulonDB. The core elements comprising the –10/–35 elements of these promoters were concatenated and subjected to phylogenetic analysis. The Levenshtein distance(Levenshtein 1966) between the focal elements and the consensus elements, which consist of TTGACA and TATAAT, was calculated. A phylogenetic tree was constructed as previously described.

Identification of the number of transcriptional regulators regulating the natural promoters

Reported regulatory interactions between genes were compiled using all interactions with experimental evidence from RegulonDB and EcoCyc(Keseler, et al. 2021) for both transcription

factors and sigma factors. The mode of regulation was classified as either activation or

inhibition, and the mode of regulation by sigma factors was classified as activation.

Numerical simulation of promoter evolution

The promoter genotype, g, was modeled by a bit string comprising either zeros or ones of length

N=10 (e.g., 1110001100). The expression level, E, was defined by the Rough Mt. Fuji (RMF)

model(Aita, et al. 2000; Neidhart, et al. 2014) as:

$$E(g) = -cD(g, g^*) + \eta(g),$$

where c is a scaling coefficient equal to one in this study, g^* is the genotype with the highest

expression level before adding noise, $D(g, g^*)$ is the Hamming distance between g and g^*, and

η is an independent and identically distributed random variable. A standard normal distribution

with a mean equal to zero and a standard deviation equal to one for η was used to generate a

rugged expression landscape. The expression landscape was then normalized using the minimal

and maximal values of $E(g)$ to range from zero to one, as follows:

$$\hat{E}(g) = \frac{E(g) - min\big(E(g)\big)}{max\big(E(g)\big) - min\big(E(g)\big)} .$$

The fitness of a given genotype g was defined as:

$$F(g) = (1 - F_b)\hat{E}(g) + F_b ,$$

where $1 \geq F_b \geq 0$. F_b represents the basal fitness when the expression level is zero and determines

the strength of the selection pressure acting on the expression level. Different values of F_b (0.3,

0.5, and 0.99) were used to demonstrate the evolution under strong positive selection, moderate

positive selection, and nearly neutral evolution, respectively. It was assumed that essential

genes underwent strong positive selection on their expression levels, while nonessential genes

underwent relatively moderate selection for their expression levels, based on the empirical fact

that knock-down of essential genes was more harmful for cell growth than nonessential

genes(Li, et al. 2016). Ten independent fitness landscapes were constructed for each F_b.

Evolutionary simulation on the fitness landscapes were performed using the standard Wright-

Fisher model(Otto and Day 2007) assuming a haploid asexual population with a population size

of 10^5 and a mutation rate of 10^{-3}. The initial genotypes were randomly selected for each

simulation, and 600 independent simulations were examined for each fitness landscape. For

each generation, mutants with genotypes differing by one Hamming distance from the current

population were generated at the mutation rate. The frequency of each genotype within the

population was updated based on the fitness landscape. Subsequently, a new population for the

next generation was formed by sampling mutants, with replacement, from the current

population using a binomial distribution with the given population size. The most frequent

mutants in a population after 30 generations were isolated in each simulation, resulting in 6,000

mutants for each F_b. Mutational changes in expression levels were calculated for each isolate

by averaging the expression levels of all nearest neighboring genotypes that differed by one

Hamming distance from the focal isolate. Custom R scripts were used for the numerical

simulations, modified from scripts constructed in a previous study(Obolski, et al. 2018; Song

and Zhang 2021).

Data visualization

All figures were generated in R and the following packages were used. Phylogenetic trees of

the promoters were visualized using the ape(Paradis and Schliep 2019) and ggtree(Yu, et al.

2017) packages. Other illustrative plots were generated using the ggplot2(Wickham 2016) and

ggpubr(Kassambara 2023) packages.

Data Availability

Custom R scripts used in the manuscript are available at GitHub (https://github.com/tsuruubi/Variability_PromoterAct_E_coli).

The raw fcs data are available at Zenodo (WT for https://doi.org/10.5281/zenodo.11108555; MT for https://doi.org/10.5281/zenodo.11108786).

Acknowledgments

The authors thank Hiroyo Koike for technical assistance of Sanger sequencing. The authors also appreciate Dr. Olin Silander and Dr. Luise Wolf for fruitful technical advice on the construction of a synthetic promoter library. The pNS2-σVL plasmid was a gift from Dr. Michael Elowitz (Addgene, plasmid # 26756).

Funding

This study was supported by the Japan Society for the Promotion of Science (JSPS) KAKENHI (18H02427, 22H05403 to S.T.;17H06389 to C.F. and S.T.; 22K21344 to C.F.), and the Japan Science and Technology Agency (JST) ERATO (JPMJER1902 to S.T. and C.F.).

References

Aita T, Uchiyama H, Inaoka T, Nakajima M, Kokubo T, Husimi Y. 2000. Analysis of a local

fitness landscape with a model of the rough Mt. Fuji-type landscape: application to prolyl

endopeptidase and thermolysin. Biopolymers 54:64-79.

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search

tool. Journal of Molecular Biology 215:403-410.

Belliveau NM, Barnes SL, Ireland WT, Jones DL, Sweredoski MJ, Moradian A, Hess S,

Kinney JB, Phillips R. 2018. Systematic approach for dissecting the molecular mechanisms

of transcriptional regulation in bacteria. Proc Natl Acad Sci U S A 115:E4796-E4805.

Benjamini Y, Hochberg Y. 1995. Controlling the False Discovery Rate: A Practical and

Powerful Approach to Multiple Testing. Journal of the Royal Statistical Society: Series B

(Methodological) 57:289-300.

Bilu Y, Barkai N. 2005. The design of transcription-factor binding sites is affected by

combinatorial regulation. Genome Biol 6:R103.

Blattner FR, Plunkett G, Bloch CA, Perna NT, Burland V, Riley M, Collado-Vides J, Glasner

JD, Rode CK, Mayhew GF, et al. 1997. The complete genome sequence of *Escherichia coli*

K-12. Science 277:1453-1462.

Bodenhofer U, Bonatesta E, Horejs-Kainrath C, Hochreiter S. 2015. msa: an R package for

multiple sequence alignment. Bioinformatics 31:3997-3999.

Cui L, Vigouroux A, Rousset F, Varet H, Khanna V, Bikard D. 2018. A CRISPRi screen in *E.*

coli reveals sequence-specific toxicity of dCas9. Nature Communications 9:1-10.

Darwin C. 1859. On the origin of species by means of natural selection, or preservation of

favoured races in the struggle for life: London : John Murray.

de Visser JA, Hermisson J, Wagner GP, Ancel Meyers L, Bagheri-Chaichian H, Blanchard JL,

Chao L, Cheverud JM, Elena SF, Fontana W, et al. 2003. Perspective: Evolution and

detection of genetic robustness. Evolution 57:1959-1972.

Denver DR, Morris K, Streelman JT, Kim SK, Lynch M, Thomas WK. 2005. The

transcriptional consequences of mutation and natural selection in *Caenorhabditis elegans*.

Nat Genet 37:544-548.

Draghi JA, Whitlock MC. 2012. Phenotypic plasticity facilitates mutational variance, genetic

variance, and evolvability along the major axis of environmental variation. Evolution

66:2891-2902.

Dunlop MJ, Cox RS, Levine JH, Murray RM, Elowitz MB. 2008. Regulatory activity

revealed by dynamic correlations in gene expression noise. Nature Genetics 40:1493-1498.

Ellis B, Haaland P, Hahne F, Meur L, Nolwenn, Gopalakrishnan N, Spidlen J, Jiang M, Finak

G, Granjeaud S. 2024. flowCore: flowCore: Basic structures for flow cytometry data. R

package version 2.14.2:https://bioconductor.org/packages/flowCore/.

Furusawa C, Kaneko K. 2018. Formation of dominant mode by evolution in biological

systems. Phys Rev E 97:042410.

Goodall ECA, Robinson A, Johnston IG, Jabbari S, Turner KA, Cunningham AF, Lund PA,

Cole JA, Henderson IR. 2018. The essential genome of *Escherichia coli* K-12. mBio 9:1-

18.

Halligan DL, Keightley PD. 2009. Spontaneous mutation accumulation studies in

evolutionary genetics. Annual Review of Ecology, Evolution, and Systematics 40:151-172.

Ho WC, Zhang J. 2016. Adaptive Genetic Robustness of *Escherichia coli* Metabolic Fluxes.

Molecular Biology and Evolution 33:1164-1176.

Ho WC, Zhang J. 2014. The genotype-phenotype map of yeast complex traits: basic

parameters and the role of natural selection. Mol Biol Evol 31:1568-1580.

Ireland WT, Beeler SM, Flores-Bautista E, McCarty NS, Roschinger T, Belliveau NM,

Sweredoski MJ, Moradian A, Kinney JB, Phillips R. 2020. Deciphering the regulatory

genome of *Escherichia coli*, one hundred promoters at a time. Elife 9.

Kaneko K. 2009. Relationship among phenotypic plasticity, phenotypic fluctuations,

robustness, and evolvability; Waddington's legacy revisited under the spirit of Einstein. J

Biosci 34:529-542.

Kassambara A. 2023. ggpubr: 'ggplot2' Based Publication Ready Plots. R package version

0.6.0:https://rpkgs.datanovia.com/ggpubr/.

Kauffman SA, Weinberger ED. 1989. The NK model of rugged fitness landscapes and its

application to maturation of the immune response. J Theor Biol 141:211-245.

Keseler IM, Gama-Castro S, Mackie A, Billington R, Bonavides-Martinez C, Caspi R,

Kothari A, Krummenacker M, Midford PE, Muniz-Rascado L, et al. 2021. The EcoCyc

Database in 2021. Front Microbiol 12:711077.

Kinney JB, Murugan A, Callan CG, Jr., Cox EC. 2010. Using deep sequencing to characterize

the biophysical mechanism of a transcriptional regulatory sequence. Proc Natl Acad Sci U

S A 107:9158-9163.

Lamoureux CR, Decker KT, Sastry AV, Rychel K, Gao Y, McConn JL, Zielinski DC, Palsson

BO. 2022. A multi-scale transcriptional regulatory network knowledge base for *Escherichia*

coli. bioRxiv:2021.2004.2008.439047.

Landry CR, Lemos B, Rifkin SA, Dickinson WJ, Hartl DL. 2007. Genetic properties

influencing the evolvability of gene expression. Science 317:118-121.

Levenshtein VI editor. Soviet physics doklady. 1966.

Li XT, Jun Y, Erickstad MJ, Brown SD, Parks A, Court DL, Jun S. 2016. tCRISPRi: tunable

and reversible, one-step control of gene expression. Sci Rep 6:39076.

Lutz R. 1997. Independent and tight regulation of transcriptional units in *Escherichia coli* via

the LacR/O, the TetR/O and AraC/I1-I2 regulatory elements. Nucleic Acids Research

25:1203-1210.

Lynch M, Walsh B. 1998. Genetics and analysis of quantitative traits: Sinauer Sunderland,

MA.

Meiklejohn CD, Hartl DL. 2002. A single mode of canalization. Trends in Ecology &

Evolution 17:468-473.

Neidhart J, Szendro IG, Krug J. 2014. Adaptation in tunably rugged fitness landscapes: the

rough Mount Fuji model. Genetics 198:699-721.

Noble DWA, Radersma R, Uller T. 2019. Plastic responses to novel environments are biased

towards phenotype dimensions with high additive genetic variation. Proc Natl Acad Sci U S

A 116:13452-13461.

Obolski U, Ram Y, Hadany L. 2018. Key issues review: evolution on rugged adaptive

landscapes. Rep Prog Phys 81:012602.

Otto SP, Day T. 2007. A Biologist's Guide to Mathematical Modeling in Ecology and

Evolution: Princeton University Press.

Paradis E, Schliep K. 2019. ape 5.0: an environment for modern phylogenetics and

evolutionary analyses in R. Bioinformatics 35:526-528.

Payne JL, Wagner A. 2015. Mechanisms of mutational robustness in transcriptional

regulation. Front Genet 6:322.

R Core Team. 2023. R: A Language and Environment for Statistical Computing.

Reynolds R, Bermúdez-Cruz RM, Chamberlin MJ. 1992. Parameters affecting transcription

termination by *Escherichia coli* RNA polymerase. Journal of Molecular Biology 224:31-51.

Rifkin SA, Houle D, Kim J, White KP. 2005. A mutation accumulation assay reveals a broad

capacity for rapid evolution of gene expression. Nature 438:220-223.

Ringquist S, Shinedling S, Barrick D, Green L, Binkley J, Stormo GD, Gold L. 1992.

Translation initiation in *Escherichia coli*: sequences within the ribosome-binding site.

Molecular Microbiology 6:1219-1229.

Rychel K, Decker K, Sastry AV, Phaneuf PV, Poudel S, Palsson BO. 2021. iModulonDB: a

knowledgebase of microbial transcriptional regulation derived from machine learning.

Nucleic Acids Res 49:D112-D120.

Sastry AV, Gao Y, Szubin R, Hefner Y, Xu S, Kim D, Choudhary KS, Yang L, King ZA,

Palsson BO. 2019. The *Escherichia coli* transcriptome mostly consists of independently

regulated modules. Nat Commun 10:5536.

Siegal ML, Bergman A. 2002. Waddington's canalization revisited: developmental stability

and evolution. Proc Natl Acad Sci U S A 99:10528-10532.

Silander OK, Nikolic N, Zaslaver A, Bren A, Kikoin I, Alon U, Ackermann M. 2012. A

genome-wide analysis of promoter-mediated phenotypic noise in *Escherichia coli*. PLoS

Genet 8:e1002443.

Smith JM, Burian R, Kauffman S, Alberch P, Campbell J, Goodwin B, Lande R, Raup D,

Wolpert L. 1985. Developmental constraints and evolution: a perspective from the

Mountain Lake conference on development and evolution. The Quarterly Review of

Biology 60:265-287.

Song S, Zhang J. 2021. Unbiased inference of the fitness landscape ruggedness from

imprecise fitness estimates. Evolution 75:2658-2671.

Szollosi GJ, Derenyi I. 2009. Congruent evolution of genetic and environmental robustness in

micro-RNA. Mol Biol Evol 26:867-874.

Tierrafria VH, Rioualen C, Salgado H, Lara P, Gama-Castro S, Lally P, Gomez-Romero L,

Pena-Loredo P, Lopez-Almazo AG, Alarcon-Carranza G, et al. 2022. RegulonDB 11.0:

Comprehensive high-throughput datasets on transcriptional regulation in *Escherichia coli*

K-12. Microb Genom 8.

Tirosh I, Weinberger A, Carmi M, Barkai N. 2006. A genetic signature of interspecies

variations in gene expression. Nat Genet 38:830-834.

Tsuru S, Furusawa C. 2024. Genetic properties underlying transcriptional variability in

Escherichia coli. bioRxiv:2024.2004.2015.589659.

Tsuru S, Ishizawa Y, Shibai A, Takahashi Y, Motooka D, Nakamura S, Yomo T. 2015.

Genomic confirmation of nutrient-dependent mutability of mutators in *Escherichia coli*.

Genes Cells 20:972-981.

Uller T, Moczek AP, Watson RA, Brakefield PM, Laland KN. 2018. Developmental Bias and

Evolution: A Regulatory Network Perspective. Genetics 209:949-966.

Vaishnav ED, de Boer CG, Molinet J, Yassour M, Fan L, Adiconis X, Thompson DA, Levin

JZ, Cubillos FA, Regev A. 2022. The evolution, evolvability and engineering of gene

regulatory DNA. Nature 603:455-463.

Waddington CH. 1942. Canalization of development and the inheritance of acquired

characters. Nature 150:563-565.

Waddington CH. 1957. The strategy of the genes: Routledge.

Walker JA. 2007. A general model of functional constraints on phenotypic evolution. Am Nat

170:681-689.

Wickham H. 2016. ggplot2: Elegant Graphics for Data Analysis.https://ggplot2.tidyverse.org.

Wittkopp PJ, Haerum BK, Clark AG. 2004. Evolutionary changes in cis and trans gene

regulation. Nature 430:85-88.

Wolf L, Silander OK, van Nimwegen E. 2015. Expression noise facilitates the evolution of

gene regulation. Elife 4:1-48.

Yona AH, Alm EJ, Gore J. 2018. Random sequences rapidly evolve into de novo promoters.

Nat Commun 9:1530.

Yu G, Smith DK, Zhu H, Guan Y, Lam TT-Y. 2017. ggtree: an r package for visualization and

annotation of phylogenetic trees with their covariates and other associated data. Methods in

Ecology and Evolution 8:28-36.

Zaslaver A, Bren A, Ronen M, Itzkovitz S, Kikoin I, Shavit S, Liebermeister W, Surette MG,

Alon U. 2006. A comprehensive library of fluorescent transcriptional reporters for

Escherichia coli. Nature Methods 3:623-628.

Zheng W, Gianoulis TA, Karczewski KJ, Zhao H, Snyder M. 2011. Regulatory variation

within and between species. Annu Rev Genomics Hum Genet 12:327-346.

Figures and Figure legends

Fig. 1: Transcriptional variability of essential genes in *E. coli*.

(**A**–**D**) Transcriptional profiles of six independent mutants of *E. coli* cultured under identical

environmental conditions, termed the Mut dataset. The mutants were created through the MA

experiment to accumulate random genomic mutations by genetic drift. (**E**–**H**) Transcriptional

profiles of a single strain, MG1655, cultured under 76 different environmental conditions,

termed the Env dataset. (**A**, **E**) Schematics of the datasets. (**B**, **F**) Relationship between the

means and standard deviations of mRNA expression levels for thousands of genes. Red and

blue dots represent essential and nonessential genes, respectively. The magenta and cyan lines

represent running medians and their smoothed splines, respectively. Spearman's rank

correlation coefficients (R) and p-values are shown. (**C**, **G**) Mean expression levels of essential

and nonessential genes. (**D**, **H**) Transcriptional variability, represented as either DM_{mut} or DM_{env},

of essential and nonessential genes. The DM values were calculated as the vertical distance of

792   the standard deviations from the smoothed splines of the running median of standard deviations

793   in panels **B** and **F**. The lower and upper edges of the boxes in panels **C**, **D**, **G**, **H** represent the

794   first (q1) and third (q3) quartiles, respectively. The horizontal lines in the boxes represent the

795   medians (m). The whiskers from the boxes extend to the most extreme observed values inside

796   inner fences, $m \pm 1.5(q3-q1)$.

Fig. 2: Experimental design to measure expression levels of mutant promoter library.

(**A**) Natural promoters of *E. coli* were categorized into nonessential and essential promoters,

termed Nes and Ess, respectively. Nonessential promoters control transcriptional units

comprising only nonessential genes, while essential promoters regulate transcriptional units

containing at least one essential gene. Gene essentiality was based on the requirement for cell

growth, as defined by Goodall et al(Goodall, et al. 2018). The promoter regions were obtained

from the *E. coli* Promoter Collection(Zaslaver, et al. 2006). (**B**) Synthetic promoters, termed

Syn, were derived from a random oligo DNA pool. Transformants harboring a plasmid on which

the random sequences were placed upstream of the *gfp* gene were analyzed using flow

cytometry. The 16 unique transformants, satisfying a defined green fluorescence (red shaded

area), were isolated. The obtained synthetic promoters were named SP1–16. (**C**) Natural and

synthetic promoters were placed upstream of *gfp* on a plasmid. Transformants harboring each

of these plasmids were analyzed using flow cytometry. Mean green fluorescence of each

population is shown as small dots. The large dots and error bars represent the means and

standard deviations within the groups, respectively. The number of promoters is indicated inside

the panel. The p-value corresponds to ANOVA. (**D**) Schematic showing construction of mutant

library from a given promoter region. The promoter region was subjected to random

mutagenesis by error-prone PCR. A mutant library (MT) was created through transformation to

yield more than 100,000 genetically-different mutants in a population. Unmutagenized plasmid

was also transformed to create wild type clones (WT). (**E**) Representative examples of $\log_{10}$-

transformed green fluorescence intensity distributions of WT and MT for the promoter region

of the *metG* gene in *E. coli* (NPmetG). The means (vertical lines) and standard deviations

(horizontal arrows) were calculated in each population.

Fig. 3: Mutational variability in promoter activity.

(**A**) Comparison of the standard deviations in the GFP distributions between WT and MT. The solid line represents y=x. (**B**, **C**) Relationship between the mean and standard deviation of the GFP distributions for WT (**B**) and MT (**C**). Red lines represent a regression line for WT and were used to compensate for the mean-dependency of the standard deviations of both WT and MT by subtraction. (**D**) Mutational variability in promoter activity, term MV_{prm}, is shown for each promoter group. The MV_{prm} was calculated by subtracting the compensated standard deviations of WT from those of MT. The large dots and error bars are the means and standard deviations, respectively. Pairwise Wilcoxon test was examined, where ns represents no significance. (**E**, **F**) The relationships between MV_{prm} and the DM values (DM_{env} for **E** and DM_{mut} for **F**). (**G**, **H**) Relationship between MV_{prm} and the number of known activation (**G**) and inhibition (**H**) of natural promoters by transcriptional regulators. The relationship between promoters and known transcriptional regulators were based on RegulonDB(Tierrafria, et al.

835   2022) and EcoCyc(Keseler, et al. 2021). The grey, blue, and red dots in each panel represent

836   synthetic (Syn), nonessential (Nes), and essential (Ess) promoters, respectively. Spearman's R

837   and p-values are shown in panes **A**–**C** and **E**–**H**.

Fig. 4: Biased mutational changes in activity of essential promoters.

(**A**) Comparison of the means in the GFP distributions between WT and MT. The solid line

represents y=x. (**B**) Relationship between $\log_{10}$-fold change of mean expression levels from WT

to MT and the expression levels of WT. The blue line represents a regression line used to

compensatie the mean-dependency of the $\log_{10}$-fold changes by subtraction. (**C**, **D**)

Relationship between activation bias in natural promoters and the unnormalized (**C**) and

normalized (**D**) $\log_{10}$-fold changes. The activation bias was calculated by subtracting the

number of known inhibitions from those of activations by transcriptional regulators.

Spearman's R and p-values are shown in panes **A–D**. Grey, blue, and red dots in each panel

represent synthetic (Syn), nonessential (Nes), and essential (Ess) promoters, respectively. (**E**,

F) Unnormalized (**E**) and normalized (**F**) $\log_{10}$-fold changes for each promoter group. The large

dots and error bars are the means and standard deviations, respectively. Asterisks represent

significance levels in pairwise Wilcoxon test (ns: $p > 0.05$; *: $p \leq 0.05$; **: $p \leq 0.01$). The p-

values were adjusted using the BH method(Benjamini and Hochberg 1995).

Fig. 5: Numerical simulation of promoter evolution on a ragged fitness landscape.

(**A–C**) Schematics of landscapes of expression levels (**A, B**) and fitness (**C**). (**A**) Expression

landscape was constructed by the rough Mt. Fuji (RNF) model. (**B**) Mutational change of each

genotype was calculated as the mean of differential expression levels of the nearest neighboring

genotypes (arrow heads, MT) from the focal genotype (colored circles, WT). The nearest

neighbors differ from the focal genotype by one Hamming distance. The color bar represents

the fraction of the genotypes with higher expression levels than the focal genotype among the

nearest neighbors. (**C**) Fitness landscape was modeled by adding basal fitness (F_b, red line) to

the expression landscape and normalizing it. (**D–F**) Representatives of expression landscape

(**D**), mutational change (**E**), and fitness landscape (**F**). Genotype was modeled as 10-bit binary

numbers. Hamming distance represents the distance from the genotype with the global optima

in expression level and fitness. (**G**) Mutational changes of the evolved isolates obtained through

evolutionary simulation using the standard Wright-Fisher model. The basal fitness was varied

by mimicking strong positive (F_b=0.3, left), moderate positive (F_b=0.5, middle), and nearly

neutral selection (F_b=0.99, right) acting upon higher expression levels. Ten independent

expression/fitness landscapes were generated for each condition of basal fitness. In each fitness

landscape, 600 independent evolutionary simulations were examined where initial genotypes

were randomly selected. Dots represents the most abundant mutants within the population after

30 generations in each run. Solid lines represent the running medians. Dashed lines are guides

for the eye. (**H**) Enlarged figure of the running medians in (**G**). (**I**) Expression level of the

evolved isolates. Asterisks represent significance levels of the adjusted p-value (the BH

method) in the Wilcoxon test (****: $p \leq 0.0001$).