Combined forces of hydrostatic pressure and actin polymerization drive endothelial tip cell

migration and sprouting angiogenesis

Igor Kondrychyn[1], Haymar Wint[1], Liqun He[2], Christer Betsholtz[2, 3] and Li-Kun Phng[1*]

[1] Laboratory for Vascular Morphogenesis, RIKEN Center for Biosystems Dynamics Research,

Kobe 650-0047, Japan

[2] Department of Immunology, Genetics and Pathology, Rudbeck Laboratory, Uppsala University,

75185 Uppsala, Sweden

[3] Department of Medicine Huddinge, Karolinska Institutet, 14157 Huddinge, Sweden

* Correspondence: likun.phng@riken.jp

SUMMARY

Cell migration is a key process in the shaping and formation of tissues. During sprouting

angiogenesis, endothelial tip cells invade avascular tissues by generating actomyosin-dependent

forces that drive cell migration and vascular expansion. Surprisingly, ECs can still invade if actin

polymerization is inhibited. In this study, we show that endothelial tip cells employ an alternative

mechanism of cell migration that is dependent on Aquaporin (Aqp)-mediated water inflow and

increase in hydrostatic pressure. In the zebrafish, ECs express *aqp1a.1* and *aqp8a.1* in newly

formed vascular sprouts in a VEGFR2-dependent manner. Aqp1a.1 and Aqp8a.1 loss-of-function

studies show an impairment in intersegmental vessels formation because of a decreased capacity

of tip cells to increase their cytoplasmic volume and generate membrane protrusions, leading to

delayed tip cell emergence from the dorsal aorta and slower migration. Further inhibition of actin

polymerization resulted in a greater decrease in sprouting angiogenesis, indicating that ECs

employ two mechanisms for robust cell migration *in vivo*. Our study highlights an important role

of hydrostatic pressure in tissue morphogenesis.

Keywords

Endothelial cell migration, Sprouting angiogenesis, Aquaporin, Water flow, Hydostatic Pressure,

Actin, Zebrafish

INTRODUCTION

The blood vascular system is a dynamic, multicellular tissue that adapts to the metabolic demands

of development, growth and homeostasis by altering its pattern and morphology. Macroscopic

transformation in the vascular network is achieved by microscopic changes in endothelial cell (EC)

behaviours. For example, vascular expansion through sprouting angiogenesis requires the

coordination of EC migration, proliferation, anastomosis, lumen formation and cell

rearrangements, all of which require specific cell shape changes. At the initial phase of sprouting

angiogenesis, endothelial tip cells generate numerous membrane protrusions such as filopodia and

lamellipodia that drive migration and anastomosis (Figueiredo et al., 2021; Gerhardt et al., 2003;

Phng et al., 2013; Sauteur et al., 2017). During tubulogenesis, apical membranes deform by

generating inverse blebs to expand lumens so that blood can be transported efficiently through the

blood vascular network (Gebala et al., 2016). However, following vessel perfusion, ECs become

less dynamic and instead develop resistance to the deforming forces of blood pressure to maintain

vessel morphology (Kondrychyn et al., 2020). EC shape is therefore dynamic and changes

depending on the morphogenetic state of vascular development.

It is well established that localized remodelling of actin cytoskeleton and non-myosin II

contractility are instrumental in driving cell shape changes during tissue morphogenesis (Clarke

and Martin, 2021; Munjal and Lecuit, 2014; Murrell et al., 2015). More recently, there is an

accumulating body of work implicating a role of water flow and the resulting changes in

hydrostatic pressure as another mechanical mechanism of cell shape change (Choudhury et al.,

2022; Chugh et al., 2022; Li et al., 2020). For example, in the osmotic engine model of cell

migration, water inflow at the leading edge and outflow at the rear leads to forward translocation

of tumour cells in confined microenvironment (Stroka et al., 2014). Additionally, hydrostatic

pressure from the extracellular environment can drive tissue morphogenesis, as demonstrated

during mouse blastocyst formation (Chan et al., 2019; Dumortier et al., 2019), development of the

otic vesicle in the zebrafish (Mosaliganti et al., 2019) and blood vessel lumen expansion (Gebala

et al., 2016). Hydrostatic pressure can therefore generate sufficient forces to shape tissues and

organs to their proper form and size during development.

We have previously discovered that EC migration persists after the inhibition of actin

polymerization and in the absence of filopodia to generate new blood vessels in the zebrafish (Phng

et al., 2013). This observation suggests the existence of an alternative mechanism of migration

independent of actin polymerization. In this study, we sought to determine whether hydrostatic

pressure regulates EC migration during sprouting angiogenesis in the zebrafish by investigating

the function of Aquaporins (Aqp), which are transmembrane water channels that increase water

permeability of cell membranes to promote transcellular water flow (Day et al., 2014; Farinas et

al., 1997; Kozono et al., 2002; Preston et al., 1992). We discovered that ECs of newly formed

intersegmental vessels (ISV) express *aqp1a.1* and *aqp8a.1* mRNA, and observed the enrichment

of Aqp1a.1 and Aqp8a.1 proteins in the leading edge of migrating tip cells. Detailed single cell

analyses showed that endothelial tip cells lacking *aqp1a.1* and *aqp8a.1* expression have reduced

cell volume, membrane protrusions and migration capacity. As a result, there is defective sprouting

angiogenesis and ISV formation in *aqp1a.1;aqp8a.1* double mutant zebrafish. Notably, when actin

polymerization is inhibited in *aqp1a.1*;*aqp8a.1* double mutant zebrafish, there is a greater

impairment in EC migration and sprouting angiogenesis, demonstrating the additive function of

actin polymerization and hydrostatic pressure (that is created by water inflow) in generating

membrane protrusions to drive EC migration *in vivo*.

RESULTS

***aqp1a.1* and *aqp8a.1* are differentially expressed by tip and stalk cells during sprouting**

angiogenesis

Of the 18 number of *aquaporin* genes that are expressed in zebrafish (Tingaud-Sequeira et al.,

2010), *aqp1a.1* and *aqp8a.1* mRNA have been detected in blood endothelial cells (ECs) (Koun et

al., 2016; Rehn et al., 2011). We confirmed the endothelial expression of *aqp1a.1* and *aqp8a.1* at

30 hours post fertilization (hpf) and 2-, 3- and 4-days post fertilization (dpf) by whole-mount *in*

situ hybridization (Fig. S1). The two endothelial *aqp* genes, however, differ in spatial expression.

While *aqp1a.1* is widely expressed in blood vessels of the head and trunk (dorsal aorta (DA),

caudal artery (CA), ISVs, dorsal longitudinal anastomotic vessels (DLAV) and caudal vein plexus)

at 30 hpf, *aqp8a.1* expression is absent in cerebral blood vessels and is restricted to the DA, CA

and the ventral regions of ISVs in the trunk. After 1 dpf, *aqp8a.1* expression expands to the entire

ISV and DLAV. Additionally, the expression of *aqp1a.1* and *aqp8a.1* gradually decreases in the

DA and CA so that both are absent by 4 dpf. Single cell RNA sequencing (scRNAseq) of ECs

isolated at 34 hpf and 3 dpf further confirmed the endothelial expression of *aqp1a.1* and *aqp8a.1*

mRNA (Fig. S2 A - D), revealed differential expression of *aqp1a.1* and *aqp8a.1* mRNA in distinct

endothelial clusters and highlighted higher *aqp1a.1* transcript expression in all endothelial

subtypes compared to *aqp8a.1* (Fig. S2 E - H).

We next examined the expression pattern of *aqp1a.1* and *aqp8a.1* at the beginning of

sprouting angiogenesis at higher resolution using RNAscope and confocal microscopy. At 20 hpf,

aqp1a.1 mRNA is highly expressed in the DA with little expression in the PCV (Fig. 1A, C and

Fig. S3). Closer inspection uncovered heterogeneous expression in the DA, with higher expression

in ECs that would be specified as tip cells (Fig. 1A, C and Fig. S3B). Indeed, tip cells of newly

formed vascular sprouts express higher levels of *aqp1a.1* compared to adjacent ECs in the DA at

20 hpf (Fig. 1A – C, Fig. S3B) and the majority of ISVs (61%) display higher *aqp1a.1* expression

in tip cells than in trailing stalk cells at 22 hpf (Fig. 1G). Although we previously observed *aqp8a.1*

mRNA in the DA at 30 hpf (Fig. S1), its expression here is largely absent at 20 hpf (Fig. 1B, D

and Fig. S3C). At 22hpf, *aqp8a.1* mRNA expression in the DA increases (Fig. 1E and F).

Interestingly, unlike *aqp1a.1*, in 78% of ISVs analyzed, *aqp8a.1* mRNA is more highly expressed

in stalk cells than tip cells (Fig. 1G). These observations highlight differential expression pattern

of *aqp1a.1* and *aqp8a.1* in newly formed vascular sprouts, with *aqp1a.1* expression enriched in tip

cells and *aqp8a.1* expression in stalk cells.

VEGF-VEGFR2 signalling induces the expression of *aqp1a.1* and *aqp8a.1* mRNA

As *aqp1a.1* and *aqp8a.1* mRNA display differential tip-stalk cell expression, we explored whether

their expression is regulated by the VEGFR2-Notch signalling axis, which controls endothelial tip-

stalk cell specification during sprouting angiogenesis (Benedito and Hellström, 2013; Phng and

Gerhardt, 2009; Siekmann et al., 2013). To inhibit VEGFR2 activity, embryos were treated with 1

µM ki8751 from 20 hpf. Relative quantitative polymerase chain reaction (qPCR) experiments

showed that *aqp1a.1* mRNA expression was suppressed by 52% after 6 hours ($p < 0.0001$) and

by 40% after 24 hours ($p < 0.0001$) of ki8751 treatment when compared to DMSO-treated embryos

(Fig. 1H). *aqp8a.1* mRNA expression was also decreased by the inhibition of VEGFR2 activity,

and its suppression was accentuated after 24 hours treatment (93% decrease, $p < 0.0001$) compared

to after 6 hours (17% decrease, $p = 0.0028$). These results demonstrate that VEGFR2 activity

upregulates the expression of *aqp1a.1* and *aqp8a.1*, but the temporal dynamics is different for

aqp1a.1 and *aqp8a.1*. To examine whether VEGFR2 activity also regulates Aquaporin expression

in human ECs, we treated cultured human aortic endothelial cells (HAECs) with ki8751 for 6 hours and analyzed *AQP1* expression by qPCR. *AQP8* expression was not examined since it is not expressed in HAECs (data not shown). Inhibition of VEGFR2 activity with 1 nM and 10 nM ki8751 decreased *AQP1* mRNA expression by 31% ($p = 0.0023$, Fig. 1I) and 40% ($p = 0.1032$, Fig. 1I), respectively, demonstrating cross-species conservation of VEGFR2 activity in inducing *Aqp1* mRNA expression.

Notch signalling was inhibited using the gamma-secretase inhibitor, Dibenzazepine (DBZ), which prevents proteolytic cleavage and consequently, activation of the Notch receptor (Groth et al., 2010). Treatment of embryos with DBZ from 20 hpf for 6 and 24 hours significantly decreased the expression of *hey1* and *hey2,* canonical targets of Notch signalling (Fig. 1J), confirming the effectiveness of the drug treatment in our experiments. When compared to DMSO-treated embryos, 6-hour treatment with DBZ did not cause a significant change in *aqp1a.1* and *aqp8a.1* expression levels (Fig. 1J). Longer treatment with DBZ for 24 hours, however, increased *aqp8a.1* expression by 98% ($p < 0.0001$) while *aqp1a.1* expression was relatively unchanged (Fig. 1J). These findings suggest that Notch signalling inhibits *aqp8a.1*, but not *aqp1.a*, expression.

Aqp1a.1 and Aqp8a.1 proteins are enriched at the leading front of migrating tip cells

To gain a better understanding of where within the cell Aquaporin protein functions, we generated stable *Tg(fli1ep:aqp1a.1-mEmerald)^rk30* and *Tg(fli1ep:aqp8a.1-mEmerald)^rk31* zebrafish lines in which ECs express Aqp1a.1 or Aqp8a.1 tagged with mEmerald, respectively. At 25 hpf, we observed an accumulation of Aqp1a.1 (Fig. 2A) and Aqp8a.1 (Fig. 2B) in mCherryCAAX-positive membrane clusters in the cytoplasm and filopodia of tip cells. Interestingly, many of the Aqp1a.1- or Aqp8a.1-positive membrane clusters are located at the migrating front of the tip cell. Time-lapse imaging capturing the process of ISV formation between 24 to 30 hpf further showed sustained enrichment of Aqp1a.1 and Aqp8a.1 in clusters at the front of the tip cell as well as in filopodia (Fig. 2C and D, Movie S1 and S2), suggesting that Aqp1a.1 and Aqp8a.1 may promote localized water flux at the migrating edge of tip cells.

Loss of Aqp1a.1 and Aqp8a.1 function leads to defective trunk blood vessel formation

We subsequently proceeded to generate zebrafish *aqp1a.1* and *aqp8a.1* mutants to investigate the function of Aquaporin-mediated water flow in blood vessel development. Using the CRISPR/Cas9

genome editing technique, we targeted the first exons of *aqp1a.1* and *aqp8a.1* genes. The CRISPR-generating allele *aqp1a.1^rk28^* carries a 14-bp deletion and an 8-bp insertion in the 5' end of the gene leading to a premature termination codon at amino acid 76 after 1 missense amino acid (Fig. S4A-B). The *aqp8a.1^rk29^* allele carries a 4-bp deletion in the 5' end of the gene that leads to a frameshift after T15 and premature termination codon at amino acid 73 after 58 missense amino acids (Fig. S5A-B). Premature termination codons are often more deleterious than missense mutations because they result in the loss of protein expression. Nonsense-mediated mRNA decay is a quality-control mechanism that selectively degrades mRNAs harboring premature termination codons (Chang et al., 2007). Using whole-mount *in situ* hybridization and qPCR analysis, we demonstrated that *aqp1a.1* and *aqp8a.1* mRNA expression is significantly decreased in *aqp1a.1^rk28/rk28^* (Fig. S4C-D) and *aqp8a.1^rk29/rk29^* (Fig. S5C-D) zebrafish embryos, respectively, suggesting a rapid degradation of mutant mRNAs and as a result, the loss of Aquaporin protein expression.

Although *aqp1a.1^rk28/rk28^;aqp8a.1^rk29/rk29^* embryos look morphologically normal when compared to wildtype or *aqp1a.1^+/rk28^;aqp8a.1^+/rk29^* embryos (Fig. S8A, B, C and D), we observed defects in blood vessel formation and patterning at 1 to 3 dpf. Compared to 28 hpf control embryos whose ISVs have reached the dorsal roof of the neural tube and started to form the DLAV, ISVs of *aqp1a.1^rk28/rk28^;aqp8a.1^rk29/rk29^* embryos are much less developed (Fig. 3A and D) with significantly reduced length (Fig. 3E and Fig. S6H and K). Single deletion of *aqp1a.1* (Fig. 3B and Fig. S6F and I) or *aqp8a.1* (Fig. 3C and Fig. S6G and J) also led to defective ISV formation, although the defects are less severe compared to the loss of both *aqp1a.1* and *aqp8a.1* expression. Comparison of ISV length in *aqp1a.1^rk28/rk28^*, *aqp8a.1^rk29/rk29^* and *aqp1a.1^rk28/rk28^;aqp8a.1^rk29/rk29^* embryos shows that the loss of *aqp1a.1* leads to a greater decrease in ISV length than the loss of *aqp8a.1* (Fig. 3E, Fig. S6I and J), and that the loss of both *aqp1a.1* and *aqp8a.1* resulted in the greatest reduction. These observations indicate an additive effect of Aqp1a.1 and Aqp8a.1 function in blood vessel development and implicate Aqp1a.1 as the more dominant Aquaporin protein in EC function, in line with the higher expression of *aqp1a.1* mRNA in tip cells.

We next examined the embryos at 3 dpf, a stage when arterial ISVs (aISVs) and venous (vISVs) are fully established and lumenized, connecting the DLAV to either the DA or PCV (Fig. 3F and G), respectively. Such fully established ISVs are significantly reduced in embryos when *aqp1a.1*, *aqp8a.1* or both *aqp1a.1* and *aqp8a.1* are deleted. Instead, these embryos display an

increased number of aISVs and vISVs that fail to establish a connection to the DLAV, DA or PCV

(Fig. 3H – J). Quantification shows that while 19.1% of wildtype embryos display at least 1

incomplete ISV per embryo, this increased to 65.8% in *aqp8a.1$^{rk29/rk29}$* embryos, 83.3% in

aqp1a.1$^{rk28/rk28}$ embryos and 92.2% in *aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$* embryos (Fig. 3K). The

number of incomplete ISVs per embryo in *aquaporin* mutants was variable, between 1 to 6, 1 to

16 and 1 to 22 incomplete ISVs in *aqp8a.1$^{rk29/rk29}$*, *aqp1a.1$^{rk28/rk28}$* and

aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$ embryos, respectively (Fig. S6A-D). We also found that incomplete

ISVs appear with higher frequency in the zebrafish trunk (ISV number 6 to 15) rather than in the

tail (ISV number 16 to 27, Fig. S6E). Detailed analysis revealed that most of the incomplete ISVs

are vISVs that are connected to the PCV but not the DLAV (Class I, 75.2%), followed by the

absence of aISV or vISV (Class II, 15.8%, Fig. 3L and Fig. S7A and B). A small fraction of the

incompletely formed vessels are aISVs that are not connected to the DA (Class IV, 2.2%, Fig.

S7D) or DLAV (Class V, 2.9%, Fig. S7E), or vISVs that are not connected to the DLAV (Class

VII, 2.68%, Fig. S7F). At some regions, there is an aISV that is not connected to the DA and a

vISV that is not connected to the DLAV (Class III, 1.2%, Fig. S7C).

As *aqp1a.1* is expressed in brain vasculature during development (Fig. S1), we examined

whether cerebral vasculature is also affected in *aqp1a.1$^{rk28/rk28}$* embryos. At 4 dpf, we found that

the length of the dorsal longitudinal vein (DLV), a main blood vessel that supplies blood to the

choroid plexus (Bill and Korzh, 2014) was reduced in *aqp1a.1$^{rk28/rk28}$* embryos ($p = 0.0098$, Fig.

S9B). Also, in 2 out of 21 *aqp1a.1$^{rk28/rk28}$* embryos, the DLV was undeveloped (Fig. S9A and C),

a defect that is also observed in 50% of *vegfab$^{-/-}$* and *vegfc$^{-/-}$* mutant zebrafish (Parab et al., 2021).

In summary, our analyses suggest that in the absence of Aqp1a.1 and/or Aqp8a.1 function,

there is an initial delay in ISV formation during primary sprouting angiogenesis. By 3 dpf, some

ISVs are completely formed (Fig. S8K and M), connecting the DA or PCV to the DLAV. However,

the number of completely formed ISVs is significantly decreased, with the loss of Aqp1a.1 having

a more severe impact compared to the loss of Aqp8a.1, and the combined loss of Aqp1a.1 or

Aqp8a.1 having a greater deleterious effect.

Aquaporins promote endothelial tip cell protrusion and migration

To gain a better understanding of how incomplete ISV formation arise, we performed timelapse

imaging in zebrafish depleted of both Aqp1a.1 and Aqp8a.1 function from 21 – 22hpf, when

sprouting angiogenesis begins. In *aqp1a.1$^{+/rk28}$;aqp8a.1$^{+/rk29}$* embryos, tip cells emerge from the

DA between 20 and 24 hpf (Fig. 4A) and migrate dorsally between the somite boundaries over the

notochord and neural tube to form primary ISVs (Fig. 4B, Movie S3). However, in

aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$ embryos, tip cell nuclei emerge between 22 and 28 hpf (Fig. 4A).

Furthermore, in some *aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$* embryos, tip cells retract whereby membrane

protrusions do not extend dorsally in the direction of migration but withdraw into the DA so that

a primary ISV does not form at this location (Fig. 4C, Movie S4). These observations indicate that

tip cell protrusion from the DA is significantly delayed in the absence of Aqp1a.1 and Aqp8a.1

function.

We have previously demonstrated that filopodia facilitate EC migration by serving as a

template for lamellipodia formation (Phng et al., 2013). Lamellipodia frequently emanate laterally

from stable filopodia at the leading edge of tip cells, leading to rapid expansion of the filopodium,

increase in volume and the formation of a stable protrusion in the direction of migration (Phng et

al., 2013). This was similarly observed in *aqp1a.1$^{+/rk28}$;aqp8a.1$^{+/rk29}$* embryos (Fig. 4D, Movie S5)

but not in *aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$* embryos (Fig. 4E, Movie S6). Comparison of filopodia

dynamics in wildtype (Fig. 4F), *aqp1a.1$^{+/rk28}$;aqp8a.1$^{+/rk29}$* and *aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$*

(Fig. 4G) embryos revealed a significant reduction in filopodia number (wildtype, 18.75 ±

5.1/100µm; *aqp1a.1$^{+/rk28}$;aqp8a.1$^{+/rk29}$*, 19.8 ± 6.3/100µm; *aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$*, 9.9 ±

5.2/100µm; Fig. 4H) and length (wildtype, 5.9 ± 3.2 µm; *aqp1a.1$^{+/rk28}$;aqp8a.1$^{+/rk29}$*, 6.2 ± 3.3 µm;

aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$, 4.8 ± 2.4 µm; Fig. 4I) in ECs lacking both Aqp1a.1 and Aqp8a.1.

function. Analysis of time-lapse images of tip cell behaviour further showed slower tip cell

membrane expansion at the leading edge (Fig. 4J), decreased nuclei displacement (Fig. 4K) and a

significant reduction in tip cell migration velocity (Fig. 4L) in *aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$*

embryos.

In conclusion, Aqp1a.1 and Aqp8a.1 are required for sprouting angiogenesis by promoting

endothelial tip cell emergence from the DA, the formation of membrane protrusions and

endothelial cell migration.

Tip cell volume regulation depends on Aquaporin-mediated water influx

Aquaporin channels permit bidirectional flow of water across the plasma membrane, with the

direction of flow determined by the osmotic gradient between the extracellular environment and

cell cytoplasm (Agre et al., 2002). We next sought to determine whether water flows in or out of

tip cells during sprouting angiogenesis. We hypothesize that water flux across the cell membrane

can lead to changes in tip cell volume, with water influx increasing the volume while outflow

decreases. To quantify tip cell volume, we injected a plasmid encoding *kdrl:mEmerald* transgene

into 1-cell stage wildtype or *aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$* embryos to label ECs in a mosaic

manner. mEmerald-positive tip cells were imaged and their volume measured at 24 to 25 hpf.

Wildtype ECs are on average 946.6 ± 299.8 µm^3 in size (Fig. 5A and C). In the absence of both

Aqp1a.1 and Aqp8a.1 function (Fig. 5B), tip cell volume significantly decreased by about 24% to

718.4 ± 308.5 µm^3 ($p = 0.0176$, Fig. 5C) when compared to wildtype tip cells. This finding

supports water influx as the direction of water flow in tip cells during migration, and that this is

mediated through Aqp1a.1 and Aqp8a.1.

It has been proposed that water influx through Aquaporin increases the diffusion of actin

monomers and increases actin polymerisation (Loitto et al., 2007; Papadopoulos et al., 2008) .

Given that filopodia formation is dependent on actin polymerization (Mallavarapu and Mitchison,

1999; Ridley, 2011), we examined whether actin polymerization is perturbed in the absence of

Aqp1a.1 and Aqp8a.1 by quantifying the growth rate of actin bundles in tip cell filopodia by

tracking Lifeact-mCherry signal in the filopodia. In wildtype embryos, the average growth rate of

actin is 0.054 ± 0.015 µm per second whereas in *aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$* embryos, the

growth rate is significant decreased by 47% to 0.028 ± 0.010 µm per second ($p < 0.0001$, Fig. 5D).

Our observations therefore support a role of zebrafish Aqp1a.1 and Aqp8a.1 in promoting actin

polymerisation and consequently, filopodia number and length, which are reduced in

aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$ embryos (Fig. 4F and G).

In summary, our results demonstrate a role of endothelial Aquaporins in increasing water

influx and as a consequence, endothelial tip cell volume and hydrostatic pressure, as well as in

promoting actin polymerization.

Additive effects of actin polymerization and water influx in driving EC migration and

sprouting angiogenesis

We have previously observed that ECs are still able to migrate to generate ISVs after the inhibition

of actin polymerization (Phng et al., 2013). In these experiments, embryos were treated with a low

concentration of Latrunculin B (Lat. B), an inhibitor of actin polymerization, that resulted in the

suppression of filopodia formation. Under such conditions, ECs can generate small membrane
protrusions and migrate more slowly in a directed manner to generate ISVs. How ECs are still able
to migrate after the inhibition of actin polymerization was unclear. As our current study
demonstrates a role of water inflow and hydrostatic pressure as another mechanism of cell
migration, we hypothesize that ECs employ water influx to migrate when actin polymerization is
compromised and that the depletion of both water influx and actin polymerization will result in a
greater inhibition of EC migration.

To test this hypothesis, we examined ISV development in embryos with decreased actin
polymerization and water inflow by treating $aqp1a.1^{rk28/rk28};aqp8a.1^{rk29/rk29}$ embryos with Lat. B.
At 28 hpf, ISVs of wildtype embryos treated with DMSO have reached the dorsal roof of the neural
tube (Fig. 6A) and show an average length of 104.1 ± 13.47 μm (Fig. 6E). The treatment of
wildtype embryos with 0.08 μg/ml Lat. B from 20 to 28 hpf (Fig. 6B) resulted in a significant
decrease in ISV length (61.53 ± 23.27 μm) compared to control embryos ($p < 0.0001$, Fig. 6E). A
similar decrease in ISV length was also observed in $aqp1a.1^{rk28/rk28};aqp8a.1^{rk29/rk29}$ embryos treated
with DMSO (63.75 ± 30.24 μm, Fig. 6C and E). When $aqp1a.1^{rk28/rk28};aqp8a.1^{rk29/rk29}$ embryos
were treated with 0.08μg/ml Lat. B from 20 to 28 hpf, there was a greater decrease in the length
of ISV formed at 28 hpf (43.10 ± 24.42 μm, Fig. 6D and E) when compared to the inhibition of
actin polymerization ($p = 0.0028$) or depletion of water influx ($p < 0.0001$) alone (Movie S7).
These results therefore demonstrate additive functions of actin polymerization and water influx in
driving EC migration during sprouting angiogenesis (Fig. 6F).

DISCUSSION
In this study, we demonstrate that endothelial tip cells utilize two mechanisms of migration
concurrently during sprouting angiogenesis in the zebrafish - actin polymerization and hydrostatic
pressure. The increase in hydrostatic pressure in endothelial tip cells is generated by Aqp1a.1- and
Aqp8a.1-mediated water inflow. By performing detailed expression analyses, we showed that
aqp1a.1 and *aqp8a.1* are differentially expressed in newly formed vascular sprouts. *Aqp1a.1* is
expressed earlier in the DA than *aqp8a.1* during development and in ECs destined to be tip cells.
In newly formed ISVs, endothelial tip cells express higher *aqp1a.1* mRNA levels while stalk cells
express higher *aqp8a.1* mRNA levels. In zebrafish harbouring mutations in both *aqp1a.1* and
aqp8a.1, there is delayed tip cell protrusion from the DA, decreased tip cell migration and

increased occurrence of truncated ISVs at 3 dpf. We further discovered that *aqp1a.1* and *aqp8a.1*

expression is upregulated by VEGFR2 activity, proposing that VEGFA-VEGFR2 signalling

employs Aquaporin function to drive sprouting angiogenesis.

Several lines of evidence implicate Aquaporins in facilitating water flow into endothelial

tip cells during sprouting angiogenesis. First, timelapse imaging showed enriched localization of

Aqp1a.1 and Aqp8a.1 proteins at the migrating front of tip cells, suggesting that water flow occurs

locally at the leading edge of the cell. Secondly, single cell analyses revealed Aqp1a.1- and

Aqp8a.1-deficient tip cells have decreased cell volume, impaired expansion of cell membranes at

the leading edge and decreased generation of stable cell protrusions. These observations implicate

water flow into tip cells, leading to increased cell volume. Due to the incompressible property of

water within an enclosed compartment such as the cell, water influx will lead to a rise in hydrostatic

pressure. At the leading edge of tip cells, increased hydrostatic pressure would expand and deform

membranes to generate protrusions. Additionally, water influx can reduce cytoplasmic viscosity

and increase spacing between the plasma membrane and the underlying actin cytoskeleton,

promoting actin monomer diffusion, actin polymerization and the generation of membrane

protrusions to facilitate cell migration (Boer et al., 2023; Loitto et al., 2009).

A key finding in this study is the demonstration that endothelial tip cells employ hydrostatic

pressure as a second mechanism of cell migration *in vivo*. In a previous study, we discovered that

ECs continue to migrate when actin polymerization is inhibited and in the absence of filopodia

(Phng et al., 2013). Here, we show that hydrostatic pressure is the driving force for EC migration

in the absence of actin-based force generation. When both mechanisms are perturbed, there is a

greater impairment in EC migration (Fig. 6D and E). This finding highlights the importance of

hydrostatic pressure as an additional mechanism that reinforces actin-based cell shape changes and

motility to ensure robust migration and formation of new blood vessels in complex three-

dimensional environments. Such dual mode migration is advantageous when blood vessels

develop in physically confined spaces such as in the zebrafish trunk, where ECs must emerge from

the DA and migrate within somite boundaries and over tissues such as the notochord and neural

tube to form ISVs. Such dependence on water flow in cell migration under confined

microenvironment has been reported in human MDA-MB-231 breast cancer cells and mouse S180

sarcoma cells, which show cell migration persistence in narrow channels after inhibition of actin

polymerization or myosin II-mediated contractility (Stroka et al., 2014). In this context, AQP5

mediates water flow and the decrease or increase in AQP5 expression suppresses or enhances, respectively, cancer cell migration (Chae et al., 2008; Jung et al., 2011; Stroka et al., 2014).

Our findings on the endothelial function of zebrafish Aqp1a.1, which is homologous to mammalian AQP1, corroborate previous studies demonstrating a function of AQP1 in promoting cell migration. In tumour bearing AQP1-null mice, there is reduced tumour angiogenesis and AQP1-deficient ECs display reduced migration in an *in vitro* wound healing assay (Maltaneri et al., 2020; Saadoun et al., 2005). In chick, AQP1 expression levels affect neural crest cell migration speed and direction by regulating filopodia length and stability (McLennan et al., 2019). The regulation of membrane protrusions has also been demonstrated for other Aquaporin proteins. For example, AQP9 weakens membrane-cytoskeleton anchorage and promote formation of membrane protrusions such as filopodia and blebs (Karlsson et al., 2013; Loitto et al., 2007). We also demonstrate for the first time a role of Aqp8a.1 in promoting EC migration and sprouting angiogenesis in the zebrafish. However, its mammalian ortholog, AQP8, was shown to inhibit migration of colorectal cancer cell lines *in vitro* (Wu et al., 2017), suggesting cell- and context-dependent function of AQP8 in the regulation of cell migration.

A central determinant of water movement is the osmotic gradient across the membrane, with water flowing in the direction of higher solute concentration. Given a crucial role of water flow in regulating cell volume, shape and migration, it is also important to understand the function and distribution of ion channels, exchangers and transporters and water channels that generate an osmotic gradient across the cell. This is reflected in a growing number of studies demonstrating a role of ion transporters in cell migration. In neutrophils, the increase in cell volume and potentiation of cell migration depend on the sodium-proton exchanger 1 (NHE1) and the chloride-bicarbonate exchanger 2 (AE2) (Nagy et al., 2023) . In T cells, chemokine-induced migration depends on ion influx via SLC12A1, and water influx via AQP3 (Boer et al., 2023). In tumour cells migrating in narrow channels, the establishment of a polarized distribution of Na^+/H^+ pumps and Aquaporin proteins in cell membranes is required for net inflow of ions and water at the leading edge and net outflow at the trailing edge (Stroka et al., 2014). We have also observed enrichment of Aqp1a.1 and Aqp8a.1 proteins in ECs, at the leading edge of migrating endothelial tip cells as well as apical membranes during lumen formation (not shown). This leads to the question of how Aquaporin localisation is regulated as this will determine the site of increased water flow and hydrostatic pressure. The rapid changes in subcellular localization of mammalian

Aquaporins upon stimulation occurs mainly via trafficking to and from the plasma membrane (Markou et al., 2022), and can lead to changes in the amount of Aquaporin protein in the plasma membrane. The regulation of AQP subcellular relocalization via calmodulin- and/or phosphorylation-dependent mechanisms has been implicated for AQP0-5 and AQP7-9 (Markou et al., 2022). For example, rapid translocation of AQP1 to plasma membrane upon a hypotonic stimulus is dependent on calmodulin activation and phosphorylation by protein kinases C (PKC) (Conner et al., 2012). Phosphorylation of AQP2 by protein kinase A (PKA) is required for vasopressin-stimulated relocalization of AQP2 from storage vesicles to the plasma membrane (Noda and Sasaki, 2005). Subcellular localization of AQP8 in hepatocytes is also regulated by PKA and PI3K signalling (Gradilone et al., 2005, 2003). In mouse neutrophils, exposure to a chemotactic gradient caused translocation of AQP9 to the leading edge of the cell that is mediated by PKC-dependent phosphorylation (Karlsson et al., 2011). Further work is needed to elucidate the mechanism of Aqp1a.1 and Aqp8a.1 protein distribution in ECs and which ion transporters are expressed in ECs to regulate the ion movement and cell migration.

In this study, we have assumed that the function of Aqp1a.1 and Aqp8a.1 is a result of water permeation based on previous studies demonstrating water molecules flowing through mammalian AQP1 (Preston et al., 1992; Zeidel et al., 1992) and AQP8 (Liu et al., 2006; Ma et al., 1997) as well as through their zebrafish orthologs Aqp1a.1 and Aqp8a.1 (Tingaud-Sequeira et al., 2010). However, AQP1 and AQP8 are not water-specific channels. AQP1 can conduct unpolar gases carbon dioxide (Nakhoul et al., 1998; Prasad et al., 1998; Wang et al., 2007) and nitric oxide (Herrera et al., 2006; Herrera and Garvin, 2007), and in zebrafish, Aqp1a.1 also mediates the transfer for small gaseous molecules such as carbon dioxide and ammonia (Talbot et al., 2015). AQP8 is permeable to ammonia (Jahn et al., 2004; Liu et al., 2006) and is present in the inner mitochondrial membrane of different cells (Calamita et al., 2005), where it is suggested to mediate ammonia transport rather than water fluxes (Soria et al., 2010). Among the three Aqp8 paralogs in teleost, Aqp8a.1 and Aqp8a.2 are also permeable to urea (Tingaud-Sequeira et al., 2010) and Aqp8b was shown to act mainly as a mitochondrial H_2O_2 channel (peroxiporin) in activated marine and freshwater teleost spermatozoa (Chauvigné et al., 2021, 2015). Nevertheless, because water movement flows up an osmotic gradient, increased entry of ions into the cell through Aquaporins will also trigger water inflow.

In summary, our study highlights the role of water influx and hydrostatic pressure as
another force-generating mechanism utilized by cells to build tissues during animal development.
We demonstrate that endothelial tip cells employ both actin polymerization and hydrostatic
pressure for robust sprouting angiogenesis. As morphogenetic events are governed by a
coordination of cell shape changes, cell migration and rearrangements, we envision that the
shaping and formation of other tissues will also depend on Aquaporin function and changes in
hydrostatic pressure.
410

METHODS
Zebrafish maintenance and stocks.
Zebrafish (*Danio rerio*) were raised and staged according to established protocols (Kimmel et al.,
1995). Transgenic lines used in this work are *Tg(kdrl:EGFP)^{s843}* (Jin et al., 2005) ,*Tg(fli1a:H2B-*
EGFP)^{ncv69} [re-named in ZFIN as *Tg(fli1:zgc:114046-EGFP)*] (Ando et al., 2019), *Tg(fli1:Lifeact-*
mCherry)^{ncv7} (Wakayama et al., 2015), *Tg(fli1:myr-mCherry)^{ncv1}* (Fukuhara et al., 2014),
Tg(fli1ep:EGFP-PLC1δPH)^{rk26} (Kondrychyn et al., 2020), *Tg(fli1:h2bc1-mCherry)^{ncv31}* (Nakajima
et al., 2017), *Tg(kdrl:ras-mCherry)^{s916}* (Hogan et al., 2009). Zebrafish were maintained on a 12-
hour light/12-hour dark cycle, and fertilized eggs were collected and raised in E3 medium at 28^{0}C.
To inhibit pigmentation in embryos older than 24 hpf, 0.003% *N*-Phenylthiourea (Sigma-Aldrich)
in E3 medium was used. All animal experiments were approved by the Institutional Animal Care
and Use Committee at RIKEN Kobe Branch (IACUC).
423

Plasmids and oligonucleotides.
All DNA constructs used in this study were generated by using In-Fusion HD Cloning kit (Takara
Bio Inc.). Plasmid *mEmerald-N1* was a gift from Michael Davidson (Addgene plasmid #53976).
Plasmids *pDsChLexOG* (Emelyanov and Parinov, 2008) containing terminal *cis*-required
sequences of the *Ds* transposon from maize was a generous gift from Sergei Parinov. A detailed
information regarding plasmids and primers used in this study can be found in Supplementary
Tables 1 and 2, respectively.
431

Cloning of *aqp1a.1* and *aqp8a.1* genes. Total RNA was isolated from 1-day old zebrafish
embryos with TRI Reagent using Direct-zol^{TM} RNA MicroPrep kit (Zymo Research) according to

the manufacturer's protocol. The first-strand cDNA was synthesized from 1 µg of a total RNA by

oligo(dT) priming using the SuperScript III First-Strand synthesis system (Invitrogen) according

to the manufacturer's protocol. Amplification of cDNAs was performed using a high-fidelity

KOD-Plus-Neo DNA polymerase (Toyobo, Japan) and resulting PCR products were cloned using

NEB® PCR Cloning kit (New England BioLabs). Positive clones and plasmids were verified by

DNA sequencing.

Mosaic expression of DNA constructs and transgenic lines generation.

To establish *Tg(fli1ep:aqp1a.1-mEmerald)^{rk30}* and *Tg(fli1ep:aqp8a.1-mEmerald)^{rk31}* transgenic

lines, *Tol2*-based plasmids (10 ng/µl) were co-injected into one-cell stage zebrafish embryos with

Tol2 transposase mRNA (100 ng/µl) transcribed from Not*I*-linearized *pCS-TP* plasmid (a gift from

Koichi Kawakami, National Institute of Genetics, Japan) using the mMESSAGE mMACHINE

SP6 kit (Invitrogen). Injected embryos were raised to adults and screened for germline

transmission to establish a stable transgenic line. For mosaic expression of DNA construct, *Ds*-

based *kdrl:mEmerald* plasmid (10 ng/µl) was co-injected into one-cell stage zebrafish embryos

with *Ac* transposase mRNA (100 ng/µl) transcribed from Bam*HI*-linearized *pAc-SP6* plasmid

(Emelyanov et al., 2006) using the mMESSAGE mMACHINE SP6 kit (Invitrogen). Embryos

were analyzed at 25-26 hpf after injections.

***In vivo* cell volume analysis.**

Single-cell labelling of ECs in ISVs was achieved by mosaic expression of *pDs-kdrl:mEmerald*

plasmid in wild type or *aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$* mutant *Tg(fli1:Lifeact-mCherry)^{ncv7}*

transgenic embryos. Embryos were imaged at 25-26 hpf with an Olympus UPLSAPO 60x/NA 1.2

water immersion objective with optical Z planes interval of 0.26 µm. Cell volume was measured

with Huygens Essential 22.10 software (Scientific Volume Imaging B.V.) using object analyzer

tool with manual adjustment of threshold.

Generation of zebrafish *aqp1a.1* and *aqp8a.1* mutants.

Aquaporins mutants were generated by CRISPR/Cas9-mediated mutagenesis. A guide RNAs

(gRNAs) targeting the first exons of *aqp1a.1* and *aqp8a.1* (Fig. S4 and S5) were generated by

using cloning-independent protocol (Gagnon et al., 2014). Briefly, to generate templates for gRNA

transcription, a 60-base oligodeoxynucleotides containing the SP6 promoter sequence, the gene-specific sequence (*aqp1a.1*: 5'-GACAGCTGGCCAGCAGACCC-3'; *aqp8a.1*: 5'-GATGTCTCCCCCATCGCCCG-3') and 23-base overlap region was annealed to an 80-base constant oligodeoxynucleotide encoding the reverse-complement of the tracrRNA tail. The ssDNA overhangs were filled-in with T4 DNA polymerase (New England BioLabs) to generate dsDNA. The gRNAs were *in vitro* transcribed using MEGAScript SP6 kit (Invitrogen), DNase treated and purified with RNA Clean and Concentrator kit (Zymo Research). The gRNAs were quality-checked by running 1 µg of the product on a 2% TBE-agarose gel. Cas9/gRNA RNP complex was assembled just prior to injection and after 5 min incubation at room temperature 200 pg of Cas9 protein (Invitrogen) and 200 pg of gRNAs (100 pg of each, *aqp1a.1* gRNA and *aqp8a.1* gRNA) were co-injected into one-cell stage $Tg(fli1:myr-mCherry)^{ncv1}$ embryos. Such dose produced over 70% embryos survival post-injection showed CRISPR/Cas9-induced somatic *aqp1a.1* and *aqp8a.1* gene mutations. F_0 founders were identified by outcrossing CRISPR/Cas9-injected fish with wild type fish and screening the offspring for mutations at 2 dpf using Sanger sequencing. We found one F_0 fish with germline transmitted mutations in both *aqp1a.1* and *aqp8a.1* genes and outcrossed it with $Tg(fli1:myr-mCherry)^{ncv1}$ and $Tg(fli1a:H2B-EGFP)^{ncv69}$ transgenic fish to establish F_1 generation of heterozygotes. A 55 F_1 fish were genotyped to identify 4 $aqp1a.1^{+/rk28}$, 11 $aqp8a.1^{+/rk29}$ and 12 $aqp1a.1^{+/rk28};aqp8a.1^{+/rk29}$ heterozygote fish. Subsequently, F_1 heterozygote fish were in-crossed to establish a single and double homozygote fish.

DNA isolation and genotyping. Genomic DNA was isolated from either embryos or fin clips using HotSHOT method (Meeker et al., 2007). To identify genomic lesions, the following primers were used: aqp1a1-fwd (5'-CGCCTCCAGATTCATTAGCAGGA-3') and aqp1a1-rev (5'-GTAAGTGAACTGCTGCCAGTGA-3') to amplify a 550 bp fragment, and aqp8a1-fwd (5-GGATCAATTGAGTTGCATAACAGAC-3') and aqp8a1-rev (5'-CTGTAATGTAGACTTGTAAAGTGGA-3') to amplify a 638 bp fragment. Mutations were assessed by direct sequencing of purified PCR products.

Quantitative real-time PCR.
Total RNA from whole embryos was isolated with TRI reagent using Direct-zol RNA MicroPrep kit (Zymo Research) according to the manufacturer's protocol, including on-column DNA

digestion. RNA was quantified using a NanoDrop 1000 spectrophotometer (ThermoFisher
Scientific, USA). cDNA was synthesized from 300 ng of purified RNA using LunaScript RT
SuperMix kit (New England BioLabs) according to the manufacturer's protocol. Amplification of
target cDNA was performed in technical triplicate of three biological replicates using the SYBR
green methods. Each qPCR reaction mixture contained 5 µL 2x Luna Universal qPCR master mix
(New England BioLabs), 1 µL cDNA (2-fold dilution), 0.2 µL antarctic thermolabile UDG (New
England BioLabs) and 250 nM each primer to a final volume of 10 µL. Reactions were run in 384-
well plates (Applied Biosystems) using the QuantStudio 5 Real-Time PCR system (Applied
Biosystems) with the following thermal cycling conditions: initial UDG treatment at 25^0C for 2
minutes, UDG inactivation at 50^0C for 5 min and denaturation at 95^0C for 1 minutes, followed by
45 cycles of 15 sec at 95^0C, 30 sec at 60^0C with a plate read at the end of the extension step.
Control reactions included a no-template control (NTC) and a no-reverse transcriptase control
(NRT). Dissociation analysis of the PCR products was performed by running a gradient from 60
to 95^0C to confirm the presence of a single PCR product. Amplification data was analyzed using
QuantStudio Design and Analysis software v1.5.3 (Applied Biosystems) and relative fold change
was calculated using Pfaffl method (Pfaffl, 2001) and normalized to *gapdh* expression (as an
internal control). Primers are listed in Supplementary Table 2. Two embryos were used in each
biological replicate to generate an average value that was used to calculate the final mean ± SD
from two or three independent experiments.
515
RNA *in situ* hybridization.
RNAscope *in situ* hybridization. RNAscope® *in situ* hybridization was conducted by using
RNAscope Multiplex Fluorescent Reagent kit v2 (Advanced Cell Diagnostics). We adapted
manufacture's protocol designed for samples mounted on slides and protocol developed for whole-
mount embryo samples (Gross-Thebing et al., 2014). For each experimental point, 5 embryos were
processed in one 1.5 ml Eppendorf tube. Briefly, 20 and 22 hpf old embryos were manually
dechorionated and fixed in freshly prepared 4% methanol-free PFA in PBS for 1 hour at room
temperature. After fixation embryos were washed three times in PBS containing 0.01% Tween-20
(PBST), dehydrated stepwise in 5-min washes in a series of increasing methanol concentrations
(25%, 50%, 75%, 100%) in PBS and then stored in 100% methanol at -20^0C before use them for
hybridization. The methanol-stored embryos were incubated with 5% H_2O_2 in methanol for 20 min

at room temperature and then rehydrated stepwise in 5-min washes in series of decreasing
methanol concentrations (75%, 50%, 25%) in PBS, followed by three times 5-min washes in PBST.
Embryos were permeabilized in 2 drops of RNAscope® Protease III for 15 min, rinsed three times
with PBST and hybridized with RNAscope target probes overnight at 40^0C in water bath. After
hybridization probes were recovered and embryos were washed three times with 0.2x SSCT
[0.01% Tween-20 in SSC] at room temperature, re-fixed with 4% PFA for 10 min, and washed
three times with 0.2x SSCT. For RNA detection, the embryos were first hybridized with three
different amplifier solution for 30 min (Amp1, and Amp2) and 15 min (Amp3) in a water bath at
40^0C. After each hybridization step, the embryos were washed three times with 0.2x SSCT for 15
min at room temperature. To develop signal of each probe, the embryos were sequentially
incubated in a water bath at 40^0C with: (i) the horseradish peroxidase (HRP) for 15 min, (ii) TSA
fluorophore for 30 min and (iii) the HRP blocker for 15 min. After each incubation step, the
embryos were washed three times with 0.2x SSCT for 15 min at room temperature. HRP step is
linked to a probe channel, we first developed C1 probe, *Dr-aqp1a.1* (Advanced Cell Diagnostics)
using HRP-C1 and TSA Vivid Fluorophore 520 (Tocris, dilution 1:1500), and then C3 probe, *Dr-*
aqp8a.1 (Advanced Cell Diagnostics) using HRP-C3 and TSA Vivid Fluorophore 650 (Tocris,
dilution 1:1500). Nuclei were counterstained with DAPI ready-to-use solution overnight at 4^0C.
Prior to imaging embryos were rinsed in PBST and kept in 70% glycerol in PBS. Images were
acquired using Olympus FV3000 confocal microscope and an Olympus UPlanXApo 60x/NA 1.42
oil immersion objective.
Chromogenic *in situ* hybridization. Chromogenic *in situ* hybridization was conducted according
to standard protocol (Thisse and Thisse, 2007) with minor modifications. For each experimental
point, 10 embryos were processed in one 1.5 ml Eppendorf tube. Briefly, 30, 48, 72 and 96 hpf old
embryos were manually dechorionated and fixed in freshly prepared 4% methanol-free PFA in
PBS (pH 7.4) at 4^0C overnight. After fixation embryos were washed three times in PBS containing
0.1% Tween-20 (PBST), dehydrated stepwise in 5-min washes in a series of increasing methanol
concentrations (25%, 50%, 75%, 100%) in PBS and then stored in 100% methanol at -20^0C before
use them for hybridization. The methanol-stored embryos were rehydrated stepwise in 5-min
washes in series of decreasing methanol concentrations (75%, 50%, 25%) in PBS followed by
three times 5-min washes in PBST, and permeabilized 20 min with 10 µg/mL proteinase K
(ThermoFisher Scientific). Hybridization was carried out in buffer [50% formamide, 5x SSC, 50

g/mL heparin, 500 g/mL tRNA (Roche) and 0.1% Tween-20] containing 5% dextran sulfate

(MW=500 kDa) at 69^0C overnight. After stringency wash, the specimens were blocked with 2%

blocking reagent (Roche) in maleic acid buffer [100 mM maleic acid, 150 mM NaCl, 50 mM

$MgCl_2$ and 0.1% Tween-20, pH 7.5] and incubated overnight with anti-digoxigenin antibody,

conjugated with alkaline phosphatase (dilution 1:5000, Roche) at 4^0C. For detection of alkaline

phosphatase, specimens were incubated in staining buffer [50 mM Tris-HCl, 50 mM NaCl, 25 mM

$MgCl_2$, 2% Polyvinyl alcohol and 0.1% Tween-20, pH 9.5] containing 375 µg/mL nitro-blue

tetrazolium chloride (Roche) and 175 µg/mL 5-bromo-4-chloro-3'-indolyl-phosphate (Roche).

Embryos were cleared in 70% glycerol overnight and imaged using Leica M205FA microscope.

RNA probe synthesis. The cDNA-containing vectors were linearized with appropriate restriction

enzymes (detailed maps of vectors will be provided upon request) and used as a template for RNA

probe synthesis. Sense and antisense RNA probes were synthesized using MEGAscript SP6 or T7

kits (Invitrogen) and digoxigenin (DIG)-labeled rNTPs (Roche) according to the manufacturer's

protocol and purified using RNA Clean and Concentrator kit (Zymo Research). The following

sense and antisense DIG-labeled riboprobes were generated: (i) *aqp1a.1*, 1.1 kb long probe

comprised 121 nt of 5'UTR, 783 nt of the open reading frame (ORF) and 207 nt of 3'UTR; (ii)

aqp8a.1, 0.8 kb long probe comprised 17 nt of 5'UTR, 783 nt of ORF and 30 nt of 3'UTR. Sense

probes showed no specific or unspecific staining (data not shown).

Image processing.

Chromogenic and RNAscope *in situ* hybridization images were processed using ImageJ software

(NIH, version 2.9.0/1.53t) with brightness and contrast adjustments. Z-stacks of fluorescent

images are presented as maximum intensity projections.

Analysis of mRNA expression in tip and stalk cells.

Embryos were mounted on slide and imaged using Olympus FV3000 confocal microscope and

Olympus UPlanXApo 60x/NA 1.42 oil immersion objective. Maximum intensity projection of z-

stacks was used to determine the level of cellular fluorescence in ImageJ. The tip and stalk cells

were selected using freehand selection tool, "mean grey value", "area" and "integrated density" in

each ROI were measured first for channel 1 (aqp1a.1). A region next to cell that has no

fluorescence was selected as background and measured. The step was repeated for channel 2

(aqp8a.1) on the same ROIs. Two parameters were calculated for each channel, (i) the corrected

total cell fluorescence, CTCF=Integrated Density – (Area of selected cell x Mean fluorescence of

background) and (ii) ratio, R=CTCF of tip cell/CTCF of stalk cell. We consider that if R <0.9,

expression is lower in tip cell, if R=0.9-1.1, expression is equal in tip and stalk cells, if R>1.1,

expression is higher in tip cell.

Imaging.

For live confocal imaging, embryos were mounted in 0.8% low-melt agarose (Bio-Rad) in E3

medium containing 0.16 mg/mL Tricaine (Sigma-Aldrich) and 0.003% phenylthiourea (Sigma-

Aldrich) in glass bottom 35-mm dishes (MatTek). Confocal Z-stacks were acquired using an

inverted Olympus IX83/Yokogawa CSU-W1 spinning disc confocal microscope equipped with a

Zyla 4.2 CMOS camera (Andor) and Olympus UPLSAPO 40x/NA 1.25 or 30x/NA 1.05 silicone

oil immersion objectives. Bright-field images were acquired on Leica M205FA microscope.

Images were processed using ImageJ software (NIH, version 2.9.0/1.53t).

Analysis of cell and nuclei migration.

aqp1a.1$^{+/rk28}$;aqp8a.1$^{+/rk29}$ and   *aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$*   embryos   on   *Tg(fli1a:H2B-*

EGFP)^{ncv69};(fli1:Lifeact-mCherry)^{ncv7} double transgenic background were imaged from 21 to 30

hpf with time interval 6 min using Olympus UPLSAPO 30x/NA 1.05 silicon oil immersion

objective. We tried to mount both heterozygote and mutant embryos on the same dish (1 het plus

2 mut, or 2 het plus 2 mut) to image embryos on the same experimental conditions. Acquired time-

lapse images were registered using HyperStackReg plugin in ImageJ, tip cell and nuclei were

tracked using Manual Tracking plugin and velocity was calculated.

Analysis of grow rate of actin bundles in filopodia.

Wild   type   and   *aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$*   embryos   on   *Tg(fli1ep:EGFP-*

PLC1δPH)^{rk26};(fli1:Lifeact-mCherry)^{ncv7} double transgenic background were imaged starting from

25 hpf for 10 min with time interval 30 sec using Olympus UPLSAPO 60x/NA 1.2 water

immersion objective. Actin bundles growth was tracked using Manual Tracking plugin and

velocity was calculated.

Analysis of filopodia number and length.

Wild type, *aqp1a.1^{+/rk28};aqp8a.1^{+/rk29}* and *aqp1a.1^{rk28/rk28};aqp8a.1^{rk29/rk29}* embryos on

Tg(fli1ep:EGFP-PLC1δPH)^{rk26};(fli1:Lifeact-mCherry)^{ncv7} double transgenic background were

imaged starting from 24 hpf for 30 min with time interval of 50 sec using Olympus UPLSAPO

60x/NA 1.2 water immersion objective. In a single image we counted filopodia number per vessel,

measured membrane length and calculated filopodia number.

Chemical treatment.

Latrunculin B (Merck Millipore) was dissolved in DMSO to 1 mg/ml and stored at -20^0C.

Dibenzazepine (YO-01027) (Selleck Chemicals) was prepared as 10 mM solution in DMSO and

stored at -80^0C. Ki8751 (Selleck Chemicals) was prepared as 5 mM solution in DMSO and stored

at -80^0C. All compounds were diluted to the desired concentration in E3 medium.

Statistical analysis.

Statistical analysis was performed using Prism software version 10.2.0 (GraphPad). The variance

between the mean values of two groups was evaluated using the unpaired Student's *t*-test. For

assessment of more than three groups, we used one-way analysis of variance (ANOVA) test. A *P*

value of <0.05 was considered statistically significant. Statistic details can be found in each figure

legend.

Cell culture.

Adult human aortic endothelial cells (HAEC) (Lonza, lot-20TL231227) were cultured in EGM

medium (Lonza) and used at passage 3. Cells were seeded in EBM medium (Lonzo) supplemented

with 2% FBS (Gibco) at 4×10^5 cells/well on 24-well plate (Trueline) coated with 5 μg/ml

fibronectin (Sigma-Aldrich). Cells were treated with either 0.01% DMSO or different

concentrations of Ki8751 inhibitor for 6 hours. Cells were lysed with TRI reagent and RNA was

isolated using Direct-zol RNA MicroPrep kit (Zymo Research).

Single-cell RNA sequencing.

ECs were isolated from *Tg(kdrl:EGFP)^{s843}* transgenic embryos. Cell sorting was carried out with

FACSAriaII Cell Sorter (BD Bioscience). Single-cell suspension was loaded into the 10X

Chromium system and cDNA libraries were constructed using Chromium Next GEM Single Cell

3′ GEM, Library and Gel Bead Kit v2 (10X Genomics) according to manufacturer's protocol.

Library was sequenced on the Illumina HiSeq 1500 Sequencer (Illumina, USA). Cell Ranger v2.1

was used to de-multiplex raw base call (BCL) files generated by Illumina sequencers into FASTQ

files, perform the alignment, barcode counting, and UMI counting. Sequencing reads were mapped

to the zebrafish genome assembly (GRCz11, Ensembl release 92). Further analyses were

performed using Seurat package49 (version 4.3.0) in R software (https://www.r-project.org). The

expression matrices were first filtered by keeping genes that are expressed in a minimum of 3 cells

and cells that expressed a minimum of 200 genes for downstream analysis. The data were then

normalized using NormalizeData function which normalizes the gene expression for each cell by

the total expression counts. To correct for batch effect between the data from the two-time points,

the rpca method in the Seurat package was applied to integrate the data. The top 2000 variable

genes identified using the vst method in FindVariableFeatures function were used for principal

component analysis in RunPCA function, and the first 30 principal components were used for

visualization analysis with Uniform Manifold Approximation and Projection (UMAP) method,

and in FindNeighbors function analysis. The cell clustering resolution was set at 0.5 in

FindClusters function.

Acknowledgements

We thank members of the Phng Lab, Y-C. Wang and S. Thukral for discussions and suggestions;

G. Chen, E. Taniguchi, RIKEN BDR Research Aquarium and RIKEN Kobe BioImaging Facilities

& Factory for technical assistance; J. Vermot, M. Francois and A. Yap for comments on the

manuscript. This work was supported by core funding from RIKEN BDR (to L.K.P.), RIKEN

BDR-Otsuka Pharmaceutical Collaboration Center (I.K.), the Naito Foundation (L.K.P.), the JSPS

Grants-in-Aid for Scientific Research (KAKENHI) grants (22H02624, 22H05168 to L.K.P;

22K06244 to I.K.), Swedish Research Council (2015-00550 to C.B.); Swedish Cancer Society

(2018/449, 2018/1154 to C.B.); Knut and Alice Wallenberg Foundation (2020.0057, 2018.0218 to

C.B.); Swedish Brain Foundation (ALZ2019-0130, ALZ2022-0005 to C.B.); the Leducq

Foundation (22CVD01, 23CVD02 to C.B.) and the Innovative Medicines Initiative (IM2PACT-

807015 to C.B.)

Authorship contributions

Conceptualization: L.K.P.; Methodology: I.K., H.W. and L.K.P.; Formal analysis: I.K. and L.H.;

Resources: C.B. and L.K.P.; Writing: I.K. and L.K.P; Supervision: L.K.P.

Declaration of interests

The authors declare no competing interests.

Figure legends

Figure 1. Differential expression of *aqp1a.1* and *aqp8a.1* mRNAs in tip and stalk cells.

(A-F) Detection of *aqp1a.1* and *aqp8a.1* mRNA by RNAscope *in situ* hybridization in 20 (A - D)

and 22 (E and F) hpf zebrafish embryos. Representative maximum intensity projection of confocal

z-stacks of tip and stalk cells sprouting from DA are shown (n = 9 embryos, 2 independent

experiments). B and F show magnified region of a tip cell from a 20 and 22 hpf zebrafish,

respectively (tip cell and EC nuclei in the DA are outlined). Scale bar, 10 μm (B and F) and 40 μm

(A, C - E).

(G) Percentage of vascular sprouts with differential expression of *aqp1a.1* and *aqp8a.1* mRNAs

in tip and stalk cells in 22 hpf zebrafish (n=23 sprouts from 8 embryos, 2 independent experiments).

(H) qPCR analysis of *aqp1a.1* and *aqp8a.1* expression in zebrafish embryos treated with 0.01%

DMSO or 1 μM ki8751 at 20 hpf for 6 or 24 hours (n = 3 independent experiments).

(I) qPCR analysis of *AQP1* gene expression in HAECs treated with 0.01% DMSO or 1 nM and 10

nM ki8751 for 6 hr (n = 2 independent experiments).

(J) qPCR analysis of *aqp1a.1* and *aqp8a.1* gene expression in the zebrafish embryos treated with

0.1% DMSO or 10 μM DBZ at 20 hpf for 6 or 24 hr (n = 2 independent experiments). In panels

H-J, gene expressions are shown relative to *gapdh* expression and data are presented as mean ±

SD; statistical significance was determined by Brown-Forsythe and Welch ANOVA tests with

Dunnett's multiple comparisons test; ns $p > 0.05$, **$p < 0.01$, ***$p < 0.001$, and ****$p < 0.0001$. DA,

dorsal aorta; PCV, posterior cardinal vein.

See also Fig. S3.

Figure 2. Aquaporin proteins are enriched at the leading front of migrating cells.

(A and B) Representative maximum intensity projection confocal z-stacks of endothelial tip cells of 25 hpf *Tg(fli1ep:aqp1a.1-mEmerald)^{rk30};(kdrl:ras-mCherry)^{s916}* (A) and *Tg(fli1ep:aqp8a.1-mEmerald)^{rk31};(kdrl:ras-mCherry)^{s916}* (B) embryos. Scale bar, 20 µm.

(C and D) Still images from time-lapse movies of migrating tip cells from *Tg(fli1ep:aqp1a.1-mEmerald)^{rk30}* (C) and *Tg(fli1ep:aqp8a.1-mEmerald)^{rk31}* (D) embryos. Movies were taken from 24 to 30 hpf. Black arrowheads, Aquaporin protein localization at the leading edge of migrating tip cells. Arrows, Aquaporin localization in filopodia. Time, h:min. Scale bar, 5 µm.

See also Movie S1 and S2.

Figure 3. Loss of Aquaporin function leads to defective trunk vessel formation.

(A-D) Representative maximum intensity projection confocal z-stacks of 28 hpf *aqp1a.1$^{+/rk28}$;aqp8a.1$^{+/rk29}$* (A), *aqp1a.1$^{rk28/rk28}$* (B), *aqp8a.1$^{rk29/rk29}$* (C) and *aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$* (D) embryos.

(E) Quantification of ISV length in 28 hpf embryos (*aqp1a.1$^{+/rk28}$;aqp8a.1$^{+/rk29}$*: n = 21 embryos; *aqp1a.1$^{rk28/rk28}$*: n = 20 embryos; *aqp8a.1$^{rk29/rk29}$*: n = 20 embryos; *aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$*: n = 19 embryos from 2 independent experiments).

(F) Illustration of aISV and vISV connections in the trunk of wild type embryo at 3 dpf.

(G-J) Representative maximum intensity projection confocal z-stacks of 3 dpf wild type (G), *aqp8a.1$^{rk29/rk29}$* (H), *aqp1a.1$^{rk28/rk28}$* (I), and *aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$* (J) embryos. Asterisk marks incomplete ISVs.

(K) Quantification of percentage of wild type and *aquaporin* mutant embryos at 3 dpf with at least one incomplete ISVs per embryo (wild type, n = 47 embryos; *aqp1a.1$^{rk28/rk28}$*, n = 30 embryos; *aqp8a.1$^{rk29/rk29}$*, n = 38 embryos; *aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$*, n = 204 embryos).

(L) A pie chart showing proportion of different classes of incomplete ISV phenotypes (I-VI) found in *aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$* mutant embryos at 3 dpf.

DA, dorsal aorta; DLAV, dorsal longitudinal anastomotic vessel; ISV, intersegmental vessel; aISV, arterial ISV; vISV, venous ISV; NC, notochord; NT, neural tube; PCV, posterior cardinal vein. Scale bar, 50 µm (G-J) and 200 µm (A-D).

See also Fig. S6, S7 and S8.

Figure 4. Aquaporins promote endothelial tip cell protrusion and migration.

(A) Timing of tip cell emergence from the DA in $aqp1a.1^{+/rk28};aqp8a.1^{+/rk29}$ (n = 23) and

$aqp1a.1^{rk28/rk28};aqp8a.1^{rk29/rk29}$ (n = 27).

(B and C) Still images from time-lapse imaging of migrating tip cells from

$aqp1a.1^{+/rk28};aqp8a.1^{+/rk29}$ (B) and $aqp1a.1^{rk28/rk28};aqp8a.1^{rk29/rk29}$ (C) embryos from 20 to 30 hpf.

White arrows, retracting tip cell. *, secondary sprouting from PCV. Scale bar, 10 μm.

(D and E) Stills images from representative time-lapse movies of migrating tip cells in

$aqp1a.1^{+/rk28};aqp8a.1^{+/rk29}$ (D, n = 2) and $aqp1a.1^{rk28/rk28};aqp8a.1^{rk29/rk29}$ (E, n = 5) embryos.

Movies were taken from 25 to 30 hpf (n = 2 independent experiments). Bracket, formation of stable

protrusions. Scale bar, 20 μm.

(F and G) Representative maximum intensity projection confocal z-stacks of tip cells from wild

type (F) and $aqp1a.1^{rk28/rk28};aqp8a.1^{rk29/rk29}$ (G) embryos at 26 hpf. Scale bar, 10 μm.

(H) Quantification of filopodia number in tip cells of wild type (n = 36 cells from 10 embryos, 2

independent experiments), $aqp1a.1^{+/rk28};aqp8a.1^{+/rk29}$ (n = 19 cells from 6 embryos, 2 independent

experiments) and $aqp1a.1^{rk28/rk28};aqp8a.1^{rk29/rk29}$ (n = 28 cells from 9 embryos, 2 independent

experiments) embryos.

(I) Quantification of filopodia length in tip cells of wild type (n = 24 cells from 7 embryos, 2

independent experiments), $aqp1a.1^{+/rk28};aqp8a.1^{+/rk29}$ (n = 16 cells from 7 embryos, 2 independent

experiments) and $aqp1a.1^{rk28/rk28};aqp8a.1^{rk29/rk29}$ (n = 11 cells from 6 embryos, 2 independent

experiments) embryos.

(J) Quantification of tip cell leading edge displacement of $aqp1a.1^{+/rk28};aqp8a.1^{+/rk29}$ (n = 19 cells

from 5 embryos, 3 independent experiments) and $aqp1a.1^{rk28/rk28};aqp8a.1^{rk29/rk29}$ (n = 47 cells from

13 embryos, 6 independent experiments) embryos at 25 - 30 hpf.

(K) Quantification of tip cell nuclei displacement of $aqp1a.1^{+/rk28};aqp8a.1^{+/rk29}$ (n = 20 cells from

6 embryos, 4 independent experiments) and $aqp1a.1^{rk28/rk28};aqp8a.1^{rk29/rk29}$ (n = 20 cells from 4

embryos, 2 independent experiments) embryos at 21 - 30 hpf.

(L) Quantification of tip cell migration velocity in $aqp1a.1^{+/rk28};aqp8a.1^{+/rk29}$ (n = 19 cells from 5

embryos, 3 independent experiments) and $aqp1a.1^{rk28/rk28};aqp8a.1^{rk29/rk29}$ (n = 47 cells from 13

embryos, 6 independent experiments) embryos at 25 - 30 hpf.

Statistical significance was determined by Brown-Forsythe and Welch ANOVA tests with

Dunnett's (H) or Sidak's (I) multiple comparisons test, and with unpaired t-test (L). ns, $p>0.05$, *,

$p < 0.05$, ***, $p < 0.001$, and ****, $p < 0.0001$.

See also Movie S3-6.

Figure 5. Aquaporin-mediated water influx increases tip cell volume and actin elongation in

filopodia.

(A and B) Representative maximum intensity projection confocal z-stacks of tip cell and its 3D-

volume rendering of 25 hpf wild type (A) and *aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$* (B) labelled with

cytosolic mEmerald. Scale bar, 10 μm.

(C) Quantification of tip cell volume in 25 hpf wild type (n = 24 cells) and

aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$ (n = 20 cells) embryos from 3 independent experiments.

(D) Quantification of growth rate of actin bundles in tip cell filopodia in 25 hpf wild type (n = 12

cells from 7 embryos, 2 independent experiments) and *aqp1a.1^{rk28};aqp8a.1^{rk29}* (n = 12 cells from

6 embryos, 2 independent experiments) embryos.

Statistical significance was determined by unpaired *t*-test with (C) or without (D) Welch's

correction.

Figure 6. Additive function of actin polymerization and hydrostatic pressure in driving EC

migration and sprouting angiogenesis.

(A-D) Representative maximum intensity projection confocal z-stacks of 28 hpf wild type (A and

B) and *aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$* (C and D) embryos treated with 0.1% DMSO (A and C) or

0.8 μg/ml Lat. B (B and D) from 20 to 28 hpf (for each condition: wild type, n = 10 embryos;

aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$, n =12 embryos, 2 independent experiments). Scale bar, 20 μm.

(E) Quantification of ISV length in 28 hpf wild type and *aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$* embryos

treated with 0.1% DMSO or 0.8 μg/ml Lat. B (wild type: n = 60 vessels from 10 control embryos,

n = 66 vessels from 10 Lat. B-treated embryos; *aqp1a.1^{rk28};aqp8a.1^{rk29}*: n = 126 vessels from 12

control embryos, n = 112 vessels from 11 Lat. B-treated embryos, 2 independent experiments).

Statistical significance was determined by Brown-Forsythe ANOVA test with Sidak's multiple

comparison test; ns, $p,>,0.05$, **$p < 0.01$, and ****, $p < 0.0001$. (F) Endothelial tip cells generate

membrane protrusions by actin polymerization and increased hydrostatic pressure via Aquaporin-

mediated water inflow. In the absence of actin polymerization and presence of Aquaporins,

hydrostatic pressure deforms membranes to generate membrane protrusions. When Aquaporin

function is lost, membrane protrusions is generated by actin polymerization. Fewer membrane

protrusions are formed when only one mechanism is utilized, resulting in slower EC migration.

See also Movie S7.

806
807   **References**
808

Agre P, King LS, Yasui M, Guggino WB, Ottersen OP, Fujiyoshi Y, Engel A, Nielsen S. 2002.
Aquaporin water channels – from atomic structure to clinical medicine. *J Physiol* **542**:3–16.
doi:10.1113/jphysiol.2002.020818

Ando K, Wang W, Peng D, Chiba A, Lagendijk AK, Barske L, Crump JG, Stainier DYR,
Lendahl U, Koltowska K, Hogan BM, Fukuhara S, Mochizuki N, Betsholtz C. 2019. Peri-
arterial specification of vascular mural cells from naïve mesenchyme requires Notch
signaling. *Development* **146**:dev165589-13. doi:10.1242/dev.165589

Baeyens N, Nicoli S, Coon BG, Ross TD, Dries K den, Han J, Lauridsen HM, Mejean CO,
Eichmann A, Thomas J-L, Humphrey JD, Schwartz MA. 2015. Vascular remodeling is
governed by a VEGFR3-dependent fluid shear stress set point. *Elife* **4**.
doi:10.7554/elife.04645

Benedito R, Hellström M. 2013. Notch as a hub for signaling in angiogenesis. *Exp Cell Res* 1–8.

Benfenati V, Caprini M, Dovizio M, Mylonakou MN, Ferroni S, Ottersen OP, Amiry-
Moghaddam M. 2011. An aquaporin-4/transient receptor potential vanilloid 4
(AQP4/TRPV4) complex is essential for cell-volume control in astrocytes. *Proc Natl Acad*
Sci USA **108**:2563–8. doi:10.1073/pnas.1012867108

Bill BR, Korzh V. 2014. Choroid plexus in developmental and evolutionary perspective. *Front*
Neurosci **8**:363. doi:10.3389/fnins.2014.00363

Boer LL de, Vanes L, Melgrati S, O'May JB, Hayward D, Driscoll PC, Day J, Griffiths A,
Magueta R, Morrell A, MacRae JI, Köchl R, Tybulewicz VLJ. 2023. T cell migration requires
ion and water influx to regulate actin polymerization. *Nat Commun* **14**:7844.
doi:10.1038/s41467-023-43423-8

Bussmann J, Wolfe SA, Siekmann AF. 2011. Arterial-venous network formation during brain
vascularization involves hemodynamic regulation of chemokine signaling. *Development*
138:1717–1726.

Calamita G, Ferri D, Gena P, Liquori GE, Cavalier A, Thomas D, Svelto M. 2005. The Inner
Mitochondrial Membrane Has Aquaporin-8 Water Channels and Is Highly Permeable to
Water. *J Biol Chem* **280**:17149–17153. doi:10.1074/jbc.c400595200

Chae YK, Woo J, Kim M-J, Kang SK, Kim MS, Lee J, Lee SK, Gong G, Kim YH, Soria JC,
Jang SJ, Sidransky D, Moon C. 2008. Expression of Aquaporin 5 (AQP5) Promotes Tumor
Invasion in Human Non Small Cell Lung Cancer. *PLoS ONE* **3**:e2162.
doi:10.1371/journal.pone.0002162

Chan CJ, Costanzo M, Ruiz-Herrero T, Mönke G, Petrie RJ, Bergert M, Diz-Muñoz A,
Mahadevan L, Hiiragi T. 2019. Hydraulic control of mammalian embryo size and cell fate.
Nature **571**:112–116. doi:10.1038/s41586-019-1309-x

Chang Y-F, Imam JS, Wilkinson MF. 2007. The Nonsense-Mediated Decay RNA Surveillance
Pathway. *Annu Rev Biochem* **76**:51–74. doi:10.1146/annurev.biochem.76.050106.093909

Chauvigné F, Boj M, Finn RN, Cerdà J. 2015. Mitochondrial aquaporin-8-mediated hydrogen
peroxide transport is essential for teleost spermatozoon motility. *Sci Rep* **5**:7789.
doi:10.1038/srep07789

Chauvigné F, Ducat C, Ferré A, Hansen T, Carrascal M, Abián J, Finn RN, Cerdà J. 2021. A
multiplier peroxiporin signal transduction pathway powers piscine spermatozoa. *Proc Natl
Acad Sci USA* **118**:e2019346118. doi:10.1073/pnas.2019346118

Choudhury MI, Benson MA, Sun SX. 2022. Trans-epithelial fluid flow and mechanics of
epithelial morphogenesis. *Semin Cell Dev Biol* **131**:146–159.
doi:10.1016/j.semcdb.2022.05.020

Chugh M, Munjal A, Megason SG. 2022. Hydrostatic pressure as a driver of cell and tissue
morphogenesis. *Semin Cell Dev Biol* **131**:134–145. doi:10.1016/j.semcdb.2022.04.021

Clarke DN, Martin AC. 2021. Actin-based force generation and cell adhesion in tissue
morphogenesis. *Curr Biol* **31**:R667–R680. doi:10.1016/j.cub.2021.03.031

Conner MT, Conner AC, Bland CE, Taylor LHJ, Brown JEP, Parri HR, Bill RM. 2012. Rapid
Aquaporin Translocation Regulates Cellular Water Flow. *J Biol Chem* **287**:11516–11525.
doi:10.1074/jbc.m111.329219

Day RE, Kitchen P, Owen DS, Bland C, Marshall L, Conner AC, Bill RM, Conner MT. 2014.
Human aquaporins: regulators of transcellular water flow. *Biochim Biophys Acta* **1840**:1492–
1506. doi:10.1016/j.bbagen.2013.09.033

Dumortier JG, Verge-Serandour ML, Tortorelli AF, Mielke A, Plater L de, Turlier H, Maître J-L.
2019. Hydraulic fracturing and active coarsening position the lumen of the mouse blastocyst.
Science **365**:465–468. doi:10.1126/science.aaw7709

Emelyanov A, Gao Y, Naqvi NI, Parinov S. 2006. Trans-Kingdom Transposition of the Maize
Dissociation Element. *Genetics* **174**:1095–1104. doi:10.1534/genetics.106.061184

Emelyanov A, Parinov S. 2008. Mifepristone-inducible LexPR system to drive and control gene
expression in transgenic zebrafish. *Dev Biol* **320**:113–121. doi:10.1016/j.ydbio.2008.04.042

Farinas J, Kneen M, Moore M, Verkman AS. 1997. Plasma Membrane Water Permeability of
Cultured Cells and Epithelia Measured by Light Microscopy with Spatial Filtering. *J Gen*
Physiol **110**:283–296. doi:10.1085/jgp.110.3.283

Figueiredo AM, Barbacena P, Russo A, Vaccaro S, Ramalho D, Pena A, Lima AP, Ferreira RR,
Fidalgo MA, El-Marjou F, Carvalho Y, Vasconcelos FF, Lennon-Duménil A-M, Vignjevic
DM, Franco CA. 2021. Endothelial cell invasion is controlled by dactylopodia. *Proc Natl*
Acad Sci USA **118**. doi:10.1073/pnas.2023829118

Fukuhara S, Zhang J, Yuge S, Ando K, Wakayama Y, Sakaue-Sawano A, Miyawaki A,
Mochizuki N. 2014. Visualizing the cell-cycle progression of endothelial cells in zebrafish.
Dev Biol **393**:10–23. doi:10.1016/j.ydbio.2014.06.015

Gagnon JA, Valen E, Thyme SB, Huang P, Akhmetova L, Ahkmetova L, Pauli A, Montague
TG, Zimmerman S, Richter C, Schier AF. 2014. Efficient mutagenesis by Cas9 protein-
mediated oligonucleotide insertion and large-scale assessment of single-guide RNAs. *PLoS*
ONE **9**:e98186. doi:10.1371/journal.pone.0098186

Gebala V, Collins R, Geudens I, Phng L-K, Gerhardt H. 2016. Blood flow drives lumen
formation by inverse membrane blebbing during angiogenesis in vivo. *Nat Cell Biol* **18**:443–
450. doi:10.1038/ncb3320

Gerhardt H, Golding M, Fruttiger M, Ruhrberg C, Lundkvist A, Abramsson A, Jeltsch M,
Mitchell C, Alitalo K, Shima D, Betsholtz C. 2003. VEGF guides angiogenic sprouting
utilizing endothelial tip cell filopodia. *J Cell Biol* **161**. doi:10.1083/jcb.200302047

Gradilone SA, Carreras FI, Lehmann GL, Marinelli RA. 2005. Phosphoinositide 3-kinase is
involved in the glucagon-induced translocation of aquaporin-8 to hepatocyte plasma
membrane. *Biol Cell* **97**:831–836. doi:10.1042/bc20040115

Gradilone SA, García F, Huebert RC, Tietz PS, Larocca MC, Kierbel A, Carreras FI, Larusso
NF, Marinelli RA. 2003. Glucagon induces the plasma membrane insertion of functional
aquaporin-8 water channels in isolated rat hepatocytes. *Hepatology* **37**:1435–41.
doi:10.1053/jhep.2003.50241

Gross-Thebing T, Paksa A, Raz E. 2014. Simultaneous high-resolution detection of multiple
transcripts combined with localization of proteins in whole-mount embryos. *BMC Biol* **12**:55.
doi:10.1186/s12915-014-0055-7

Groth C, Alvord WG, Quiñones OA, Fortini ME. 2010. Pharmacological Analysis of Drosophila
melanogaster γ-Secretase with Respect to Differential Proteolysis of Notch and APP. *Mol*
Pharmacol **77**:567–574. doi:10.1124/mol.109.062471

Herrera M, Garvin JL. 2007. Novel role of AQP-1 in NO-dependent vasorelaxation. *Am J*
Physiol Ren Physiol **292**:F1443–F1451. doi:10.1152/ajprenal.00353.2006

Herrera M, Hong NJ, Garvin JL. 2006. Aquaporin-1 Transports NO Across Cell Membranes.
Hypertension **48**:157–164. doi:10.1161/01.hyp.0000223652.29338.77

Hogan BM, Bos FL, Bussmann J, Witte M, Chi NC, Duckers HJ, Schulte-Merker S. 2009. ccbe1
is required for embryonic lymphangiogenesis and venous sprouting. *Nat Genet* **41**.
doi:10.1038/ng.321

Hu J, Verkman AS, Hu J, Verkman AS. 2006. Increased migration and metastatic potential of
tumor cells expressing aquaporin water channels. *FASEB J* **20**:1892–1894. doi:10.1096/fj.06-
5930fje

Jahn TP, Møller ALB, Zeuthen T, Holm LM, Klærke DA, Mohsin B, Kühlbrandt W, Schjoerring
JK. 2004. Aquaporin homologues in plants and mammals transport ammonia. *FEBS Lett*
574:31–36. doi:10.1016/j.febslet.2004.08.004

Jin S-W, Beis D, Mitchell T, Chen J-N, Stainier DY. 2005. Cellular and molecular analyses of
vascular tube and lumen formation in zebrafish. *Development* **132**. doi:10.1242/dev.02087

Jung HJ, Park J-Y, Jeon H-S, Kwon T-H. 2011. Aquaporin-5: A Marker Protein for Proliferation
and Migration of Human Breast Cancer Cells. *PLoS ONE* **6**:e28492.
doi:10.1371/journal.pone.0028492

Karlsson T, Bolshakova A, Magalhães MAO, Loitto VM, Magnusson K-E. 2013. Fluxes of
Water through Aquaporin 9 Weaken Membrane-Cytoskeleton Anchorage and Promote
Formation of Membrane Protrusions. *PLoS ONE* **8**:e59901.
doi:10.1371/journal.pone.0059901

Karlsson T, Glogauer M, Ellen RP, Loitto V, Magnusson K, Magalhães MAO. 2011. Aquaporin
9 phosphorylation mediates membrane localization and neutrophil polarization. *J Leukoc Biol*
90:963–973. doi:10.1189/jlb.0910540

Kimmel CB, Ballard WW, Kimmel SR, Ullmann B, Schilling TF. 1995. Stages of embryonic
development of the zebrafish. *Dev Dyn* **203**:253--310.

Kondrychyn I, Kelly DJ, Carretero NT, Nomori A, Kato K, Chong J, Nakajima H, Okuda S,
Mochizuki N, Phng L-K. 2020. Marcksl1 modulates endothelial cell mechanoresponse to
haemodynamic forces to control blood vessel shape and size. *Nat Commun* **11**:5476–18.
doi:10.1038/s41467-020-19308-5

Koun S, Kim J-D, Rhee M, Kim M-J, Huh T-L. 2016. Spatiotemporal expression pattern of the
zebrafish aquaporin 8 family during early developmental stages. *Gene Expr Patterns* **21**:1–6.
doi:10.1016/j.gep.2016.06.001

Kozono D, Yasui M, King LS, Agre P. 2002. Aquaporin water channels: atomic structure
molecular dynamics meet clinical medicine. *J Clin Invest* **109**:1395–1399.
doi:10.1172/jci15851

Li Y, Konstantopoulos K, Zhao R, Mori Y, Sun SX. 2020. The importance of water and
hydraulic pressure in cell dynamics. *J Cell Sci* **133**:jcs240341. doi:10.1242/jcs.240341

Liu K, Nagase H, Huang CG, Calamita G, Agre P. 2006. Purification and functional
characterization of aquaporin-8. *Biol Cell* **98**:153–161. doi:10.1042/bc20050026

Loitto VM, Huang C, Sigal YJ, Jacobson K. 2007. Filopodia are induced by aquaporin-9
expression. *Exp Cell Res* **313**:1295–1306. doi:10.1016/j.yexcr.2007.01.023

Loitto VM, Karlsson T, Magnusson K. 2009. Water flux in cell motility: Expanding the
mechanisms of membrane protrusion. *Cell Motil Cytoskeleton* **66**:237–247.
doi:10.1002/cm.20357

Ma T, Yang B, Verkman AS. 1997. Cloning of a Novel Water and Urea-Permeable Aquaporin
from Mouse Expressed Strongly in Colon, Placenta, Liver, and Heart. *Biochem Biophys Res*
Commun **240**:324–328. doi:10.1006/bbrc.1997.7664

Mallavarapu A, Mitchison T. 1999. Regulated Actin Cytoskeleton Assembly at Filopodium Tips
Controls Their Extension and Retraction. *J Cell Biol* **146**:1097–1106.
doi:10.1083/jcb.146.5.1097

Maltaneri RE, Schiappacasse A, Chamorro ME, Nesse AB, Vittori DC. 2020. Aquaporin-1 plays
a key role in erythropoietin-induced endothelial cell migration. *Biochim Biophys Acta Mol*
Cell Res **1867**:118569. doi:10.1016/j.bbamcr.2019.118569

Marikawa Y, Alarcon VB. 2012. Mouse Development, From Oocyte to Stem Cells. *Results*
Probl Cell Differ **55**:165–184. doi:10.1007/978-3-642-30406-4_9

Markou A, Unger L, Abir-Awan M, Saadallah A, Halsey A, Balklava Z, Conner M, Törnroth-
Horsefield S, Greenhill SD, Conner A, Bill RM, Salman MM, Kitchen P. 2022. Molecular
mechanisms governing aquaporin relocalisation. *Biochim Biophys Acta Biomembr*
1864:183853. doi:10.1016/j.bbamem.2021.183853

McLennan R, McKinney MC, Teddy JM, Morrison JA, Kasemeier-Kulesa JC, Ridenour DA,
Manthe CA, Giniunaite R, Robinson M, Baker RE, Maini PK, Kulesa PM. 2019. Neural crest
cells bulldoze through the microenvironment using Aquaporin 1 to stabilize filopodia.
Development **147**:dev185231. doi:10.1242/dev.185231

Meeker ND, Hutchinson SA, Ho L, Trede NS. 2007. Method for isolation of PCR-ready
genomic DNA from zebrafish tissues. *Biotech* **43**:610–614.

Mosaliganti KR, Swinburne IA, Chan CU, Obholzer ND, Green AA, Tanksale S, Mahadevan L,
Megason SG. 2019. Size control of the inner ear via hydraulic feedback. *Elife* **8**:e39596.
doi:10.7554/elife.39596

Munjal A, Lecuit T. 2014. Actomyosin networks and tissue morphogenesis. *Development*
141:1789-1793. doi:10.1242/dev.091645

Murrell M, Oakes PW, Lenz M, Gardel ML. 2015. Forcing cells into shape: the mechanics of
actomyosin contractility. *Nat Cell Biol* **16**:486–498. doi:10.1038/nrm4012

Nagy TL, Strickland J, Weiner OD. 2023. Neutrophils actively swell to potentiate rapid
migration. *bioRxiv* 2023.05.15.540704. doi:10.1101/2023.05.15.540704

Nakajima H, Yamamoto K, Agarwala S, Terai K, Fukui H, Fukuhara S, Ando K, Miyazaki T,
Yokota Y, Schmelzer E, Belting H-G, Affolter M, Lecaudey V, Mochizuki N. 2017. Flow-
Dependent Endothelial YAP Regulation Contributes to Vessel Maintenance. *Dev Cell* **40**:523-
536.e6. doi:10.1016/j.devcel.2017.02.019

Nakhoul NL, Davis BA, Romero MF, Boron WF. 1998. Effect of expressing the water channel
aquaporin-1 on the CO2 permeability of Xenopus oocytes. *Am J Physiol Cell Physiol*
274:C543–C548. doi:10.1152/ajpcell.1998.274.2.c543

Noda Y, Sasaki S. 2005. Trafficking mechanism of water channel aquaporin-2. *Biol Cell*
97:885–892. doi:10.1042/bc20040120

Papadopoulos MC, Saadoun S, Verkman AS. 2008. Aquaporins and cell migration. *Pflugers*
Arch **456**:693–700. doi:10.1007/s00424-007-0357-5

Parab S, Quick RE, Matsuoka RL. 2021. Endothelial cell-type-specific molecular requirements
for angiogenesis drive fenestrated vessel development in the brain. *Elife* **10**:e64295.
doi:10.7554/elife.64295

Pfaffl MW. 2001. A new mathematical model for relative quantification in real-time RT–PCR.
Nucleic Acids Res **29**:e45–e45. doi:10.1093/nar/29.9.e45

Phng LK, Gerhardt H. 2009. Angiogenesis: A Team Effort Coordinated by Notch. *Dev Cell*
16:196–208. doi:10.1016/j.devcel.2009.01.015

Phng LK, Stanchi F, Gerhardt H. 2013. Filopodia are dispensable for endothelial tip cell
guidance. *Development* **140**:4031--4040.

Phng L-K, Stanchi F, Gerhardt H. 2013. Filopodia are dispensable for endothelial tip cell
guidance. *Development* **140**:4031–4040. doi:10.1242/dev.097352

Prasad GV, Coury LA, Finn F, Zeidel ML. 1998. Reconstituted aquaporin 1 water channels
transport CO2 across membranes. *J Biol Chem* **273**:33123–6. doi:10.1074/jbc.273.50.33123

Preston GM, Carroll TP, Guggino WB, Agre P. 1992. Appearance of water channels in Xenopus
oocytes expressing red cell CHIP28 protein. *Science* **256**:385–7.
doi:10.1126/science.256.5055.385

Rehn K, Wong KS, Balciunas D, Sumanas S. 2011. Zebrafish enhancer trap line recapitulates
embryonic aquaporin 1a expression pattern in vascular endothelial cells. *Int J Dev Biol*
55:613–618. doi:10.1387/ijdb.103249kp

Ridley AJ. 2011. Life at the Leading Edge. *Cell* **145**:1012–1022. doi:10.1016/j.cell.2011.06.010

Saadoun S, Papadopoulos MC, Hara-Chikuma M, Verkman AS. 2005. Impairment of
angiogenesis and cell migration by targeted aquaporin-1 gene disruption. *Nature* **434**:786–
792. doi:10.1038/nature03460

Sato Y, Shigematsu M, Shibata-Kanno M, Maejima S, Tamura C, Sakamoto H. 2023. Aquaporin
regulates cell rounding through vacuole formation during endothelial-to-hematopoietic
transition. *Development* **150**. doi:10.1242/dev.201275

Sauteur L, Affolter M, Belting H-G. 2017. Distinct and redundant functions of Esama and VE-
cadherin during vascular morphogenesis. *Development* **144**:1554–1565.
doi:10.1242/dev.140038

Siekmann AF, Affolter M, Belting H-G. 2013. The tip cell concept 10 years after: New players
tune in for a common theme. *Exp Cell Res* **319**:1255–1263.

Son S, Kang JH, Oh S, Kirschner MW, Mitchison TJ, Manalis S. 2015. Resonant microchannel
volume and mass measurements show that suspended cells swell during mitosis. *J Cell Biol*
211:757–763. doi:10.1083/jcb.201505058

Soria LR, Fanelli E, Altamura N, Svelto M, Marinelli RA, Calamita G. 2010. Aquaporin-8-
facilitated mitochondrial ammonia transport. *Biochem Biophys Res Commun* **393**:217–221.
doi:10.1016/j.bbrc.2010.01.104

Stewart MP, Helenius J, Toyoda Y, Ramanathan SP, Muller DJ, Hyman AA. 2011. Hydrostatic
pressure and the actomyosin cortex drive mitotic cell rounding. *Nature* **469**:226–230.
doi:10.1038/nature09642

Stroka KM, Jiang H, Chen S-H, Tong Z, Wirtz D, Sun SX, Konstantopoulos K. 2014. Water
permeation drives tumor cell migration in confined microenvironments. *Cell* **157**:611–623.
doi:10.1016/j.cell.2014.02.052

Talbot K, Kwong RWM, Gilmour KM, Perry SF. 2015. The water channel aquaporin-1a1
facilitates movement of CO_2 and ammonia in zebrafish (Danio rerio) larvae. *J Exp Biol*
218:3931–3940. doi:10.1242/jeb.129759

Thisse C, Thisse B. 2007. High-resolution in situ hybridization to whole-mount zebrafish
embryos. *Nat Protoc* **3**:59–69. doi:10.1038/nprot.2007.514

Tingaud-Sequeira A, Calusinska M, Finn RN, Chauvigné F, Lozano J, Cerdà J. 2010. The
zebrafish genome encodes the largest vertebrate repertoire of functional aquaporins with dual
paralogy and substrate specificities similar to mammals. *BMC Evol Biol* **10**:38.
doi:10.1186/1471-2148-10-38

Wakayama Y, Fukuhara S, Ando K, Matsuda M, Mochizuki N. 2015. Cdc42 Mediates Bmp-
Induced Sprouting Angiogenesis through Fmnl3-Driven Assembly of Endothelial Filopodia in
Zebrafish. *Dev Cell* **32**:109–122. doi:10.1016/j.devcel.2014.11.024

Wang Y, Cohen J, Boron WF, Schulten K, Tajkhorshid E. 2007. Exploring gas permeability of
cellular membranes and membrane channels with molecular dynamics. *J Struct Biol* **157**:534–
544. doi:10.1016/j.jsb.2006.11.008

Wu DQ, Yang ZF, Wang KJ, Feng XY, Lv ZJ, Li Y, Jian ZX. 2017. AQP8 inhibits colorectal
cancer growth and metastasis by down-regulating PI3K/AKT signaling and PCDH7
expression. *Am J Cancer Res* **8**:266–279.

Zeidel ML, Ambudkar SV, Smith BL, Agre P. 1992. Reconstitution of functional water channels
in liposomes containing purified red cell CHIP28 protein. *Biochemistry* **31**:7436–7440.
doi:10.1021/bi00148a002

1056

Figure 1

Figure 2

Figure 3

A *aqp1a.1$^{+/rk28}$;aqp8a.1$^{+/rk29}$

B *aqp1a.1$^{rk28/rk28}$*

C *aqp8a.1$^{rk29/rk29}$*

D *aqp1a.1$^{rk28/rk28}$; aqp8a.1$^{rk29/rk29}$*

E

ISV length (μm) vs ISV position

- *aqp1a.1$^{+/rk28}$;aqp8a.1$^{+/rk29}$*
- *aqp1a.1$^{rk28/rk28}$*
- *aqp8a.1$^{rk29/rk29}$*
- *aqp1a.1$^{rk28/rk28}$; aqp8a.1$^{rk29/rk29}$*

K

% of embryos with at least one incomplete ISV (%)

Wildtype, *aqp1a.1$^{rk28/rk28}$*, *aqp8a.1$^{rk29/rk29}$*, *aqp1a.1$^{rk28/rk28}$; aqp8a.1$^{rk29/rk29}$*

F aISV — DLAV, NT, NC, DA, PCV; vISV — DLAV, NT, NC, DA, PCV

G *aqp1a.1$^{+/+}$;aqp8a.1$^{+/+}$* Tg(fli1a:H2B-EGFP) Tg(fli1:Lifeact-mCherry)

H *aqp8a.1$^{rk29/rk29}$* Tg(fli1a:H2B-EGFP) Tg(fli1:myr-mCherry)

I *aqp1a.1$^{rk28/rk28}$* Tg(fli1a:H2B-EGFP) Tg(fli1:Lifeact-mCherry)

J *aqp1a.1$^{rk28/rk28}$; aqp8a.1$^{rk29/rk29}$* Tg(fli1a:H2B-EGFP) Tg(fli1:Lifeact-mCherry)

L I II III IV V VII

- 75.20% Class I
- 15.79% Class II
- 1.16% Class III
- 2.23% Class IV
- 2.94% Class V
- 2.68% Class VI

n=1123 ISVs (ISVs no. 5-28)

Figure 4

Figure 5

A — aqp1a.1$^{+/+}$; aqp8a.1$^{+/+}$

mEmerald
Tg(fli1:myr-mCherry)   Tip cell volume

B — aqp1a.1$^{rk28/rk28}$; aqp8a.1$^{rk29/rk29}$

C

Volume (μm³)

0.0176

aqp1a.1$^{+/+}$;
aqp8a.1$^{+/+}$ aqp1a.1$^{rk28/rk28}$;
aqp8a.1$^{rk29/rk29}$

D

Rate of actin elongation (μm/s)

<0.0001

aqp1a.1$^{+/+}$;
aqp8a.1$^{+/+}$ aqp1a.1$^{rk28/rk28}$;
aqp8a.1$^{rk29/rk29}$

Figure 6

Membrane protrusion generated by actin polymerization and increased hydrostatic pressure.

Membrane protrusion generated by increased hydrostatic pressure.

Membrane protrusion generated by actin polymerization.