Combined forces of hydrostatic pressure and actin polymerization drive endothelial tip cell

migration and sprouting angiogenesis

Igor Kondrychyn[1], Liqun He[2], Haymar Wint[1], Christer Betsholtz[2, 3] and Li-Kun Phng[1, 4]

[1] Laboratory for Vascular Morphogenesis, RIKEN Center for Biosystems Dynamics Research,

Kobe 650-0047, Japan

[2] Department of Immunology, Genetics and Pathology, Rudbeck Laboratory, Uppsala University,

75185 Uppsala, Sweden

[3] Department of Medicine Huddinge, Karolinska Institutet, 14157 Huddinge, Sweden

[4] Lead contact

* Correspondence: likun.phng@riken.jp

SUMMARY

Cell migration is a key process in the shaping and formation of tissues. During sprouting

angiogenesis, endothelial tip cells invade avascular tissues by generating actomyosin-dependent

forces that drive cell migration and vascular expansion. Surprisingly, ECs can still invade if actin

polymerization is inhibited. In this study, we show that endothelial tip cells employ an alternative

mechanism of cell migration that is dependent on Aquaporin (Aqp)-mediated water inflow and

increase in hydrostatic pressure. In the zebrafish, ECs express *aqp1a.1* and *aqp8a.1* in newly

formed vascular sprouts in a VEGFR2-dependent manner. Aqp1a.1 and Aqp8a.1 loss-of-function

studies show an impairment in intersegmental vessels formation because of a decreased capacity

of tip cells to increase their cytoplasmic volume and generate membrane protrusions, leading to

delayed tip cell emergence from the dorsal aorta and slower migration. Further inhibition of actin

polymerization resulted in a greater decrease in sprouting angiogenesis, indicating that ECs

employ two mechanisms for robust cell migration *in vivo*. Our study thus highlights an important

role of hydrostatic pressure in tissue morphogenesis.

Keywords

Endothelial cell migration, Sprouting angiogenesis, Aquaporin, Water flow, Hydostatic Pressure,

Actin, Zebrafish

INTRODUCTION

The blood vascular system is a dynamic, multicellular tissue that adapts to the metabolic demands of development, growth and homeostasis by altering its pattern and morphology. Macroscopic transformation in the vascular network is achieved by microscopic changes in endothelial cell (EC) behaviours. For example, vascular expansion through sprouting angiogenesis requires the coordination of EC migration, proliferation, anastomosis, lumen formation and cell rearrangements, all of which require specific cell shape changes. At the initial phase of sprouting angiogenesis, endothelial tip cells generate numerous membrane protrusions such as filopodia and lamellipodia that drive migration and anastomosis (Figueiredo et al., 2021; Gerhardt et al., 2003; Phng et al., 2013; Sauteur et al., 2017). During tubulogenesis, apical membranes deform by generating inverse blebs to expand lumens so that blood can be transported efficiently through the blood vascular network (Gebala et al., 2016). However, following vessel perfusion, ECs become less dynamic and instead develop mechanoresistance to the deforming forces of blood pressure to maintain vessel morphology (Kondrychyn et al., 2020). EC shape is therefore dynamic and changes depending on the morphogenetic state of vascular development.

It is well established that localized remodelling of actin cytoskeleton and non-myosin II contractility are instrumental in driving cell shape changes during tissue morphogenesis (Clarke and Martin, 2021; Munjal and Lecuit, 2014; Murrell et al., 2015). More recently, there is an accumulating body of work implicating a role of water flow and the resulting increase in hydrostatic pressure as another mechanical mechanism of cell shape change (Choudhury et al., 2022; Chugh et al., 2022; Li et al., 2020). For example, in the osmotic engine model of cell migration, polarized distribution of ion channels and transporters generate an osmotic gradient that drives water inflow at the leading edge and outflow at the rear, leading to forward translocation of tumour cells in confined microenvironment (Stroka et al., 2014). Additionally, hydrostatic pressure from the extracellular environment can drive tissue morphogenesis, as demonstrated during mouse blastocyst formation (Chan et al., 2019; Dumortier et al., 2019), development of the otic vesicle in the zebrafish (Mosaliganti et al., 2019) and blood vessel lumen expansion (Gebala et al., 2016). Hydrostatic pressure can therefore generate sufficient forces to shape tissues and organs to their proper form and size during development.

We have previously discovered that EC migration persists after the inhibition of actin polymerization and in the absence of filopodia to generate new blood vessels in the zebrafish

(Phng et al., 2013). This observation suggests the existence of an alternative mechanism of
migration independent of actin polymerization. In this study, we sought to determine whether
hydrostatic pressure regulates EC migration during sprouting angiogenesis in the zebrafish by
investigating the function of Aquaporins (Aqp), which are transmembrane water channels that
increase water permeability of cell membranes to promote transcellular water flow (Day et al.,
2014; Farinas et al., 1997; Kozono et al., 2002; Preston et al., 1992). We discovered that ECs of
newly formed intersegmental vessels (ISV) express *aqp1a.1* and *aqp8a.1* mRNA, and observed
the enrichment of Aqp1a.1 and Aqp8a.1 proteins in the leading edge of migrating tip cells.
Detailed single cell analyses showed that endothelial tip cells lacking *aqp1a.1* and *aqp8a.1*
expression have reduced cell volume, membrane protrusions and migration capacity. As a result,
there is defective sprouting angiogenesis and ISV formation in *aqp1a.1;aqp8a.1* double mutant
zebrafish. Notably, when actin polymerization is inhibited in *aqp1a.1*;*aqp8a.1* double mutant
zebrafish, there is a greater impairment in EC migration and sprouting angiogenesis, demonstrating
the additive function of actin polymerization and hydrostatic pressure (that is elevated by water
inflow) in generating membrane protrusions to drive EC migration *in vivo*.

RESULTS

***aqp1a.1* and *aqp8a.1* are differentially expressed by tip and stalk cells during sprouting**
angiogenesis
Of the 18 *aquaporin* genes that are expressed in zebrafish (Tingaud-Sequeira et al., 2010), *aqp1a.1*
and *aqp8a.1* mRNA have been detected in blood endothelial cells (ECs) (Koun et al., 2016; Rehn
et al., 2011). We confirmed the endothelial expression of *aqp1a.1* and *aqp8a.1* at 30 hours post
fertilization (hpf) and 2-, 3- and 4-days post fertilization (dpf) by whole-mount *in situ* hybridization
(WISH, Fig. S1). The two endothelial *aqp* genes, however, differ in spatial expression. While
aqp1a.1 is widely expressed in blood vessels of the head and trunk (dorsal aorta (DA), caudal
artery (CA), ISVs, dorsal longitudinal anastomotic vessels (DLAV) and caudal vein plexus) at 30
hpf, *aqp8a.1* expression is absent in cerebral blood vessels and is restricted to the DA, CA and the
ventral regions of ISVs in the trunk. After 1 dpf, *aqp8a.1* expression expands to the entire ISV and
DLAV. Additionally, the expression of *aqp1a.1* and *aqp8a.1* gradually decreases in the DA and
CA so that both are absent by 4 dpf. Single cell RNA sequencing (scRNAseq) of ECs isolated at

24 hpf, 34 hpf and 3 dpf further confirmed the endothelial expression of *aqp1a.1* and *aqp8a.1*
mRNA (Fig. S2 A - F), revealed differential expression of *aqp1a.1* and *aqp8a.1* mRNA in distinct
endothelial clusters and highlighted higher *aqp1a.1* transcript expression in all endothelial
subtypes compared to *aqp8a.1* (Fig. S2 E - H).

We next examined the expression pattern of *aqp1a.1* and *aqp8a.1* at the beginning of
sprouting angiogenesis at higher resolution using RNAscope and confocal microscopy. At 20 hpf,
aqp1a.1 mRNA is highly expressed in the DA with little expression in the PCV (Fig. 1A, C and
Fig. S3). Closer inspection uncovered heterogeneous expression in the DA, with higher expression
in ECs that would be specified as tip cells (Fig. 1A, C and Fig. S3B). Indeed, tip cells of newly
formed vascular sprouts express higher levels of *aqp1a.1* compared to adjacent ECs in the DA at
20 hpf (Fig. 1A – C, Fig. S3B) and the majority of ISVs (61%) display higher *aqp1a.1* expression
in tip cells than in trailing stalk cells at 22 hpf (Fig. 1G). *Aqp8a.1* expression is at first largely
absent in the DA at 20 hpf (Fig. 1B, D and Fig. S3C) but becomes detectable at 22hpf (Fig. 1E and
F). Interestingly, unlike *aqp1a.1*, *aqp8a.1* expression is undetectable in tip cells of emerging ISVs
at 20 hpf. At 22 hpf, it is expressed in 22 out of 23 tip cells analyzed with 78% of ISVs exhibiting
higher *aqp8a.1* mRNA expression in stalk cells than tip cells (Fig. 1G). These observations
highlight differential expression pattern of *aqp1a.1* and *aqp8a.1* in newly formed vascular sprouts,
with *aqp1a.1* expression enriched in tip cells and *aqp8a.1* expression in stalk cells.

VEGF-VEGFR2 signalling induces the expression of *aqp1a.1* and *aqp8a.1* mRNA
We next explored whether *aqp1a.1* and *aqp8a.1* expression is regulated by VEGFR2 signalling,
which is highly activated in tip cells during sprouting angiogenesis (Gerhardt et al., 2003;
Siekmann et al., 2013). To inhibit VEGFR2 activity, embryos were treated with 1 µM ki8751 from
20 hpf. Relative quantitative polymerase chain reaction (qPCR) experiments showed that *aqp1a.1*
and *aqp8a.1* mRNA expression was suppressed by 52% ($p < 0.0001$) and 17% ($p = 0.0028$),
respectively, after 6 hours of ki8751 treatment when compared to DMSO-treated embryos (Fig.
1H). These results demonstrate that VEGFR2 activity upregulates the expression of *aqp1a.1* and
aqp8a.1. To examine whether VEGFR2 activity also regulates Aquaporin expression in human
ECs, we treated cultured human aortic endothelial cells (HAECs) with ki8751 for 6 hours and
analyzed *AQP1* expression by qPCR. *AQP8* expression was not examined since it is not expressed
in HAECs (data not shown). Inhibition of VEGFR2 activity with 1 nM and 10 nM ki8751

decreased *AQP1* mRNA expression by 31% ($p = 0.0023$, Fig. 1I) and 40% ($p = 0.1032$, Fig. 1I),

respectively, demonstrating cross-species conservation of VEGFR2 activity in inducing *Aqp1*

mRNA expression.

Aqp1a.1 and Aqp8a.1 proteins are enriched at the leading edge of migrating tip cells

To gain a better understanding of where within the cell Aquaporin protein functions, we generated

stable *Tg(fli1ep:aqp1a.1-mEmerald)^{rk30}* and *Tg(fli1ep:aqp8a.1-mEmerald)^{rk31}* zebrafish lines in

which ECs express Aqp1a.1 or Aqp8a.1 tagged with mEmerald, respectively. At 25 hpf, we

observed an accumulation of Aqp1a.1 (Fig. 2A) and Aqp8a.1 (Fig. 2B) in mCherryCAAX-positive

membrane clusters in the cytoplasm and filopodia of tip cells. Notably, many of the Aqp1a.1- or

Aqp8a.1-positive membrane clusters are located at the migrating front of the tip cell. Time-lapse

imaging capturing the process of ISV formation between 24 to 30 hpf further showed sustained

enrichment of Aqp1a.1 and Aqp8a.1 in clusters at the front of the tip cell as well as in filopodia

(Fig. 2C and D, Movie S1 and S2), suggesting that Aqp1a.1 and Aqp8a.1 may promote localized

water flux at the migrating edge of tip cells.

Loss of Aqp1a.1 and Aqp8a.1 function leads to defective trunk blood vessel formation

We subsequently proceeded to generate zebrafish *aqp1a.1* and *aqp8a.1* mutants to investigate the

function of Aquaporin-mediated water flow in blood vessel development. Using CRISPR/Cas9

genome editing, we targeted the first exons of *aqp1a.1* and *aqp8a.1* genes. The CRISPR-

generating allele *aqp1a.1^{rk28}* carries a 14-bp deletion and an 8-bp insertion in the 5' end of the gene

leading to a premature termination codon at amino acid 76 after 1 missense amino acid (Fig. S4A-

B). The *aqp8a.1^{rk29}* allele carries a 4-bp deletion in the 5' end of the gene that leads to a frameshift

after Thr15 and premature termination codon at amino acid 73 after 58 missense amino acids (Fig.

S5A-B). Using whole-mount *in situ* hybridization and qPCR analysis, we demonstrated that

aqp1a.1 and *aqp8a.1* mRNA expression is significantly decreased in *aqp1a.1$^{rk28/rk28}$* (Fig. S4C-D)

and *aqp8a.1$^{rk29/rk29}$* (Fig. S5C-D) zebrafish embryos, respectively, suggesting a rapid degradation

of mutant mRNAs and as a result, the loss of Aquaporin protein expression.

Although *aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$* embryos look morphologically normal when

compared to wildtype or *aqp1a.1$^{+/rk28}$;aqp8a.1$^{+/rk29}$* embryos (Fig. S8A-D), we observed defects

in blood vessel formation and patterning at 1 to 3 dpf. Compared to 28 hpf control embryos whose

ISVs have reached the dorsal roof of the neural tube and started to form the DLAV, ISVs of *aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$* embryos are much less developed (Fig. 3A and D) with significantly reduced length (Fig. 3E and Fig. S6H and K). Single deletion of *aqp1a.1* (Fig. 3B and Fig. S6F and I) or *aqp8a.1* (Fig. 3C and Fig. S6G and J) also led to defective ISV formation, although the defects are less severe compared to the loss of both *aqp1a.1* and *aqp8a.1* expression. Comparison of ISV length in *aqp1a.1$^{rk28/rk28}$*, *aqp8a.1$^{rk29/rk29}$* and *aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$* embryos shows that the loss of *aqp1a.1* leads to a greater decrease in ISV length than the loss of *aqp8a.1* (Fig. 3E, Fig. S6I and J), and that the loss of both *aqp1a.1* and *aqp8a.1* results in the greatest reduction. These observations indicate an additive effect of Aqp1a.1 and Aqp8a.1 function in blood vessel development and implicate Aqp1a.1 as the more dominant Aquaporin protein in EC function, in line with the higher expression of *aqp1a.1* mRNA in tip cells.

We next examined the embryos at 3 dpf, a stage when arterial ISVs (aISVs) and venous (vISVs) are fully established and lumenized, connecting the DLAV to either the DA or PCV (Fig. 3F and G), respectively. Such fully established ISVs are significantly reduced in embryos when *aqp1a.1*, *aqp8a.1* or both *aqp1a.1* and *aqp8a.1* are deleted. Instead, these embryos display an increased number of aISVs and vISVs that fail to establish a connection to the DLAV, DA or PCV (Fig. 3H – J). The number of incompletely formed or truncated ISVs per embryo in *aquaporin* mutants was variable, between 1 to 6, 1 to 16 and 1 to 22 truncated ISVs in *aqp8a.1$^{rk29/rk29}$*, *aqp1a.1$^{rk28/rk28}$* and *aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$* embryos, respectively (Fig. S6A-D). We also found that truncated ISVs appear with higher frequency in the zebrafish trunk (ISV number 5 to 15) rather than in the tail (ISV number 16 to 27, Fig. S6E). Detailed analysis revealed that most of the truncated ISVs are vISVs that are connected to the PCV but not the DLAV (Class I in Fig. 3K), followed by the absence of aISV or vISV (Class II in Fig. 3K, Fig. S7A and B). A small fraction of the truncated vessels are aISVs that are not connected to the DA (Class IV in Fig. 3K, Fig. S7D) or DLAV (Class V in Fig. 3K, Fig. S7E), or vISVs that are not connected to the DLAV (Class VI in Fig. 3K , Fig. S7F). At some regions, there is a truncated aISV that is not connected to the DA and an incomplete vISV that is not connected to the DLAV (Class III in Fig. 3K, Fig. S7C).

As *aqp1a.1* is expressed in the brain vasculature during development (Fig. S1), we examined whether cerebral vasculature is also affected in *aqp1a.1$^{rk28/rk28}$* embryos. At 4 dpf, we found that the length of the dorsal longitudinal vein (DLV), a main blood vessel that supplies blood to the choroid plexus (Bill and Korzh, 2014) was reduced in *aqp1a.1$^{rk28/rk28}$* embryos (p = 0.0098,

Fig. S9A, B and E). Also, in 2 out of 21 *aqp1a.1^{rk28/rk28}* embryos, the DLV was undeveloped (Fig.

S9C, D and F), a defect that is also observed in 50% of *vegfab^{-/-}* and *vegfc^{-/-}* mutant zebrafish

(Parab et al., 2021).

In summary, our analyses show that in the absence of Aqp1a.1 and/or Aqp8a.1 function,

there is an initial delay in ISV formation during primary sprouting angiogenesis. By 3 dpf, some

ISVs are completely formed (Fig. S8K and M), connecting the DA or PCV to the DLAV. However,

the number of completely formed ISVs is significantly decreased, with the loss of Aqp1a.1 having

a more severe impact compared to the loss of Aqp8a.1, and the combined loss of Aqp1a.1 or

Aqp8a.1 having a greater deleterious effect.

Aquaporins promote endothelial tip cell protrusion and migration

To gain a better understanding of why ISVs do not develop normally, we assessed EC number. By

counting the number of nuclei in aISVs and vISVs, we found a significant reduction in EC number

per vessel in 2 and 3 dpf *aqp1a.1^{rk28/rk28};aqp8a.1^{rk29/rk29}* embryos when compared to wildtype

embryos (Fig. S10A and B), suggesting that the defective formation of ISVs may result from

insufficient EC number. Quantification of endothelial mitotic events between 21 and 30 hpf shows

that there is a similar number of tip and stalk cell division in *aqp1a.1^{+/rk28};aqp8a.1^{+/rk29}* embryos

and *aqp1a.1^{rk28/rk28};aqp8a.1^{rk29/rk29}* embryos (Fig. S10C), indicating that decreased EC division is

not the cause of reduced EC number in ISVs. We next explored whether EC migration from the

DA is impaired by examining tip cell behaviour from 21 – 22hpf, when sprouting angiogenesis

begins. In *aqp1a.1^{+/rk28};aqp8a.1^{+/rk29}* embryos, tip cells emerge from the DA between 20 and 24

hpf (Fig. 4A) and migrate dorsally between the somite boundaries over the notochord and neural

tube to form primary ISVs (Fig. 4B, Movie S3). However, in *aqp1a.1^{rk28/rk28};aqp8a.1^{rk29/rk29}*

embryos, tip cell nuclei emerge between 22 and 28 hpf (Fig. 4A). Furthermore, in some

aqp1a.1^{rk28/rk28};aqp8a.1^{rk29/rk29} embryos, tip cell membrane protrusions do not extend dorsally in

the direction of migration but retract back to the DA so that a primary ISV does not form at this

location (Fig. 4C, Movie S4), potentially giving rise to the Class II phenotype at 3 dpf (Fig. 3K

and Fig. S7B). These observations indicate that tip cell protrusion from the DA is significantly

delayed in the absence of Aqp1a.1 and Aqp8a.1 function.

We have previously demonstrated that filopodia facilitate EC migration by serving as a

template for lamellipodia formation (Phng et al., 2013). Lamellipodia frequently emanate laterally

from stable filopodia at the leading edge of tip cells, leading to rapid expansion of the filopodium, increase in volume and the formation of a stable protrusion in the direction of migration (Phng et al., 2013). This was similarly observed in *aqp1a.1$^{+/rk28}$;aqp8a.1$^{+/rk29}$* embryos (Fig. 4D and Movie S5) but not in *aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$* embryos (Fig. 4E and Movie S6). Comparison of filopodia dynamics in wildtype (Fig. 4F), *aqp1a.1$^{+/rk28}$;aqp8a.1$^{+/rk29}$* and *aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$* (Fig. 4G) embryos revealed a significant reduction in filopodia number (wildtype, 18.73 ± 5.1/100μm; *aqp1a.1$^{+/rk28}$;aqp8a.1$^{+/rk29}$*, 20.4 ± 6.5/100μm; *aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$*, 9.8 ± 5.5/100μm; Fig. 4H) and length (wildtype, 5.5 ± 2.8 μm; *aqp1a.1$^{+/rk28}$;aqp8a.1$^{+/rk29}$*, 6.0 ± 3.4 μm; *aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$*, 4.5 ± 2.2 μm; Fig. 4I) in ECs lacking both Aqp1a.1 and Aqp8a.1 function that can be accounted for by a decrease in actin polymerization (Fig. 4J). By tracking Lifeact-mCherry signal in filopodia, we determined that the average elongation rate of actin as 0.055 ± 0.013 μm per second in wildtype embryos and that this is significantly decreased by 47% to 0.029 ± 0.011 μm per second ($p < 0.0001$) in *aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$* embryos. Analysis of time-lapse images of tip cell behaviour further showed slower tip cell membrane expansion at the leading edge (Fig. 4K), decreased nuclei displacement (Fig. 4L) and a significant reduction in tip cell migration velocity (Fig. 4M) in *aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$* embryos. To determine whether the slower EC migration is caused by a loss of front-rear polarity, we examined Golgi position relative to the nucleus in wildtype, *aqp1a.1$^{rk28/rk28}$* and *aqp8a.1$^{rk29/rk29}$* embryos at 24 - 26 hpf (Fig. S10D). This analysis shows that tip cell polarity is unperturbed in the absence of Aqp1a.1 or Aqp8a.1 function.

In conclusion, our findings demonstrate that Aqp1a.1 and Aqp8a.1 regulate sprouting angiogenesis by promoting endothelial tip cell emergence from the DA, the formation and expansion of membrane protrusions and EC migration.

Tip cell volume regulation depends on Aquaporin-mediated water influx

Aquaporin channels permit bidirectional flow of water across the plasma membrane, with the direction of flow determined by the osmotic gradient between the extracellular environment and cell cytoplasm (Agre et al., 2002). We next sought to determine whether water flows in or out of tip cells during sprouting angiogenesis. We hypothesize that water flux across the cell membrane can lead to changes in tip cell volume, with water influx increasing the volume while outflow decreases. To quantify tip cell volume in the absence of Aquaporin function, we injected a plasmid

encoding *kdrl:mEmerald* into 1-cell stage *aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$* embryos to label ECs in

a mosaic manner. We also investigated the effects of increasing Aqp1a.1 expression on tip cell

volume by injecting a plasmid encoding *Aqp1a.1-P2A-EGFP* into wildtype embryos. As control,

wildtype embryos were injected with the plasmid encoding *kdrl:mEmerald*. mEmerald/EGFP-

positive tip cells were imaged and their volume measured at 24 to 25 hpf from 3D reconstructions

of labelled cells. Wildtype ECs are on average 964.6 ± 270.2 μm³ in size (Fig. 5A and D). In the

absence of both Aqp1a.1 and Aqp8a.1 function (Fig. 5B), tip cell volume significantly decreased

by 26% to 709.8 ± 293.5 μm³ (*p* = 0.015, Fig. 5D) when compared to wildtype tip cells. The

overexpression of Aqp1a.1 increased tip cell volume by 36% to 1313 ± 527.4 μm³ when compared

to wildtype tip cells (Fig. 5C and D, *p* = 0.0069) and by 85% when compared to

aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$ tip cells (*p* < 0.0001). These results therefore support water influx

as the direction of water flow in tip cells during migration, and that this is mediated through

Aqp1a.1 and Aqp8a.1.

In summary, our results demonstrate a role of endothelial Aquaporins in promoting water

influx to increase endothelial tip cell volume and cytoplasmic hydrostatic pressure, as well as

accelerate actin polymerization in filopodia (Fig. 4J).

The anion channel, SWELL1, promotes EC migration and sprouting angiogenesis

As the direction of water flow is dictated by an osmotic gradient, with water flowing from low to

high osmolarity, we next examined the function of ion channels in establishing an osmotic gradient.

SWELL1, or LRRC8A, is a component of volume-regulated anion channel (VRAC) that mediates

ions, especially chloride anions, and osmolyte efflux (Qiu et al., 2014; Voss et al., 2014). Recently,

SWELL1 has been demonstrated to polarize at the trailing edge of migrating breast cancer cells to

direct water efflux and confer confined migration direction (Zhang et al., 2022). In the zebrafish,

scRNAseq data shows that the gene encoding SWELL1, *lrrc8aa*, is expressed in a subset of ECs,

with 12.32% of ECs coexpressing *lrrc8aa* and *aqp1a.1* (Fig. 6A) and 10.16% coexpressing *lrrc8aa*

and *aqp8a.1* (Fig. 6B). To determine whether *lrrc8aa* is expressed in endothelial tip cells, we

performed RNAscope and confocal microscopy. Although *lrrc8aa* is expressed in surrounding

somites and notochord (Fig. 6C), we could also detect its expression in endothelial tip cells (Fig.

6D and E), suggesting that SWELL1 can establish an osmotic gradient in the vicinity of the

developing ISVs and within tip cells to direct water flow. To address the function of SWELL1-

generated osmotic gradient, we disrupted its activity by treating zebrafish with 5μM DCPIB
(Gunasekar et al., 2022) for 6 hours from 20 hpf and observed a similar impairment in ISV
formation (Fig. 6F and G) as in *aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$* embryos (Fig. 3D and E). SWELL1
inhibition resulted in shorter ISVs formed at 26 to 27 hpf (Fig. 6H) due to impaired leading edge
displacement of tip cells (Fig. 6I) and decreased tip cell velocity (Fig. 6J). Further analysis of tip
cell morphology showed a decrease in the number (Fig. 6K) and length (Fig. 6L) of filopodia
formed and compromised elongation of tip cells (Fig. 6M and N, Movie S7) in DCPIB-treated
embryos. In summary, these results suggest that chloride efflux through SWELL1 establishes an
osmotic gradient, either within ECs or between EC and the surrounding interstitial tissue, to direct
water flow and thereby modulate EC migration and sprouting angiogenesis.
290

Additive effects of actin polymerization and water influx in driving EC migration and
sprouting angiogenesis
We have previously observed that ECs are still able to migrate and form ISVs after the inhibition
of actin polymerization (Phng et al., 2013). In these experiments, embryos were treated with a low
concentration of Latrunculin B (Lat. B), an inhibitor of actin polymerization, that resulted in the
suppression of filopodia formation. Under such conditions, ECs can generate small membrane
protrusions and migrate in a directed manner at a reduced speed. How ECs are still able to migrate
after the inhibition of actin polymerization was unclear. As our current study demonstrates a role
of water inflow and hydrostatic pressure as another mechanism of cell migration, we hypothesize
that ECs employ water influx to migrate when actin polymerization is compromised and that the
depletion of both water influx and actin polymerization will result in a greater inhibition of EC
migration.
To test this hypothesis, we examined ISV development in embryos with decreased actin
polymerization and water inflow by treating *aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$* embryos with Lat. B.
At 28 hpf, ISVs of wildtype embryos treated with DMSO have reached the dorsal roof of the neural
tube (Fig. 7A) and show an average length of 104.1 ± 13.47 μm (Fig. 7E). The treatment of
wildtype embryos with 0.08 μg/ml Lat. B from 20 to 28 hpf (Fig. 7B) resulted in a significant
decrease in ISV length (61.53 ± 23.27 μm) compared to control embryos ($p < 0.0001$, Fig. 7E). A
similar decrease in ISV length was also observed in *aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$* embryos treated
with DMSO (63.75 ± 30.24 μm, Fig. 7C and E). When *aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$* embryos

were treated with 0.08µg/ml Lat. B from 20 to 28 hpf, there was a greater decrease in the length

of ISV formed at 28 hpf (43.10 ± 24.42 µm, Fig. 7D and E) when compared to the inhibition of

actin polymerization (p = 0.0028) or depletion of water influx (p < 0.0001) alone (Movie S8).

These results therefore demonstrate additive functions of actin polymerization and water influx in

driving EC migration during sprouting angiogenesis (Fig. 7F).

DISCUSSION

In this study, we demonstrate that endothelial tip cells utilize two mechanisms of migration

concurrently – actin polymerization and hydrostatic pressure - during sprouting angiogenesis in

the zebrafish. The increase in hydrostatic pressure in endothelial tip cells is generated by Aqp1a.1-

and Aqp8a.1-mediated water inflow. By performing detailed expression analyses, we showed that

aqp1a.1 and *aqp8a.1* are differentially expressed in newly formed vascular sprouts. *Aqp1a.1* is

expressed earlier in the DA than *aqp8a.1* during development and in ECs destined to be tip cells.

In newly formed ISVs, endothelial tip cells express higher *aqp1a.1* mRNA levels while stalk cells

express higher *aqp8a.1* mRNA levels. In zebrafish harbouring mutations in both *aqp1a.1* and

aqp8a.1, there is delayed tip cell protrusion from the DA, decreased tip cell migration and

increased occurrence of truncated ISVs at 3 dpf. We further discovered that *aqp1a.1* and *aqp8a.1*

expression is upregulated by VEGFR2 activity, suggesting that VEGFA-VEGFR2 signalling

employs Aquaporin function as one mechanism to drive sprouting angiogenesis.

Several lines of evidence implicate Aquaporins in facilitating water flow into endothelial

tip cells during sprouting angiogenesis. First, timelapse imaging showed polarized localization of

Aqp1a.1 and Aqp8a.1 proteins at the leading edge of migrating tip cells, suggesting that water flow

occurs locally at the migrating front of the cell. Secondly, single cell analyses revealed Aqp1a.1-

and Aqp8a.1-deficient tip cells have reduced cell volume, impaired expansion of cell membranes

at the leading edge and decreased generation of stable cell protrusions. These observations

implicate water flow into tip cells, leading to increased cell volume. This is confirmed by the

overexpression of Aqp1a.1 in tip cells that resulted in increased cell volume. Due to the

incompressible property of water within an enclosed compartment such as the cell, water influx

will lead to elevation in hydrostatic pressure. At the leading edge of tip cells, increased hydrostatic

pressure would expand and deform membranes to generate protrusions. Additionally, water influx

can reduce cytoplasmic viscosity and increase spacing between the plasma membrane and the

underlying actin cytoskeleton, promoting actin monomer diffusion, actin polymerization and the

generation of membrane protrusions to facilitate cell migration (Boer et al., 2023; Loitto et al.,

2009).

A key finding in this study is the demonstration that endothelial tip cells employ hydrostatic

pressure as a second mechanism of cell migration *in vivo*. In a previous study, we discovered that

ECs continue to migrate when actin polymerization is inhibited and in the absence of filopodia

(Phng et al., 2013). Here, we show that hydrostatic pressure can generate force for EC migration

in the absence of actin-based force generation. When both mechanisms are perturbed, there is a

greater impairment in EC migration (Fig. 7D and E). This finding highlights the importance of

hydrostatic pressure as an additional mechanism that reinforces actin-based cell shape changes and

motility to ensure robust migration and formation of new blood vessels in complex three-

dimensional environments. Such dual mode migration is advantageous when blood vessels

develop in physically confined spaces such as in the zebrafish trunk, where ECs must emerge from

the DA and migrate between somite boundaries and over tissues such as the notochord and neural

tube to form ISVs. Such dependence on water flow in cell migration under confined

microenvironment has been reported in human MDA-MB-231 breast cancer cells and mouse S180

sarcoma cells, which show cell migration persistence in narrow channels after inhibition of actin

polymerization or myosin II-mediated contractility (Stroka et al., 2014). In this context, AQP5

mediates water flow and the decrease or increase in AQP5 expression suppresses or enhances,

respectively, cancer cell migration (Chae et al., 2008; Jung et al., 2011; Stroka et al., 2014).

Our findings on the endothelial function of zebrafish Aqp1a.1, which is homologous to

mammalian AQP1, corroborate previous studies demonstrating a function of AQP1 in promoting

cell migration. In tumour bearing AQP1-null mice, there is reduced tumour angiogenesis and

AQP1-deficient ECs display reduced migration in an *in vitro* wound healing assay (Maltaneri et

al., 2020; Saadoun et al., 2005). In chick, AQP1 expression levels affect neural crest cell migration

speed and direction by regulating filopodia length and stability (McLennan et al., 2019). The

regulation of membrane protrusions has also been demonstrated for other Aquaporin proteins. For

example, AQP9 weakens membrane-cytoskeleton anchorage and promote formation of membrane

protrusions such as filopodia and blebs (Karlsson et al., 2013; Loitto et al., 2007). We also

demonstrate for the first time a role of Aqp8a.1 in promoting EC migration and sprouting

angiogenesis in the zebrafish. However, its mammalian ortholog, AQP8, was shown to inhibit

colorectal cancer cell lines *in vitro* (Wu et al., 2017), suggesting cell- and context-dependent

function of AQP8 in the regulation of cell migration. The role of Aqp1a.1 and Aqp8a.1 has also

been recently investigated in the zebrafish. However, the authors did not report defects in EC

migration but instead observed altered ISV diameter in Aqp1a.1 and Aqp8a.1 knockout and

overexpression experiments (Chen et al., 2024). While we corroborate a decrease in the diameter

of aISVs and vISVs in *aqp1a.1$^{rk28/rk28}$* and *aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$* zebrafish, we observed

a slight increase in the diameter in *aqp8a.1$^{rk29/rk29}$* zebrafish at 2 dpf (Fig. S11A). Furthermore,

unlike described in Chen et al, we could not detect a difference in ISV diameter between control

and Aqp1a.1- or Aqp8a.1-overexpressing ECs (Fig. S11B). The disparity in phenotypes observed

between our groups are likely a result of differences in the *aqp1a.1* and *aqp8a.1* mutations and the

level of Aqp1a.1 or Aqp8a. overexpression generated in our respective zebrafish lines.

A central determinant of water movement is the osmotic gradient across the membrane,

with water flowing in the direction of higher osmotic concentration to build up hydrostatic pressure.

Given a crucial role of water flow in regulating cell volume, shape and migration, it is also

important to understand the function and distribution of ion channels, exchangers and transporters

and water channels that generate an osmotic gradient across the cell. This is reflected in a growing

number of studies demonstrating a role of ion transporters in cell migration. In neutrophils, the

increase in cell volume and potentiation of cell migration depend on the sodium-proton exchanger

1 (NHE1) and the chloride-bicarbonate exchanger 2 (AE2) (Nagy et al., 2023). In T cells,

chemokine-induced migration depends on ion influx via SLC12A1, and water influx via AQP3

(Boer et al., 2023). In tumour cells migrating in narrow channels, the establishment of a polarized

distribution of Na^+/H^+ pumps and Aquaporin proteins in cell membranes is required for net inflow

of ions and water at the leading edge and net outflow at the trailing edge (Stroka et al., 2014).

Although our study identifies a function of Aqp1a.1 and Aqp8a.1 in facilitating water inflow, it

remains unknown which ion channels and transporters control ion flux across the cell membrane

to modulate water flow in tip cells. Results from our study suggest a role of SWELL1 in generating

an osmotic gradient that controls EC migration, since its inhibition resulted in reduced membrane

protrusions, slower tip cell migration and defective sprouting angiogenesis. Due to the limitations

of our experimental approach, we do not know whether the effect of inhibiting SWELL1 function

is cell autonomous or a consequence of altered chloride flux in the neighbouring cells that would

alter tissue osmolarity. It also remains to be clarified where SWELL1 is localized within migrating

ECs, how it regulates the osmotic gradient within the EC (cellular) and between the EC and the surrounding interstitial tissue (extracellular) to control the direction of water flow, and whether other ion channels also work in concert to modulate overall cellular and extracellular osmolarity to direct EC migration.

We have also observed enrichment of Aqp1a.1 and Aqp8a.1 proteins in specific compartments of ECs, at the leading edge of migrating endothelial tip cells as well as apical membranes during lumen formation (not shown). This leads to the question of how Aquaporin localization is regulated as this will determine the site of increased water inflow and hydrostatic pressure. It has been shown that rapid changes in subcellular localization of mammalian Aquaporins upon stimulation occurs mainly via trafficking to and from the plasma membrane (Markou et al., 2022) to modulate the amount of protein in the plasma membrane. The regulation of AQP subcellular relocalization via calmodulin- and/or phosphorylation-dependent mechanisms has been implicated for AQP0-5 and AQP7-9 (Markou et al., 2022). For example, rapid translocation of AQP1 to plasma membrane upon a hypotonic stimulus is dependent on calmodulin activation and phosphorylation by protein kinases C (PKC) (Conner et al., 2012) and subcellular localization of AQP8 in hepatocytes is also regulated by PKA and PI3K signalling (Gradilone et al., 2005, 2003). Further work is needed to elucidate the mechanism of Aqp1a.1 and Aqp8a.1 protein distribution in ECs.

In this study, we have assumed that Aqp1a.1 and Aqp8a.1 regulate EC behaviours through their ability to control water permeation based on previous studies demonstrating that water molecules flow through mammalian AQP1 (Preston et al., 1992; Zeidel et al., 1992) and AQP8 (Liu et al., 2006; Ma et al., 1997) as well as through zebrafish orthologs Aqp1a.1 and Aqp8a.1 (Tingaud-Sequeira et al., 2010). However, AQP1 and AQP8 are not water-specific channels. AQP1 can conduct unpolar gases such as carbon dioxide (Nakhoul et al., 1998; Prasad et al., 1998; Wang et al., 2007) and nitric oxide (Herrera et al., 2006; Herrera and Garvin, 2007), and in zebrafish, Aqp1a.1 also mediates the transfer for small gaseous molecules such as carbon dioxide and ammonia (Talbot et al., 2015). AQP8 is permeable to ammonia (Jahn et al., 2004; Liu et al., 2006) and is present in the inner mitochondrial membrane (Calamita et al., 2005), where it is suggested to mediate ammonia transport rather than water fluxes (Soria et al., 2010). Nevertheless, because water movement flows up an osmotic gradient, increased entry of ions into the cell through Aquaporins will also trigger water inflow.

435   In summary, our study highlights the role of water influx and hydrostatic pressure as
436 another force-generating mechanism utilized by cells to build tissues during animal development.
437 We demonstrate that endothelial tip cells employ both actin polymerization and hydrostatic
438 pressure for robust sprouting angiogenesis. As morphogenetic events are governed by a
439 coordination of cell shape changes, cell migration and rearrangements, we envision that the
440 shaping and formation of other tissues will also depend on Aquaporin function and changes in
441 hydrostatic pressure.

443 **METHODS**

445 **Zebrafish maintenance and stocks.**

446 Zebrafish (*Danio rerio*) were raised and staged according to established protocols (Kimmel et al.,
447 1995). Transgenic lines used in this work are *Tg(kdrl:EGFP)^{s843}* (Jin et al., 2005), *Tg(fli1a:H2B-*
448 *EGFP)^{ncv69}* [re-named in ZFIN as *Tg(fli1:zgc:114046-EGFP)*] (Ando et al., 2019), *Tg(fli1:Lifeact-*
449 *mCherry)^{ncv7}* (Wakayama et al., 2015), *Tg(fli1:myr-mCherry)^{ncv1}* (Fukuhara et al., 2014),
450 *Tg(fli1ep:EGFP-PLC1δPH)^{rk26}* (Kondrychyn et al., 2020), *Tg(fli1:h2bc1-mCherry)^{ncv31}*
451 (Nakajima et al., 2017), *Tg(kdrl:ras-mCherry)^{s916}* (Hogan et al., 2009), *Tg(kdrl:nls-*
452 *EGFP)^{ubs1};(fli1:Has.B4GALT1-mCherry)^{bns9}* (Kwon et al., 2016), *Tg(fli1ep:GAL4FF)^{ubs2}*
453 (Herwig et al., 2011). Zebrafish were maintained on a 12-hour light/12-hour dark cycle, and
454 fertilized eggs were collected and raised in E3 medium at 28^{0}C. To inhibit pigmentation in embryos
455 older than 24 hpf, 0.003% *N*-Phenylthiourea (Sigma-Aldrich) in E3 medium was used. All animal
456 experiments were approved by the Institutional Animal Care and Use Committee at RIKEN Kobe
457 Branch (IACUC).

459 **Plasmid construction and transgenesis.**

460 Plasmid *mEmerald-N1* was a gift from Michael Davidson (Addgene plasmid #53976), plasmids
461 *pMDS6* containing terminal *cis*-required sequences of the *Ds* transposon from maize and *pAc-SP6*
462 containing an ORF of *Ac* transposase were a gift from Sergei Parinov. Other plasmids used in this
463 study were generated by using In-Fusion HD Cloning kit (Takara Bio Inc.). A detailed information
464 regarding plasmids and primers used in this study can be found in Supplementary Tables 1 and 2,
465 respectively. *Tg(fli1ep:aqp1a.1-mEmerald)^{rk30}* and *Tg(fli1ep:aqp8a.1-mEmerald)^{rk31}* transgenic

zebrafish were generated by injecting *Tol2*-based *fli1ep:aqp1a.1-mEmerald* or *fli1ep:aqp8a.1-mEmerald* plasmids (10 ng/μl), respectively, along with *Tol2 transposase* mRNA (100 ng/μl) into one-cell stage AB embryos. *Tol2* transposase mRNA was transcribed from Not*I*-linearized *pCS-TP* plasmid (a gift from Koichi Kawakami, National Institute of Genetics, Japan) using the mMESSAGE mMACHINE SP6 kit (Invitrogen). Injected embryos were raised to adult and screened for founders.

Cloning of *aqp1a.1* and *aqp8a.1* genes.
Total RNA was isolated from 1-day old zebrafish embryos with TRI Reagent using Direct-zol™ RNA MicroPrep kit (Zymo Research) according to the manufacturer's protocol. The first-strand cDNA was synthesized from 1 μg of a total RNA by oligo(dT) priming using the SuperScript III First-Strand synthesis system (Invitrogen) according to the manufacturer's protocol. Amplification of cDNAs was performed using a high-fidelity KOD-Plus-Neo DNA polymerase (Toyobo, Japan) and resulting PCR products were cloned using NEB® PCR Cloning kit (New England BioLabs). Positive clones and plasmids were verified by DNA sequencing.

Mosaic expression of DNA constructs.
To avoid unnecessary *Tol2* re-transposition from the donor site in transgenic zebrafish lines generated using *Tol2* transposon (Kondrychyn et al., 2009), *Ac*/*Ds* transposon system from maize (Emelyanov et al., 2006) was used for mosaic expression of DNA constructs. *Ds*-based plasmids (10 pg) were co-injected with 100 pg of *Ac* transposase mRNA, transcribed from Bam*HI*-linearized *pAc-SP6* plasmid using the mMESSAGE mMACHINE SP6 kit (Invitrogen), into one-cell stage zebrafish wildtype or *aquaporin* mutant embryos. Embryos were analyzed at 25-26 hpf after injections.

***In vivo* cell volume analysis.**
Single-cell labelling of ECs in ISVs was achieved by mosaic expression of *pDs-kdrl:mEmerald* plasmid in wild type or *aqp1a.1[rk28/rk28];aqp8a.1[rk29/rk29]* mutant *Tg(fli1:Lifeact-mCherry)[ncv7]* transgenic embryos. To visualize single ECs with Aqp1a.1 overexpression, we injected *pDs-6xUAS:aqp1a.1-P2A-EGFP-T2A-mKate2-CAAX* plasmid into *Tg(fli1:GAL4FF)[ubs2]; Tg(fli1:myr-mCherry)[ncv1]* embryos. Embryos were imaged at 25-26 hpf with an Olympus UPLSAPO 60x/NA

1.2 water immersion objective with optical Z planes interval of 0.26 μm. Cell volume was

measured with Huygens Essential 22.10 software (Scientific Volume Imaging B.V.) using object

analyzer tool with manual adjustment of threshold.

Generation of zebrafish *aqp1a.1* and *aqp8a.1* mutants.

Aquaporins mutants were generated by CRISPR/Cas9-mediated mutagenesis. Guide RNAs

(gRNAs) targeting the first exons of *aqp1a.1* and *aqp8a.1* (Fig. S4 and S5) were generated by

using cloning-independent protocol (Gagnon et al., 2014). Briefly, to generate templates for gRNA

transcription, a 60-base oligodeoxynucleotides containing the SP6 promoter sequence, the gene-

specific   sequence   (*aqp1a.1*:   5'-GACAGCTGGCCAGCAGACCC-3';   *aqp8a.1*:   5'-

GATGTCTCCCCATCGCCCG-3') and 23-base overlap region was annealed to an 80-base

constant oligodeoxynucleotide encoding the reverse-complement of the tracrRNA tail. The ssDNA

overhangs were filled-in with T4 DNA polymerase (New England BioLabs) to generate dsDNA.

The gRNAs were *in vitro* transcribed using MEGAScript SP6 kit (Invitrogen), DNase treated and

purified with RNA Clean and Concentrator kit (Zymo Research). The gRNAs were quality-

checked by running 1 μg of the product on a 2% TBE-agarose gel. Cas9/gRNA RNP complex was

assembled just prior to injection and after 5 min incubation at room temperature 200 pg of Cas9

protein (Invitrogen) and 200 pg of gRNAs (100 pg of each, *aqp1a.1* gRNA and *aqp8a.1* gRNA)

were co-injected into one-cell stage *Tg(fli1:myr-mCherry)^{ncv1}* embryos. Such dose produced over

70% embryos survival post-injection showed CRISPR/Cas9-induced somatic *aqp1a.1* and

aqp8a.1 gene mutations. F_0 founders were identified by outcrossing CRISPR/Cas9-injected fish

with wild type fish and screening the offspring for mutations at 2 dpf using Sanger sequencing.

We found one F_0 fish with germline transmitted mutations in both *aqp1a.1* and *aqp8a.1* genes and

outcrossed it with *Tg(fli1:myr-mCherry)^{ncv1}* and *Tg(fli1a:H2B-EGFP)^{ncv69}* transgenic fish to

establish F_1 generation of heterozygotes. A 55 F_1 fish were genotyped to identify 4 *aqp1a.1$^{+/rk28}$*,

11 aqp8a.1$^{+/rk29}$ and 12   *aqp1a.1$^{+/rk28}$;aqp8a.1$^{+/rk29}$* heterozygote fish. Subsequently, F_1

heterozygote fish were in-crossed to establish a single and double homozygote fish.

DNA isolation and genotyping.

Genomic DNA was isolated from either embryos or fin clips using HotSHOT method (Meeker et

al., 2007). To identify genomic lesions, the following primers were used: aqp1a1-fwd (5'-

CGCCTCCAGATTCATTAGCAGGA-3') and aqp1a1-rev (5'-
GTAAGTGAACTGCTGCCAGTGA-3') to amplify a 550 bp fragment, and aqp8a1-fwd (5-
GGATCAATTGAGTTGCATAACAGAC-3') and aqp8a1-rev (5'-
CTGTAATGTAGACTTGTAAAGTGGA-3') to amplify a 638 bp fragment. Mutations were
assessed by direct sequencing of purified PCR products.

Quantitative real-time PCR.

Total RNA from whole embryos was isolated with TRI reagent using Direct-zol RNA MicroPrep

kit (Zymo Research) according to the manufacturer's protocol, including on-column DNA

digestion. RNA was quantified using a NanoDrop 1000 spectrophotometer (ThermoFisher

Scientific, USA). cDNA was synthesized from 300 ng of purified RNA using LunaScript RT

SuperMix kit (New England BioLabs) according to the manufacturer's protocol. Amplification of

target cDNA was performed in technical triplicate of three biological replicates using the SYBR

green methods. Each qPCR reaction mixture contained 5 µL 2x Luna Universal qPCR master mix

(New England BioLabs), 1 µL cDNA (2-fold dilution), 0.2 µL antarctic thermolabile UDG (New

England BioLabs) and 250 nM each primer to a final volume of 10 µL. Reactions were run in 384-

well plates (Applied Biosystems) using the QuantStudio 5 Real-Time PCR system (Applied

Biosystems) with the following thermal cycling conditions: initial UDG treatment at 25^0C for 2

minutes, UDG inactivation at 50^0C for 5 min and denaturation at 95^0C for 1 minutes, followed by

45 cycles of 15 sec at 95^0C, 30 sec at 60^0C with a plate read at the end of the extension step.

Control reactions included a no-template control (NTC) and a no-reverse transcriptase control

(NRT). Dissociation analysis of the PCR products was performed by running a gradient from 60

to 95^0C to confirm the presence of a single PCR product. Amplification data was analyzed using

QuantStudio Design and Analysis software v1.5.3 (Applied Biosystems) and relative fold change

was calculated using Pfaffl method (Pfaffl, 2001) and normalized to *gapdh* expression (as an

internal control). Primers are listed in Supplementary Table 2. Two embryos were used in each

biological replicate to generate an average value that was used to calculate the final mean ± SD

from two or three independent experiments.

RNA *in situ* hybridization.

RNAscope *in situ* hybridization. RNAscope® *in situ* hybridization was conducted by using RNAscope Multiplex Fluorescent Reagent kit v2 (Advanced Cell Diagnostics). We adapted manufacture's protocol designed for samples mounted on slides and protocol developed for whole-mount embryo samples (Gross-Thebing et al., 2014). For each experimental point, 5 embryos were processed in one 1.5 ml Eppendorf tube. Briefly, 20 and 22 hpf old embryos were manually dechorionated and fixed in freshly prepared 4% methanol-free PFA in PBS for 1 hour at room temperature. After fixation, embryos were washed three times in PBS containing 0.01% Tween-20 (PBST), dehydrated stepwise in 5-min washes in a series of increasing methanol concentrations (25%, 50%, 75%, 100%) in PBS and then stored in 100% methanol at -20^{0}C before use them for hybridization. The methanol-stored embryos were incubated with 5% H_2O_2 in methanol for 20 min at room temperature and then rehydrated stepwise in 5-min washes in series of decreasing methanol concentrations (75%, 50%, 25%) in PBS, followed by three times 5-min washes in PBST. Embryos were permeabilized in 2 drops of RNAscope® Protease III for 15 min, rinsed three times with PBST and hybridized with RNAscope target probes overnight at 50^{0}C in water bath. After hybridization probes were recovered and embryos were washed three times with 0.2x SSCT [0.01% Tween-20 in SSC] at room temperature, re-fixed with 4% PFA for 10 min, and washed three times with 0.2x SSCT. For RNA detection, the embryos were first hybridized with three different amplifier solution for 30 min (Amp1, and Amp2) and 15 min (Amp3) in a water bath at 40^{0}C. After each hybridization step, the embryos were washed three times with 0.2x SSCT for 10 min at room temperature. To develop signal of each probe, the embryos were sequentially incubated in a water bath at 40^{0}C with: (i) the horseradish peroxidase (HRP) for 15 min, (ii) TSA fluorophore for 30 min and (iii) the HRP blocker for 15 min. After each incubation step, the embryos were washed three times with 0.2x SSCT for 10 min at room temperature. HRP step is linked to a probe channel, we first developed C1 probe, *Dr-aqp1a.1* (Advanced Cell Diagnostics) using HRP-C1 and TSA Vivid Fluorophore 520 (Tocris, dilution 1:1500), then C2 probe, *Dr-lrrc8aa* (Advance Cell Diagnostics) using HRP-C2 and TSA Vivid Fluorophore 570 (Tocris, dilution 1:1500), and finally C3 probe, *Dr-aqp8a.1* (Advanced Cell Diagnostics) using HRP-C3 and TSA Vivid Fluorophore 650 (Tocris, dilution 1:1500). Nuclei were counterstained with DAPI ready-to-use solution overnight at 4^{0}C in the dark. Prior to imaging embryos were rinsed in PBST and kept in 70% glycerol in PBS at 4^{0}C in the dark. Images were acquired using Olympus FV3000 confocal microscope and an Olympus UPlanXApo 60x/NA 1.42 oil immersion objective.

Chromogenic *in situ* hybridization. Chromogenic *in situ* hybridization was conducted according to standard protocol (Thisse and Thisse, 2007) with minor modifications. For each experimental point, 10 embryos were processed in one 1.5 ml Eppendorf tube. Briefly, 30, 48, 72 and 96 hpf old embryos were manually dechorionated and fixed in freshly prepared 4% methanol-free PFA in PBS (pH 7.4) at 4^0C overnight. After fixation embryos were washed three times in PBS containing 0.1% Tween-20 (PBST), dehydrated stepwise in 5-min washes in a series of increasing methanol concentrations (25%, 50%, 75%, 100%) in PBS and then stored in 100% methanol at -20^0C before use them for hybridization. The methanol-stored embryos were rehydrated stepwise in 5-min washes in series of decreasing methanol concentrations (75%, 50%, 25%) in PBS followed by three times 5-min washes in PBST, and permeabilized 20 min with 10 µg/mL proteinase K (ThermoFisher Scientific). Hybridization was carried out in buffer [50% formamide, 5x SSC, 50 g/mL heparin, 500 g/mL tRNA (Roche) and 0.1% Tween-20] containing 5% dextran sulfate (MW=500 kDa) at 69^0C overnight. After stringency wash, the specimens were blocked with 2% blocking reagent (Roche) in maleic acid buffer [100 mM maleic acid, 150 mM NaCl, 50 mM $MgCl_2$ and 0.1% Tween-20, pH 7.5] and incubated overnight with anti-digoxigenin antibody, conjugated with alkaline phosphatase (dilution 1:5000, Roche) at 4^0C. For detection of alkaline phosphatase, specimens were incubated in staining buffer [50 mM Tris-HCl, 50 mM NaCl, 25 mM $MgCl_2$, 2% Polyvinyl alcohol and 0.1% Tween-20, pH 9.5] containing 375 µg/mL nitro-blue tetrazolium chloride (Roche) and 175 µg/mL 5-bromo-4-chloro-3'-indolyl-phosphate (Roche). Embryos were cleared in 70% glycerol overnight and imaged using Leica M205FA microscope.

RNA probe synthesis. The cDNA-containing vectors were linearized with appropriate restriction enzymes (detailed maps of vectors will be provided upon request) and used as a template for RNA probe synthesis. Sense and antisense RNA probes were synthesized using MEGAscript SP6 or T7 kits (Invitrogen) and digoxigenin (DIG)-labeled rNTPs (Roche) according to the manufacturer's protocol and purified using RNA Clean and Concentrator kit (Zymo Research). The following sense and antisense DIG-labeled riboprobes were generated: (i) *aqp1a.1*, 1.1 kb long probe comprised 121 nt of 5'UTR, 783 nt of the open reading frame (ORF) and 207 nt of 3'UTR; (ii) *aqp8a.1*, 0.8 kb long probe comprised 17 nt of 5'UTR, 783 nt of ORF and 30 nt of 3'UTR. Sense probes showed no specific or unspecific staining (data not shown).

618   **Image processing.** Chromogenic and RNAscope *in situ* hybridization images were processed
619   using ImageJ software (NIH, version 2.9.0/1.53t) with brightness and contrast adjustments. Z-
620   stacks of fluorescent images are presented as maximum intensity projections.
621
Analysis of mRNA expression in tip and stalk cells.
RNAscope processed embryos were mounted on slide and imaged using Olympus FV3000
confocal microscope and Olympus UPlanXApo 60x/NA 1.42 oil immersion objective. Maximum
intensity projection of z-stacks was used to determine the level of cellular fluorescence in ImageJ.
The tip and stalk cells were selected using freehand selection tool, "mean grey value", "area" and
"integrated density" in each ROI were measured first for channel 1 (aqp1a.1). A region next to cell
that has no fluorescence was selected as background and measured. The step was repeated for
channel 2 (aqp8a.1) on the same ROIs. Two parameters were calculated for each channel, (i) the
corrected total cell fluorescence, CTCF=Integrated Density – (Area of selected cell x Mean
fluorescence of background) and (ii) ratio, R=CTCF of tip cell/CTCF of stalk cell. We consider
that if R <0.9, expression is lower in tip cell, if R=0.9-1.1, expression is equal in tip and stalk cells,
if R>1.1, expression is higher in tip cell.
634
Imaging.
For live confocal imaging, embryos were mounted in 0.8% low-melt agarose (Bio-Rad) in E3
medium containing 0.16 mg/mL Tricaine (Sigma-Aldrich) and 0.003% phenylthiourea (Sigma-
Aldrich) in glass bottom 35-mm dishes (MatTek). Confocal Z-stacks were acquired using an
inverted Olympus IX83/Yokogawa CSU-W1 spinning disc confocal microscope equipped with a
Zyla 4.2 CMOS camera (Andor) and Olympus UPLSAPO 40x/NA 1.25 or 30x/NA 1.05 silicone
oil immersion objectives. Bright-field images were acquired on Leica M205FA microscope.
Images were processed using ImageJ software (NIH, version 2.9.0/1.53t).
643
Analysis of cell migration.
aqp1a.1+/rk28;aqp8a.1+/rk29 and *aqp1a.1rk28/rk28;aqp8a.1rk29/rk29* embryos on *Tg(fli1a:H2B-*
EGFP)ncv69;(fli1:Lifeact-mCherry)ncv7 double transgenic background were imaged from 21 to 30
hpf with time interval 6 min using Olympus UPLSAPO 30x/NA 1.05 silicon oil immersion
objective. We tried to mount both heterozygote and mutant embryos on the same dish (1 het plus

2 mut, or 2 het plus 2 mut) to image embryos on the same experimental conditions. Acquired time-

lapse images were registered using HyperStackReg plugin in ImageJ, tip cell and nuclei were

tracked using Manual Tracking plugin and velocity was calculated.

Analysis of grow rate of actin bundles in filopodia.

Wild type and *aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$* embryos on *Tg(fli1ep:EGFP-*

PLC1δPH)^{rk26};(fli1:Lifeact-mCherry)^{ncv7} double transgenic background were imaged starting from

25 hpf for 10 min with time interval 30 sec using Olympus UPLSAPO 60x/NA 1.2 water

immersion objective. Actin bundles growth was tracked using Manual Tracking plugin and

velocity was calculated.

Analysis of filopodia number and length.

Wild type, *aqp1a.1$^{+/rk28}$;aqp8a.1$^{+/rk29}$* and *aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$* embryos on

Tg(fli1ep:EGFP-PLC1δPH)^{rk26};(fli1:Lifeact-mCherry)^{ncv7} double transgenic background were

imaged starting from 24 hpf for 30 min with time interval of 50 sec using Olympus UPLSAPO

60x/NA 1.2 water immersion objective. In a single image we counted filopodia number per vessel,

measured membrane length and calculated filopodia number.

Assessment of tip cell polarization.

Tg(kdrl:nls-EGFP)^{ubs1};(fli1:Has.B4GALT1-mCherry)^{bns9Tg} wildtype and *aqp8a.1$^{rk29/rk29}$* embryos

were imaged at 24-26 hpf using an Olympus UPLSAPO 40x/NA 1.25 silicon oil immersion

objective. To analyze endothelial tip cell polarity in *aqp1a.1$^{rk28/rk29}$* embryos, Golgi apparatus and

nuclei were labelled by mosaic expression of *pDs-fli1ep:nls-EGFP-P2A-mKate2-GM130* plasmid.

Images were analyzed using Fiji (NIH). To assess polarity, the nucleus was fit into an ellipsoid

shape and the angle between the primary axis of the ellipse and the center of the Golgi was

measured (Fig. S10D). Polarization of ECs is defined as follows: (1) front (polarized), if the Golgi

is located within $-30^0 \sim +30^0$; (2) none, if the Golgi is located within $+30^0 \sim +150^0$ or $-30^0 \sim -150^0$;

or (3) rear, if the Golgi is located on the downstream side of the nucleus and angle is $-150^0 \sim +150^0$.

Tip cells in ISVs no. 5-15 were analyzed.

Quantification of EC number and ISV diameter.

Measurements were performed in wildtype, *aqp1a.1[rk28/rk28]*, *aqp8a.1[rk29/rk29]* and

aqp1a.1[rk28/rk28];aqp8a.1[rk29/rk29] zebrafish raised in *Tg(fli1:Lifeact-mCherry)[ncv7];(fli1a:H2B-*

EGFP)[ncv69] transgenic line or in *Tg(fli1ep:aqp1a.1-mEmerald)[rk30];(fli1:myr-mCherry)[ncv1]* and

Tg(fli1ep:aqp8a.1-mEmerald)[rk31]; (fli1:myr-mCherry)[ncv1] zebrafish. Confocal z-stacks images of

embryos were taken at 50 - 54 hpf and 3 dpf using Olympus UPLSAPO 40x/NA 1.25 silicone oil

immersion objective. ISVs no. 5 to 15 were used for quantification. For ISV diameter, 5

measurements were made along the length of each ISV using ImageJ (NIH) and the average was

plotted.

Chemical treatment.

Latrunculin B (Merck Millipore) was dissolved in DMSO to 1 mg/ml and stored at -20^0C. Ki8751

(Selleck Chemicals) was prepared as 5 mM solution in DMSO and stored at -80^0C. DCPIB (Tocris)

was prepared as 10 mM solution in EtOH and stored at -20^0C. All compounds were diluted to the

desired concentration in E3 medium (see in figure legends). Embryos were treated from 20 hpf for

6 (Ki8751 and DCPIB) or 8 hours (Latrunculin B). For imaging experiments, the same

concentrations of chemicals were added to the agarose and E3 medium.

Cell culture.

Adult human aortic endothelial cells (HAEC) (Lonza, lot-20TL231227) were cultured in EGM

medium (Lonza) and used at passage 3. Cells were seeded in EBM medium (Lonzo) supplemented

with 2% FBS (Gibco) at 4×10^5 cells/well on 24-well plate (Trueline) coated with 5 μg/ml

fibronectin (Sigma-Aldrich). Cells were treated with either 0.01% DMSO or different

concentrations of Ki8751 inhibitor for 6 hours. Cells were lysed with TRI reagent and RNA was

isolated using Direct-zol RNA MicroPrep kit (Zymo Research).

Single-cell RNA sequencing.

ECs were isolated from *Tg(kdrl:EGFP)[s843]* transgenic embryos. Cell sorting was carried out with

FACSAriaII Cell Sorter (BD Bioscience). Single-cell suspension was loaded into the 10X

Chromium system and cDNA libraries were constructed using Chromium Next GEM Single Cell

3′ GEM, Library and Gel Bead Kit v2 (10X Genomics) according to manufacturer's protocol.

Library was sequenced on the Illumina HiSeq 1500 Sequencer (Illumina, USA). Cell Ranger v2.1

was used to de-multiplex raw base call (BCL) files generated by Illumina sequencers into FASTQ

files.

Single-cell RNA sequencing data analysis.

The raw sequence data from above mentioned internal sequencing and also a public zebrafish

embryo data at 24 hpf stage (Gurung et al., 2022) were processed. The public dataset was obtained

from NCBI GEO database (accession number GSE202912). The raw sequencing reads from both

datasets were mapped to the zebrafish genome assembly (GRCz11, Ensembl release 112). Further

analyses were performed using Seurat package (version 4.3.0) in R software (https://www.r-

project.org). The expression matrices were first filtered by keeping genes that are expressed in a

minimum of 3 cells and cells that expressed a minimum of 200 genes for downstream analysis.

The data were then normalized using NormalizeData function which normalizes the gene

expression for each cell by the total expression counts (with a scale factor 10,000). To correct for

batch effect between the data from the two-time points, the rpca method in the Seurat package was

applied to integrate the data. The top 2000 variable genes identified using the vst method in

FindVariableFeatures function were used for principal component analysis in RunPCA function,

and the first 30 principal components were used for visualization analysis with Uniform Manifold

Approximation and Projection (UMAP) method, and in FindNeighbors function analysis. The cell

clustering resolution was set at 0.5 in FindClusters function. The Seurat object was then processed

using the ShinyCell package (version: 2.1.0) for gene expression visualizations.

Statistical analysis.

Statistical analysis was performed using Prism software version 10.2.0 (GraphPad). The variance

between the mean values of two groups was evaluated using the unpaired Student's *t*-test. For

assessment of more than three groups, we used one-way analysis of variance (ANOVA) test. A *P*

value of <0.05 was considered statistically significant. Statistic details can be found in each figure

legend.

Data availability.

The scRNA-seq raw data from this study is available at NCBI's Gene Expression Omnibus
database (accession number GSE262232). The data supporting the fundings of this study are
available from the corresponding author upon reasonable request.

Acknowledgements
We thank members of the Phng Lab, Y-C. Wang and S. Thukral for discussions and suggestions;
G. Chen, E. Taniguchi, RIKEN BDR Research Aquarium and RIKEN Kobe BioImaging Facilities
& Factory for technical assistance; J. Vermot, M. Francois and A. Yap for comments on the
manuscript. This work was supported by core funding from RIKEN BDR (to L.K.P.), RIKEN
BDR-Otsuka Pharmaceutical Collaboration Center (I.K.), the Naito Foundation (L.K.P.), the JSPS
Grants-in-Aid for Scientific Research (KAKENHI) grants (22H02624, 22H05168 to L.K.P;
22K06244 to I.K.), Swedish Research Council (2015-00550 to C.B.); Swedish Cancer Society
(2018/449, 2018/1154 to C.B.); Knut and Alice Wallenberg Foundation (2020.0057, 2018.0218 to
C.B.); Swedish Brain Foundation (ALZ2019-0130, ALZ2022-0005 to C.B.); the Leducq
Foundation (22CVD01, 23CVD02 to C.B.) and the Innovative Medicines Initiative (IM2PACT-
807015 to C.B.)

Authorship contributions
Conceptualization: L.K.P.; Methodology: I.K., L.H., H.W. and L.K.P.; Formal analysis: I.K. and
L.H.; Resources: C.B. and L.K.P.; Writing: I.K. and L.K.P; Supervision: L.K.P.

Declaration of interests
The authors declare no competing interests.

Figure legends

Figure 1. Differential expression of *aqp1a.1* and *aqp8a.1* mRNAs in tip and stalk cells.
(**A-F**) Detection of *aqp1a.1* and *aqp8a.1* mRNA by RNAscope *in situ* hybridization in 20 (A - D)
and 22 (E and F) hpf embryos. Representative maximum intensity projection of confocal z-stacks
of tip and stalk cells sprouting from DA are shown (n = 9 embryos, 2 independent experiments).

B and F show magnified region of a tip cell from a 20 and 22 hpf zebrafish, respectively (tip cell and EC nuclei in the DA are outlined). Scale bar, 10 μm (B and F) and 40 μm (A, C - E).

(**G**) Percentage of vascular sprouts with differential expression of *aqp1a.1* and *aqp8a.1* mRNA in tip and stalk cells in 22 hpf embryos (n=23 sprouts from 8 embryos, 2 independent experiments).

(**H**) qPCR analysis of *aqp1a.1* and *aqp8a.1* expression in zebrafish embryos treated with 0.01% DMSO or 1 μM ki8751 at 20 hpf for 6 hours (n = 3 independent experiments).

(**I**) qPCR analysis of *AQP1* gene expression in HAECs treated with 0.01% DMSO, 1 or 10 nM ki8751 for 6 hours (n = 2 independent experiments).

In panels H and J, gene expressions are shown relative to *gapdh* expression and data are presented as mean ± SD; statistical significance was determined by Brown-Forsythe and Welch ANOVA tests with Dunnett's multiple comparisons test; ns $p>0.05$, **$p<0.01$, ***$p<0.001$, and ****$p<0.0001$.

DA, dorsal aorta; PCV, posterior cardinal vein.

See also Fig. S1-S3.

Figure 2. Aquaporin proteins are enriched at the leading edge of migrating tip cells.

(**A-B**) Representative maximum intensity projection confocal z-stacks of endothelial tip cells of 25 hpf *Tg(fli1ep:aqp1a.1-mEmerald)^{rk30};(kdrl:ras-mCherry)^{s916}* (A) and *Tg(fli1ep:aqp8a.1-mEmerald)^{rk31};(kdrl:ras-mCherry)^{s916}* (B) embryos. White serrated line outlines tip-stalk cell border. Scale bar, 20 μm.

(**C-D**) Still images from time-lapse movies of migrating tip cells from *Tg(fli1ep:aqp1a.1-mEmerald)^{rk30}* (C) and *Tg(fli1ep:aqp8a.1-mEmerald)^{rk31}* (D) embryos. Movies were taken from 24 to 30 hpf. Arrowheads, Aquaporin protein localization at the leading edge of migrating tip cells. Arrows, Aquaporin localization in filopodia. Time, h:min. Scale bar, 5 μm.

See also Movies S1 and S2.

Figure 3. Loss of Aquaporin function leads to defective trunk vessel formation.

(**A-D**) Representative maximum intensity projection confocal z-stacks of 28 hpf *aqp1a.1$^{+/rk28}$;aqp8a.1$^{+/rk29}$* (A), *aqp1a.1$^{rk28/rk28}$* (B), *aqp8a.1$^{rk29/rk29}$* (C) and *aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$* (D) embryos.

(**E**) Quantification of ISV length at different positions along the trunk of 28 hpf embryos

(*aqp1a.1$^{+/rk28}$;aqp8a.1$^{+/rk29}$*: n = 21 embryos; *aqp1a.1$^{rk28/rk28}$*: n = 20 embryos; *aqp8a.1$^{rk29/rk29}$*: n =

20 embryos; *aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$*: n = 19 embryos from 2 independent experiments).

(**F**) Illustration of aISV and vISV connections in the trunk of wild type embryo at 3 dpf.

(**G-J**) Representative maximum intensity projection confocal z-stacks of 3 dpf wild type (G),

aqp8a.1$^{rk29/rk29}$ (H), *aqp1a.1$^{rk28/rk28}$* (I), and *aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$* (J) embryos. Asterisk

marks incomplete ISVs, arrows point EC nuclei on the contralateral side.

(**K**) Pie chart showing proportion of different classes of truncated ISV phenotypes (I-VI) found in

aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$ mutant embryos at 3 dpf.

DA, dorsal aorta; DLAV, dorsal longitudinal anastomotic vessel; ISV, intersegmental vessel; aISV,

arterial ISV; vISV, venous ISV; NC, notochord; NT, neural tube; PCV, posterior cardinal vein.

Scale bar, 50 µm (G-J) and 200 µm (A-D).

See also Fig. S6-S8.

Figure 4. Aquaporins promote endothelial tip cell protrusion and migration.

(**A**) Timing of tip cell emergence from the DA in *aqp1a.1$^{+/rk28}$;aqp8a.1$^{+/rk29}$* (n = 23 cells) and

aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$ (n = 27 cells).

(**B-C**) Still images from time-lapse imaging of migrating tip cells from *aqp1a.1$^{+/rk28}$;aqp8a.1$^{+/rk29}$*

(B) and *aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$* (C) embryos from 20 to 30 hpf. White arrows, retracting

tip cell. *, secondary sprouting from PCV. Scale bar, 10 µm.

(**D-E**) Stills images from representative time-lapse movies of migrating tip cells in

aqp1a.1$^{+/rk28}$;aqp8a.1$^{+/rk29}$ (D, n = 2) and *aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$* (E, n = 5) embryos.

Movies were taken from 25 to 30 hpf (n = 2 independent experiments). Bracket, formation of stable

protrusions. Scale bar, 20 µm.

(**F-G**) Representative maximum intensity projection confocal z-stacks of tip cells from wild type

(F) and *aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$* (G) embryos at 26 hpf. Scale bar, 10 µm.

(**H**) Quantification of filopodia number in tip cells of wild type (n = 36 cells from 10 embryos, 2

independent experiments), *aqp1a.1$^{+/rk28}$;aqp8a.1$^{+/rk29}$* (n = 19 cells from 6 embryos, 2 independent

experiments) and *aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$* (n = 28 cells from 9 embryos, 2 independent

experiments) embryos.

(**I**) Quantification of filopodia length in tip cells of wild type (n = 24 cells from 7 embryos, 2

independent experiments), *aqp1a.1$^{+/rk28}$;aqp8a.1$^{+/rk29}$* (n = 16 cells from 7 embryos, 2 independent

experiments) and *aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$* (n = 11 cells from 6 embryos, 2 independent

experiments) embryos.

(**J**) Quantification of growth rate of actin bundles in tip cell filopodia in 25 hpf wild type (n = 12

cells from 7 embryos, 2 independent experiments) and *aqp1a.1^{rk28};aqp8a.1^{rk29}* (n = 12 cells from

6 embryos, 2 independent experiments) embryos.

(**K**) Quantification of tip cell leading edge displacement of *aqp1a.1$^{+/rk28}$;aqp8a.1$^{+/rk29}$* (n = 19 cells

from 5 embryos, 3 independent experiments) and *aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$* (n = 47 cells from

13 embryos, 6 independent experiments) embryos at 25 - 30 hpf.

(**L**) Quantification of tip cell nuclei displacement of *aqp1a.1$^{+/rk28}$;aqp8a.1$^{+/rk29}$* (n = 20 cells from

6 embryos, 4 independent experiments) and *aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$* (n = 20 cells from 4

embryos, 2 independent experiments) embryos at 21 - 30 hpf.

(**M**) Quantification of tip cell migration velocity in *aqp1a.1$^{+/rk28}$;aqp8a.1$^{+/rk29}$* (n = 19 cells from

5 embryos, 3 independent experiments) and *aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$* (n = 47 cells from 13

embryos, 6 independent experiments) embryos at 25 - 30 hpf.

Statistical significance was determined by Brown-Forsythe and Welch ANOVA tests with

Dunnett's (H) or Sidak's (I) multiple comparisons test, and with unpaired *t*-test (J, M). ns, $p > 0.05$,

**, $p < 0.01$, ***, $p < 0.001$, and ****, $p < 0.0001$.

See also Fig. S10, Movies S3-6.

Figure 5. Aquaporin-mediated water influx increases tip cell volume.

(**A-C**) Representative maximum intensity projection confocal z-stacks of tip cell and its 3D-

volume rendering of 25 hpf wildtype (A), *aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$* (B) and wildtype

embryos overexpressing Aqp1a.1 (C) labelled with cytosolic mEmerald or EGFP. Scale bar, 10

μm.

(**D**) Quantification of tip cell volume in 25 hpf wild type (n = 24 cells) and

aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$ (n = 20 cells) and wildtype overexpressing Aqp1a.1 (n = 32 cells)

embryos from 3 independent experiments. Statistical significance was determined by Brown-

Forsythe and Welch ANOVA tests. *, $p < 0.05$, **, $p < 0.01$, and ****, $p < 0.0001$.

Figure 6. The chloride channel SWELL1 promotes EC migration and sprouting angiogenesis.

(**A-B**) UMAP plots showing co-expression of *lrrc8aa* and *aqp1a.1* (A), and *lrrc8aa* and *aqp8a.1* (B) in endothelial cells from 24 hpf, 34 hpf and 3 dpf embryos.

(**C-E**) Detection of *aqp1a.1*, *aqp8a.1* and *lrrc8aa* mRNA by RNAscope *in situ* hybridization in 22 hpf zebrafish embryo. Representative maximum intensity projection rendering of confocal *z*-stacks of tip and stalk cells (C). D and E show surface rendering of image C where only endothelial expression of *lrrc8aa* is shown after masking. Scale bar, 10 µm.

(**F-G**) Representative maximum intensity projection confocal z-stacks of 26 hpf wildtype embryos treated with 0.05% EtOH (F) or 5 µM DCPIB (G) from 20 to 26 hpf (control, n = 16 embryos; DCPIB, n = 18 embryos; 2 independent experiments). Scale bar, 50 µm.

(**H**) Quantification of ISV length in 26 hpf wildtype embryos treated with 0.05% EtOH or 5 µM DCPIB (control, n = 133 vessels from 16 embryos; DCPIB, 157 vessels from 18 embryos; 2 independent experiments). Statistical significance was determined with unpaired *t*-test; ****, $p < 0.0001$.

(**I**) Quantification of tip cell leading edge displacement of wildtype embryos treated with 0.05% EtOH (n = 29 cells from 7 embryos, 2 independent experiments) or 5 µM DCPIB (n = 29 cells from 8 embryos, 3 independent experiments) at 25 - 30 hpf.

(**J**) Quantification of tip cell migration velocity in wildtype treated with 0.05% EtOH (n = 29 cells from 7 embryos, 2 independent experiments) or 5 µM DCPIB (n = 29 cells from 8 embryos, 3 independent experiments) at 25 - 30 hpf. Statistical significance was determined with unpaired *t*-test; ****, $p < 0.0001$.

(**K**) Quantification of filopodia number in tip cells of wildtype embryos treated with 0.05% EtOH (n = 34 vessels from 9 embryos, 3 independent experiments) or 5 µM DCPIB (n = 53 vessels from 16 embryos, 3 independent experiments). Statistical significance was determined with unpaired *t*-test; ****, $p < 0.0001$.

(**M-N**) Still images from representative time-lapse movies of migrating tip cells in wildtype embryos treated with 0.05% EtOH (M, n = 7) or 5 µM DCPIB (N, n = 8). Movies were taken from 25 to 30 hpf (n = 3 independent experiments). Bracket, formation of stable protrusions. Scale bar, 20 µm.

Figure 7. Additive function of actin polymerization and hydrostatic pressure in driving EC migration and sprouting angiogenesis.

(**A-D**) Representative maximum intensity projection confocal z-stacks of 28 hpf wild type (A and B) and $aqp1a.1^{rk28/rk28};aqp8a.1^{rk29/rk29}$ (C and D) embryos treated with 0.1% DMSO (A and C) or 0.8 µg/ml Lat. B (B and D) from 20 to 28 hpf (for each condition: wild type, n = 10 embryos; $aqp1a.1^{rk28/rk28};aqp8a.1^{rk29/rk29}$, n =12 embryos, 2 independent experiments). Scale bar, 20 µm.

(**E**) Quantification of ISV length in 28 hpf wild type and $aqp1a.1^{rk28/rk28};aqp8a.1^{rk29/rk29}$ embryos treated with 0.1% DMSO or 0.8 µg/ml Lat. B (wild type: n = 60 vessels from 10 control embryos, n = 66 vessels from 10 Lat. B-treated embryos; $aqp1a.1^{rk28};aqp8a.1^{rk29}$: n = 126 vessels from 12 control embryos, n = 112 vessels from 11 Lat. B-treated embryos, 2 independent experiments). Statistical significance was determined by Brown-Forsythe ANOVA test with Sidak's multiple comparison test; ns, $p,>,0.05$, $^{**},p < 0.01$, and $^{****},p < 0.0001$. (**F**) Model of endothelial tip cell migration. Tip cells generate membrane protrusions by actin polymerization and increased hydrostatic pressure via Aquaporin-mediated water inflow. In the absence of actin polymerization and presence of Aquaporins, hydrostatic pressure deforms membranes to generate membrane protrusions. When Aquaporin function is lost, membrane protrusions is generated by actin polymerization. Fewer membrane protrusions are formed when only one mechanism is utilized, resulting in slower EC migration.

See also Movie S8.

References

Agre P, King LS, Yasui M, Guggino WB, Ottersen OP, Fujiyoshi Y, Engel A, Nielsen S. 2002. Aquaporin water channels – from atomic structure to clinical medicine. *Journal of Physiology* **542**:3–16. doi:10.1113/jphysiol.2002.020818

Ando K, Wang W, Peng D, Chiba A, Lagendijk AK, Barske L, Crump JG, Stainier DYR, Lendahl U, Koltowska K, Hogan BM, Fukuhara S, Mochizuki N, Betsholtz C. 2019. Peri-arterial specification of vascular mural cells from naïve mesenchyme requires Notch signaling. *Development* **146**:dev165589-13. doi:10.1242/dev.165589

Bill BR, Korzh V. 2014. Choroid plexus in developmental and evolutionary perspective. *Frontiers in Neuroscience* **8**:363. doi:10.3389/fnins.2014.00363

Boer LL de, Vanes L, Melgrati S, O'May JB, Hayward D, Driscoll PC, Day J, Griffiths A, Magueta R, Morrell A, MacRae JI, Köchl R, Tybulewicz VLJ. 2023. T cell migration requires

ion and water influx to regulate actin polymerization. *Nature Communications* **14**:7844.
doi:10.1038/s41467-023-43423-8

Calamita G, Ferri D, Gena P, Liquori GE, Cavalier A, Thomas D, Svelto M. 2005. The Inner
Mitochondrial Membrane Has Aquaporin-8 Water Channels and Is Highly Permeable to
Water*. *Journal of Biological Chemistry* **280**:17149–17153. doi:10.1074/jbc.c400595200

Chae YK, Woo J, Kim M-J, Kang SK, Kim MS, Lee J, Lee SK, Gong G, Kim YH, Soria JC,
Jang SJ, Sidransky D, Moon C. 2008. Expression of Aquaporin 5 (AQP5) Promotes Tumor
Invasion in Human Non Small Cell Lung Cancer. *PLoS ONE* **3**:e2162.
doi:10.1371/journal.pone.0002162

Chan CJ, Costanzo M, Ruiz-Herrero T, Mönke G, Petrie RJ, Bergert M, Diz-Muñoz A,
Mahadevan L, Hiiragi T. 2019. Hydraulic control of mammalian embryo size and cell fate.
Nature **571**:112–116. doi:10.1038/s41586-019-1309-x

Chen C, Qin Y, Xu Y, Wang X, Lei W, Shen X, Chen L, Wang L, Gong J, Wang Y, Hu S, Liu
D. 2024. Aquaporins enriched in endothelial vacuole membrane regulate the diameters of
microvasculature in hyperglycaemia. *Cardiovascular Research* **120**:1065-1080.
doi:10.1093/cvr/cvae085

Choudhury MI, Benson MA, Sun SX. 2022. Trans-epithelial fluid flow and mechanics of
epithelial morphogenesis. *Seminars in Cell & Developmental Biology* **131**:146–159.
doi:10.1016/j.semcdb.2022.05.020

Chugh M, Munjal A, Megason SG. 2022. Hydrostatic pressure as a driver of cell and tissue
morphogenesis. *Seminars in Cell & Developmental Biology* **131**:134–145.
doi:10.1016/j.semcdb.2022.04.021

Clarke DN, Martin AC. 2021. Actin-based force generation and cell adhesion in tissue
morphogenesis. *Current Biology* **31**:R667–R680. doi:10.1016/j.cub.2021.03.031

Conner MT, Conner AC, Bland CE, Taylor LHJ, Brown JEP, Parri HR, Bill RM. 2012. Rapid
Aquaporin Translocation Regulates Cellular Water Flow. *Journal of Biological Chemistry*
287:11516–11525. doi:10.1074/jbc.m111.329219

Day RE, Kitchen P, Owen DS, Bland C, Marshall L, Conner AC, Bill RM, Conner MT. 2014.
Human aquaporins: regulators of transcellular water flow. *Biochimica et Biophysica Acta*
1840:1492–1506. doi:10.1016/j.bbagen.2013.09.033

Dumortier JG, Verge-Serandour ML, Tortorelli AF, Mielke A, Plater L de, Turlier H, Maître J-L.
2019. Hydraulic fracturing and active coarsening position the lumen of the mouse blastocyst.
Science **365**:465–468. doi:10.1126/science.aaw7709

Emelyanov A, Gao Y, Naqvi NI, Parinov S. 2006. Trans-Kingdom Transposition of the Maize
Dissociation Element. *Genetics* **174**:1095–1104. doi:10.1534/genetics.106.061184

Farinas J, Kneen M, Moore M, Verkman AS. 1997. Plasma Membrane Water Permeability of
Cultured Cells and Epithelia Measured by Light Microscopy with Spatial Filtering. *Journal of*
General Physiology **110**:283–296. doi:10.1085/jgp.110.3.283

Figueiredo AM, Barbacena P, Russo A, Vaccaro S, Ramalho D, Pena A, Lima AP, Ferreira RR,
Fidalgo MA, El-Marjou F, Carvalho Y, Vasconcelos FF, Lennon-Duménil A-M, Vignjevic
DM, Franco CA. 2021. Endothelial cell invasion is controlled by dactylopodia. *Proceedings*
of the National Academy of Sciences **118**:e2023829118. doi:10.1073/pnas.2023829118

Fukuhara S, Zhang J, Yuge S, Ando K, Wakayama Y, Sakaue-Sawano A, Miyawaki A,
Mochizuki N. 2014. Visualizing the cell-cycle progression of endothelial cells in zebrafish.
Developmental Biology **393**:10–23. doi:10.1016/j.ydbio.2014.06.015

Gagnon JA, Valen E, Thyme SB, Huang P, Akhmetova L, Ahkmetova L, Pauli A, Montague
TG, Zimmerman S, Richter C, Schier AF. 2014. Efficient mutagenesis by Cas9 protein-
mediated oligonucleotide insertion and large-scale assessment of single-guide RNAs. *PLoS*
ONE **9**:e98186. doi:10.1371/journal.pone.0098186

Gebala V, Collins R, Geudens I, Phng L-K, Gerhardt H. 2016. Blood flow drives lumen
formation by inverse membrane blebbing during angiogenesis in vivo. *Nature Cell Biology*
18:443–450. doi:10.1038/ncb3320

Gerhardt H, Golding M, Fruttiger M, Ruhrberg C, Lundkvist A, Abramsson A, Jeltsch M,
Mitchell C, Alitalo K, Shima D, Betsholtz C. 2003. VEGF guides angiogenic sprouting
utilizing endothelial tip cell filopodia. *Journal of Cell Biology* **161**:1163–1177.
doi:10.1083/jcb.200302047

Gradilone SA, Carreras FI, Lehmann GL, Marinelli RA. 2005. Phosphoinositide 3-kinase is
involved in the glucagon-induced translocation of aquaporin-8 to hepatocyte plasma
membrane. *Biology of the Cell* **97**:831–836. doi:10.1042/bc20040115

Gradilone SA, García F, Huebert RC, Tietz PS, Larocca MC, Kierbel A, Carreras FI, Larusso
NF, Marinelli RA. 2003. Glucagon induces the plasma membrane insertion of functional
aquaporin-8 water channels in isolated rat hepatocytes. *Hepatology* **37**:1435–1441.
doi:10.1053/jhep.2003.50241

Gross-Thebing T, Paksa A, Raz E. 2014. Simultaneous high-resolution detection of multiple
transcripts combined with localization of proteins in whole-mount embryos. *BMC Biology*
12:55. doi:10.1186/s12915-014-0055-7

Gunasekar SK, Xie L, Kumar A, Hong J, Chheda PR, Kang C, Kern DM, My-Ta C, Maurer J,
Heebink J, Gerber EE, Grzesik WJ, Elliot-Hudson M, Zhang Y, Key P, Kulkarni CA, Beals
JW, Smith GI, Samuel I, Smith JK, Nau P, Imai Y, Sheldon RD, Taylor EB, Lerner DJ,
Norris AW, Klein S, Brohawn SG, Kerns R, Sah R. 2022. Small molecule SWELL1 complex
induction improves glycemic control and nonalcoholic fatty liver disease in murine Type 2
diabetes. *Nature Communications* **13**:784. doi:10.1038/s41467-022-28435-0

Gurung S, Restrepo NK, Chestnut B, Klimkaite L, Sumanas S. 2022. Single-cell transcriptomic
analysis of vascular endothelial cells in zebrafish embryos. *Scientific Reports* **12**:13065.
doi:10.1038/s41598-022-17127-w

Herrera M, Garvin JL. 2007. Novel role of AQP-1 in NO-dependent vasorelaxation. *American*
Journal of Physiology-Renal Physiology **292**:F1443–F1451. doi:10.1152/ajprenal.00353.2006

Herrera M, Hong NJ, Garvin JL. 2006. Aquaporin-1 Transports NO Across Cell Membranes.
Hypertension **48**:157–164. doi:10.1161/01.hyp.0000223652.29338.77

Herwig L, Blum Y, Krudewig A, Ellertsdottir E, Lenard A, Belting H-G, Affolter M. 2011.
Distinct cellular mechanisms of blood vessel fusion in the zebrafish embryo. *Current*
Biology **21**:1942-1948. doi:10.1016/j.cub.2011.10.016

Hogan BM, Bos FL, Bussmann J, Witte M, Chi NC, Duckers HJ, Schulte-Merker S. 2009. ccbe1
is required for embryonic lymphangiogenesis and venous sprouting. *Nature Genetics* **41**:396-
398. doi:10.1038/ng.321

Jahn TP, Møller ALB, Zeuthen T, Holm LM, Klærke DA, Mohsin B, Kühlbrandt W, Schjoerring
JK. 2004. Aquaporin homologues in plants and mammals transport ammonia. *FEBS Letters*
574:31–36. doi:10.1016/j.febslet.2004.08.004

Jin S-W, Beis D, Mitchell T, Chen J-N, Stainier DY. 2005. Cellular and molecular analyses of
vascular tube and lumen formation in zebrafish. *Development* **132**: 5199-5209.
doi:10.1242/dev.02087

Jung HJ, Park J-Y, Jeon H-S, Kwon T-H. 2011. Aquaporin-5: A Marker Protein for Proliferation
and Migration of Human Breast Cancer Cells. *PLoS ONE* **6**:e28492.
doi:10.1371/journal.pone.0028492

Karlsson T, Bolshakova A, Magalhães MAO, Loitto VM, Magnusson K-E. 2013. Fluxes of
Water through Aquaporin 9 Weaken Membrane-Cytoskeleton Anchorage and Promote
Formation of Membrane Protrusions. *PLoS ONE* **8**:e59901.
doi:10.1371/journal.pone.0059901

Kimmel CB, Ballard WW, Kimmel SR, Ullmann B, Schilling TF. 1995. Stages of embryonic
development of the zebrafish. *Developmental dynamics* **203**:253--310.
doi:10.1002/aja.1002030302

Kondrychyn I, Garcia-Lecea M, Emelyanov A, Parinov S, Korzh V. 2009. Genome-wide
analysis of Tol2 transposon reintegration in zebrafish. *BMC Genomics* **10**:418–418.
doi:10.1186/1471-2164-10-418

Kondrychyn I, Kelly DJ, Carretero NT, Nomori A, Kato K, Chong J, Nakajima H, Okuda S,
Mochizuki N, Phng L-K. 2020. Marcksl1 modulates endothelial cell mechanoresponse to

haemodynamic forces to control blood vessel shape and size. *Nature Communications*
11:5476–18. doi:10.1038/s41467-020-19308-5

Koun S, Kim J-D, Rhee M, Kim M-J, Huh T-L. 2016. Spatiotemporal expression pattern of the
zebrafish aquaporin 8 family during early developmental stages. *Gene Expression Patterns*
21:1–6. doi:10.1016/j.gep.2016.06.001

Kozono D, Yasui M, King LS, Agre P. 2002. Aquaporin water channels: atomic structure
molecular dynamics meet clinical medicine. *Journal of Clinical Investigation* **109**:1395–1399.
doi:10.1172/jci15851

Kwon H-B, Wang S, Helker CSM, Rasouli SJ, Maischein H-M, Offermanns S, Herzog W,
Stainier DYR. 2016. In vivo modulation of endothelial polarization by Apelin receptor
signalling. *Nature Communications* **7**:11805–12. doi:10.1038/ncomms11805

Li Y, Konstantopoulos K, Zhao R, Mori Y, Sun SX. 2020. The importance of water and
hydraulic pressure in cell dynamics. *Journal of Cell Science* **133**:jcs240341.
doi:10.1242/jcs.240341

Liu K, Nagase H, Huang CG, Calamita G, Agre P. 2006. Purification and functional
characterization of aquaporin-8. *Biology of the Cell* **98**:153–161. doi:10.1042/bc20050026

Loitto VM, Huang C, Sigal YJ, Jacobson K. 2007. Filopodia are induced by aquaporin-9
expression. *Experimental Cell Research* **313**:1295–1306. doi:10.1016/j.yexcr.2007.01.023

Loitto VM, Karlsson T, Magnusson K. 2009. Water flux in cell motility: Expanding the
mechanisms of membrane protrusion. *Cell Motility and the Cytoskeleton* **66**:237–247.
doi:10.1002/cm.20357

Ma T, Yang B, Verkman AS. 1997. Cloning of a Novel Water and Urea-Permeable Aquaporin
from Mouse Expressed Strongly in Colon, Placenta, Liver, and Heart. *Biochemical and*
Biophysical Research Communications **240**:324–328. doi:10.1006/bbrc.1997.7664

Maltaneri RE, Schiappacasse A, Chamorro ME, Nesse AB, Vittori DC. 2020. Aquaporin-1 plays
a key role in erythropoietin-induced endothelial cell migration. *Biochimica et Biophysica Acta*
(BBA) - Molecular Cell Research **1867**:118569. doi:10.1016/j.bbamcr.2019.118569

Markou A, Unger L, Abir-Awan M, Saadallah A, Halsey A, Balklava Z, Conner M, Törnroth-
Horsefield S, Greenhill SD, Conner A, Bill RM, Salman MM, Kitchen P. 2022. Molecular
mechanisms governing aquaporin relocalisation. *Biochimica et Biophysica Acta (BBA) -*
Biomembranes **1864**:183853. doi:10.1016/j.bbamem.2021.183853

McLennan R, McKinney MC, Teddy JM, Morrison JA, Kasemeier-Kulesa JC, Ridenour DA,
Manthe CA, Giniunaite R, Robinson M, Baker RE, Maini PK, Kulesa PM. 2019. Neural crest
cells bulldoze through the microenvironment using Aquaporin 1 to stabilize filopodia.
Development **147**:dev185231. doi:10.1242/dev.185231

Meeker ND, Hutchinson SA, Ho L, Trede NS. 2007. Method for isolation of PCR-ready
genomic DNA from zebrafish tissues. *Biotechniques* **43**:610–614. doi:10.2144/000112619

Mosaliganti KR, Swinburne IA, Chan CU, Obholzer ND, Green AA, Tanksale S, Mahadevan L,
Megason SG. 2019. Size control of the inner ear via hydraulic feedback. *eLife* **8**:e39596.
doi:10.7554/elife.39596

Munjal A, Lecuit T. 2014. Actomyosin networks and tissue morphogenesis. *Development*
141:1789-1793. doi: 10.1242/dev.091645

Murrell M, Oakes PW, Lenz M, Gardel ML. 2015. Forcing cells into shape: the mechanics of
actomyosin contractility. *Nature Cell Biology* **16**:486–498. doi:10.1038/nrm4012

Nagy TL, Strickland J, Weiner OD. 2023. Neutrophils actively swell to potentiate rapid
migration. *bioRxiv* 2023.05.15.540704. doi:10.1101/2023.05.15.540704

Nakajima H, Yamamoto K, Agarwala S, Terai K, Fukui H, Fukuhara S, Ando K, Miyazaki T,
Yokota Y, Schmelzer E, Belting H-G, Affolter M, Lecaudey V, Mochizuki N. 2017. Flow-
Dependent Endothelial YAP Regulation Contributes to Vessel Maintenance. *Developmental*
Cell **40**:523-536.e6. doi:10.1016/j.devcel.2017.02.019

Nakhoul NL, Davis BA, Romero MF, Boron WF. 1998. Effect of expressing the water channel
aquaporin-1 on the CO_2 permeability of Xenopus oocytes. *American Journal of Physiology-*
Cell Physiology **274**:C543–C548. doi:10.1152/ajpcell.1998.274.2.c543

Parab S, Quick RE, Matsuoka RL. 2021. Endothelial cell-type-specific molecular requirements
for angiogenesis drive fenestrated vessel development in the brain. *eLife* **10**:e64295.
doi:10.7554/elife.64295

Pfaffl MW. 2001. A new mathematical model for relative quantification in real-time RT–PCR.
Nucleic Acids Research **29**:e45–e45. doi:10.1093/nar/29.9.e45

Phng L-K, Stanchi F, Gerhardt H. 2013. Filopodia are dispensable for endothelial tip cell
guidance. *Development* **140**:4031–4040. doi:10.1242/dev.097352

Prasad GV, Coury LA, Finn F, Zeidel ML. 1998. Reconstituted aquaporin 1 water channels
transport CO_2 across membranes. *Journal of Biological Chemistry* **273**:33123–33126.
doi:10.1074/jbc.273.50.33123

Preston GM, Carroll TP, Guggino WB, Agre P. 1992. Appearance of water channels in Xenopus
oocytes expressing red cell CHIP28 protein. *Science* **256**:385–387.
doi:10.1126/science.256.5055.385

Qiu Z, Dubin AE, Mathur J, Tu B, Reddy K, Miraglia LJ, Reinhardt J, Orth AP, Patapoutian A.
2014. SWELL1, a Plasma Membrane Protein, Is an Essential Component of Volume-
Regulated Anion Channel. *Cell* **157**:447–458. doi:10.1016/j.cell.2014.03.024

Rehn K, Wong KS, Balciunas D, Sumanas S. 2011. Zebrafish enhancer trap line recapitulates
embryonic aquaporin 1a expression pattern in vascular endothelial cells. *International*
Journal of Developmental Biology **55**:613–618. doi:10.1387/ijdb.103249kp

Saadoun S, Papadopoulos MC, Hara-Chikuma M, Verkman AS. 2005. Impairment of
angiogenesis and cell migration by targeted aquaporin-1 gene disruption. *Nature* **434**:786–
792. doi:10.1038/nature03460

Sauteur L, Affolter M, Belting H-G. 2017. Distinct and redundant functions of Esama and VE-
cadherin during vascular morphogenesis. *Development* **144**:1554–1565.
doi:10.1242/dev.140038

Siekmann AF, Affolter M, Belting H-G. 2013. The tip cell concept 10 years after: New players
tune in for a common theme. *Experimental Cell Research* **319**:1255–1263.
doi:10.1016/j.yexcr.2013.01.019

Soria LR, Fanelli E, Altamura N, Svelto M, Marinelli RA, Calamita G. 2010. Aquaporin-8-
facilitated mitochondrial ammonia transport. *Biochemical and Biophysical Research*
Communications **393**:217–221. doi:10.1016/j.bbrc.2010.01.104

Stroka KM, Jiang H, Chen S-H, Tong Z, Wirtz D, Sun SX, Konstantopoulos K. 2014. Water
permeation drives tumor cell migration in confined microenvironments. *Cell* **157**:611–623.
doi:10.1016/j.cell.2014.02.052

Talbot K, Kwong RWM, Gilmour KM, Perry SF. 2015. The water channel aquaporin-1a1
facilitates movement of CO2 and ammonia in zebrafish (Danio rerio) larvae. *Journal of*
Experimental Biology **218**:3931–3940. doi:10.1242/jeb.129759

Thisse C, Thisse B. 2007. High-resolution in situ hybridization to whole-mount zebrafish
embryos. *Nature Protocols* **3**:59–69. doi:10.1038/nprot.2007.514

Tingaud-Sequeira A, Calusinska M, Finn RN, Chauvigné F, Lozano J, Cerdà J. 2010. The
zebrafish genome encodes the largest vertebrate repertoire of functional aquaporins with dual
paralogy and substrate specificities similar to mammals. *BMC Evolutionary Biology* **10**:38.
doi:10.1186/1471-2148-10-38

Voss FK, Ullrich F, Münch J, Lazarow K, Lutter D, Mah N, Andrade-Navarro MA, Kries JP
von, Stauber T, Jentsch TJ. 2014. Identification of LRRC8 Heteromers as an Essential
Component of the Volume-Regulated Anion Channel VRAC. *Science* **344**:634–638.
doi:10.1126/science.1252826

Wakayama Y, Fukuhara S, Ando K, Matsuda M, Mochizuki N. 2015. Cdc42 Mediates Bmp-
Induced Sprouting Angiogenesis through Fmnl3-Driven Assembly of Endothelial Filopodia in
Zebrafish. *Developmental Cell* **32**:109–122. doi:10.1016/j.devcel.2014.11.024

Wang Y, Cohen J, Boron WF, Schulten K, Tajkhorshid E. 2007. Exploring gas permeability of
cellular membranes and membrane channels with molecular dynamics. *Journal of Structural*
Biology **157**:534–544. doi:10.1016/j.jsb.2006.11.008

Wu DQ, Yang ZF, Wang KJ, Feng XY, Lv ZJ, Li Y, Jian ZX. 2017. AQP8 inhibits colorectal
cancer growth and metastasis by down-regulating PI3K/AKT signaling and PCDH7
expression. *American Journal of Cancer Research* **8**:266–279.

Zeidel ML, Ambudkar SV, Smith BL, Agre P. 1992. Reconstitution of functional water channels
in liposomes containing purified red cell CHIP28 protein. *Biochemistry* **31**:7436–7440.
doi:10.1021/bi00148a002

Zhang Y, Li Y, Thompson KN, Stoletov K, Yuan Q, Bera K, Lee SJ, Zhao R, Kiepas A, Wang
Y, Mistriotis P, Serra SA, Lewis JD, Valverde MA, Martin SS, Sun SX, Konstantopoulos K.
2022. Polarized NHE1 and SWELL1 regulate migration direction, efficiency and metastasis.
Nature Communications **13**:6128. doi:10.1038/s41467-022-33683-1

Figure 1

Figure 2

Figure 3

A *aqp1a.1$^{+/rk28}$;aqp8a.1$^{+/rk29}$*

B *aqp1a.1$^{rk28/rk28}$*

C *aqp8a.1$^{rk29/rk29}$*

D *aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$*

E

ISV length (µm) vs ISV position

- ■ *aqp1a.1$^{+/rk28}$;aqp8a.1$^{+/rk29}$*
- □ *aqp1a.1$^{rk28/rk28}$*
- ▨ *aqp8a.1$^{rk29/rk29}$*
- ■ *aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$*

F

aISV: DLAV, NT, NC, DA, PCV
vISV: DLAV, NT, NC, DA, PCV

G *aqp1a.1$^{+/+}$;aqp8a.1$^{+/+}$*

H *aqp8a.1$^{rk29/rk29}$*

I *aqp1a.1$^{rk28/rk28}$*

J *aqp1a.1$^{rk28/rk28}$;aqp8a.1$^{rk29/rk29}$*

Tg(fli1a:H2B-EGFP)^{ncv69} Tg(fli1:Lifeact-mCherry)^{ncv7}

K I II III IV V VI

- ■ 75.20% Class I
- ■ 15.79% Class II
- ■ 1.16% Class III
- ■ 2.23% Class IV
- ■ 2.94% Class V
- ■ 2.68% Class VI

n=1123 ISVs
(ISVs no. 5-28)

Figure 4

Figure 5

Figure 6

Tg(fli1:Lifeact-mCherry)^ncv7 Tg(fli1ep:EGFP-PLCδ1PH)^rk26

Figure 7

Tg(fli1ep:EGFP-PLCd1PH)^rk26 *Tg(fli1:Lifeact-mCherry)^ncv7*

Membrane protrusions generated by actin polymerization and increased hydrostatic pressure.

Membrane protrusions generated by increased hydrostatic pressure.

Membrane protrusions generated by actin polymerization.