1

N-cadherin facilitates trigeminal sensory neuron outgrowth and target

tissue innervation

Caroline Halmi[1], Alec McIntosh[2], Carrie E. Leonard[1], Lisa A. Taneyhill[1],*

[1]Department of Animal and Avian Sciences, University of Maryland, College Park, MD
20742 USA
[2]Lombardi Comprehensive Cancer Center, Georgetown University, Washington, DC,
20007 USA

Email: ltaney@umd.edu

*Corresponding author

ABSTRACT

During the development of the sensory nervous system, various cell types converge to form different tissues. The trigeminal ganglion houses the cell bodies for the trigeminal nerve and exemplifies these intercellular interactions, as it is formed from the condensation of two diverse precursor cell populations, neural crest cells and placode cells. The dual origin of the trigeminal ganglion has been understood for decades, but the molecules that orchestrate this process remain relatively unknown. Assembling the trigeminal ganglion is mediated by cell adhesion molecules including the protein neural cadherin (N-cadherin), expressed first in placode-derived neurons and later in all trigeminal sensory neurons. Prior studies have shown that N-cadherin knockdown in chick trigeminal placode cells leads to early defects in trigeminal ganglion assembly by impacting the ability of placode-derived neurons to properly condense with undifferentiated neural crest cells. Later functions for N-cadherin in chick trigeminal gangliogenesis, however, are unknown. Using morpholino-mediated knockdown of N-cadherin in chick trigeminal placode cells, we examined trigeminal ganglion development at later developmental stages, when neural crest cells are differentiating into neurons. Through these experiments, we uncovered a sustained negative impact on the trigeminal ganglion, leading to decreases in ganglion size, nerve outgrowth, and branching to target tissues *in vivo*. Further, blocking the adhesive function of N-cadherin reveals its importance in the outgrowth ability for some, but not all, trigeminal neurons *in vitro*. These deficits reflect potential cell and non-cell autonomous effects on placodal and neural crest-derived neurons, respectively, and point to the importance of N-cadherin-mediated adhesion among trigeminal sensory neurons. Our findings reveal continued adhesion-dependent functions for N-cadherin in the trigeminal ganglion, which will aid in the understanding of its tissue-specific roles.

INTRODUCTION

The development of an advanced nervous system has contributed, in part, to the evolution of more complex organisms like vertebrates that can sense their surroundings and respond to environmental cues. To this end, neurons of the peripheral nervous system transmit sensory information from the body back to the brain. A portion of these neurons condense into sensory ganglia, and specifically within the head, are known as cranial ganglia. While some of the cranial nerves function to transmit only sensory information or motor information, a subset of them conveys both. Of the five cranial nerves (trigeminal, facial, glossopharyngeal, vagus, and spinal accessory), the paired trigeminal nerves are unique in that they are the largest of all the cranial nerves and extensively innervate many facial regions. The trigeminal nerve is comprised of three major branches - the ophthalmic, maxillary, and mandibular - that innervate the forehead and eye region, upper jaw, and lower jaw, respectively. While all three branches relay sensory information, only the mandibular branch has motor neurons that aid in mastication.

The chick embryo has proved to be an exemplary model in understanding many developmental processes, including discerning the cell types that comprise the trigeminal ganglion[1,2]. Both neural crest cells and neurogenic placode cells contribute neurons to the trigeminal ganglion[1], while the neural crest also form glial cells. These embryonic cell populations are derived from the ectoderm and arise at the neural plate border, located between the non-neural ectoderm and the neural plate[3]. During early neural development, neural crest cells reside in the future neural folds, where they subsequently undergo an epithelial-to-mesenchymal transition to delaminate from the dorsal neural folds and become motile. Neural crest cells that contribute to the trigeminal ganglion originate from the midbrain and anterior hindbrain[4,5]; however, the neural crest as a cell population are found throughout the body axis and are capable of differentiating into numerous different cell types including cranial bone and cartilage, melanocytes, chromaffin cells, cardiac myocytes, and neurons of the peripheral nervous system[6]. Cranial neural crest cells collectively migrate through the embryonic head using a combination of chemoattraction and repulsion and mechanical sensing[7,8]. Trunk neural crest cells differ from the cranial neural crest in a number of ways including their derivatives, multipotent abilities, and migratory methods[7].

Cranial neurogenic placodes are specialized regions of thickened ectoderm that
differentiate into neurons prior to ingression[9]. Initially, there is slight overlap between the
different placodes that comprise the cranial ganglia; however, as development
progresses, the placodes become more segregated and distinct[10]. The trigeminal
placode, which consists of the ophthalmic and maxillomandibular placodes, gives rise to
all the placode-derived neurons of the trigeminal ganglion. Although these two placodes
work together to produce one structure, the placodes differ from one another in two
important ways. First, the ophthalmic and maxillomandibular placodes contribute neurons
to distinct lobes of the trigeminal ganglion (ophthalmic, and maxillary and mandibular,
respectively). Second, the majority of the cells to which the ophthalmic placode gives rise
are post-mitotic, traveling to the trigeminal ganglion as differentiated neurons, while the
maxillomandibular placode generates neuroblasts that do not differentiate until they have
reached the trigeminal ganglion[11–14]. Notably, in the chick, placodal neurons are the first
neurons present within the trigeminal ganglion, and are located distally compared to
neural crest-derived neurons, which are more proximal and form later during
development[2,15].

Prior studies have led to great advancements in understanding the complex
interactions that occur between neural crest and placodal neurons during their early
migration and the subsequent onset of gangliogenesis. Like neural crest cells, placodal
neurons are also capable of collective migration[16,17]. While the molecular mechanisms
behind these movements are poorly understood, the cellular basis of this involves neural
crest cells forming corridors to facilitate this process[18], as they are a more permissive
substrate compared to the surrounding mesoderm. Furthermore, neural crest and
placode cells exhibit "chase and run" behavior, where neural crest cells promote the
migration of placode cells to their terminal destinations[16]. These two reports, in addition
to other studies, indicate an intricate dynamic between these two cell types that point to
important reciprocal interactions[1,19,20]. Although much has been understood about the
influence of neural crest cells on surrounding cell types, placode cells included, there is
still limited knowledge about how placodal neurons influence the migration and
differentiation of neural crest cells.

The earliest populations of neurons within a system are known as pioneer neurons,
as they migrate to targets first and provide a scaffold for later born neurons to grow
along[21]. While there is some understanding of the function of pioneer neurons within
vertebrates[22–24], much of our knowledge in this area has come from studies in
invertebrates[25–28]. Still, there is conflicting data about whether pioneer neurons are
necessary and unique cells, and additionally, their importance appears to be system-
dependent. Within the trigeminal ganglion, there is very limited information surrounding
the function and relevance of pioneer neurons. One study in mouse found the earliest
trigeminal neurons of the ophthalmic lobe reaching their targets by embryonic day (E)
9.5[29]. Within the chick, the use of a monoclonal antibody against beta-tubulin III (Tubb3)
has aided in understanding that placode cells express Tubb3 before they leave the
ectoderm, while neural crest cells express Tubb3 later, after they differentiate into
neurons[30]. This is in keeping with findings on the timeline of trigeminal sensory neuron
differentiation in the chick – placode- and neural crest-derived neurons become post-
mitotic and differentiate within 2-5 days and 4-7 days of incubation, respectively[15]. During
this time, the placodal neurons are beginning to grow towards the periphery and are
making their initial connections with their targets. This cumulative knowledge suggests
that placodal neurons are pioneers that serve as templates for the later differentiating
neural crest-derived neurons to grow and extend axons[30]; however, this has never been
definitively proved. Furthermore, molecules orchestrating this process remain unknown.
The migratory abilities of neural crest and placode cells, as well as the interactions
between the two to form the trigeminal ganglion[31], are mediated by cell-cell adhesion.
More specifically, cadherins, which are transmembrane proteins that function in
strengthening adhesion between adjacent cells, have been shown to play an important
role in gangliogenesis. The extracellular domain of a cadherin can participate in *cis* or
trans interactions. Although cadherins generally engage in homophilic interactions, they
are capable of interacting heterophilically as well[32–34]. The cadherin intracellular domain
contains two binding sites for catenin proteins. The interaction between a cadherin and
p120-catenin regulates the availability of the cadherin at the membrane and its ability to
engage in adhesion, while cadherin binding to β-catenin and α-catenin serves to link the
cadherin to the actin cytoskeleton[35].

The cadherin profile of a cell is dynamic and can change depending on the needs
of the cell. Premigratory cranial neural crest cells initially express Cadherin-6B prior to
their delamination and subsequent epithelial-to-mesenchymal transition, but once
migratory, begin expressing Cadherin-7[36], a type-II cadherin which aids in transient cell-
cell adhesion. In the chick, the first expression of neural cadherin (N-cadherin) is seen as
early as 24 hours into development and is localized to the neural plate. N-cadherin
expression persists as the neural plate grows and becomes the neural tube[37]. Further,
placode cells and their neuronal derivatives express N-cadherin prior to delaminating from
the ectoderm[38]. This is not surprising given that prior studies have shown that N-cadherin
possesses important roles during neurodevelopment including regulating neurite
outgrowth[39], axonal fasciculation[40], and proper synapse formation[41,42]. All of the listed
examples highlight adhesion-dependent roles of N-cadherin during neurodevelopment;
however, N-cadherin can also play adhesion-independent roles such as controlling neural
differentiation and acting as a modulator of various signaling pathways[43,44].

While N-cadherin expression in trigeminal placode cells is critical for initial stages
of gangliogenesis and the ability for early trigeminal neurons to condense[38], whether there
is a need for N-cadherin throughout the development of the trigeminal ganglion and its
nerves remains unknown. Furthermore, how the early expression of N-cadherin within
placodal neurons impacts all the neurons of the trigeminal ganglion is not understood.
Here, we show that placodal expression of N-cadherin is important throughout
gangliogenesis, with defects to morphology observed in both the ganglion as well as
projections to target tissues upon N-cadherin depletion from placode cells. Additionally,
we show a reduction to axonal projections both *in vivo* and *in vitro* that can be attributed
to a loss of N-cadherin-mediated adhesion. These results suggest that trigeminal neurons
depend on the adhesive function of N-cadherin, initially expressed in placodal neurons,
to facilitate their outgrowth.

MATERIALS AND METHODS

Chick embryos

Fertilized chicken eggs (*Gallus gallus*) were obtained from the Department of Animal and Avian Sciences at the University of Maryland and Centurion Poultry Incorporated (GA) and incubated at 37 °C in humidified incubators on their sides. Embryos were incubated to reach desired Hamburger Hamilton (HH) stages based on times provided therein[45]. Embryos were staged with this method or by counting the number of somite pairs (somite stage, ss).

Morpholinos

To reduce N-cadherin protein expression, a validated lissamine-tagged translation-blocking morpholino antisense oligonucleotide (MO) against N-cadherin was used (GeneTools, LLC)[38], with a sequence of 5'-GCGGCGTTCCCGCTATCCGGCACAT-3'. For control experiments, a 5 base pair mismatch N-cadherin control MO was used[38], with a sequence of 5'-GCGcCcTTCCCcCTATgCcGCACAT-3' (altered base pairs in lowercase). Both MOs were used at a final concentration of 500 μM and were diluted with ultrapure water and 10% sucrose (in 1X phosphate-buffered saline (PBS)).

***In ovo* unilateral placode electroporation**

To target trigeminal placode cells prior to their delamination, a unilateral electroporation was performed at HH10-11. Briefly, the N-cadherin MO or N-cadherin mismatch MO (MM) was expelled on top of the ectoderm on one side of the embryo using a fine glass needle. Next, a 30-gauge needle was used to puncture a small hole through the vitelline membrane so that electrodes could lay vertically above and below the ectoderm over which the MO had been placed. Three, 9V pulses were delivered over 50 ms at 200 ms intervals. After electroporations, eggs were resealed with tape and parafilm and grown at 37 °C to desired stages of interest. Due to the inherent variations that arise during normal development, even when embryos are staged identically, the contralateral side of the electroporated embryo was a more appropriate control and was used for all *in vivo* studies. Initial experiments comparing N-cadherin MO-electroporated embryos to N-cadherin control MO-electroporated embryos were done to ensure none of the reported

201 phenotypes was due to the electroporation method. All embryos were screened for
202 successful MO electroporation based on the red fluorescence of the MO prior to collection
203 and use in subsequent assays.
204
Immunostaining

Electroporated embryos were grown to desired stages of interest, collected, and
fixed in 4% paraformaldehyde (PFA) overnight at 4°C with agitation. Following fixation,
embryos were washed 3 x 10 minutes in 1X PBS and stored in 1X PBS at 4°C until further
processing. Primary antibodies used included Tubb3 (Abcam, 1:500 for section, 1:250 for
whole-mount) and Sox10 (GeneTex, 1:250). Secondary antibodies used included
goat@rabbit 405 (ThermoFisher) and goat@mouse IgG$_{2a}$ 555 or 647 (SouthernBiotech).
Tissue sections: Embryos were processed through an increasing sucrose
gradient (5% and 15% sucrose (Fisher) in 1X PBS) either overnight at 4°C or until they
sank to the bottom of the vials. Next, they were processed through 7.5% gelatin (Sigma)
in 15% sucrose and 20% gelatin in 1X PBS, both at 37°C, either overnight or until the
tissue sank. Embryos were embedded in 20% gelatin, flash-frozen using liquid nitrogen,
and stored at -80°C until ready for use. Embedded embryos were then cryo-sectioned at
12μm on charged slides (VWR). Slides were stored at -20°C until ready for use. For
immunohistochemistry, slides were de-gelatinized by incubation in 1X PBS at 42°C for
15-20 minutes. Following 3 x 5-minute washes in 1X PBS, the slides were blocked with
10% heat-treated sheep serum (HTSS) in PBST (Sigma, 1X PBS + 0.1% Triton) at room
temperature for 1-2 hours in a humidified chamber, then rinsed in PBST (0.1%). Primary
antibodies were diluted in PBST + 5% HTSS, added to the slides, covered with a glass
coverslip (Fisher) to prevent sections from drying out, and incubated overnight at 4°C.
The following day, the coverslips were removed, and the slides were washed 4 x 30
minutes in PBST. Secondary antibodies were then diluted in PBST + 5% HTSS, added to
the slides, covered with a glass coverslip, and incubated overnight at 4°C. The next day,
slides were again washed 4 x 30 minutes in PBST and mounted with DAPI-fluoromount
media (SouthernBiotech).
Standard whole-mount: Embryos were blocked with 10% HTSS in PBST (0.1%)
for 2 hours at room temperature. Following a rinse with PBST, primary antibodies were

diluted in PBST + 5% HTSS and incubated overnight at 4°C with agitation. The following

day, the embryos were washed 4 x 30 minutes in PBST. Secondary antibodies were

diluted in PBTS + 5% HTSS and incubated overnight at 4°C with agitation. The next day,

embryos were again washed 4 x 30 minutes in PBST and stored in 1X PBS at 4°C until

they were either imaged or FRUIT cleared.

237   **Modified whole-mount for iDISCO clearing:** Fixed embryos were washed 2

times for 1 hour in 1X PBS and then were taken through an increasing methanol (MeOH

in 1X PBS) gradient (50% (1 x 1 hour), 80% (1 x 1 hour), 100% (2 x 1 hour)), before they

were bleached overnight in 5% H_2O_2 in 20% DMSO (Fisher)/methanol at 4°C. The next

day, samples were taken through the following series of washes: 2 x 1 hour in 100%

MeOH, 2 x 1 hour in 20% DMSO/MeOH, 1 x 1 hour in 80% MeOH, 1 x 1 hour in 50%

MeOH, 2 x 1 hour in 1X PBS, and 2 x 1 hour in PBST (0.2%). Samples were incubated

overnight in 20% DMSO/0.3M glycine (Fisher)/PBST at 4°C. The next day, samples were

blocked in 10% HTSS/10% DMSO/PBST overnight at 4°C. Embryos were washed 2 x 1

hour in 0.2% Tween-20 (Sigma)/1X PBS with 10 µg/mL heparin (Fisher, PTwH) at room

temperature before primary antibodies were added and incubated for 72 hours at 4°C.

Primary antibodies were diluted in 5% DMSO/5% HTSS/PTwH. Samples were washed in

PTwH for 24 hours before secondary antibodies were added (diluted in the same manner

as the primary antibodies), and samples were incubated with secondary antibodies for 48

hours at 4°C. Samples were then washed for 48 hours in PTwH before iDISCO clearing.

252   **Explant immunocytochemistry:** Explant slides were washed once with 1X PBS,

followed by fixation in 4% PFA for 1 hour at room temperature with agitation. Following

fixation, slides were rinsed with 1X PBS and stored in 1X PBS at 4°C until ready for

immunocytochemistry. A hydrophobic perimeter was drawn around the slides to prevent

dehydration of tissue using an ImmEdge Pen (Vector Laboratories), and explants were

taken through blocking, primary antibody incubation, and secondary antibody incubation

in the absence of coverslips as described in the tissue section methods. Slides were

mounted with DAPI-fluoromount media (SouthernBiotech) and covered with a glass

coverslip.

Tissue clearing

As embryos progress through development and tissue structures surrounding the trigeminal ganglion become more complex, visualization of the trigeminal ganglion after whole-mount immunohistochemistry became increasingly difficult. Therefore, clearing methods to render the tissues transparent were performed. At early stages (up to HH22), the FRUIT[46] tissue clearing protocol was used, whereas later stages required iDISCO[47] tissue clearing. With both protocols, MO signal was quenched during the clearing processes.

FRUIT: Following the standard whole-mount immunohistochemistry protocol, embryos were progressed through increasing concentrations of a fructose/urea cocktail (35%, 40%, 60% and 80%) for a minimum of 8 hours, up to 24 hours at room temperature, depending on the stage of the embryo. The time spent in a certain concentration of FRUIT buffer was determined based on how quickly the sample was rendered transparent. A sample was only moved up to the next FRUIT buffer once it had visibly changed transparency. The cocktail, containing 8M urea (Sigma), 0.5% (v/v) alpha-thioglycerol (Fisher), and increasing concentrations of fructose (Sigma), was made by first dissolving the fructose in ultrapure water at 37 °C. Once dissolved, the urea was added, and the solution was rocked at room temperature until homogenous. Lastly, alpha-thioglycerol was added, and the buffers were stored at 4°C until use. Following clearing, embryos were imaged in 60% or 80% (depending on the stage of the embryo) FRUIT buffer in glass-bottomed petri dishes. Typically, embryos up to HH19 were imaged in 60% FRUIT while HH20-22 staged embryos were imaged in 80% FRUIT.

iDISCO: Following the modified whole-mount immunohistochemistry protocol, embryos were washed in PBST (0.2%) for a minimum of 48 hours. Samples were incubated in 50% (v/v) tetrahydrofuran (THF, Sigma)/water overnight at room temperature in glass vials, making sure that over 90% of the vial was filled with the solution. The following day, embryos were then incubated in 80% THF/water for one hour at room temperature, followed by 2 x 1-hour incubations in 100% THF. Next, samples were incubated in 100% dichloromethane (Sigma) for 1-2 hours (or until they sank to the bottom of the vial). Lastly, embryos were incubated in dibenzyl ether (DBE, Sigma) overnight and imaged the following day in DBE. Embryos were stored in DBE at room temperature.

iDISCO-cleared embryos were imaged in a glass petri dish containing a 3D chamber[47],

which was printed in Terrapin Works, part of the School of Engineering at the University

of Maryland.

TUNEL assay

To assess apoptosis, the *In Situ* Cell Death Detection Kit, Fluorescein (Roche) was

used according to manufacturer's instructions. Briefly, following secondary antibody

washes (4 x 30 minutes in PBST (0.1%)), slides were washed once for 30 minutes in 1X

PBS. Slides were re-fixed with 4% PFA for 20 minutes at room temperature, followed by

one 30-minute wash in 1X PBS. The TUNEL reaction mixture was added to the slides (75

µL per slide), and a glass coverslip was placed over them. Slides were incubated in the

dark for 1 hour at 37°C. Next, the coverslips were removed, and the slides were washed

3 x 5 minutes in 1X PBS. Lastly, slides were mounted with DAPI-fluoromount media

(SouthernBiotech).

Trigeminal ganglion explant assay

Wild-type HH13/14 embryos were dissected from eggs and stored on ice in 1X

PBS until dissections. Trigeminal ganglia were dissected on a sylgard-coated dish by first

bisecting the embryo and then carefully taking the area of the forming trigeminal ganglion,

making sure to exclude the eye and neural tube. Explants were cultured on 2-well glass

chamber slides (ThermoFisher) coated with Poly-L-lysine (0.01% PLL, Sigma) and

fibronectin (0.01%, Sigma). To make the coated slides, 1 mL of PLL (diluted in ultrapure

water) was added to each chamber on the glass slides and incubated inside a biosafety

cabinet for 30 minutes at room temperature. The PLL was removed, and slides were

washed 3 times with water. Next, 1 mL of fibronectin (diluted in DMEM media

(ThermoFisher)) was added to each chamber and incubated at 37°C for a minimum of 2

hours. The fibronectin solution was removed, and slides were stored at 4°C until needed.

Explants were grown in DMEM media supplemented with N2 (1:100, ThermoFisher) and

Penicillin/Streptomycin (1:100, Fisher).

N-cadherin function-blocking antibody experiments

After 24 hours of incubation, fresh media containing either the MNCD2 function blocking antibody (DSHB, 0.1µg/mL), or a normal rat IgG control (Santa Cruz Biotechnology, 1.6 µg/mL), was added to the explants to replace the current media, and explants were incubated overnight. The following morning, the slides were briefly rinsed with 1X PBS, and then fixed in 4% PFA for one hour at room temperature. The explants were washed 3 times in 1X PBS, and then processed for immunocytochemistry.

Confocal imaging

All images were acquired with the LSM Zeiss 800 confocal microscope with Airyscan detection (Carl Zeiss Microscopy, Thornwood, NY, USA). Laser power, gain, and offset were kept consistent for the different channels during all experiments where possible. ImageJ (NIH)[48,49] and Adobe Photoshop CC 2023 were used for image processing. For whole-mount imaging with Z-stacks, all embryos were imaged at a 5µm interval. For sections, images of at least five serial transverse sections were acquired and used for analysis. For all experiments, a minimum of five replicates were obtained.

Quantifications and statistical analyses

Surface area: For modified surface area measurements of the whole trigeminal ganglion, the OpV lobe, and the MmV lobes, maximum intensity Z-projections were first generated from Z-stacks taken through the trigeminal ganglion using ImageJ. The polygon tool was used to outline the perimeter of Tubb3 signal and area measurements were generated. Individual area measurements for each developmental stage grouping were averaged together and plotted to generate graphs.

Sox10 and TUNEL cell counting: N-cadherin MO-treated embryos were processed for section immunohistochemistry for Sox10 and Tubb3, followed by the TUNEL staining protocol. Once imaged, the individual channels were opened on ImageJ. With the Tubb3 channel image, the polygon tool was used to trace the outline of the trigeminal ganglion. The outline was added to Region of Interest (ROI) manager, and then used to crop both the TUNEL channel and Sox10 channel to only include the area of the image that contained neurons of the trigeminal ganglion. Once images from all

experiments were processed, they were analyzed using the Squassh[50] plugin from the
MOSAICsuite in ImageJ, as described[34]. Segmentation parameters were kept uniform for
all images in one experimental condition (Sox10 or TUNEL) and run as a batch function.
Following batch analysis, the quality of image segmentation was confirmed visually to
ensure accurate segmentation. If batch parameters were not sufficient, they were
adjusted on a case-by-case basis and the segmentation analysis was rerun. For TUNEL
analysis, the number of puncta (observed as objects) was calculated per image and
averaged to obtain an average number per replicate. Individual averages for each
developmental stage grouping were averaged together and plotted to generate graphs.
For Sox10 nuclei counting, a further adjustment was made to accurately account for all
nuclei in an image. After the segmentation was run, two values were found: the median
area of nuclear objects and the total area of all objects. The total area was divided by the
median nuclei size to generate an adjusted nuclei count. The adjusted count was found
per image and averaged to obtain an average number per replicate. Individual averages
for each developmental stage grouping were averaged together and plotted to generate
graphs.

Neuron branching: Z-stack images of the OpV branch of the trigeminal ganglion
were analyzed in ImageJ using custom ImageJ Macro language (IJM) scripts (available
upon request). Maximum intensity Z-projection images were saved as an 8-bit image.
Each image was input into the ImageJ macro as an 8-bit image where the user was
prompted to broadly and manually create a closed polygon selection that encompassed
the entire ophthalmic branch of interest and its distal neuronal extensions. Once the
region of interest was selected, the ImageJ macro iterated over the entirety of the image
with a vertical line selection and ran the command, *getProfile()* at 30µm intervals. Results
were exported at each interval and served as input into a separate script created in R (R
core team, Vienna, Austria, 2021) using the R programming language (available upon
request).

The peaks obtained via *getProfile()* at each interval across an image identified
unique neuronal branches. To identify individual peaks in each iteration, the
smooth.spline() function was used in base R to fit a curve with known maxima and minima
to the profile outputted from the ImageJ macro in each iteration. A subsequent results

387 table was produced for each image documenting the number of maxima at 30μm intervals
388 from the sample edge. Data were transformed, if necessary, so that for all images
389 assessed, increasing distance from baseline represented an increased distance from the
390 trigeminal ganglion. Data for individual replicates for each condition were aligned in
391 PRISM (GraphPad Software, Inc.) so that quantification of neuronal branching over a
392 distance concordantly began with the first measurement found. Data were displayed as
393 the mean quantity of neuronal branches ± SEM and plotted to generate graphs.

394   **Bundle width:** Using maximum intensity Z-projections, bundle width
395 measurements for both the lateral and medial nasal nerves were generated using the
396 straight line tool on FIJI. The diameter of individual nerve bundles (prior to the point where
397 nerve fascicles branched into individual neurons) were measured based on Tubb3 signal
398 and summed together to yield one value per embryo. Individual bundle widths were
399 averaged together for N-cadherin MO-treated embryos and their contralateral control
400 sides and plotted to generate graphs.

401   **Innervation area:** Using maximum intensity Z-projections, innervation area
402 measurements for both the lateral and medial nasal nerves were generated using the
403 polygon tool on FIJI. A border was drawn around the Tubb3 signal following extensive
404 neuron branching. Individual area measurements were averaged together for N-cadherin
405 MO-treated embryos and their contralateral control sides and plotted to generate graphs.

406   **Explant analysis**: Each low magnification, tiled image of the explants was first
407 converted to 8-bit images on FIJI. Using the polygon function, the explant tissue piece,
408 as well any background signal not associated with axons, was cleared for the images.
409 Next, the ridge detection plug-in was used to highlight neurons to obtain median axon
410 length and the detected segment version of the image was saved. Using the detected
411 segment images and the concentric circle plugin, 20 concentric circles were overlayed on
412 the images, with the center circle being placed in the center of the explant. The number
413 of axonal intersections across each circle of increasing diameter was counted and
414 compiled into a spreadsheet where the averages ± SEM were plotted in 175μm intervals.
415
416

RESULTS

N-cadherin knockdown in placode cells disrupts trigeminal ganglion development

While the expression of N-cadherin in cranial placode cells as well as sensory and motor neurons has been established[51–54], along with N-cadherin function in early trigeminal ganglion assembly[38], the role of N-cadherin in later trigeminal ganglion development in both placode cell- and neural crest cell-derived neurons is unknown. To elucidate the function of N-cadherin-expressing placodal neurons in the formation and maturation of the trigeminal ganglion, a validated N-cadherin morpholino (MO) was unilaterally electroporated into trigeminal placode cells prior to their delamination[38]. The electroporated embryos were grown for up to five days post-electroporation and assembled into stage groupings based on 24-hour increments. Trigeminal ganglion and nerve morphology was analyzed following whole-mount immunohistochemistry with an antibody to Tubb3 to label neurons and tissue clearing to render samples transparent for imaging. Based on previous literature[15,30], the majority of neural crest cells begin differentiating and contributing to the neuronal population of the trigeminal ganglion beginning at E5, with some differentiating earlier at E4. Therefore, up to E4, it is assumed that the Tubb3 antibody is only labeling placode cell-derived neurons, and after labels neuronal contributions from both cell types indistinguishably.

During initial gangliogenesis (Figure 1A-A', E2.5), N-cadherin depletion from trigeminal placode cells leads to increased dispersal of neurons (Figure 1E) due to the inability of placodal neurons to properly intermix with undifferentiated neural crest cells. These results recapitulate what has been shown previously in that perturbation of N-cadherin during early gangliogenesis impacts proper aggregation of trigeminal placodal neurons[38]. Following the condensation of the trigeminal ganglion 24 hours later, placodal neurons can assemble into the forming trigeminal ganglion; however, surprisingly, the opposite phenotype is now observed. We note a statistically significant reduction in the size of the trigeminal ganglion on the electroporated side (Figure 1B-B', E, E3.5) compared to the contralateral control (untreated side) of the same embryo (and to control MO-treated embryos, Supplemental Figure 1). One day later (Figure 1C-C', E4.5), as neural crest cells start differentiating into neurons, the reduction of the trigeminal ganglion persists and it is significantly smaller (Figure 1E). Although the trigeminal ganglia that

were electroporated are visually still smaller at E5.5, they no longer show statistically
significant differences in area (Figure 1E). Lastly, at E6.5, the surface area between the
contralateral control and the electroporated trigeminal ganglia are not significantly
different (data not shown, Figure 1E).
Following the results from the whole trigeminal ganglion surface area
measurements, we next sought to discern whether N-cadherin had differential importance
within the OpV and the MmV placodes, seeing as these placodes arise distinctly from one
another[12,13]. Thus, surface area measurements were repeated, this time separately
measuring the OpV and MmV lobes and nerves from the trigeminal ganglion.
Interestingly, N-cadherin knockdown leads to a statistically significant increase in
dispersal of MmV neurons, with no impact to OpV neurons, at E2.5 when the ganglion is
beginning to assemble (Figure 1F, G). However, after condensation (E3.5-E5.5), N-
cadherin depletion did not impact the MmV lobe (Figure 1G) but did lead to a statistically
significant reduction in OpV lobe size (Figure 1F), correlating with what we observed
qualitatively. At E6.5, the size of both lobes was approximately the same. Given the drastic
effects of N-cadherin knockdown on OpV compared to MmV (Figure 1F, G), we chose to
focus on the development of the OpV lobe in subsequent experiments.
465
N-cadherin knockdown in placode cells does not lead to changes in cell death or
the number of neural crest cells within the trigeminal ganglion
To assess whether the decreased size of the trigeminal ganglion was due to an
increase in cell death and/or a change in the contribution of neural crest cells, a TUNEL
assay and Sox10 immunohistochemistry, respectively, were performed on tissue sections
take through the trigeminal ganglion. Only the stage groupings that showed significant
differences in size from Figure 1 were examined (HH16-18, HH19-22, HH23-25). These
experiments revealed no differences in cell death or the number of neural crest cells
between the contralateral side of the embryo (Figure 2A-A''') and the N-cadherin MO-
treated side (Figure 2B-B''', C, D). Apoptosis within the trigeminal ganglion was evident
on both the contralateral control (Figure 2A') as well as the N-cadherin MO (Figure 2B')
side, with no significant differences in the amount of TUNEL signal between the two
(Figure 2C). Furthermore, there is little co-localization of MO-positive cells and the TUNEL

signal, indicating that MO-expressing cells are not dying (Figure 2B', arrowheads). When examining TUNEL distribution, the TUNEL signal did not preferentially co-localize with either Tubb3 (marking neurons; Figure 2A", B"), or Sox10 (marking undifferentiated neural crest cells and newly differentiated glia).

Intriguingly, these data revealed less overall cell death occurring in the first stage grouping (initial trigeminal ganglion assembly) compared to the later stage groupings. Following condensation during these later stage groupings, however, it does not appear that there is a trend in the amount of cell death, as the average number of TUNEL-positive cells is not significantly different between the HH19-H22 and H23-H25 group (Figure 2C). Regarding the number of Sox10-positive nuclei present within the trigeminal ganglion after N-cadherin knockdown, there is also no statistically significant difference between the contralateral control and the electroporated sides (Figure 2D). These results indicate that the decrease in trigeminal ganglion size is not due to increased cell death or a reduction in the number of neural crest cells present in the trigeminal ganglion.

Depletion of N-cadherin from trigeminal placode cells disrupts OpV target innervation by the lateral nasal branch

Our TUNEL and Sox10 data suggest there is no impact on the neuron cell body and glial contribution to the ganglion upon N-cadherin knockdown in placode cells. As such, we sought to discern whether the neuronal projections, or axons, to target tissues might be impacted instead. We first examined the lateral nasal branch, one of the nerve branches of the OpV lobe, at embryonic stage group HH23-25, as by this point, the branch is a prominent structure within the embryonic head[55]. To this end, we used whole-mount immunohistochemistry with an antibody to Tubb3, which labels both placodal and neural crest-derived neurons at this stage. These experiments revealed a substantial reduction in the axons of this branch after N-cadherin MO treatment (Figure 3A-A'). The corresponding nerve on the contralateral control side appeared to have more nerve bundles as well as individual axons growing towards the rostral side of the embryo (Figure 3A), while the experimental side often had only one nerve bundle and many fewer individual axons (Figure 3A').

To quantify this apparent decrease in nerve outgrowth, the number of axons was
counted at 30µm intervals across each image for the contralateral and experimental
sides. While initially there is no significant change between the groups, as the main axon
bundle splits into smaller bundles, there are significant reductions on the N-cadherin MO-
treated side (Figure 3B). Next, bundle width and innervation area were also measured to
evaluate fasciculation (Figure 3C) and target innervation (Figure 3D), respectively. For
both bundle width and innervation area, the values were significantly reduced after N-
cadherin knockdown, indicating that N-cadherin is needed not only for proper nerve
outgrowth and fasciculation, but also for at least some of the individual neurons to send
axons to their targets (Figure 3C, D).

Depletion of N-cadherin from trigeminal placode cells leads to specific defects in
OpV innervation by the medial nasal nerve

Next, we examined another branch of the OpV lobe, the medial nasal branch, using
the same types of analyses performed in Figure 3. Although the medial nasal nerve is
beginning to form in HH23-25 staged embryos[55], there was more variability in the
development of this branch during these stages of development. Therefore, we chose to
examine the HH26-27 stage grouping due to more consistent morphological development
among embryos. Again, whole-mount immunohistochemical staining with Tubb3 revealed
a decrease in axonal projections between the control and experimental sides of the
embryos (Figure 4A-A'), indicating that N-cadherin is likely needed in a similar manner for
these branches of the OpV lobe. Interestingly, when the axon branches were counted
between the two groups, there was no significant difference at any point (Figure 4B).
Furthermore, both bundle width and innervation area were decreased for this branch, in
keeping with the results from the lateral nasal branch (Figure 4C, D). For both
measurements, the medial nasal branch was impacted to lesser extents compared to the
lateral nasal nerve (bundle width p=0.0009 for lateral vs p=0.0055 for medial, innervation
area: p=0.0029 for lateral vs p=0.0374 for medial). These results reveal that N-cadherin
controls the proper development of at least some nerve branches of the OpV lobe in
distinct ways.

N-cadherin is important for proper axon outgrowth of trigeminal neurons *in vitro*

Given the phenotypes observed *in vivo* that could not be attributed to changes in
cell number, we hypothesized that trigeminal axon outgrowth was facilitated by N-
cadherin in an adhesion-dependent manner. If trigeminal neurons were expressing less
N-cadherin, they may not be adhering to each other as well and would not be able to grow
out as effectively, leading to reduced outgrowth and target innervation. To test this, we
chose to utilize a function-blocking antibody against the extracellular domain of N-
cadherin, known as MNCD2[54,56], and test our hypothesis *in vitro* using explanted
trigeminal ganglion tissue. Explants were cultured *in vitro*, treated with MNCD2 or a rat
IgG control antibody for 24 hours, followed by Tubb3 immunocytochemistry and analysis.
Qualitatively, more axons appeared to grow out in control explants (Figure 5A) compared
to MNCD2-treated explants (Figure 5A'). Further processing of images was then
performed to detect individual axons (Figure 5B-B'), and then a modified intersection
analysis was run to measure changes to axon outgrowth (Figure 5C-C'). We discovered
a significant reduction to the amount of Tubb3 signal in the MNCD2-treated samples
(Figure 5D), indicating fewer trigeminal neurons after blocking N-cadherin adhesion.
Furthermore, our intersection analysis showed diminished axonal intersections overall
within the MNCD2-treated explants, pointing to reduced axonal outgrowth and complexity
(Figure 5E). Interestingly, when looking at the median length of a single axon between
the two treatments, there was no significant difference (Figure 5F). This indicates that
some trigeminal neurons are not dependent on N-cadherin and can properly grow out to
the periphery in the absence of N-cadherin. Taken together, these findings reveal that
blocking the adhesive function of N-cadherin reduces the overall outgrowth of some, but
not all, trigeminal neurons.

DISCUSSION

During neurodevelopment, cells experience several environmental changes as they migrate to their terminal locations and differentiate into specialized neurons. The first neurons to grow to target tissues, termed pioneers, set the path that all later born neurons will use as a template. Here we demonstrate a sustained role for N-cadherin during trigeminal gangliogenesis that shows the importance of this cell adhesion molecule during the formation and maturation of the trigeminal ganglion as it innervates its target tissues. Our findings show that reducing N-cadherin from trigeminal placode cells and their neuronal derivatives leads to defects in overall trigeminal ganglion morphology and negatively impacts the lateral and medial nasal branches of the ophthalmic lobe. Intriguingly, these phenotypes cannot be attributed to increased apoptosis or changes to the number of neural crest cells present within the trigeminal ganglion. We have found that N-cadherin-mediated adhesion is necessary for the outgrowth of the majority but not all trigeminal neurons. Our results suggest that a subpopulation of placodal neurons serve as the pioneers for the trigeminal ganglion and function independently of N-cadherin, while the remaining follower neurons of both cellular origins rely on N-cadherin for proper outgrowth and target innervation.

Depletion of N-cadherin in placodal neurons leads to morphological defects of the trigeminal ganglion at certain developmental timepoints

Although a role for N-cadherin-expressing placode cells has been shown during the coalescence of neural crest and placodal neurons as they initially generate the trigeminal ganglion[57], how these early N-cadherin-expressing neurons impact trigeminal gangliogenesis, and the ability of the trigeminal ganglion to properly innervate its targets as embryogenesis progresses, is not understood. We first sought to understand whether the phenotypes reported during early gangliogenesis[57] influenced trigeminal ganglion morphology at later developmental stages. Using relative surface area measurements we found that during early gangliogenesis N-cadherin knockdown leads to increased dispersal of trigeminal neurons and a larger surface area than controls, replicating previously published results[57]. Although there is increased dispersal of trigeminal neurons during early gangliogenesis, these neurons are eventually able to coalesce to form the

trigeminal ganglion as development progresses. Interestingly, once cells condense, N-
cadherin MO-treated trigeminal ganglia were significantly smaller up to 4.5 days into
development. At both 5.5 and 6.5 days, there were no statistical differences between the
N-cadherin knockdown and control groups when looking at the entire trigeminal ganglion.
We believe there are two likely reasons for this trend. First, MOs are typically only stable
for up to 4-5 days *in vivo*, reflecting a limitation of our technique[58]. Therefore, it is possible
that at the later stages that were examined, the MO was getting degraded, leading to
reduced phenotypic effects. This is further supported by the presence of MO-positive
puncta in the cytoplasm of placodal neurons, representing vesicles like lysosomes (not
shown). Second, it is possible that (additional) neural crest cells differentiate into neurons
between E4-5[2] to compensate for effects on placodal neurons after N-cadherin
knockdown.

The placode contribution of the trigeminal ganglion can be separated into two
distinct placodes, the ophthalmic and the maxillomandibular. While both placodes will give
rise to placodal neurons within the trigeminal ganglion, they express specific markers and
undergo differentiation at different times[59,60]. Ophthalmic placode cells express neuron-
specific markers earlier than maxillomandibular placode cells, and there is evidence that
a subset of ophthalmic placode cells differentiate and become post-mitotic while still in
the ectoderm. Conversely, the majority of maxillomandibular placode cells remain as
neuroblasts until after delamination from the ectoderm. Additionally, previous work has
reported more drastic phenotypes in the OpV lobe compared to the MmV lobe after N-
cadherin knockdown in placode cells[57]. Taking this into consideration, we next chose to
separately measure the surface areas of the OpV lobe and MmV lobe to determine
whether the requirement for N-cadherin was different between the two placodes.
Interestingly, the results were near complete inverses of each other. While there was no
significant difference during the first stage grouping measured for the OpV lobe, this time
point was the only place that showed significant results for the MmV lobe. Although it is
possible that N-cadherin is differentially required in the two placodes as they have unique
origins, we hypothesize that this difference is most likely due to the MO getting diluted out
over time in MmV. These neuroblasts continue to undergo cell division before
differentiating into neurons and becoming post-mitotic, whereas the OpV placode cells

are either already post-mitotic or about to differentiate when the electroporation is
performed, leading to more discernable phenotypes. When analyzing the OpV lobe, we
noticed that E3.5 showed the greatest difference in size of all the time points measured.
Together with previous data, our results indicate that knockdown of N-cadherin does not
impact the ability of the placode cells to migrate to the location of the trigeminal ganglion[57],
nor does it lead to increased apoptosis that could explain the changes to size. This time
point represents when the majority of placodal neurons have initiated growth to their
target tissues[30], and therefore might explain why the phenotype is the most drastic at this
stage. If N-cadherin is needed by some placodal neurons for axon outgrowth, then its
reduction would provide a molecular explanation for the results we observed.
638
Morphological changes after N-cadherin knockdown from placode cells cannot be
attributed to increased cell death or changes to neural crest cells within the
trigeminal ganglion
Given the changes to trigeminal ganglion morphology observed upon N-cadherin
depletion from placode cells, we next sought to discern whether these phenotypes were
due to changes in the number of cells present within the trigeminal ganglion. As depletion
of N-cadherin from placode cells does not impact ingression[57], we sought to evaluate the
other critical cell population, the neural crest, and cell death in the trigeminal ganglion. To
this end, we examined the number of neural crest cells and apoptosis, one or both of
which could also decrease the size of the trigeminal ganglion. Our results indicate no
change in the number of neural crest cells within the trigeminal ganglion nor no increase
in apoptosis within the trigeminal ganglion after N-cadherin knockdown. The TUNEL
signal did not preferentially co-localize with MO-positive cells and neural crest and
placode cells were equally apoptotic, supporting the notion that one cell type is not
undergoing more apoptosis than the other during trigeminal gangliogenesis. Therefore,
apoptosis is not a driving factor behind the reduction in size of the trigeminal ganglion
after N-cadherin knockdown. Interestingly, we observed that the majority of TUNEL signal
was found adjacent to Sox10-positive nuclei, which may be identifying future glial cells.
This phenomenon, seen previously[61,62] suggests that glial cells aid in the removal of
debris during gangliogenesis. When looking at general trends of cell death occurring

within the trigeminal ganglion over development, the first time point had decreased apoptosis compared to the other two time points. This can be explained by the presence of a larger number of neurons initially that decreases over time as trigeminal neurons receiving target-derived growth factors specialize and differentiate, while the ones that do not eventually perish[63]. Given this, we anticipate observing more cell death in the embryo at later stages.

Due to the reciprocal manner in which placodal neurons and neural crest cells engage with each other, it was important to examine potential changes to neural crest cell number, as it was possible that disrupting a cadherin on one cell type could impact the interactions with the other cell type. However, our results showed no difference to the number of neural crest cells at any of the stages examined. When evaluating the overall number of Sox10-positive cells within the trigeminal ganglion, we noticed a steady increasing trend as the embryo progressed through development. This represents the continued contribution of undifferentiated migratory neural crest cells to the ganglion and their eventual differentiation into neurons (Sox10-negative) and glia (Sox10-positive). It is important to note that we did not examine much later stages of trigeminal gangliogenesis when more neurons within the trigeminal ganglion would be neural crest-derived. As neural crest cells turn off Sox10 once they have differentiated into neurons, it is likely that the number of Sox10-positive cells would decrease at later stages of development, as it would then only mark glial cells.

Reducing N-cadherin from placodal neurons impacts OpV nerve branch innervation to target tissues

Traveling away from the cell bodies housed in the trigeminal ganglion, we next sought to understand how N-cadherin knockdown from placodal neurons impacted axonal outgrowth to target tissues. We chose to examine two branches of the OpV lobe, the lateral and medial nasal nerve branches. Although both branches arise from the cell bodies of the OpV lobe, we found that N-cadherin is needed in different manners between the two, with the lateral medial branch exhibiting more drastic changes. When examining the number of axons traveling to target tissues through intersection analyses, we found there were multiple points that were significantly reduced for the lateral nasal branch.

Additionally, we observed a decrease in bundling of the nerve prior to extensive branching
and a reduction in innervation area. While the medial nasal nerve exhibited similar trends
for bundle width and innervation area, albeit to a lesser degree, the results from the
intersection analysis were not changed at any of the points measured. This contrasting
result - a smaller nerve bundle having similar axonal branching - suggests that N-cadherin
functions to adhere individual axons at targets to each other for the medial nasal branch
of OpV. Similar phenotypes after N-cadherin perturbations have been observed in other
systems as well[53,64,65].

Although both of these branches begin developing a similar stages during
development[55] we noticed much more variation in the medial nasal nerve in the HH23-25
embryo group. To this end, we chose to examine the next group, HH26-27, for more
consistency among the embryos. While it is possible that the difference in the stage of
embryo examined between the two nerve branches could explain the differences in
significance, we believe it is equally plausible that N-cadherin-expressing placodal
neurons are needed to different extents in the two branches. Since the contribution of
neural crest-derived neurons and placodal neurons has not been examined for these
nerve branches before, it is possible that neural crest- and placode-derived neurons
comprise the bulk of the medial nasal and lateral nasal nerves, respectively. Another
potential explanation for the differences observed between the nerves could be due to
when the branches develop. Our results suggest that the medial nasal branch develops
slightly later than the lateral nasal branch. If this is true, then the early knockdown of N-
cadherin could preferentially impact nerve branches developing earlier versus later.

While the majority of embryos analyzed showed reduction to the nerve branches,
a subset of embryos actually showed increased measurements for bundle width and
innervation area measurements due to increased nerve bundles and increased individual
axons at targets. This suggests that while the overt phenotype of N-cadherin knockdown
leads to a reduced ability of axons to grow along other neurons and properly reach
targets, a subset of neurons, potentially the earliest ones to extend axons, is still able to
grow. However, these neurons, also lacking N-cadherin, cannot adhere to surrounding
axons in a proper manner, leading to these phenotypes.

Homophilic N-cadherin-mediated interactions facilitate the outgrowth of some trigeminal neurons

To understand whether the phenotypes seen *in vivo* were due to an adhesion-dependent role of N-cadherin during trigeminal gangliogenesis, we chose to utilize a function-blocking antibody that binds to the extracellular domain of N-cadherin and prevents its interactions with other cadherins. Multiple conditions, including concentration of antibody and incubation times of explants, were tested prior to beginning experimentation. We chose a concentration of MNCD2 that showed phenotypic changes to the explants without eliciting cellular responses indicative of apoptosis. Additionally, we noted that regardless of the incubation time of the explants prior to incorporating the MNCD2 antibody into the media, there was some axon outgrowth occurring. This suggests that some trigeminal axons are dependent on N-cadherin for outgrowth, while others are not.

Analysis of the N-cadherin function-blocking *in vitro* experiments led to three major conclusions. First, overall, fewer trigeminal neurons are able to migrate out of the explant and towards the periphery of the dishes in which they were grown. This was evident both visually as well as when the fluorescent intensity of each image was measured. Second, and in line with this, our intersection analyses indicate reduced outgrowth and less branching of neurons when N-cadherin adhesion is abrogated. Lastly, when looking at individual axon lengths, there were no significant differences in the median (and average, not shown) axon length when comparing rat IgG control-treated explants to MNCD2-treated explants. This result indicates that the subset of trigeminal neurons that are unaffected by blocking N-cadherin function are reaching the periphery of the explants in the same manner as their control counterparts, suggesting N-cadherin is not important for the outgrowth ability of these neurons. This conclusion models what was observed *in vivo* when examining target innervation by the lateral and medial nasal nerve branches. In both instances, N-cadherin knockdown did not completely inhibit target innervation but instead reduced the number of axons successfully reaching target tissues. While this phenotype could be attributed to the mosaic nature of the MO electroporation, our *in vitro* results now offer another explanation, in that some neurons are not reliant on N-cadherin for their outgrowth.

Altogether, these findings reveal that some trigeminal axons rely on N-cadherin-

mediated adhesion to facilitate their outgrowth to targets, and mirrors what was observed

in vivo when N-cadherin was reduced from placodal neurons. We hypothesize that the

neurons that do not rely on N-cadherin are the true pioneers of the trigeminal ganglion.

These neurons, which migrate through surrounding tissues using guidance cues, appear

to be less reliant on cell adhesion compared to followers, which commonly utilize cell

adhesion molecules to grow along axon tracts established by the pioneers[66].

Furthermore, the cranial mesenchyme is also known to express N-cadherin[67], so it is

possible that interactions between neurons and surrounding cells also facilitate this

process. Patterned substrate assays for N-cadherin would be beneficial for addressing

this question.

CONCLUSION

      In this study, we have shown a sustained negative impact on the trigeminal ganglion after N-cadherin is depleted from placodal neurons. Changes to the size of the trigeminal ganglion, as well as reduced axonal outgrowth to target tissues, are observed at early stages of development as well as later stages when neural crest cells have differentiated into neurons. These morphological changes cannot be explained by increased apoptosis or changes to neural crest cell number, suggesting an issue with the ability of neurons to properly grow out in the absence of N-cadherin. Blocking the extracellular domain of N-cadherin, which prevents the protein from interacting with other cadherins, leads to reduced axonal outgrowth, replicated results seen *in vivo*. Collectively, our results highlight a long-term, critical role for N-cadherin during trigeminal gangliogenesis.

FIGURE LEGENDS

Figure 1. Knockdown of N-cadherin in placode cells disrupts trigeminal ganglion development. (A-D') Representative images of the lateral view of the chick embryo head after immunostaining for Tubb3 and FRUIT (A-B') or iDISCO (C-D') clearing methods from stage groupings (HH16-18: A-A' (n=10), HH19-22: B-B' (n=10), HH23-25: C-C' (n=14), HH26-27: D-D' (n=13), not shown: HH28-30 (n=6)). Scale bar in (A) is 250μm and applies to (A'-B'). Scale bar in (C) is 250μm and applies to (C'-D'). (E) Quantifications of relative surface area of trigeminal ganglia taken from maximum intensity Z projections. (F) Quantifications of relative surface area of only the OpV lobe taken from maximum intensity Z projections. (G) Quantifications of relative surface area of only the MmV or MxV/MnV taken from maximum intensity Z projections. Average surface areas were graphed as histograms with error bars representing SEM. Arrowheads in (A'), (B'), (C'), and (D') indicate reduction in OpV lobe size in N-cad MO samples compared to controls. Paired t-tests were performed using GraphPad Prism and adjusted for multiple comparisons. (*) $p < 0.05$, (**) $p < 0.01$, (***) $p < 0.001$, (****) $p < 0.0001$. Abbreviations: OpV: ophthalmic; MmV: maxillomandibular; MxV: maxillary; MnV: mandibular; ns: not significant.

Figure 2. Morphology changes in the trigeminal ganglion after N-cadherin knockdown in placode cells is not due to increased cell death or changes to neural crest cells. (A-B''') Transverse sections through the ophthalmic lobe of the trigeminal ganglion from representative HH19 embryos with fluorescent immunohistochemistry for Sox10 (blue, marks neural crest cells) and Tubb3 (purple, marks neurons) and TUNEL staining (green, marks double stranded DNA breaks). Scale bar in (A) is 50μm and applies to (B). Scale bar in (A') is 50μm and applies to (A'', A''', B'-B'''). (C, D) Five serial sections per embryo from stage groupings HH16-18 (n=6), HH19-22 (n=8), and HH23-25 (n=11) were imaged and used for quantification analysis. ROIs were drawn to label the border of the trigeminal ganglion using Tubb3 staining. Within each section, individual TUNEL-positive cells and Sox10-positive nuclei were counted and averages were generated for each embryo. Total average counts for stage groupings were graphed as histograms with

error bars representing SEM. Arrowheads in (B') mark MO-positive, TUNEL-negative

cells. Paired t-tests were performed using GraphPad Prism and adjusted for multiple

comparisons. Abbreviations: e, ectoderm; ns: not significant.

Figure 3. N-cadherin depletion from placode cells negatively impacts lateral nasal

nerve outgrowth and target tissue innervation. (A-A") Representative images of the

lateral view of chick embryo head following immunostaining for Tubb3 (marks neurons)

and clearing with iDISCO (n=12). Scale bar in (A) is 250µm and applies to (A'). (B) Axon

branch counts were generated using custom scripts on FIJI to measure the signal within

the ROI in 30µm intervals. Averages were plotted with error bars representing SEM. ** at

300µm, 420µm, 450µm, 480µm, and 510µm, **** at 330µm, 360µm, and 390µm. (C)

Bundle width was calculated as a summation of the width of the nerve bundle for each

embryo prior to extensive axon branching. Sums were averaged and plotted as

histograms with error bars representing SEM. (D) Innervation area of the Tubb3-positive

axons was calculated by measuring the ROI area where the axons terminate for each

embryo. Areas were averaged and plotted as histograms with error bars representing

SEM. A repeated measure ANOVA was used to calculate the statistical significance for

intersection analysis in (B) using GraphPad Prism. Paired t-tests were performed for (C,

D).

Figure 4. N-cadherin knockdown in placode cells negatively impacts medial nasal

nerve outgrowth and target tissue innervation. (A-A") Representative images of the

lateral view of chick embryo head following immunostaining for Tubb3 (marks neurons)

and clearing with iDISCO (n=13). Scale bar in (A) is 250µm and applies to (A'). (B) Axon

branch counts were generated using custom scripts on FIJI to measure the signal within

the ROI in 30µm intervals. Averages were plotted with error bars representing SEM. ns

at any of the distances measured. (C) Bundle width was calculated as a summation of

the width of the nerve bundle for each embryo prior to extensive axon branching. Sums

were averaged and plotted as histograms with error bars representing SEM. (D)

Innervation area of the Tubb3-positive axons was calculated by measuring the ROI area

where the axons terminate for each embryo. Areas were averaged and plotted as

histograms with error bars representing SEM. A repeated measure ANOVA was used to
calculate the statistical significance for intersection analysis in (B). Paired t-tests were
performed for (C, D).

Figure 5. Blocking the extracellular domain of N-cadherin reduces trigeminal
neuron axon outgrowth. (A-C') Work-flow process for *in vitro* analyses. (A-A')
Representative images of trigeminal ganglion explants after treatment with either control
rat IgG (n=7) or MNCD2 (n=9) antibody, respectively, followed by immunostaining for
Tubb3 (marks neurons). Scale bar in (A) is 1mm and applies to (A'). (B-B') Generated
images after ridge detection processing on FIJI for rat IgG or MNCD2 treated explants,
respectively. (C-C') Processed images overlaid with concentric circle function to count the
number of axonal intersections. (D) Mean gray values of explant images after rat IgG or
MNCD2 treatment. Averages were plotted with error bars representing SEM. (E)
Intersection analysis graph calculating the number of axonal intersections across
concentric circles of increasing diameter after function-blocking experiments. Averages
were graphed with error bars representing SEM. Intersections with statistical differences:
* at 175μm, 350μm, and 700μm, ** at 525μm. (F) Median length of axons of each explant
image after rat IgG or MNCD2 treatment. Averages were plotted with error bars
representing SEM. Unpaired t-tests were run for all graphs, with adjusted made for
multiple comparisons for (E).

Supplemental Figure 1. N-cadherin control MO-treated embryos show no
alterations in trigeminal ganglion morphology. (A-A') Representative images of the
lateral view of the chick embryo head after immunostaining for Tubb3 and iDISCO
clearing from stage group HH23-25: n=4. Scale bar in (A) is 250μm and applies to (A').
(B) Quantifications of relative surface area of trigeminal ganglia taken from maximum
intensity Z projections. Average surface areas were graphed as histograms with error bars
representing SEM. A paired t-test was performed using GraphPad Prism. Abbreviations:
OpV: ophthalmic; MmV: maxillomandibular; MM: mismatch control MO; ns: not significant.

Figure 1. Knockdown of N-cadherin in placode cells disrupts trigeminal ganglion

development.

Figure 2. Morphology changes in the trigeminal ganglion after N-cadherin knockdown in

placode cells are not due to increased cell death or changes to neural crest cells.

904   **Figure 3. N-cadherin depletion from placode cells negatively impacts lateral nasal nerve**

905   **outgrowth and target tissue innervation.**

Figure 4. N-cadherin knockdown in placode cells negatively impacts medial nasal nerve outgrowth and target tissue innervation.

935   **Figure 5. Blocking the extracellular domain of N-cadherin reduces trigeminal neuron**

936   **axon outgrowth.**

951   **Supplemental Figure 1. N-cadherin control MO-treated embryos show no alterations in**

952   **trigeminal ganglion morphology.**

ACKNOWLEDGMENTS

The authors would like to thank the National Institute of Dental and Craniofacial Research (R01DE024217) and the University of Maryland Graduate School for financially supporting this research. Additionally, we would like to thank Ms. Zoe Davidson and Ms. Reagan Armstrong for their technical assistance.

REFERENCES

1. Hamburger, V. Experimental analysis of the dual origin of the trigeminal ganglion in the chick embryo. *J. Exp. Zool.* **148**, 91–123 (1961).

2. d'Amico-Martel, A. & Noden, D. M. An autoradiographic analysis of the development of the chick trigeminal ganglion. *J. Embryol. Exp. Morphol.* **55**, 167–182 (1980).

3. Dubey, A., Rose, R. E., Jones, D. R. & Saint-Jeannet, J. Generating retinoic acid gradients by local degradation during craniofacial development: One cell's cue is another cell's poison. *genesis* **56**, e23091 (2018).

4. Noden, D. M. The migratory behavior of neural crest cells. *Symp. Oral Sens. Percept.* 9–36 (1973).

5. Lumsden, A., Sprawson, N. & Graham, A. Segmental origin and migration of neural crest cells in the hindbrain region of the chick embryo. *Development* **113**, 1281–1291 (1991).

6. Rodrigues-Da-Silva, M. A., De Espindola Da Silveira, G., Taufer, C. R. & Calloni, G. W. The mesenchymal potential of trunk neural crest cells. *Int. J. Dev. Biol.* **66**, 317–331 (2022).

7. Shellard, A. & Mayor, R. Integrating chemical and mechanical signals in neural crest cell migration. *Curr. Opin. Genet. Dev.* **57**, 16–24 (2019).

8. Canales Coutiño, B. & Mayor, R. Neural crest mechanosensors: Seeing old proteins in a new light. *Dev. Cell* **57**, 1792–1801 (2022).

9. Schlosser, G. Making senses development of vertebrate cranial placodes. *Int. Rev. Cell Mol. Biol.* **283**, 129–234 (2010).

10. Bellairs, R. & Osmond, M. *The Atlas of Chick Development*. (Elsevier/AP, Academic Press is an imprint of Elsevier, Amsterdam, 2014).

11. Graham, A. & Shimeld, S. M. The origin and evolution of the ectodermal placodes. *J.*

Anat. **222**, 32–40 (2013).

12. Begbie, J. Early Steps in the Production of Sensory Neurons by the Neurogenic Placodes.

Mol. Cell. Neurosci. **21**, 502–511 (2002).

13. McCabe, K. L., Sechrist, J. W. & Bronner-Fraser, M. Birth of ophthalmic trigeminal

neurons initiates early in the placodal ectoderm. *J. Comp. Neurol.* **514**, 161–173 (2009).

14. Shiau, C. E., Das, R. M. & Storey, K. G. An effective assay for high cellular resolution

time-lapse imaging of sensory placode formation and morphogenesis. *BMC Neurosci.* **12**, 37

(2011).

15. D'amico-Martel, A. & Noden, D. M. Contributions of placodal and neural crest cells to

avian cranial peripheral ganglia. *Am. J. Anat.* **166**, 445–468 (1983).

16. Theveneau, E. *et al.* Chase-and-run between adjacent cell populations promotes

directional collective migration. *Nat. Cell Biol.* **15**, 763–772 (2013).

17. Breau, M. A. & Schneider-Maunoury, S. Cranial placodes: Models for exploring the

multi-facets of cell adhesion in epithelial rearrangement, collective migration and neuronal

movements. *Dev. Biol.* **401**, 25–36 (2015).

18. Freter, S., Fleenor, S. J., Freter, R., Liu, K. J. & Begbie, J. Cranial neural crest cells form

corridors prefiguring sensory neuroblast migration. *Dev. Camb. Engl.* **140**, 3595–3600 (2013).

19. Moody, S. A. & Heaton, M. B. Developmental relationships between trigeminal ganglia

and trigeminal motoneurons in chick embryos. I. Ganglion development is necessary for

motoneuron migration. *J. Comp. Neurol.* **213**, 327–343 (1983).

20. Shiau, C. E., Lwigale, P. Y., Das, R. M., Wilson, S. A. & Bronner-Fraser, M. Robo2-

Slit1 dependent cell-cell interactions mediate assembly of the trigeminal ganglion. *Nat.*

Neurosci. **11**, 269–276 (2008).

21. Bate, C. M. Pioneer neurones in an insect embryo. *Nature* **260**, 54–56 (1976).

22. Eisen, J. S., Myers, P. Z. & Westerfield, M. Pathway selection by growth cones of

identified motoneurones in live zebra fish embryos. *Nature* **320**, 269–271 (1986).

23. Ghosh, A., Antonini, A., McConnell, S. K. & Shatz, C. J. Requirement for subplate

neurons in the formation of thalamocortical connections. *Nature* **347**, 179–181 (1990).

24. Gong, Q. & Shipley, M. T. Evidence that pioneer olfactory axons regulate telencephalon

cell cycle kinetics to induce the formation of the olfactory bulb. *Neuron* **14**, 91–101 (1995).

25. Klose, M. & Bentley, D. Transient pioneer neurons are essential for formation of an

embryonic peripheral nerve. *Science* **245**, 982–984 (1989).

26. Lin, D. M., Auld, V. J. & Goodman, C. S. Targeted neuronal cell ablation in the

drosophila embryo: Pathfinding by follower growth cones in the absence of pioneers. *Neuron*

14, 707–715 (1995).

27. Hidalgo, A. & Brand, A. H. Targeted neuronal ablation: the role of pioneer neurons in

guidance and fasciculation in the CNS of *Drosophila*. *Development* **124**, 3253–3262 (1997).

28. Clarke, A. *et al.* Dynamic morphogenesis of a pioneer axon in *Drosophila* and its

regulation by Abl tyrosine kinase. *Mol. Biol. Cell* **31**, 452–465 (2020).

29. Stainier, D. Y. & Gilbert, W. Pioneer neurons in the mouse trigeminal sensory system.

Proc. Natl. Acad. Sci. U. S. A. **87**, 923–927 (1990).

30.   Moody, S. A., Quigg, M. S. & Frankfurter, A. Development of the peripheral trigeminal

system in the chick revealed by an isotype-specific anti-beta-tubulin monoclonal antibody. *J.*

Comp. Neurol. **279**, 567–580 (1989).

31.   Steventon, B., Mayor, R. & Streit, A. Neural crest and placode interaction during the

development of the cranial sensory system. *Dev. Biol.* **389**, 28–38 (2014).

32.   Straub, B. K. *et al.* E–N-cadherin heterodimers define novel adherens junctions

connecting endoderm-derived cells. *J. Cell Biol.* **195**, 873–887 (2011).

33.   Basu, R. *et al.* Heterophilic Type II Cadherins Are Required for High-Magnitude

Synaptic Potentiation in the Hippocampus. *Neuron* **96**, 160-176.e8 (2017).

34.   Halmi, C. A., Wu, C.-Y. & Taneyhill, L. A. Neural crest cell-placodal neuron interactions

are mediated by Cadherin-7 and N-cadherin during early chick trigeminal ganglion assembly.

F1000Research **11**, 741 (2022).

35.   Niessen, C. M., Leckband, D. & Yap, A. S. Tissue organization by cadherin adhesion

molecules: dynamic molecular and cellular mechanisms of morphogenetic regulation. *Physiol.*

Rev. **91**, 691–731 (2011).

36.   Wu, C. & Taneyhill, L. A. Cadherin-7 mediates proper neural crest cell–placodal neuron

interactions during trigeminal ganglion assembly. *genesis* **57**, e23264 (2019).

37.   Hatta, K. & Takeichi, M. Expression of N-cadherin adhesion molecules associated with

early morphogenetic events in chick development. *Nature* **320**, 447–449 (1986).

38.   Shiau, C. E. & Bronner-Fraser, M. N-cadherin acts in concert with Slit1-Robo2 signaling

in regulating aggregation of placode-derived cranial sensory neurons. *Dev. Camb. Engl.* **136**,

4155–4164 (2009).

39. Riehl, R. *et al*. Cadherin Function Is Required for Axon Outgrowth in Retinal Ganglion

Cells In Vivo. *Neuron* **17**, 837–848 (1996).

40. Masai, I. *et al*. N-cadherin mediates retinal lamination, maintenance of forebrain

compartments and patterning of retinal neurites. *Development* **130**, 2479–2494 (2003).

41. Benson, D. L. & Tanaka, H. N-Cadherin Redistribution during Synaptogenesis in

Hippocampal Neurons. *J. Neurosci*. **18**, 6892–6904 (1998).

42. Jontes, J. D., Emond, M. R. & Smith, S. J. *In Vivo* Trafficking and Targeting of N-

Cadherin to Nascent Presynaptic Terminals. *J. Neurosci*. **24**, 9027–9034 (2004).

43. Punovuori, K. *et al*. N-cadherin stabilises neural identity by dampening anti-neural

signals. *Development* dev.183269 (2019) doi:10.1242/dev.183269.

44. Aubert, J., Dunstan, H., Chambers, I. & Smith, A. Functional gene screening in

embryonic stem cells implicates Wnt antagonism in neural differentiation. *Nat. Biotechnol*.

20, 1240–1245 (2002).

45. Hamburger, V. & Hamilton, H. L. A series of normal stages in the development of the

chick embryo. *Dev. Dyn*. **195**, 231–272 (1992).

46. Hou, B. *et al*. Scalable and DiI-compatible optical clearance of the mammalian brain.

Front. Neuroanat. **9**, 19 (2015).

47. Renier, N. *et al*. iDISCO: A Simple, Rapid Method to Immunolabel Large Tissue

Samples for Volume Imaging. *Cell* **159**, 896–910 (2014).

48. Schindelin, J. *et al*. Fiji: an open-source platform for biological-image analysis. *Nat.

Methods* **9**, 676–682 (2012).

49. Schneider, C. A., Rasband, W. S. & Eliceiri, K. W. NIH Image to ImageJ: 25 years of

image analysis. *Nat. Methods* **9**, 671–675 (2012).

50.   Rizk, A. *et al*. Segmentation and quantification of subcellular structures in fluorescence

1091   microscopy images using Squassh. *Nat. Protoc*. **9**, 586–596 (2014).

1092   51.   Itoh, K., Ozaki, M., Stevens, B. & Fields, R. D. Activity-dependent regulation of N-

1093   cadherin in DRG neurons: differential regulation of N-cadherin, NCAM, and L1 by distinct

1094   patterns of action potentials. *J. Neurobiol*. **33**, 735–748 (1997).

1095   52.   Inuzuka, H., Redies, C. & Takeichi, M. Differential expression of R- and N-cadherin in

1096   neural and mesodermal tissues during early chicken development. *Development* **113**, 959–967

1097   (1991).

1098   53.   Tan, Z.-J., Peng, Y., Song, H.-L., Zheng, J.-J. & Yu, X. N-cadherin-dependent neuron–

1099   neuron interaction is required for the maintenance of activity-induced dendrite growth. *Proc*.

1100   *Natl. Acad. Sci*. **107**, 9873–9878 (2010).

1101   54.   Redies, C., Inuzuka, H. & Takeichi, M. Restricted expression of N- and R-cadherin on

1102   neurites of the developing chicken CNS. *J. Neurosci. Off. J. Soc. Neurosci*. **12**, 3525–3534

1103   (1992).

1104   55.   Higashiyama, H. & Kuratani, S. On the maxillary nerve: The Maxillary Nerve in

1105   Vertebrates. *J. Morphol*. **275**, 17–38 (2014).

1106   56.   Bronner-Fraser, M., Wolf, J. J. & Murray, B. A. Effects of antibodies against N-cadherin

1107   and N-CAM on the cranial neural crest and neural tube. *Dev. Biol*. **153**, 291–301 (1992).

1108   57.   Shiau, C. E. & Bronner-Fraser, M. N-cadherin acts in concert with Slit1-Robo2 signaling

1109   in regulating aggregation of placode-derived cranial sensory neurons. *Development* **136**,

1110   4155–4164 (2009).

1111   58.   Bill, B. R., Petzold, A. M., Clark, K. J., Schimmenti, L. A. & Ekker, S. C. A Primer for

1112   Morpholino Use in Zebrafish. *Zebrafish* **6**, 69–77 (2009).

59.   Stark, M. R., Sechrist, J., Bronner-Fraser, M. & Marcelle, C. Neural tube-ectoderm

interactions are required for trigeminal placode formation. *Development* **124**, 4287–4295

(1997).

60.   Lassiter, R. N. T., Stark, M. R., Zhao, T. & Zhou, C. J. Signaling mechanisms controlling

cranial placode neurogenesis and delamination. *Dev. Biol.* **389**, 39–49 (2014).

61.   Wu, H.-H. *et al.* Glial precursors clear sensory neuron corpses during development via

Jedi-1, an engulfment receptor. *Nat. Neurosci.* **12**, 1534–1541 (2009).

62.   Leonard, C. E., Quiros, J., Lefcort, F. & Taneyhill, L. A. Loss of Elp1 disrupts trigeminal

ganglion neurodevelopment in a model of familial dysautonomia. *eLife* **11**, e71455 (2022).

63.   Basch, M. L., Bronner-Fraser, M. & García-Castro, M. I. Specification of the neural crest

occurs during gastrulation and requires Pax7. *Nature* **441**, 218–222 (2006).

64.   Reis, S. A. *et al.* Striatal neurons expressing full-length mutant huntingtin exhibit

decreased N-cadherin and altered neuritogenesis. *Hum. Mol. Genet.* **20**, 2344–2355 (2011).

65.   Shikanai, M. *et al.* Caveolin-1 Promotes Early Neuronal Maturation via Caveolae-

Independent Trafficking of N-Cadherin and L1. *iScience* **7**, 53–67 (2018).

66.   Jontes, J. D. The Cadherin Superfamily in Neural Circuit Assembly. *Cold Spring Harb.*

Perspect. Biol. **10**, a029306 (2018).

67.   Dady, A., Blavet, C. & Duband, J.-L. Timing and kinetics of E- to N-cadherin switch

during neurulation in the avian embryo. *Dev. Dyn.* **241**, 1333–1349 (2012).

1132

1133

1134

1135
1136
1137